Regional estimates of gross primary production applying the Process-Based Model 3D-CMCC-FEM vs. multiple Remote-Sensing datasets

Authors: D. Dalmonech[1,2,*], E.Vangi[1], M. Chiesi[3], G. Chirici[4], L. Fibbi[3], F. Giannetti[4], G. Marano[5], C. Massari[6], A. Nolè[7], J. Xiao[8], A. Collalti[1,2]

[1]Forest Modelling Laboratory, Institute for Agriculture and Forestry Systems in the Mediterranean, National Research Council of Italy (CNR-ISAFOM), 06128 Perugia, Italy

[2]National Biodiversity Future Center (NBFC), 90133 Palermo, Italy

[3]National Research Council of Italy, Institute of BioEconomy, (CNR-IBE), 50019 Sesto Fiorentino, Italy

[4]GeoLab - Laboratory of Forest Geomatics, Dept. of Agriculture, Food, Environment and Forestry, Università degli Studi di Firenze, 50145 Firenze.

[5]Forest Ecology, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland

[6]Research Institute for Geo-Hydrological Protection, National Research Council (CNR-IRPI), 06128 Perugia, Italy

[7]School of Agricultural, Forest, Food and Environmental Sciences University of Basilicata, 85100 Potenza, Italy

[8] Earth Systems Research Center, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824, USA

ORCID:

Dalmonech D.:	0000-0002-1932-5011
Chiesi M.:	0000-0003-3459-6693
Chirici G.:	0000-0002-0669-5726
Fibbi L.:	0000-0001-6985-6809
Giannetti F.:	0000-0002-4590-827X
Marano G.:	0000-0003-2600-984X
Massari C.:	0000-0003-0983-1276
Nolè A.:	0000-0002-5144-3421
Vangi E.:	0000-0002-9772-2258
Xiao J.:	0000-0002-0622-6903
Collalti A.:	0000-0002-4980-8487

Corresponding author: Daniela Dalmonech daniela.dalmonech@cnr.it

e-mail coauthors:

Chiesi M.:   marta.chiesi@ibe.cnr.it
Chirici G.:   gherardo.chirici@unifi.it
Fibbi L.:   fibbi@lamma.toscana.it
Giannetti F.:   francesca.giannetti@unifi.it
Marano G.:   gina.marano@usys.ethz.ch
Massari C.:   christian.massari@irpi.cnr.it
Nolè A.:   angelo.nole@unibas.it
Vangi E.:   elia.vangi@isafom.cnr.it
Xiao J.:   j.xiao@unh.edu
Collalti A.:   alessio.collalti@cnr.it

Key Points:

• We apply a process-based forest model on a regular grid at 1 km spatial resolution in a
Mediterranean region.

• Initial forest state is estimated using spatially explicit input data derived from remote
sensing and national forest inventory data.

• The 3D-CMCC-FEM shows comparably estimates in simulating both spatial and
temporally the gross primary production, when compared to independent satellite-based
products.

Running title: Regional estimates of gross primary production

***Key words*:** process-based forest model; wall-to-wall map; gross primary production; national

forest inventory; Mediterranean region;

Abstract

Process-based Forest Models (PBFMs) offer the possibility to capture important spatial and

temporal patterns of both carbon fluxes and stocks in forests, accounting for ecophysiological,

climate and geographical variability. Yet, their predictive capacity should be demonstrated not

only at the stand-level but also in the context of large spatial and temporal heterogeneity. For the

first time, we apply a stand scale process-based model (3D-CMCC-FEM) in a spatially explicit

manner at 1 km spatial resolution in a Mediterranean region in southern Italy. Specifically, we

developed a methodology to initialize the model that comprehends the use of spatial information

derived from the integration of remote sensing (RS) data, the national forest inventory data and

regional forest maps to characterize structural features of the main forest species. Gross primary

production (GPP) is simulated over the period 2005-2019 and the multiyear predictive capability

of the model in simulating GPP is evaluated both aggregated as at species-level by means of

independent multiple data sources based on different RS-based products. We show that the

model is able to reproduce most of the spatial (~2800 km^2) and temporal (32 years in total)

patterns of the observed GPP at both seasonal, annual and interannual time scales, even at the

species-level. These new very promising results open the possibility of applying the 3D-CMCC-

FEM confidently and robustly to investigate the forests' behavior under climate and

environmental variability over large areas across the highly variable ecological and bio-

geographical heterogeneity of the Mediterranean region.

1 Introduction

Forest ecosystems absorb globally ~2 Gigatonnes of Carbon (C) stocking the carbon in their biomass and soil, thus acting as a net carbon sink. In Europe alone, forest ecosystems, which cover about a 40%, currently act as a net carbon sink for ~315 Megatonnes of CO_2eq and compensate for about 8% of EU-27's total greenhouse emissions (Verkerk et al., 2022). However, adverse climate impacts such as heat waves and drought (Allen et al., 2015; D'Andrea et al., 2020, 2021; Schuldt et al., 2020) and increasing natural disturbance rates (Grünig et al., 2023; Patacca et al., 2023) are all stressors which have potentially significant effects on current and future forest dynamics, jeopardizing the European forest ecosystems functioning and their carbon mitigation potential under future climate change (Schuldt et al., 2020; Senf et al., 2020; De Marco et al., 2022).

Nevertheless, ground data scarcity and short-term monitoring efforts still represent major challenges in studying the effects of climate change on forest dynamics in Mediterranean areas because are characterized by a large ecosystem heterogeneity and a biogeographically diverse structure shaped by human activity (Gauquelin et al., 2018; Médail et al., 2019; Peñuelas et al., 2017). As the Mediterranean region are known as a climate change 'hotspot' (Dubrovský et al., 2014; Noce et al., 2016), experiencing already increasing frequency in extreme events such heat waves and droughts (Vogel et al., 2021) it is thus crucial in these areas to provide large-scale forest monitoring and eventually predict the future state of forest ecosystems. Recent efforts have addressed the shortage of ancillary or ground data by integrating National Forest Inventories (NFI) data and high-resolution remote-sensing (RS) data. These initiatives produced comprehensive wall-to-wall maps of various forest variables, such as growing stocks volumes or biomass (Nord-Larsen & Schumacher, 2012; Waser et al., 2017; Chirici et al., 2020; Giannetti et

al., 2022; Vangi et al., 2023). These maps represent meaningful data for and carbon cycle

assessment (e.g. Vangi et al., 2023). In parallel, process-based forest models (PBFMs) are

analytical tools developed and tested over a wide range of applications, because of their

capability in simulating forest ecosystem even on long-term dynamics (Vacchiano et al., 2012;

Bugmann & Seidl, 2022), carbon fluxes exchange and stocks (Chiesi et al., 2010; Reyer et al.,

2014; Reyer, 2015; Dalmonech et al., 2022; Mahnken et al., 2022) under external environmental

variability by accounting for population dynamics and inner physiological processes

mechanistically (Pretzsch et al., 2008; Vacchiano et al., 2012; Maréchaux et al., 2021). On the

other side, given the large amount of requested data for their initialization and parameterization,

such models are mostly run at a very local scale, i.e. site level (one hectare or a bit more), where

high-quality/measured ancillary data and meteorological data are available (Collalti et al., 2016;

Suárez-Muñoz et al., 2023). Yet, initializing stand scale PBFMs from actual, measured forest

state variables (or close to the observed states), rather than from equilibrium conditions, is the

desirable option to implicitly take into account the climate and management history of the site,

and to more realistically simulate the response of forests and their resilience, even in the context

of climate change, and natural and anthropogenic disturbances (e.g. Pretzsch et al., 2008;

Kannenberg et al., 2020; Zampieri et al., 2021).

Despite their undoubted utility, only recently, stand-scale PBFMs were applied on a regular grid

(Sanchez-Ruiz et al., 2018; Minunno et al., 2019) with the purpose of estimating aggregated,

country-level, carbon stocks and wood products. Yet, in southern Europe and in the

Mediterranean the ability of a PBFMs to simulate the GPP at large scale is crucial, but largely

overlooked, because photosynthesis respond sensitively to both meteorological and climate

variability and spatial heterogeneity at the daily to decadal scales (e.g. Mahnken et al., 2022;

Fernández-Martínez et al., 2023), therefore GPP can be, and has been, considered a good proxy

of the ecosystem physiological functionality especially in Mediterranean forests (Collalti et al.,

2018; Chen et al., 2023). The objective of this study is, thus, the application and the testing of

the biochemical, biophysical, process-based forest model 3D-CMCC-FEM (Collalti et al., 2014,

2016, 2018), on a regular grid at 1 km spatial resolution in the Mediterranean area, initializing

the model through the use of spatial information derived by integrating data of different nature:

NFI data, RS-based wall-to-wall map, and regional forest maps to characterize structural features

of the main forest species. The final aim is to simulate GPP at regional scale. As a case study, the

model was tested over one of the southernmost regions in Italy, the Basilicata Region, which, as

most regions in the Mediterranean basin, spans over a multitude of ecological, morphological

and soil-type gradients and climate conditions. The capability of the model to simulate the GPP

in terms of mean annual, seasonal and interannual variability is evaluated by comparing model

results against a portfolio of different independent RS-based GPP-estimates. The 3D-CMCC-

FEM GPP vs. RS-based GPP data agreement is presented and sources of uncertainties and

challenges of applying a PBFMs with the presented modeling strategy are also discussed.

2. Materials and Methods

2.1 Study Area

The Basilicata region has a spatial extent of about 10,000 km^2 and is located in southern Italy

(Figure 1a). It is characterized by typical Mediterranean climate conditions with hot and dry

summers and wet and mild winters. The region has been chosen to test the model in a complex

biogeographic area with pronounced environmental gradients. The territory is characterized by

about 47% of mountain areas represented by the Apennines Mountains, followed by hilly areas,

about 45%, and then plain. Average annual precipitation varies between ~500 and ~2000

millimeters (mm) per year, mirroring the orographic complexity of the region and the proximities

to the sea, with a west-to-east gradient from humid to dry sub-humid areas.

According to the last NFI (INFC 2015), forest vegetation and other wooded lands occupy

392,412 hectares (ha), about 39% of the region. Deciduous species cover 54% of the forest area

and are represented mainly by oaks spp (*Q. cerris* L., *Q. ilex* L.), which dominated the hilly areas

between 400 and 1200 m above sea level (a.s.l.), and European beech (*Fagus sylvatica* L.) the

main species above 1000 m a.s.l.. Coniferous species are less abundant and are represented

mainly by pines spp (*P. halepensis* Mill., and *P. nigra* J.F. Arnold), often used for reforestation

purposes (Figure 1b).

The study area was tessellated into a 1 km spatial resolution regional grid whose pixel area

represent the best compromise between the forcing variables and the operative model resolution,

for a total of ~10,073 pixels. The regional grid served as a spatial reference grid for resampling

the input data needed for the model initialization to 1 km spatial resolution.

2.2 The Process-based model 3D-CMCC-FEM

The 3D-CMCC-FEM ('*Three Dimensional – Couple Model Carbon Cycle – Forest Ecosystem*

Module') is an ecophysiological, biogeochemical, biophysical process-based model which

simulates the dynamic of carbon, water and nitrogen and the allocation through a cohort-

structured forest stands (Collalti et al., 2014, 2016, 2017, 2018, 2020, 2022; Marconi et al., 2017;

Dalmonech et al., 2022; Testolin et al., 2023), providing detailed output from daily to annual

time scale of carbon fluxes and stocks. The model simulates forest growth and structural

development at varying environmental conditions and different climate, atmospheric CO_2

concentrations and forest management scenarios (Collalti et al., 2018; Dalmonech et al., 2022;

Testolin et al., 2023).

The daily gross photosynthesis is simulated through the Farquhar–von Caemmerer–Berry

biochemical model (Farquhar et al., 1980), modified for sun and shaded leaves (De Pury &

Farquhar, 1997), and acclimated for temperature (Kattge & Knorr, 2007). The carbon and

nitrogen allocation schemes are described extensively in Collalti et al. (2016, 2019, 2020) and

Merganičová et al. (2019). Tree removal can occur via management (Testolin et al., 2023) or by

natural mortality (Collalti et al., 2018). Self-thinning, age-related, carbon starvation, and

background mortality represent the different types of mortalities simulated by the model (Collalti

et al., 2016). Soil hydrology is simulated by means of a one-soil layer bucket model with free

drainage. The plant water availability in the model is thus modulated by the soil depth (i.e.

rooting depth until which the water is uptake to sustain leaf transpiration), because of the zero-

dimensional soil model. Soil water stress operates on canopy exchange processes *via* stomatal

and biochemical pathways (e.g. photosynthesis). An in-depth description of the model's

underlying characteristics, effects of climate change and model parameter sensitivity and

uncertainty, as well as model limitations, is reported in Collalti et al. (2019).

2.3 Forest data source

The model requires the description of the forest structural characteristics: i.e. diameter at breast

height (DBH), tree height (H), stand density (number of trees per cell) and age class, in order to

be initialized and to run the simulations. In this work, to initialize the model for each grid cell of

the matrix, we used data from the second NFI for 2005 (www.inventarioforestale.org). The NFI

is based on a three-phase, systematic, unaligned sampling design with 1 km grid cells. In the first

phase, 301,300 points were extracted and classified using aerial orthophotos into forest/non-

forest categories. In the second phase, a field survey was carried out in a sub-sample of the first-

phase points falling in the forest category, to collect qualitative information such as forest type,

management, and property. Finally, in the third phase, for a sub-sample of 6782 points extracted

from the second-phase points, a dendrometric survey was carried out for circular plots of a 13 m

radius. All tree stems with DBH of at least 2.5 cm were callipered, and for a subsample, height

was measured.

Field-survey data from the NFI were used to produce a 'wall-to-wall' map of the forest basal

area at a 23 m spatial resolution. This map consists of random forests predictions of basal area

per hectare for all 23 m spatial resolution forest pixels. The random forests model was trained

using NFI plot-level data and Landsat and other RS-based datasets as predictors, including

climate information such as minimum, mean, and maximum temperature, and daily

precipitations from the E-OBS dataset, the land-only gridded daily observational dataset for

Europe (see section 2.4). The statistical model fitting and tuning steps were carried out using the

'*randomForest*' package in the statistical software R 4.0.5 (Liaw & Wiener, 2002). More

information about the procedure can be found in Chirici et al. (2020), Vangi et al. (2021) and

Giannetti et al. (2022). The pixel-level estimations of the basal area range between 5 and 43 m^2

ha^{-1} with a mean value of 12 $m^2 ha^{-1}$, that is in line with the range reported in the context of NFI

for the field plots data estimation.

The basal area data were then resampled to the regional grid at 1 km resolution. The resampled

amp was then masked according to the regional forest map by Constantini et al. (2006). This

map provided the forest type according to Barbati et al. (2007) and the development stage of the

forests. Estimation of the forest structural data used to initialize the model in each regional grid

cell, is described in section 2.5.1.

2.4 Forcing and soil data

The 3D-CMCC-FEM was forced with daily maximum (Tmax, °C) and minimum (Tmin, °C) air

temperatures, precipitation (P, mm day^{-1}), downward short-wave radiation at the surface (SW

MJ m^{-2} day^{-1}) and relative humidity (RH, %). Meteorological data for the period 2005-2019

were retrieved from the E-OBS v.23.1e gridded dataset (Cornes et al., 2018), which is provided

at 0.1° decimal degree resolution The E-OBS dataset has already been used in environmental

impact studies (e.g. Rita et al., 2020), climate scenarios bias correction (Dosio & Paruolo, 2011;

Rojas et al., 2011) and benchmark activities (Moreno & Hasenauer, 2016; Herrera et al., 2019;

Lorenz et al., 2019; Massari et al., 2020).

All the physical variables were bilinearly interpolated to the regional grid at 1 km resolution and

the temperature data were corrected for the topographic effect by applying a lapse rate correction

based on elevation differences between the E-OBS reference elevation and the finest and most

accurate DEM (Digital Elevation Model) currently available in Italy, obtained in the framework

of the TINITALY project. The TINITALYDEM is a national DEM of Italy at 10 m resolution

(Tarquini et al., 2009) and for the lapse rate correction it was resampled at 1 km resolution of the

regional grid. The lapse rate estimates of –5 °C km^{-1} for Tmax and –3 °C km^{-1} for Tmin, were

derived from termo-pluviometric ground station measurements over an elevation transect in the

Basilicata region.

The model was forced by global annual atmospheric CO_2 concentrations from Dlugokencky and

Tans (https://www.esrl.noaa.gov/gmd/ccgg/trends/), covering in total the years 2005-2019.

The model requires information on soil depth as well as soil texture for each grid cell. As a proxy

of soil depth, we used the estimated depth available to root from the European Soil Database

Derived Data product ESBD v2 (Hiederer, 2013) provided at 1 km resolution. The lower depth

value of each class of the map was attributed, and a maximum of 1 m for the rooting depth was

set. This boundary value can be considered a good approximation for European forests (Schenk

& Jackson, 2005). Soil texture as a percentage of clay, silt, and sand was estimated from the

pedological map of the region (year 2005).

The meteorological and geographic information were all re-projected using the same coordinate

reference system WGS84/UTM zone 33 North (EPSG: 32633), and then resampled onto the

regional grid at 1 km resolution. The main data analyses were performed using the computing

language *R* (R Core Team 2021). Key packages used for data preprocessing included '*terra'*

(Hijmans et al., 2022) and '*rgdal'* (Sumner & Hijmans, 2023)

2.5 Model simulations

2.5.1 Model initialization

The 3D-CMCC-FEM model (v.5.6) was applied on the regional grid at 1 km spatial resolution to

simulate the forest carbon dynamic starting in January 2005 until December 2019. The model

requires the description of the forest attributes at the beginning of simulation in order to be

initialized. The initial forest state was set according to a simplified model initialization of the

forest aboveground structural complexity, i.e. for each grid cell, we determined the dominant

forest species and estimated the average structural data: the average tree diameter at breast height

(DBH), the average tree height (H), the stand density, and the average age class which represent

the mandatory initial data for model initialization.

The following six key species were considered: European beech (*F. sylvatica* L.), Black pine (*P.*

nigra J.F. Arnold), Sweet chestnut (*C. sativa* Mill.), Turkey oak (*Q. cerris* L.), Aleppo pine (*P.*

halepensis Mill.) and Holm oak (*Q. ilex* L.) as representative of the most common forest types in

the study area. The regional forest map was used to define for each 1 km grid cell the dominant

forest species as the one covering the highest forest fraction (Figure 1b) and the average age

class based on the development stage (provided by the regional forest map along with the forest

class). Areas with dominance of maquis and other minor forest species (the latest accounts for

~3.9% of the region and are not currently parameterized in the model) were masked out from the

regional gird. The final dominant forest age classes, result in a total of ~2800 km^2, corresponding

to ~80 % of the entire regional forested area.

Tree density data of the NFI field plots were used to provide a representative estimate of the

forest density in each grid cell. To do so, we zonally averaged the density data according to the

regional grid and the dominant forest classes. The basal area map was then used in combination

with the density map data to calculate an average cell-level DBH and to provide a 1 km

resolution DBH map for each dominant species (data not shown).

To be consistent with the model processing and inherent logic, the H was calculated from the

average DBH by applying the calibrated Chapmann-Richard equation (Richards, 1959), which

links DBH and H, for each forest class with each cell. Starting from these mandatory structural

variables, i.e. DBH and H, the model self–initializes the other state variables: i.e. leaf, stem,

branches, coarse and fine root, reserves (non-structural carbon, NSC; which includes starch and

sugars) carbon and nitrogen pools using a species-specific parameterization at the beginning of

the model simulations. Species-specific model parameters (e.g. specific leaf area or maximum

stomatal conductance, see Table S1) were retrieved and calibrated from literature data.

Specifically, the species-specific allometric equations linking DBH to H and DBH to stem

biomass, were calibrated in this study Using the second NFI tree-level data.

2.5.2 Simulations settings

Due to the relatively short time period covered by the simulations, i.e. 2005-2019, we did not

consider any change in dominant species or land use. For the same reason, and because of the

spatial resolution, a constant thinning rate implemented each year was considered as the only

silvicultural intervention as similarly as in Gutsch et al. (2018). The thinning rate was an

approximation derived from the forest management guidelines of the region as set to a yearly

removal rate of 1%, corresponding to ~20% of biomass removed in 20 years. Forests in protected

areas of the Natura2000 network of the region are characterized by lower disturbance extension

compared to the other forested areas according to the disturbance maps produced by Francini et

al. (2021), and are assumed to be interested by a lower level of tree harvesting. For simplicity,

we assume, thus, that mortality in protected area is mainly caused by natural and background

mortality alone. Grid cells interested in fire events over the period 2005-2019, were identified

using the national dataset of burnt areas from forest fires, produced by the Italian Forest Service

(Comando Unità Forestali, Ambientali e Agroalimentari of Carabinieri). This dataset is acquired

through a ground survey using Global Navigation Satellite System receiver (GNSS) and is

available from 2005 to 2019. Grid cells where more than 30% of the forest area was interested in

fire events were excluded from the analysis as the model does not simulate extended natural

disturbances (i.e. fire and pests). This threshold was a compromise between excluding too many

grid cells and including too many fire-disturbed areas.

2.6 Remote sensing - GPP datasets

Datasets based on RS-based data (or modeled by forcing with remote sensed data) are the most suitable candidate to assess the overall model capability at reproducing the GPP over large areas, due to their continuous spatial and temporal coverage. In order to make a more complete and comprehensive agreement assessment, we selected gridded GPP estimates from different independent sources as reference datasets, which are:

2.6.1 GOSIF-GPP

The GOSIF GPP dataset (Li & Xiao, 2019b) is a recently developed GPP product that is based on the global OCO-2 based SIF product (Li & Xiao, 2019a). The GOSIF setup combines in a data-driven approach the remotely sensed sun-induced fluorescence (SIF), observed by the Orbiting Carbon Observatory-2 (OCO-2), the enhanced vegetation index (EVI) from the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite data, meteorological data, i.e. photosynthetically active radiation (PAR), vapor pressure deficit (VPD), and air temperature obtained from the NASA reanalysis MERRA-2 data set to return a gridded SIF dataset (Li & Xiao, 2019a). Established relationships between the original OCO-2 SIF and flux tower GPP (Li et al., 2018; Xiao et al., 2019) were then used to provide the final gridded GPP product. For the model-data comparison in this study we used the monthly and annually aggregated ensemble mean of eight different GPP estimates resulting from different GPP-SIF relationships, (http://globalecology.unhedu). This GPP product, hereinafter simply referred as 'GOSIF', provides GPP estimates at 0.05 degree (corresponding to ~5 km) spatial resolution aggregated on a monthly time step, over the years 2005-2019.

2.6.2 CFIX-GPP

The CFIX dataset provides gridded GPP values covering Italy. The estimates are obtained

combining meteorological and remotely sensed data within a Light Use Efficiency modeling

approach (Veroustraete et al., 2002; Maselli et al., 2006). The original model version was further

modified to simulate the GPP in water-limited, Mediterranean forest ecosystems by Maselli et al.

(2009), who introduced a short-term water stress factor based on daily meteorological data. In

particular, the authors utilized the 1-km normalized difference vegetation index (NDVI) from the

Spot-VEGETATION imagery to linearly retrieve the fraction of absorbed photosynthetically

active radiation (APAR), and the downscaled E-OBS meteorological data (both air temperature

and precipitation)(Maselli et al., 2012; Fibbi et al., 2016) to retrieve the GPP of all Italian forests

at 1-km spatial resolution. The accuracy of the product was assessed by comparison to Eddy-

Covariance GPP data (Pastorello et al., 2020) collected at several EC sites spread over the Italian

peninsula and for different forest types, showing satisfactory performances (Chirici et al., 2016).

The currently utilized product, provides forest GPP estimates at 1 km spatial resolution

aggregated on a monthly time step, over the years 2005-2013.

2.6.3 FLUXCOM-GPP

The FLUXCOM dataset is an upscaled product derived from different Machine Learning-based

approaches (ML) combining data from the FLUXNET network of eddy covariance towers with

RS and meteorological data as predictors (Jung et al., 2019; Pastorello et al., 2020). In this study

we used the FLUXCOM-RS dataset which embeds in its statistical processing, RS land products

at 8-day resolution from the MODIS instrument, such as EVI, fAPAR, and the land surface

temperature (see Jung et al. (2019) for a thorough description of the FLUXCOM products).

Compared to other products of the FLUXCOM-database, the FLUXCOM-RS dataset has a finer

spatial resolution, 0.083 degree (corresponding to ~8 km) spatial resolution, allowing to better

deal with the complex topography of the region under study.

The FLUXCOM-RS dataset, hereinafter simply referred to as FLUXCOM, has been widely used

as a reference dataset in several inter-comparison studies with both model and other reference

datasets (O'Sullivan et al., 2020; Wang et al., 2021; Zhang & Ye, 2021). The selected dataset is

provided as the ensemble mean of monthly data computed from multiple ML algorithms

covering the period 2000-2015 (**www.fluxcom.org**).

2.7   3D-CMCC-FEM GPP versus RS-based GPP

For comparability with the 3D-CMCC-FEM GPP results, all the RS-based datasets were

resampled to the regional grid at 1 km resolution and reprojected using the WGS84/ UTM zone

33 North coordinate reference system (EPSG: 326633) and compared at both spatial and

temporal level against the 3D-CMCC-FEM GPP data. Results at species-level are first shown as

aggregated data over regional level and then shown at species-level.

2.7.1 Spatial variability analyses

The spatial agreement between the 3D-CMCC-FEM GPP and the RS-based GPP data was

assessed by considering: Root Mean Square Error (RMSE), Relative Difference (RD, expressed

as: (modelled – observed) / observed * 100, where 'observed' stands for RS-based data and

'modeled' for 3D-CMCC-FEM GPP), Pearson's correlation (r), and the SPAtial EFficiency

(SPAEF) metrics. The SPAEF (see Eq. 1) is an integrated evaluation index which considers the

Pearson's spatial correlation (r), the fraction of the coefficient of variation

$\beta = (\sigma_{sim}/\mu_{sim})/(\sigma_{obs}/\mu_{obs})$ and the model-data histogram intersection

$\gamma = \sum_{j=1}^{n} \min(K_j/L_j)/\sum_{j=1}^{n} K$, where n is the number of bins in the histogram and K and L are

the histogram for observed and simulated data (Koch et al., 2018). SPAEF is equal to 1 in the

case of a perfect match and 0 in case of mismatch.

$$SPAEF = 1 - \sqrt{(r-1)^2 + (\beta-1)^2 + (\gamma-1)^2}$$ Eq. 1)

The statistics were computed on the mean annual GPP and seasonal GPP: i.e. winter (December,

January, February: DJF), spring (March, April, May: MAM), summer (June, July, August: JJA)

and autumn (September, October, November: SON) for 3D-CMCC-FEM and RS-based data.

2.7.2 Temporal variability analyses

The temporal agreement between the 3D-CMCC-FEM GPP and the RS-based GPP data was

assessed by considering RMSE, RD, the Fractional Variance (FV), which returns values bounded

between –2 and 2 and it is equal to 0 when 'modelled' and 'observed' have the same variance

(Janssen & Heuberger, 1995). The Pearson's correlation coefficient r was also used at grid-cell

level, which returns the direction of the correlation i.e. model-data correspondence of the sign of

year to year variability. Conversely to the spatial analysis the mean seasonal cycle (MSC) was

also considered here. Anomalies were calculated for the model and data by first removing the

long-term linear trend, and normalized by their standard deviation.

Both the 3D-CMCC-FEM and the RS-based GPP climate sensitivity in the summer period has

been evaluated through the Standardized Precipitation Evaporation Index, SPEI (Vicente-Serrano

et al., 2010). The SPEI drought index helps to highlight periods of wetter or drier conditions.

This is a multi-scalar meteorological drought index based on a statistical transformation of the

climatic water balance, i.e. precipitation *minus* potential evapotranspiration. We computed the

SPEI using the simplistic Hargreaves equations for the potential evapotranspiration calculation

and considered different time scales, with the aim to cover the growing season period before the

month of August. Following a similar approach as in Mahnken et al. (2022), we regressed the

residuals of the 3D-CMCC-FEM and RS-based datasets anomalies against the SPEI values as the

predictor, and the slope of the linear regression computed. Values of the slope close to 0 indicate

that the 3D-CMCC-FEM shows a GPP-sensitivity to SPEI like the RS-based dataset.

The overall analyses where carried out considering the entire RS-based dataset available years,

thus 2005-2013 for CFIX, 2005-2015 for FLUXCOM and 2005-2019 for GOSIF (i.e. 32 years in

total).

3 Results

3.1 Spatial variability analysis

The 3D-CMCC-FEM simulated average annual GPP for the period 2005-2019 are shown in

Figure 2, with overall values ranging from ~600 up to ~2200 gC m^{-2} yr^{-1}. 3D-CMCC-FEM GPP

follows a west-east gradient with higher productivity over the western side of the region in

correspondence with the more humid and the more productive beech-dominated areas; the

forested areas over the plain zones are mainly dominated by Mediterranean pine species, which

show the lowest GPP values (Figure 2). At annual level (multiyear mean annual) better

correlations between 3D-CMCC-FEM and RS-based GPP data are with GOSIF and FLUXCOM

($r = 0.77$ and 0.67, respectively; Table 1) as also for SPAEF metric (SPAEF = 0.62 and 0.59,

respectively) while lower RMSE and RD are when 3D-CMCC-FEM GPP is compared with

GOSIF and CFIX (RMSE = 235.8 and 221.9 gC m^{-2} yr^{-1}, and RD = –4.3 and 0.01%,

respectively). At seasonal level, 3D-CMCC-FEM GPP better correlates with GOSIF and CIFX

during summer (RMSE = 181.3 and 123.8 g C m^{-2} yr^{-1}, RD = –13.7 and 10.7% and $r = 0.79$ and

0.8, respectively). Similarly, also in spring 3D-CMCC-FEM better correlates, although with

lower values, with GOSIF and CIFX (RMSE = 104.4 and 103.6 gC m^{-2} yr^{-1}, RD = 9 and –7.9%

and r = 0.4 and 0.28, respectively). In winter 3D-CMCC-FEM GPP better correlates with CIFX

and FLUXCOM (r = 0.55 and 0.5, and SPAEF = –1.26 and 0.53, respectively) but with slightly

lower RMSE values when compared with GOSIF and FLUXCOM (RMSE = 54.3 and 39.97 gC

m^{-2} yr^{-1}). During autumn 3D-CMCC-FEM is in agreement with GOSIF and CIFX (RMSE =

64.47 and 52.41 gC m^{-2} yr^{-1}, RD = 2.4 and 15% and r = 0.74 and 0.67, respectively).

3D-CMCC-FEM GPP *vs.* GOSIF GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	SPAEF	*r*
ANNUAL	235.81	–4.30	**0.62**	**0.77**
DJF	54.34	–31.3	–1.50	0.39
MAM	**104.43**	9.0	0.22	**0.40**
JJA	181.35	–13.7	**0.61**	**0.79**
SON	64.47	**2.4**	0.46	**0.74**
3D-CMCC-FEM GPP *vs.* CFIX GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	SPAEF	*r*
ANNUAL	**221.91**	**0.01**	–0.07	0.59
DJF	65.26	–47.8	**–1.26**	0.55
MAM	103.61	**–7.9**	–0.31	0.28
JJA	**123.78**	**10.7**	0.38	0.80
SON	**52.41**	15.1	**0.48**	0.67
3D-CMCC-FEM GPP *vs.* FLUXCOM GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	SPAEF	*r*
ANNUAL	477.42	40.0	0.59	0.67
DJF	**39.97**	**–13.4**	–0.53	**0.50**
MAM	198.47	45.9	**–0.45**	–0.09
JJA	203.44	37.2	0.57	0.72
SON	118.25	57.7	0.40	0.68

423 **Table 1 Spatial comparison of the 3D-CMCC-FEM GPP vs. RS–based GPP. RMSE= root mean square error**
424 **(gC m^{-2} yr^{-1}), RD = relative difference (%), SPAEF = SPatial EFficiency, r = Pearson's correlation. DJF,**
425 **winter months; MAM, spring months, JJA, summer months; SON, autumn months. In bold values with the**
426 **best agreement between 3D-CMCC-FEM GPP and RS-based GPP.**

3.2 Temporal variability analysis

Generally, the 3D-CMCC-FEM GPP shows overall lower correlations of the temporal (which corresponds to the interannual variability, IAV) vs. spatial comparison for the mean annual GPP. Similarly, as in the spatial analysis, 3D-CMCC-FEM GPP shows better agreement with GOSIF (RMSE = 231.6 gC m^{-2} yr^{-1}, RD = −4.6% and r = 0.4, see Table 2 for overall statistics and Figure 3, 4, 5, 6 for the maps) and the lowest Fractional Variance between the RS-based dataset considered (FV = 0.83, see also Standard Deviation analysis Figure S1). At seasonal level, during summer, 3D-CMCC-FEM GPP correlates better with the FLUXCOM (r = 0.62 and FV = 1.64) although with the highest RMSE (182.53 gC m^{-2} yr^{-1}) among the RS-based dataset considered. Conversely, in spring 3D-CMCC-FEM GPP correlates better with CFIX (r = 0.31) and with the lowest RMSE (86.81 gC m^{-2} yr^{-1}) and RD (−7.9%). In winter modeled GPP values from 3D-CMCC-FEM better correlates with CFIX (r = 0.69) but with lower RMSE, RD and FV values when compared with FLUXCOM (RMSE = 36.24 gC m^{-2} yr^{-1}, RD = −24.1 and FV = 1.77). Also during autumn season, the 3D-CMCC-FEM GPP better correlates with CFIX (r = 0.3) and with the lowest RMSE (57.97 gC m^{-2} yr^{-1}) although the lowest Relative Difference is when compared with GOSIF (RD = 1.7%).

At the Mean Seasonal Cycle level (thus comparing month per month average GPP values) 3D-CMCC-FEM GPP well correlates with all RS-based dataset with r varying from 0.95 (when compared with CFIX) to 0.91 (when compared with GOSIF)(see Figure 7, Table 2 and Figure S2)

3D-CMCC-FEM GPP *vs.* GOSIF GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	FV	*r*
ANNUAL	231.60	–4.6	**0.83**	**0.40**
DJF	54.64	–40.6	1.70	0.51
MAM	105.61	10.4	**0.77**	0.15
JJA	146.72	–12.7	0.97	0.53
SON	77.33	**1.7**	**0.38**	0.17
MSC	/	/	/	0.91
3D-CMCC-FEM GPP *vs.* CFIX GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	FV	*r*
ANNUAL	**179.55**	**–1.3**	1.17	0.12
DJF	66.67	–58.7	**1.09**	**0.69**
MAM	**86.81**	**–7.9**	1.36	**0.31**
JJA	**86.12**	**7.5**	0.68	**0.65**
SON	**57.97**	15.2	0.65	**0.30**
MSC	/	/	/	**0.95**
3D-CMCC-FEM GPP *vs.* FLUXCOM GPP				
	RMSE (gC m^{-2} yr^{-1})	RD (%)	FV	*r*
ANNUAL	486.65	40.6	1.69	0.39
DJF	**36.24**	**–24.1**	1.77	0.55
MAM	209.14	49.4	1.66	0.26
JJA	182.53	37.9	1.64	0.62
SON	120.33	54.7	1.41	0.17
MSC	/	/	/	0.92

Table 2 Temporal comparison of the 3D-CMCC-FEM GPP vs. RS–based GPP. RMSE= root mean square error (gC m^{-2} yr^{-1}); RD = relative difference (%); FV = fractional variance; *r* = Pearson's correlation. Metrics are first computed at grid cell level and reported as the median value. DJF, winter months; MAM, spring months, JJA, summer months; SON, autumn months; MSC, mean seasonal cycle. In bold values with the best agreement between 3D-CMCC-FEM GPP and RS-based GPP.

3.3 SPEI

We first computed the SPEI at a time-scale of 5 months (SPEI5) representing the spring and summer period where the correlation between the RS-based summer GPP and SPEI were stronger. We estimate whether the 3D-CMCC-FEM data and RS-based data differ in GPP

sensitivity toward interannual variation in SPEI by computing the slope of the regression

between 3D-CMCC-FEM vs. RS-based differences of GPP and the SPEI. Generally, results

show how, when considering CFIX and FLUXCOM, the slope is not significantly different from

0 for almost the entire simulation domain (Figure 8). Such a behavior indicates that 3D-CMCC-

FEM and RS-based GPP summer anomalies respond to SPEI5 similarly. When comparing with

the GOSIF the slope is still not significant over large areas. However, in clustered areas,

corresponding to about a 20% of the simulation domain, the slope is positive and significant

showing that the 3D-CMCC-FEM response to aridity is stronger than the GOSIF data (Figure 8).

3.4 Species-level comparison

Data analysis at species level reveals as 3D-CMCC-FEM model GPP tends to correlate better for

some species for some RS-based datasets than other (Table 3 and Figure 9). 3D-CMCC-FEM

GPP correlates well for *Q. cerris* and *Q. ilex* with GOSIF ($r = 0.73$ and 0.82) but with low

RMSE and RD for CFIX (RMSE = 141.09 and 141.99 gC m^{-2} yr^{-1}, and RD = –2.69 and 3.39%,

respectively). In all cases 3D-CMCC-FEM GPP for *F. sylvatica* are slightly far from results from

all RS-based datasets with better correlations with CFIX ($r = 0.43$) and lower RMSE and RD

values for GOSIF (193.35 gC m^{-2} yr^{-1} and 3.43%). Satisfactorily correlations are shown for *C.*

sativa with the GPP values of FLUXCOM ($r = 0.61$) but with lower RMSE and RD with GOSIF

(200.45 gC m^{-2} yr^{-1} and –1.71%). For Pinus species (both as *P. halpensis* and *P. nigra*), 3D-

CMCC-FEM GPP shows to better correlates and with lower RMSE and RD values with GOSIF

($r = 0.84$, RMSE = 170.84 gC m^{-2} yr^{-1} and RD = –7.91% for *P. halepensis*; and $r = 0.65$, RMSE

= 223.92 gC m^{-2} yr^{-1} and RD = 7.85% for *P. nigra*) than with CFIX and FLUXCOM (Table 3

and Figure 9).

3D-CMCC-FEM GPP *vs.* GOSIF GPP			
	RMSE (gC m^{-2} yr^{-1})	RD (%)	*r*
Quercus cerris	245.66	–6.33	**0.73**
Fagus sylvatica	**193.35**	**3.43**	0.28
Quercus ilex	267.65	3.96	**0.83**
Pinus halepensis	170.84	–7.91	**0.85**
Pinus nigra	**223.92**	**7.86**	**0.66**
Castanea sativa	**200.45**	**–1.72**	0.25
3D-CMCC-FEM GPP *vs.* CFIX GPP			
	RMSE (gC m^{-2} yr^{-1})	RD (%)	*r*
Quercus cerris	**141.1**	**–2.7**	0.60
Fagus sylvatica	477.4	31.1	**0.43**
Quercus ilex	**141.9**	**3.4**	0.70
Pinus halepensis	355.1 (28%)	–26.8	0.65
Pinus nigra	314.5 (21%)	16.9	0.25
Castanea sativa	203.3	7.0	0.36
3D-CMCC-FEM GPP *vs.* FLUXCOM GPP			
	RMSE (gC m^{-2} yr^{-1})	RD (%)	*r*
Quercus cerris	440.4	39.1	0.66
Fagus sylvatica	777.8	63.1	0.10
Quercus ilex	462.1	40.0	0.75
Pinus halepensis	**140.7**	**7.2**	0.68
Pinus nigra	656.1	57.4	0.57
Castanea sativa	584.1	49.0	**0.61**

Table 3 Comparison of the 3D-CMCC-FEM GPP vs. RS-based GPP at single species level. RMSE= root mean square error (gC m^{-2} yr^{-1}), RD = relative differences (%), r = Pearson's correlation. Metrics are first computed at grid cell level and reported as the median value. In bold values with the best agreement between 3D-CMCC-FEM GPP and RS-based GPP.

4 Discussion

In this study, we applied the 3D-CMCC-FEM spatial explicitly over a Mediterranean region characterized by elevated bio-geographical and climatological heterogeneity. Initializing the forest structure, combining ground data from the NFI and the RS-based wall-to-wall basal area map, allows having a more realistic initial state of the carbon pools and forest structure,

reducing, thus, uncertainties related to spin-up procedures which often translate into significant

differences in the carbon fluxes (Carvalhais et al., 2008, 2010; Massoud et al., 2019; Lindeskog

et al., 2021). The calibration of the allometric equations, to which the model has shown to be

sensitive (Collalti et al., 2019), sets an additional constraint on the modeled initial growing

biomass, which in turn influences the simulated GPP *via* the amount of sapwood. By means of

the use of the basal area map to build the forest characteristics at the beginning of simulation, the

model is able to retain part of the spatial information of the initial aboveground biomass and

average forests structure, contributing, for instance, to the very satisfactorily spatial correlations

of the modeled annual GPP with 2 out of 3 RS-based datasets. While aggregating basal area data

at the 1 km resolution might reduce random uncertainties in basal area, additional uncertainty

might stem from the tree density data, a data which is less constrained. Yet, performed tests (not

shown here) indicate no significant sensitivity of the simulated GPP to stand density.

Overall, 3D-CMCC-FEM performances in simulating GPP when compared to other large scale

and independent RS-based data, are shown to be generally satisfactory at both spatial as also

temporal scales. In particular in the summer period across the seasons 3D-CMCC-FEM GPP

show satisfactorily correlations and low relative differences and RMSE values when most of the

vegetation activity takes place and it is thus the most robust signal across model and RS-based

datasets. Yet, the comparative analyzed pinpointed some important challenges in applying the

3D-CMCC-FEM in Mediterranean areas and highlighted sources of uncertainties explaining the

residual 3D-CMCC-FEM and RS-based data differences across RS-based datasets, which are

grouped as follows:

4.1 3D-CMCC-FEM GPP vs. RS-based GPP

The 3D-CMCC-FEM GPP is close to the GOSIF and CFIX GPP estimates, while a general systematic positive overestimation emerge when comparing the 3D-CMCC-FEM GPP to FLUXCOM GPP (although for not all species, Figure 9). The 3D-CMCC-FEM and RS-based data differences resulting from the analyses are here discussed in relation to the different nature of the RS-based datasets used and their underlying algorithms and adopted approaches. FLUXCOM is an up-scaled product of the local EC-tower GPP estimates which is often used as benchmark RS-based dataset in model evaluation analyses (e.g. Byrne et al., 2018). Part of its worldwide application relies on the capability to sample in the entire climate-vegetation space. However, some particular areas at the rear edge of the Mediterranean forests might be not covered by this product (Jung et al., 2020). In addition, FLUXCOM operates also at coarse spatial resolution (Zheng et al., 2020; Zhang et al., 2022). Compared to GOSIF and CFIX, the spatial, seasonal and interannual variability in FLUXCOM relies on the RS-based dataset alone, without including any climatic drivers. Unfortunately, climate has been shown to play a significant role in the local, regional and global GPP. The lack of climatic drivers in FLUXCOM, might, thus, partly explain the systematic differences observed when compared to the 3D-CMCC-FEM GPP, which, at the opposite, showed to be sensitive to climate (Mahnken et al., 2022). However, the noticeable differences found, although some good correlations for some species, between the 3D-CMCC-FEM GPP and FLUXCOM, is common to other process-based and dynamic vegetation models comparative studies (Li & Xiao, 2019b; Jung et al., 2020; Zhang & Ye, 2021), and recent studies found that the GPP may be underestimated in the FLUXCOM-GPP in temperate areas (e.g. Bacour et al., 2019; Norton et al., 2019; Wild et al., 2022).

The GOSIF and the CFIX datasets, which 3D-CMCC-FEM GPP seems to reply better than

FLUXCOM, have the advantage of having a finer spatial resolution (~5 and 1 km versus ~ 8 km

in FLUXCOM) and, thus, contain more information, with the GOSIF having a globally higher

continuous coverage *via* the original SIF data, which is the proxy of the photosynthetic activity at

canopy scales (Sun et al., 2017; Li & Xiao, 2019a). CFIX is instead driven by a Light Use

Efficiency model driven by NDVI, which is a vegetation index more suitable to investigate plant

greenness rather than purely photosynthetic activities, as highlighted by Camps-Valls et al.

(2021) and Walther et al. (2019). This inherent characteristic for NDVI might explain the lack of

spatial correlation but also lower spatial correlations when compared to FLUXCOM and GOSIF.

In addition, the Light Use Efficiency models are known to not saturate at increasing solar

radiation, but that would lead CFIX having higher GPP values than 3D-CMCC-FEM GPP,

which uses the biochemical model of Farquhar et al. (1980). However, while in some species

CFIX has higher GPP values than 3D-CMCC-FEM GPP (e.g. *F. sylvatica* and *Q. Ilex*) in other

species values are lower (*P. halepensis* and *Q. cerris*)(Figure 9). Differences between 3D-

CMCC-FEM and CFIX may thus more probably rely on different models' parameterization and

not just on the different approach used to simulate photosynthesis. Indeed, as outlined in other

studies carried at global scales there is a higher uncertainties in interannual variability even

across different RS-based datasets, (Butterfield et al., 2020; Zhang & Ye, 2021) and even in its

correct (of GPP) calculation across different models (Dunkl et al., 2023).

The differences in standard deviation between the 3D-CMCC-FEM and the RS-based GPP (see

Figure S1), despite high in absolute terms, are still comparable to the range shown for instance in

Zhang & Ye (2021) and Zheng et al. (2020). Similarly, the partial lack of agreement in the winter

(but this is worth also for the spring and autumn) months, that we found in this study, it has been

observed also in other RS-based data and process-based models comparative analyses (Zhang

and Ye 2020). However, this it is not surprising, at least for deciduous species, given that 3D-

CMCC-FEM simulates dominant vegetation only and no photosynthesis when there are no

leaves on at the opposite to RS-based data which may account for underneath vegetation and this

might explain the better correlations with evergreen species than for the deciduous for some RD-

based datasets. Spring and autumn also depend on the spatial and temporal variability of bud

breaks and leaf falls which control photosynthetic activity during and across the years. However,

mismatches and asynchronies for the beginning and the end of the growing season largely vary

between species and RS-based datasets considered as also found in the literature for other models

(Peano et al., 2019, 2021). In addition, the RS-based datasets capture the fluxes embedding the

entire sub-grid variability, including the contribution of the vegetation not simulated by the

model (e.g. crops, *maquis* and understory which in any case comprise about one fifth of the

entire vegetation area only), in particular in areas where the forest cover is indeed low because of

low stand density, such as in the plain areas of the region under study or in degraded areas.

Spatial scale mismatch between data and models may explain, thus, likely part of the low

performances in some areas of the region. Zhao and Zu (2022) showed how GOSIF has

apparently a similar interannual variability and trends when compared to the TRENDY

simulations (an ensemble of simulations from Dynamic Global Vegetation Models; Sitch et al.,

2015), as similarly as we found here using a process-based, stand-level, forest model.

On the other hand, some have raised doubts and concerns on the robustness of the relationship

between GPP and remotely sensed sun-induced fluorescence (SIF)(Wohlfahrt et al., 2018;

Bacour et al., 2019; Chen et al., 2021) showing that the SIF might overestimate the GPP values

at least in temperate broadleaved forests (Qiu et al., 2020). However, to our knowledge, there are

no specific indication that this retains also for Mediterranean forests. Yet the SIF data from RS

remain one of the best proxies for photosynthesis over large areas to date (Sun et al., 2017; Li &

Xiao, 2019a), making the GOSIF estimates potentially the more robust estimates compared to

both CFIX and FLUXCOM.

Differences in the original spatial resolution of RS-based data or land cover used to drive the

GOSIF and CFIX datasets, might additionally contribute to explain the 3D-CMCC-FEM GPP

and RS-based datasets – but also between RS-based datasets – differences. Interestingly, when

compared RS-based dataset vs. RS-based dataset GPP (e.g. GOSIF vs. FLUXCOM), and not just

3D-CMCC-FEM GPP vs. RS-based dataset, low correlations and discrepancies (both in the

relative as in the absolute sense), although minor than against 3D-CMCC-FEM, across the

results (see Table S3 and Table S4), both at the spatial as the temporal scale (including at the

species level, see for example the large RMSE for all deciduous species in Table S4), have

found. That was, however, expected given that RS-based methods are diagnostic and of similar

nature tools rather than inherently prognostic (and with greater uncertainties) tools as potentially

all models are. A recent large review on RS-based products intercomparison confirmed the large

variability and different sensitivity to climatic factors between different approaches and criteria

when compared at site-level (Sun et al., 2019).

4.2 3D-CMCC-FEM-related uncertainties

The 3D-CMCC-FEM model has been extensively evaluated at the site level all over Europe and

evaluated against measurements of carbon fluxes from the FLUXNET database (Collalti et al.,

2016, 2018; Marconi et al., 2017). The evaluation has been carried out in the past over a broad

spectrum of climate, forest management, and species at stand level (e.g. Dalmonech et al., 2022;

Testolin et al., 2023). In some recent model inter-comparison studies, the 3D-CMCC-FEM was

shown to be able to simulate, among other things, C fluxes, e.g. GPP, and key structural

variables, e.g. diameter, basal area, etc, with very good performance when compared to

measured data and to other state-of-the-art forest models (Engel et al., 2020; Mahnken et al.,

2022). However, when applying the 3D-CMCC-FEM at a different spatial scales model

parameters, structural, and input-related uncertainties might amplify or even buffer the error in

simulating GPP as observed for other models (e.g. Dalmonech et al., 2015; Zheng et al., 2022;

Dunkl et al., 2023). In any case, the overall, 3D-CMCC-FEM GPP (~600 - ~2200 gC m^{-2} yr^{-1}) is

well in the bounds of the ~600 - ~2500 gC m^{-2} yr^{-1} GPP values described in Collalti and Prentice

(2019) for temperate deciduous and coniferous forests in Europe.

The seasonal cycle of GPP describes the seasonal pattern of carbon gross assimilation by plants

and the beginning and end of the growing season for the deciduous forests. For two out of the

three datasets used here the 3D-CMCC-FEM may simulate a slightly earlier beginning of the

growing season (e.g. for *Q. cerris*)(Figure S2). This might be partially attributable to the

budburst parameterization, which is based on the Thermic sum and the growing degree days

metric which is a trigger for the leaf flushing. Such a parameter is kept constant (as in many

other models, see Peano et al., 2019; Peaucelle et al., 2019; Collalti et al., 2019) as a species-

specific parameter, irrespective of any local climatic adaptation. The observed discrepancy in the

beginning of the growing season, compared to the delay projected by the 3D-CMCC-FEM

model, may be partly attributed also to the influence of undergrowth vegetation (such as grass or

shrubs) on RS-based data. These understory plants frequently begin photosynthesis earlier, a

phenomenon not accounted for by the 3D-CMCC-FEM model. The seasonality of the GPP in the

evergreen species is instead apparently more shaped by the direct environmental effect on the

photosynthetic process at a seasonal time scale (and less affected by underneath vegetation),

suggesting that the 3D-CMCC-FEM GPP and RS-based datasets differences might be also

attributable to some bias in the meteorological data used or for the other physiological

parameters adopted or in the below-canopy vegetation. Indeed, model ecophysiological

parameter values are derived from the literature, yet not calibrated for the specific sites or

regions, in order to allow the 3D-CMCC-FEM general applicability as reported in Mahnken et al.

(2022). In addition, other potential source of uncertainty, although minor, for the apparent

mismatches in the beginning and in the end of the growing season between 3D-CMCC-FEM and

RS-based datasets may stem from the different temporal resolutions, indeed, e.g. FLUXCOM is

an 8-day product while 3D-CMCC-FEM is a daily one.

To capture the summer GPP sensitivity to hydrological variability, i.e. wetter or drier conditions,

we use a simple GPP-SPEI diagnostics. The 3D-CMCC-FEM was first shown to be able to

simulate comparably to the RS-based datasets the interannual variability (see Table 2) and then

of the summer GPP and aridity signal over a large extent (see Figure 8). In large areas of the

region, the 3D-CMCC-FEM and the RS-based datasets have similar responses in terms of GPP-

SPEI5. Yet, in some areas, a too-strong model response to SPEI5 also emerges, indicating that

the 3D-CMCC-FEM has a higher summer GPP-response to negative SPEI5 (i.e. drier conditions)

compared to the RS-based datasets. In particular, results indicate a higher model sensitivity of

GPP to SPEI for oaks-dominated forests clustered on higher elevations. Oak's species are known

to be an isohydric species, i.e. they weakly regulate stomatal openness under drought conditions

and are more resilient to drought (Ripullone et al., 2020; Castellaneta et al., 2022), and it is

possible that the high sensitivity of GPP to SPEI5 in the 3D-CMCC-FEM is a result either of a

too strong control of VPD on stomatal conductance (Grossiord et al., 2020) or of a too strong soil

649 moisture control on the photosynthesis (Crow et al., 2020; Fang et al., 2021). Summer period is

650 the season where most of the annual vegetation activity takes place and reaches its maximum,

651 and which variability is the most prominent feature in Mediterranean areas. As a matter of fact,

652 in a previous study (see Collalti et al., 2016) was shown how, at site level in beech and

653 coniferous forests, the 3D-CMCC-FEM was able to sufficiently simulate the sign of the year-to-

654 year variability of the annual GPP anomalies, and in particular in years affected by important

655 summer drought (i.e. 2003). Yet interannual variability is the most uncertain signal event when

656 comparing different data and often considered as an 'acid test' in vegetation modeling (Keenan

657 et al., 2012; Collalti et al., 2016; Dunkl et al., 2023).

658 However, stand-level studies showed how the 3D-CMCC-FEM model realistically simulates the

659 response of GPP to atmospheric VPD and temperature (Mahnken et al., 2022). In the model soil

660 hydrology is simulated *via* a single-layer bucket model and this might be another factor

661 contributing to the stronger than observed modeled response, as it has been shown as overall that

662 the majority of models with a soil bucket hydrology tend to limit GPP more than models with

663 other soil conceptual e.g. multilayered schemes (Hanson et al., 2004). This higher sensitivity and

664 the apparent stronger drop of modeled summer GPP might partially explain the higher standard

665 deviation of the modeled signal compared to RS-based datasets (see Figure S1). One potential

666 additional explanation to the modeled stronger GPP sensitivity to SPEI, would be the detected,

667 although the narrow, earlier onset of the growing season in the model compared to the GOSIF

668 dataset. In fact, leaf development not only drives how earlier or later the uptake of carbon from

669 the atmosphere starts in the season, but also how earlier/later other processes, such as the leaf

670 transpiration occurs in spring, potentially affecting the summer GPP response (Peano et al.,

671 2019; Bastos et al., 2021; Chen et al., 2023).

4.3 Meteorological and soil data uncertainties

Low accuracy in the reconstructed meteorological forcing (see Bandhauer et al., 2022) might play an additional important role in studies that are conducted at high spatial resolution. The impact of the meteorological data used in data-driven and PBFMs was already shown to be important to accurately simulate GPP (e.g. Wu et al., 2017; F. Zhang et al., 2022; Y. Zhang & Ye, 2022). The 3D-CMCC-FEM was shown to be able to simulate GPP closer to observations at the site level in northern, central and southern Europe, where the downscaling and bias correction of the climate data was facilitated by the local geography (Collalti et al., 2018; Testolin et al., 2023).

Potential positive biases in the original E-OBS temperature dataset, as its accuracy relies also on the density of the termopluviometric stations and altitudinal cover, might translate into higher temperatures at higher elevations where, indeed, there are few stations as in southern Italy. A positive bias in temperature over areas where the temperature should be a limiting factor for vegetation, may partially explains the high modeled GPP values in beech forests leading to an anticipated onset or at thermic stress during the summer. Beech in the Italian Apennine Mountain regions represents the upper limit of forest vegetation and other studies (J. Fang & Lechowicz, 2006; Marchand et al., 2023) showed that the most limiting factor for the presence and growth of the beech in Europe is, indeed, temperature. The simulated values of GPP in beech-dominated forests are, on average higher than the three reference dataset, with simulated values in some areas larger than 2000 $gCm^{-2}yr^{-1}$. However these values are overestimated as observed average annual values of GPP in a beech forest in central Italy, a monitoring site with fluxes measured by means of the eddy covariance technique, are in the order of 1600 $gC\ m^{-2}yr^{-1}$ with range between ~550 and 2500 $gC\ m^{-2}\ yr^{-1}$ from Reyer et al. (2020). When applied in the same beech forest, the

3D-CMCC-FEM was shown to be able to fall within the observed range of GPP (Mahnken et al.,

2022), indicating how the source of bias might be indeed attributable to uncertainties stemming

from the up-scaling and affecting mostly the medium to higher elevation areas.

Simulated values for the deciduous oaks are indeed more comparable to EC-estimates in a site in

central Italy (Roccarespampani, www.fluxnet.com, Tedeschi et al., 2006), with annual values of

the order of 1600 gC m^{-2} yr^{-1}for model and observations. However, a west-east gradient in the

model-data difference is apparent at the regional level when comparing the 3D-CMCC-FEM to

CFIX and GOSIF. In the former case, differences correlate with elevation and temperature, very

likely a results of the lack of accuracy in the E-OBS dataset. Spatial differences in the 3D-

CMCC-FEM GPP emerge in this study, which cannot be explained only by uncertainties in the

modeling setup alone. The amount of water available for plants is determined in the 3D-CMCC-

FEM by the balance between the precipitation (the inflow) and the Evapotranspiration (the

outflow) as well as the soil characteristics, such as soil depth and texture. The water availability

and the soil texture translate into the matric potential, which directly affects the stomatal

regulation, leaf transpiration and the photosynthesis. When the model is applied spatially in a

Mediterranean area with orographic and geographical complexity, the capability of the climate to

retain the spatial and temporal variability of precipitation might have important effects in

modulating the modeled spatial and temporal GPP variability (see Y. Zhang & Ye, 2022), and

this will be the object of further investigations.

The last source of uncertainties in the 3D-CMCC-FEM GPP and RS-based GPP might stem from

the soil characteristics and, in particular, the soil depth. The dataset used in the analyses as a

proxy of depth available to root does not show any sensible spatial variability, e.g. with

elevation, which is not realistic considering the topography and plasticity of plants roots (Fan et

al., 2017; Stocker et al., 2023). The 3D-CMCC-FEM at this scale of application would benefit

from coupling with more advanced hydrological scheme or a soil depth parameterization, to

better couple the vegetation with the effective soil water availability (Kollet & Maxwell, 2008;

Niu et al., 2011) or from soil moisture assimilation from remote sensing products (Kumar et al.,

2014; Crow et al., 2020).

4.4 Limitations and further considerations

In this study we did not consider the land use change and the model does not simulate any spatial

interaction from neighboring grid cells, not allowing for example simulating forest expansion.

We additionally aggregated the forest species according to macro classes, hence neglecting intra-

specific differences, if any. However, some compromises between input data requirements and

time for model runs was needed to keep simulations and calculations feasible. In particular, at

the spatial resolution of the study, we were more interested in capturing trends and spatial

variability in GPP than the finest sub-grid variability within the forest ecosystem.

No Nitrogen cycle is included, however, the precipitation is commonly identified as the main

limiting factor for photosynthesis in Mediterranean areas (Keenan et al., 2009; Flexas et al.,

2014; Fyllas et al., 2017).

5 Conclusions and outlook

PBFMs offer a complementary tool to ground forest inventory networks and satellite-based

records in monitoring carbon sequestration forest growth and the climate change impacts on

forests, on many other variables which are otherwise difficult to measure extensively or monitor

continuously. However, any model's reliability needs to be verified and tested even in high

heterogeneity contexts and not only to site-level ones.
With this aim, we used RS-based data to initialize, apply, and test for the first time the
biogeochemical, biophysical, process–based model 3D-CMCC-FEM on a regular grid at high
spatial resolution under typical Mediterranean climate. We compare the obtained gross primary
production over a large area (~80% of the Basilicata region) against a suite of different in nature
independent RS-based data. In spite of the simplified initial forest setup and the underlined
uncertainties, the 3D-CMCC-FEM was shown capable of capturing both the spatial and the
temporal variability of the RS-based data, even at species-level. Further tests are needed, yet the
very promising results open the possibility of using the PBFMs to investigate the spatio-temporal
dynamics of forest growth over larger spatial scales and under drought conditions and future
climate scenarios, shaped by the spatial climatic and ecological heterogeneity such as in the
Mediterranean areas.

Acknowledgments

The work was carried out in the framework of the project 'Advanced EO Technologies for
studying Climate Change impacts on the environment – OT4CLIMA' (D.D. 2261 - 6.9.2018,
PON R&I 2014–2020 and FSC). This work has been partially supported by MIUR Project (PRIN
2020) "Unraveling interactions between WATER and carbon cycles during drought and their
impact on water resources and forest and grassland ecosySTEMs in the Mediterranean climate
(WATERSTEM)"(Project number: 20202WF53Z), "WAFER" at CNR (Consiglio Nazionale
delle Ricerche) and D.D., E.V., A.C. were supported by PRIN 2020 (cod 2020E52THS) -
Research Projects of National Relevance funded by the Italian Ministry of University and
Research entitled: "Multi-scale observations to predict Forest response to pollution and climate
change" (MULTIFOR, project number 2020E52THS). D.D., E.V., A.C. acknowledge also
funding by the project OptForEU H2020 research and innovation programme under grant
agreement No. 101060554. We also acknowledge the project funded under the National
Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender

No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian
Ministry of University and Research funded by the European Union – NextGenerationEU under
award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022
adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project
title "National Biodiversity Future Centre - NBFC". We acknowledge the E-OBS dataset from
the EU-FP6 project UERRA (https://www.uerra.eu) and the Copernicus Climate Change Service,
and the data providers in the ECA&D project (https://www.ecad.eu)". The FLUXCOM products
were obtained from the Data Portal at https://www.fluxcom.org/. D.D. thanks C. Trotta for
valuable discussions about tree allometric relationships and thanks M. Willeit and E. Grieco for
providing comments on a previous draft of the manuscript.

Conflict of Interest

The authors have no conflicts of interest to declare.

Author contribution

Daniela Dalmonech: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resources, Software, Validation, Visualization, Writing - original draft. **Elia Vangi, Gherardo Chirici,** and **Francesca Giannetti:** Resources, Writing - review & editing. **Jingfend Xiao**: Resources, Writing - review & editing; **Marta Chiesi** and **Luca Fibbi:** Resources, Writing - review & editing**. Gina Marano:** Software, Writing - review & editing **Christian Massari:** Writing - review & editing. **Angelo Nole**: Resources, Writing - review & editing. **Alessio Collalti**: Conceptualization, Formal analysis, Investigation, Methodology, Resources, Software, Validation, Visualization, Writing - review & editing.

Data Availability Statement

The 3D-CMCC-FEM model code version 5.6 is publicly available under the GNU General Public Licence v3.0 (GPL) and can be found on the GitHub platform at: https://github.com/Forest-Modelling-Lab/3D-CMCC-FEM). All data and model executable, and scripts to perform analyses and figures presented in this work are provided open access in the

794 Zenodo server (https://doi.org/10.5281/zenodo.8060401). Correspondence and requests for
795 additional materials should be addressed to the corresponding author.

References

Allen, C. D., Breshears, D. D., & McDowell, N. G. (2015). On underestimation of global
vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene.
Ecosphere, *6*(8), 1–55. https://doi.org/10.1890/ES15-00203.1

Amato, V., Lanza, G. G., Foresta, D. La, & Simonetti, L. (2008). *Comunità montane . Soggetti*
propulsori dello sviluppo o enti inefficaci ?

Bacour, C., Maignan, F., MacBean, N., Porcar-Castell, A., Flexas, J., Frankenberg, C., Peylin, P.,
Chevallier, F., Vuichard, N., & Bastrikov, V. (2019). Improving Estimates of Gross Primary
Productivity by Assimilating Solar-Induced Fluorescence Satellite Retrievals in a Terrestrial
Biosphere Model Using a Process-Based SIF Model. *Journal of Geophysical Research:*
Biogeosciences, *124*(11), 3281–3306. https://doi.org/10.1029/2019JG005040

Bandhauer, M., Isotta, F., Lakatos, M., Lussana, C., Båserud, L., Izsák, B., Szentes, O., Tveito,
O. E., & Frei, C. (2022). Evaluation of daily precipitation analyses in E-OBS (v19.0e) and
ERA5 by comparison to regional high-resolution datasets in European regions.
International Journal of Climatology, *42*(2), 727–747. https://doi.org/10.1002/joc.7269

Barbati A., Corona P., Marchetti M. (2007). A forest typology for monitoring sustainable forest
management: The case of European Forest Types. *Plant Biosystems - An International*
Journal Dealing with all Aspects of Plant Biology, 141:1, 93-
103, https://doi.org/10.1080/11263500601153842

Bastos, A., Orth, R., Reichstein, M., Ciais, P., Viovy, N., Zaehle, S., Anthoni, P., Arneth, A.,
Gentine, P., Joetzjer, E., Lienert, S., Loughran, T., McGuire, P. C., Sungmin, O., Pongratz,
J., & Sitch, S. (2021). Vulnerability of European ecosystems to two compound dry and hot
summers in 2018 and 2019. *Earth System Dynamics*, *12*(4), 1015–1035.
https://doi.org/10.5194/esd-12-1015-2021

Byrne, B., Wunch D., Jones D.B.A. et al. (2018). Evaluating GPP and respirstion estimates over
northern midlatitude ecosystems using-solar-induced fluorescence and atmospheric CO2
measurements. J*ournal of Geophysical Research: Biogeosciences.*
Https://doi.org/10.1029/2018JG004472

Bugmann, H., & Seidl, R. (2022). The evolution, complexity and diversity of models of long-
term forest dynamics. *Journal of Ecology*, *110*(10), 2288–2307.
https://doi.org/10.1111/1365-2745.13989

Butterfield, Z., Buermann, W., & Keppel-Aleks, G. (2020). Satellite observations reveal seasonal
redistribution of northern ecosystem productivity in response to interannual climate
variability. *Remote Sensing of Environment*, *242*(June 2019).

https://doi.org/10.1016/j.rse.2020.111755

Carvalhais, N., Reichstein, M., Ciais, P., Collatz, G. J., Mahecha, M. D., Montagnani, L., Papale,
D., Rambal, S., & Seixas, J. (2010). Identification of vegetation and soil carbon pools out of
equilibrium in a process model via eddy covariance and biometric constraints. *Global*
Change Biology, *16*(10), 2813–2829. https://doi.org/10.1111/j.1365-2486.2010.02173.x

Carvalhais, N., Reichstein, M., Seixas, J., Colltaz, G. J., Pereira, J. S., Berbigier, P., Carrara, A.,
Granier, A., Montagnani, L., Papale, D., Rambal, S., Sanz, M. J., & Valentini, R. (2008).
Implications of the carbon cycle steady state assumption for biogeochemical modeling
performance and inverse parameter retrieval. *Global Biogeochemical Cycles*, *22*(2), 1–16.
https://doi.org/10.1029/2007GB003033

Castellaneta, M., Rita, A., Camarero, J. J., Colangelo, M., & Ripullone, F. (2022). Declines in
canopy greenness and tree growth are caused by combined climate extremes during
drought-induced dieback. *Science of the Total Environment*, *813*, 152666.
https://doi.org/10.1016/j.scitotenv.2021.152666

Chen, A., Mao, J., Ricciuto, D., Lu, D., Xiao, J., Li, X., Thornton, P. E., & Knapp, A. K. (2021).
Seasonal changes in GPP/SIF ratios and their climatic determinants across the Northern
Hemisphere. *Global Change Biology*, *27*(20), 5186–5197.
https://doi.org/10.1111/gcb.15775

Chen, Keski-Saari, S., Kontunen-Soppela, S., Zhu, X., Zhou, X., Hänninen, H., Pumpanen, J.,
Mola-Yudego, B., Wu, D., & Berninger, F. (2023). Immediate and carry-over effects of
late-spring frost and growing season drought on forest gross primary productivity capacity
in the Northern Hemisphere. *Global Change Biology*, *October 2022*, 1–17.
https://doi.org/10.1111/gcb.16751

Chiesi, M., Fibbi, L., Genesio, L., Gioli, B., Magno, R., Maselli, F., Moriondo, M., & Vaccari, F.
P. (2011). Integration of ground and satellite data to model Mediterranean forest processes.
International Journal of Applied Earth Observation and Geoinformation, *13*(3), 504–515.
https://doi.org/10.1016/j.jag.2010.10.006

Chiesi, M., Moriondo, M., Maselli, F., Gardin, L., Fibbi, L., Bindi, M., & Running, S. W.
(2010). Simulation of mediterranean forest carbon pools under expected environmental
scenarios. *Canadian Journal of Forest Research*, *40*(5), 850–860.
https://doi.org/10.1139/X10-037

Chirici, G., Giannetti, F., McRoberts, R. E., Travaglini, D., Pecchi, M., Maselli, F., Chiesi, M., &
Corona, P. (2020). Wall-to-wall spatial prediction of growing stock volume based on Italian
National Forest Inventory plots and remotely sensed data. *International Journal of Applied*
Earth Observation and Geoinformation, *84*(September 2019).
https://doi.org/10.1016/j.jag.2019.101959

Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., & Valentini, R. (2014).
A process-based model to simulate growth in forests with complex structure: Evaluation
and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central Italy.
Ecological Modelling, *272*, 362–378. https://doi.org/10.1016/j.ecolmodel.2013.09.016

Collalti, A., Thornton, P. E., Cescatti, A., Rita, A., Borghetti, M., Nolè, A., Trotta, C., Ciais, P.,
& Matteucci, G. (2019). The sensitivity of the forest carbon budget shifts across processes

along with stand development and climate change. *Ecological Applications*, *29*(2), 1–18. https://doi.org/10.1002/eap.1837

Collalti, A., Tjoelker, M. G., Hoch, G., Mäkelä, A., Guidolotti, G., Heskel, M., Petit, G., Ryan, M. G., Battipaglia, G., Matteucci, G., & Prentice, I. C. (2020). Plant respiration: Controlled by photosynthesis or biomass? *Global Change Biology*, *26*(3), 1739–1753. https://doi.org/10.1111/gcb.14857

Collalti, Marconi, S., Ibrom, A., Trotta, C., Anav, A., D'andrea, E., Matteucci, G., Montagnani, L., Gielen, B., Mammarella, I., Grünwald, T., Knohl, A., Berninger, F., Zhao, Y., Valentini, R., & Santini, M. (2016). Validation of 3D-CMCC Forest Ecosystem Model (v.5.1) against eddy covariance data for 10 European forest sites. *Geoscientific Model Development*, *9*(2), 479–504. https://doi.org/10.5194/gmd-9-479-2016

Collalti, Trotta, C., Keenan, T. F., Ibrom, A., Bond-Lamberty, B., Grote, R., Vicca, S., Reyer, C. P. O., Migliavacca, M., Veroustraete, F., Anav, A., Campioli, M., Scoccimarro, E., Šigut, L., Grieco, E., Cescatti, A., & Matteucci, G. (2018). Thinning Can Reduce Losses in Carbon Use Efficiency and Carbon Stocks in Managed Forests Under Warmer Climate. *Journal of Advances in Modeling Earth Systems*, *10*(10), 2427–2452. https://doi.org/10.1029/2018MS001275

Collalti A., Biondo C., Buttafuoco G., Maesano M., Caloiero T., Lucà F., Pellicone G., Ricca N., Salvati R., Veltri A., Scarascia-Mugnozza G., Matteucci G. (2017). Simulation, calibration and validation protocols for the model 3D-CMCC-CNR-FEM: a case study in the Bonis watershed (Calabria), Italy. *Forest@*, 14(14): 247-256, https://doi.org/10.3832/efor2368-014, 2017

Collalti A., Dalmonech D., Marano G., Vangi E., Puchi P., Grieco E., Orrico M. (2022) "3D-CMCC-FEM User's Guide". ISBN 978-88-8080-573-1, https://doi.org/10.32018/3D-CMCC-FEM-2022

Collalti A. & Prentice I.C. (2019). Is NPP proportional to GPP? Waring's hypothesis twenty years on. *Tree Physiology*, 39(8): 1473-1483, https://doi.org/10.1093/treephys/tpz034, 2019.

Cornes, R. C., van der Schrier, G., van den Besselaar, E. J. M., & Jones, P. D. (2018). An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets. *Journal of Geophysical Research: Atmospheres*, *123*(17), 9391–9409. https://doi.org/10.1029/2017JD028200

Corona, P. (2008). La carta forestale della Toscana. *Forest@ - Rivista Di Selvicoltura Ed Ecologia Forestale*, *3*(3), 41.

Crow, W. T., Gomez, C. A., Sabater, J. M., Holmes, T., Hain, C. R., Lei, F., Dong, J., Alfieri, J. G., & Anderson, M. C. (2020). Soil moisture–evapotranspiration overcoupling and l-band brightness temperature assimilation: Sources and forecast implications. *Journal of Hydrometeorology*, *21*(10), 2359–2374. https://doi.org/10.1175/JHM-D-20-0088.1

D'Andrea, E., Rezaie, N., Prislan, P., Gričar, J., Collalti, A., Muhr, J., & Matteucci, G. (2020). Frost and drought: Effects of extreme weather events on stem carbon dynamics in a Mediterranean beech forest. *Plant Cell and Environment*, *43*(10), 2365–2379. https://doi.org/10.1111/pce.13858

D'Andrea, E., Scartazza, A., Battistelli, A., Collalti, A., Proietti, S., Rezaie, N., Matteucci, G., &

Moscatello, S. (2021). Unravelling resilience mechanisms in forests: role of non-structural carbohydrates in responding to extreme weather events. *Tree Physiology*, *41*(10), 1808–1818. https://doi.org/10.1093/treephys/tpab044

Dalmonech, D., Marano, G., Amthor, J. S., Cescatti, A., Lindner, M., Trotta, C., & Collalti, A. (2022). Feasibility of enhancing carbon sequestration and stock capacity in temperate and boreal European forests via changes to management regimes. *Agricultural and Forest Meteorology*, *327*(April), 109203. https://doi.org/10.1016/j.agrformet.2022.109203

De Marco A., Sicard, P., Feng, Z., Agathokleous, E., Alonso, R., Araminiene, V., Augustatis, A., Badea, O., Beasley, J. C., Branquinho, C., Bruckman, V. J., Collalti, A., David-Schwartz, R., Domingos, M., Du, E., Garcia Gomez, H., Hashimoto, S., Hoshika, Y., Jakovljevic, T. … Paoletti, E. (2022). Strategic roadmap to assess forest vulnerability under air pollution and climate change. Global Change Biology, 28, 5062–5085. https://doi.org/10.1111/gcb.16278

De Pury, D. G. G., & Farquhar, G. D. (1997). Simple scaling of photosynthesis from leaves to canopies without the errors of big-leaf models. *Plant, Cell and Environment*. https://doi.org/10.1111/j.1365-3040.1997.00094.x

Dosio, A., & Paruolo, P. (2011). Bias correction of the ENSEMBLES high-resolution climate change projections for use by impact models: Evaluation on the present climate. *Journal of Geophysical Research Atmospheres*, *116*(16), 1–22. https://doi.org/10.1029/2011JD015934

Dubrovský, M., Hayes, M., Duce, P., Trnka, M., Svoboda, M., & Zara, P. (2014). Multi-GCM projections of future drought and climate variability indicators for the Mediterranean region. *Regional Environmental Change*, *14*(5), 1907–1919. https://doi.org/10.1007/s10113-013-0562-z

Dunkl, I., Lovenduski, N., Collalti, A., Arora, V. K., Ilyina, T., and Brovkin, V. (2023). Gross primary productivity and the predictability of CO_2: more uncertainty in what we predict than how well we predict it. *Biogeosciences*, 20, 3523–3538, https://doi.org/10.5194/bg-20-3523-2023

Engel M., Vospernik S., Toigo M., Morin X., Tomao A., Trotta C., Steckel M., Barbati A., Nothdurft A. Pretzsch H., del Rio M., Skrzyszewski J., Ponette Q., Lof M., Jansons A., Brazaitis G. (2021). Simulating the effects of thinning and species mixing on stands of oak (*Quercus petrea* (Matt.) Liebl. / *Quercus robur* L.) and pine (*Pinus sylvestris* L.) across Europe. *Ecological Modelling*, 442(15):109406.https://doi.org/10.1016/j.ecolmodel.2020.109406

Fan, Y., Miguez-Macho, G., Jobbágy, E. G., Jackson, R. B., & Otero-Casal, C. (2017). Hydrologic regulation of plant rooting depth. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(40), 10572–10577. https://doi.org/10.1073/pnas.1712381114

Fang, J., & Lechowicz, M. J. (2006). Climatic limits for the present distribution of beech (Fagus L.) species in the world. *Journal of Biogeography*, *33*(10), 1804–1819. https://doi.org/10.1111/j.1365-2699.2006.01533.x

Fang, Y., Leung, L. R., Wolfe, B. T., Detto, M., Knox, R. G., McDowell, N. G., Grossiord, C., Xu, C., Christoffersen, B. O., Gentine, P., Koven, C. D., & Chambers, J. Q. (2021).

Disentangling the Effects of Vapor Pressure Deficit and Soil Water Availability on Canopy Conductance in a Seasonal Tropical Forest During the 2015 El Niño Drought. *Journal of Geophysical Research: Atmospheres*, *126*(10), 1–20. https://doi.org/10.1029/2021JD035004

Farquhar, G. D., Caemmerer, S., & Berry, J. A. (1980). A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species. *Planta*, *149*(1), 78-90–90. http://dx.doi.org/10.1007/BF00386231

Fernández-Martínez, M., Peñuelas, J., Chevallier, F., Ciais, P., Obersteiner, M., Rödenbeck, C., Sardans, J., Vicca, S., Yang, H., Sitch, S., Friedlingstein, P., Arora, V. K., Goll, D. S., Jain, A. K., Lombardozzi, D. L., McGuire, P. C., & Janssens, I. A. (2023). Diagnosing destabilization risk in global land carbon sinks. *Nature*, *615*(November 2021). https://doi.org/10.1038/s41586-023-05725-1

Fibbi, L., Chiesi, M., Moriondo, M., Bindi, M., Chirici, G., Papale, D., Gozzini, B., & Maselli, F. (2016). Correction of a 1 km daily rainfall dataset for modelling forest ecosystem processes in Italy. *Meteorological Applications*, *23*(2), 294–303. https://doi.org/10.1002/met.1554

Flexas, J., Diaz-Espejo, A., Gago, J., Gallé, A., Galmés, J., Gulías, J., & Medrano, H. (2014). Photosynthetic limitations in Mediterranean plants: A review. *Environmental and Experimental Botany*, *103*, 12–23. https://doi.org/10.1016/j.envexpbot.2013.09.002

Francini, S., McRoberts, R. E., Giannetti, F., Marchetti, M., Scarascia Mugnozza, G., & Chirici, G. (2021). The Three Indices Three Dimensions (3I3D) algorithm: a new method for forest disturbance mapping and area estimation based on optical remotely sensed imagery. *International Journal of Remote Sensing*, *42*(12), 4697–4715. https://doi.org/10.1080/01431161.2021.1899334

Fyllas, N. M., Christopoulou, A., Galanidis, A., Michelaki, C. Z., Dimitrakopoulos, P. G., Fulé, P. Z., & Arianoutsou, M. (2017). Tree growth-climate relationships in a forest-plot network on Mediterranean mountains. *Science of the Total Environment*, *598*, 393–403. https://doi.org/10.1016/j.scitotenv.2017.04.145

Gauquelin, T., Michon, G., Joffre, R., Duponnois, R., Génin, D., Fady, B., Bou Dagher-Kharrat, M., Derridj, A., Slimani, S., Badri, W., Alifriqui, M., Auclair, L., Simenel, R., Aderghal, M., Baudoin, E., Galiana, A., Prin, Y., Sanguin, H., Fernandez, C., & Baldy, V. (2018). Mediterranean forests, land use and climate change: a social-ecological perspective. *Regional Environmental Change*, *18*(3), 623–636. https://doi.org/10.1007/s10113-016-0994-3

Giannetti, F., Chirici, G., Vangi, E., Corona, P., Maselli, F., Chiesi, M., D'Amico, G., & Puletti, N. (2022). Wall-to-Wall Mapping of Forest Biomass and Wood Volume Increment in Italy. *Forests*, *13*(12). https://doi.org/10.3390/f13121989

Grossiord, C., Buckley, T. N., Cernusak, L. A., Novick, K. A., Poulter, B., Siegwolf, R. T. W., Sperry, J. S., & McDowell, N. G. (2020). Plant responses to rising vapor pressure deficit. *New Phytologist*, *226*(6), 1550–1566. https://doi.org/10.1111/nph.16485

Grünig, M., Seidl, R., & Senf, C. (2023). Increasing aridity causes larger and more severe forest fires across Europe. *Global Change Biology*, *29*(6), 1648–1659. https://doi.org/10.1111/gcb.16547

Gutsch, Lasch-Born, P., Kollas, C., Suckow, F., & Reyer, C. P. O. (2018). Balancing trade-offs

between ecosystem services in Germany's forests under climate change. *Environmental Research Letters*, *13*(4). https://doi.org/10.1088/1748-9326/aab4e5

Hanson, P. J., Amthor, J. S., Wullschleger, S. D., Wilson, K. B., Grant, R. F., Hartley, A., Hui, D., Hunt, E. R., Johnson, D. W., Kimball, J. S., King, A. W., Luo, Y., McNulty, S. G., Sun, G., Thornton, P. E., Wang, S., Williams, M., Baldocchi, D. D., & Cushman, R. M. (2004). Oak forest carbon and water simulations: Model intercomparisons and evaluations against independent data. *Ecological Monographs*, *74*(3), 443–489. https://doi.org/10.1890/03-4049

Härkönen, S., Neumann, M., Mues, V., Berninger, F., Bronisz, K., Cardellini, G., Chirici, G., Hasenauer, H., Koehl, M., Lang, M., Merganicova, K., Mohren, F., Moiseyev, A., Moreno, A., Mura, M., Muys, B., Olschofsky, K., Del Perugia, B., Rørstad, P. K., … Mäkelä, A. (2019). A climate-sensitive forest model for assessing impacts of forest management in Europe. *Environmental Modelling and Software*, *115*(February), 128–143. https://doi.org/10.1016/j.envsoft.2019.02.009

Haylock, M. R., Hofstra, N., Klein Tank, A. M. G., Klok, E. J., Jones, P. D., & New, M. (2008). A European daily high-resolution gridded data set of surface temperature and precipitation for 1950-2006. *Journal of Geophysical Research Atmospheres*, *113*(20), 1–12. https://doi.org/10.1029/2008JD010201

Herrera, S., Margarida Cardoso, R., Matos Soares, P., Espírito-Santo, F., Viterbo, P., & Gutiérrez, J. M. (2019). Iberia01: A new gridded dataset of daily precipitation and temperatures over Iberia. *Earth System Science Data*, *11*(4), 1947–1956. https://doi.org/10.5194/essd-11-1947-2019

Hiederer, R. (2013). Mapping Soil Properties for Europe - Spatial Representation of Soil Database Attributes. In *JRC Technical Reports*. http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Mapping+Soil+Properties+for+Europe+-+Spatial+Representation+of+Soil+Database+Attributes#0

Hijmans, R. J., Bivand, R., Forner, K., Ooms, J., Pebesma, E., & Sumner, M. D. (2022). *Package 'terra .' 3*(500), 1–6.

Janssen, P. H. M., & Heuberger, P. S. C. (1995). Calibration of process-oriented models. *Ecological Modelling*, *83*(1–2), 55–66. https://doi.org/10.1016/0304-3800(95)00084-9

Jung, M., Koirala, S., Weber, U., Ichii, K., Gans, F., Camps-Valls, G., Papale, D., Schwalm, C., Tramontana, G., & Reichstein, M. (2019). The FLUXCOM ensemble of global land-atmosphere energy fluxes. *Scientific Data*, *6*(1), 1–14. https://doi.org/10.1038/s41597-019-0076-8

Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard, S., Bodesheim, P., Carvalhais, N., Chevallier, F., Gans, F., S Goll, D., Haverd, V., Köhler, P., Ichii, K., K Jain, A., Liu, J., Lombardozzi, D., … Reichstein, M. (2020). Scaling carbon fluxes from eddy covariance sites to globe: Synthesis and evaluation of the FLUXCOM approach. *Biogeosciences*, *17*(5), 1343–1365. https://doi.org/10.5194/bg-17-1343-2020

Kannenberg, S. A., Schwalm, C. R., & Anderegg, W. R. L. (2020). Ghosts of the past: how drought legacy effects shape forest functioning and carbon cycling. *Ecology Letters*, *23*(5), 891–901. https://doi.org/10.1111/ele.13485

Kattge, J., & Knorr, W. (2007). Temperature acclimation in a biochemical model of photosynthesis: A reanalysis of data from 36 species. *Plant, Cell and Environment*. https://doi.org/10.1111/j.1365-3040.2007.01690.x

Keenan, T., García, R., Friend, A. D., Zaehle, S., Gracia, C., & Sabate, S. (2009). Improved understanding of drought controls on seasonal variation in mediterranean forest canopy CO_2 and water fluxes through combined in situ measurements and ecosystem modelling. *Biogeosciences*, 6(8), 1423–1444. https://doi.org/10.5194/bg-6-1423-2009

Keenan, T., Baker, I., Barr, A., Ciais, P., Davis, K., Dietze, M., Dragoni, D., Gough, C. M., Grant, R., Hollinger, D., Hufkens, K., Poulter, B., McCaughey, H., Raczka, B., Ryu, Y., Schaefer, K., Tian, H., Verbeeck, H., Zhao, M., and Richardson, A. D. (2012). Terrestrial biosphere model performances for inter-annual variability of land-atmosphere CO_2 exchange, *Glob. Change Biol.*, 18, 1971–1987

Koch, J., Demirel, M. C., & Stisen, S. (2018). The SPAtial EFficiency metric (SPAEF): Multiple-component evaluation of spatial patterns for optimization of hydrological models. *Geoscientific Model Development*, 11(5), 1873–1886. https://doi.org/10.5194/gmd-11-1873-2018

Kollet, S. J., & Maxwell, R. M. (2008). Capturing the influence of groundwater dynamics on land surface processes using an integrated, distributed watershed model. *Water Resources Research*, 44(2), 1–18. https://doi.org/10.1029/2007WR006004

Kumar, S. V., Peters-Lidard, C. D., Mocko, D., Reichle, R., Liu, Y., Arsenault, K. R., Xia, Y., Ek, M., Riggs, G., Livneh, B., & Cosh, M. (2014). Assimilation of remotely sensed soil moisture and snow depth retrievals for drought estimation. *Journal of Hydrometeorology*, 15(6), 2446–2469. https://doi.org/10.1175/JHM-D-13-0132.1

Li, X., & Xiao, J. (2019a). A global, 0.05-degree product of solar-induced chlorophyll fluorescence derived from OCO-2, MODIS, and reanalysis data. *Remote Sensing*, 11(5). https://doi.org/10.3390/rs11050517

Li, X., & Xiao, J. (2019b). Mapping photosynthesis solely from solar-induced chlorophyll fluorescence: A global, fine-resolution dataset of gross primary production derived from OCO-2. *Remote Sensing*, 11(21). https://doi.org/10.3390/rs11212563

Li, X., Xiao, J., He, B., Altaf Arain, M., Beringer, J., Desai, A. R., Emmel, C., Hollinger, D. Y., Krasnova, A., Mammarella, I., Noe, S. M., Ortiz, P. S., Rey-Sanchez, A. C., Rocha, A. V., & Varlagin, A. (2018). Solar-induced chlorophyll fluorescence is strongly correlated with terrestrial photosynthesis for a wide variety of biomes: First global analysis based on OCO-2 and flux tower observations. *Global Change Biology*, 24(9), 3990–4008. https://doi.org/10.1111/gcb.14297

Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News*, 2(3), 18–22.

Lindeskog, M., Smith, B., Lagergren, F., Sycheva, E., Ficko, A., Pretzsch, H., & Rammig, A. (2021). Accounting for forest management in the estimation of forest carbon balance using the dynamic vegetation model LPJ-GUESS (v4.0, r9710): Implementation and evaluation of simulations for Europe. *Geoscientific Model Development*, 14(10), 6071–6112. https://doi.org/10.5194/gmd-14-6071-2021

Lorenz, R., Stalhandske, Z., & Fischer, E. M. (2019). Detection of a Climate Change Signal in Extreme Heat, Heat Stress, and Cold in Europe From Observations. *Geophysical Research Letters*, *46*(14), 8363–8374. https://doi.org/10.1029/2019GL082062

Mahnken, M., Cailleret, M., Collalti, A., Trotta, C., Biondo, C., D'Andrea, E., Dalmonech, D., Marano, G., Mäkelä, A., Minunno, F., Peltoniemi, M., Trotsiuk, V., Nadal-Sala, D., Sabaté, S., Vallet, P., Aussenac, R., Cameron, D. R., Bohn, F. J., Grote, R., … Reyer, C. P. O. (2022). Accuracy, realism and general applicability of European forest models. *Global Change Biology*, *28*(23), 6921–6943. https://doi.org/10.1111/gcb.16384

Marchand, W., Buechling, A., Rydval, M., Čada, V., Stegehuis, A. I., Fruleux, A., Poláček, M., Hofmeister, J., Pavlin, J., Ralhan, D., Dušátko, M., Janda, P., Mikoláš, M., Vostarek, O., Bače, R., Frankovič, M., Kozák, D., Roibu, C. C., Chaskovskyy, O., … Svoboda, M. (2023). Accelerated growth rates of Norway spruce and European beech saplings from Europe's temperate primary forests are related to warmer conditions. *Agricultural and Forest Meteorology*, *329*(March 2022). https://doi.org/10.1016/j.agrformet.2022.109280

Marconi, S., Chiti, T., Nolè, A., Valentini, R., & Collalti, A. (2017). The role of respiration in estimation of net carbon cycle: Coupling soil carbon dynamics and canopy turnover in a novel version of 3D-CMCC forest ecosystem model. *Forests*, *8*(6). https://doi.org/10.3390/f8060220

Maréchaux, I., Langerwisch, F., Huth, A., Bugmann, H., Morin, X., Reyer, C. P. O., Seidl, R., Collalti, A., Dantas de Paula, M., Fischer, R., Gutsch, M., Lexer, M. J., Lischke, H., Rammig, A., Rödig, E., Sakschewski, B., Taubert, F., Thonicke, K., Vacchiano, G., & Bohn, F. J. (2021). Tackling unresolved questions in forest ecology: The past and future role of simulation models. *Ecology and Evolution*, *11*(9), 3746–3770. https://doi.org/10.1002/ece3.7391

Maselli, Barbati, A., Chiesi, M., Chirici, G., & Corona, P. (2006). Use of remotely sensed and ancillary data for estimating forest gross primary productivity in Italy. *Remote Sensing of Environment*, *100*(4), 563–575. https://doi.org/10.1016/j.rse.2005.11.010

Maselli, F., Argenti, G., Chiesi, M., Angeli, L., & Papale, D. (2013). Simulation of grassland productivity by the combination of ground and satellite data. *Agriculture, Ecosystems and Environment*, *165*, 163–172. https://doi.org/10.1016/j.agee.2012.11.006

Maselli, F., Pasqui, M., Chirici, G., Chiesi, M., Fibbi, L., Salvati, R., & Corona, P. (2012). Modeling primary production using a 1 km daily meteorological data set. *Climate Research*, *54*(3), 271–285. https://doi.org/10.3354/cr01121

Maselli, Papale, D., Puletti, N., Chirici, G., & Corona, P. (2009). Combining remote sensing and ancillary data to monitor the gross productivity of water-limited forest ecosystems. *Remote Sensing of Environment*, *113*(3), 657–667. https://doi.org/10.1016/j.rse.2008.11.008

Massari, C., Brocca, L., Pellarin, T., Abramowitz, G., Filippucci, P., Ciabatta, L., Maggioni, V., Kerr, Y., & Fernandez Prieto, D. (2020). A daily 25 km short-latency rainfall product for data-scarce regions based on the integration of the Global Precipitation Measurement mission rainfall and multiple-satellite soil moisture products. *Hydrology and Earth System Sciences*, *24*(5), 2687–2710. https://doi.org/10.5194/hess-24-2687-2020Massoud, E. C., Xu, C., Fisher, R. A., Knox, R. G., Walker, A. P., Serbin, S. P., Christoffersen, B. O., Holm, J. A., Kueppers, L. M., Ricciuto, D. M., Wei, L., Johnson, D.

J., Chambers, J. Q., Koven, C. D., McDowell, N. G., and Vrugt, J. A. (2019). Identification of key parameters controlling demographically structured vegetation dynamics in a land surface model: CLM4.5(FATES), *Geosci. Model Dev.*, 12, 4133–4164, https://doi.org/10.5194/gmd-12-4133-2019

Médail, F., Monnet, A. C., Pavon, D., Nikolic, T., Dimopoulos, P., Bacchetta, G., Arroyo, J., Barina, Z., Albassatneh, M. C., Domina, G., Fady, B., Matevski, V., Mifsud, S., & Leriche, A. (2019). What is a tree in the mediterranean basin hotspot? A critical analysis. *Forest Ecosystems*, 6(1). https://doi.org/10.1186/s40663-019-0170-6

Merganičová, K., Merganič, J., Lehtonen, A., Vacchiano, G., Sever, M. Z. O., Augustynczik, A. L. D., Grote, R., Kyselová, I., Mäkelä, A., Yousefpour, R., Krejza, J., Collalti, A., & Reyer, C. P. O. (2019). Forest carbon allocation modelling under climate change. *Tree Physiology*. https://doi.org/10.1093/treephys/tpz105

Minunno, F., Peltoniemi, M., Härkönen, S., Kalliokoski, T., Makinen, H., & Mäkelä, A. (2019). Bayesian calibration of a carbon balance model PREBAS using data from permanent growth experiments and national forest inventory. *Forest Ecology and Management*, 440(March), 208–257. https://doi.org/10.1016/j.foreco.2019.02.041

Moreno, A., & Hasenauer, H. (2016). Spatial downscaling of European climate data. *International Journal of Climatology*, 36(3), 1444–1458. https://doi.org/10.1002/joc.4436

Niu, G. Y., Yang, Z. L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., & Xia, Y. (2011). The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements. *Journal of Geophysical Research Atmospheres*, 116(12), 1–19. https://doi.org/10.1029/2010JD015139

Noce, S., Collalti, A., Valentini, R., & Santini, M. (2016). Hot spot maps of forest presence in the Mediterranean basin. *IForest*, 9(5), 766–774. https://doi.org/10.3832/ifor1802-009

Nord-Larsen, T., & Schumacher, J. (2012). Estimation of forest resources from a country wide laser scanning survey and national forest inventory data. *Remote Sensing of Environment*. https://doi.org/10.1016/j.rse.2011.12.022

Norton, A. J., Rayner, P. J., Koffi, E. N., Scholze, M., Silver, J. D., & Wang, Y. P. (2019). Estimating global gross primary productivity using chlorophyll fluorescence and a data assimilation system with the BETHY-SCOPE model. *Biogeosciences*, 16(15), 3069–3093. https://doi.org/10.5194/bg-16-3069-2019

Noto, L. V., Cipolla, G., Francipane, A., & Pumo, D. (2022). Climate Change in the Mediterranean Basin (Part I): Induced Alterations on Climate Forcings and Hydrological Processes. *Water Resources Management*, 2287–2305. https://doi.org/10.1007/s11269-022-03400-0

O'Sullivan, M., Smith, W. K., Sitch, S., Friedlingstein, P., Arora, V. K., Haverd, V., Jain, A. K., Kato, E., Kautz, M., Lombardozzi, D., Nabel, J. E. M. S., Tian, H., Vuichard, N., Wiltshire, A., Zhu, D., & Buermann, W. (2020). Climate-Driven Variability and Trends in Plant Productivity Over Recent Decades Based on Three Global Products. *Global Biogeochemical Cycles*, 34(12). https://doi.org/10.1029/2020GB006613

Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y. W., Poindexter, C.,

Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., Ardö, J., … Papale, D. (2020). The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data. *Scientific Data*, 7(1), 225. https://doi.org/10.1038/s41597-020-0534-3

Patacca, M., Lindner, M., Lucas-Borja, M. E., Cordonnier, T., Fidej, G., Gardiner, B., Hauf, Y., Jasinevičius, G., Labonne, S., Linkevičius, E., Mahnken, M., Milanovic, S., Nabuurs, G. J., Nagel, T. A., Nikinmaa, L., Panyatov, M., Bercak, R., Seidl, R., Ostrogović Sever, M. Z., … Schelhaas, M. J. (2023). Significant increase in natural disturbance impacts on European forests since 1950. *Global Change Biology*, 29(5), 1359–1376. https://doi.org/10.1111/gcb.16531

Peano, D., Materia, S., Collalti, A., Alessandri, A., Anav, A., Bombelli, A., & Gualdi, S. (2019). Global Variability of Simulated and Observed Vegetation Growing Season. *Journal of Geophysical Research: Biogeosciences*, 124(11), 3569–3587. https://doi.org/10.1029/2018JG004881

Peano, D., Hemming, D., Materia, S., Delire, C., Fan, Y., Joetzjer, E., Lee, H., Nabel, J. E. M. S., Park, T., Peylin, P., Wårlind, D., Wiltshire, A., and Zaehle, S. (2021). Plant phenology evaluation of CRESCENDO land surface models – Part 1: Start and end of the growing season, *Biogeosciences*, 18, 2405–2428, https://doi.org/10.5194/bg-18-2405-2021,

Peaucelle, M., Janssens, I. A., Stocker, B. D., Descals Ferrando, A., Fu, Y. H., Molowny-Horas, R., Ciais, P., & Peñuelas, J. (2019). Spatial variance of spring phenology in temperate deciduous forests is constrained by background climatic conditions. *Nature Communications*, 10(1), 1–10. https://doi.org/10.1038/s41467-019-13365-1

Peñuelas, J., Sardans, J., Filella, I., Estiarte, M., Llusià, J., Ogaya, R., Carnicer, J., Bartrons, M., Rivas-Ubach, A., Grau, O., Peguero, G., Margalef, O., Pla-Rabés, S., Stefanescu, C., Asensio, D., Preece, C., Liu, L., Verger, A., Barbeta, A., … Terradas, J. (2017). Impacts of global change on Mediterranean forests and their services. *Forests*, 8(12), 1–37. https://doi.org/10.3390/f8120463

Pretzsch, H., Grote, R., Reineking, B., Rötzer, T., & Seifert, S. (2008). Models for forest ecosystem management: A European perspective. *Annals of Botany*, 101(8), 1065–1087. https://doi.org/10.1093/aob/mcm246

Qiu, R., Han, G., Ma, X., Xu, H., Shi, T., & Zhang, M. (2020). A comparison of OCO-2 SIF, MODIS GPP, and GOSIF data from gross primary production (GPP) estimation and seasonal cycles in North America. *Remote Sensing*, 12(2). https://doi.org/10.3390/rs12020258

Reyer. (2015). Forest productivity under environmental change—A review of stand-scale modeling studies. *Current Forestry Reports*, 1(2), 53–68. https://doi.org/10.1007/s40725-015-0009-5

Reyer, C., Lasch-Born, P., Suckow, F., Gutsch, M., Murawski, A., & Pilz, T. (2014). Projections of regional changes in forest net primary productivity for different tree species in Europe driven by climate change and carbon dioxide. *Annals of Forest Science*, 71(2), 211–225. https://doi.org/10.1007/s13595-013-0306-8

Reyer, Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P.,

Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., … Frieler, K. (2020). The PROFOUND Database for evaluating vegetation models and simulating climate impacts on European forests. *Earth System Science Data*, *12*(2), 1295–1320. https://doi.org/10.5194/essd-12-1295-2020

Richards, F. J. (1959). A flexible growth function for empirical use. *Journal of Experimental Botany*, *10*(2), 290–301. https://doi.org/10.1093/jxb/10.2.290

Ripullone, F., Camarero, J. J., Colangelo, M., & Voltas, J. (2020). Variation in the access to deep soil water pools explains tree-to-tree differences in drought-triggered dieback of mediterranean oaks. *Tree Physiology*, *40*(5), 591–604. https://doi.org/10.1093/treephys/tpaa026

Rita, A., Camarero, J. J., Nolè, A., Borghetti, M., Brunetti, M., Pergola, N., Serio, C., Vicente-Serrano, S. M., Tramutoli, V., & Ripullone, F. (2020). The impact of drought spells on forests depends on site conditions: The case of 2017 summer heat wave in southern Europe. *Global Change Biology*, *26*(2), 851–863. https://doi.org/10.1111/gcb.14825

Rojas, R., Feyen, L., Dosio, A., & Bavera, D. (2011). Improving pan-European hydrological simulation of extreme events through statistical bias correction of RCM-driven climate simulations. *Hydrology and Earth System Sciences*, *15*(8), 2599–2620. https://doi.org/10.5194/hess-15-2599-2011

Sanchez-Ruiz, S., Chiesi, M., Fibbi, L., Carrara, A., Maselli, F., & Gilabert, M. A. (2018). Optimized Application of Biome-BGC for Modeling the Daily GPP of Natural Vegetation Over Peninsular Spain. *Journal of Geophysical Research: Biogeosciences*, *123*(2), 531–546. https://doi.org/10.1002/2017JG004360

Schenk, H. J., & Jackson, R. B. (2005). Mapping the global distribution of deep roots in relation to climate and soil characteristics. *Geoderma*, *126*(1-2 SPEC. ISS.), 129–140. https://doi.org/10.1016/j.geoderma.2004.11.018

Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M., Grams, T. E. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, E., Hoch, G., Holloway-Phillips, M., Körner, C., Larysch, E., Lübbe, T., Nelson, D. B., Rammig, A., … Kahmen, A. (2020). A first assessment of the impact of the extreme 2018 summer drought on Central European forests. *Basic and Applied Ecology*, *45*, 86–103. https://doi.org/10.1016/j.baae.2020.04.003

Senf, C., Buras, A., Zang, C. S., Rammig, A., & Seidl, R. (2020). Excess forest mortality is consistently linked to drought across Europe. *Nature Communications*, *11*(1), 1–8. https://doi.org/10.1038/s41467-020-19924-1

Stocker, B. D., Tumber-Dávila, S. J., Konings, A. G., Anderson, M. C., Hain, C., & Jackson, R. B. (2023). Global patterns of water storage in the rooting zones of vegetation. *Nature Geoscience*, *16*(March). https://doi.org/10.1038/s41561-023-01125-2

Suárez-Muñoz, M., Bonet-García, F. J., Navarro-Cerrillo, R., Herrero, J., & Mina, M. (2023). Forest management scenarios drive future dynamics of Mediterranean planted pine forests under climate change. *Landscape Ecology*, *0123456789*. https://doi.org/10.1007/s10980-023-01678-y

Sumner, M., & Hijmans, R. (2023). *Package 'rgdal.'*

Sun, Y., Frankenberg, C., Wood, J. D., Schimel, D. S., Jung, M., Guanter, L., Drewry, D. T., Verma, M., Porcar-Castell, A., Griffis, T. J., Gu, L., Magney, T. S., Köhler, P., Evans, B., & Yuen, K. (2017). OCO-2 advances photosynthesis observation from space via solar-induced chlorophyll fluorescence. *Science*, *358*(6360). https://doi.org/10.1126/science.aam5747

Sun Z., Wang W.,Zhag X., Tai H. Guo H., Yin S. Zhag T., (2019). A recent large review on RS-based products intercomparison confirmed the large variability and different sensitivity to climatic factors between different approaches and criteria when compared at site-level. *Science of The Total Environment*, 668, 696-713, 2019.

Tarquini, T., I. Isola, M. Favalli, F. Mazzarini, M. Bisson, M. T. Pareschi, & E. Boschi. (2009). TINITALY/01: a new Triangular Irregular Network of Italy. *Annals of Geophysics*, *50*(3). https://doi.org/10.4401/ag-4424

Tedeschi, V., Rey, A., Manca, G., Valentini, R., Jarvis, P. J., & Borghetti, M. (2006). Soil respiration in a Mediterranean oak forest at different developmental stages after coppicing. *Global Change Biology*, *12*(1), 110–121. https://doi.org/10.1111/j.1365-2486.2005.01081.x

Testolin, R., Dalmonech, D., Marano, G., Bagnara, M., D'Andrea, E., Matteucci, G., Noce, S., & Collalti, A. (2023). Simulating diverse forest management options in a changing climate on a Pinus nigra subsp. laricio plantation in Southern Italy. *Science of the Total Environment*, *857*(October 2022), 159361. https://doi.org/10.1016/j.scitotenv.2022.159361

Turco, M., Von Hardenberg, J., AghaKouchak, A., Llasat, M. C., Provenzale, A., & Trigo, R. M. (2017). On the key role of droughts in the dynamics of summer fires in Mediterranean Europe. *Scientific Reports*, *7*(1), 1–10. https://doi.org/10.1038/s41598-017-00116-9

Vacchiano G., Magnani F., Collalti A., "Modeling Italian forests: state of the art and future challenges", iForest – Biogeosciences and Forestry, 5: 113-120, https://doi.org/10.3832/ifor0614-005, 2012

Vangi, E., D'Amico, G., Francini, S., Borghi, C., Giannetti, F., Corona, P., Marchetti, M., Travaglini, D., Pellis, G., Vitullo, M., & Chirici, G. (2023). LARGE-SCALE high-resolution yearly modeling of forest growing stock volume and above-ground carbon pool. *Environmental Modelling and Software*. https://doi.org/10.1016/j.envsoft.2022.105580

Vangi, E., D'amico, G., Francini, S., Giannetti, F., Lasserre, B., Marchetti, M., McRoberts, R. E., & Chirici, G. (2021). The effect of forest mask quality in the wall-to-wall estimation of growing stock volume. *Remote Sensing*, *13*(5), 1–20. https://doi.org/10.3390/rs13051038

Verkerk, P. J., Delacote, P., Hurmekoski, E., Kunttu, J., Matthews, R., Mäkipää, R., Mosley, F., Perugini, L., Reyer, C. P. O., Roe, S., & Trømborg, E. (2022). *Forest-based climate change mitigation and adaptation in Europe.*

Veroustraete, F., Sabbe, H., & Eerens, H. (2002). Estimation of carbon mass fluxes over Europe using the C-Fix model and Euroflux data. *Remote Sensing of Environment*, *83*(3), 376–399. https://doi.org/10.1016/S0034-4257(02)00043-3

Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *Journal of Climate*, *23*(7), 1696–1718. https://doi.org/10.1175/2009JCLI2909.1

Vogel, J., Paton, E., Aich, V., & Bronstert, A. (2021). Increasing compound warm spells and droughts in the Mediterranean Basin. Weather and Climate Extremes, 32, 100312.

Walther, S., Duveiller, G., Jung, M., Guanter, L., Cescatti, A., & Camps-Valls, G. (2019). Satellite observations of the contrasting response of trees and grasses to variations in water availability, Geophysical Research Letters, 46, 1429–1440. https://doi.org/10.1029/2018GL080535

Wang, S., Zhang, Y., Ju, W., Qiu, B., & Zhang, Z. (2021). Tracking the seasonal and inter-annual variations of global gross primary production during last four decades using satellite near-infrared reflectance data. *Science of the Total Environment*, *755*. https://doi.org/10.1016/j.scitotenv.2020.142569

Waser, L. T., Ginzler, C., & Rehush, N. (2017). Wall-to-Wall tree type mapping from countrywide airborne remote sensing surveys. *Remote Sensing*. https://doi.org/10.3390/rs9080766

Wild, B., Teubner, I., Moesinger, L., Zotta, R., & Forkel, M. (2022). Gross Primary Production Dataset From Microwave Remote Sensing. *Earth System Science Data*, *14*, 1063–1085.

Wohlfahrt, G., Gerdel, K., Migliavacca, M., Rotenberg, E., Tatarinov, F., Müller, J., Hammerle, A., Julitta, T., Spielmann, F. M., & Yakir, D. (2018). Sun-induced fluorescence and gross primary productivity during a heat wave. *Scientific Reports*, *8*(1), 1–9. https://doi.org/10.1038/s41598-018-32602-z

Wu, Z., Ahlström, A., Smith, B., Ardö, J., Eklundh, L., Fensholt, R., & Lehsten, V. (2017). Climate data induced uncertainty in model-based estimations of terrestrial primary productivity. *Environmental Research Letters*, *12*(6). https://doi.org/10.1088/1748-9326/aa6fd8

Xiao, J., Li, X., He, B., Arain, M. A., Beringer, J., Desai, A. R., Emmel, C., Hollinger, D. Y., Krasnova, A., Mammarella, I., Noe, S. M., Serrano Ortiz, P., Rey-Sanchez, C., Rocha, A. V., & Varlagin, A. (2019). Solar-induced chlorophyll fluorescence exhibits a universal relationship with gross primary productivity across a wide variety of biomes. *Global Change Biology*, *25*(4), e4–e6. https://doi.org/10.1111/gcb.14565

Zampieri M., Grizzetti B., Toreti A., De Palma P., Collalti A., "Rise and fall of vegetation annual primary production resilience to climate variability projected by a large ensemble of Earth System Models' simulations", Environmental Research Letter, 16, 105001, https://doi.org/10.1088/1748-9326/ac2407, 2021

Zhang, F., Lu, X., Huang, Q., & Jiang, F. (2022). Impact of different ERA reanalysis data on GPP simulation. *Ecological Informatics*, *68*(August 2021), 101520. https://doi.org/10.1016/j.ecoinf.2021.101520

Zhang, Y., & Ye, A. (2022). Uncertainty analysis of multiple terrestrial gross primary productivity products. *Global Ecology and Biogeography*, *31*(11), 2204–2218. https://doi.org/10.1111/geb.13578

Zhang, & Ye, A. (2021). Would the obtainable gross primary productivity (GPP) products stand up? A critical assessment of 45 global GPP products. *Science of the Total Environment*, *783*, 146965. https://doi.org/10.1016/j.scitotenv.2021.146965

Zheng, Y., Shen, R., Wang, Y., Li, X., Liu, S., Liang, S., Chen, J. M., Ju, W. (2020). Zhang, L., and Yuan, W.: Improved estimate of global gross primary production for reproducing its long-term variation, 1982–2017, *Earth Syst. Sci. Data*, 12, 2725–2746, https://doi.org/10.5194/essd-12-2725-2020

1377

1378 **Figure 1a) Study area in the Italian peninsula and elevation map of the Region Basilicata. The red line**
1379 **indicates the administrative limits of the region, b) distribution of the dominant forest class at 1x1 km spatial**
1380 **resolution.**

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

Figure 2 3D-CMCC-FEM mean annual GPP values (gC m⁻² yr⁻¹) for the period 2005-2019 at 1 km spatial resolution. White areas indicate areas not simulated by the 3D-CMCC-FEM.

1403
Figure 2 3D-CMCC-FEM mean annual GPP values (gC m^{-2} yr^{-1}) for the period 2005-2019 at 1 km spatial
1404
resolution. White areas indicate areas not simulated by the 3D-CMCC-FEM.

1405

1406

1407

1408

1409

1410

1411

1412

1413

Figure 3 Root Mean Square Error (RMSE, gC m^{-2} yr^{-1}) between mean annual 3D-CMCC-FEM GPP (gC m^{-2} yr^{-1}) and the RS-based GPP: a) 3D-CMCC-FEM GPP vs. GOSIF GPP for the period 2005-2019, b) 3D-CMCC-FEM GPP vs. CFIX GPP for the period 2005-2013, c) 3D-CMCC-FEM GPP vs. FLUXCOM GPP for the period 2005-2015; d) histogram of the relative difference, dashed lines indicate the median value. White areas on the maps indicate areas not simulated by the 3D-CMCC-FEM.

1419

1420

1421

1422

1423

1424

Figure 4 Relative Difference (RD, %) between mean annual 3D-CMCC-FEM GPP and the RS-based GPP: a) 3D-CMCC-FEM GPP vs. GOSIF GPP for the period 2005-2019, b) 3D-CMCC-FEM GPP vs. CFIX GPP for the period 2005-2013, c) 3D-CMCC-FEM GPP vs. FLUXCOM GPP for the period 2005-2015; d) histogram of the relative difference, dashed lines indicate the median value. White areas on the maps indicate areas not simulated by the 3D-CMCC-FEM.

1430

1431

1432

1433

1434

1435

1436

Figure 5 Fractional Variance (FV) between mean annual 3D-CMCC-FEM GPP and the RS-based GPP: a)
3D-CMCC-FEM GPP vs. GOSIF GPP for the period 2005-2019, b) 3D-CMCC-FEM GPP vs. CFIX GPP for
the period 2005-2013, c) 3D-CMCC-FEM GPP vs. FLUXCOM GPP for the period 2005-2015; d) histogram
of the relative difference, dashed lines indicate the median value. White areas on the maps indicate areas not
simulated by the 3D-CMCC-FEM.

1442

1443

1444

1445

1446

1447

1448

1449

Figure 6 Pearson's correlation (*r*) between mean annual 3D-CMCC-FEM GPP and the RS-based GPP: a) 3D-CMCC-FEM GPP vs. GOSIF GPP for the period 2005-2019, b) 3D-CMCC-FEM GPP vs. CFIX GPP for the period 2005-2013, c) 3D-CMCC-FEM GPP vs. FLUXCOM GPP for the period 2005-2015; d) histogram of the relative difference, dashed lines indicate the median value. Grey areas on the maps indicate where correlations are not significant. White areas on the maps indicate areas not simulated by the 3D-CMCC-FEM.

1466
1467

Figure 7 Pearson's correlation (*r*) between Mean Seasonal Cycle (MSC) 3D-CMCC-FEM GPP and the RS-based GPP: a) 3D-CMCC-FEM GPP vs. GOSIF GPP for the period 2005-2019, b) 3D-CMCC-FEM GPP vs. CFIX GPP for the period 2005-2013, c) 3D-CMCC-FEM GPP vs. FLUXCOM GPP for the period 2005-2015, d) histogram of the r, dashed lines indicate the median value. White areas on the maps indicate areas not simulated by the 3D-CMCC-FEM.

1473

1474

1475

1476

1477

1478

1479

1480

Figure 8 Spatial distribution of the slope of the linear regression of the summer GPP residuals and SPEI5 for: a) 3D-CMCC-FEM vs. GOSIF, b) 3D-CMCC-FEM vs. CFIX, c) 3D-CMCC-FEM vs. FLUXCOM. Grey areas indicate the slope is not significantly different from 0 at p<0.05. White areas on the maps indicate areas not simulated by the 3D-CMCC-FEM.

1485

1486

1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

Figure 9 Mean annual 3D-CMCC-FEM GPP vs. RS-based GPP scatter plot (gC m^{-2} yr^{-1}) at the species-level: a) the 3D-CMCC-FEM GPP vs. GOSIF b) 3D-CMCC-FEM GPP vs. CFIX and c) 3D-CMCC-FEM GPP vs. FLUXCOM estimates (black line is the 1:1 line, dashed line is the linear fit). Each point represents data at grid cell level, different colors indicate the different species considered.

1510