Modeling soybean growth: A mixed model approach

Maud Delattre [1*], Yusuke Toda[2], Jessica Tressou[2,3], Hiroyoshi Iwata[2].

1 Université Paris-Saclay, INRAE, MaIAGE, Jouy-en-Josas, France
2 Graduate School of Agricultural and Life Sciences,The University of Tokyo, Tokyo, Japan
3 Paris-Saclay University-AgroParisTech-INRAE, UMR MIA-Paris-Saclay, Palaiseau, France

* maud.delattre@inrae.fr

Abstract

The evaluation of plant and animal growth, separately for genetic and environmental effects, is necessary for genetic understanding and genetic improvement of environmental responses of plants and animals. We propose an original approach that combines nonlinear mixed-effects model (NLME) and the stochastic approximation of the Expectation-Maximization algorithm (SAEM) to analyze genetic and environmental effects on plant growth. These tools are widely used in many fields but very rarely in plant biology. During model formulation, a nonlinear mechanistic function describes the shape of growth, and random effects describe genetic and environmental effects and their variability. Genetic relationships among the varieties were also integrated into the model using a kinship matrix. The SAEM algorithm was chosen as an efficient alternative to MCMC methods, which are more commonly used in the domain. It was implemented to infer the expected growth patterns in the analyzed population and the expected curves for each variety through a maximum-likelihood and a maximum-a-posteriori approaches, respectively. The obtained estimates can be used to predict the growth curves for each variety. We illustrate the strengths of the proposed approach using simulated data and soybean plant growth data obtained from a soybean cultivation experiment conducted

at the Arid Land Research Center, Tottori University. In this experiment, plant height was measured daily using drones, and the growth was monitored for approximately 200 soybean cultivars for which whole-genome sequence data were available. The NLME approach improved our understanding of the determinants of soybean growth and can be successfully used for the genomic prediction of growth pattern characteristics.

Author summary

Nonlinear mechanistic models are useful for modeling animal and plant growth; however, their parameters are influenced by both genetic and environmental factors. If the same model can be applied to data with different genetic and environmental factors by allowing parameter variations, it can be used to understand, predict, and control the genetic and environmental influences of growth models based on parameter variation. In this study, we propose a statistical method for integrating a nonlinear mixed-effects model with a nonlinear mechanistic model. The simulation and real data analysis results show that the proposed method was effective in modeling the growth of genetically different soybean varieties under different drought conditions. The usefulness of the proposed method is expected to increase, as high-throughput measurements provide growth data for a large number of genotypes in various environments.

1 Introduction

Controlling and predicting the growth patterns of plants and animals are important for growing crops and raising livestock. For example, in crops, growth patterns are known to be closely related to the amount of carbon fixation and thus harvested product [1]. Plant growth patterns are determined by genetics and the environment, and if we could model the relationship between genetics, environment, and growth patterns, we can control and predict the growth based on that model; for example, by properly managing the cultivation of varieties with desirable growth patterns, which can have a positive impact on the yield and quality of the harvest. From this perspective, many recent attempts have been made to measure the growth over time and to model the growth data obtained. In particular, in crop science, the development of remote-sensing

technologies, such as unmanned aerial vehicles [2] and proximal sensing [3,4], has enabled the modeling of the growth curves of many genotypes simultaneously [5–7].

As mentioned earlier, growth patterns are determined not only by genetics and the environment but also by physiological constraints. Owing to physiological constraints, growth patterns often exhibit a nonlinear pattern in which growth is slow in the early stages, accelerates in the mid-term, but slows down after a certain period, and reaches a plateau in the late stages. Such patterns are often approximated by nonlinear functions such as a logistic curve [8] or a Gompertz curve [9]. Because growth patterns are determined by genetic and environmental factors, the parameters of the nonlinear functions describing the growth patterns change under the influence of these factors. Based on this perspective, modeling was conducted in a two-step process: the growth patterns were first approximated by a nonlinear function, and thereafter the estimated parameters of the function were modeled based on the effects of genetic and environmental factors [10–13]. For example, Toda et al. (2022) applied a two-step process to growth data (changes in leaf area index) of various soybean cultivars grown under different growing conditions [13]. They first approximated the observed growth using a nonlinear function and then constructed a model in which the estimated parameters were explained by the genetic effects of genome-wide marker genotypes and variations due to blocks. Modeling the genetic effects of genome-wide marker genotypes is a technique known as genomic prediction [14], and these predictions are used to select superior genotypes in plant and animal breeding. Toda et al. (2022) combined the genomic prediction potential of nonlinear functional parameters with early growth data under a Bayesian framework and showed that such a combination could accurately predict growth patterns in later growth stages.

The two-step approach described above is relatively easy to apply but has a drawback. The first step approximates the growth of each individual or plots separately with a nonlinear function; however, this approximation does not consider the factors behind the parameters of the nonlinear model, that is, genetic and environmental effects. For example, given a pair of genetically related samples, these samples are likely to exhibit similar growth and the parameters of the nonlinear model are likely to exhibit similar values. In the second stage of modeling, such factors that are ignored in the first stage are also considered [15]. If the underlying genetic and environmental factors are

considered during nonlinear curve fitting, one sample can "borrow information" from other genetically related samples during the estimation, preventing over-fitting to individual growth patterns resulting in a more robust model.

Therefore, quantitative trait locus (QTL) analysis models have been developed using a one-step approach in which parameters are represented by a linear combination of locus effects [16,17]. However, this model assumes that the effects of a single QTL are estimated and tested individually as fixed effects, and it cannot be used to make genomic predictions for traits that are controlled by multiple medium or small gene effects. In genomic selection, the effects of genome-wide markers are generally modeled as random effects [14,18,19]. Attempting to model the variation in the parameters of a nonlinear function with both random and fixed effects requires a framework that integrates mixed effects into a nonlinear function, that is, a nonlinear mixed-effects model (NLMEM). A few examples of NLMEMs have been applied to the genomic prediction of growth patterns [6,20,21], but they have not yet been fully explored.

NLMEMs are statistical models typically used to describe dynamic phenomena from repeated measurements of several individuals [22,23]. They allow the simultaneous analysis of the data of several subjects in a single model by describing the variability of the phenomenon of interest in the studied population and the specificity of each individual at the same time. During many applications, such as in pharmacokinetics or in the study of growth curves, these models are generally utilized as mechanistic models with random effects on the mechanistic parameters. Statistical inferences in these models provide access to multiple pieces of information. Estimation of the mean values of the mechanistic parameters and their variances allowed us to describe the phenomenon of global interest in the population. Some estimates of the individual parameters could also be obtained from these models, making prediction at the individual level as well as the comparison of the individual responses, for example, to select plants with high genetic abilities, possible. The advantage of NLMEMs is that all information can be inferred from the same statistical model. Studies on plant growth and genetic variation among cultivars are generally based on measuring the same phenotypic traits multiple times over time in plants of different cultivars. Therefore, it was appropriate to use mixed-effects models for this purpose. Although this approach was proposed several years ago in the field of animal genetics [12,24], it is not widely

used. One of the main objectives of this study was to illustrate the relevance of ⁹⁶

NLMEM in plant breeding when available data were repeated from several plants of ⁷⁷

different varieties. ⁷⁸

In this study, NLMEM was applied to remotely sensed data of soybean growth ⁷⁹

during cultivation trials under different irrigation conditions. In the experiments used ⁸⁰

for the analysis, the whole-genome sequences of all the varieties were decoded, allowing ⁸¹

their genetic relationships to be incorporated into the model. The influence of genetic ⁸²

relationships on the irrigation effects was also modeled. Specific tools are required to ⁸³

estimate and predict parameters from the NLMEM. Here, we used a framework that ⁸⁴

employs an algorithm derived from the Expectation-Maximization (EM) algorithm [25], ⁸⁵

that is, the stochastic Approximation EM (SAEM) algorithm [26], which is widely used ⁸⁶

in the NLMEM community and is very efficient in terms of computation time compared ⁸⁷

with those of standard MCMC methods. The effectiveness of NLMEM was examined by ⁸⁸

applying it to simulated data generated for a cultivation trial and observed data. ⁸⁹

2 Materials and methods ⁹⁰

2.1 Data description ⁹¹

A field trial was conducted from 2017 to 2022 in an experimental field with sandy soil at ⁹²

the Arid Land Research Center, Tottori University (35 °32 ′ N latitude, 134 °12′ E ⁹³

longitude, 14 m. above sea level). Soybean accessions registered in the National ⁹⁴

Agriculture and Food Research Organization Genebank ⁹⁵

(`https://www.gene.affrc.go.jp/databases-core_collections_jw_en.php` and ⁹⁶

`https://www.gene.affrc.go.jp/databases-core_collections_wg_en.php`) were ⁹⁷

used in this study. A total of 198 accessions consisted of 96 Japanese accessions, 96 ⁹⁸

world accessions from mini-core collections [27], and six additional accessions ⁹⁹

(GmWMC160, GmWMC157, GmWMC163, GmJMC002, GmJMC003, and ¹⁰⁰

GmJMC033). In 2017, 188 out of the 198 accessions were used for the analysis. Each ¹⁰¹

plot contained four plants. Sowing was performed at the beginning of July each year, ¹⁰²

followed by thinning after two weeks. Two watering treatment levels, control (C) and ¹⁰³

drought (D), were used to evaluate the genetic variations in the responses to water ¹⁰⁴

stress. Irrigation was applied daily for 5 h (7:00–9:00, 12:00–14:00, and 16:00–17:00), 105

starting the day after the thinning in treatment C, while no water was applied in 106

treatment D. Plant growth was measured both manually and by remote sensing for 107

some plants, which allowed the calibration of the UAV data. The R package missForest 108

was used to perform nonparametric missing value imputation using random forest. In 109

the following section, we describe the analysis of UAV data collected in 2017. In 2017, 110

there were up to 10 UAV measurements per plant (1-56 days after sowing). Weather 111

data were used to convert the time units, expressed in terms of the number of days after 112

sowing, into heat units. 113

2.2 Modeling 114

We used NLMEMs with two levels of random effects to study genetic variability among 115

soybean varieties and to perform varietal selection. The general model is first described 116

before presenting the model used in this study. 117

2.2.1 Description of nonlinear mixed models with genetic effects 118

We assume N_p plots from N_v varieties $N_v < N_p$ were observed. Let n_i be the number of 119

observations for plot $i = 1, \ldots, N_p$, and $y_{ij} \in \mathbf{R}$ denote the jth observation of plot i 120

obtained under the time condition t_{ij}. The most generic way to formulate a NLMEM is 121

as a hierarchical model, with each stratum describing a different level of variability in 122

the data (see *e.g.*, [22,23]). As in [24], the following form is used: 123

$$
\begin{aligned}
y_{ij} &= g(\varphi_i, t_{ij}) + \epsilon_{ij} \ , \ \epsilon_{ij} \underset{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2), &(1)\\
\varphi_i &= X_i\beta + Z_i u + p_i \ , \ u \sim \mathcal{N}(0, K \otimes G) \ , \ p_i \underset{i.i.d.}{\sim} \mathcal{N}(0, P). &(2)
\end{aligned}
$$

The first stratum, given by Equation (1), describes the data for each plot with the 124

same statistical model but different parameters from one plot to another. $g(\varphi, t)$ is a 125

known (nonlinear) function parameterized by $\varphi \in \mathbf{R}^m$, which describes the evolution of 126

the growth process over time in an appropriate and often mechanistic manner. The 127

second stratum, stated by Equation (2), aims at describing the differences between plots 128

by means of covariates (*i.e.*, explanatory measured variables characterizing the plots or 129

their cultural conditions, arranged in $m \times p$ matrix X_i and $m \times q.N_v$ matrix Z_i with q the number of genetic effects per variety), genetic varietal effects $u \in \mathbf{R}^{q.N_v}$ and residual effects $p_i \in \mathbf{R}^m$. More precisely, the term $X_i\beta$ describes the covariate effects common to all varieties using the fixed-effects vector $\beta \in \mathbf{R}^p$, whereas the term Z_iu describes the covariate effects that differ from one variety to another using the genetic effect u. Therefore, estimating u is of interest for varietal selection, whereas the random effects p_i represent a part of the variability between plots, which is not due to covariates or genetics. The random varietal effect (u) and plot effect (p_i) are assumed to be independent centered Gaussian random variables whose respective variances are specified by means of covariance matrices G and P and the kinship matrix K between the varieties. Note that this model is very flexible as it can be easily adapted according to the observed growth patterns simply by choosing an appropriate function g while still allowing an analysis of the genetic variability between varieties.

To simplify the notation below, we note that θ is the set of unknown parameters in the model (1)-(2), composed of β, P, G, and σ^2.

2.2.2 Examples

We now consider specific cases that were used in the simulation study and the analysis of soybean experimental data.

The logistic growth curve   is a commonly used growth curve where function g is given by:

$$g(\varphi_i, t_{ij}) = \frac{\varphi_{i1}}{1 + \exp\left(-\gamma(\varphi_{i2} - t_{ij})\right)}$$

We considered a logistic growth curve with a constant and known scale parameter (γ), random asymptotic value (φ_{i1}), and the time of half-maximum height (φ_{i2}). We considered no covariate effects or simple genetic effects on these parameters and thus defined $\varphi_i = (\varphi_{i1}, \varphi_{i2})^\top$ as

$$\varphi_{i1} = \mu_A + u_{v_i,A} + p_{i1},$$

$$\varphi_{i2} = \mu_x + u_{v_i,x} + p_{i2},$$

where $v_i \in \{1, \dots, N_v\}$ is the variety of plot i, vector $u_v = (u_{v,A}, u_{v,x})^\top$ is the vector of genetic effects for variety v and $p_i = (p_{i1}, p_{i2})^\top \underset{i.i.d.}{\sim} \mathcal{N}(0, P)$. The vector u was obtained by the concatenation of the N_v genetic effect vectors u_v's such that $u \sim \mathcal{N}(0, K \otimes G)$ with the genetic covariance matrix G and the kinship matrix between the N_v observed varieties K. To connect with the general notations introduced in equations (1) and (2), the fixed effects vector in this model was $\beta = (\mu_A, \mu_x)^T$ and the covariate matrices are respectively $X_i = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and the N_v-block matrix $Z_i = \begin{pmatrix} \mathbf{0} & \dots & Z_{v_i} & \dots & \mathbf{0} \end{pmatrix}$ with $Z_{v_i} = X_i$. Matrices P and G were both 2 by 2 matrices.

The asymptotic regression curve is another commonly used growth curve. Function g was defined as follows:

$$g(\varphi_i, t_{ij}) = \varphi_{i1}(1 - \exp(-\exp(\varphi_{i2})t_{ij})). \tag{3}$$

During the analysis of the soybean growth data presented in Section 3.2, we considered the cross-effect between genetics and irrigation conditions on both parameters φ_{i1} and φ_{i2} by defining $\varphi_i = (\varphi_{i1}, \varphi_{i2})^\top$ as

$$\varphi_{i1} = (\mu_A + u_{v_i,A}) + x_i(\delta_A + u_{v_i,\delta_A}) + p_{i1},$$
$$\varphi_{i2} = (\mu_B + u_{v_i,B}) + x_i(\delta_B + u_{v_i,\delta_B}) + p_{i2}, \tag{4}$$

where x_i denotes the water condition plot i is subject to ($x_i = 1$ normal, $x_i = 0$ dry), v_i denotes the variety of plot i, and $p_i = (p_{i1}, p_{i2})^\top \underset{i.i.d.}{\sim} \mathcal{N}(0, P)$. The vector of genetic effects for variety $v \in \{1, \dots, N_v\}$ was composed of four components: $u_v = (u_{v,A}, u_{v,B}, u_{v,\delta_A}, u_{v,\delta_B})^\top$. The vector u obtained by the concatenation of the u_v's is such that $u \sim \mathcal{N}(0, K \otimes G)$ with the genetic covariance matrix G and the kinship matrix between the N_v observed varieties K. Here, the fixed effects vector was $\beta = (\mu_A, \mu_B, \delta_A, \delta_B)^T$ and the covariate matrices were $X_i = \begin{pmatrix} 1 & 0 & x_i & 0 \\ 0 & 1 & 0 & x_i \end{pmatrix}$ and the N_v-block matrix $Z_i = \begin{pmatrix} \mathbf{0} & \dots & Z_{v_i} & \dots & \mathbf{0} \end{pmatrix}$ with $Z_{v_i} = X_i$. P and G were respectively a 2 by 2 and a 4 by 4 covariance matrices.

2.3   Algorithms

We intended to process the growth data by performing both a maximum likelihood estimation of the parameters and a maximum a posteriori estimation of the predicted genetic effects by using a unique statistical model for both tasks.

Model (1)-(2) is a nonlinear model with the random effects u and p_i as latent variables. Because of these two features, neither the likelihood of observations nor the posterior density of genetic effects has an explicit form. The inferences in the model (1)-(2) fall within the scope of EM-like algorithms. Here, owing to the model nonlinearity, we propose to tackle both parameter estimation and genetic effects prediction using the SAEM algorithm [26] instead of the EM algorithm.

2.3.1   General overview of the SAEM algorithm

The EM algorithm [25] is an iterative algorithm in which the estimate of an unknown parameter is updated by maximizing the expectation of the log-likelihood of the complete data (*i.e.*, observed and latent data) conditional on the observations. Considering a general latent variable model with y and z as the observed and the unobserved variables respectively, iteration $[k]$ of the EM algorithm requires the computation of $\mathbf{E}[\log p(y, z; \theta)|y, \theta^{[k-1]}]$ with $\theta^{[k-1]}$ the estimation of parameter θ obtained at the previous iteration $[k-1]$. When $\mathbf{E}[\log p(y, z; \theta)|y, \theta^{[k-1]}]$ is not available in a closed form, the SAEM algorithm replaces its exact calculation with a stochastic approximation as follows:

$$\mathcal{Q}(\theta)^{[k]} = \mathcal{Q}(\theta)^{[k-1]} + \gamma_k \left(\log p(y, z^{[k]}; \theta) - \mathcal{Q}(\theta)^{[k-1]} \right),$$

where $z^{[k]}$ is a random draw from the conditional distribution $p(.|y, \theta^{[k-1]})$, either directly or using Monte Carlo methods, if the exact simulation under $p(.|y, \theta^{[k-1]})$ is unfeasible [28]. $(\gamma_k)_{k>0}$ is a sequence of positive decreasing step sizes, such that $\gamma_1 = 1$, $\sum \gamma_k = \infty$ and $\sum \gamma_k^2 < \infty$. Hence, the calculation of the complete log-likelihood and simulation of the latent data are two central points in the implementation of the algorithm.

2.3.2 Parameter estimation

Within the model (1)-(2), the observations are the available height measures from the N_p plots, $y = \{y_i\}_{1 \leqslant i \leqslant N_p}$ where $y_i = (y_{i1}, \ldots, y_{in_i})^\top$ is the vector of observations for plot i, and as suggested in [24], a possible and convenient choice for the missing data is $z = (\varphi, u)$, where $\varphi = \{\varphi_i\}_{1 \leqslant i \leqslant N_p}$ is the set of plant growth parameters and u is the vector of genetic effects. Therefore, the likelihood of the complete data can therefore be decomposed into:

$$p(y, \varphi, u) = p(y \mid \varphi, u)p(\varphi \mid u)p(u),$$

where each element of the right-hand term represents an easily computable Gaussian density. An exact simulation under $p(\varphi, u|y; \theta)$ is not feasible, but the Metropolis within Gibbs [29] is easy to implement by exploiting the hierarchical dependency between y, φ, and u and using that the exact simulation under $p(u|\varphi; \theta)$ is feasible. Iteration $[k]$ of the algorithm can thus be implemented as follows:

- *Simulation step:*

  - Draw $\varphi^{[k]}$ under $q_{\theta[k-1]}(\cdot)$ a transition kernel of a Monte Carlo Markov Chain algorithm targeting $p(\cdot|u^{[k-1]}, y, \theta^{[k-1]})$ as a stationary distribution

  - Draw $u^{[k]}$ under $p(u|\varphi^{[k]}, \theta^{[k-1]})$

- *Stochastic Approximation step:* compute

$$\mathcal{Q}(\theta)^{[k]} = \mathcal{Q}(\theta)^{[k-1]} + \gamma_k \left(\log p(y, \varphi^{[k]}, u^{[k]}; \theta) - \mathcal{Q}(\theta)^{[k-1]} \right),$$

  where $\gamma_1 = 1$, $\sum \gamma_k = \infty$ and $\sum \gamma_k^2 < \infty$.

- *Maximization step:* compute

$$\theta^{[k]} = \arg\max_\theta \mathcal{Q}(\theta)^{[k]}.$$

The succession of these three steps is repeated until the algorithm convergences. Implementation details and justifications are provided in Appendix S1 (see Algorithm 2).

2.3.3 Prediction of the varietal genetic effects

The varietal genetic effects can be predicted by computing

$$\hat{u} = \arg\max_u p(u|y, \hat{\theta}),$$

where $\hat{\theta}$ denotes the estimated parameter value obtained at the convergence of the algorithm described in the previous section. The conditional density $p(u|y, \theta)$ has no explicit form in model (1)-(2), but as $p(u|y, \theta) = \int p(u|\varphi; \theta)p(\varphi|y, \theta)d\varphi$, it is possible to use the SAEM algorithm again by considering φ as the latent variable and u as the element to be estimated. Iteration $[k]$ of the obtained algorithm is decomposed as follows:

- *Simulation step:* draw $\varphi^{[k]}$ under $p(\varphi|y, u^{[k-1]}, \hat{\theta})$ under $q_{\hat{\theta}}(\cdot)$ a transition kernel of a Monte Carlo Markov Chain algorithm targeting $p(\cdot|u^{[k-1]}, y, \hat{\theta})$ as a stationary distribution

- *Stochastic Approximation step:* compute

$$\mathcal{Q}(u)^{[k]} = \mathcal{Q}(u)^{[k-1]} + \gamma_k \left(\log p(y, \varphi^{[k]}|u; \hat{\theta}) - \mathcal{Q}(u)^{[k-1]} \right),$$

  where $\gamma_1 = 1$, $\sum \gamma_k = \infty$ and $\sum \gamma_k^2 < \infty$.

- *Maximization step:* compute

$$u^{[k]} = \arg\max_u \mathcal{Q}(u)^{[k]}.$$

Implementation details are provided in the Appendix (see Algorithm 2).

2.4 Numerical study design

We used the two-parameter logistic growth model introduced in section 2.2.2 with the following parameter values: $\mu_A = 50$, $\mu_x = 13$, $P = \begin{pmatrix} 6 & 0 \\ 0 & 0.4 \end{pmatrix}$, $G = \begin{pmatrix} 25 & 0 \\ 0 & 1.5 \end{pmatrix}$, and $\sigma^2 = 4$. Growth data are simulated on time interval $[1, 45]$ according to different scenarios to illustrate the performance of our approach based on nonlinear mixed-effects

models and the SAEM algorithm in terms of estimation and prediction, and to make recommendations for data collection.

1. *Scenario 1* is used to study the effect of the number of observed varieties and of the number of plots per variety on the performances of our methodology. The total number of plots is $N_p = 2000$, and the number of observed varieties is either $N_v = 50$ or $N_v = 100$. We consider a high frequency of data collection over the curves by simulating one data point at each unit time step leading to 45 observations per plot.

2. *Scenario 2* is used to study the effect of sampling frequency along the curves. We use for N_v and N_p a similar value as for the real data, $N_p = 800$ and $N_v = 200$. We simulate regularly spaced observations along the plant growth curves, with a total of either 45 (rich design) or 10 (sparse design) observations per curve.

50 datasets are simulated for each scenario and for each combination of N_p, N_v, and sampling frequency along the curves. The SAEM algorithm was applied to each of them to estimate the population parameters and then predict the genetic values of the N_v varieties. Both algorithms are run for 1500 iterations in total, with step sizes for the stochastic approximation defined as $\gamma_k = 1$ if $k < 1300$ and $\gamma_k = (k - 1300)^{-2/3}$ otherwise. This setting corresponds to defining a burn-in phase of 1300 iterations aimed at converging quickly to the neighborhood of the maximum likelihood estimator, followed by a refining phase in which the estimation is smoothed by decreasing the step sizes. Note that we use here a large number of iterations to guarantee good convergence of the algorithms on all simulated datasets; however, in practice, we observe that the convergence is reached far before the end of the 1500 iterations.

3 Results

3.1 Simulations

3.1.1 Parameter estimation

The simulation results related to parameter estimation are depicted in Fig 1, 2, 3 for Scenario 1, and on Fig 4, 5, 6 for Scenario 2. We see that all the parameters are

correctly estimated, *i.e.*, with very limited bias,regardless of the number of varieties, 252

plots per variety, and time points per curve used for the data simulation. 253

From Fig 1, 2, and 3, it appears that for the values considered for the number of 254

varieties considered in Scenario 1, changing N_v primarily affects the variance of the 255

estimates. Specifically, increasing N_v increases the precision of the intercept and genetic 256

variance estimates. However, increasing N_v reduces the precision of the estimate of the 257

covariance matrix P. The rationale for this behavior could be that, because this 258

experiment is conducted on a fixed total number of simulated plots (N_p), increasing N_v 259

resulted in a decrease in the number of observed plots per variety. Increasing the 260

number of observed plots per variety improved the quality of the estimations of P. 261

Fig 4, 5, and 6 show that for simulated data with similar values of N_v and N_p to 262

the real data analyzed in the next section, the model parameters are also estimated 263

with very limited bias. For these values of N_v and N_p, decreasing the number of 264

observations per curve does not bias the estimates, but increases their variance. Thus, 265

the experiments conducted under the second simulation scenario thus show that it is 266

possible to estimate satisfactorily the parameters of an NLMEM of the form (1)-(2) 267

under sampling conditions similar to those of the real data collected in Tottori in 2017. 268

[Fig 1 about here.] 269

[Fig 2 about here.] 270

[Fig 3 about here.] 271

[Fig 4 about here.] 272

[Fig 5 about here.] 273

[Fig 6 about here.] 274

3.1.2   Prediction of genetic effects
275

The results obtained for the predictions of the genetic effects under both experimental 276

scenarios are listed in Table 1. To quantify the effect of the parameter estimation error 277

on the quality of the prediction, we computed both the predictions obtained using the 278

true values of the parameters (column *θ known*) and those obtained with our complete 279

methodology, which requires estimating the parameters by SAEM in the first step (column θ *estimated*). In both cases, the average squared differences between true genetic effects and their predictions were computed for the 50 simulated datasets. We observed that the quality of predictions is always better when θ is known, that is, when the prediction is not affected by any parameter estimation error. Nevertheless, in all cases, the increase in the prediction error is minor because of the small estimation error (Figs 1 to 6). The results obtained in scenarios 1 and 2 show the significant impacts of the number of plots observed per variety (N_p/N_v) and of the sampling frequency along the growth curves, respectively. The predictions improved as N_p/N_v and the number of measurements along the curves increased.

[Table 1 about here.]

3.2 Real data analysis

The dataset was composed of $N_p = 752$ plots of $N_v = 188$ varieties observed at 10 time-points. Some data transformation was required before applying the NLMEM approach to the 2017 Tottori UAV data. The growth dynamics represented in Fig 7 correspond to that of logistic growth, but the observations show strong heteroscedasticity over time. To avoid strong numerical instabilities, the logarithms of the measured heights rather than on the heights themselves were used. The log growth curves are represented in Fig 8. Their shape suggests the use of an asymptotic regression curve for their analysis, as defined in Equation (3).

[Fig 7 about here.]

[Fig 8 about here.]

In the growth model, because it is applied to the logarithm of height measures, φ_{i1} represents the maximum expected log height of the plant, and $\exp(-\varphi_{i2})$ is the growth rate. The effects of irrigation conditions and genotypes (varietal differences) on the growth mechanisms were studied using model (4) on $(\varphi_{i1}, \varphi_{i2})$. In this model, μ_A and μ_B are the averages of φ_{i1} and φ_{i2}, respectively, under drought conditions, and δ_A and δ_B are the average effects of irrigation on these two quantities, respectively. As mentioned in Section 2.2.2, the genetic effects of each variety in model (4) have four

components: $u_v = (u_{v,A}, u_{v,B}, u_{v,\delta_A}, u_{v,\delta_B})^\top$, which measures the specificity of each variety with respect to the logarithm of the maximum height and growth rate under drought conditions as well as the effects of irrigation. G is the covariance matrix of genetic effect u_v.

The SAEM algorithm for parameter estimation was run for 400 iterations, of which 300 was run for burn-in with step sizes defined as $\gamma_k = (k - 300)^{-2/3}$ in the refining phase. The algorithm for computing the genetic effects was run on 1000 iterations of which 800 for burn-in and step sizes defined as $\gamma_k = (k - 800)^{-2/3}$ after burn-in. Sample convergence graphs of the SAEM algorithm for both parameter estimation and prediction are provided in Appendix S2. Both the algorithms converged satisfactorily.

We obtained the following parameter estimates:

$$\hat{\mu}_A = 3.87, \hat{\mu}_B = -5.96, \hat{\delta}_A = 0.54, \hat{\delta}_B = -0.16, \widehat{\sigma^2} = 0.01,$$

$$\hat{P} = \begin{pmatrix} 0.02 & -0.02 \\ -0.02 & 0.05 \end{pmatrix}, \hat{G} = \begin{pmatrix} 0.03 & 0.02 & 0.02 & 0.01 \\ 0.02 & 0.03 & 0.01 & 0.01 \\ 0.02 & 0.01 & 0.05 & 0.02 \\ 0.01 & 0.01 & 0.02 & 0.03 \end{pmatrix}.$$

The estimates suggest that the soybean log-growth curves reached a plateau with a value of 3.87 on average under drought conditions and 4.41 under normal irrigation conditions. The growth rate parameter φ_{i2} had an average value of -5.96 among soybean varieties under drought conditions and -6.12 under normal irrigation conditions, corresponding to slower growth under drought conditions. These results are consistent with what is observed in Fig 8.

The estimations of matrices P and G provide a quantification of the different sources of variability in growth patterns and allow for the evaluation of genetic and marginal correlations between varietal effects or between growth parameters. The genetic

correlation matrix, obtained by converting $\hat{G}$ into a correlation matrix, is estimated to

$$
\begin{pmatrix}
1.00 & 0.59 & 0.47 & 0.41 \\
0.59 & 1.00 & 0.35 & 0.33 \\
0.47 & 0.35 & 1.00 & 0.43 \\
0.41 & 0.33 & 0.43 & 1.00
\end{pmatrix}.
$$

This allowed us to assess the correlations between genetic effects in u_v. All genetic correlations in the above matrix are positive, implying that an increase in the genetic effect on the maximum height under drought conditions φ_{i1} would be associated with an increase in the genetic effect on φ_{i2}, and thus on the growth rate, and a more pronounced genetic effect on the sensitivity to irrigation variations.

Correlations between the growth patterns φ_{i1} and φ_{i2}, rather than the genetic effects, were also investigated. According to Equation (2) used in our modeling approach, the marginal covariance matrix of $\varphi_i = (\varphi_{i1}, \varphi_{i2})^\top$ is

$$
Z_i(K \otimes G)Z_i^\top + P. \tag{5}
$$

This equation includes the contribution of two parts: one that is exclusively due to genetics given by $Z_i(K \otimes G)Z_i^\top$, and the other P due to uncontrolled environmental conditions. Notably, because of the kinship matrix K in the above formula, the covariance of φ_i changes slightly from one variety to the other, in addition to its variation between the irrigation conditions because of the changes in Z_i (*i.e.*, drought and normal irrigation). Using formula (5) and the estimated values $\hat{P}$ and $\hat{G}$, by averaging over the $N_v = 188$ varieties and then converting them into correlation matrices, we found that the average genetic correlation between φ_{i1} and φ_{i2} is 0.002 under normal irrigation conditions and 0.04 under drought conditions, whereas the average marginal correlations are -0.18 and -0.25 under normal and drought conditions, respectively. This difference in sign between the genetic and marginal correlations suggests a stronger effect of the environment than that of genetics on the growth behavior of soybeans. The respective contributions of genetics and the environment to the variance of each parameter were also quantified using heritability values. The heritability of each of the mechanistic parameters of the growth curves *i.e.*,

the maximum log-height (φ_{i1}) and the growth rate (φ_{i2}) under drought conditions, denoted by h_A and h_B respectively, and the heritability of the irrigation effect on each of these parameters, denoted by h_{δ_A} and h_{δ_B}, can be deduced by calculating the following ratios

$$h_A = \frac{\widehat{G^2}_A}{\widehat{P^2}_A + \widehat{G^2}_A}, h_B = \frac{\widehat{G^2}_B}{\widehat{P^2}_B + \widehat{G^2}_B}, h_{\delta_A} = \frac{\widehat{G^2}_A}{\widehat{P^2}_A + \widehat{G^2}_{\delta_A}}, h_{\delta_B} = \frac{\widehat{G^2}_B}{\widehat{P^2}_B + \widehat{G^2}_{\delta_B}}.$$

$\widehat{P^2}_A$ and $\widehat{P^2}_B$ are the diagonal terms of $\hat{P}$, and $\widehat{G^2}_A$, $\widehat{G^2}_B$, $\widehat{G^2}_{\delta_A}$, and $\widehat{G^2}_{\delta_B}$ are 333

diagonal terms of $\hat{G}$. We obtained the following values for heritability h_A, h_B, h_{δ_A}, and 334

h_{δ_B}: 52%, 66%, 36%, and 41%, respectively. The maximum plant height and growth 335

rate under the reference conditions showed more than half of the variability explained 336

by genetics. This justifies considering the prediction of varietal effects on these 337

parameters for potential varietal selection. In contrast, the contribution of genetics to 338

the effects of irrigation on growth patterns,*i.e.*, the contribution of genetic and 339

irrigation interactions, appears to be relatively small; thus, genomic selection based on 340

these effects would be less accurate for this soybean population. Broadly speaking, the 341

predicted parameters $\widehat{\varphi}_i$ under the two conditions could be used in practice to select 342

varieties that perform better (*e.g.*, those with larger plant size or faster growth rate) 343

under drought conditions or are less sensitive to climate change. The predicted varietal 344

growth parameters are shown in Fig 9, by the water condition or by crossing the 345

normal and drought conditions. The color of the dots represents the type of variety as 346

defined in [30] ("Japan", "World" and "Primitive"). Overall, the Japanese varieties had 347

a lower growth rate than those of the "Word" and "Primitive" varieties under both 348

conditions. The "Primitive" varieties grew faster in both conditions, attaining high 349

maximum sizes among all the varieties. The "World" varieties had intermediate growth 350

rates and reach the highest heights. 351

The robustness of the predictions of genetic effects was assessed by running 5-fold 352

cross-validation four times, allowing 20 repetitions of the parameter estimation from a 353

training sample composed of 80% of the varieties, followed by the prediction of the 354

genetic effects on a test sample composed of the remaining varieties. The predictions 355

were then compared with those calculated using the full sample. For the four 356

replications of the 5-fold cross-validation, the average coefficient of correlation between 357

the cross-predictions and the genetic effects estimated from the full sample was 0.91 and the average square difference between both quantities was 0.06.

<p style="text-align:center">[Fig 9 about here.]</p>

In the present study, only 4 plots were observed for each variety. As suggested by the simulation study, the overall robustness of the analysis could be improved by increasing the number of plots per variety to provide better information on the differences between the varieties.

4 Discussion

Nonlinear mechanistic models have been used in the genomic prediction modeling of plant and animal growth [6, 12, 13, 20, 21]. In genome prediction modeling, a fusion of nonlinear functions and mixed-effects models, that is, the NLMEM framework, is required. Two approaches have been used: a two-step approach in which a nonlinear function is fitted beforehand and its parameters are modeled by a mixed-effects model [12, 13], and a one-step approach in which these are performed in a single step [6, 20, 21]. The two-step approach requires estimating the parameters of the nonlinear function for each plant and plot, sometimes resulting in overfitting. The one-step approach, in contrast, has the advantage of robust estimation because the parameters are estimated simultaneously for all plants and plots, considering genetic relationships and differences in cultivation treatments. In this study, we applied the NLMEM framework using a one-step approach and modeled the growth of numerous soybean varieties under different environmental conditions using a single model. The NLMEM framework allowed us to separate the variation in the mechanistic growth model parameters into genetic and environmental variations as well as the effect of growing conditions (irrigation or no irrigation) on these variations. Whole-genome DNA polymorphisms of soybean varieties have been decoded and classified into three groups based on the genetic similarities between varieties [30]. The genetic variation in the parameters estimated in this study clearly showed that there were differences in growth patterns among these classified groups. The "Primitive" varieties, which were considered close to the soybean varieties when they were first introduced to cultivation,

showed fast and vigorous growth, while the varieties belonging to the "Japan" group exhibited the opposite patterns. However, these patterns are average trends, and some varieties in the same group showed different patterns. For example, some of the "Japan" varieties showed fast and vigorous growth. The growth pattern of soybeans is also related to weed suppression [31], which is an important trait in breeding. Characterization of genetic variations in growth patterns using the method proposed in this study is expected to be effective in soybean breeding.

The proposed method has the following characteristics compared to previous one-step approaches [6, 20, 21]. First, the parameters of the nonlinear model, which can be regarded as latent variables, can be calculated with high speed and efficiency using an efficient algorithm called the SAEM, which is an extension of the EM algorithm. In contrast, previous methods based on the one-step approach used Markov chain Monte Carlo (MCMC) and variational Bayesian methods. The proposed inference methodology with SAEM has several advantages over competing methods. First, the SAEM algorithm is less computationally intensive and much faster than the MCMC implementations. A thorough comparison between SAEM and MCMC was performed by [32] for Bayesian variable selection in NLMEs, which highlighted the faster execution times associated with SAEM than with MCMC. In practice, the computation times vary with sample characteristics and model complexity. For the soybean data example processed with R software with a 2GHz intel core i5 processor, parameter estimation lasted 17 min and genetic effects prediction lasted 47 min, which is very promising compared with the computation times reported in the literature for similar analyses.

Another feature of the proposed method is that it considers not only the genetic correlation among parameters but also the correlation structure in the error. In the previous one-step approach-based methods, estimation was performed assuming that the errors were independent of the parameters [6, 20]. In this study, by assuming non-independence of errors among parameters. Based on the estimated genetic and environmental variance-covariance matrices of the growth model parameters, we were able to determine not only the heritability of each parameter but also the genetic and phenotypic correlations (correlations due to both genetic and environmental effects) among the parameters. Such information can be used to determine the degree of genetic control of growth and predict selection responses when selection is based on growth

patterns. Genetic correlation can also be an indicator of the difficulty of simultaneously modifying two parameters in a certain direction. By comparing phenotypic and genetic correlations, it is possible to ascertain the extent to which the correlation that appears as phenotypic is due to inheritance.

In addition to the logistic growth and asymptotic regression curves used in this study, a variety of nonlinear mechanistic models are available [12, 33]. Various covariance matrices representing the genetic similarity among varieties have also been used in addition to the additive genomic relationship matrix used in this study, including a pedigree-based relationship matrix and a nonlinear relationship matrix, such as a dominance relationship matrix and a Gaussian kernel [19]. Further, in this study, we used two levels of irrigation as part of the cultivation management treatment, but various treatments and levels are possible [34]. The proposed methodology can be straightforwardly adapted to different contexts without any major impact on the general implementation of the SAEM algorithm, by changing either the nonlinear model function $g(\cdot)$ to better reflect the curve profiles or the definition of matrices X_i and Z_i depending on the number and the potential fixed and genetic effects of the available covariates. In particular, it is very easy to consider more than two environmental conditions using the proposed approach. However, it should be highlighted that, as shown in our data analysis, a careful choice of $g(\cdot)$ is crucial to ensure good estimates and predictions. Users should also note that the performance of the modeling approach based on NLMEM depends on the structure of the collected data, as demonstrated in the simulation study.

As shown in the simulated data study, by varying the number of varieties and plots and the data collection scheme, observing enough varieties, plots per variety, and points on the growth curves is essential to ensure accurate parameter estimations and predictions of the genetic effects. While one can reasonably trust the results obtained on the soybean growth data analyzed in this paper, as suggested by the simulated results on datasets with similar structures, the data collection must be streamlined beforehand to be able to collect data of sufficient quality and ensure that the NLME approach provides reliable and robust results. Note that during the simulations, data from 2000 plots were used simultaneously to estimate parameters. This also demonstrates the scalability of the proposed method for large data-sets.

Recently, remote sensing has enabled the highly efficient measurement of plant and animal growth over time. Furthermore, the determination of genome-wide markers has become more efficient, making it possible to measure both genome-wide polymorphisms and growth for a large number of genotypes. Combining the measurements obtained from such technological advances with the methods proposed in this study could genetically model the growth patterns of plants and animals, and help to understand, control, and predict their inheritance. The method proposed in this study is scalable to larger data sets owing to its computational speed. The codes for the method are available at `https://github.com/madelattre/Scripts-soybean-paper` and can be extended according to the conditions of the application in each research project.

Supporting information

Computer codes The codes are available at

`https://github.com/madelattre/Scripts-soybean-paper`

S1 Appendix. Algorithmic details. The objective of this appendix is to provide the main technical elements for implementing the SAEM algorithm in model (1)-(2). First, we introduce the compact matrix notations:

- $\varphi = \begin{pmatrix} \varphi_1 \\ \vdots \\ \varphi_{N_p} \end{pmatrix}$: the $m.N_p$ vector obtained by concatenation of the φ_i's,

- $X = \begin{pmatrix} X_1 \\ \vdots \\ X_{N_p} \end{pmatrix}$ (resp. $Z = \begin{pmatrix} Z_1 \\ \vdots \\ Z_{N_p} \end{pmatrix}$): the $m.N_p \times p$ (resp. $m.N_p \times q.N_v$) matrix obtained by vertical concatenation of the X_i's (resp. Z_i's),

1. **Key distributions in model (1)-(2)**

   (a) **Complete data log-likelihood.**

   Using that in model (1)-(2), $y_{ij}|\varphi_i, u \sim \mathcal{N}(g(t_{ij}, \varphi_i), \sigma^2)$,

$$\varphi|u \sim \mathcal{N}(X\beta + Zu, P_{N_p}) \text{ and } u \sim \mathcal{N}(\mathbf{0}, K \otimes G), \text{ we get}$$

$$\log p(y, \varphi, u; \theta) = \log p(y \mid \varphi, u; \theta) + \log p(\varphi \mid u; \theta) + \log p(u; \theta)$$

$$= A - \frac{N_{tot}}{2} \log \sigma^2 - \frac{1}{2\sigma^2} \sum_{i=1}^{N_p} \sum_{j=1}^{n_i} (y_{ij} - g(t_{ij}, \varphi_i))^2 - \frac{N_p}{2} \log(|P|)$$

$$- \frac{1}{2}(\varphi - Zu - X\beta)'(I_{N_p} \otimes P^{-1})(\varphi - Zu - X\beta) - \frac{1}{2} \log(|K \otimes G|)$$

$$- \frac{1}{2} u'(K \otimes G)^{-1} u, \quad (6)$$

where A is a constant term which does not depend on θ, I_{N_p} is the identity

matrix of size N_p, n_i is the number of observations for plant $i \in \{1, \ldots, N_p\}$

and $N_{tot} = \sum_{i=1}^{N_p} n_i$ is the total number of observations available in the

sample (*i.e.* all observations of all plants).

(b) **Conditional distribution of the genetic effects u given φ.**

Using the Bayes formula where $p(u|\varphi; \theta) = p(\varphi|u; \theta)p(u; \theta)/p(\varphi; \theta)$ and that

in model (1)-(2), $\varphi|u \sim \mathcal{N}(X\beta + Zu, P_{N_p})$, $u \sim \mathcal{N}(\mathbf{0}, K \otimes G)$ and

$\varphi \sim \mathcal{N}(X\beta, Z(K \otimes G)Z^\top + I_{N_p} \otimes P)$, by elementary calculations, we get

that $u|\varphi \sim \mathcal{N}(m_u(\theta), \Sigma_u(\theta))$, where

$$\Sigma_u(\theta)^{-1} = Z^\top(I_{N_p} \otimes P^{-1})Z + (K \otimes G)^{-1}$$

$$m_u(\theta) = \Sigma_u(\theta)Z(I_{N_p} \otimes P^{-1})(\varphi - X\beta) \qquad (7)$$

2. **SAEM algorithm**

Algorithm 1 describes the algorithm used to estimate parameters. K is the total

number of iterations, which is tuned by the user such that the algorithm converges

at the end of K iterations. The stochastic approximation step uses model (1)-(2)

belonging to the curved exponential family, and thus consists of the computation

of the stochastic approximations of the sufficient statistics (see [26]).

Remarks:

- To better manage the acceptance proportions at the simulation step of the

  algorithm, it is possible to perform several (but a finite number of) iterations

of Metropolis-Hastings rather than one, and define $\varphi_i^{[k]}$ as the last accepted value after these iterations.

- Choosing $q_{i,\theta}(\cdot)$ as $p(\cdot|u;\theta)$ allows to simplify the expression of the acceptance ratio $\alpha_{i,k}$.

- In some situations, in particular when the number of observed plants per variety is small, the convergence of the algorithm can be improved by using M Monte-Carlo Markov chains instead of only one as described in Algorithm 1. This amounts to simulating M values $\tilde{\varphi}_i^{[k,m]}$, $m \in 1,\ldots,M$, at each iteration and replacing the sufficient statistics by their means over the M simulated values.

Algorithm 2 presents algorithm for genetic effects prediction. This should be used with $\theta = \hat{\theta}^{[K]}$ obtained at the convergence of the first algorithm. In general, Algorithm 2 is written for any value of θ. It also exploits the fact that the model (1)-(2) belongs to the curved exponential family.

S2 Appendix. Convergence graphs.

[Fig 10 about here.]

[Fig 11 about here.]

[Fig 12 about here.]

[Fig 13 about here.]

Acknowledgments

This study was supported by JST CREST (grant no. JPMJCR16O2) and JSPS KAKENHI (grant no. JP22H02306). The funders had no role in the study design, data collection and analysis, decision to publish, or manuscript preparation.
We are grateful to the technical staff of the Arid Land Research Center, Tottori University, and Izumi Higashida for the management of the field experiments. We would like to thank Yoshihiro Ohmori, Yuji Yamasaki, Hirokazu Takahashi, Hideki Takanashi, Mai Tsuda, Yuji Sawada, Hisashi Tsujimoto, Akito Kaga, Mikio Nakazono,

---

Algorithm 1 SAEM algorithm for parameter estimation

Input: $(\gamma_k)_{k \geqslant 0}$

Initialization: $\theta^{[0]}$, $u^{[0]}$, $\varphi^{[0]}$

for $k \in \{1, \dots, K\}$ **do**

    1. **S-step**

    for $i \in \{1, \dots, N_p\}$ **do**

        Draw $\tilde{\varphi}_i^{[k]}$ from a proposal kernel $q_{i,\theta^{[k-1]}}(\cdot)$ and set $\varphi_i^{[k]} = \tilde{\varphi}_i^{[k]}$ with probability

$$
\begin{aligned}
\alpha_{i,k} &= \min\left(1, \frac{p(\tilde{\varphi}_i^{[k]}|u^{[k-1]}, y_i; \theta^{[k-1]}) q_{i,\theta^{[k-1]}}(\varphi_i^{[k-1]})}{p(\varphi_i^{[k-1]}|u^{[k-1]}, y_i; \theta^{[k-1]}) q_{i,\theta^{[k-1]}}(\tilde{\varphi}_i^{[k]})}\right) \\
&= \min\left(1, \frac{p(y_i|\tilde{\varphi}_i^{[k]}; \theta^{[k-1]}) p(\tilde{\varphi}_i^{[k]}|u^{[k-1]}; \theta^{[k-1]}) q_{i,\theta^{[k-1]}}(\varphi_i^{[k-1]})}{p(y_i|\varphi_i^{[k-1]}; \theta^{[k-1]}) p(\varphi_i^{[k-1]}|u^{[k-1]}; \theta^{[k-1]}) q_{i,\theta^{[k-1]}}(\tilde{\varphi}_i^{[k]})}\right)
\end{aligned}
$$

        and $\varphi_i^{[k]} = \varphi_i^{[k-1]}$ with probability $1 - \alpha_{i,k}$.

    end for

    Draw $u^{[k]}$ under a Gaussian distribution with mean $m_u(\theta^{[k-1]})$ and covariance matrix $\Sigma_u(\theta^{[k-1]})$ whose formula are given in (7).

    2. **SA-step**

    Compute stochastic approximations of the sufficient statistics

$$
\begin{aligned}
s_1^{[k]} &= (1 - \gamma_k) s_1^{[k-1]} + \gamma_k (\varphi^{[k]} - Z u^{[k]}) \\
s_2^{[k]} &= (1 - \gamma_k) s_2^{[k-1]} + \gamma_k (u^{[k]})^\top K^{-1} u^{[k]} \\
s_3^{[k]} &= (1 - \gamma_k) s_3^{[k-1]} + \gamma_k (\varphi^{[k]} - Z u^{[k]})^\top (\varphi^{[k]} - Z u^{[k]}) \\
s_4^{[k]} &= (1 - \gamma_k) s_4^{[k-1]} + \gamma_k \sum_{i=1}^{N_p} \sum_{j=1}^{n_i} (y_{ij} - g(t_{ij}, \varphi_i^{[k]}))^2
\end{aligned}
$$

    3. **M-step**

    Update the parameter estimates according to

$$
\begin{aligned}
\hat{\beta}^{[k]} &= (X^\top (I_{N_p} \otimes P^{-1}) X)^{-1} X^\top (I_{N_p} \otimes P^{-1}) s_1^{[k]} \\
\hat{P}^{[k]} &= \frac{1}{N_p} \left(s_3^{[k]} - 2(X\hat{\beta}^{[k]})^\top s_1^{[k]} + (X\hat{\beta}^{[k]})^\top X\hat{\beta}^{[k]} \right) \\
\hat{G}^{[k]} &= \frac{s_2^{[k]}}{N_v} \\
\widehat{\sigma^2}^{[k]} &= \frac{s_4^{[k]}}{N_{tot}}
\end{aligned}
$$

Algorithm 2 SAEM algorithm for genetic effects prediction

Input: $(\gamma_k)_{k \geqslant 0}$, $\theta = (\beta, P, G, \sigma^2)$

Initialization: $u^{[0]}$, $\varphi^{[0]}$

for $k \in \{1, \ldots, K\}$ **do**

  1. S-step

  for $i \in \{1, \ldots, N_p\}$ **do**

    Draw $\tilde{\varphi}_i^{[k]}$ from a proposal kernel $q_{i,\theta}(\cdot)$ and set $\varphi_i^{[k]} = \tilde{\varphi}_i^{[k]}$ with probability

$$
\begin{aligned}
\alpha_{i,k} &= \min\left(1, \frac{p(\tilde{\varphi}_i^{[k]}|u^{[k-1]}, y_i; \theta)q_{i,\theta}(\varphi_i^{[k-1]})}{p(\varphi_i^{[k-1]}|u^{[k-1]}, y_i; \theta)q_{i,\theta}(\tilde{\varphi}_i^{[k]})}\right) \\
&= \min\left(1, \frac{p(y_i|\tilde{\varphi}_i^{[k]}; \theta)p(\tilde{\varphi}_i^{[k]}|u^{[k-1]}; \theta)q_{i,\theta}(\varphi_i^{[k-1]})}{p(y_i|\varphi_i^{[k-1]}; \theta)p(\varphi_i^{[k-1]}|u^{[k-1]}; \theta)q_{i,\theta}(\tilde{\varphi}_i^{[k]})}\right)
\end{aligned}
$$

    and $\varphi_i^{[k]} = \varphi_i^{[k-1]}$ with probability $1 - \alpha_{i,k}$.

  end for

  2. SA-step

  Compute stochastic approximations of the sufficient statistics

$$
s^{[k]} = (1 - \gamma_k)s_1^{[k-1]} + \gamma_k \varphi^{[k]}
$$

  3. M-step

  Update the genetic values according to

$$
\hat{u}^{[k]} = \left(Z^\top(I_{N_p} \otimes P^{-1})Z + (K \otimes G)^{-1}\right)^{-1} Z^\top(I_{N_p} \otimes P^{-1})(s^{[k]} - X\beta)
$$

end for

Output: $\hat{u}^{[K]}$.

Toru Fujiwara for being involved in the field experiments, Hiromi Kajiya-Kanegae and Kosuke Hamazaki for curating whole-genome marker data, and Sawako Maruyama for curating the phenotype data.

References

1. Soltani A. Modeling physiology of crop development, growth and yield. CABi; 2012.

2. Yang G, Liu J, Zhao C, Li Z, Huang Y, Yu H, et al. Unmanned Aerial Vehicle Remote Sensing for Field-Based Crop Phenotyping: Current Status and Perspectives. Frontiers in Plant Science. 2017;8:1111. doi:10.3389/fpls.2017.01111.

3. White JW, Andrade-Sanchez P, Gore MA, Bronson KF, Coffelt TA, Conley MM, et al. Field-based phenomics for plant genetics research. Field Crops Research. 2012;133:101–112. doi:10.1016/j.fcr.2012.04.003.

4. Andrade-Sanchez P, Gore MA, Heun JT, Thorp KR, Carmo-Silva AE, French AN, et al. Development and evaluation of a field-based high-throughput phenotyping platform. Functional Plant Biology. 2013;41(1):68–79. doi:10.1071/fp13126.

5. Campbell M, Walia H, Morota G. Utilizing random regression models for genomic prediction of a longitudinal trait derived from high-throughput phenotyping. Plant Direct. 2018;2(9):e00080. doi:10.1002/pld3.80.

6. Campbell MT, Grondin A, Walia H, Morota G. Leveraging genome-enabled growth models to study shoot growth responses to water deficit in rice. Journal of experimental botany. 2020;71(18):5669–5679.

7. Toda Y, Kaga A, Kajiya-Kanegae H, Hattori T, Yamaoka S, Okamoto M, et al. Genomic prediction modeling of soybean biomass using UAV-based remote sensing and longitudinal model parameters. The Plant Genome. 2021; p. e20157. doi:10.1002/tpg2.20157.

8. Nelder J. The fitting of a generalization of the logistic curve. Biometrics. 1961;17(1):89–110.

9. Winsor CP. The Gompertz curve as a growth curve. Proceedings of the national academy of sciences. 1932;18(1):1–8.

10. Crispim AC, Kelly MJ, Guimaraes SEF, Silva FFe, Fortes MRS, Wenceslau RR, et al. Multi-Trait GWAS and New Candidate Genes Annotation for Growth Curve Parameters in Brahman Cattle. PLOS ONE. 2015;10(10):e0139906. doi:10.1371/journal.pone.0139906.

11. Baker R, Leong W, Welch S, Weinig C. Mapping and predicting non-linear Brassica rapa growth phenotypes based on Bayesian and frequentist complex trait estimation. G3: Genes, Genomes, Genetics. 2018;8(4):1247–1258.

12. Yin T, König S. Genomic predictions of growth curves in Holstein dairy cattle based on parameter estimates from nonlinear models combined with different kernel functions. Journal of dairy science. 2020;103(8):7222–7237.

13. Toda Y, Sasaki G, Ohmori Y, Yamasaki Y, Takahashi H, Takanashi H, et al. Genomic prediction of green fraction dynamics in soybean using unmanned aerial vehicles observations. Frontiers in Plant Science. 2022;13.

14. Meuwissen TH, Hayes BJ, Goddard ME. Prediction of total genetic value using genome-wide dense marker maps. Genetics. 2001;157(4):1819–29.

15. Onogi A. Connecting mathematical models to genomes: Joint estimation of model parameters and genome-wide marker effects on these parameters. Bioinformatics. 2020;36(10):3169–3176. doi:10.1093/bioinformatics/btaa129.

16. Ma CX, Casella G, Wu R. Functional Mapping of Quantitative Trait Loci Underlying the Character Process: A Theoretical Framework. Genetics. 2002;161(4):1751–1762. doi:10.1093/genetics/161.4.1751.

17. Wu R, Ma CX, Lin M, Casella G. A General Framework for Analyzing the Genetic Architecture of Developmental Characteristics. Genetics. 2004;166(3):1541–1551. doi:10.1534/genetics.166.3.1541.

18. Gianola D. Priors in Whole-Genome Regression: The Bayesian Alphabet Returns. Genetics. 2013;194(3):573–596. doi:10.1534/genetics.113.151753.

19. Morota G, Gianola D. Kernel-based whole-genome prediction of complex traits: a review. Frontiers in Genetics. 2014;5:363. doi:10.3389/fgene.2014.00363.

20. Onogi A, Ogino A, Sato A, Kurogi K, Yasumori T, Togashi K. Development of a structural growth curve model that considers the causal effect of initial phenotypes. Genetics Selection Evolution. 2019;51(1):19.

21. Yu H, van Milgen J, Knol E, Fernando R, Dekkers J. A Bayesian hierarchical model to integrate a mechanistic growth model in genomic prediction. In: 12. World congress on genetics applied to livestock production (WCGALP); 2022.

22. Pinheiro J C , Bates D M . Mixed-Effects Models in S and S-PLUS. 1st ed. Statistics and Computing. Springer-Verlag New York Inc.; 2000.

23. Lavielle M . Mixed Effects Models for the Population Approach. Models, Tasks, Methods and Tools. Chapman and Hall/CRC Biostatistics Series. Chapman and Hall/CRC; 2014.

24. Jaffrézic F, Meza C, Lavielle M, Foulley JL. Genetic analysis of growth curves using the SAEM algorithm. Genetics Selection Evolution. 2006;38(6):583–600.

25. Dempster A P , Laird N M , Rubin D B . Maximum Likelihood from Incomplete Data via the EM Algorithm. Journal of Royal Statistical Society Series B. 1977;39(1):1–38.

26. Delyon B , Lavielle M , Moulines E . Convergence of a stochastic approximation version of the EM algorithm. The Annals of Statistics. 1999;27(1):94–128.

27. Kaga A, Shimizu T, Watanabe S, Tsubokura Y, Katayose Y, Harada K, et al. Evaluation of soybean germplasm conserved in NIAS genebank and development of mini core collections. Breeding science. 2012;61(5):566–592.

28. Kuhn E , Lavielle M . Coupling a stochastic approximation version of EM with an MCMC procedure. ESAIM:PS. 2004;8:115–131.

29. Robert CP, Casella G, Casella G. Monte Carlo statistical methods. vol. 2. Springer; 1999.

30. Kajiya-Kanegae H, Nagasaki H, Kaga A, Hirano K, Ogiso-Tanaka E, Matsuoka M, et al. Whole-genome sequence diversity and association analysis of 198 soybean accessions in mini-core collections. DNA Research. 2021;28(1).

31. Kurokawa S, Kaga A, Tsuda M, Sekine D, Shibuya T. Evaluation of validity and limitations of the soybean canopy height-to-row spacing ratio as an onsite index to control weeds using diverse soybean accessions. Weed Biology and Management. 2019;19(3):103–110.

32. Naveau M, King GKK, Rincent R, Sansonnet L, Delattre M. Bayesian high-dimensional covariate selection in non-linear mixed-effects models using the SAEM algorithm; 2022.

33. Hossein-Zadeh NG. Modelling growth curve in Moghani sheep: comparison of non-linear mixed growth models and estimation of genetic relationship between growth curve parameters. The Journal of Agricultural Science. 2017;155(7):1150–1159.

34. Sakurai K, Toda Y, Kajiya-Kanegae H, Ohmori Y, Yamasaki Y, Takahashi H, et al. Time-series multispectral imaging in soybean for improving biomass and genomic prediction accuracy. The Plant Genome. 2022;15(4):e20244.

List of Figures

Fig 1. Effect of the number of observed varieties and of the number of plants per variety on the estimation of the fixed-effects parameters and of the residual variance

Fig 2. Effect of the number of observed varieties and of the number of plants per variety on the estimation of the covariance matrix P

Fig 3. Effect of the number of observed varieties and of the number of plants per variety on the estimation of the covariance matrix G

Fig 4. Effect of the sampling frequency along the curves on the estimation of the fixed-effects parameters and of the residual variance

Fig 5. Effect of the sampling frequency along the curves on the estimation of the covariance matrix P

Fig 6. Effect of the sampling frequency along the curves on the estimation of the covariance matrix G

Fig 7. 2017 Tottori UAV data by water condition. C: normal water condition, D: dry condition.

Fig 8. 2017 Tottori UAV data by water condition. C: normal water condition, D: dry condition.

Fig 9. Varietal predicted parameters under both water conditions per groups of varieties.

Fig 10. Real data analysis: Parameter estimation. Evolution of β's and σ^2's estimations over iterations of the SAEM algorithm.

Fig 11. Real data analysis: Parameter estimation. Evolution of P's estimation over iterations of the SAEM algorithm.

Fig 12. Real data analysis: Parameter estimation. Evolution of the estimation of the diagonal terms of G over iterations of the SAEM algorithm.

Fig 13. Real data analysis: Parameter estimation. Evolution of the estimation of the extra-diagonal terms of G over iterations of the SAEM algorithm.

List of Tables

Table 1. Mean square differences between the simulated genetic effects and their predictions.

	N_v	N_p	t	θ known	θ estimated
Scenario 1	50	2000	Rich	0.08	0.12
	100	2000	Rich	0.15	0.17
Scenario 2	200	800	Rich	0.60	0.61
	200	800	Sparse	0.76	0.78