CD22L Conjugation to Insulin Attenuates Insulin-Specific B cell Activation

Kyle Apley*: Department of Pharmaceutical Chemistry, University of Kansas, Lawrence, Kansas 66047, United States; orcid.org/0000-0002-7689-2215

Amber Griffith*: Department of Medicine, Division of Allergy and Immunology, Washington University School of Medicine, St. Louis, Missouri, 63110, United States; orcid.org/0000-0003-3510-9797

Grant Downes: Bioengineering Program, University of Kansas, Lawrence, Kansas 66045, United States; orcid.org/0000-0001-9063-7678

Patrick Ross: Department of Medicinal Chemistry, University of Kansas, Lawrence, Kansas 66047, United States; orcid.org/0000-0003-1264-2273

Mark P. Farrell: Department of Medicinal Chemistry, University of Kansas, Lawrence, Kansas 66047, United States; orcid.org/0000-0002-6866-428X

Peggy Kendall: Department of Medicine, Division of Allergy and Immunology, Washington University School of Medicine, St. Louis, Missouri, 63110, United States; orcid.org/0000-0002-0516-654X

Cory Berkland: Department of Pharmaceutical Chemistry, University of Kansas, Lawrence, Kansas 66047, United States; Department of Chemical and Petroleum Engineering, University of Kansas, Lawrence, Kansas 66045, United States; Bioengineering Program, University of Kansas, Lawrence, Kansas 66045; Orcid.org/0000-0002-9346-938X

*authors contributed equally

Abstract

Pancreatic islet-reactive B lymphocytes promote Type 1 diabetes (T1D) by presenting antigen to islet-destructive T cells. Teplizumab, an anti-CD3 monoclonal, delays T1D onset in patients at risk, but additional therapies are needed to prevent disease entirely. Therefore, bifunctional molecules were designed to selectively inhibit T1D-promoting anti-insulin B cells by conjugating a ligand for the B cell inhibitory receptor CD22 (i.e., CD22L) to insulin, which permit these molecules to concomitantly bind to anti-insulin B cell receptors (BCRs) and CD22. Two prototypes were synthesized: 2:2 insulin-CD22L conjugate on a 4-arm PEG backbone, and 1:1 insulin-CD22L direct conjugate. Transgenic mice ($125Tg^{SD}$) expressing anti-insulin BCRs provided cells for *in vitro* testing. Cells were cultured with constructs for three days then assessed by flow cytometry. Duplicate wells with anti-CD40 simulated T cell help. Surprisingly, a 2-insulin 4-arm PEG control caused robust proliferation and activation-induced CD86 upregulation. Anti-CD40 further boosted these effects. This was unexpected, as soluble insulin alone has no effect, and may indicate that BCR-crosslinking occurs when antigens are tethered by the PEG backbone. Addition of CD22L via the 2:2 insulin-CD22L conjugate restored B cell properties to that of controls, without additional beneficial effect. In contrast, the 1:1 insulin-CD22L direct conjugate significantly reduced anti-insulin B cell proliferation, CD86 upregulation, and cell number, even in the presence of anti-CD40. CD22L alone had no effect, and the constructs did not affect WT B cells. Thus, high valency constructs activate anti-insulin B cells, while low-valency antigen-CD22L conjugates co-ligate BCR and CD22, reducing pathogenic B cell numbers and function without harming normal cells. Thus, insulin-CD22L direct conjugate is a promising candidate for preclinical trials to prevent T1D without inducing immunodeficiency.

Introduction

Autoreactive B cells contribute to aberrant immune responses in autoimmune disease through autoantibody production, autoantigen presentation to T cells, and inflammatory cytokine production.[1, 2] One therapeutic strategy to restore B cell tolerance in autoimmunity is to block B cell activation by co-localizing the B cell receptor (BCR) with intracellular immunoreceptor tyrosine-based inhibitory motif (ITIM)-bearing receptors using antigen-ligand or anti-BCR antibody-ligand conjugates.[3-15] ITIM-bearing receptors expressed on B cells investigated in this approach include the siglecs CD33 and CD22 and the antibody-Fc receptor FcγRIIB. Notably, targeting of CD22 may contribute to selective B cell immunomodulation due to its sole expression on B cells.[16, 17] Co-localization of CD22 with the BCR complex results in phosphorylation of the CD22 ITIM by the Src-family protein tyrosine kinase LYN followed by activation of the protein tyrosine phosphatase SHP-1.[18, 19] SHP-1 then negatively regulates B cell activation by suppressing downstream effectors including MAP kinases and phospholipase C responsible for intracellular calcium release.[20]

Prior tolerization studies with BCR – CD22 co-localization focused on disease models in which antibodies are key effectors of disease including rheumatoid arthritis and alloantibody-mediated hemophilia, and these studies primarily reported *in vivo* antibody titer data.[5-10] In type 1 diabetes (T1D), however, autoantibodies do not have such a clear role in disease although their presence indicates risk of disease development. On the other hand, effector anti-islet antigen B cells are indeed pathogenic, propagating autoimmunity via autoantigen presentation and inflammatory cytokine excretion.[21-31] In addition, these prior B cell tolerization studies relied on liposome and large polymer scaffolds bearing hundreds of copies of antigen and CD22 ligand

(CD22L), and it is unknown if low valency antigen-CD22L conjugates have the potential to tolerize B cells in a similar manner.

Currently, there is little data on antigen-CD22L conjugates' ability to block B-cell activation in relation to T1D. For these studies, we developed therapeutic bifunctional molecules designed to target anti-insulin B cells that promote T1D. These bifunctional molecules use insulin autoantigen to bind the BCR and are conjugated with murine CD22L (mCD22L), developed by Paulson and co-workers to bind murine CD22 with micromolar affinity.[32] Transgenic anti-insulin (125Tg) B cells were used to study effects of discrete, low valency (1:1, 2:2) insulin/mCD22L conjugates. The 1:1 insulin-mCD22L conjugate was synthesized by direct conjugation of an azide-functionalized mCD22L to insulin(B29Lys)-alkyne, and the 2:2 insulin-mCD22L conjugate utilized a 20 kDa 4-arm PEG scaffold. Following characterization, the conjugates were incubated with 125Tg splenocytes with and without anti-CD40 to investigate the effect of the conjugates on B cell survival, proliferation and CD86 expression. The data demonstrate the ability of CD22L and antigen conjugation to attenuate anti-insulin B cell activation *in vitro.*

Methods

Insulin-azide, insulin-alkyne, and mCD22L-azide synthesis

Insulin-azide and insulin-alkyne were prepared from recombinant human insulin (EMD Millipore) by selective acylation of the B29Lys residue as previously reported.[33, 34] The synthesis of mCD22L-azide is detailed in the supporting information.

4-arm PEG-insulin synthesis

To a solution of 40 mg of 4-arm PEG-alkyne 20kDa (MW = 20000, 2 µmoles, Advanced Biochemicals) in 3.4 mL of 50 mM phosphate pH 7.4 buffer, 24.3 mg of insulin-azide (MW = 6080.92, 4 µmoles) in 1 mL of DMSO was added followed by 1 mg of CuSO₄ •7H₂O (MW = 249.69, 4 µmoles) and 8.7 mg of THPTA (MW = 434.50, 20 µmoles) dissolved in 400 µL of water. Lastly, 15.85 mg of sodium ascorbate (MW = 198.11, 80 µmoles) dissolved in 200 µL of water was added to the reaction. The reaction was stirred at RT for 1 hour. The resulting 4-arm PEG-insulin products with 1, 2, 3, or 4 insulin per PEG were isolated by preparative reverse-phase HPLC (Prep-LC) using a Waters XBridge Prep C18 5 µm OBD 19 x 250 mm column at a flow rate of 14 ml/min with a 10-minute gradient from 36-42%B (solvent A: water with 0.05% trifluoroacetic acid (TFA), solvent B: acetonitrile with 0.05% TFA). Desired fractions were concentrated by rotary evaporation under reduced pressure, frozen, then lyophilized to give a white powder. Yields of 7.1, 10.1, 11.7, and 8.2 mgs were attained. Product identity was confirmed by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry using a HABA matrix. Expected: 26,000, 32,000, 38,000, 44,000 Da; found: 26,081, 32,332, 38,096, 44,130 m/z [M+H]$^+$.

4-arm PEG-insulin(2)/mCD22L(2) synthesis

To a solution of 4.3 mg of 4-arm PEG-insulin(2) (MW = 32,000, 0.134 µmoles) dissolved in 1 mL of 50 mM phosphate pH 7.4 buffer with 250 µL of DMSO, 0.54 mg of mCD22L-azide (MW = 999.97, 0.54 µmoles) in 81 µL of buffer was added followed by 67 µg of CuSO4 •7H2O (MW = 249.69, 0.268 µmoles) and 587 µg of THPTA (MW = 434.50, 1.34 µmoles) dissolved in 27 µL of water. Lastly, 1.07 mg of sodium ascorbate (MW = 198.11, 5.36 µmoles) dissolved in 13.5 µL of water was added to the reaction. The reaction was stirred at RT for 2 hours. The

desired product was isolated by Prep-LC using a Waters XBridge Prep C18 5 μm OBD 19 x 250

mm column at a flow rate of 14 ml/min with a 10-minute gradient from 35-45%B (solvent A:

water with 0.05% trifluoroacetic acid (TFA), solvent B: acetonitrile with 0.05% TFA). Desired

fractions were concentrated by rotary evaporation under reduced pressure, frozen, then

lyophilized to give a white powder. Yield: 2.45 mgs (55%). Product identity was confirmed by

matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry

using a DHAP matrix. Expected: 34,000 Da. Found: 34,140 m/z [M+H]+.

Insulin-mCD22L synthesis

To a solution of 4 mg of insulin-alkyne (MW = 5916.77, 0.68 μmoles) dissolved in 340

μL of DMSO and 1.2 mL of 50 mM phosphate pH 7.4 buffer, 0.81 mg of mCD22L-azide (MW =

999.97, 0.81 μmoles) in 81 μL of buffer was added followed by 170 μg of CuSO4 •7H2O (MW

= 249.69, 0.68 μmoles) and 1.47 mg of THPTA (MW = 434.50, 3.4 μmoles) dissolved in 68 μL

of water. Lastly, 2.68 mg of sodium ascorbate (MW = 198.11, 13.6 μmoles) dissolved in 34 μL

of water was added to the reaction. The reaction was stirred at RT for 1 hour. The desired

product was isolated by Prep-LC using a Waters XBridge Prep C18 5 μm OBD 19 x 250 mm

column at a flow rate of 14 ml/min with a 10-minute gradient from 20-40%B (solvent A: water

with 0.05% trifluoroacetic acid (TFA), solvent B: acetonitrile with 0.05% TFA). Desired

fractions were concentrated by rotary evaporation under reduced pressure, frozen, then

lyophilized to give a white powder. Yield: 3.54 mgs (76%). Product identity was confirmed by

matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry

using a DHAP matrix. Expected: 6917.7 Da. Found: 6918.06 m/z [M+H]+.

RP-HPLC Analysis

Lyophilized Prep-LC fractions dissolved in DMSO/water at 0.5 mg/ml were injected onto a Waters XBridge BEH C18 2.5 μm 130 Å 4.6 x 150 mm column equilibrated at 50 °C using a 10 minute 30-50%B gradient (solvent A: water with 0.05% trifluoroacetic acid (TFA), solvent B: acetonitrile with 0.05% TFA). Detection was performed at 214 and 280 nm.

B Cell Proliferation and Flow Cytometry

Splenocytes from 125Tg B6 and WT B6 control mice were stained with CellTrace Violet and cultured for 3 days[35] in complete RPMI alone or stimulated with 5 ug/mL anti-mouse anti-CD40 (BD Pharmingen clone HM40-3) in the presence or absence of conjugates described above at 150 nM concentrations. The cells were then analyzed by flow cytometry using fluorochrome-conjugated antibodies against B220 (RA3-6B2, BUV395), CD86 (GL-1 PE), and CD22 (Cy34.1 FITC) from BD Biosciences. LiveDead Blue, CellTrace Violet and were obtained from Invitrogen. APC conjugated IgMa was obtained from Biolegend. Samples were collected on an Aurora Flow Cytotmeter (Cytek Biosciences), and data was analyzed using Flowjo software (Tree Star) and GraphPad Prism.

Results

Insulin-mCD22L Conjugate Synthesis

Insulin-mCD22L was prepared by copper-catalyzed azide-alkyne cycloaddition (CuAAC) involving insulin-alkyne and mCD22L-azide (**Figure 1**). The desired product was isolated by preparatory RP-HPLC and product identity was confirmed by MALDI-TOF mass

Figure 1: Synthesis of insulin-mCD22L; a direct conjugate between insulin-alkyne and mCD22L-azide. A) Reaction scheme. B) HPLC analysis of reactants and reaction products. C) MALDI-TOF-MS analysis of insulin-alkyne and isolated insulin-mCD22L.

Figure 2: Reaction scheme of 4-arm PEG-insulin(2)/mCD22L(2). First, insulin-azide is conjugated with 4-arm PEG-alkyne by copper-catalyzed azide-alkyne cycloaddition (CuAAC) to yield a mixture of 4-arm PEG-insulin(1-4) products, including 4-arm PEG-insulin(2). Second, isolated 4-arm PEG-insulin(2) is conjugated to mCD22L-azide by CuAAC to yield 4-arm PEG-insulin(2)/mCD22L(2).

spectrometry. The 4-arm PEG-insulin(2)/mCD22L(2) was prepared in a two-step process

(**Figure 2**). First, insulin-azide was conjugated to 4-arm PEG-alkyne by CuAAC yielding a

mixture of 4-arm PEG-insulin products with 1 to 4 insulin molecules attached per PEG backbone

(**Figure 3A, B**). The individual species were isolated using a preparatory reverse-phase HPLC

Figure 3: Analysis of 4-arm PEG-insulin products. A) Representation and molecular weights (MW) of the reaction products formed in the CuAAC reaction between 4-arm PEG-alkyne and insulin-azide. B) HPLC analysis of four product fractions following preparatory RP-HPLC purification of 4-arm PEG-insulin. The number in parenthesis indicates the number of insulin molecules conjugated to the 4-arm PEG. C) MALDI-MS analysis of four product fractions following preparatory RP-HPLC purification of 4-arm PEG-insulin. The mode m/z ratio of the major peak is indicated.

(Prep-LC) method with a shallow gradient. Each fraction was analyzed by HPLC, and the four major species were resolved by using a 2.5 µm C18 column at 50 °C. The purity of each species was greater than 90% following Prep-LC purification. The individual 4-arm PEG-insulin species were identified by MALDI mass spectrometry (**Figure 3C**). Due to the inherent MW distribution of the 4-arm PEG-alkyne (20 kDa) backbone, the 4-arm PEG-insulin species also exist with a similar distribution of MWs. Even so, the mode molecular weight for each 4-arm PEG-insulin species closely matched the theoretical MWs (**Figure 3A**).

Second, the 4-arm PEG-insulin(2) was reacted with mCD22L-azide by CuAAC to prepare a 4-arm PEG-insulin(2)/mCD22L(2) conjugate (**Figure 2**). Comparison of HPLC traces for pre- and post-reaction samples show a distinct shift in retention time for the product

Figure 4: Analysis of 4-arm PEG-insulin/mCD22L products. A) HPLC analysis of the CuAAC reaction between 4-arm PEG-insulin(2) with mCD22L-azide before and 2 hours after the addition of sodium ascorbate. Two equivalents of mCD22L-azide per alkyne were used. B) MALDI-MS analysis of the 4-arm PEG-insulin(2)/mCD22L(2) product following purification by RP-HPLC. The mode m/z ratio of the major peak is indicated. A representation and molecular weights (MW) of the starting material and product are shown. C) Proton NMR analysis of 4-arm PEG-insulin(2)/mCD22L(2) and precursors. The aromatic region of the spectra is shown due to the distinctive peaks from both the aromatic sidechains of insulin-azide and the bisphenyl group on mCD22L-azide.

(**Figure 4A**). MALDI mass spectrometry analysis of the product identified a 2 kDa increase in

mode MW, matching the expected 1 kDa increase per mCD22L-azide conjugated (**Figure 4B**).

Proton NMR characterization was performed for 4-arm Peg-Insulin(2)/mCD22L(2) (**Figure 4C**).

Focusing on the aromatic region, peaks from the aromatic amino acids of insulin (6.5-7.5 ppm)

are present along with peaks from the bisphenyl-aryl group on the mCD22L (7.5-7.8 ppm).

Additionally, a peak corresponding to the sole proton of the 1,2,3-N triazole formed by CuAAC

is present (8.0 ppm). Integration of the mCD22L aromatic peak area and comparison to the 1,2,3-

N triazole proton peak and PEG-backbone proton peak indicate the presence of two mCD22Ls

per backbone (**Table S1**). The broadening of the insulin aromatic amino acid peaks made quantitative assessment of the insulins per backbone infeasible for the 4-arm PEG-insulin(2)/mCD22L(2) species, but integration of the insulin aromatic amino acid peak area in the 4-arm PEG-insulin(2) species and comparison to the 1,2,3-N triazole proton peak and PEG-backbone proton peak indicate the presence of two insulins per backbone, along with the MALDI data (**Figure 4C**). Together, the HPLC, MS, and NMR data confirm the 4-arm PEG-insulin(2)/mCD22L(2) product to be a discrete entity with negligible presence of products possessing undesired quantities of insulin or mCD22L molecules.

Effect of Insulin-mCD22L Conjugates on B cells

In 125Tg mice, the transgene is expressed by over 95% of B cells, providing a large population of anti-insulin B cells for study.[30] Splenocytes from 125Tg B6 mice were pre-labeled with the proliferation dye Cell Trace Violet then incubated with insulin constructs at 150 nM concentration.[35] Following 72 hours of incubation, flow cytometry was used to measure construct effects on B cell survival, proliferation and activation. Surprisingly, the 4-arm PEG-insulin(2) control construct had a strong activating effect on anti-insulin B cells (**Figure 5A**), resulting in a significantly higher proportion remaining in wells compared to those in control wells (11.99% ± 8.24% SD; media only 3.45% ± 2.53% SD; mCD22L 3.99% ± 2.34% SD). Addition of mCD22L to the 4-arm PEG modulated this effect, holding B cell proportions at control levels (4-arm PEG-insulin(2)/CD22L(2) 3.40% ±3.05% SD). The insulin-CD22L construct had no effect on B cell proportions (3.11% ± 2.31% SD).

B cells incubated with the 4-arm PEG-insulin (2) construct also had significantly higher proliferation as measured by dye dilution, higher total B cell numbers, higher upregulation of the

Figure 5: Anti-insulin B cell responses to the incubation of insulin and mCD22L conjugates in 125Tg B6 mouse splenocytes after 72 hours. A) The percentage of total live cells that are B cells. B) Total live B cell number. C) CD22 expression levels on B cells as a ratio over untreated anti-insulin B cell CD22 expression. D) Number of proliferating B cells determined by dye-dilution assay. E) CD86 expression levels on B cells as a ratio over untreated anti-insulin B cell CD86 expression. n≥ 6 mice per condition, standard deviation is indicated by error bars. *p<0.05 **p<0.01 ***p<0.001 ****P<0.0001 by one-way Anova with Tukey's multiple comparison.

costimulatory antigen-presenting molecule CD86, and down regulation of the inhibitory

molecule CD22 (**Figure 5B-E, Table 1**). These findings suggest that 4-arm PEG-insulin(2) has

stimulatory effects on anti-insulin B cells (**Figure 5B-E, Table 1, Figure S2**), likely due to

crosslinking and strong signaling via the BCR. The addition of mCD22L to the PEG backbone in

4-arm PEG-insulin(2)/CD22L(2) antagonizes the stimulatory effects of 4-arm PEG-insulin(2),

restoring proliferation, total numbers of B cells and levels of CD86 and CD22 expression back to

levels comparable to control wells with media only (**Figure 5B-E, Table 1**). Insulin-CD22L did

not have any effects on B cell proportion, total number, number of B cells proliferated, or

expression of CD86 and CD22 (**Figure 5, Table 1**). WT B cells were not affected by any of the

constructs, indicating that the stimulatory effect of the 4-arm PEG-insulin(2) is antigen-specific.

(**Figure S3**).

Table 1: Average B cell proportion, number and MFI for anti-insulin B cells incubated with insulin constructs

Condition	Average Total Live B cells	SD	Average B Cell Proportion	SD	Average Number of B Cells Proliferated	SD	Average CD86 MFI Ratio	SD	Average CD22 MFI Ratio	SD
Media Only	2304	2130	3.44	2.53	204	145	0.98	0.38	1.07	0.39
mCD22L-azide	2397	1476	3.99	2.34	144	43	0.98	1.29	0.89	0.22
4-arm PEG-Insulin(2)	13105	7393	11.99	8.24	5458	3008	44.29	16.95	0.51	0.06
4-arm PEG-Insulin(2)/mCD22L(2)	2972	2819	3.4	3.05	172	216	6.56	6.34	0.7	0.42
Insulin-mCD22L	2997	2614	3.11	2.31	206	243	0.94	0.54	0.72	0.16

Effect of Insulin-mCD22L Conjugates on Anti-insulin B cell activation in Response to Anti-CD40

125Tg B6 mouse splenocytes were stimulated with anti-CD40 to simulate T cell help for

three days in the presence of insulin conjugates. Insulin-CD22L significantly reduced anti-insulin

B cell proliferation by 29% compared to anti-CD40 alone or in the presence of unconjugated

mCD22L (51.1% ±11.4% anti-CD40 alone vs 50.5% ± 9.0% anti-CD40 with mCD22L-azide p=

0.9926; 51.1% ±11.4% anti-CD40 alone vs 36.3% ±11.8% anti_CD40 with insulin-mCD22L

p=0.0313) (**Figure 6A, Table 2**). Once again, the 4-arm PEG-insulin(2) was highly activating,

inducing 81.2% more proliferation compared to anti-CD40 alone. The addition of mCD22L to

the PEG backbone in 4-arm PEG-insulin(2)/mCD22L(2) again antagonized excess stimulatory

effects, returning proliferation to that of B cells stimulated with anti-CD40 alone (51.1%±11.4%

anti-CD40 alone vs 46.9%±10.3% anti-CD40 with 4-arm PEG-insulin(2)/mCD22L(2) p=0.6280)

Figure 6: Effects of insulin/mCD22L conjugates on anti-CD40 B cell stimulation in 125Tg B6 mouse splenocytes at 3 days. A) Representative histograms and quantification of B cell proliferation by dye-dilution assay following stimulation with Anti-CD40 and treatment with mCD22L-azide or the insulin-mCD22L direct conjugate (* p<0.05 by one-way Anova with Tukey's multiple comparison) n≥6 mice per condition. B) Representative histograms and quantification of B cell proliferation by dye-dilution assay following stimulation with Anti-CD40 and treatment with 4-arm PEG-insulin(2) or 4-arm PEG-insulin(2)/mCD22L(2) (*p<0.05 **** p<0.0001by one-way Anova with Tukey's multiple comparison). n=8 mice per condition.

(**Figure 6B**). Anti-CD40 stimulated WT B6 splenocytes were not affected by the insulin

constructs (**Figure S4**).

CD86 expression in response to anti-CD40 simulation was indistinguishable from

stimulated anti-insulin B cells treated with insulin-mCD22L (**Figure 7A, Table 2**). However,

CD86 expression was strongly upregulated in response to anti-CD40 stimulation when treated

with 4-arm PEG-insulin(2) and 4-arm PEG-insulin(2)/mCD22L(2) compared with no treatment

Figure 7: Surface marker expression on anti-insulin B cells after 3 days of anti-CD40 stimulation. A) Representative histograms of CD86 expression levels of anti-insulin B cells and quantification as a ratio over anti-CD40 stimulated WT B cells for treatment with mCD22L-azide, 4-arm PEG-insulin(2), 4-arm PEG-insulin(2)/mCD22L(2) or the insulin-mCD22L direct conjugate following three days of anti-CD40 stimulation. B) Representative histograms of IgM expression levels of anti-insulin B cells and quantification as a ratio over anti-CD40 stimulated WT B cells for treatment with mCD22L-azide or insulin-mCD22L following three days of anti-CD40 stimulation. C) Representative histograms of CD22 expression levels of anti-insulin B cells and quantification as a ratio over anti-CD40 stimulated WT B cells for treatment with unconjugated mCD22L-azide, 4-arm PEG-insulin(2), 4-arm PEG-insulin(2)/mCD22L(2) or the insulin-mCD22L direct conjugate following three days of anti-CD40 stimulation. n≥6 mice per condition. Dotted line represents mean MFI of stimulated anti-insulin B cells with no construct. (*p<0.05 ** p<0.01***p<0.001 **** p<0.0001 by one-way Anova with Tukey's multiple comparison).

(**Figure 7A**). IgM expression was downregulated by 52.4% in B cells stimulated with anti-CD40

when treated with insulin-mCD22L (**Figure 7B**). Accurate measurement of IgM expression was

not feasible with B cells treated with 4-arm PEG-insulin(2) and 4-arm PEG-insulin(2)/

mCD22L(2) potentially due to the treatment occluding the binding of anti-IgM. In WT mice, the

insulin conjugates did not significantly affect CD22 or IgM expression, but 4-arm PEG-

insulin(2) did increase CD86 expression (**Figure S4**). WT B cell numbers and percent

proliferated were not affected (**Figure S4**). CD22 expression was reduced by 26.5% and 28% for

4-arm PEG-insulin(2)/mCD22L(2) and insulin-mCD22L respectively, whereas 4-arm PEG-

insulin(2) saw a reduction of 54.7% (**Figure 7C**).

Table 2: Average Proliferated B cell proportions and MFI for anti-insulin B cells stimulated with anti-CD40 and
treated with insulin constructs

Condition	Average B Cell Proportion Proliferated	SD	Average IgM MFI Ratio	SD	Average CD86 MFI Ratio	SD	Average CD22 MFI Ratio	SD
Media Only	51.08	11.38	0.97	0.16	0.8	0.2	1.1	0.09
mCD22L-azide	50.45	9.02	0.98	0.15	0.89	0.21	1.13	0.08
4-arm PEG-Insulin(2)	92.59	3.45			3.57	0.84	0.5	0.11
4-arm PEG-Insulin(2)/mCD22L(2)	46.86	10.25			1.65	0.4	0.79	0.15
Insulin-mCD22L	36.28	11.75	0.46	0.08	0.72	0.16	0.8	0.04

Discussion

The 4-arm PEG scaffold enables isolation of discrete, multivalent insulin-mCD22L conjugates

Prior studies investigating BCR-CD22 co-ligation have focused on large liposome and

polyacrylamide or polynorbornene polymer scaffolds that have antigen and CD22L valences in

the range of 100-1000 or more.[3-9] Additionally, by nature of their synthesis, these scaffolds result

in non-discrete ligand valences. Because of the size constraints of liposomes and the

heterogeneity of antigen and CD22L valences achieved with linear polymers, a 4-arm PEG-

alkyne scaffold was selected to synthesize discrete antigen-CD22L conjugates to investigate the

ability of lower-valency antigen-CD22L conjugates to crosslink BCR and CD22 to prevent B

cell activation. Here, the 20 kDa 4-arm PEG-alkyne proved to be a capable scaffold to synthesize

and isolate discrete insulin-mCD22L conjugates as demonstrated by the significant changes in RP-HPLC retention time and mass distributions on MALDI-TOF mass spectra for the intermediate 4-arm PEG-insulin(1-4) products. The increase in product homogeneity is substantial when the analytical data is compared to that of 20 kDa 4-arm PEG-insulin conjugates with different valency prepared by varying the ratio of insulin to 4-arm PEG scaffold.[33] Although there is 10-20% MW variation in the 20 kDa 4-arm PEG-alkyne, the discrete number of terminal function groups make it a valuable platform to achieve discrete bifunctional conjugates.

The direct insulin-mCD22L conjugate attenuates B cell activation in response to Anti-CD40 stimulation

Treatment of 125Tg mouse splenocytes with insulin-mCD22L significantly reduced B cell proliferation in response to anti-CD40 stimulation. The ability of a monomeric antigen-CD22L construct to limit B cell activation has not been previously demonstrated to our knowledge, and this data supports the potential of antigen-CD22L constructs to limit T-dependent B cell activation. Previous studies demonstrated that CD22 knock-out or CD22 mutation to prevent ligand binding result in hyperproliferation following anti-CD40 stimulation in murine B cells, illustrating how CD22 can attenuate CD40 signaling.[36, 37] The mechanism is not entirely clear, as CD22 does not associate with CD40, and anti-CD40 stimulation alone is incapable of inducing CD22 phosphorylation.[38, 39] However, anti-CD40 stimulation synergistically increases CD22 ITIM phosphorylation with anti-IgM stimulation via Lyn, further suggesting that CD22 plays a negative feedback role in CD40 activation.[39] The need for BCR stimulation in tandem with anti-CD40 stimulation to attenuate B cell activation in a CD22-dependent manner is in congruence with our data showing a lack of attenuation when cells were

treated with mCD22L-azide. CD40-mediated activating signals are propagated by TRAF1-6 and

JAK3 which mediate downstream pathways including the canonical and non-canonical NFkB

signaling; p38, Akt and JNK activation; and STAT5 phosphorylation.[40] These pathways may be

attenuated by CD22-activated protein tyrosine phosphatase SHP-1 which can dephosphorylate

TRAF3 and associate with TRAF2/6 to regulate downstream signals or by SHP-1

dephosphorylation of downstream effectors such as Syk and BLNK which play a role in both

BCR- and CD40-mediated B cell activation.[41-46] Further study is needed to elucidate the

mechanisms by which BCR-CD22 co-ligation attenuates CD40 pro-inflammatory stimulation.

The addition of 2 mCD22Ls to 4-arm PEG-insulin(2) neutralizes B cell activation

125Tg splenocyte treatment with bivalent 4-arm PEG-insulin(2) increased B cell

activation with or without anti-CD40 co-stimulation. The ability of 4-arm PEG-insulin(2) to

crosslink BCRs in a fashion similar to anti-IgM Fab2' is the likely cause of this response, but the

large magnitude of B cell activation was unexpected based on prior investigations into the effects

of lower-valency insulin-polymer conjugates on insulin-specific B cell activation.[33, 47] However,

the addition of 2 mCD22Ls to 4-arm PEG-insulin(2), to make 4-arm PEG-

insulin(2)/mCD22L(2), abrogated the excessive proliferative effects and resulted in CD86 and

proliferation levels indistinguishable from those of untreated cells. This also occurred in the

presence of anti-CD40 co-simulation. However, unlike the direct insulin-mCD22L conjugate, 4-

arm PEG-insulin(2)/mCD22L(2) was not able to reduce B cell activation resulting from anti-

CD40 stimulation. Therefore, the inclusion of an equal number of mCD22L with bivalent

antigen was only capable of negating the excessive proliferation resulting from BCR cross-

linking. In comparison, high-valency antigen and CD22L-bearing liposomes required 10-30-fold

more CD22L than antigen to mitigate the inflammatory stimulation of the multi-valent antigen to baseline levels.[5] BCR and CD40 stimulation is synergistic in B cell activation; therefore, avoiding BCR crosslinking is important to achieve tolerogenic effects with antigen-CD22L constructs in the presence of T cell help.[45, 48, 49] As demonstrated by prior antigen and CD22L-bearing liposomes, increasing antigen valency may require an exponential increase in CD22L valency to overcome the pro-inflammatory signal resulting from BCR crosslinking.

Low valency antigen-CD22L conjugates can successfully co-ligate BCR and CD22

High valency antigen-CD22L polymers and liposomes have previously demonstrated the ability of BCR/CD22 co-ligation to attenuate B cell activation, but emphasis has been placed on high CD22L valency being necessary in these constructs to outcompete binding with endogenous cis-ligands of CD22 such as sialic acids present on CD22 molecules.[3-5, 32] The high-affinity mCD22L used in our insulin-mCD22L conjugates has an approximate K_D of 1 μM for mCD22; much higher than the 150 nM concentration of insulin-mCD22L used in the assays.[32] Additionally, the surface concentration of cis-ligands on the B cells is estimated to be greater than 10 mM although the binding affinity of native CD22Ls is in the 100 μM range.[50] From these values, it is understood that binding of monovalent CD22L alone to B cells is improbable. However, conjugation of the CD22L to an antigen possessing sufficient affinity for the BCR can immobilize the antigen-CD22L conjugate on the cell surface through BCR-antigen binding. This mechanism, in which the antigen binds first, results in a high local concentration of the CD22L promoting CD22 binding and BCR/CD22 co-ligation. We postulate the same effect occurs with the 4-arm PEG-insulin(2)/mCD22L(2) although BCR cross-linking also takes place. The exact degree of BCR/CD22 co-ligation achieved by the insulin-mCD22L and the 4-arm PEG-

insulin(2)/mCD22L(2) is uncertain, and we agree with the hypothesis that a greater extent of BCR/CD22 co-ligation could be achieved by increasing the CD22L valency of each respective construct, but our data does show that conjugating a single CD22L to a single copy of antigen is capable of BCR/CD22 co-ligation in spite of competition from cis-ligands.

Conclusion

High valency antigen-CD22L conjugates have previously demonstrated the ability to prevent B cell activation. Here we demonstrated that 1:1 insulin-mCD22L attenuates B cell activation from mimicked T-dependent stimulation. However, a 2:2 insulin-mCD22L conjugate was not able to reduce B cell activation from anti-CD40 stimulation, demonstrating that competing inflammatory and anti-inflammatory signals stemming from BCR crosslinking versus BCR and CD22 co-ligation are considerable. The data suggest that antigen-CD22L constructs possessing a single copy of antigen are well-suited to achieve immunosuppressive effects, making it a promising agent for future preclinical trials.

References

1.	Hampe, C. S., B Cells in Autoimmune Diseases. *Scientifica* **2012,** *2012*, 215308.

2.	Shlomchik, M. J., Sites and Stages of Autoreactive B Cell Activation and Regulation. *Immunity* **2008,** *28* (1), 18-28.

3.	Courtney, A. H.; Puffer, E. B.; Pontrello, J. K.; Yang, Z.-Q.; Kiessling, L. L., Sialylated multivalent antigens engage CD22 in trans and inhibit B cell activation. *Proceedings of the National Academy of Sciences* **2009,** *106* (8), 2500.

4.	Duong, B. H.; Tian, H.; Ota, T.; Completo, G.; Han, S.; Vela, J. L.; Ota, M.; Kubitz, M.; Bovin, N.; Paulson, J. C.; Nemazee, D., Decoration of T-independent antigen with ligands for CD22 and Siglec-G can suppress immunity and induce B cell tolerance in vivo. *Journal of Experimental Medicine* **2009,** *207* (1), 173-187.

5.	Macauley, M. S.; Pfrengle, F.; Rademacher, C.; Nycholat, C. M.; Gale, A. J.; von Drygalski, A.; Paulson, J. C., Antigenic liposomes displaying CD22 ligands induce antigen-specific B cell apoptosis. *The Journal of Clinical Investigation* **2013,** *123* (7), 3074-3083.

6.	Bednar, K. J.; Shanina, E.; Ballet, R.; Connors, E. P.; Duan, S.; Juan, J.; Arlian, B. M.; Kulis, M. D.; Butcher, E. C.; Fung-Leung, W.-P.; Rao, T. S.; Paulson, J. C.; Macauley, M. S., Human CD22 Inhibits Murine B Cell Receptor Activation in a Human CD22 Transgenic Mouse Model. *The Journal of Immunology* **2017,** *199* (9), 3116-3128.

7.	Orgel, K. A.; Duan, S.; Wright, B. L.; Maleki, S. J.; Wolf, J. C.; Vickery, B. P.; Burks, A. W.; Paulson, J. C.; Kulis, M. D.; Macauley, M. S., Exploiting CD22 on antigen-specific B cells to prevent allergy to the major peanut allergen Ara h 2. *Journal of Allergy and Clinical Immunology* **2017,** *139* (1), 366-369.e2.

8.	Bednar, K. J.; Nycholat, C. M.; Rao, T. S.; Paulson, J. C.; Fung-Leung, W.-P.; Macauley, M. S., Exploiting CD22 To Selectively Tolerize Autoantibody Producing B-Cells in Rheumatoid Arthritis. *ACS Chemical Biology* **2019,** *14* (4), 644-654.

9.	Srivastava, A.; Arlian, B. M.; Pang, L.; Kishimoto, T. K.; Paulson, J. C., Tolerogenic Nanoparticles Impacting B and T Lymphocyte Responses Delay Autoimmune Arthritis in K/BxN Mice. *ACS Chemical Biology* **2021,** *16* (10), 1985-1993.

10.	Islam, M.; Arlian, B. M.; Pfrengle, F.; Duan, S.; Smith, S. A.; Paulson, J. C., Suppressing Immune Responses Using Siglec Ligand-Decorated Anti-receptor Antibodies. *Journal of the American Chemical Society* **2022,** *144* (21), 9302-9311.

11.	Horton, H. M.; Chu, S. Y.; Ortiz, E. C.; Pong, E.; Cemerski, S.; Leung, I. W.; Jacob, N.; Zalevsky, J.; Desjarlais, J. R.; Stohl, W., Antibody-mediated coengagement of FcγRIIb and B cell receptor complex suppresses humoral immunity in systemic lupus erythematosus. *The Journal of Immunology* **2011,** *186* (7), 4223-4233.

12.	Chu, S. Y.; Horton, H. M.; Pong, E.; Leung, I. W. L.; Chen, H.; Nguyen, D.-H.; Bautista, C.; Muchhal, U. S.; Bernett, M. J.; Moore, G. L.; Szymkowski, D. E.; Desjarlais, J. R., Reduction of total IgE by targeted coengagement of IgE B-cell receptor and FcγRIIb with Fc-engineered antibody. *Journal of Allergy and Clinical Immunology* **2012,** *129* (4), 1102-1115.

13.	Chu, S. Y.; Yeter, K.; Kotha, R.; Pong, E.; Miranda, Y.; Phung, S.; Chen, H.; Lee, S.-H.; Leung, I.; Bonzon, C.; Desjarlais, J. R.; Stohl, W.; Szymkowski, D. E., Suppression of Rheumatoid Arthritis B Cells by XmAb5871, an Anti-CD19 Antibody That Coengages B Cell Antigen Receptor

Complex and Fcγ Receptor IIb Inhibitory Receptor. *Arthritis & Rheumatology* **2014,** *66* (5), 1153-1164.

14.   Szili, D.; Cserhalmi, M.; Bankó, Z.; Nagy, G.; Szymkowski, D. E.; Sármay, G., Suppression of innate and adaptive B cell activation pathways by antibody coengagement of FcγRIIb and CD19. *MAbs* **2014,** *6* (4), 991-999.

15.   Merrill, J.; June, J.; Koumpouras, F.; Machua, W.; Khan, M. F.; Askanase, A.; Khosroshahi, A.; Sheikh, S.; James, J. A.; Guthridge, J.; Rathi, G.; Burington, B.; Foster, P.; Zack, D., FRI0176 Phase 2, Double-Blind, Randomized, Placebo-Controlled Study of a Reversible B cell Inhibitor, XmAb 5871, in Systemic Lupus Eruthematosus (SLE). *Annals of the Rheumatic Diseases* **2019,** *78* (Suppl 2), 761-762.

16.   Crocker, P. R.; Paulson, J. C.; Varki, A., Siglecs and their roles in the immune system. *Nature Reviews Immunology* **2007,** *7* (4), 255-266.

17.   Smith, K. G. C.; Clatworthy, M. R., FcγRIIB in autoimmunity and infection: evolutionary and therapeutic implications. *Nature Reviews Immunology* **2010,** *10* (5), 328-343.

18.   Smith, K. G. C.; Tarlinton, D. M.; Doody, G. M.; Hibbs, M. L.; Fearon, D. T., Inhibition of the B Cell by CD22: A Requirement for Lyn. *Journal of Experimental Medicine* **1998,** *187* (5), 807-811.

19.   Doody, G. M.; Justement, L. B.; Delibrias, C. C.; Matthews, R. J.; Lin, J.; Thomas, M. L.; Fearon, D. T., A Role in B Cell Activation for CD22 and the Protein Tyrosine Phosphatase SHP. *Science* **1995,** *269* (5221), 242-244.

20.   Tedder, T. F.; Poe, J. C.; Haas, K. M., CD22: A Multifunctional Receptor That Regulates B Lymphocyte Survival and Signal Transduction. In *Advances in Immunology*, Academic Press: 2005; Vol. 88, pp 1-50.

21.   Willcox, A.; Richardson, S. J.; Bone, A. J.; Foulis, A. K.; Morgan, N. G., Analysis of islet inflammation in human type 1 diabetes. *Clinical & Experimental Immunology* **2009,** *155* (2), 173-181.

22.   Wong, F. S.; Wen, L.; Tang, M.; Ramanathan, M.; Visintin, I.; Daugherty, J.; Hannum, L. G.; Janeway, C. A., Jr; Shlomchik, M. J., Investigation of the Role of B-Cells in Type 1 Diabetes in the NOD Mouse. *Diabetes* **2004,** *53* (10), 2581-2587.

23.   Xiu, Y.; Wong, C. P.; Bouaziz, J.-D.; Hamaguchi, Y.; Wang, Y.; Pop, S. M.; Tisch, R. M.; Tedder, T. F., B Lymphocyte Depletion by CD20 Monoclonal Antibody Prevents Diabetes in Nonobese Diabetic Mice despite Isotype-Specific Differences in FcγR Effector Functions. *The Journal of Immunology* **2008,** *180* (5), 2863-2875.

24.   Leete, P.; Willcox, A.; Krogvold, L.; Dahl-Jørgensen, K.; Foulis, A. K.; Richardson, S. J.; Morgan, N. G., Differential Insulitic Profiles Determine the Extent of β-Cell Destruction and the Age at Onset of Type 1 Diabetes. *Diabetes* **2016,** *65* (5), 1362-1369.

25.   Serreze, D. V.; Fleming, S. A.; Chapman, H. D.; Richard, S. D.; Leiter, E. H.; Tisch, R. M., B Lymphocytes Are Critical Antigen-Presenting Cells for the Initiation of T Cell-Mediated Autoimmune Diabetes in Nonobese Diabetic Mice. *The Journal of Immunology* **1998,** *161* (8), 3912-3918.

26.   Mariño, E.; Tan, B.; Binge, L.; Mackay, C. R.; Grey, S. T., B-Cell Cross-Presentation of Autologous Antigen Precipitates Diabetes. *Diabetes* **2012,** *61* (11), 2893-2905.

27.   Serreze, D. V.; Chapman, H. D.; Varnum, D. S.; Hanson, M. S.; Reifsnyder, P. C.; Richard, S. D.; Fleming, S. A.; Leiter, E. H.; Shultz, L. D., B lymphocytes are essential for the

initiation of T cell-mediated autoimmune diabetes: analysis of a new "speed congenic" stock of NOD.Ig mu null mice. *Journal of Experimental Medicine* **1996,** *184* (5), 2049-2053.

28.   Noorchashm, H.; Lieu, Y. K.; Noorchashm, N.; Rostami, S. Y.; Greeley, S. A. S.; Schlachterman, A.; Song, H. K.; Noto, L. E.; Jevnikar, A. M.; Barker, C. F.; Naji, A., I-Ag7-Mediated Antigen Presentation by B Lymphocytes Is Critical in Overcoming a Checkpoint in T Cell Tolerance to Islet β Cells of Nonobese Diabetic Mice. *The Journal of Immunology* **1999,** *163* (2), 743-750.

29.   Hulbert, C.; Riseili, B.; Rojas, M.; Thomas, J. W., Cutting Edge: B Cell Specificity Contributes to the Outcome of Diabetes in Nonobese Diabetic Mice. *The Journal of Immunology* **2001,** *167* (10), 5535-5538.

30.   Acevedo-Suárez, C. A.; Hulbert, C.; Woodward, E. J.; Thomas, J. W., Uncoupling of Anergy from Developmental Arrest in Anti-Insulin B Cells Supports the Development of Autoimmune Diabetes. *The Journal of Immunology* **2005,** *174* (2), 827-833.

31.   Henry-Bonami, R. A.; Williams, J. M.; Rachakonda, A. B.; Karamali, M.; Kendall, P. L.; Thomas, J. W., B Lymphocyte "Original Sin" in the Bone Marrow Enhances Islet Autoreactivity in Type 1 Diabetes–Prone Nonobese Diabetic Mice. *The Journal of Immunology* **2013,** *190* (12), 5992-6003.

32.   Collins, B. E.; Blixt, O.; Han, S.; Duong, B.; Li, H.; Nathan, J. K.; Bovin, N.; Paulson, J. C., High-Affinity Ligand Probes of CD22 Overcome the Threshold Set by Cis Ligands to Allow for Binding, Endocytosis, and Killing of B Cells. *The Journal of Immunology* **2006,** *177* (5), 2994-3003.

33.   Johnson, S. N.; Griffin, J. D.; Hulbert, C.; DeKosky, B. J.; Thomas, J. W.; Berkland, C. J., Multimeric Insulin Desensitizes Insulin-Specific B Cells. *ACS Applied Bio Materials* **2020,** *3* (9), 6319-6330.

34.   Apley, K. D.; Laflin, A. D.; Johnson, S. N.; Batrash, N.; Griffin, J. D.; Berkland, C. J.; DeKosky, B. J., Optimized Production of Fc Fusion Proteins by Sortase Enzymatic Ligation. *Industrial & Engineering Chemistry Research* **2021,** *60* (47), 16839-16853.

35.   Nyhoff, L. E.; Griffith, A. S.; Clark, E. S.; Thomas, J. W.; Khan, W. N.; Kendall, P. L., Btk Supports Autoreactive B Cell Development and Protects against Apoptosis but Is Expendable for Antigen Presentation. *The Journal of Immunology* **2021,** *207* (12), 2922-2932.

36.   Poe, J. C.; Fujimoto, Y.; Hasegawa, M.; Haas, K. M.; Miller, A. S.; Sanford, I. G.; Bock, C. B.; Fujimoto, M.; Tedder, T. F., CD22 regulates B lymphocyte function in vivo through both ligand-dependent and ligand-independent mechanisms. *Nature Immunology* **2004,** *5* (10), 1078-1087.

37.   Poe, J. C.; Haas, K. M.; Uchida, J.; Lee, Y.; Fujimoto, M.; Tedder, T. F., Severely Impaired B Lymphocyte Proliferation, Survival, and Induction of the c-Myc:Cullin 1 Ubiquitin Ligase Pathway Resulting from CD22 Deficiency on the C57BL/6 Genetic Background. *The Journal of Immunology* **2004,** *172* (4), 2100-2110.

38.   MoyrOn-QuirOz, J. E.; Partida-Sánchez, S.; Donís-Hernández, R.; Sandoval-Montes, C.; Santos-Argumedo, L., Expression and Function of CD22, a B-cell Restricted Molecule*. *Scandinavian Journal of Immunology* **2002,** *55* (4), 343-351.

39.   Fujimoto, M.; Kuwano, Y.; Watanabe, R.; Asashima, N.; Nakashima, H.; Yoshitake, S.; Okochi, H.; Tamaki, K.; Poe, J. C.; Tedder, T. F.; Sato, S., B Cell Antigen Receptor and CD40

Differentially Regulate CD22 Tyrosine Phosphorylation. *The Journal of Immunology* **2006,** *176* (2), 873-879.

40.   Elgueta, R.; Benson, M. J.; De Vries, V. C.; Wasiuk, A.; Guo, Y.; Noelle, R. J., Molecular mechanism and function of CD40/CD40L engagement in the immune system. *Immunological reviews* **2009,** *229* (1), 152-172.

41.   Zhang, Z.; Jimi, E.; Bothwell, A. L. M., Receptor Activator of NF-κB Ligand Stimulates Recruitment of SHP-1 to the Complex Containing TNFR-Associated Factor 6 That Regulates Osteoclastogenesis. *The Journal of Immunology* **2003,** *171* (7), 3620-3626.

42.   Hao, D.; Wang, Y.; Li, L.; Qian, G.; Liu, J.; Li, M.; Zhang, Y.; Zhou, R.; Yan, D., SHP-1 suppresses the antiviral innate immune response by targeting TRAF3. *The FASEB Journal* **2020,** *34* (9), 12392-12405.

43.   Busca, A.; Konarski, Y.; Gajanayaka, N.; O'Hara, S.; Angel, J.; Kozlowski, M.; Kumar, A., cIAP1/2–TRAF2–SHP-1–Src–MyD88 Complex Regulates Lipopolysaccharide-Induced IL-27 Production through NF-κB Activation in Human Macrophages. *The Journal of Immunology* **2018,** *200* (5), 1593-1606.

44.   Peng, L.; Zhou, Y.; Jiang, N.; Wang, T.; Zhu, J.; Chen, Y.; Li, L.; Zhang, J.; Yu, S.; Zhao, Y., DJ-1 exerts anti-inflammatory effects and regulates NLRX1-TRAF6 via SHP-1 in stroke. *Journal of Neuroinflammation* **2020,** *17* (1), 81.

45.   Ying, H.; Li, Z.; Yang, L.; Zhang, J., Syk mediates BCR- and CD40-signaling integration during B cell activation. *Immunobiology* **2011,** *216* (5), 566-570.

46.   Ren, L.; Chen, X.; Luechapanichkul, R.; Selner, N. G.; Meyer, T. M.; Wavreille, A.-S.; Chan, R.; Iorio, C.; Zhou, X.; Neel, B. G.; Pei, D., Substrate Specificity of Protein Tyrosine Phosphatases 1B, RPTPα, SHP-1, and SHP-2. *Biochemistry* **2011,** *50* (12), 2339-2356.

47.   Leon, M. A.; Wemlinger, S. M.; Larson, N. R.; Ruffalo, J. K.; Sestak, J. O.; Middaugh, C. R.; Cambier, J. C.; Berkland, C., Soluble Antigen Arrays for Selective Desensitization of Insulin-Reactive B Cells. *Molecular Pharmaceutics* **2019,** *16* (4), 1563-1572.

48.   Haxhinasto, S. A.; Hostager, B. S.; Bishop, G. A., Cutting Edge: Molecular Mechanisms of Synergy Between CD40 and the B Cell Antigen Receptor: Role for TNF Receptor-Associated Factor 2 in Receptor Interaction. *The Journal of Immunology* **2002,** *169* (3), 1145-1149.

49.   Bishop, G. A.; Warren, W. D.; Berton, M. T., Signaling via major histocompatibility complex class II molecules and antigen receptors enhances the B cell response to gp39/CD40 ligand. *European Journal of Immunology* **1995,** *25* (5), 1230-1238.

50.   Collins, B. E.; Blixt, O.; DeSieno, A. R.; Bovin, N.; Marth, J. D.; Paulson, J. C., Masking of CD22 by cis ligands does not prevent redistribution of CD22 to sites of cell contact. *Proceedings of the National Academy of Sciences* **2004,** *101* (16), 6104-6109.