Aberrant cortical activity, functional connectivity, and neural assembly

architecture after photothrombotic stroke in mice

Mischa V. Bandet,[1,2,3] and Ian R. Winship[1,2,3]*

[1]Neuroscience and Mental Health Institute

[2]Neurochemical Research Unit

[3]Department of Psychiatry

University of Alberta, Edmonton, Alberta T6G 2B7, Canada

14   **Keywords:** stroke**;** photothrombosis; two-photon; calcium imaging; gcamp; somatosensory

15   cortex; neural assemblies; ensembles; limb-associated

18   **Running Title:** Aberrant cortical activity, functional connectivity and neural assembly
19   architecture after stroke
20
21
22
23

*Corresponding Author:
Dr. Ian R. Winship
Professor
Department of Psychiatry
12-127 Clinical Sciences Building
University of Alberta
Edmonton, AB
Canada, T6G 2G3
Phone: 780-492-8658
Fax: 780-492-6841
iwinship@ualberta.ca

Abstract

Despite substantial recent progress in mapping the trajectory of network plasticity resulting from focal ischemic stroke, there is mixed evidence for changes in neuronal excitability and activity within the peri-infarct cortex of mice. Most of these data have been acquired from anesthetized animals, acute tissue slices, or immunoassays on extracted tissue, and thus may not reflect cortical activity dynamics in the intact cortex of an awake animal. Here, *in vivo* two-photon calcium imaging in awake, behaving mice was used to longitudinally track cortical activity, network functional connectivity, and neural assembly architecture for 2 months following photothrombotic stroke targeting the forelimb somatosensory cortex. This model is associated with sensorimotor recovery over the weeks following stroke, allowing us to relate network changes to behavior. Our data revealed spatially restricted but long-lasting alterations in somatosensory neural networks. Specifically, we demonstrate significant and long-lasting disruptions in neural assembly architecture concurrent with a deficit in functional connectivity between individual neurons. Reductions in neuronal spiking in peri-infarct cortex were transient but predictive of impairment in skilled locomotion measured in the tapered beam task. Notably, altered neural networks were highly localized, with assembly architecture and neural connectivity relatively unaltered a distance outside the peri-infarct cortex, even in regions demonstrated to be the site of functional remapping of the forelimb somatosensory representation in anaesthetized preparations. Thus, using longitudinal two photon microscopy in awake animals, these data show a complex spatiotemporal relationship between peri-infarct neuronal network function and behavioral recovery that is more nuanced than functional remapping in response to strong sensory stimulation under anesthesia.

Introduction

Investigations of the excitability of neurons in the peri-infarct cortex during recovery from ischemic stroke have yielded a complex and at times contradictory data set. Peri-infarct hyper-excitability was suggested by data from anesthetized rats that exhibited elevated spontaneous multiunit firing within peri-infarct regions 3-7 days after stroke [1]. Similarly, downregulated GABAergic inhibition in peri-infarct cortex [1–6], degeneration of parvalbumin-positive inhibitory interneurons [3,7], decreased paired pulse inhibition [8–10], increased NMDA receptor mediated excitation [11,12], and a downregulation of KCC2 [13–17] have suggested that the peri-infarct cortex is hyper-excitable during recovery, contributing to increased risk of epileptogenesis in the post stroke brain [13,18]. Conversely, several groups have reported that the sensory-evoked responsiveness of the peri-infarct cortex is diminished during recovery from focal photothombotic stroke [19–25]. Even after months of recovery following forelimb (FL) representation targeted stroke, the remapped representation of the FL has been shown to display prolonged modes of activation with lower amplitude [19] and reduced temporal fidelity [23].

Almost all studies that have examined post-stroke cortical responsiveness have used surgical preparations with anesthetized animals. Anesthesia disrupts cortical activity dynamics and functional connectivity between cortical areas [26–30], and has been shown to reduce the magnitude and spread of cortical activation from sensory stimuli [31–39]. It further reduces the potential contribution of corollary discharge and reafference due to voluntary action in modulating sensory input and cortical responses [40]. It has also recently been shown that the propagation pattern of cortical activity differs between evoked and spontaneous activity, with spontaneous activity showing more complex trajectories and lower activity amplitudes [41].

Somatosensory responses are known to be modulated by the relevance of stimuli to behavior and

task performance [42–47], and by corollary feedback during [48], or even before [49–51], the

onset of movement. Together, these studies point to a complex, interconnected system that

modulates the activity of somatosensory networks in the awake behaving animal that is either not

present, or is altered, in studies of evoked cortical activity under the anesthetized state. While

monitoring cortical activity in awake mobile animals presents methodological complexity, it

avoids confounds associated with the anesthetized state and provides unique insight into

evolving patterns of network activity that may be unique to the awake state.

To investigate neural activity patterns during stroke recovery while avoiding the potential

confounds of anesthesia, the present work used two photon Ca^{2+} imaging within and distal to the

peri-infarct region of a focal photothrombotic stroke lesioning the FL primary somatosensory

cortex in mice. We used awake, freely behaving but head-fixed mice in a mobile homecage to

longitudinally measure cortical activity, then used computational methods to assess functional

connectivity and neural assembly architecture at baseline and each week for 2 months following

stroke. Behavioral recovery from stroke was measured in the same animals on a tapered beam

task and string pull task to determine the time course of sensorimotor deficits related to changes

in cortical activity and network architecture. Our goal was to use longitudinal imaging and

advanced computational analyses to define network changes in somatosensory cortex associated

with sensorimotor recovery. We show that despite disturbances to the widefield topology and

amplitude of the sensory-evoked FL map (acquired under anesthetic), it is only within the

immediate peri-infarct region after FL targeted stroke that neural activity, functional

connectivity, and neural assembly architecture are disrupted. We demonstrate that the peri-

infarct somatosensory cortex displays a marked reduction in neural assembly number, an

increase in neural assembly membership, and persistent alterations in assembly-assembly

correlations, and that this occurs alongside transient deficits in both functional connectivity and

neural activity. Notably, we show that reduced neuronal spiking is strongly correlated with a

deficit in motor performance on the tapered beam task. Surprisingly, we also show that

measurable deficits in neuronal activity, functional connectivity, and neural assembly

architecture are absent within distal cortex as little as 750 μm from the stroke border, even in

areas identified by regional imaging as locations of sensory-evoked FL representational

remapping.

Results

The experimental timeline is illustrated in Fig 1A. Adult Thy1-GCaMP6S mice were

implanted with chronic cranial windows and habituated on the floating homecage, tapered beam

and string pull task (Fig 1A-C). The cFL and cHL somatosensory areas were mapped on the

cortex using widefield Ca^{2+} imaging of stimulus-evoked activity (Fig 1D, 2A, see methods).

Regions of interest for longitudinal awake two-photon Ca^{2+} imaging of cellular activity (Fig 1E)

were chosen based on pre-stroke widefield Ca^{2+} limb mapping (Fig 2) acquired under isoflurane

anaesthesia. The first imaging region of interest was located at the boundary of the pre-stroke

cFL and cHL somatosensory maps (termed "peri-infarct" region based on its proximal location

to the photothrombotic infarct after stroke) and the second region as lateral to the pre-stroke cHL

map (termed "distal" region due to its distance from the stroke boundary). These regions were

imaged at baseline, and each week following photothrombotic stroke. Notably, these regions of

interest incorporated the predicted areas of remapping for the limb associated somatosensory

representations after focal cFL cortex stroke based on previous studies [25,52]. Photothrombosis

was directed to the cFL somatosensory representation identified by regional imaging. Resulting

infarcts lesioned this region, and borders could be defined by a region of hyperfluorescence 1

week post-stroke (Fig 1D, Top). Within the peri-infarct imaging region, cellular dysmorphia and

swelling was visually apparent in some cells during two photon imaging 1 week after stroke, but

recovered over the 2 month post-stroke imaging timeframe (Fig 1D, Bottom). Morphological

changes did not occur in the more distal imaging region lateral to the cHL.

Altered sensory-evoked widefield Ca^{2+} maps after stroke

Regional remapping occurs over the weeks following focal cortical stroke [25,52]. Here,

widefield Ca^{2+} imaging of the cranial window in isoflurane anaesthetized mice was performed

during piezoelectric stimulation of the contralateral limbs prior to stroke, and at the 8 week post-

stroke timepoint (Fig 2A, see methods), to define regional sites for cellular Ca^{2+} imaging and to

monitor mesoscale changes to somatosensory evoked cortical representations. Fig 2A shows

representative montages from a stroke animal illustrating the cortical cFL and cHL Ca^{2+}

responses to 1s, 100Hz limb stimulation of the contralateral limbs at the pre-stroke and 8 week

post-stroke timepoints. The location and magnitude of the cortical responses changes drastically

between timepoints, with the cFL remapping posterior to its pre-stroke location into the area

lateral to the cHL map at 8 weeks post-stroke. We observed a significant decrease in the cFL

evoked Ca^{2+} response amplitude in the stroke group at 8 weeks post-stroke compared with the

pre-stroke timepoint (Fig 2B). This is in agreement with past studies [19–25], and suggests that

cFL targeted stroke reduces forelimb evoked activity across the somatosensory cortex in

anaesthetized animals even at late time points 2 months after stroke. We observed a concurrent

increase in cFL evoked response area in the stroke group 8 weeks post-stroke (Fig 2C),

suggesting broader regional activation in response to cFL stimulation with significant posterior

remapping of the FL representation (Fig 2D). There were no group differences between stroke

and sham groups in cHL evoked intensity, area, or map position (data not shown).

Impaired performance on tapered beam but not string pull after focal somatosensory

stroke

Post-stroke sensorimotor impairment was measured on the tapered beam task and string

pull task. In the tapered beam, errors (slips off the beam), indicative of post-stroke impairment,

were tracked as the number of left (affected) side slips, right (non-affected) side slips, and the

distance to first slip (Fig 3A). We observed a significant interaction between stroke and time

after stroke in the number of left side (contralesional) slips (Fig 3B), with *post-hoc* tests

confirming a significant increase in left side slips at the 1 week timepoint for stroke animals

compared the pre-stroke timepoint ($P = 0.0125$) and in stroke animals compared to sham ($P =$

0.0435). Consistent with the unilateral nature of the photothrombotic damage to the cortex in this

study, no main effect or interaction was seen in the number of right side slips (Fig 3C). A

statistical trend ($P = .0979$) towards a main effect of group was observed in the normalized

distance to first slip (Fig 3D). These results are consistent with a transient deficit in motor

behavior on the tapered beam during the first week after stroke induced by focal cortical lesion to

the sensorimotor cortex [53,54], as opposed to more sustained deficits observed in models of

middle cerebral artery occlusion [55,56]. In contrast to previous studies investigating string pull

after stroke [57], we did not detect a significant effect of photothrombosis on any kinematic

parameters during string pulling (Fig S1).

Firing rate of neurons in the peri-infarct cortex correlates with performance on the tapered beam task

The firing rate of cortical somatosensory neurons in our awake behaving animals (Fig 4A) was calculated using a custom Matlab algorithm to identify 'significant' calcium transients [58,59] and the mean firing rate of the sampled neural population during periods of movement and periods of rest in the mobile homecage. The normalized average firing rate in the peri-infarct region during recovery during movement and rest is illustrated in Figures 4B and 4C, respectively. A significant interaction between stroke group and recovery time on firing rate was detected at rest (Fig 4C), though a qualitative (but not statistically significant) reduction in firing rate at 1 week was also apparent during movement (Fig 4B). Notably, normalized firing rate at 1 week post-stroke during movement and at rest were significantly negatively correlated with the number of left slips in the tapered beam at the same 1 week timepoint ($R^2 = 0.5326$, $P = 0.002$ and $R^2 = 0.4346$, $P = 0.0075$, respectively) (Fig 4D and 4E). No main effects or significant interactions were observed in the distal imaging region, though a statistical trend ($P=0.0667$) towards a main effect of group was observed during animal movement (Fig 4F). Normalized firing rate during movement at 1 week within the distal region was not significantly correlated with left slips on the tapered beam at the same timepoint ($R^2 = 0.0637$, $P = 0.3286$) (Fig 4H). A statistical trend suggesting a negative correlation between normalized firing rate at rest and left slips was observed in the distal cortex ($R^2 = 0.2320$, $P = 0.0503$) (Fig 4I). These results run contrary to previous reports of increased multi-unit activity as early as 3-7 days after focal photothrombotic stroke in the peri-infarct cortex [1], and indicate decreased neuronal spiking 1 week after stroke in regions immediately adjacent the infarct that is strongly related to sensorimotor impairment. While the majority of neuronal activity occurred during movement,

196   correlations between firing rate and animal movement were low and did not vary between stroke

197   and sham groups (Fig S2).

199   **Altered functional connectivity in the peri-infarct somatosensory cortex**

200   Studies in humans and animals using mesoscale imaging methods indicate that stroke

disrupts functional connectivity between widespread cortical regions [26,60–70]. However, few

studies have examined functional connectivity at the level of local neural populations after focal

stroke [71,72], and none have examined the primary somatosensory cortex. To define functional

connectivity in local neural networks within the peri-infarct somatosensory cortex, we plotted the

functional connectivity of the neural population in terms of the strength of their cell-cell

correlations (Fig 5) and quantified the properties of these connections (Fig 6). We observed a

significant loss of functional connectivity as early as 1 week after stroke within the peri-infarct

region. This was visually apparent in the stroke group functional connectivity plots at the 1 week

timepoint (Fig 5A and 5B) compared to the same location in sham animals (Fig 5E and 5F).

Notably, no profound loss of average functional connectivity is visually apparent in the distal

region map in the stroke (Fig 5C and 5D) or sham (Fig 5G and 5H) groups. Within the peri-

infarct region, we observed a statistical trend (P=.0551) towards a main effect of stroke in the

normalized average number of all significant functional connections per neuron (Fig 6A) and a

main effect of stroke group on strong connections (r>.03) per neuron (Fig 6B). Significant

connections were reduced by stroke, and *post-hoc* tests confirmed fewer significant connections

1 week after stroke relative to sham controls. No main effects or interactions between stroke and

time were identified in connectivity or significant connections in distal regions of interest (Fig

6D and 6E, respectively), highlighting the regional specificity of these network changes relative

to the infarct and the original forelimb functional representation. The alterations in significant

connections between neurons in peri-infarct parallel a persistent deficit in functional connectivity

lasting up to 30 days that was identified within the primary and supplementary motor cortex of

mice using in vivo imaging after focal motor cortex stroke [71,72]. Surprisingly, network

density, defined as the ratio of measured significant functional network connections to the

number of total possible network connections if all neurons were interconnected, was not

decreased in either peri-infarct or distal regions in our study (Fig 6C and 6F, respectively). This

is in contrast with previous research in the primary and supplementary motor cortex after focal

motor cortex stroke that has demonstrated deficits in network density lasting up to 3 weeks post-

stroke [72].

Neural assembly structure is disrupted in the peri-infarct region after stroke

The neural assembly hypothesis defines coordinated activity in groups of neurons as the

basis for the representation of external and internal stimuli within the brain [73–75]. Likewise,

perturbations within the structure and function of assemblies may signify aberrant processing of

stimuli within neural networks, and is believed to play a significant role in the dysfunction

observed in brain damage and neuropsychiatric diseases [74]. Although the use of sophisticated

methods for detecting neural assemblies in large populations of neurons has gained increasing

popularity with the advent of large population recordings using Ca^{2+} imaging [76], no previous

studies have determined the longitudinal effects of stroke on the architecture of neural

assemblies in cortex. To determine if focal stroke to the cFL cortex affects the properties of

neural assemblies, we used a PCA-Promax procedure [58,59,76] for neural assembly detection to

determine assembly size, population membership, assembly quantity, and overlap in assembly

membership. The PCA-Promax method for determining neural assemblies allows neurons to

participate in more than a single assembly [58,59], a property necessary to examine the ability of

neural assemblies to dynamically form with varying members of the population for only brief

periods of time [58,59,76]. Fig 7 shows representative cellular Ca^{2+} fluorescence images with

neural assemblies color coded and overlaid for each timepoint for the peri-infarct imaging region

and for the distal imaging region for an example stroke and sham animal (Fig 7A and 7B,

respectively). We observed a significant main effect of stroke group on the number of assemblies

(Fig. 8A) after stroke in the peri-infarct imaging region, representing an overall reduction in

assemblies. *Post-hoc* comparisons confirmed significantly reduced number of assemblies 1 and 4

weeks after stroke relative to sham controls (Fig 8A). A significant main effect on the

normalized number of neurons per assembly was detected, reflecting increased membership in

each assembly that peaked 1 week after stroke (Fig 8B). We observed a statistical trend (P =

0.0729) suggesting a higher average percent of the population per assembly (Fig 8C). These data

suggest that after stroke, neurons in peri-infarct cortex are functionally grouped into few distinct

assemblies, with more neurons per assembly and more of the local network in the same

assembly. For the distal imaging region, no differences in neural assembly properties were

observed between the stroke and sham groups (Fig 8D-E).

259
Correlation between neural assembly activations is increased after stroke

If differential activation of neural assemblies in the naïve brain relates to representation

of different aspects of the external world [58,73,75,77], then changes in the relationship between

the activation patterns of assemblies may predict disrupted processing of sensory information

within neural networks and how these networks differentially represent external stimuli. An

increase in assembly size and reduction in assembly number might reflect a mechanism to

replace lost function and increase somatic sensitivity, but may come at the cost of reduced

stimulus specificity/fidelity within the network. The activation magnitude of assemblies can be

correlated between assemblies over the recording period (Fig 9A). A significant main effect of

stroke was detected, reflecting an increase in the average correlation coefficient for significantly

correlated assembly-assembly pairs after stroke in the peri-infarct, but not distal cortex, when

compared to sham (Fig 9B, 9C). This increased tendency for functionally connected assemblies

to be co-active after stroke may reduce the signal to noise ratio for stimulus representation within

the network and may reduce stimulus specificity within peri-infarct cortex. While assembly

activations were often correlated with movement or rest periods (Fig S3), and 60% of assemblies

in both the peri-infarct and distal regions were significantly correlated with animal movement at

all time points (S3C and S3E, respectively), stroke did not change the average correlation

between assembly activation and animal movement in either the peri-infarct (S3B Fig) or distal

region (S3D Fig).

Discussion

Here, we used longitudinal two photon calcium imaging of awake, head-fixed mice in the

mobile homecage to examine how focal photothrombotic stroke to the forelimb sensorimotor

cortex alters the activity and connectivity of neurons adjacent and distal to the infarct. Consistent

with previous studies using intrinsic optical signal imaging, mesoscale imaging of regional

calcium responses (reflecting bulk neuronal spiking in that region) showed that targeted stroke to

the somatosensory cFL representation completely disrupts the sensory evoked forelimb

representation in the infarcted region. Functional remapping of this representation occurred over

8 weeks after injury, with new sensory evoked spiking found primarily in a region lateral to the

cHL representation. Mesoscale calcium imaging allows direct measurement of aggregate

neuronal spiking, and remapped cortex exhibited reduced amplitudes of sensory-evoked activity

with a broader, less defined cortical map relative to sham. Longitudinal two-photon calcium

imaging in awake animals was used to probe single neuron and local network changes adjacent

the infarct and in this distal remapped region. This imaging revealed a decrease in firing rate at 1

week post-stroke of neurons only in the peri-infarct regions that was significantly negatively

correlated with the number of errors made with the stroke-affected limbs on the tapered beam

task. Peri-infarct cortical networks also exhibited a reduction in the number of functional

connections per neuron and a sustained disruption in neural assembly structure, including a

reduction in the number of assemblies and an increased recruitment of neurons into functional

assemblies. Elevated correlation between assemblies within the peri-infarct region peaked 1

week after stroke and was sustained throughout recovery. Notably, distal networks in the region

associated with functional remapping in anaesthetized preparations was largely undisturbed.

Cortical plasticity after stroke

Plasticity within and between cortical regions contributes to partial recovery of function

and is proportional to both the extent of damage, as well as the form and quantity of

rehabilitative therapy post-stroke [78,79]. A critical period of highest plasticity begins shortly

after the onset of stroke, is greatest during the first few weeks, and progressively diminishes over

the weeks to months after stroke [19,80–84]. Functional recovery after stroke is thought to

depend largely on the adaptive plasticity of surviving neurons that reinforce existing connections

and/or replace the function of lost networks [25,52,85–87]. This neuronal plasticity leads to

somatosensory and motor functional remapping to adjacent areas of the cortex and altered

topographical organization, as observed here. The driver for this process has largely been

ascribed to a complex cascade of intra- and extra-cellular signaling that ultimately leads to

plastic re-organization of the microarchitecture and function of surviving peri-infarct tissue

[52,78,82,86,88–90]. Consistent with our findings, structural and functional remodeling is

dependent on the distance from the stroke core, with closer tissue undergoing greater re-

organization than more distant tissue (for review, see [52]).

Previous research examining the region at the border between the cFL and cHL

somatosensory maps has shown this region to be a primary site for functional remapping after

cFL directed photothrombotic stroke, resulting in a region of cFL and cHL map functional

overlap [25]. Within this overlapping area, neurons have been shown to lose limb selectivity 1

month post-stroke [25]. This is followed by the acquisition of more selective responses 2 months

post-stroke and is associated with reduced regional overlap between cFL and cHL functional

maps [25]. Notably, this functional plasticity at the cellular level was assessed using strong

vibrotactile stimulation of the limbs in anaesthetized animals. Our findings using longitudinal

imaging in awake animals show an initial reduction in firing rate at 1 week post-stroke within

this region that was predictive of functional impairment in the tapered beam task. This transient

reduction may be associated with reduced or dysfunctional thalamic connectivity [91–93] and

reduced transmission of signals from hypo-excitable thalamo-cortical projections [94].

Importantly, the strong negative correlation we observed between the normalized firing rate of

the neural population within the peri-infarct cortex and the number of errors on the affected side,

as well as the quick recovery of firing rate and tapered beam performance, suggests this peri-

infarct cortex activity within the peri-infarct region contributes to the impairment and recovery.

This common timescale of neuronal and functional recovery also coincides with angiogenesis

and re-establishment of vascular support for peri-infarct tissue [81,95–98].

Altered network connectivity after stroke

A highly distributed network of neural circuits forms the basis of information flow in the

mammalian brain [99,100]. Many disease states are thought to result in disturbances in neural

dynamics and connectivity of these networks due to a process of randomization that affects the

network nodes and connections, leading to degraded functional performance of the networks

[100,101]. Most studies looking at the functional connectivity of cortical networks after stroke

have focused on connectivity between cortical regions and few have looked at functional

connectivity within neural populations at the single neuron level within a region of cortex. Our

finding of a decreased number of functional connections per neuron in the peri-infarct

somatosensory cortex are consistent with recent research that demonstrated a decrease in the

total number of functional connections for both inhibitory cells [71] and excitatory cells [71,72]

in the motor and premotor cortex approximately 1 week after focal sensorimotor stroke.

Reductions in strong functional connections between neurons in our experiments peaked 1 week

after stroke, with only partial recovery over the next 7 weeks. These changes were restricted to

peri-infarct cortex, and were not observed in regions associated with functional remapping in

response to vibrotactile stimulation under anesthesia. As the cumulative total number of

functional connections within a neural network is dependent on the number of cells measured

from the population, it is possible that a decrease in the number of connections at early

timepoints in both studies may either reflect a loss of neurons and/or the functional silencing of

neurons within the imaged areas. Unlike Bechay et al. [72], connection density (as % of max

possible connections) did not decrease in either the peri-infarct somatosensory cortex or distal

somatosensory cortex after stroke in our study. Thus, we observed a significant reduction in the

number of strong functional connections between neurons, but not a reduction in the density of

connections. It is notable that the peak deficit in peri-infarct functional connectivity in our study

occurs simultaneously with reduced neuronal firing and sensorimotor impairment. Early

disruption has been reported in anaesthetized animals, and our data in awake animals

demonstrates that early network changes may be a key contributor to dysfunction, and a

restoration of spiking in peri-infarct regions is related to recovery [19–25].

Altered network dynamics after stroke

According to the cell assembly hypothesis, transient synchronous activation of distributed

groups of neurons organize into "neural assemblies" that underly the representation of both

external and internal stimuli in the brain [73]. These assemblies have been demonstrated in the

mammalian [102–111] and zebrafish [58,112–114] brain, and are often similar in presentation

between spontaneous and sensory evoked activations [58,102,107,115,116]. Indeed, the

similarity between spontaneous assemblies and sensory evoked assemblies tends to increase

during development [102], potentially illustrating the "sculpting" of neural circuits towards

commonly encountered sensory stimuli. To date, however, no study has examined changes to the

architecture of neural assemblies on the level of single cells within neural networks of the

somatosensory cortex in the post-stroke brain. Our data shows a transient decrease in the number

of assemblies and an increase in the number of neurons per assembly for the peri-infarct cortex.

If assemblies within peri-infarct somatosensory cortex contribute to differential processing of

distinct elements of sensory experience, a loss in the number of assemblies may contribute to a

loss in sensory range, as is observed in human stroke patients [117–121]. Although there have

been calculations on the typical size of cell assemblies in several regions of the brain as a

percentage of the network population (for review, see [73]), whether there is an optimal number

of assemblies or optimal membership size of a neural assembly for a particular neural population

is currently unknown. It is possible that a decrease in the number of assemblies and an increase

in the membership of neurons in remaining assemblies may signify inefficient or ineffective

forms of sensory processing. Likewise, it has been shown that electrical stimulation of single

somatosensory neurons is sufficient to bias animal behavior towards a desired behavioral

response [122,123], thereby suggesting that a sparse neural code may be sufficient for sensation.

It has also been argued that minimizing correlation within the neural population serves to reduce

representational redundancy and improves representational efficiency [124,125], thereby

suggesting that fewer assemblies, with each containing more members, may be an inefficient

form of representational coding for sensory information. Furthermore, we observed a persistent

increase in the correlation between the activations of assemblies in the peri-infarct cortex that

peaks 1 week post-stroke. Co-activation of multiple assemblies simultaneously may decrease the

signal-to-noise ratio of the representational information that each specific assembly holds. It may

further reduce efficient information transfer within and between cortical networks, and may

relate to prolonged modes of somatosensory activation [19] and reduced fidelity in response to

sensory stimuli observed in mouse models of focal FL stroke [23].

Consistent with previous research [25], we show that at the 8 week timepoint after cFL

photothrombotic stroke the cFL representation, identified by mechanical limb stimulation under

anaesthesia, is shifted posterior from its pre-stroke location into the area lateral to the cHL map.

Notably, our distal imaging region was directly within this remapped cFL area. Despite this, we

found little change in the normalized firing rate in either moving or resting states in this region.

This is in contrast to past research in anesthetized animals that indicated an increase in sensory-

evoked activity within this area lateral to the cHL representation 1-2 months after focal cFL

stroke [25]. Moreover, there was no indication that the cFL remapping into this distal area lateral

to the cHL map resulted in any detectable change to the level of population correlation,

functional connectivity, assembly architecture or assembly activations as compared to sham.

Notably, sensory-evoked activity for regional mapping was evoked via strong vibrotactile (1s,

100Hz) limb stimulation. The combination of anesthetic and strong stimulation may unmask

connectivity in distal regions during anesthesia that is not apparent during more naturalistic

stimulation during awake locomotion or rest. Nonetheless, these results support a limited spatial

distance over which the peri-infarct area exhibits network functional deficits. These results are

also consistent with a spatial gradient of plasticity factors that are generally enhanced with closer

proximity to the infarct core [82,86,88,89].

In summary, this work demonstrates that the peri-infarct region near to the stroke core

displays decreased neuronal activity 1 week after stroke that is predictive of poor performance on

the tapered beam task. Moreover, local neuronal networks adjacent the stroke exhibited

reductions in functional connectivity and sustained alterations in the structure of neuronal

assemblies, suggesting that focal injury leads to long lasting alterations in functional connectivity

between neurons in local neuronal networks. Interestingly, altered connectivity and neuronal

assembly structure was not detected in more distal cortical regions, even though those regions

were associated with functional remapping of somatosensory-evoked activity in response to

vibrotactile limb stimulation under anaesthesia. The persistence of these distal networks during

recovery highlights the importance of awake imaging and naturalistic stimuli to fully define

functional network changes during recovery from nervous system injury. The restriction of these

network changes to peri-infarct cortex could reflect the small penumbra associated with

photothrombotic stroke, and future studies could make use of stroke models with larger

penumbral regions, such as the middle cerebral artery occlusion model. Larger injuries induce

more sustained sensorimotor impairment, and the relationship between neuronal firing,

connectivity, and neuronal assemblies could be further probed relative to recovery or sustained

impairment in these models.

Materials and Methods

Animals

Three to nine month old Thy1-GCaMP6S mice (Strain GP4.3, Jax Labs) (N=16 stroke

(average age: 5.4 months), 5 sham (average age: 6 months)) were used in this study. Mice were

group housed in standard laboratory cages in a temperature-controlled room (23°C), maintained

on a 12 hr light/dark cycle, and given standard laboratory diet and water *ad libitum*. Cages were

randomly assigned to stroke or sham groups such that stroke and sham mice did not co-habitate

the same cages. Animal weight was monitored daily during the entirety of the experiment. All

experiments were approved by the University of Alberta's Health Sciences Animal Care and Use

Committee and adhered to the guidelines set by the Canadian Council for Animal Care. Animals

were euthanized through decapitation under deep urethane anesthesia following the end of the

final imaging session.

Chronic cranial window implantation

Mice were implanted with a chronic cranial window [126] four weeks prior to the first imaging time point (Fig 1B). A surgical plane of anesthesia was achieved with 1.5% isofluorane. Body temperature was measured using a rectal probe and maintained at $37\pm0.5^{\circ}$C. Mice were administered 0.15 mL of saline subcutaneously to maintain hydration levels. Dexamethasone (2 ug/g) was given subcutaneously to prevent cortical swelling and hemorrhaging during the craniotomy procedure. The skull was exposed by midline scalp incision and the skin retracted. The skull was gently scraped with a scalpel to remove the periosteum. Grooves were scraped into the skull surface to improve adhesion of dental cement to the skull. A 4 x 4 mm region of the skull overlying the right hemisphere somatosensory region was thinned to 25-50% of original thickness using a high-speed dental drill (~1-5mm lateral, +2 to -2 mm posterior to bregma). A dental drill was used to progressively thin the overlying skull until the bone could be removed with forceps, leaving the dura intact. The exposed cortical surface was bathed in sterile saline solution. A 5mm diameter coverslip was held in place over the craniotomy and its edges attached to the skull using cyanoacrylate glue. A metal headplate was positioned and secured to the skull using dental cement. Animals were injected subcutaneously with buprenorphine (1.0 mg/kg), removed from the isofluorane, and allowed to recovery in a temperature-controlled recovery cage. Mice were returned to their home cage once recovered and monitored daily for weight and post-surgical signs for the duration of the chronic experiment. Mice were allowed to recover for a period of two weeks after window implantation prior to beginning behavior and task habituation (Fig 1C). Animals were excluded at the 2 week post-implantation timepoint if their cranial window became cloudy or non-imageable.

Tapered beam task habituation, testing and analysis

Methods for tapered beam habituation and automated recording and analysis have been

described previously [53], with minor modification as follows. Nesting material from the

homecage was placed inside of a dark box at the narrow end of the beam to motivate animals to

cross the beam. For the first three sessions, mice were placed at the wide end of the beam and

allowed to freely explore the beam for a period of 2 minutes, after which the experimenter placed

a sheet of paper behind the animal as it crossed the beam to block its return path to the wide end

and motivate it to cross to the narrow end with dark box. Once the mouse had reached the dark

box with bedding material at the narrow end, they were given 60 seconds within the dark box to

associate crossing the beam with reaching the safety of the dark box. The dark box was

transported to their respective cage for the animal to further associate reaching the dark box with

a safe transition to their cage. On subsequent days, mice were continuously run through the

tapered beam for a period of 2 mins each, with a return to their cage after each crossing. On each

of the 3 days prior to the first Ca^{2+} imaging session, mice were tested with three crossings of the

beam per day to determine baseline crossing performance. Left and right side slips were captured

automatically by Raspberry Pi computer attached to touch sensors on the tapered beam, and

Python scripts run to determine number of slips for each side and distance to first slip.

Performance on the 9 baseline trials were average to determine a singular average baseline

performance level. Mice were tested with three trials on each day prior to each weekly post-

imaging timepoint to measure changes in performance.

String pull task habituation, testing and analysis

Methods for string pull habituation and testing have previously been described [127,128], with modification as follows. For habituation, mice were individually placed in a transparent rectangular cage without bedding and allowed to freely explore for a period of 5 minutes. 20 strings with variable length were hung over the edge of the cage. Half of the strings were baited with chocolate flavored sucrose pellets. Mice were removed from the apparatus once all of the strings had been pulled into the cage, or once 20 minutes had elapsed. On subsequent days 2 and 3, mice were again placed in the cage with 20 strings to pull. Following this 3-day habituation, from days 4-14 mice were habituated in an alternate transparent string pull box with high sides and transparent front face for video recording. Within this second habituation box, mice received 3 trials with baited strings hung facing the recording camera. The session was terminated when the animal had pulled all three strings or once 20 minutes had elapsed, and the apparatus prepared for the next mouse. On the last day prior to the first Ca^{2+} imaging session, 3 trials of string pull from each mouse were recorded with the use of a GoPro Hero 7 Black (60 fps, 1920×1080 pixels). Mice were re-tested with this same 3 string protocol the day prior to each of their weekly imaging timepoints. Video recordings were analyzed using a semi-automated Matlab string pull package [128]. Reach and withdraw movement scaling for the left and right paw was calculated by running a Pearson's correlation between the series of Euclidian distances that the paw travelled during each of the reach or withdraw movements during the string pull and correlating with the peak speed of the paw during each of the respective reaches or withdraws [129]. A value of 1 indicates a strong linear relationship between longer reach/withdraw paw movement distance and greater peak speed that the paw obtained. Paw reach distance was calculated as the average distance that the paw travelled during each reach movement. Path

circuity is used as a measure of how direct of a path a reach or withdraw movement takes

between its starting and end points and is calculated as the total distance travelled divided by the

Euclidian distance between the start and end points [129]. The greater the path circuity value is

above 1, the more the path of the paw deviated from the ideal direct path between the start and

end points of the reach/withdraw. A bimanual correlation coefficient was calculated by first

detrending the timeseries of Y-axis paw position to remove general changes in the posture of the

animal over the course of the string pull attempt, then running a Pearson's correlation between

the Y-axis paw position measurements for the right and left paw.

Mobile floating homecage habituation and measurement of movement parameters

Methods for floating homecage (Neurotar, Finland) habituation have previously been

described [130], with modification as follows. Two weeks after cranial window implantation,

each mouse was handled for 5 minutes 3 times per day for the first 2 days to habituate them to

handling. For days 3-4, each mouse was handled and repeatedly wrapped and released with a soft

cloth for a period of 5 mins 3 times per day in order to habituate to the wrapping procedure. Mice

were also given 1 period of 5 minutes each per day to freely explore the floating homecage

without being head-restrained. For days 5-8, twice per day the animals were head-restrained in

the floating homecage and allowed to move around the floating homecage for a period of 15

minutes with the room lights off before being returned to their cage. For days 8-14, twice per day

the mice were head-restrained in the floating homecage for 25 minutes with the floating

homecage attached to the microscope in the same conditions in which the animals would be

imaged on day 15 onwards. On day 14 (one day prior to their first imaging session) baseline

performance on the string pull and tapered beam tasks was measured prior to their last

habituation in the floating homecage. Animal movement within the homecage during each Ca^{2+}

imaging sessions was tracked to determine animal speed and position. Animal speed was

thresholded at 30mm/s to determine periods of time in which the animal was moving and periods

of rest were determined as speed below 30mm/s.

Identification of forelimb and hindlimb somatosensory cortex representations using

widefield Ca^{2+} imaging of sensory-evoked responses

On day 10 of habituation, mice were anesthetized with 1.25% isofluorane after they had

completed all daily task training and floating homecage habituation. Mice were head-fixed on a

stereotaxic frame with continuous isofluorane anesthesia. Body temperature was measured using

a rectal probe and maintained at 37±0.5°C. The cortical surface was illuminated with a xenon

lamp (excitation band-pass filtered at 450-490nm). Fluorescent emissions were long-pass filtered

at 515nm and captured in 12-bit format by a Dalsa Pantera 1M60 camera mounted on a Leica

SP5 confocal microscope using a 2.5X objective. The depth of focus was set at 200 μm below

the cortical surface. Custom-made piezoceramic mechanical bending actuators were used to elicit

oscillatory limb stimulation during widefield Ca^{2+} imaging. The piezo bending actuators

comprised a piezo element (Piezo Systems # Q220-A4-203YB) attached to an electrically

insulated metal shaft holding a metal U-bend at its end. The palm of the mouse paw was placed

within the U-bend, and the U-bend bent to shape to lightly secure the palm. The metal U-bend

made contact across a vertical rectangular area of approximately 3x1mm on the palmar and

dorsal surface of the hand. Stimulators were driven with square-wave signals from an A-M

Systems Model 2100 Isolated Pulse Stimulator. Stimulation alternated between contralateral

forelimb (cFL) and contralateral hindlimb (cHL) for up to 40 trials of stimulation of each limb.

Placement of actuators was on the glabrous skin of the forepaw or hindpaw, with consistent

alignment relative to the flexion of wrist and ankle. Images were captured for 5.0s at 10 Hz (1s

before and 4s after stimulus onset; interstimulus interval = 20s). The trials for each limb were

averaged in ImageJ software (NIH). 10 imaging frames (1s) after stimulus onset were averaged

and divided by the 10 baseline frames 1s before stimulus onset to generate a response map for

each limb. Response maps were thresholded at 95% maximal response to determine limb

associated response boundaries, merged, and overlaid on an image of surface vasculature to

delineate the cFL and cHL somatosensory areas.

Calcium imaging

On the day following testing for behavioral performance on the string pull and tapered

beam tasks, animals were head-fixed within the mobile homecage and Ca^{2+} imaging was

performed. Awake in-vivo two-photon imaging was performed using a Leica SP5 MP confocal

microscope equipped with a Ti:Sapphire Coherent Chameleon Vision II laser tuned to 940 nm

for GCaMP6S excitation. A Leica HCX PL APO L 20x 1.0NA water immersion objective was

used. Images were acquired using Leica LAS AF using no averaging, a zoom of 1.7x and a

frame-rate of 25Hz. Images were acquired at 512x512 pixels over an area of 434x434μm,

yielding a resolution of 0.848μm per pixel. Two imaging regions were chosen for each animal,

with one corresponding to the half-way point between the pre-stroke cFL and cHL

somatosensory maps previously determined with widefield Ca^{2+} imaging (the "peri-infarct"

imaging region), and the other an imaging region located just lateral to the cHL sensory map (the

"distal" imaging region). These regions can be seen in Fig 1D and Fig 2A. Each imaging region

was recorded for 15 minutes while simultaneously tracking the animal's movement and position

within the mobile homecage. Imaging depth was set between 100-180um below the cortical

surface. There were no differences between the stroke group and sham group for movement

within the mobile homecage, though both groups displayed an equivalent decrease in the amount

of time they spent moving within the 15 minute recording sessions over the course of the 8 week

experimental timeline, potentially reflecting continuing habituation over this period.

Photothrombotic stroke and sham procedure

After baseline cellular Ca^{2+} imaging, mice were anesthetized using 1.25% isofluorane

and head-fixed on a stereotaxic frame with continuous isofluorane anesthesia. Body temperature

was measured using a rectal probe and maintained at 37±0.5ºC. The mapped cFL area was used

as a guide for a targeted photothrombosis procedure. Briefly, a 2mm diameter hole was punched

in black electrical tape and attached onto the glass window to block stray light from illuminating

cortical areas outside of the desired region. Rose Bengal (a photosensitive dye) was dissolved in

0.01M sterile phosphate buffered saline (Sigma) and injected intraperitoneal (30mg/kg). The cFL

cortical area visible through the punched electrical tape was illuminated using a collimated beam

of green laser light (532 nm, 17mW) for 20 minutes to photoactivate the Rose Bengal and cause

a focal ischemic lesion. Sham controls were treated in the same manner as stroke, however

illumination of the laser was omitted.

Two-photon image processing and analysis of neuronal activity

Timeseries images (15 minutes per imaging region, 22500 frames per recording, 25fps)

were group averaged in groups of 2, then motion corrected using the TurboReg plugin in FiJi

with translation only registration [131]. Z-projections within FiJi of the average and standard

deviation were used to define neuronal ROI through manual tracing. Using custom written

scripts in Matlab 2020b, ROIs were imported and converted into a format suitable for the Ca^{2+}

imaging toolbox used for subsequent steps [59]. Neuropil for each ROI was determined by

expanding an annular donut around each ROI to calculate the neuropil deltaF/Fo surrounding

each ROI. During computation of neuron ROI deltaF/Fo, fluorescence was corrected for neuropil

contamination using the formula ($F_{corrected} = F_{raw} - \alpha * F_{neuropil}$) where α was set to 0.4 based on

previous studies suggesting that α values between 0.3-0.5 were optimal [132], and with $F_{neuropil}$

determined by the peri-somatic donut neuropil fluorescence closely associated with each neuron.

A smoothed estimate of the baseline fluorescence was calculated by taking a 30s running average

of the 8^{th} percentile of the raw fluorescence, which was then subtracted from the raw

fluorescence to remove baseline drift.

Ca^{2+} trace deconvolution to determine neuronal transients

To determine significant fluorescent Ca^{2+} transients from the deltaF/Fo of each neuron, a

dynamic threshold implementing a Bayesian odds ratio estimation framework for noise

estimation was used to determine transients that met the condition of being greater than 98% of

the confidence interval for the calculated fluorescent baseline noise, and were compatible with a

tau of ~1.8s for GCaMP6S [58,59]. Determination of firing was performed on the neuronal

deltaF/Fo traces, as implemented in [58,59], in order to define the Ca^{2+} transient rate and to

generate raster plots. This information was used to determine neuronal activity "firing rate" as

shown in Fig 4.

Neuron-neuron correlation analysis and functional connectivity

Using custom written scripts in Matlab, Pearson product-moment correlation coefficients were calculated between the z-scored Ca^{2+} traces derived from each of the neuronal ROIs in a given optical section. Distance between neuron pairs was calculated from the Euclidean distance between the central points of the neuron ROIs of the image frame. Functional connectivity plots (Fig 5) were generated by plotting neuron ROI centroids as red dots, and lines between them with line weight and color determined by the strength of the correlation between them. Stationary variable block bootstrapping (5000 iterations) was performed as a statistical test for the significance of each pairwise correlation and only correlations that were greater than the 99th percentile of the bootstrap were deemed statistically significant and plotted. Average number of significant connections per neuron was calculated by dividing the number of functional connections that met the bootstrapping threshold by the total number of neurons in the population. Connection density was calculated as percent of max by taking the total number of significant functional connections and dividing by the total potential number of functional connections if every neuron within the population was functionally connected with all other neurons.

Determination of neural assemblies and their activity patterns

To evaluate the co-activity patterns of neurons that form putative neural assemblies, a PCA-Promax procedure was applied as previously described [58,59], with a zMax threshold manually selected by the first clear minimum in the distribution of z-scored maximal ROI loadings. Notably, the PCA-Promax procedure relaxes the PCA orthogonality condition using a PROMAX oblique rotation of the PC axes [133], such that assemblies can contain neurons found

within other assemblies. Due to this relaxation of the PCA orthogonality condition, assemblies

with high overlap whose dot product exceeded 0.6 were merged [133]. All assemblies were

compared to surrogate control datasets, and only those assemblies whose members were

significantly correlated and synchronous were kept ($p < 0.05$). To determine the co-activity of

neurons within assemblies, a matching index was calculated [58,59,134,135]. The matching

index quantifies the proportion of neurons within the assembly that are co-activated

simultaneously over the timeseries, with a maximal value of 1 indicating perfect overlap in

assembly member activation. The significance of each assembly activation over the timeseries

was determined by comparing the activation events to a probability distribution based on the size

of the assembly, the size of the total population of N ROIs, and with a threshold p-value of <0.05

considered to be a statistically significant activation. For cross-correlation of assembly

activations with speed (Fig S3) and assembly-assembly correlations (Fig 9), the matching index

was first multiplied by the Boolean timeseries of individual assembly activation significance to

generate matching index timeseries of significant assembly activations only (presented in Fig S3

and Fig 9 as percent of maximal assembly activation). ROIs of assembly members were color

coded according to their assembly membership, and overlaid on an averaged fluorescence image

to generate assembly plots (Fig 7).

Statistical analysis

Multivariate comparisons were made using a mixed-effects model for repeated measures

based on a restricted maximum likelihood generalized linear mixed model as implemented in

Graphpad Prism 9.0.0, with Bonferroni-Sidak corrections used for *post-hoc* comparisons.

Bonferroni-Sidak *post-hoc* testing was used to compare the means of the stroke vs. the sham

group at each timepoint, as well as to identify within group differences by comparing different

timepoints within each group. Normalized data was analyzed using multivariate mixed-effects

models based on a restricted maximum likelihood generalized linear mixed model, with the "pre"

timepoint removed from calculation of main effects, interaction, and *post-hoc* statistical testing.

Grubb's test was used to remove data outliers with alpha set to 0.05. All zero time-lag cross-

correlations were computed using Pearson's *r*. A stationary bootstrapping procedure was used as

a test of the significance of the calculated Pearson's r of each pairwise timeseries comparison.

Within the stationary bootstrapping procedure, 5000 iterations were run and an average block

length of 23 frames (1.84s) was used based on previous studies indicating ~1.8s as the decay

time for GCaMP6S Ca^{2+} sensor within neurons [136,137]. Pairwise r values greater than the 99th

percentile of the stationary bootstrap were deemed significant. Non-significant pairwise r values

were excluded from analyses of average correlation values. For all statistical comparisons, a *p*

value of ≤0.05 was considered statistically significant. A *p* value of ≤0.10 was considered a

statistical trend. Data are expressed as the mean ± SEM.

Author Contributions

693   MVB and IRW designed all experiments. MVB performed experiments and analyzed

694   data. MVB and IRW wrote the manuscript together.

Competing Interests

697   The authors declare no competing interests.

Data Availability

All data generated or analyzed during this study will be made publicly available on OSF (https://osf.io/wam9j/).

REFERENCES

1. Schiene K, Bruehl C, Zilles K, Qü M, Hagemann G, Kraemer M, et al. Neuronal hyperexcitability and reduction of GABAA-receptor expression in the surround of cerebral photothrombosis. J Cereb Blood Flow Metab Off J Int Soc Cereb Blood Flow Metab. 1996;16: 906–914. doi:10.1097/00004647-199609000-00014

2. Mittmann T, Qü M, Zilles K, Luhmann HJ. Long-term cellular dysfunction after focal cerebral ischemia: in vitro analyses. Neuroscience. 1998;85: 15–27. doi:10.1016/S0306-4522(97)00638-6

3. Neumann-Haefelin T, Staiger JF, Redecker C, Zilles K, Fritschy JM, Möhler H, et al. Immunohistochemical evidence for dysregulation of the GABAergic system ipsilateral to photochemically induced cortical infarcts in rats. Neuroscience. 1998;87: 871–879. doi:10.1016/s0306-4522(98)00124-9

4. Neumann-Haefelin T, Bosse F, Redecker C, Müller HW, Witte OW. Upregulation of GABAA-receptor alpha1- and alpha2-subunit mRNAs following ischemic cortical lesions in rats. Brain Res. 1999;816: 234–237. doi:10.1016/s0006-8993(98)01162-7

5. Qü M, Buchkremer-Ratzmann I, Schiene K, Schroeter M, Witte OW, Zilles K. Bihemispheric reduction of GABAA receptor binding following focal cortical photothrombotic lesions in the rat brain. Brain Res. 1998;813: 374–380. doi:10.1016/s0006-8993(98)01063-4

6. Redecker C, Wang W, Fritschy J-M, Witte OW. Widespread and Long-Lasting Alterations in GABAA-Receptor Subtypes after Focal Cortical Infarcts in Rats: Mediation by NMDA-Dependent Processes. J Cereb Blood Flow Metab. 2002;22: 1463–1475. doi:10.1097/01.WCB.0000034149.72481.BD

7. Luhmann HJ, Mudrick-Donnon LA, Mittmann T, Heinemann U. Ischaemia-induced long-term hyperexcitability in rat neocortex. Eur J Neurosci. 1995;7: 180–191. doi:10.1111/j.1460-9568.1995.tb01054.x

8. Buchkremer-Ratzmann I, Witte OW. Extended brain disinhibition following small photothrombotic lesions in rat frontal cortex. NeuroReport. 1997;8: 519–522.

9.   Domann R, Hagemann G, Kraemer M, Freund H-J, Witte OW. Electrophysiological
changes in the surrounding brain tissue of photochemically induced cortical infarcts in the
rat. Neurosci Lett. 1993;155: 69–72. doi:10.1016/0304-3940(93)90675-B

10.   Fujioka H, Kaneko H, Suzuki SS, Mabuchi K. Hyperexcitability-associated rapid
plasticity after a focal cerebral ischemia. Stroke. 2004;35: e346-348.
doi:10.1161/01.STR.0000130990.28734.9c

11.   Qü M, Mittmann T, Luhmann HJ, Schleicher A, Zilles K. Long-term changes of
ionotropic glutamate and GABA receptors after unilateral permanent focal cerebral
ischemia in the mouse brain. Neuroscience. 1998;85: 29–43. doi:10.1016/s0306-
4522(97)00656-8

12.   Que M, Schiene K, Witte OW, Zilles K. Widespread up-regulation of N-methyl-D-
aspartate receptors after focal photothrombotic lesion in rat brain. Neurosci Lett.
1999;273: 77–80. doi:10.1016/s0304-3940(99)00598-4

13.   Jaenisch N, Liebmann L, Guenther M, Hübner CA, Frahm C, Witte OW. Reduced tonic
inhibition after stroke promotes motor performance and epileptic seizures. Sci Rep.
2016;6: 26173. doi:10.1038/srep26173

14.   Jin X, Huguenard JR, Prince DA. Impaired Cl- extrusion in layer V pyramidal neurons of
chronically injured epileptogenic neocortex. J Neurophysiol. 2005;93: 2117–2126.
doi:10.1152/jn.00728.2004

15.   Khirug S, Soni S, Saez Garcia M, Tessier M, Zhou L, Kulesskaya N, et al. Protective Role
of Low Ethanol Administration Following Ischemic Stroke via Recovery of KCC2 and
p75NTR Expression. Mol Neurobiol. 2021;58: 1145–1161. doi:10.1007/s12035-020-
02176-x

16.   Martín-Aragón Baudel MAS, Poole AV, Darlison MG. Chloride co-transporters as
possible therapeutic targets for stroke. J Neurochem. 2017;140: 195–209.
doi:10.1111/jnc.13901

17.   Schulte JT, Wierenga CJ, Bruining H. Chloride transporters and GABA polarity in
developmental, neurological and psychiatric conditions. Neurosci Biobehav Rev. 2018;90:
260–271. doi:10.1016/j.neubiorev.2018.05.001

18.   Witte OW, Freund HJ. Neuronal dysfunction, epilepsy, and postlesional brain plasticity.
Adv Neurol. 1999;81: 25–36.

19.   Brown CE, Aminoltejari K, Erb H, Winship IR, Murphy TH. In Vivo Voltage-Sensitive
Dye Imaging in Adult Mice Reveals That Somatosensory Maps Lost to Stroke Are
Replaced over Weeks by New Structural and Functional Circuits with Prolonged Modes of
Activation within Both the Peri-Infarct Zone and Distant Sites. J Neurosci. 2009;29:
1719–1734. doi:10.1523/JNEUROSCI.4249-08.2009

20.   Chen S, Mohajerani MH, Xie Y, Murphy TH. Optogenetic analysis of neuronal
769   excitability during global ischemia reveals selective deficits in sensory processing
770   following reperfusion in mouse cortex. J Neurosci Off J Soc Neurosci. 2012;32: 13510–
771   13519. doi:10.1523/JNEUROSCI.1439-12.2012

772   21.   Lim DH, LeDue JM, Mohajerani MH, Murphy TH. Optogenetic mapping after stroke
773   reveals network-wide scaling of functional connections and heterogeneous recovery of the
774   peri-infarct. J Neurosci Off J Soc Neurosci. 2014;34: 16455–16466.
775   doi:10.1523/JNEUROSCI.3384-14.2014

776   22.   Sigler A, Mohajerani MH, Murphy TH. Imaging rapid redistribution of sensory-evoked
777   depolarization through existing cortical pathways after targeted stroke in mice. Proc Natl
778   Acad Sci. 2009;106: 11759–11764.

779   23.   Sweetnam DA, Brown CE. Stroke induces long-lasting deficits in the temporal fidelity of
780   sensory processing in the somatosensory cortex. J Cereb Blood Flow Metab Off J Int Soc
781   Cereb Blood Flow Metab. 2013;33: 91–96. doi:10.1038/jcbfm.2012.135

782   24.   Sweetnam D, Holmes A, Tennant KA, Zamani A, Walle M, Jones P, et al. Diabetes
783   Impairs Cortical Plasticity and Functional Recovery Following Ischemic Stroke. J
784   Neurosci. 2012;32: 5132–5143. doi:10.1523/JNEUROSCI.5075-11.2012

785   25.   Winship IR, Murphy TH. In vivo calcium imaging reveals functional rewiring of single
786   somatosensory neurons after stroke. J Neurosci Off J Soc Neurosci. 2008;28: 6592–6606.
787   doi:10.1523/JNEUROSCI.0622-08.2008

788   26.   Cramer JV, Gesierich B, Roth S, Dichgans M, Düring M, Liesz A. In vivo widefield
789   calcium imaging of the mouse cortex for analysis of network connectivity in health and
790   brain disease. NeuroImage. 2019;199: 570–584. doi:10.1016/j.neuroimage.2019.06.014

791   27.   Grandjean J, Schroeter A, Batata I, Rudin M. Optimization of anesthesia protocol for
792   resting-state fMRI in mice based on differential effects of anesthetics on functional
793   connectivity patterns. NeuroImage. 2014;102: 838–847.
794   doi:10.1016/j.neuroimage.2014.08.043

795   28.   Jonckers E, Delgado y Palacios R, Shah D, Guglielmetti C, Verhoye M, Van der Linden
796   A. Different anesthesia regimes modulate the functional connectivity outcome in mice.
797   Magn Reson Med. 2014;72: 1103–1112. doi:10.1002/mrm.24990

798   29.   Kalthoff D, Po C, Wiedermann D, Hoehn M. Reliability and spatial specificity of rat brain
799   sensorimotor functional connectivity networks are superior under sedation compared with
800   general anesthesia. NMR Biomed. 2013;26: 638–650. doi:10.1002/nbm.2908

801   30.   Nasrallah FA, Tay H-C, Chuang K-H. Detection of functional connectivity in the resting
802   mouse brain. NeuroImage. 2014;86: 417–424. doi:10.1016/j.neuroimage.2013.10.025

31.   Chapin JK, Lin CS. Mapping the body representation in the SI cortex of anesthetized and

awake rats. J Comp Neurol. 1984;229: 199–213. doi:10.1002/cne.902290206

32. Chen LM, Friedman RM, Roe AW. Optical imaging of SI topography in anesthetized and
awake squirrel monkeys. J Neurosci Off J Soc Neurosci. 2005;25: 7648–7659.
doi:10.1523/JNEUROSCI.1990-05.2005

33. Devonshire IM, Grandy TH, Dommett EJ, Greenfield SA. Effects of urethane anaesthesia
on sensory processing in the rat barrel cortex revealed by combined optical imaging and
electrophysiology. Eur J Neurosci. 2010;32: 786–797. doi:10.1111/j.1460-
9568.2010.07322.x

34. Lissek T, Obenhaus HA, Ditzel DAW, Nagai T, Miyawaki A, Sprengel R, et al. General
Anesthetic Conditions Induce Network Synchrony and Disrupt Sensory Processing in the
Cortex. Front Cell Neurosci. 2016;10: 64. doi:10.3389/fncel.2016.00064

35. Ebner FF, Kaas JH. Chapter 24 - Somatosensory System. In: Paxinos G, editor. The Rat
Nervous System (Fourth Edition). San Diego: Academic Press; 2015. pp. 675–701.
doi:10.1016/B978-0-12-374245-2.00024-3

36. Friedberg MH, Lee SM, Ebner FF. Modulation of Receptive Field Properties of Thalamic
Somatosensory Neurons by the Depth of Anesthesia. J Neurophysiol. 1999;81: 2243–
2252. doi:10.1152/jn.1999.81.5.2243

37. Faggin BM, Nguyen KT, Nicolelis MA. Immediate and simultaneous sensory
reorganization at cortical and subcortical levels of the somatosensory system. Proc Natl
Acad Sci U S A. 1997;94: 9428–9433. doi:10.1073/pnas.94.17.9428

38. Moore CI, Nelson SB. Spatio-Temporal Subthreshold Receptive Fields in the Vibrissa
Representation of Rat Primary Somatosensory Cortex. J Neurophysiol. 1998;80: 2882–
2892. doi:10.1152/jn.1998.80.6.2882

39. Nicolelis MA, Chapin JK. Spatiotemporal structure of somatosensory responses of many-
neuron ensembles in the rat ventral posterior medial nucleus of the thalamus. J Neurosci.
1994;14: 3511–3532. doi:10.1523/JNEUROSCI.14-06-03511.1994

40. Crapse TB, Sommer MA. Corollary discharge across the animal kingdom. Nat Rev
Neurosci. 2008;9: 587–600. doi:10.1038/nrn2457

41. Afrashteh N, Inayat S, Bermudez-Contreras E, Luczak A, McNaughton BL, Mohajerani
MH. Spatiotemporal structure of sensory-evoked and spontaneous activity revealed by
mesoscale imaging in anesthetized and awake mice. Cell Rep. 2021;37: 110081.
doi:10.1016/j.celrep.2021.110081

42. Dionne JK, Legon W, Staines WR. Crossmodal influences on early somatosensory
processing: interaction of vision, touch, and task-relevance. Exp Brain Res. 2013;226:
503–512. doi:10.1007/s00221-013-3462-z

43.   Johansen-Berg H, Christensen V, Woolrich M, Matthews PM. Attention to touch
modulates activity in both primary and secondary somatosensory areas. Neuroreport.
2000;11: 1237–1241.

44.   Nelson AJ, Staines WR, Graham SJ, McIlroy WE. Activation in SI and SII: the influence
of vibrotactile amplitude during passive and task-relevant stimulation. Brain Res Cogn
Brain Res. 2004;19: 174–184. doi:10.1016/j.cogbrainres.2003.11.013

45.   Staines WR, Graham SJ, Black SE, McIlroy WE. Task-relevant modulation of
contralateral and ipsilateral primary somatosensory cortex and the role of a prefrontal-
cortical sensory gating system. NeuroImage. 2002;15: 190–199.
doi:10.1006/nimg.2001.0953

46.   Gomez-Ramirez M, Hysaj K, Niebur E. Neural mechanisms of selective attention in the
somatosensory system. J Neurophysiol. 2016;116: 1218–1231. doi:10.1152/jn.00637.2015

47.   Scaglione A, Foffani G, Moxon KA. Spike count, spike timing and temporal information
in the cortex of awake, freely moving rats. J Neural Eng. 2014;11: 046022.
doi:10.1088/1741-2560/11/4/046022

48.   Yu J, Gutnisky DA, Hires SA, Svoboda K. Layer 4 fast-spiking interneurons filter
thalamocortical signals during active somatosensation. Nat Neurosci. 2016;19: 1647–
1657. doi:10.1038/nn.4412

49.   London BM, Miller LE. Responses of somatosensory area 2 neurons to actively and
passively generated limb movements. J Neurophysiol. 2013;109: 1505–1513.
doi:10.1152/jn.00372.2012

50.   Nelson RJ. Activity of monkey primary somatosensory cortical neurons changes prior to
active movement. Brain Res. 1987;406: 402–407. doi:10.1016/0006-8993(87)90815-8

51.   Bensmaia S, Helms Tillery SI. Tactile Feedback from the Hand. In: Balasubramanian R,
Santos VJ, editors. The Human Hand as an Inspiration for Robot Hand Development.
Cham: Springer International Publishing; 2014. pp. 143–157. doi:10.1007/978-3-319-
03017-3_7

52.   Winship IR, Murphy TH. Remapping the somatosensory cortex after stroke: insight from
imaging the synapse to network. Neurosci Rev J Bringing Neurobiol Neurol Psychiatry.
2009;15: 507–524. doi:10.1177/1073858409333076

53.   Ardesch DJ, Balbi M, Murphy TH. Automated touch sensing in the mouse tapered beam
test using Raspberry Pi. J Neurosci Methods. 2017;291: 221–226.
doi:10.1016/j.jneumeth.2017.08.030

54.   Zhao C, Puurunen K, Schallert T, Sivenius J, Jolkkonen J. Effect of cholinergic
medication, before and after focal photothrombotic ischemic cortical injury, on
histological and functional outcome in aged and young adult rats. Behav Brain Res.

2005;156: 85–94. doi:10.1016/j.bbr.2004.05.011

55.   Lipsanen A, Hiltunen M, Jolkkonen J. Chronic ibuprofen treatment does not affect the
secondary pathology in the thalamus or improve behavioral outcome in middle cerebral
artery occlusion rats. Pharmacol Biochem Behav. 2011;99: 468–474.
doi:10.1016/j.pbb.2011.04.019

56.   Schallert T, Woodlee MT, Fleming SM. Disentangling multiple types of recovery from
brain injury. Pharmacol Cereb Ischemia. 2002; 201–216.

57.   Blackwell AA, Widick WL, Cheatwood JL, Whishaw IQ, Wallace DG. Unilateral
forelimb sensorimotor cortex devascularization disrupts the topographic and kinematic
characteristics of hand movements while string-pulling for food in the rat. Behav Brain
Res. 2018;338: 88–100. doi:10.1016/j.bbr.2017.10.014

58.   Romano SA, Pietri T, Pérez-Schuster V, Jouary A, Haudrechy M, Sumbre G. Spontaneous
Neuronal Network Dynamics Reveal Circuit's Functional Adaptations for Behavior.
Neuron. 2015;85: 1070–1085. doi:10.1016/j.neuron.2015.01.027

59.   Romano SA, Pérez-Schuster V, Jouary A, Boulanger-Weill J, Candeo A, Pietri T, et al. An
integrated calcium imaging processing toolbox for the analysis of neuronal population
dynamics. PLoS Comput Biol. 2017;13: e1005526. doi:10.1371/journal.pcbi.1005526

60.   Bauer AQ, Kraft AW, Wright PW, Snyder AZ, Lee J-M, Culver JP. Optical imaging of
disrupted functional connectivity following ischemic stroke in mice. NeuroImage.
2014;99: 388–401. doi:10.1016/j.neuroimage.2014.05.051

61.   Blaschke SJ, Hensel L, Minassian A, Vlachakis S, Tscherpel C, Vay SU, et al. Translating
Functional Connectivity After Stroke: Functional Magnetic Resonance Imaging Detects
Comparable Network Changes in Mice and Humans. Stroke. 2021;52: 2948–2960.
doi:10.1161/STROKEAHA.120.032511

62.   Carter AR, Astafiev SV, Lang CE, Connor LT, Rengachary J, Strube MJ, et al. Resting
interhemispheric functional magnetic resonance imaging connectivity predicts
performance after stroke. Ann Neurol. 2010;67: 365–375. doi:10.1002/ana.21905

63.   Carter AR, Shulman GL, Corbetta M. Why use a connectivity-based approach to study
stroke and recovery of function? NeuroImage. 2012;62: 2271–2280.
doi:10.1016/j.neuroimage.2012.02.070

64.   Hakon J, Quattromani MJ, Sjölund C, Tomasevic G, Carey L, Lee J-M, et al. Multisensory
stimulation improves functional recovery and resting-state functional connectivity in the
mouse brain after stroke. NeuroImage Clin. 2017;17: 717–730.
doi:10.1016/j.nicl.2017.11.022

65.   Longo V, Barbati SA, Re A, Paciello F, Bolla M, Rinaudo M, et al. Transcranial Direct
Current Stimulation Enhances Neuroplasticity and Accelerates Motor Recovery in a

Stroke Mouse Model. Stroke. 2022; STROKEAHA121034200.
doi:10.1161/STROKEAHA.121.034200

66.   Olafson ER, Jamison KW, Sweeney EM, Liu H, Wang D, Bruss JE, et al. Functional
connectome reorganization relates to post-stroke motor recovery and structural and
functional disconnection. NeuroImage. 2021;245: 118642.
doi:10.1016/j.neuroimage.2021.118642

67.   Rehme AK, Grefkes C. Cerebral network disorders after stroke: evidence from imaging-
based connectivity analyses of active and resting brain states in humans. J Physiol.
2013;591: 17–31. doi:10.1113/jphysiol.2012.243469

68.   Siegel JS, Ramsey LE, Snyder AZ, Metcalf NV, Chacko RV, Weinberger K, et al.
Disruptions of network connectivity predict impairment in multiple behavioral domains
after stroke. Proc Natl Acad Sci U S A. 2016;113: E4367–E4376.
doi:10.1073/pnas.1521083113

69.   Silasi G, Murphy TH. Stroke and the Connectome: How Connectivity Guides Therapeutic
Intervention. Neuron. 2014;83: 1354–1368. doi:10.1016/j.neuron.2014.08.052

70.   Urbin MA, Hong X, Lang CE, Carter AR. Resting-state functional connectivity and its
association with multiple domains of upper-extremity function in chronic stroke.
Neurorehabil Neural Repair. 2014;28: 761–769. doi:10.1177/1545968314522349

71.   Latifi S, Mitchell S, Habibey R, Hosseini F, Donzis E, Estrada-Sánchez AM, et al.
Neuronal Network Topology Indicates Distinct Recovery Processes after Stroke. Cereb
Cortex N Y NY. 2020;30: 6363–6375. doi:10.1093/cercor/bhaa191

72.   Bechay KR, Abduljawad N, Latifi S, Suzuki K, Iwashita H, Carmichael ST. PDE2A
Inhibition Enhances Axonal Sprouting, Functional Connectivity, and Recovery after
Stroke. J Neurosci. 2022;42: 8225–8236. doi:10.1523/JNEUROSCI.0730-22.2022

73.   Buzsáki G. Neural syntax: cell assemblies, synapsembles, and readers. Neuron. 2010;68:
362–385. doi:10.1016/j.neuron.2010.09.023

74.   Buzsáki G, Watson BO. Brain rhythms and neural syntax: implications for efficient coding
of cognitive content and neuropsychiatric disease. Dialogues Clin Neurosci. 2012;14:
345–367. doi:10.31887/DCNS.2012.14.4/gbuzsaki

75.   Palm G, Knoblauch A, Hauser F, Schüz A. Cell assemblies in the cerebral cortex. Biol
Cybern. 2014;108: 559–572. doi:10.1007/s00422-014-0596-4

76.   Mölter J, Avitan L, Goodhill GJ. Detecting neural assemblies in calcium imaging data.
BMC Biol. 2018;16: 143. doi:10.1186/s12915-018-0606-4

77.   Deolindo CS, Kunicki ACB, da Silva MI, Lima Brasil F, Moioli RC. Neuronal Assemblies
Evidence Distributed Interactions within a Tactile Discrimination Task in Rats. Front

Neural Circuits. 2017;11: 114. doi:10.3389/fncir.2017.00114

78.   Murphy TH, Corbett D. Plasticity during stroke recovery: from synapse to behaviour. Nat
Rev Neurosci. 2009;10: 861–872. doi:10.1038/nrn2735

79.   Xu X, Ye L, Ruan Q. Environmental enrichment induces synaptic structural modification
after transient focal cerebral ischemia in rats. Exp Biol Med Maywood NJ. 2009;234:
296–305. doi:10.3181/0804-RM-128

80.   Biernaskie J, Chernenko G, Corbett D. Efficacy of rehabilitative experience declines with
time after focal ischemic brain injury. J Neurosci Off J Soc Neurosci. 2004;24: 1245–
1254. doi:10.1523/JNEUROSCI.3834-03.2004

81.   Brown CE, Li P, Boyd JD, Delaney KR, Murphy TH. Extensive turnover of dendritic
spines and vascular remodeling in cortical tissues recovering from stroke. J Neurosci Off J
Soc Neurosci. 2007;27: 4101–4109. doi:10.1523/JNEUROSCI.4295-06.2007

82.   Carmichael ST, Archibeque I, Luke L, Nolan T, Momiy J, Li S. Growth-associated gene
expression after stroke: evidence for a growth-promoting region in peri-infarct cortex. Exp
Neurol. 2005;193: 291–311. doi:10.1016/j.expneurol.2005.01.004

83.   Cheatwood JL, Emerick AJ, Kartje GL. Neuronal plasticity and functional recovery after
ischemic stroke. Top Stroke Rehabil. 2008;15: 42–50. doi:10.1310/tsr1501-42

84.   Ploughman M, Windle V, MacLellan CL, White N, Doré JJ, Corbett D. Brain-derived
neurotrophic factor contributes to recovery of skilled reaching after focal ischemia in rats.
Stroke. 2009;40: 1490–1495. doi:10.1161/STROKEAHA.108.531806

85.   Carmichael ST. Plasticity of Cortical Projections after Stroke. The Neuroscientist. 2003;9:
64–75. doi:10.1177/1073858402239592

86.   Carmichael ST. Cellular and molecular mechanisms of neural repair after stroke: Making
waves. Ann Neurol. 2006;59: 735–742. doi:10.1002/ana.20845

87.   Dancause N. Extensive Cortical Rewiring after Brain Injury. J Neurosci. 2005;25: 10167–
10179. doi:10.1523/JNEUROSCI.3256-05.2005

88.   Carmichael ST. Gene expression changes after focal stroke, traumatic brain and spinal
cord injuries. Curr Opin Neurol. 2003;16: 699–704.
doi:10.1097/01.wco.0000102621.38669.77

89.   Li S, Carmichael ST. Growth-associated gene and protein expression in the region of
axonal sprouting in the aged brain after stroke. Neurobiol Dis. 2006;23: 362–373.
doi:10.1016/j.nbd.2006.03.011

90.   Li S, Overman JJ, Katsman D, Kozlov SV, Donnelly CJ, Twiss JL, et al. An age-related
sprouting transcriptome provides molecular control of axonal sprouting after stroke. Nat

Neurosci. 2010;13: 1496–1504. doi:10.1038/nn.2674

91. Paz JT, Christian CA, Parada I, Prince DA, Huguenard JR. Focal cortical infarcts alter
intrinsic excitability and synaptic excitation in the reticular thalamic nucleus. J Neurosci
Off J Soc Neurosci. 2010;30: 5465–5479. doi:10.1523/JNEUROSCI.5083-09.2010

92. Staines WR, Black SE, Graham SJ, McIlroy WE. Somatosensory gating and recovery
from stroke involving the thalamus. Stroke. 2002;33: 2642–2651.
doi:10.1161/01.str.0000032552.40405.40

93. Carmichael ST, Wei L, Rovainen CM, Woolsey TA. New patterns of intracortical
projections after focal cortical stroke. Neurobiol Dis. 2001;8: 910–922.
doi:10.1006/nbdi.2001.0425

94. Tennant KA, Taylor SL, White ER, Brown CE. Optogenetic rewiring of thalamocortical
circuits to restore function in the stroke injured brain. Nat Commun. 2017;8: 15879.
doi:10.1038/ncomms15879

95. Brown CE, Murphy TH. Livin' on the Edge: Imaging Dendritic Spine Turnover in the
Peri-Infarct Zone during Ischemic Stroke and Recovery. The Neuroscientist. 2008;14:
139–146. doi:10.1177/1073858407309854

96. Ergul A, Alhusban A, Fagan SC. Angiogenesis: a harmonized target for recovery after
stroke. Stroke. 2012;43: 2270–2274. doi:10.1161/STROKEAHA.111.642710

97. Greenberg DA. Cerebral angiogenesis: a realistic therapy for ischemic disease? Methods
Mol Biol Clifton NJ. 2014;1135: 21–24. doi:10.1007/978-1-4939-0320-7_2

98. Hatakeyama M, Ninomiya I, Kanazawa M. Angiogenesis and neuronal remodeling after
ischemic stroke. Neural Regen Res. 2020;15: 16–19. doi:10.4103/1673-5374.264442

99. Bullmore E, Sporns O. Complex brain networks: graph theoretical analysis of structural
and functional systems. Nat Rev Neurosci. 2009;10: 186–198. doi:10.1038/nrn2575

100. Sporns O. The non-random brain: efficiency, economy, and complex dynamics. Front
Comput Neurosci. 2011;5: 5. doi:10.3389/fncom.2011.00005

101. Stam CJ. Modern network science of neurological disorders. Nat Rev Neurosci. 2014;15:
683–695. doi:10.1038/nrn3801

102. Berkes P, Orbán G, Lengyel M, Fiser J. Spontaneous cortical activity reveals hallmarks of
an optimal internal model of the environment. Science. 2011;331: 83–87.
doi:10.1126/science.1195870

103. Cossart R, Aronov D, Yuste R. Attractor dynamics of network UP states in the neocortex.
Nature. 2003;423: 283–288. doi:10.1038/nature01614

104.   Harris KD, Csicsvari J, Hirase H, Dragoi G, Buzsáki G. Organization of cell assemblies in
1014   the hippocampus. Nature. 2003;424: 552–556. doi:10.1038/nature01834

1015   105.   Hyman JM, Whitman J, Emberly E, Woodward TS, Seamans JK. Action and outcome
1016   activity state patterns in the anterior cingulate cortex. Cereb Cortex N Y N 1991. 2013;23:
1017   1257–1268. doi:10.1093/cercor/bhs104

1018   106.   Malvache A, Reichinnek S, Villette V, Haimerl C, Cossart R. Awake hippocampal
1019   reactivations project onto orthogonal neuronal assemblies. Science. 2016;353: 1280–1283.
1020   doi:10.1126/science.aaf3319

1021   107.   Miller JK, Ayzenshtat I, Carrillo-Reid L, Yuste R. Visual stimuli recruit intrinsically
1022   generated cortical ensembles. Proc Natl Acad Sci U S A. 2014;111: E4053-4061.
1023   doi:10.1073/pnas.1406077111

1024   108.   Sakata S, Harris KD. Laminar structure of spontaneous and sensory-evoked population
1025   activity in auditory cortex. Neuron. 2009;64: 404–418. doi:10.1016/j.neuron.2009.09.020

1026   109.   See JZ, Atencio CA, Sohal VS, Schreiner CE. Coordinated neuronal ensembles in primary
1027   auditory cortical columns. eLife. 7: e35587. doi:10.7554/eLife.35587

1028   110.   Truccolo W, Hochberg LR, Donoghue JP. Collective dynamics in human and monkey
1029   sensorimotor cortex: predicting single neuron spikes. Nat Neurosci. 2010;13: 105–111.
1030   doi:10.1038/nn.2455

1031   111.   Villette V, Malvache A, Tressard T, Dupuy N, Cossart R. Internally Recurring
1032   Hippocampal Sequences as a Population Template of Spatiotemporal Information.
1033   Neuron. 2015;88: 357–366. doi:10.1016/j.neuron.2015.09.052

1034   112.   Avitan L, Pujic Z, Mölter J, Van De Poll M, Sun B, Teng H, et al. Spontaneous Activity in
1035   the Zebrafish Tectum Reorganizes over Development and Is Influenced by Visual
1036   Experience. Curr Biol CB. 2017;27: 2407-2419.e4. doi:10.1016/j.cub.2017.06.056

1037   113.   Pietri T, Romano SA, Pérez-Schuster V, Boulanger-Weill J, Candat V, Sumbre G. The
1038   Emergence of the Spatial Structure of Tectal Spontaneous Activity Is Independent of
1039   Visual Inputs. Cell Rep. 2017;19: 939–948. doi:10.1016/j.celrep.2017.04.015

1040   114.   Thompson AW, Scott EK. Characterisation of sensitivity and orientation tuning for
1041   visually responsive ensembles in the zebrafish tectum. Sci Rep. 2016;6: 34887.
1042   doi:10.1038/srep34887

1043   115.   Luczak A, Barthó P, Harris KD. Spontaneous events outline the realm of possible sensory
1044   responses in neocortical populations. Neuron. 2009;62: 413–425.
1045   doi:10.1016/j.neuron.2009.03.014

1046   116.   MacLean JN, Watson BO, Aaron GB, Yuste R. Internal dynamics determine the cortical
1047   response to thalamic stimulation. Neuron. 2005;48: 811–823.

doi:10.1016/j.neuron.2005.09.035

117. Carey LM, Matyas TA. Frequency of discriminative sensory loss in the hand after stroke
in a rehabilitation setting. J Rehabil Med. 2011;43: 257–263. doi:10.2340/16501977-0662

118. Connell LA, Lincoln NB, Radford KA. Somatosensory impairment after stroke: frequency
of different deficits and their recovery. Clin Rehabil. 2008;22: 758–767.
doi:10.1177/0269215508090674

119. Kwakkel G, Kollen B, Lindeman E. Understanding the pattern of functional recovery after
stroke: facts and theories. Restor Neurol Neurosci. 2004;22: 281–299.

120. Patel AT, Duncan PW, Lai SM, Studenski S. The relation between impairments and
functional outcomes poststroke. Arch Phys Med Rehabil. 2000;81: 1357–1363.
doi:10.1053/apmr.2000.9397

121. Tyson SF, Hanley M, Chillala J, Selley AB, Tallis RC. Sensory loss in hospital-admitted
people with stroke: characteristics, associated factors, and relationship with function.
Neurorehabil Neural Repair. 2008;22: 166–172. doi:10.1177/1545968307305523

122. Brecht M, Schneider M, Sakmann B, Margrie TW. Whisker movements evoked by
stimulation of single pyramidal cells in rat motor cortex. Nature. 2004;427: 704–710.
doi:10.1038/nature02266

123. Houweling AR, Brecht M. Behavioural report of single neuron stimulation in
somatosensory cortex. Nature. 2008;451: 65–68. doi:10.1038/nature06447

124. Simoncelli EP, Olshausen BA. Natural image statistics and neural representation. Annu
Rev Neurosci. 2001;24: 1193–1216. doi:10.1146/annurev.neuro.24.1.1193

125. Barlow H. Redundancy reduction revisited. Netw Bristol Engl. 2001;12: 241–253.

126. Holtmaat A, Bonhoeffer T, Chow DK, Chuckowree J, De Paola V, Hofer SB, et al. Long-
term, high-resolution imaging in the mouse neocortex through a chronic cranial window.
Nat Protoc. 2009;4: 1128–1144. doi:10.1038/nprot.2009.89

127. Blackwell AA, Banovetz MT, Qandeel, Whishaw IQ, Wallace DG. The structure of arm
and hand movements in a spontaneous and food rewarded on-line string-pulling task by
the mouse. Behav Brain Res. 2018;345: 49–58. doi:10.1016/j.bbr.2018.02.025

128. Inayat S, Singh S, Ghasroddashti A, Qandeel, Egodage P, Whishaw IQ, et al. A Matlab-
based toolbox for characterizing behavior of rodents engaged in string-pulling. Kleinfeld
D, Wassum KM, editors. eLife. 2020;9: e54540. doi:10.7554/eLife.54540

129. Blackwell AA, Köppen JR, Whishaw IQ, Wallace DG. String-pulling for food by the rat:
Assessment of movement, topography and kinematics of a bilaterally skilled forelimb act.
Learn Motiv. 2018;61: 63–73. doi:10.1016/j.lmot.2017.03.010

130. Kislin M, Mugantseva E, Molotkov D, Kulesskaya N, Khirug S, Kirilkin I, et al. Flat-
floored Air-lifted Platform: A New Method for Combining Behavior with Microscopy or
Electrophysiology on Awake Freely Moving Rodents. JoVE J Vis Exp. 2014; e51869.
doi:10.3791/51869

131. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.
doi:10.1038/nmeth.2019

132. Peron S, Chen T-W, Svoboda K. Comprehensive imaging of cortical networks. Curr Opin
Neurobiol. 2015;32: 115–123. doi:10.1016/j.conb.2015.03.016

133. Hendrickson AE, White PO. Promax: A Quick Method for Rotation to Oblique Simple
Structure. Br J Stat Psychol. 1964;17: 65–70. doi:10.1111/j.2044-8317.1964.tb00244.x

134. Hilgetag CC, Kötter R, Stephan KE, Sporns O. Computational Methods for the Analysis
of Brain Connectivity. In: Ascoli GA, editor. Computational Neuroanatomy: Principles
and Methods. Totowa, NJ: Humana Press; 2002. pp. 295–335. doi:10.1007/978-1-59259-
275-3_14

135. Sporns O, Honey CJ, Kötter R. Identification and Classification of Hubs in Brain
Networks. PLOS ONE. 2007;2: e1049. doi:10.1371/journal.pone.0001049

136. Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, et al. Ultra-sensitive
fluorescent proteins for imaging neuronal activity. Nature. 2013;499: 295–300.
doi:10.1038/nature12354

137. Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, et al. High-performance
calcium sensors for imaging activity in neuronal populations and microcompartments. Nat
Methods. 2019;16: 649–657. doi:10.1038/s41592-019-0435-6

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

Fig 1. Experimental timeline and methods. (**A**) Timeline of experimental procedures, imaging times, and behavioral tests. (**B**) Glass cranial windows and headplates were implanted 2 weeks prior to beginning habituation protocols in Thy1-GCaMP6S mice. (**C**) Mice were habituated on the floating homecage, string pull and tapered beam for a period of 2 weeks prior to baseline measurements for behavior. Widefield Ca2+ response maps were used to determine locations for the imaging site located between the cFL and cHL sensory maps at baseline ("peri-infarct" imaging region) and imaging site lateral to the cHL at baseline ("distal" imaging region). For each Ca^{2+} imaging session, both imaging regions were imaged for a period of 15 minutes while simultaneously tracking animal movement within the mobile homecage. After baseline Ca^{2+} imaging, the somatosensory cFL map was targeted with photothrombosis. Behavioral testing on the string pull task and tapered beam task were performed weekly 1 day prior to the Ca^{2+} imaging session. After the final cellular Ca^{2+} imaging session at the 8 week time, stimulus-evoked widefield Ca^{2+} imaging was once again performed to determine the cFL and cHL somatosensory maps. (**D**) Representative widefield and two photon calcium imaging at each weekly time. The area of hyper-fluorescence due to stroke damage is apparent, with its surface area decreasing over time. (**E**) Neuron Ca^{2+} traces (ΔF) for 100 neurons from one example animal pre-stroke (left panel) and 1 week post-stroke (right panel) in the peri-infarct somatosensory imaging region. Lower line graphs demonstrate the tracked movement (mm/s) within the floating homecage corresponding to the Ca^{2+} traces above.

Fig 2. Photothrombotic stroke results in forelimb representation remapping onto adjacent areas of cortex and altered sensory-evoked widefield Ca^{2+} response properties at 8 weeks post-stroke. (**A**) Pseudocolored (%DeltaFoF) montages of representative cFL and cHL responses in S1 of an anesthetized animal pre-stroke and 8 week post-stroke resulting from oscillatory stimulation (1s, 100 Hz) of the cFL and cHL, respectively. At the 8 week post-stroke time, the cFL map has remapped posterior to its pre-stroke location into the area lateral to the cHL. (**B**) Mean peak forelimb Ca^{2+} transient response intensity (%DeltaFoF), measured from the thresholded cortical map area of the cFL. A significant interaction between time and stroke group was found, with *post-hoc* tests showing a significant decrease in the peak cFL response at 8-week post-stroke compared to pre-stroke (P = 0.0143). Mixed Effects Model, Time $F_{(1, 12)}$ = 0.8570, P = 0.3728; Stroke Group $F_{(1, 12)}$ = 0.2092, P = 0.6556. Interaction $F_{(1, 12)}$ = 8.632, P = 0.0124). (**C**) Thresholded cFL response area (mm^2). A significant main effect of group and significant interaction was found, with the stroke group showing a significant enlargement in the cFL response area at 8 weeks post-stroke compared to pre-stroke (P = 0.0152), and a greater response area at 8 weeks in the stroke group compared to sham (P = 0.0036). Mixed Effects Model, Time $F_{(1, 12)}$ = 2.273, P = 0.1575. Group $F_{(1, 12)}$ = 7.328, P = 0.0191. Interaction $F_{(1, 12)}$ = 5.372, P = 0.0389). (**D**) The stroke group was found to have significantly greater total distance shifted for the forelimb map at the 8 week time compared to sham (P = 0.0002). Stroke N = 9, Sham N = 5. *p < 0.05; **p < 0.01; ***p < 0.001

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

Fig 3. Impaired performance on tapered beam task after stroke. (**A**) Illustration of the
tapered beam test with the three elements measured; left (contralesional) side slips, right side
slips, and distance to first slip. (**B**) Mean number of left side slips. A significant interaction
between time and stroke group was observed, with *post-hoc* tests showing a greater number of
left side slips at 1 week in the stroke group relative to the sham group ($P = 0.0435$), and in the
stroke group at 1 week relative to baseline ($P = 0.0125$). Mixed Effects Model, Time $F_{(2.007, 38.13)}$
$= 2.178$, $P = 0.1270$; Stroke Group $F_{(1, 19)} = 2.260$; $P = 0.1492$. Interaction $F_{(5, 95)} = 2.914$, $P =$
0.0172. (**C**) Mean number of right-side slips. No main effects or interactions were observed. (**D**)
Normalized distance to first slip (as % of baseline). A trend towards a main effect of group was
observed, with no trends found in *post-hoc* tests. Mixed Effects Model, Time $F_{(3.135, 59.56)} =$
0.6447, $P = 0.5960.$; Stroke Group $F_{(1, 19)} = 3.030$, $P = 0.0979$; Interaction $F_{(4, 76)} = 1.242$, $P =$
0.4596). Stroke N = 16, Sham N = 5. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$

Fig 4. Firing rate of neurons in the peri-infarct cortex correlates with performance on the tapered beam task. (**A**) Example Ca^{2+} traces for 40 example neurons at each time point from the peri-infarct imaging region for one example animal in the stroke group. Significant Ca^{2+} transients are highlighted by red segments. A decrease in the number of Ca^{2+} transients is visually apparent at 1 week post-stroke. (**B**) In the peri-infarct imaging region, no main effects or interaction was seen in the normalized average firing rate during animal movement, however *post-hoc* analysis indicates a trend towards decreased normalized average firing rate at 1 week in the stroke group relative to sham ($P = 0.0593$). Mixed Effects Model, Time $F_{(2.448, 29.99)} = 0.9487$, $P = 0.4144$; Stroke Group $F_{(1, 13)} = 0.04256$, $P = 0.8397$; Interaction $F_{(4, 49)} = 1.852$, $P = 0.1339$. A significant interaction was seen in the normalized average firing rate at rest (**C**), with *post-hoc* tests showing a trend towards a decrease in firing rate at rest between the stroke and sham groups at the 1 week post time ($P = 0.0793$, Mixed Effects Model, Time $F_{(2.240, 27.44)} = 1.048$, $P = 0.3712$; Stroke Group $F_{(1, 13)} = 0.006936$, $P = 0.9349$; Interaction $F_{(4, 49)} = 2.604$, $P = 0.0471$). Normalized firing rate 1 week post-stroke was negatively correlated with the number of left slips in the tapered beam at the same 1 week time during both movement ($R^2 = 0.5326$, $P = 0.002$) (**D**) and rest ($R^2 = 0.4346$, $P = 0.0075$) (**E**). In the distal imaging region, no main effects or interaction were found in the normalized average firing rate for movement (**F**) or at rest (**G**). No correlation was observed between the normalized moving firing rate at 1 week in the distal region and the number of left slips on the tapered beam ($R^2 = 0.0637$, $P = 0.3286$) (**H**). A statistical trend was observed towards a negative correlation between normalized firing rate at rest at 1 week with left slips on the tapered beam at the same 1 week time ($R^2 = 0.2320$, $P = 0.0503$) (**I**). Peri-infarct region Stroke N = 11, Sham N = 4. Distal region Stroke N = 13, Sham N = 4. *p < 0.05; **p < 0.01; ***p < 0.001

1301

1302

1303

1304

1305

1306

Fig 5. Example cellular GCaMP6S Ca^{2+} fluorescence images and population functional connectivity plots. (**A-D**) Ca^{2+} fluorescence images and functional connectivity plots from the peri-infarct imaging region and the distal imaging region for one example stroke group animal. Population functional connectivity plots demonstrate neurons as red dots, and the functional connection strength as lines with color and line width determined by the strength of the correlation between neurons. Notable cortical damage is visible in the upper portion of the 1 week post-stroke Ca^{2+} fluorescence image in the peri-infarct imaging region (**A**), which notably improves over the 2-4 week times. A notable loss of functional connections and loss in the number of connections with strong correlation (>0.3) is visible in the functional connectivity plots at 1 week post-stroke in the peri-infarct imaging region (**B**). In the distal imaging region post-stroke, less structural change is visible in the cellular Ca^{2+} fluorescence images (**C**), and less change to functional connectivity of the population over time (**D**). (**E-H**) Ca^{2+} fluorescence images and functional connectivity plots from the peri-infarct imaging region and the distal imaging region for one example sham group animal. Less structural changes within the cellular Ca^{2+} fluorescence images and in the functional connectivity plots are visually apparent relative to the example stroke animal.

Fig 6. Altered functional connectivity in the peri-infarct somatosensory cortex. In the peri-infarct imaging region, a trend towards a main effect of group was found in the normalized average number of significant connections per neuron (**A**), with *post-hoc* tests showing a significant decrease at the 1 week post time in the stroke group relative to sham (P = 0.0440, Mixed Effects Model, Time $F_{(4.132, 47.11)}$ = 2.217, P = 0.1498; Stroke Group $F_{(1, 13)}$ = 4.505, P = 0.0551; Interaction $F_{(4, 45)}$ = 1.585, P = 0.1947). A significant main effect of group was found in the normalized number of significant connections per neurons with correlation greater than 0.3 (**B**), with *post-hoc* tests indicating a significant decrease at the 1 week post time in the stroke group relative to sham (P = 0.0363). (Time $F_{(2.102, 22.60)}$ = 0.3885, P = 0.6925. Group $F_{(1, 13)}$ = 6.423, P = 0.0262. Interaction $F_{(4, 43)}$ = 0.6512, P = 0.6292). (**C**) No main effects or interactions are seen in the connection density for the peri-infarct imaging region. (Time $F_{(3.251, 40.31)}$ = 0.3546, P = 0.8015. Group $F_{(1, 13)}$ = 1.257, P = 0.2824. Interaction $F_{(5, 62)}$ = 1.146, P = 0.3462). In the distal imaging region, no significant main effects or interactions are seen in the normalized number of significant connections per neuron (**D**) (Time $F_{(3.173, 46.32)}$ = 0.5674, P = 0.6487. Group $F_{(1, 15)}$ = 0.3172, P = 0.5816. Interaction $F_{(5, 73)}$ = 1.525, P = 0.1925), normalized amount of significant connections with correlation greater than 0.3 (**E**) (Time $F_{(2.228, 32.86)}$ = 2.196, P = 0.1224. Group $F_{(1, 15)}$ = 1.036, P = 0.3249. Interaction $F_{(4, 59)}$ = 0.7713, P = 0.5482), or in the connection density (**F**) (Time $F_{(3.467, 52.01)}$ = 1.299, P = 0.2841. Group $F_{(1, 15)}$ = 2.062, P = 0.1715. Interaction $F_{(5, 75)}$ = 1.251, P = 0.2942). Peri-infarct region Stroke N = 11, Sham N = 4. Distal region Stroke N = 13, Sham N = 4. *p < 0.05; **p < 0.01; ***p < 0.001

A

| | Pre | 1 Week Post | 2 Week Post | 3 Week Post | 4 Week Post | 8 Week Post |

Stroke Animal — Peri-Infarct / Distal

Sham Animal — Peri-Infarct / Distal

Fig 7. Color coded neural assembly plots depict altered neural assembly architecture after
stroke in the peri-infarct region. (A) Representative cellular Ca^{2+} fluorescence images with
neural assemblies color coded and overlaid for each time depict a loss in the number of neural
assemblies after stroke in the peri-infarct region, and a concurrent increase in the number of
neurons for each remaining assembly. (B) Representative sham animal displays no visible
change in the number of assemblies or number of neurons per assembly.

Fig 8. Neural assembly structure is disrupted in the peri-infarct region after stroke. In the peri-infarct imaging region, a significant main effect of group was found for the number of assemblies (**A**) (Time $F_{(2.883, 35.75)} = 0.8231$, $P = 0.4857$. Group $F_{(1, 13)} = 11.90$, $P = 0.0043$. Interaction $F_{(5, 62)} = 1.496$, $P = 0.2040$), with *post-hoc* tests showing a significant decrease in the number of assemblies at the 1 week and 3 week times for the stroke group compared to sham ($P = 0.0114$ & 0.0058, respectively). A statistical trend is also observed at the 2 week time for the stroke group compared to sham ($P = 0.0589$). A trend towards a main effect of group was found in the normalized number of neurons per assembly (**B**, Mixed Effects Model, Time $F_{(2.361, 28.80)} = 2.185$, $P = 0.1234$; Stroke Group $F_{(1, 13)} = 3.694$, $P = 0.0768$; Interaction $F_{(5, 61)} = 1.202$, $P = 0.3192$), with *post-hoc* tests showing a significant increase in the normalized number of neurons per assembly at 1 week for the stroke group compared to sham ($P = 0.0124$). A trend towards a main effect of group was found in the average percent of the population per assembly (**C**, Time $F_{(1.711, 21.22)} = 1.738$, $P = 0.2023$; Stroke Group $F_{(1, 13)} = 3.807$, $P = 0.0729$. Interaction $F_{(5, 62)} = 1.765$, $P = 0.1333$), with *post-hoc* tests showing a trend towards an increase in the stroke group relative to sham at the 1 week post time ($P = 0.0863$). No main effects or interaction was observed in the distal imaging region for number of assemblies (**D**), normalized number of neurons per assembly (**E**), or average percent of the population per assembly (**F**). Peri-infarct region Stroke N = 11, Sham N = 4. Distal region Stroke N = 13, Sham N = 4. *p < 0.05; **p < 0.01; ***p < 0.001

Fig 9. Correlation among neural assemblies is increased in the peri-infarct imaging region after stroke. (**A**) Illustration of three example color coded neural assemblies from a neural population and their corresponding time series activations that were correlated. In the peri-infarct imaging region, a significant main effect of group was found in the correlation coefficient for significantly correlated assembly-assembly pairs (**B**, Mixed Effects Model, Time $F_{(3.718, 42.39)}$ = 2.014, P = 0.1141; Stroke Group $F_{(1, 13)}$ = 9.155, P = 0.0097. Interaction $F_{(5, 57)}$ = 2.348, P = 0.0524), with *post-hoc* tests indicating a significantly higher correlation at 2 week (P = 0.029) and 4 weeks (P = 0.0253) in the stroke group relative to sham, and at 1 week (P = 0.0084) and 4 week (P = 0.0261) in the stroke group compared to pre-stroke. In the distal imaging region, no significant main effects or interaction was found for the correlation coefficient for significantly correlated assembly-assembly pairs (**C**). Peri-infarct region Stroke N = 11, Sham N = 4. Distal region Stroke N = 13, Sham N = 4. *p < 0.05; **p < 0.01; ***p < 0.001