Enhancing *Escherichia coli* Abiotic Stress Resistance through Ornithine Lipid Formation

Leidy Patricia Bedoya-Pérez[1], Alejandro Aguilar-Vera[1], Mishael Sánchez-Pérez, José Utrilla[1*], Christian Sohlenkamp[1*]

[1]Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Av. Universidad s/n Col. Chamilpa, C.P. 62210, Cuernavaca, Mor., México.

*Corresponding authors

Name: José Utrilla: utrilla@ccg.unam.mx

Name: Christian Sohlenkamp: chsohlen@ccg.unam.mx

Address: Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Av. Universidad s/n, Col. Chamilpa, C.P. 62210, Cuernavaca, Mor., México.

Abstract

Escherichia coli is a common host for biotechnology and synthetic biology applications. During growth and fermentation, the microbes are often exposed to stress conditions, such as variations in pH or solvent concentrations. Bacterial membranes play a key role in response to abiotic stresses. Ornithine lipids (OLs) are a group of membrane lipids whose presence and synthesis have been related to stress resistance in bacteria. We wondered if this stress resistance could be transferred to bacteria not encoding the capacity to form OLs in their genome, such as *E. coli*. In this study, we engineered different *E. coli* strains to produce unmodified OLs and hydroxylated OLs by expressing the synthetic operon *olsFC*. Our results showed that OL formation improved pH resistance and increased biomass under phosphate limitation. Transcriptome analysis revealed that OL-forming strains differentially expressed stress- and membrane-related genes. OL-producing strains also showed better growth in the presence of the ionophore carbonyl cyanide 3-chlorophenylhydrazone (CCCP), suggesting reduced proton leakiness in OL-producing strains. Furthermore, our engineered strains showed improved heterologous violacein production at phosphate limitation and also at low pH. Overall, this study demonstrates the potential of engineering the *E. coli* membrane composition for constructing robust hosts with an increased abiotic stress resistance for biotechnology and synthetic biology applications.

Keypoints:

- The *E. coli* membrane composition was engineered by producing ornithine lipids
- Ornithine lipid production increase biomass yield under phosphate limitation
- Engineered strains show enhanced production phenotype under low pH stress
- Transcriptome analysis and CCCP experiments revealed reduced proton leakage

48   **Graphical abstract**

INTRODUCTION

Microorganisms are crucial in biotechnology for producing bulk chemicals, proteins, biofuels, and pharmaceuticals. During fermentation, these microbes often face stress conditions, such as pH variations, changing temperature, and nutrient levels (Cordell et al. 2023). Media components can significantly contribute to the overall cost of a bioprocess (Cardoso et al. 2020), and phosphate is a scarce and costly component. To improve industrial production, it is desirable to cultivate strains that can handle stress and different growth conditions (Calero and Nikel 2019). Such adapted microorganisms are called "microbial chassis" (Beites and Mendes 2015) and can be engineered using synthetic biology tools (Marquez-Zavala and Utrilla 2023).

During the response of bacteria to abiotic stresses, the membrane has an important function, with bacteria frequently adapting their lipid composition (Sohlenkamp 2017). A major component of *Escherichia coli* membranes are glycerophospholipids, which are composed of two fatty acids, a glycerol moiety, a phosphate group, and a variable head group (Sohlenkamp and Geiger 2015). Although glycerophospholipids are the best-known membrane lipids, many other lipids can be present in membranes of different bacteria, such as hopanoids, ornithine lipids (OLs), betaine lipids, sulfonolipids, or sphingolipids (Sohlenkamp and Geiger 2015). Most importantly, the lipid composition of a bacterial membrane is not static and the presence of some membrane lipids, specifically hopanoids and OLs, seems to form part of a stress response to (changing) environmental conditions in some bacteria (Sohlenkamp and Geiger 2015; López-Lara and Geiger 2017).

OLs are phosphorus-free membrane lipids that are relatively common in bacteria but are absent in eukaryotes and archaea (Kim et al. 2018). They are composed of a 3-hydroxy fatty acid amide bound to the α-amino group of ornithine and a second fatty acyl group ester-linked to the 3-hydroxy position of the first fatty acid. This basic OL structure can be modified by hydroxylation at different positions, by *N*-methylation, or by taurine transfer. It was predicted that approximately 50% of sequenced bacterial species can synthesize OLs under certain growth conditions. These

conditions include limiting phosphate concentrations, acidic pH, and increased temperatures and

several bacteria such as *Pseudomonas fluorescens*, *Serratia proteamaculans*, and *Vibrio cholerae*

only form OLs under phosphate-limited conditions. In *Burkholderia cepacia,* an increased

formation of hydroxylated OLs (OLs-OH) was observed in cells grown at 40 °C, and in the

symbiotic alpha-proteobacterium *Rhizobium tropici* CIAT899 the presence of OLs-OH improves

growth at increased temperatures and under acidic conditions (Vences-Guzmán et al. 2011;

Vences-Guzmán et al. 2015)(Sohlenkamp and Geiger 2015; Córdoba-Castro et al. 2021). In

particular, the hydroxylation introduced by the OL hydroxylase OlsC in the C2 position of the

secondary fatty acid in OLs may increase hydrogen bonding between the lipid headgroups,

thereby increasing acid stress resistance (Nikaido 2003). In recent years, many enzymes involved

in the synthesis and modification of OLs have been described (Sohlenkamp 2017). OlsF is a

bifunctional OL synthase that was first described in *S. proteamaculans*. When OlsF is expressed

in *E. coli*, the resulting strains can form OLs in one step (Vences-Guzmán et al. 2015; Escobedo-

Hinojosa et al. 2015)

E. coli is among the most frequently used bacterial chassis because of its easy genetic

manipulation, fast growth to high cell densities, and safe use (Chi et al. 2019). Recently, the

computational tool ReProMin (Regulation-based Proteome Minimization) was designed to

identify genetic interventions required to divert cellular resources invested in the expression of

non-essential genes toward the increase in biosynthetic potential. Two generated mutants (PFC,

and PYC strains) showed higher cellular resources to produce recombinant proteins, plasmid

DNA, and heterologous metabolites (Lastiri-Pancardo et al. 2020; de la Cruz et al. 2020).

However, these mutants lack the phosphate regulon transcriptional activator encoding gene *phoB*,

which might limit their use of alternative phosphate sources.

Wild-type *E. coli* cannot form OLs and in its inner membrane the glycerophospholipids,

phosphatidylethanolamine (PE), phosphatidylglycerol (PG), and cardiolipin (CL) can be detected.

This composition is relatively stable under different growth conditions. In the present study, we

hypothesized that the formation of OLs and OLs-OH in *E. coli* strains might improve their

resistance to different abiotic stress conditions. We constructed the synthetic operon pSEVA*olsFC* (p*olsFC*) comprising the genes o*lsF* from *S. proteamaculans* and *olsC* from *R. tropici* CIAT899 required to produce unmodified and hydroxylated OLs, respectively, and expressed it in different genetic backgrounds of *E. coli* (Rojas-Jiménez et al. 2005; Escobedo-Hinojosa et al. 2015). We observed increased biomass formation under phosphate limitation and improved growth with a shortened lag phase at reduced pH values. Transcriptome analysis revealed that OL-forming strains differentially expressed stress- and membrane-related genes, indicating reduced proton leakiness. Consistently, OL-producing strains showed better growth in the presence of the ionophore carbonyl cyanide 3-chlorophenylhydrazone (CCCP). Finally, we showed that the formation of OLs-OH in *E. coli* strains improved violacein formation in a heterologous expression system compared with *E. coli* strains not forming OLs.

MATERIALS AND METHODS

Strains and culture media

Bacterial strains used in this study are listed in Table S1. *E. coli* strains were routinely grown in

LB medium and MOPS/MES minimal medium supplemented with 0.01% casamino acids (CAA)

(Sigma-Aldrich), 4 g/L glucose, and 2.6 mM K_2HPO_4 as phosphate source (0.4 mM in phosphate

limited experiments). For the low pH experiments, MES buffer was used. To evaluate bacterial

tolerance to the proton motive force (PMF) uncoupler CCCP, MOPS/glucose medium was used

with 10μM CCCP (Sigma-Aldrich). Antibiotics were used at the following concentrations: 25

μg/mL gentamicin (Gm), 100 μg/mL ampicillin (Amp), and 50 μg/mL kanamycin (km).

Construction of pSEVA336*olsFC* and pAJM336*vioABCDE* plasmids by Gibson assembly

Plasmids and oligonucleotides used in this study are listed in Tables S1 and S2, respectively. The

olsF gene from *S. proteamaculans* was PCR amplified from plasmid pET9a2569 (Vences-

Guzmán et al. 2015) using primers Spro2669_olsF FW and Spro2669_olsF RV. Plasmid

pSEVA631 (Amp^r) (Silva-Rocha et al. 2013)was used as the cloning backbone, it was PCR

amplified using primers pSEVAolsF FW and pSEVAolsF RV. All backbone vector PCR products

were digested with DpnI prior to the assembly reactions. The PCR products were gel purified

(Zymoclean Gel DNA Recovery Kits) and assembled into plasmid pSEVA631*olsF* (NEBuilder

HiFi DNA Assembly Master Mix) following the manufacturer's recommendations. Then the *R.*

tropici olsC gene was PCR amplified from plasmid pCCS98 (Vences-Guzmán et al. 2011) using

the oligonucleotides pSEVAolsFC_FW (which contains the sequence of a ribosome binding site)

and pSEVAolsFC_RV. The pSEVA631olsF was PCR amplified and assembled with the *olsC*

PCR product to generate pSEVA*olsFC*. The violacein production plasmid was constructed

following a similar approach, the *vioABCD* operon was constructed as a three-part assembly

sourcing coding gene from pSEVA63-hvio (Darlington et al. 2018). The two-part operon was

assembled into pAJM336 which is an IPTG inducible expression vector. The *vioA* fragment was

PCR amplified using oligonucleotides vioA_Fw and vioA_Rv, while genes *vioB*, *vioC*, *vioD*, and

vioE were amplified in a single fragment with oligonucleotides vioB_E-Fw and vioB_E-Rv. All

plasmids were verified by Sanger sequencing of the cloned products. Also, the presence of OLs-

OH was verified by TLC.

Growth phenotype characterization.

The bacterial strains were cultured overnight in 125-mL flasks, each containing 25 mL of

MOPS/glucose medium supplemented with 2.6 mM K_2HPO_4. These cultures were incubated at

37 °C with continuous agitation at 250 rpm. On the following day, the cells were harvested by

centrifugation at 5000 rpm for 10 minutes at 4°C. Subsequently, cells were washed three times

with 25 mL of culture medium tailored to match the specific experimental conditions (phosphate

limitation, low pH, or 10 µM CCCP treatment).

For the growth experiments, 600-µL aliquots of each culture with an OD_{600} of approximately 0.05

were prepared using fresh medium. From each cellular suspension, 150 µL were pipetted into

individual wells in transparent flat-bottom 96-well plates (Corning), with triplicate wells for each

sample. The plates were then incubated at 30 °C with fast linear shaking in a microplate reader

(ELx808, BioTek), where the OD_{600} was measured at 20-minute intervals over a 24-hour period.

To determine cell biomass concentration (in dry weight mg/mL), we employed a calibration curve

established through linear regression (y = 0.01997 + 1.92194 * x, R^2 = 0.99) correlating OD_{600}

values to dry weight measurements. The analysis of growth kinetics was conducted across three

independent experiments, each with triplicate measurements (n = 9).

For the assessment of growth kinetics in 96-well plates, we employed the algorithm developed by

Swain et al. (Swain et al. 2016) with default parameters, running it on Python. This analysis

provided us with data on the magnitude and timing of the maximum growth rate and maximum

OD_{600} values. To assess statistical significance, we calculated *p*-values using a two-tailed

Student's t-test as the reliability index for statistical significance (Hidalgo et al. 2022).

***In vivo* labeling of *E. coli* strains and qualitative analysis of lipid extracts**

The lipid compositions of bacterial strains were analyzed by labeling cells with $[^{14}C]$ acetate

(Amersham Biosciences) as previously described (Vences-Guzmán et al. 2015). The strains were

grown overnight in 2.6 mM K_2HPO_4 MOPS/glucose medium supplemented with 0.01% CAA and

Amp. Then, the cells were harvested by centrifugation at 4000 rpm at room temperature and

washed three times with culture medium (phosphate-limited MOPS/glucose or MES/glucose pH

5.8). 125-mL flasks containing 25 mL of the respective culture medium were inoculated with

aliquots of each preculture to obtain a final OD_{600} of approximately 0.1. When an OD_{600} of

approximately 0.3 was reached, 1 mL of each culture was transferred into a 15-mL tube and mixed

with 0.5 µCi of $[^{14}C]$ acetate (60 µCi mmol^{-1}), and the cultures were further incubated overnight

at 37 °C. The cells were then harvested by centrifugation, washed once with 500 µL of water, and

resuspended in 100 µL of water. Lipid extracts were obtained according to the Bligh and Dyer

protocol (Bligh and Dyer 1959). Aliquots of the lipid extracts were spotted on high-performance

TLC silica gel 60 plates (Merck, Poole, UK) and separated in one dimension using chloroform/

methanol/water (130:50:8, v/v) as the mobile phase. The one-dimensional TLC plates were

developed for 4-6h and were exposed to a PhosphorImager screen (Amersham Biosciences) for

24 to 72 h. The relative quantities of individual lipids were determined using ImageQuant

software (Amersham Biosciences).

Primary amine-containing lipids (PE, OL, OL-OH) were visualized by spraying the plates with a

solution of 0.2% (w/v) ninhydrin in acetone and subsequently heated at 120°C for 10 minutes.

Quantification of violacein production in *E. coli* strains.

Strains were grown in MOPS/glucose medium containing Gm and Amp and incubated at 37°C

overnight. The next day, 1 mL of each culture was harvested by centrifugation, washed three

times with culture medium (MOPS/glucose phosphate limitation or MES/glucose), and

resuspended in 1 mL of the same solution. Microtubes (1.5 mL) containing 0.6 mL (final volume)

of either MOPS/glucose or MES/glucose medium supplemented with 2 g/L tryptophan, 0.01%

CAA, and 800 nM IPTG were inoculated with their corresponding preculture to obtain an OD_{600}

of approximately 0.05. Then, 150 μL of each culture (in triplicate) were transferred into a 96-well

plate and incubated as described above. After the incubation period (24 h), the plates were

centrifuged for 10 min at $13,000 \times g$, and the supernatant was discarded. For each well, violacein

was extracted by resuspending the cell pellets in 200 μL absolute ethanol. The plates

were incubated at 95 °C for 10 min, followed by pelleting the cell debris by centrifugation

($13,000 \times g$, 10 min). The ethanol phase containing violacein was collected and transferred to

another 96-well plate, and its absorbance was measured at OD_{575} in a microplate reader (Synergy

2.0, BioTek). The violacein concentration was determined by extrapolation from the violacein

standard (Sigma-Aldrich) calibration curve and calculated as mg/L (Lastiri-Pancardo et al. 2020).

RNA extraction

E. coli strains K12pSEVA, and K12polsFC were pre-grown in 125-mL flasks containing 25 mL

of MOPS minimal medium with 4 g/L glucose at 37°C overnight, 250 rpm. The next day, 1 mL

of cultures was harvested by centrifugation at 4000 rpm, 4°C, and washed three times with

MES/glucose medium pH 5.8. Cultures were inoculated in 125-mL flasks containing 25 mL of

MES/glucose medium pH 5.8. When the cultures reached the exponential growth phase (11h), 3

mL of cultures were transferred to a sterile 15-mL tube and mixed with 3 mL of RNA protect

reagent. Cells were harvested by centrifugation, and the pellets were immediately frozen in liquid

nitrogen, and stored at -70°C until use. Total RNA was prepared using the RNeasy mini kit

(QIAGEN, Hildesheim, Germany). Pellets were resuspended in RLT buffer supplemented with

lysozyme (20 mg/mL; QIAGEN, Hildesheim, Germany) and Zirconium Oxide Beads (0.5 mm,

Next Advance). Cells were disrupted using a Bullet Blender Tissue Homogenizer (Next-Advance,

Troy, NY, USA) in impact-resistant 2-mL tubes. Genomic DNA was eliminated by digestion with

RNase-free DNase (QIAGEN, Hildesheim, Germany) for 20 min at room temperature. Final RNA

concentrations were determined using a NanoDrop (Thermo Scientific). The typical OD_{260} to

OD_{280} ratio of the RNA samples was approximately 2.0. The integrity of the RNA samples was

verified by determining the RNA integrity number (RINe) in a TapeStation 2200 instrument

(Agilent Technologies). Two independent total RNA extractions were obtained for each condition

and each one was analyzed separately (Guerrero-Castro et al. 2018).

RNA-Seq

The RNA-seq samples were analyzed by Novogene (Sacramento, CA, USA). A total amount of

1 μg RNA per sample was used as input material for the RNA sample preparations. Sequencing

libraries were generated using the NEBNext ® UltraTM RNA Library Prep Kit for Illumina®

(NEB, USA) according to the manufacturer's recommendations, and index codes were added to

attribute sequences to each sample. After rRNA removal, fragmentation was performed using

divalent cations under elevated temperature in NEBNext First Strand Synthesis Reaction Buffer

(5X). First-strand cDNA was synthesized using random hexamer primers and M-MuLV reverse

transcriptase (RNase H-). Second-strand cDNA synthesis was subsequently performed using

DNA polymerase I and RNase H. The remaining overhangs were converted into blunt ends via

exonuclease/polymerase activities. After the adenylation of 3' ends of DNA fragments, the

NEBNext adaptor with a hairpin loop structure was ligated to prepare for hybridization. To select

cDNA fragments of preferentially 150–200 bp in length, the library fragments were purified with

the AMPure XP system (Beckman Coulter, Beverly, USA). Then, 3 μL USER Enzyme (NEB,

USA) was used with size-selected, adaptor-ligated cDNA at 37 °C for 15 min followed by 5 min

at 95 °C before PCR. PCR was then performed with Phusion High-Fidelity DNA polymerase,

Universal PCR primers, and Index (X) Primer. Finally, PCR products were purified (AMPure XP

system) and library quality was assessed using the Agilent Bioanalyzer 2100 system. Clustering

of the index-coded samples was performed on an Illumina Novaseq sequencer according to the

manufacturer's instructions. After cluster generation, the libraries were sequenced on the same

machine, and paired-end reads were generated.

Data processing and differential expression

The raw reads were processed for adapter trimming and quality control filtering using Trim

Galore (v.0.6.6). The reads were aligned using the R package Rsubread (v.2.2.6) and mapped to

the reference genome (GCF_000005845.2). For gene expression quantification, the function

feature Counts from the R subread package were used, generating the read counts matrix. We

used batch correction ARSyseq and TMM for normalization. Differential expression analysis was

performed using DESeq2 (v.1.28.1). The filters used for the DEGs were: absolute

log2FoldChange >= 1 and *p*-value adjust <= 0.05. The iModulon analysis consists of adding the

differential expression data to each iModulon, and the order within the graphs was made based

on the "system category" field (Rychel et al. 2021; Lamoureux et al. 2023). The accession number

for the RNAseq data reported in this study is GEO: GSE233657.

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE233657).

 RESULTS

Construction of *E. coli* strains synthesizing ornithine lipids and hydroxylated ornithine lipids

The success of many synthetic biology and biotechnology applications depends on the presence of a robust microbial chassis. We wanted to test if and how the presence of OLs in *E. coli* would affect its stress resistance and its performance as a host for synthetic biology applications.

First, we constructed a plasmid-borne synthetic operon containing the genes *olsF* and *olsC* required for the synthesis of OLs and hydroxylated OLs (OLs-OH) (Rojas-Jiménez et al. 2005; Escobedo-Hinojosa et al. 2015) (Fig. 1a). This plasmid called p*olsFC* and the empty control plasmid pSEVA631 (pSEVA) (Silva-Rocha et al. 2013)were transformed into four different *E. coli* strains: the wild-type strain *E. coli* K12 (BW25113), the triple mutants PFC (*phoB, fhlC* and *cueR*) and PYC (*phoB, yedW, cusR*), which have a reduced cellular proteome(Lastiri-Pancardo et al. 2020), and a *phoB*-deficient strain (Baba et al. 2006). Second, we wanted to determine if the *E. coli* strains expressing the synthetic operon formed OLs and OLs-OH. The strains were grown MOPS medium without phosphate-limited, [^{14}C] acetate was added for lipid labeling, and total lipids were extracted and analyzed by thin-layer chromatography (TLC). In all strains, the glycerophospholipids PE, PG, and CL were detected. In addition to these lipids, the *E. coli* strains expressing the synthetic operon *olsFC* produced both unmodified OLs and OLs-OH (Fig. 1b). A few lipids of minor abundance that were not identified were observed in the TLC (labeled with asterisks). Interestingly, in *E. coli* strains accumulating OLs and OLs-OH, the formation of PE was reduced relative to the control strains with the empty plasmid (Fig. 1b).

Figure 1. Heterologous production of OLs and OLs-OH in _E. coli_ strains. (a) Synthetic operon _olsFC_

under the control of a C6 Acyl homoserine Lactone (AHL) inducible promoter in the plasmid p_olsFC_. (b)

TLC analysis of total membrane lipids of _E. coli_ strains K12, PYC, PFC, and _phoB_ harboring pSEVA or

p_olsFC_. Total lipids were labeled with [^{14}C]acetate, extracted, and separated by one-dimensional TLC. PE,

phosphatidylethanolamine; PG, phosphatidylglycerol; CL, cardiolipin; OLs, unmodified ornithine lipid,

OLs-OH: hydroxylated ornithine lipid, *: unidentified lipid.

OLs-OH production increases biomass yield under phosphate limited

In bacteria such as _S. proteamaculans_ or _V. cholerae,_ OLs are not formed under phosphate-replete

conditions, and their synthesis is induced by phosphate-limited growth conditions (Barbosa et al.

2018). It is thought that the zwitterionic OLs replace the zwitterionic phospholipid PE under

conditions of phosphate limitation, reducing the requirement for phosphate for the membrane and

thereby allowing an alternative use for this nutrient. We observed a decrease in PE formation

when OLs were accumulated in _E. coli_ strains expressing the synthetic operon (Fig 1b). Therefore,

we hypothesized that an _E. coli_ strain able to replace a fraction of its phospholipids with OLs

could have a growth advantage under phosphate-limited conditions because phosphate normally

used for membrane synthesis would be liberated for other cellular processes.

We performed growth experiments of all *E. coli* strains in MOPS/glucose medium with

phosphate-limited concentrations (0.4 mM) and phosphate-replete concentrations (2.6 mM). We

used 0.4 mM phosphate concentration where we observed a substantial reduction in biomass yield

but it did not severely affect the growth rates of the evaluated strains (Fig.S1a). In this set of

experiments, we included host strains and their derivatives carrying empty plasmids or expressing

the synthetic operon *olsFC*. We determined the effect of the presence of OLs-OH on the biomass

production of *E. coli* grown under phosphate-limited (0.4 mM) and phosphate-replete (2.6 mM)

concentrations. In MOPS/glucose medium with 2.6 mM phosphate, no difference in biomass

formation was observed between the different strains analyzed (Fig. S1a). In contrast, in

phosphate-limited MOPS/glucose medium, biomass production in the strains that synthesized

OLs increased by 23 to 41 % compared with the strains carrying the empty plasmid (Fig. 2a). In

all cases, the biomass formation of the OL-forming strains was higher than that of the wild-type

strains lacking the plasmid p*olsFC* (Fig. 2a, S1a and S2). These results suggest that the synthesis

of OLs provides *E. coli* with an advantage when growing under growth-limited phosphate

concentrations.

OLs-OH production improves growth phenotype at low pH

We wondered if the presence of hydroxylated OLs would improve growth in MES/glucose

medium without phosphate limitation at pH 5.8 in our engineered *E. coli* strains. As above

experiments, we evaluated the four host strains and their derivatives carrying either the

pSEVA631*olsFC* plasmid or the empty vector. The presence of OLs significantly enhanced the

maximum growth rate of *E. coli* strains at pH 5.8, resulting in an approximately 35% increase

compared with the vector control strains in three out of the four genetic backgrounds examined

(Fig. 2b). When strains were grown under phosphate-replete conditions, no differences in growth

were observed. (Fig S1a and S1b). On the other hand, during the first 8 h of the growth kinetic,

all strains (with or without OLs) exhibited similar biomass. However, a noticeable difference

emerged when comparing the plasmid-harboring strains in the subsequent hours and after 24h, all

four OL-forming strains accumulated more biomass than the vector control strains. The OL-

forming strains showed a shortened lag phase compared with the four wild-type strains lacking

plasmids (Fig. S3).

Figure 2. The presence of ornithine lipids in *E. coli* strains improves their growth phenotype under phosphate limitation and at pH 5.8. (a) Growth rates of the different *E. coli* strains were calculated from the growth curves in MES/glucose medium with 2.6 mM phosphate at pH 5.8. (b) Dry weight concentration (mgDW/mL) of cultures after 24-hour growth in MOPS/glucose medium with low phosphate concentration (0.4 mM). The data represent the average of three independent experiments. Error bars indicate the standard deviation using two-tailed unpaired Student 's t-test. *p*-values $*p \leq 0.05$, $****p \leq 0.0001$.

Transcriptional response to OLs production at pH 7.4 and pH 5.8

Abiotic stresses, including acidic pH, have been reported to affect the transcriptional response in

E. coli (Aquino et al. 2017).Therefore, we wanted to determine if the positive phenotypic effects

at low pH caused by the presence of OLs-OH in *E. coli* could be explained by transcriptional

changes. Given that PYC and PFC strains lack transcription factor genes that rewire their

transcriptional regulatory network (Lastiri-Pancardo et al. 2020), we used the wild-type K12

strain to evaluate the specific effect of the OL presence on the transcriptome of *E. coli* grown at

both pH 7.4 and pH 5.8 without phosphate limitation.

Total RNA was extracted from exponentially growing cultures (11h) as at this time point, a clear

difference in growth rate was observed between the strains forming OLs and those lacking them

(Fig. 2a). We conducted RNA-Seq analysis on two independent biological samples and sequenced

the libraries to approximately 10 million read depths. We then compared the expression profiles

of these samples based on their response to different perturbations: A) response to OLs production

at neutral pH, B) response to OLs production at low pH, C) response of the control strain to low

pH compared with neutral pH, D) response of the OLs- producing strain to low pH compared with

neutral pH, and E) differential response to low pH and OLs formation (Table 1).

Table 1. Transcriptional response to OLs formation at neutral and low pH. Perturbations and descriptions

PERTURBATION	DESCRIPTION
A	Response to the presence of OLs at pH 7.4. Comparison between K12pSEVA pH 7.4 and K12pOlsFC pH 7.4.
B	Response to the presence of OLs at pH 5.8. Comparison between K12pSEVA pH 5.8 and K12pOlsFC pH 5.8.
C	Response of the wild-type strain to low pH. Comparison between K12pSEVA pH 7.4 and K12pSEVA pH 5.8.
D	Response of the OL-forming strain to low pH. Comparison between K12pOlsFC pH 7.4 and K12pOlsFC pH 5.8.
E	Differential response to low pH challenge and OLs formation. Comparison between K12pSEVA pH 7.4 and K12pOlsFC pH 5.8.

For the transcriptomic analyses, we used the iModulon classification of the Differentially Expressed Genes (DEGs) data, which consists of the clustering of genes sharing an independently modulated signal. iModulons were obtained by independent component analysis (ICA) of 815 sample transcriptional profiles from 422 conditions for *E. coli* . For further analyses, we considered only those iModulons that covered 80% or more of the DEGs with a *P*-value < 0.05 and Log2 fold change of ± 1; that is, from the 204 classified iModulons(Rychel et al. 2021), 48 passed the filter (Fig. 3a and Fig. S4). Overall, the results show a mild transcriptional response to OLs production (perturbations A and B), and a large transcriptional response to low pH conditions (C and D) (Fig 3b).

The presence of OLs in *E. coli* grown at pH 7.4 (perturbation A) affected the gene expression level of 25 iModulons, 13 of which were upregulated, including those related to stress response (Curli, PspF, Hydrogenase 1, ProVWX, BlurR, asr, ZntR and Fur 1-2) and metabolism (GlcC, BirA, AHAS I and PurR). Some of the iModulons with reduced activity in perturbation A are associated with the maintenance of membrane and proton motive force (PMF) (*pspF*), and the integrity of the outer membrane under moderate acid stress (*ars*) (Fig 3b). In perturbation B

(presence of OLs at pH 5.8), five and eight iModulons showed reduced and increased activity,

respectively. Of note, the expression of genes clustered in the *pspF* iModulon were also decreased

as in perturbation A (Fig 3b). For perturbations C and D (effect of pH 5.8 in the absence and

presence of OLs, respectively), a similar tendency was observed in the class of iModulons

affected (stress and metabolism). However, in perturbation D, the nutrition limitation-associated

pyR iModulon was upregulated while it did not change in perturbation C (Fig 3b).

To better understand the specific transcriptional responses that benefit the growth phenotype of

the OLs-producing strains under low pH, we compared the expression profile of the control strain

at neutral pH with that of the OLs-producing strain at low pH (perturbation E). Two of the 48

filtered iModulons showed changes in their activity in this perturbation. The differential response

to low pH when expressing OLs showed a few key differences compared with the response of the

control strain at neutral pH (Fig 3b). For instance, reduced activity of the *pspF* iModulon was

detected in perturbation E (as in A and B), suggesting that the presence of OLs specifically

reduces the expression of genes related to the maintenance of PMF. In addition, expression of the

kdpE iModulon (potassium two-component system) was reduced in the presence of OLs

(perturbations, B, C, D, and E). In particular, a reduction in the activity of the *pdhR* iModulon,

which contains genes associated with the formation of the pyruvate dehydrogenase (PDH)

multienzyme complex and respiratory electron transport system (Ogasawara et al. 2007), was

only detected by the analysis of perturbation E (Fig 3b).

An increase in the activity of the *curli* iModulon in the perturbations A, B and E, indicates that

OLs production positively affects the expression of genes related to curli fiber synthesis and

biofilm formation in *E. coli*. In addition, the proVWX iModulon (transport system for the

osmoprotectant glycine betaine) was also induced in these perturbations (Fig 3b).

Besides the above-mentioned analyses by iModulons, we also analyzed individual genes that are

related to ornithine synthesis, the synthesis and turnover of membrane lipids, and the response to

acid stress in the four defined perturbations (Fig 3c). Genes *adiY* and *adiC*, which encode an

arginine/agmatine antiporter of the arginine-dependent extreme acid resistance (XAR), were

induced at low pH (Fig. 3c; perturbations C, D, and E). Genes involved in the acid stress response

such as *gadA*, *hdeD*, and *yhiM* were induced by the formation of OLs (perturbations A, B and E).

In contrast, the expression of *asr* that encodes a periplasmic intrinsically disordered protein

induced under acid shock conditions was downregulated in perturbations A, C, D and E (Fig 3c).

Collectively, these results showed that modifications of the membrane composition by OLs

change the regulatory response to low pH in *E. coli*.

Figure 3. Low pH conditions cause a large transcriptional response, whereas OL formation causes a

mild transcriptional response. (a) 48 iModulons with coverage greater than 80% in at least one

perturbation were used for this analysis. The functional categories are denoted according to color, yellow:

metabolism, black: stress response, and white: other functions. (b) Differential expression of each analyzed

iModulon. (c) Expression changes of individual genes involved in arginine metabolism, membrane and

lipid synthesis, and stress response to low pH not classified by iModulons. (value= DEGup - DEGdown),

blue color, reduced expression, and red color, induced expression.

OLs-OH producing strains are more tolerant to the proton motive force uncoupler CCCP

The consistently reduced expression of the *pspF* iModulon in strains producing OLs-OH (Fig. 3;

perturbations A, B, and E), suggests that changes in membrane lipid composition could affect

how engineered strains manage proton membrane permeability, potentially providing a

mechanism for their increased tolerance to low pH. This iModulon comprises genes known as

Phage Shock Proteins, which are membrane proteins that play a critical role in regulating the

membrane permeability to protons. Recent research by Wang et al. (2021) indicated that *pspA*

mutants have a decreased ability to maintain the PMF during periods of starvation as well as its

participation in long-term stationary phase survival was shown (Takano et al. 2023).To

investigate this, we determined the effect of the presence of OLs-OH on the biomass production

of *E. coli* growing in the presence of the ionophore carbonyl cyanide m-chlorophenyl hydrazine

(CCCP), a compound that dissipates the proton gradient across the inner membrane (Kane et al.

2018). The strain producing OLs-OH exhibited better growth than the *E. coli* K12 strain carrying

the empty vector (Fig. 4a). Furthermore, the OL-producing strain showed enhanced tolerance to

CCCP, as it grew 35% faster than the control strain (Fig. 4b). These results indicate that the

presence of OLs-OHs in the *E. coli* membranes makes the cells membrane less proton permeable,

providing a possible mechanism for the low pH resistance of the engineered strains evaluated in

this work.

Figure 4. The presence of OLs in *E. coli* increases resistance to the ionophore CCCP. *E. coli* strains harboring the empty control plasmid or the plasmid containing *olsFC* were grown at pH 7.4 in MOPS/glucose medium without phosphate limitation. The growth curves are shown in (a), and the maximum specific growth rates are shown in (b). The data represent the average of three independent experiments. Error bars indicate the standard deviation using two-tailed unpaired Student's t-test. *p*-values *$p \leq 0.05$

OLs-OH production increases heterologous violacein production

We have shown that the presence of OLs in *E. coli* benefits its growth under both phosphate-limited and acidic pH growth conditions. These improved properties could enhance the performance of *E. coli* strains during heterologous expression, for example, during industrial applications. Previously, we reported that the proteome-reduced *E. coli* PFC strain produced higher levels of violacein than the wild-type strain, a pigment of industrial interest naturally synthesized by *Chromobacterium violaceum* (Lastiri-Pancardo et al. 2020). Therefore, and as a

proof-of-concept experiment, we tested the ability of our OLs- and OLs-OH-producing

engineered strains to synthesize violacein. The biosynthetic operon *vioABCDE* was cloned into

the plasmid pAJM336 under the control of an IPTG-inducible promoter. The resulting plasmid

pAJM336:*vioABCDE* (Fig 5a) was transformed into OL-producing derivatives of the proteome-

reduced mutants PFC and PYC. First, we determined that these strains, transformed with both

plasmids, produced unmodified and hydroxylated OLs (Fig S5). Then, we quantified violacein

production in the evaluated strains after 24 h at 30 °C in MOPS/glucose medium (pH 7.4) in

phosphate-limited conditions (0.4 mM phosphate), and MES/glucose medium (pH 5.8) under

phosphate-replete conditions (2.6 mM phosphate). At pH 7.4 and phosphate-limited conditions in

the absence of OLs, the triple mutant PFC accumulated more violacein than the triple mutant PYC

(Fig. 5b). Both strains, PFC and PYC, produced more than twice the violacein levels under

phosphate limitation when synthesizing OLs than the control strains (Fig 5b). At pH 5.8 with 2.6

mM phosphate, again comparing the strains harboring the empty plasmid with the strains

expressing *olsFC*, violacein production was more than 10-fold higher in strains forming OLs (Fig.

5c). These results demonstrate that our membrane engineering approach is beneficial for

producing costly metabolites via a heterologous pathway in a metabolic engineering application.

Figure 5. *E. coli* strains forming OLs-OH show an increase in total violacein production. All strains harbored the plasmid pJAM336:*vioABCDE* (a) in addition to pSEVA or p*olsFC*. Strains were grown in MOPS/glucose medium at pH 7.4 with 0.4 mM phosphate (b) and in MES/glucose medium at pH 5.8 with 2.6 mM phosphate (c), supplemented with 2 g/L tryptophan, 800 µM IPTG and 0.01% casamino acids (mean +/- s.d., n=9).

470
471

DISCUSSION

One of the main interests in synthetic biology is the design of robust microbial chassis resistant to different stressing conditions during biotechnological processes. One important mechanism of stress resistance is the specific modification of membrane composition(Sohlenkamp 2017). OLs are a group of phosphorus-free bacterial membrane lipids that have been described to have an important function during resistance to abiotic stress. The formation of OLs or an increase in their synthesis and modifications have been described in different bacteria under acid stress, increased temperatures, and phosphate limitation (Vences-Guzmán et al. 2011; Córdoba-Castro et al. 2021). Wild-type *E. coli* strains are unable to form OLs; therefore, in the present study, we proposed to modify the membrane lipid composition of *E. coli* by the expression of an OL synthase and an OL hydroxylase. A synthetic operon containing the genes *olsF* and *olsC* was constructed in a plasmid, transformed into different *E. coli* genetic backgrounds, and tested for its capacity to improve phenotypic characteristics of *E. coli* under conditions resembling an industrial process.

The expression of the synthetic operon *olsFC* led to the formation of OLs and OLs-OH in all four tested genetic backgrounds. The accumulation of OLs and OLs-OH caused a reduction in the concentration of the zwitterionic glycerophospholipid PE, indicating that *E. coli* cells have a mechanism for sensing and regulating the amount of these lipids in the membrane (Fig 1). Such reciprocal behavior of OL and PE concentrations has also been described in *S. proteamaculans*, *V. cholerae*, and *P. fluorescens*, among others, when they are grown under phosphate-limited conditions (Vences-Guzmán et al. 2015; Escobedo-Hinojosa et al. 2015; Barbosa et al. 2018).

Three of the four strains used in this study are deficient in the gene *phoB*; the *phoB* mutant*,* and the two proteome-reduced PYC and PFC strains (Lastiri-Pancardo et al. 2020), which are therefore unable to induce a proper phosphate scavenging response. In defined media such as those used here, these scavenging responses should have a cost and no benefit, as no other phosphate sources are present. Phosphate is the fifth most important element in the cell and one of the most expensive nutrients in bacterial cultures at an industrial level (Santos-Beneit 2015).

Our results show that by replacing phospholipids with ornithine lipids the engineered strains

change their cellular phosphate requirements yielding more biomass under this nutrient-limited

condition.

Bacteria are frequently exposed to acid stress. To survive these unfavorable conditions, they have

evolved a set of resistance mechanisms, which include the pumping of protons out of the

cytoplasm, the production of ammonia, proton-consuming decarboxylation reactions, and

modifications of the membrane lipid composition (Sohlenkamp 2017). *R. tropici* mutants

deficient in the OL hydroxylase *olsC* and lacking the hydroxylation of the C2 of the secondary

fatty acid of OLs are strongly affected in their growth at pH 4.5 (Vences-Guzmán et al. 2011). The

formation of OLs induced the expression of genes related to the arginine synthesis (Fig 3c,

perturbations C, D, and E). Because ornithine is an intermediate of arginine synthesis in *E. coli*

and is also used as a precursor for OL synthesis (Warneke et al. 2023), we hypothesize that the

expression of OlsFC may cause arginine limitation. This perceived low concentration of

intracellular arginine could explain the induction of arginine synthesis genes in the presence of

OLs. Furthermore, a lower amount of free arginine would also reduce the efficacy of the XAR

system(Iyer et al. 2003), whose coding genes *adiY* and *adiC* were upregulated at low pH and in

the presence of OLs (Fig 3c). Our results in transcriptional profiles showed that there is a mild

transcriptional response to the presence of OLs in *E. coli*, but specific changes provide the

possible mechanisms to improve the phenotype at low pH. In strains producing OLs-OH,

consistent under expression of the *pspF* iModulon and increased tolerance to CCCP (Fig 4a),

suggest that changes in membrane composition affected how engineered strains manage proton

membrane permeability, potentially providing a mechanism for their increased tolerance to low

pH.

To be able to design specific membrane properties in the future, further experiments are needed

to describe and understand the changes in membrane characteristics, such as permeability and

fluidity, etc., caused by the presence of OLs. Another area of opportunity is how the engineering

of the bacterial membrane affects the resistance to other types of stresses, such as osmotic

pressure, shear stress, or temperature changes that can be present in large-scale

fermentations(Wehrs et al. 2019). The abiotic stress response involves cellular mechanisms

beyond the change in membrane composition, and the combination of our approach with others

should aid in the generation of highly tolerant microbial cell factories.

Our study provides a solid basis to demonstrate that engineering the membrane composition is a

viable option to increase bacterial stress resistance, such as to phosphate limitation and to low pH

stress. We demonstrate the advantage of the approach by expressing a heterologous metabolite

from a costly pathway in proteome-reduced strains of *E. coli*. Finally, we show for the first time

that the presence of OLs-OH seems to reduce the permeability of the internal membrane to

protons, and we provide a possible mechanism for the enhanced tolerance to acid stress by

modifying the membrane composition. Changes in membrane permeability, such as those studied

here, should be further investigated. Other modified ornithine lipids can be further investigated

for this and other membrane properties.

Acknowledgments

We thank Miguel Angel Vences-Guzmán for technical support. LPBP was a recipient of the Postdoctoral

Scholarship Program in the UNAM-DGAPA. This work was supported by grants from PAPIIT-UNAM

IN214923 to JU and by grants from CONACyT (237713 and 425886) and PAPIIT UNAM (IN208116 and

IN208319) to CS.

Author contributions

The project was conceived, supervised, and guided by JU and CS. The experiments, data processing,

statistical analyses, and figures were carried out by LPBP. RNAseq data processing was carried out by

MSP and AAV, RNAseq data analysis and figures were carried out by AAV. Critical analysis of

experimental data and RNAseq categorization were performed by LPBP, JU, and CS. The manuscript was

written by LPBP, JU, and CS.

Conflict of interest

The authors declare no competing interests.

Data availability

The data that support the findings of this study are available within the article and in its supplementary

information. The accession number for the RNAseq data reported in this study is GEO: GSE233657.

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE233657).

REFERENCES

Aquino P, Honda B, Jaini S, Lyubetskaya A, Hosur K, Chiu JG, Ekladious I, Hu D, Jin L, Sayeg MK,
Stettner AI, Wang J, Wong BG, Wong WS, Alexander SL, Ba C, Bensussen SI, Bernstein DB,
Braff D, Cha S, Cheng DI, Cho JH, Chou K, Chuang J, Gastler DE, Grasso DJ, Greifenberger JS,
Guo C, Hawes AK, Israni D V., Jain SR, Kim J, Lei J, Li H, Li D, Li Q, Mancuso CP, Mao N,
Masud SF, Meisel CL, Mi J, Nykyforchyn CS, Park M, Peterson HM, Ramirez AK, Reynolds DS,
Rim NG, Saffie JC, Su H, Su WR, Su Y, Sun M, Thommes MM, Tu T, Varongchayakul N, Wagner
TE, Weinberg BH, Yang R, Yaroslavsky A, Yoon C, Zhao Y, Zollinger AJ, Stringer AM, Foster
JW, Wade J, Raman S, Broude N, Wong WW, Galagan JE (2017) Coordinated regulation of acid
resistance in Escherichia coli. BMC Syst Biol 11. https://doi.org/10.1186/s12918-016-0376-y

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori
H (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio
collection. Mol Syst Biol 2. https://doi.org/10.1038/msb4100050

Barbosa LC, Goulart CL, Avellar MM, Bisch PM, von Kruger WMA (2018) Accumulation of ornithine
lipids in Vibrio cholerae under phosphate deprivation is dependent on VC0489 (OlsF) and PhoBR
system. Microbiology (United Kingdom) 164:395–399. https://doi.org/10.1099/mic.0.000607

Beites T, Mendes M V. (2015) Chassis optimization as a cornerstone for the application of synthetic
biology based strategies in microbial secondary metabolism. Front Microbiol 6.
https://doi.org/10.3389/fmicb.2015.00906

Bligh EG, Dyer WJ (1959) A RAPID METHOD OF TOTAL LIPID EXTRACTION AND
PURIFICATION. Can J Biochem Physiol 37:911–917. https://doi.org/10.1139/o59-099

Calero P, Nikel PI (2019) Chasing bacterial *chassis* for metabolic engineering: a perspective review from
classical to non-traditional microorganisms. Microb Biotechnol 12:98–124.
https://doi.org/10.1111/1751-7915.13292

Cardoso VM, Campani G, Santos MP, Silva GG, Pires MC, Gonçalves VM, de C. Giordano R, Sargo CR,
Horta ACL, Zangirolami TC (2020) Cost analysis based on bioreactor cultivation conditions:
Production of a soluble recombinant protein using Escherichia coli BL21(DE3). Biotechnology
Reports 26:e00441. https://doi.org/10.1016/j.btre.2020.e00441

Chi H, Wang X, Shao Y, Qin Y, Deng Z, Wang L, Chen S (2019) Engineering and modification of
microbial chassis for systems and synthetic biology. Synth Syst Biotechnol 4:25–33

Cordell WT, Avolio G, Takors R, Pfleger BF (2023) Milligrams to kilograms: making microbes work at
scale. Trends Biotechnol. https://doi.org/10.1016/J.TIBTECH.2023.05.002

Córdoba-Castro LA, Salgado-Morales R, Torres M, Martínez-Aguilar L, Lozano L, Vences-Guzmán MÁ,
Guan Z, Dantán-González E, Serrano M, Sohlenkamp C (2021) Ornithine Lipids in Burkholderia
spp. Pathogenicity. Front Mol Biosci 7. https://doi.org/10.3389/fmolb.2020.610932

Darlington APS, Kim J, Jiménez JI, Bates DG (2018) Dynamic allocation of orthogonal ribosomes
facilitates uncoupling of co-expressed genes. Nat Commun 9. https://doi.org/10.1038/s41467-018-
02898-6

de la Cruz M, Ramírez EA, Sigala J-C, Utrilla J, Lara AR (2020) Plasmid DNA Production in Proteome-
Reduced Escherichia coli. Microorganisms 8:1444.
https://doi.org/10.3390/microorganisms8091444

Escobedo-Hinojosa WI, Vences-Guzmán MÁ, Schubotz F, Sandoval-Calderón M, Summons RE, López-
Lara IM, Geiger O, Sohlenkamp C (2015) OlsG (Sinac_1600) Is an Ornithine Lipid N-
Methyltransferase from the Planctomycete Singulisphaera acidiphila. Journal of Biological
Chemistry 290:15102–15111. https://doi.org/10.1074/jbc.M115.639575

Guerrero-Castro J, Lozano L, Sohlenkamp C (2018) Dissecting the Acid Stress Response of Rhizobium tropici CIAT 899. Front Microbiol 9. https://doi.org/10.3389/fmicb.2018.00846

Hidalgo D, Martínez-Ortiz CA, Palsson BO, Jiménez JI, Utrilla J (2022) Regulatory perturbations of ribosome allocation in bacteria reshape the growth proteome with a trade-off in adaptation capacity. iScience 25. https://doi.org/10.1016/j.isci.2022.103879

Iyer R, Williams C, Miller C (2003) Arginine-Agmatine Antiporter in Extreme Acid Resistance in *Escherichia coli*. J Bacteriol 185:6556–6561. https://doi.org/10.1128/JB.185.22.6556-6561.2003

Kane MS, Paris A, Codron P, Cassereau J, Procaccio V, Lenaers G, Reynier P, Chevrollier A (2018) Current mechanistic insights into the CCCP-induced cell survival response. Biochem Pharmacol 148:100–110

Kim SK, Park SJ, Li XH, Choi YS, Im DS, Lee JH (2018) Bacterial ornithine lipid, a surrogate membrane lipid under phosphate-limiting conditions, plays important roles in bacterial persistence and interaction with host. Environ Microbiol 20:3992–4008. https://doi.org/10.1111/1462-2920.14430

Lamoureux CR, Decker KT, Sastry A V, Rychel K, Gao Y, McConn JL, Zielinski DC, Palsson BO (2023) A multi-scale expression and regulation knowledge base for *Escherichia coli*. Nucleic Acids Res. https://doi.org/10.1093/nar/gkad750

Lastiri-Pancardo G, Mercado-Hernández JS, Kim J, Jiménez JI, Utrilla J (2020) A quantitative method for proteome reallocation using minimal regulatory interventions. Nat Chem Biol 16:1026–1033. https://doi.org/10.1038/s41589-020-0593-y

López-Lara IM, Geiger O (2017) Bacterial lipid diversity. Biochim Biophys Acta Mol Cell Biol Lipids 1862:1287–1299

Marquez-Zavala E, Utrilla J (2023) Engineering resource allocation in artificially minimized cells: Is genome reduction the best strategy? Microb Biotechnol 16:990–999. https://doi.org/10.1111/1751-7915.14233

Nikaido H (2003) Molecular Basis of Bacterial Outer Membrane Permeability Revisited. Microbiology and Molecular Biology Reviews 67:593–656. https://doi.org/10.1128/MMBR.67.4.593-656.2003

Ogasawara H, Ishida Y, Yamada K, Yamamoto K, Ishihama A (2007) PdhR (Pyruvate Dehydrogenase Complex Regulator) Controls the Respiratory Electron Transport System in *Escherichia coli*. J Bacteriol 189:5534–5541. https://doi.org/10.1128/JB.00229-07

Rojas-Jiménez K, Sohlenkamp C, Geiger O, Martínez-Romero E, Werner D, Vinuesa P (2005) A ClC Chloride Channel Homolog and Ornithine-Containing Membrane Lipids of *Rhizobium tropici* CIAT899 Are Involved in Symbiotic Efficiency and Acid Tolerance. Molecular Plant-Microbe Interactions® 18:1175–1185. https://doi.org/10.1094/MPMI-18-1175

Rychel K, Decker K, Sastry A V, Phaneuf P V, Poudel S, Palsson BO (2021) iModulonDB: a knowledgebase of microbial transcriptional regulation derived from machine learning. Nucleic Acids Res 49:D112–D120. https://doi.org/10.1093/nar/gkaa810

Santos-Beneit F (2015) The Pho regulon: a huge regulatory network in bacteria. Front Microbiol 6. https://doi.org/10.3389/fmicb.2015.00402

Silva-Rocha R, Martínez-García E, Calles B, Chavarría M, Arce-Rodríguez A, de las Heras A, Páez-Espino AD, Durante-Rodríguez G, Kim J, Nikel PI, Platero R, de Lorenzo V (2013) The Standard European Vector Architecture (SEVA): a coherent platform for the analysis and deployment of complex prokaryotic phenotypes. Nucleic Acids Res 41:D666–D675. https://doi.org/10.1093/nar/gks1119

Sohlenkamp C (2017) Membrane Homeostasis in Bacteria upon pH Challenge. In: Biogenesis of Fatty Acids, Lipids and Membranes. Springer International Publishing, pp 1–13

Sohlenkamp C, Geiger O (2015) Bacterial membrane lipids: Diversity in structures and pathways. FEMS
Microbiol Rev 40:133–159

Swain PS, Stevenson K, Leary A, Montano-Gutierrez LF, Clark IBN, Vogel J, Pilizota T (2016) Inferring
time derivatives including cell growth rates using Gaussian processes. Nat Commun 7:13766.
https://doi.org/10.1038/ncomms13766

Takano S, Takahashi H, Yama Y, Miyazaki R, Furusawa C, Tsuru S (2023) Inference of transcriptome
signatures of Escherichia coli in long-term stationary phase. Sci Rep 13:5647.
https://doi.org/10.1038/s41598-023-32525-4

Vences-Guzmán MÁ, Guan Z, Escobedo-Hinojosa WI, Bermúdez-Barrientos JR, Geiger O, Sohlenkamp
C (2015) Discovery of a bifunctional acyltransferase responsible for ornithine lipid synthesis in
Serratia proteamaculans. Environ Microbiol 17:1487–1496. https://doi.org/10.1111/1462-
2920.12562

Vences-Guzmán MÁ, Guan Z, Ormeño-Orrillo E, González-Silva N, López-Lara IM, Martínez-Romero
E, Geiger O, Sohlenkamp C (2011) Hydroxylated ornithine lipids increase stress tolerance in
Rhizobium tropici CIAT899. Mol Microbiol 79:1496–1514. https://doi.org/10.1111/j.1365-
2958.2011.07535.x

Warneke R, Garbers TB, Herzberg C, Aschenbrandt G, Ficner R, Stülke J (2023) Ornithine is the central
intermediate in the arginine degradative pathway and its regulation in Bacillus subtilis. Journal of
Biological Chemistry 299:104944. https://doi.org/10.1016/j.jbc.2023.104944

Wehrs M, Tanjore D, Eng T, Lievense J, Pray TR, Mukhopadhyay A (2019) Engineering Robust
Production Microbes for Large-Scale Cultivation. Trends Microbiol 27:524–537.
https://doi.org/10.1016/j.tim.2019.01.006

Supplementary material

S1. Synthesis of ornithine lipids (OLs) in *E. coli* increases biomass production under phosphate-
limited conditions but the growth rate does not change. (a) dry weight concentration (mgDW/mL) and
(b) maximum specific growth rate are shown. Cells were grown in 96-well plates for 24 hours at pH 7.4 in
MOPS/glucose medium without phosphate limitation (2.6 mM) and with phosphate limited (0.4 mM). Each
strain was grown harboring the empty control plasmid or the plasmid containing *olsFC*. The data represent
the average of three independent experiments. The error bars indicate the standard deviation using a two-
tailed unpaired student's t-test.

Biomass yield Phosphate-Limited

688 S2. **The presence of ornithine lipids in *E. coli* strains improves their growth characteristics under**
689 **phosphate limitation.** Growth curves of *E. coli* K12 (a), *E. coli* mutant *phoB* (b), *E. coli* triple mutant PFC
690 (c), *E. coli* triple mutant PYC (d), in MOPS/glucose medium with 0.4 mM phosphate at pH 7.4.

Low pH

692   S3. **The presence of ornithine lipids in *E. coli* strains improves their growth characteristics at pH 5.8.**
693   Growth curves of *E. coli* K12 (a), *E. coli* mutant *phoB* (b), *E. coli* triple mutant PFC (c), *E. coli* triple mutant
694   PYC (d) in MES/glucose medium with 2.6 mM phosphate at pH 5.8.

696
697
S4. Numbers of iModulon in the different perturbations. This classification consists in the clustering of
genes that share an independent modulated signal. IModulons coverage is represented in dark green when
is ≥80% of genes, light green <80%, and beige when no changes in expression were observed.

S5. *E. coli* strains forming OLs and synthesizing violacein. Thin layer chromatography (TLC) analysis
of total lipids from *E. coli* strains forming OLs and harboring the violacein synthesis pathway. The TLC
plate was stained with 0.2% ninhydrin to detect primary amine-containing lipids which are in this case PE,
OL, and OLs-OH. All strains analyzed for their lipid composition harbored the plasmid
pJAM336:*vioABCDE* in addition to a second plasmid. Strains PFC and PYC with pSEVA (empty) and
p*olsFC* with pJAM336:*vioABCDE* grown in MOPS medium with 0.4 mM phosphate and pH 7.4 (left)
and MES 2.6 mM and pH 5.8 (right). PE: phosphatidylethanolamine; OL: unmodified ornithine lipid; OLs-
OH: hydroxylated ornithine lipid.
710

 Table S1. Strains and plasmids used in this study

STRAINS	GENOTYPE OR DESCRIPTION	REFERENCE
PFC	*E. coli* K12BW25113 with mutations in *phoB, yedW, cusR*	Lastiri-Pancardo *et al.*, 2020
PYC	*E. coli* K12BW25113 with mutations in *phoB, fhlC y cueR*	Lastiri-Pancardo *et al.,* 2020
PhoB	*E. coli* K12BW25113 with mutation in *phoB*	KEIO collection
E. coli K12 BW25113	*E. coli* K12BW25113 wild-type strain	Laboratory stock
E. coli K12 BW25113pSEVA	*E. coli* K12BW25113 wild-type strain harboring pSEVA631 Gm^r	This study
phoBpSEVA	*phoB* strain harboring pSEVA631 Gm^r	This study
PYCpSEVA	PYC strain harboring pSEVA631 Gm^r	This study
PFCpSEVA	PFC strain harboring pSEVA631 Gm^r	This study
PFCp*olsFC*	PFC strain harboring pSEVA:*olsFC* Amp^r	This study
PYCp*olsFC*	PYC strain harboring pSEVA:*olsFC* Amp^r	This study
PhoBp*olsFC*	*pho* strain harboring pSEVA:*olsFC* Amp^r	This study
K12p*olsFC*	*E. coli* K12BW25113 wild-type strain harboring pSEVA:*olsFC* Amp^r	This study
PYCpSEVA:*vio*	PYC strain harboring pSEVA Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PFCpSEVA:*vio*	PFC strain harboring pSEVA Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PYC*polsFC*:*vio*	PYC strain harboring pSEVA:*olsFC* Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PFC*polsFC*:*vio*	PFC strain harboring pSEVA:*olsFC* Amp^r and pAJM.336*pvioABCDE* Km^r	This study
pCCS98	*olsC* of *R. tropici* in pET9a	Vences *et al.* 2011
pET9a-2569	*olsF2569* of *Serratia proteomaculans* in pET9a	Vences *et al.* 2015
pSEVA63-Hvio	*vioABCDE* operon of *Chromobacterium* in pSEVA631	Darlintong *et al.* 2018

720   **Table S2. Oligonucleotides used in this study.** All sequences are written in 5' to 3' direction.

pSEVA*olsF*	
pSEVA_*olsF* REV	AGAGAGTCCAAGCTAAACATCTAGTATTTCTCCTCTTTCTCTAGTATTTAT
pSEVA_ *olsF* FWD	AGGACACCTGTATAGAGTAATACTAGAGCCAGGCATCGCA
Spro_2569*olsF* FWD	AGAAAGAGGAGAAATACTAGATGTTTAGCTTGGACTCTCT
Spro_2569*olsF* REV	TGCGATGCCTGGCTCTAGTATTACTCTATACAGGTGTCCTGT
pSEVA*olsFC*	
pSEVA_*olsFC* FWD	GAAACCGCCCGAAGCCCTGACGCAAAAAACCCCGCTTCGG
pSEVA_*olsFC* REV	TCCGTCATTCCAAACCTCCTATGCCTGGCTCTAGTATTACTCTATAC
pCCS98_*olsC* FWD	GTAATACTAGAGCCAGGCATAGGAGGTTTGGAATGACGGAGA
pCCS98_*olsC* REV	CCGAAGCGGGGTTTTTTGCGTCAGGGCTTCGGGCGGTTTC
pAJM.336:*vioABCDE*	
pAJM.336_*vioE* FWD	TTTTTTCTGCCTCGTGATACGCCTACTCGGTACCAAATTCCAGAAAAGAG
pAJM.336_*vioA* REV	TGCAGATATCGGAAGAATGCTTCATCTAGTATTTCCCCTCTTTCTCTAGTATTAAAC
vioA FWD	ACTAGAGAAAGAGGGGAAATACTAGATGAAGCATTCTTCCGATATCTGCATTGTCG
vioA REV	TAGTATTTCTCCTCTTTCTCTAGTATAGAGGATCCCCGGGTACCGAGCTC
vioB_E FWD	GAGCTCGGTACCCGGGGATCCTCTATACTAGAGAAAGAGGAGAAATACTAGATGAG
vioB_E REV	CTCTTTTCTGGAATTTGGTACCGAGTAGGCGTATCACGAGGCAGAAAAAA