Engineering of Robust Host Strains: Enhancing *Escherichia coli* Abiotic Stress Resistance through Ornithine Lipid Formation

Leidy Patricia Bedoya-Pérez[1], Alejandro Aguilar-Vera[1], José Utrilla[1*], Christian Sohlenkamp[1*]

[1]Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Av. Universidad s/n Col. Chamilpa, C.P. 62210, Cuernavaca, Mor., Mexico.

*Corresponding authors

Name: José Utrilla: utrilla@ccg.unam.mx

Name: Christian Sohlenkamp: chsohlen@ccg.unam.mx

Address: Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Av. Universidad s/n, Col. Chamilpa, C.P. 62210, Cuernavaca, Mor., Mexico.

Abstract

Escherichia coli is a common host for biotechnology and synthetic biology applications. During growth and fermentation, the microbes are often exposed to stress conditions, like variations in pH or solvent concentrations and it has been described that bacterial membranes play a key role in response to abiotic stresses. Ornithine lipids (OLs) are a group of membrane lipids whose presence and synthesis have been related to stress resistance in different bacteria. In this study, we engineered different *E. coli* strains to produce unmodified OLs and hydroxylated OLs by expressing the synthetic operon *olsFC*. Our results showed that OL formation improved pH resistance and increased biomass under phosphate limitation. Transcriptome analysis revealed that OL-forming strains differentially expressed stress and membrane-related genes. OL-producing strains also showed better growth in the presence of the ionophore carbonyl cyanide 3-chlorophenylhydrazone (CCCP), suggesting reduced proton leakiness. Furthermore, our engineered strains showed improved heterologous violacein production at phosphate limitation and low pH. Overall, this study demonstrates the potential of engineering the *E. coli* membrane composition for constructing robust hosts for biotechnology and synthetic biology applications.

Keywords: synthetic biology, ornithine lipids, membrane engineering, acid stress response, phosphate limitation response, violacein production, proton motive force.

Graphical abstract. *Escherichia coli* strains forming ornithine lipids in addition to glycerophospholipids show increased resistance to acidic pH, improved growth under phosphate-limiting conditions, and allowed a higher production of violacein in a heterologous expression system. Transcriptional profiling by RNAseq showed expression changes that suggest reduced proton leakiness explained by CCCP tolerance.

INTRODUCTION

The use of microorganisms as cell factories plays an important role in biotechnology to produce bulk chemicals, proteins, biofuels, pharmaceuticals, etc. During these fermentations, the microbes are often exposed to stress conditions, like variations in pH conditions or solvent concentrations[1]. It can also be desirable to grow the production strain under conditions different from standard growth conditions, for example, to use less phosphate in the fermentation broth or to grow the microbe at an increased temperature. Microbes with both efficient production and tolerance to stress conditions are desirable for industrial production processes, however, strains combining different properties are rare to occur in nature [2]. Such adapted microorganisms are named "microbial chassis" [3] and could be engineered using synthetic biology tools [4].

Escherichia coli is among the most frequently used bacteria because of its easy genetic manipulation, fast growth to high cell densities, and safe use [5]. Efforts have been made to construct improved *E. coli* strains for biotechnological applications. Recently, the computational tool ReProMin (Regulation-based Proteome Minimization) was designed to identify genetic interventions required to divert cellular resources invested in the expression of non-essential genes toward the increase in biosynthetic potential [6]. These predictions allowed the design of two triple mutants called PFC and PYC, with mutations in *phoB, fhlC y cueR* genes, and mutations in *phoB, yedW, cusR* genes, respectively, allowing a theoretical release of 0.5% of the *E. coli* proteome. These strains exhibited increased availability of cellular resources that may be of interest in the biotech industry allowing a larger production of recombinant proteins [6]. Both triple

mutants, PFC, and PYC, lack the *phoB* gene encoding PhoB, a transcription

factor central to the bacterial phosphate starvation response. PhoB activates the

expression of 33 genes, mainly related to the use of alternative sources of

phosphorus [7] [8]. Its absence in the two triple mutants could imply limitations in

their use on a large scale in minimal media with low phosphate concentration.

On the other side, the bacterial membrane has an important function during the

response of bacteria to abiotic stresses, with bacteria frequently adapting their

membrane lipid compositions [9]. A major component of *E. coli* membranes are

glycerophospholipids, which are composed of two fatty acids, a glycerol moiety,

a phosphate group, and a variable head group [10]. In recent years, a much wider

diversity of lipid structures has become evident in different bacteria. Although

glycerophospholipids are the best-known membrane lipids, it is now known that

many other lipids can be present in bacterial membranes, examples being

hopanoids, ornithine lipids, betaine lipids, sulfonolipids or sphingolipids [10]. Most

importantly, the lipid composition of a bacterial membrane in many cases is not

static and the presence of some membrane lipids, specifically hopanoids and

ornithine lipids, seems to form part of a stress response to (changing)

environmental conditions in some bacteria.

Ornithine lipids (OLs) are phosphorus-free membrane lipids relatively common in

bacteria but absent from eukaryotes and archaea [11]. They are composed of a 3-

hydroxy fatty acid amide-bound to the α-amino group of ornithine and of a second

fatty acyl group ester-linked to the 3-hydroxy position of the first fatty acid. This

basic OL structure can be modified by hydroxylation in different positions, by *N*-

methylation, or by taurine transfer. It was predicted that around 50% of

sequenced bacterial species can synthesize OLs under certain growth

conditions. These conditions include limiting phosphate concentrations, acidic

pH, and increased temperatures [10] [12,13] and several bacteria such as

Pseudomonas fluorescens, *Serratia proteamaculans* and *Vibrio cholerae* only

form OLs under phosphate-limiting conditions. In *Burkholderia cepacia,* an

increased formation of hydroxylated OLs (OLs-OH) was observed in cells grown

at 40 °C, and in the symbiotic alpha-proteobacterium *Rhizobium tropici* CIAT899

it has been described that the presence of OLs-OH improves growth at increased

temperatures and under acidic conditions [12,14]. Especially, the hydroxylation

introduced by the OL hydroxylase OlsC in the C2 position of the secondary fatty

acid in OLs may increase hydrogen bonding between the lipid headgroups

thereby increasing acid stress resistance [15]. In recent years, many enzymes

involved in the synthesis and modification of OLs have been described [9]. One of

them is OlsF, a bifunctional OL synthase that was first described in *S.*

proteamaculans. When OlsF is expressed in *E. coli*, the resulting strains can form

OLs in one step [13,16].

Wild-type *E. coli* cannot form OLs and in its inner membrane the

glycerophospholipids, phosphatidylethanolamine (PE), phosphatidylglycerol

(PG), and cardiolipin (CP) can be detected. This composition is relatively stable

under different growth conditions. In the present study, we hypothesized that the

formation of OLs and hydroxylated OLs (OLs-OH) in *E. coli* strains might improve

the resistance of the hosts to different abiotic stress conditions. We constructed

the synthetic operon pSEVA*olsFC* (p*olsFC*) composed of the genes o*lsF* from *S.*

proteamaculans and *olsC* from *R. tropici* CIAT899 required to produce

unmodified and hydroxylated OLs, respectively [16,17] and expressed it in different

genetic backgrounds of *E. coli*. We observed increased biomass formation under

phosphate limitation and improved growth with a shortened lag phase at reduced

pH values. Transcriptome analysis revealed that OL-forming strains differentially

expressed stress and membrane-related genes, indicating a reduced proton

leakiness. Consistently, it was observed that OL-producing strains showed better

growth in the presence of the ionophore carbonyl cyanide 3-

chlorophenylhydrazone (CCCP). Finally, we could show that the formation of

OLs-OH in *E. coli* strains improved violacein formation in a heterologous

expression system compared to *E. coli* strain not forming OLs.

RESULTS AND DISCUSSION

Construction of *E. coli* strains synthesizing ornithine lipids and hydroxylated ornithine lipids

The success of many synthetic biology and biotechnology applications depends on the presence of a robust microbial chassis. The modification of bacterial membrane lipid composition is a promising way to obtain strains with improved properties. OLs are a group of phosphorus-free bacterial membrane lipids that have an important function during bacterial resistance to abiotic stress, including acid stress. *E. coli* lacks genes encoding enzymes required for OL synthesis, but *E. coli* strains forming OLs can be constructed [13,16]. We wanted to test how the presence of OLs in *E. coli* would affect its stress resistance and its performance as a host for synthetic biology applications. First, we constructed a plasmid-borne synthetic operon containing the genes *olsF* and *olsC* required for the synthesis of OLs and hydroxylated OLs (OLs-OH) [16,17] (Fig. 1A). This plasmid called p*olsFC* and the empty control plasmid pSEVA631 (pSEVA) [18] were transformed into four different *E. coli* strains: The wild-type strain *E. coli* K12 (BW25113), the triple mutants PFC (*phoB, fhlC* and *cueR*) and PYC (*phoB, yedW, cusR*), which have a reduced cellular proteome [6], and a *phoB*-deficient strain [19].

First, we wanted to determine if the *E. coli* strains expressing the synthetic operon formed OLs and OLs-OH. The strains were grown in a phosphate-limited MOPS medium, [^{14}C] acetate was added for lipid labeling, and total lipids were extracted and analyzed by thin-layer chromatography (TLC). In all strains, the glycerophospholipids phosphatidylethanolamine (PE), phosphatidylglycerol (PG) and cardiolipin (CL) were detected. In addition to the lipids PG, PE, and CL, the

E. coli strains expressing the synthetic operon produced both, unmodified OLs

and OLs-OH (Fig. 1B). A few lipids of minor abundance which were not identified

were observed in the TLC (labeled with asterisks). Interestingly, in *E. coli* strains

accumulating OLs and OLs-OH the formation of phosphatidylethanolamine (PE)

was reduced relative to the control strains with the empty plasmid (Fig. 1B).

Figure 1. Heterologous production of OLs and OLs-OH in *E. coli* strains. (A)

Synthetic operon containing *olsFC* under the control of a C6 Acyl homoserine Lactone

(AHL) inducible promoter in plasmid p*olsFC*. (B) TLC analysis of total membrane lipids

of *E. coli* strains K12, PYC, PFC, and *phoB* harboring pSEVA or p*olsFC*. Total lipids were

labeled with [^{14}C]acetate, extracted and separated by one-dimensional TLC. PE:

phosphatidylethanolamine, PG: phosphatidylglycerol, CL: cardiolipin, OLs: unmodified

ornithine lipid, OLs-OH: hydroxylated ornithine lipid, *: undetermined lipid.

OLs-OH production increases biomass yield under phosphate limitation

In some bacteria such as *S. proteamaculans* or *V. cholerae,* OLs are not formed

under phosphate-replete conditions, and their synthesis is induced by phosphate-

limiting growth conditions [20]. It is thought that the zwitterionic OLs replace the

zwitterionic phospholipid PE under conditions of phosphate limitation, reducing

the requirement for phosphate for the membrane, and thereby allowing an

alternative use for the phosphate. Phosphorus is a scarce mineral and one of the

most expensive nutrients in bacterial cultures at an industrial level [21]. Three of

the four strains we used in this study are deficient in the gene *phoB* and are

therefore unable to induce a proper phosphate scavenging response. As

described earlier, we observed a decrease in PE formation when OLs were

accumulated in the *E. coli* strains expressing the synthetic operon. Therefore, we

hypothesized that an *E. coli* strain being able to replace a fraction of its

phospholipids with OLs could have a growth advantage under phosphate-limiting

conditions, because phosphate normally used for membrane synthesis would be

liberated for other cellular processes.

We performed growth kinetics of all *E. coli* strains in MOPS/glucose medium with

different phosphate concentrations to define a phosphate concentration that

affected the growth of *E. coli* strains without severely limiting it. In MOPS/glucose

medium with 0.4 mM phosphate, slight growth inhibition was observed when

compared to non-limiting phosphate concentrations of standard MOPS/glucose

medium (2.6 mM). Phosphate concentrations below 0.4 mM severely affected the

growth of *E. coli* strains. Therefore, a phosphate concentration of 0.4 mM was

chosen for further experiments. In this set of experiments, we included the host

strains, the host strains with the empty plasmid, and the host strains expressing

the synthetic operon. We determined the effect of the presence of OLs-OH on

biomass production of *E. coli* grown under phosphate-limiting (0.4 mM) and

phosphate-replete (2.6 mM) concentrations. In MOPS/glucose medium with 2.6

mM phosphate, no difference in biomass formation was observed between the

different strains analyzed (Fig. S1A). In the phosphate-limited MOPS/glucose

medium, the biomass production in the strains that synthesize OLs increased by

23 to 39 % compared to the strains with the empty plasmid (Fig. 2A). In all cases,

the biomass formation of the OL-forming strains was higher than the wild-type

strains lacking a plasmid. These results suggest that the synthesis of OLs

provides *E. coli* an advantage when growing under growth-limiting phosphate

concentrations.

OLs-OH production improves growth phenotype at low pH

Bacteria are frequently exposed to acid stress[22]. To survive these unfavorable

conditions, they have evolved various resistance mechanisms, which include the

pumping of protons out of the cytoplasm, the production of ammonia, proton-

consuming decarboxylation reactions, and modifications of the membrane lipid

composition [21]. The presence of hydroxylated OLs has been related to increased

resistance to acid growth conditions, for example, the growth of a *R. tropici*

mutant deficient in the OL hydroxylase OlsC and lacking the hydroxylation of the

C2 of the secondary fatty acid of OLs is strongly affected in its growth at pH 4.5.

We wondered if the presence of OLs and hydroxylated OLs would improve growth

in MES/glucose medium without phosphate limitation at pH 5.8 in our engineered

E. coli strains. Again, we included the four host strains, the host strains with the

empty plasmid, and the host strains expressing the synthetic operon. During the

initial 6 hours, all strains (with or without OLs) exhibited similar growth kinetics.

However, a noticeable difference emerged when comparing the plasmid-

harboring strains in the subsequent hours. The presence of OLs in the *E. coli*

strains (triangles) significantly enhanced the maximum growth rate of the bacteria

at pH 5.8, resulting in approximately a 40% increase compared to three out of the

four genetic backgrounds examined (Fig. 2B to E). Furthermore, all four OL-

forming strains accumulated more biomass than the vector control strains (Fig.

2F). The *E. coli* strains harboring the empty vector controls reached the lowest

final optical density of all groups of strains. The OL-forming strains showed a

shortened lag phase compared to the four wild-type strains lacking plasmids. As

a control, we compared the maximum growth rate of the strains grown without

phosphate limitation at pH 7.4. No statistically significant differences were

observed between the strains at pH 7.4 (Fig S2).

Figure 2. The presence of ornithine lipids in *E. coli* strains improves their growth characteristics under phosphate limitation and at pH 5.8 (A) Dry weight concentration (mgDW/mL) of cultures after 24-hour growth in MOPS/glucose medium with low phosphate concentration (0.4 mM). Growth curves of *E. coli* K12 (B), *E. coli* triple mutant PFC (C), *E. coli* triple mutant PYC (D), and *E. coli* mutant *phoB* (E) in MES/glucose medium with 2.6 mM phosphate at pH 5.8. (F) Growth rates of the different *E. coli* strains were calculated from the growth curves in MES/glucose medium with 2.6

mM phosphate at pH 5.8. The data represent the average of three independent

experiments. The error bars indicate the standard deviation using a two-tailed unpaired

Student's t-test. P-values *P ≤ 0.05, ****P ≤0.0001.

4.- Transcriptional response to OLs expression at pH 7.4 and pH 5.8

We wondered if part of the positive phenotypic effects of the presence of OLs-

OH could be explained by transcriptional changes. To evaluate this, we compared

transcriptomes of *E. coli* K12.pSEVA and *E. coli* K12.p*olsFC* strains grown at pH

7.4 and pH 5.8 without phosphate limitation. Total RNA was extracted from

exponential growing cultures (11h). At this time point, a clear difference in growth

rate could be observed between the strains forming OLs and the strains lacking

them (Fig. 2B). We conducted RNA-Seq analysis on two independent biological

samples and sequenced the libraries to approximately 10 million read depth. We

then compared the expression profiles of these samples based on their response

to different perturbations: Low pH with and without OLs, response to OLs at low

pH, and response to OLs at neutral pH (Table 1).

Table 1. Transcriptional response to OLs expression at neutral and low pH.

Perturbations and descriptions

PERTURBATION	DESCRIPTION
A	Response of wild-type strain to low pH. Comparison between K12pSEVA pH 7.4 and K12pSEVA pH 5.8.
B	Differential response to low pH challenge and OLs formation. Comparison between K12pSEVA pH 7.4 and K12pOlsFC pH 5.8.
C	Response of OL-forming strain to low pH. Comparison between K12pOlsFC pH 7.4 and K12pOlsFC pH 5.8.
D	Response to the presence of OLs at pH 7.4. Comparison between K12pSEVA pH 7.4 and K12pOlsFC pH 7.4.
E	Response to the presence of OLs at pH 5.8. Comparison between K12pSEVA pH 5.8 and K12pOlsFC pH 5.8.

Overall, the results show that there is a large transcriptional response to low pH conditions (A, B, C), and a mild transcriptional response to OLs expression (B, D, E). We used the iModulon classification of the DEGs (differentially expressed genes) data, which consists of the clustering of genes sharing an independently modulated signal. The iModulons were obtained by independent component analysis (ICA) of 815 sample transcriptional profiles from 422 conditions for *E. coli* [23]. We considered only those iModulons that covered 80% or more of the DEGs for further analysis (fig. 3A). We observed changes in iModulons related to metabolism, stress response, two-component systems, energy production, and transcription/translation. Of special interest to us were the specific transcriptional changes observed in OLs expressing strains (fig. 3B, perturbations B, D, E) which occurred in the iModulons *pspF*, the *kdpE, proW* and *curli*. The iModulon *pspF* was of special interest because its involvement in membrane and proton motive

force maintenance. It is composed of genes encoding the so-called Phage Shock Proteins, which are membrane proteins that play a critical role in regulating the permeability of the membrane to protons. Recent research by Wang et al. (2021) indicated that *pspA* mutants had a decreased ability to maintain the Proton Motive Force (PMF) during periods of starvation [24]. Recently, also its participation in long-term stationary phase survival was shown[25]. Expression of the potassium two-component system *kdpE* was reduced in the presence of OLs, genes from efflux pumps such as *kdpE* were recently shown to be overrepresented in bacterial taxa with low pH preferences [26]. On the contrary, the expression of the glycine betaine *proVWX* and the *curli* iModulon were consistently overexpressed in OLs-producing strains. Furthermore, we analyzed individual genes, not classified by the iModulon analysis but related to ornithine synthesis, synthesis and turnover of membrane lipids, and response to acid stress as shown in Figure 3C. Of particular interest are the downregulation of *asr* (Fig 3C, perturbations A, B, and C), a highly charged, stress-induced, intrinsically disordered protein that contributes to the maintenance of outer membrane integrity under moderate acid stress [27]. Again, of special interest to us were the specific transcriptional changes observed in OLs expressing strains (fig. 3C, pertubations B, D, E). The genes *adiY/adiC* encoding an arginine/agmatine antiporter, of the arginine-dependent extreme acid resistance (XAR) were induced in the transcriptional response to low pH. AdiY/AdiC is one of several, decarboxylase-antiporter systems that protect *E. coli* from exposure to the acid environment [28,29]. Ornithine is an intermediate of arginine synthesis in *E. coli* and we hypothesized that the expression of OlsFC which uses ornithine as a precursor for OL synthesis might cause arginine limitation. This perceived low concentration of intracellular

arginine might explain the induction of arginine synthesis genes in pertubations

B and D. A lower amount of free arginine would also reduce the efficacy of the

XAR system [28]. Interestingly, some genes known to be involved in the acid stress

response are induced by the formation of OLs, such as *gadA*, *hdeD* and *yhiM*.

This differential expression might possibly contribute to the shortening of the lag

phase at pH 5.8 in OL-forming strains. By performing a comprehensive

transcriptional profiling experiment, we showed that modifications of the

membrane composition by OLs change the regulatory response to low pH.

Figure 3. Low pH conditions cause a large transcriptional response whereas OL formation causes a mild transcriptional response. (A) 48 iModulons with coverage greater than 80% in at least one perturbation. The functional categories are denoted according to color, yellow: metabolism, black: stress response, and white: other functions. (B) Differential expression of each analyzed iModulon. (C) Expression changes of individual genes involved in arginine metabolism, membrane and lipid synthesis, and stress response to low pH not classified by iModulons. (value= DEGup - DEGdown), blue color, reduced expression, and red color, induced expression.

OLs-OH producing strains are more tolerant to the proton motive force uncoupler CCCP

The consistently reduced expression of the *pspF* iModulon in strains producing OLs suggests that changes in membrane composition could affect how engineered strains manage proton membrane permeability, potentially providing a mechanism for their increased tolerance to low pH. To investigate this possible mechanism, we performed growth experiments using the ionophore carbonyl cyanide m-chlorophenyl hydrazine (CCCP), a compound that dissipates the proton gradient across the inner membrane. Our results show that strains producing OLs can grow 36% faster than strains harboring the empty plasmid (Fig. 4). The growth profiles of the strains are similar to those observed in the low pH experiments where the OLs-OH producing strains start growing at the same rate as the control but then they separate around the 5th hour, and they ultimately have an overall faster growth rate and higher biomass production. This result indicates that the presence of OLs-OHs in the *E. coli* membranes makes the cells

less susceptible to the ionophore CCCP providing a mechanism for the low pH

resistance of the engineered strains.

Figure 4. The presence of OLs in _E. coli_ increases resistance to the ionophore

CCCP. _E. coli_ strains harboring the empty control plasmid or the plasmid containing

olsFC were grown at pH 7.4 in MOPS/glucose medium without phosphate limitation. The

growth curves are shown in (A), and the maximum specific growth rates in (B). The data

represent the average of three independent experiments. The error bars indicate the

standard deviation using a two-tailed unpaired Student's t-test. P-values *P ≤ 0.05

6.- OLs-OH expression increases heterologous violacein production

We have shown here that the presence of OLs in _E. coli_ benefits its growth under

both phosphate-limiting and acidic pH growth conditions. These improved

properties could enhance the behavior of _E. coli_ strains during heterologous

expression, for example during an industrial application. As a proof-of-concept

experiment, we tested the ability of our engineered strains to synthesize violacein.

Violacein is a pigment naturally synthesized by _Chromobacterium violaceum_ with

antiviral, antiparasitic, and antibacterial activities, and it has several applications at the industrial level. This compound is synthesized in a five-step metabolic pathway using tryptophan as a precursor [30]. The biosynthetic operon *vioABCDE* was cloned into the plasmid pAJM336 under the control of an IPTG inducible promoter. The resulting plasmid pAJM336:*vioABCDE* was transformed into OLs-producing derivatives of the triple mutants PFC and PYC which had been obtained by the ReProMin analysis. First, we showed that these strains, transformed with both plasmids, were producing unmodified and hydroxylated OLs (Fig S2). We quantified violacein production in the evaluated strains after 24 hours at 30 °C in MOPS/glucose medium (pH 7.4) and MES/glucose medium (pH 5.8) under phosphate-replete conditions (2.6 mM phosphate) and in MOPS/glucose medium (pH 7.4) in phosphate-limiting conditions (0.4 mM phosphate). At pH 7.4 in the absence of OLs, the triple mutant PFC accumulated more violacein than the triple mutant PYC (Fig. 5A). The presence of OLs at pH 7.4, increased the amount of violacein formed by the triple mutant PYC (Fig. 5A). At pH 7.4 under phosphate-limiting conditions, more violacein formation can be detected in the OL-forming strains than in the vector control strains (Fig. 5B). At pH 5.8 with 2.6 mM phosphate, again comparing the strains harboring the empty plasmid with the strains expressing *olsFC* violacein formation was more than 10-fold higher in strains forming OLs (Fig. 5C).

In both low-phosphate and low-pH conditions, the presence of OLs-OH increased violacein production compared to the control. These results show that our membrane engineering approach is beneficial for producing complicated metabolites via a heterologous pathway.

Figure 5. *E. coli* **strains forming OLs-OH show an increase in total violacein production.** All strains harbored the plasmid pJAM336:*vioABCDE* (A) in addition to pSEVA or p*olsFC*. Strains were grown in MOPS/glucose medium at pH 7.4 with 0.4 mM phosphate (B) and in MES/glucose medium at pH 5.8 with 2.6 mM phosphate (C), supplemented with 2 g/L tryptophan, 800 µM IPTG and 0.01% casamino acids (mean +/- s.d., n=9).

386
387

CONCLUSIONS

Our study provides a solid basis to demonstrate the applicability of engineering of the membrane composition to reduce phosphate need, to increase tolerance to low pH stress and we demonstrate the advantage of the approach by expressing a heterologous metabolite from a costly pathway in proteome-reduced strains of *E. coli.*

We show for the first time that the presence of OLs-OH seems to reduce the permeability of the internal membrane to protons, and we provide a possible mechanism for the enhanced tolerance to acid stress by the modification of the membrane composition.

To be able to design specific membrane properties in the future, further experiments are needed to describe and understand the changes in membrane characteristics such as permeability and fluidity, etc. caused by the presence of OLs. Another area of opportunity is how the engineering of the bacterial membrane affects the resistance to other kinds of stresses, such as osmotic pressure, shear stress, or temperature changes that can be present in large-scale fermentations [31]. The abiotic stress response involves cellular mechanisms beyond the change in membrane composition and the combination of our approach with others should aid in the generation of highly tolerant microbial cell factories.

MATERIALS AND METHODS

Strain, plasmids and growth conditions

Bacterial strains and plasmids used are listed in supplementary table 1 and the oligonucleotides used are listed in supplementary table 2.

Plasmid pSEVA63 confers ampicillin resistance and has the replication system of the plasmid pBBR1 which is regarded as a "medium copy" plasmid. It was used to construct pSEVA*OlsFC* containing the *olsF* gene (1722 bp) from *Serratia proteamaculans* and the *olsC* gene (846 bp) from *Rhizobium tropici.* The *synthetic olsFC* operon has a ribosome binding site of 12 bp between both genes. The transcription of this operon is regulated by C6-.AHL (acyl homoserine lactone) and LuxR, but the system has some basal level expression even without induction. Therefore, we decided to use the empty plasmid pSEVA63 as negative control for our studies.

The plasmid pAJM.336Vio confers kanamycin resistance and contains the genes *vioA* (1257 bp), *vioB* (2994 bp), *vioC* (1290 bp), *vioD* (1122 bp), *vioE* (576 bp) from *Chromobacterium violaceum* encoding the enzymes required for violacein synthesis using tryptophan as a precursor.

Gibson assembly was used for plasmid construction. 10 µL of 2X NEBuilder HiFi DNA Assembly Master Mix were incubated with one or more DNA fragments, 0.05 pmol each, in a final volume of 20 µL at 50°C for 60 minutes. TOP10 chemically competent *E. coli* were transformed with 2 µL of the assembled products and incubated overnight at 37°C after transformation.

LB medium (Sigma), LB medium supplemented with 1 g/L glucose and MOPS

minimal media supplemented with 0.01% casamino acids (CAA) (Sigma), and 4

g/L glucose and K_2HPO_4 (0.4 or 2.6 mM) were used[32]. For low pH medium MOPS

buffer was replaced with the same concentration of MES buffer (MES/glucose

medium). Antibiotics were used in the following concentrations: gentamicin 25

µg/mL, spectinomycin (Spc) 50 µg/mL, ampicillin 100 µg/mL, and kanamycin 50

µg/mL.

Bacterial growth experiments

For the evaluation of bacterial growth at different phosphate concentrations, we

used MOPS/glucose medium. For growth experiments at pH 5.8 MES/glucose

medium was used. Cells were cultured overnight in MOPS/glucose medium

without phosphate limitation at 37 °C and 250 rpm. The next day cells were

washed three times with the receptor medium (phosphate limitation or low pH)

and cells were again cultured overnight. Then, strains were diluted to an OD_{600}

of 0.05 in fresh medium and 150 µL of the fresh culture were transferred to a

transparent 96-well plate (Corning) and incubated at 30 °C with fast linear shaking

in a microplate reader (Synergy 2.0, BioTek) for 24 hours, taking measurements

of the OD_{600} every 20 min. In all growth experiments, three replicates were

included, and the experiment was repeated independently on three different

days.

For the evaluation of bacterial tolerance to the proton motive force uncoupler

CCCP, we grew the strains in MOPS/glucose medium with 10µM CCCP in

transparent 96-well plates as described above.

The analysis of the growth kinetics of the plate experiments was conducted using the algorithm developed by Swain *et al.*, with the default parameters running on Python [33]. From this analysis, the magnitude and time of the maximum growth rate and the maximum OD_{600} were obtained. The p-value was determined using a two-tailed Student t-test as the statistical significance reliability index [34].

Lipid analysis

The lipid compositions of bacterial strains were determined following labeling with [^{14}C]acetate (Amersham Biosciences). Cultures (1 mL) of wild-type and mutant strains were inoculated from pre-cultures grown in the same medium. After the addition of 0.5 mCi of [^{14}C]acetate (60 mCi mmol^{-1}) to each culture, the cultures were incubated for 4 h. The cells were harvested by centrifugation, washed with 500 μL of water, and resuspended in 100 μL of water. Lipid extracts were obtained according to Bligh and Dyer (1959). Aliquots of the lipid extracts were spotted on high-performance TLC silica gel 60 (Merck, Poole, UK) plates and were separated in one dimension using chloroform/ methanol/water (150:30:8, v/v) as mobile phase. To visualize the membrane lipids, developed one-dimensional TLC plates were exposed to a PhosphorImager screen (Amersham Biosciences). The individual lipids were quantified using ImageQuant software (Amersham Biosciences). Primary amine-containing lipids (PE, OL, OL-OH) were visualized by spraying the plates with a solution of 0.2% (w/v) ninhydrin in acetone and subsequent treatment at 120°C for 10 minutes [12].

Characterization of violacein-producing strains.

Strains were inoculated into MOPS/glucose medium supplemented with gentamicin (20 µL/mL) and grown overnight. The next day cells were washed three times with the receptor medium (MOPS/glucose, phosphate limitation or MES/glucose) and then inoculated at an OD_{600} of 0.05 in fresh receptor medium supplemented with tryptophan (2.0 g/L). Then 150 µL of the fresh culture were transferred to a 96-well plate and incubated as described above. After 24 h the plates were centrifuged (13,000×g, 10 min), and the supernatant of each well was discarded. Violacein was extracted by suspending the pellet in each well in 200 µL absolute ethanol and incubating the plate at 95 °C for 10 min followed by pelleting cell debris (13,000×g, 10 min). The violacein present in the extract was determined spectrophotometrically at 575 nm in a microplate reader (Synergy 2.0, BioTek) [6].

RNA extraction for transcriptome analysis

E. coli K12pSEVA, and K12pOlsFC cells were pre-grown at 37°C in 25 mL of MOPS minimal medium with 4 g/L glucose overnight. The cells were washed three-time and cultured in MES/glucose medium. During exponential growth (11h), 3 mL were transferred to a tube, and 3 mL of RNA protect reagent were added and mixed. Cells were harvested by centrifugation and the pellets were immediately frozen in liquid nitrogen. Total RNA was prepared using the RNeasy mini kit (QIAGEN, Hildesheim, Germany). Pellets were resuspended in RLT buffer supplemented with lysozyme (20 mg per mL; QIAGEN, Hildesheim, Germany) and containing Zirconium Oxide Beads (0.5 mm, Next Advance). Cells

were disrupted using the Bullet Blender Tissue Homogenizer (Next-Advance,

Troy, NY, USA) in impact-resistant 2mL tubes. Genomic DNA was eliminated by

digestion with RNase-free DNase (QIAGEN, Hildesheim, Germany) for 20 min at

room temperature. Final RNA concentrations were determined using a NanoDrop

(Thermo Scientific). The typical OD_{260} to OD_{280} ratio of RNA samples was

approximately 2.0. The integrity of RNA samples was verified by determining the

RNA integrity number (RINe) in a TapeStation 2200 instrument (Agilent

Technologies). Two independent total RNA extractions were obtained for each

condition and each one was analyzed separately [35][28].

RNA-Seq

The samples of RNA-seq were analyzed by Novogene (Sacramento, CA, USA).

A total amount of 1 μg RNA per sample was used as input material for the RNA

simple preparations. Sequencing libraries were generated using NEBNext®

UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) following the

manufacturer's recommendations and index codes were added to attribute

sequences to each sample. After rRNA removal, fragmentation was carried out

using divalent cations under elevated temperature in NEBNext First Strand

Synthesis Reaction Buffer (5X). First strand cDNA was synthesized using random

hexamer primer and M-MuLV Reverse Transcriptase (RNase H-). Second-strand

cDNA synthesis was subsequently performed using DNA Polymerase I and

RNase H. Remaining overhangs were converted into blunt ends via

exonuclease/polymerase activities. After adenylation of 3' ends of DNA

fragments, NEBNext Adaptor with hairpin loop structure were ligated to prepare

for hybridization. In order to select cDNA fragments of preferentially 150~200 bp

in length, the library fragments were purified with AMPure XP system (Beckman

Coulter, Beverly, USA). Then 3 µL USER Enzyme (NEB, USA) was used with

size-selected, adaptor- ligated cDNA at 37 °C for 15 min followed by 5 min at 95

°C before PCR. Then PCR was performed with Phusion High-Fidelity DNA

polymerase, Universal PCR primers and Index (X) Primer. At last, PCR products

were purified (AMPure XP system) and library quality was assessed on the

Agilent Bioanalyzer 2100 system. The clustering of the index-coded samples was

performed on an Illumina Novaseq sequencer according to the manufacturer's

instructions. After cluster generation, the libraries were sequenced on the same

machine and paired-end reads were generated.

Data processing and differential expression

The raw reads were processed for adapter trimming and quality control filtering

using Trim Galore (v.0.6.6). The reads were aligned using R package Rsubread

(v.2.2.6) and mapped to the reference genome (GCF_000005845.2). For gene

expression quantification the function featureCounts from Rsubread package

were used, generating the read counts matrix. We used the batch correction

ARSyseq and TMM for normalization. The differential expression analysis was

performed using DESeq2 (v.1.28.1). The filters used for the DEGs were: absolute

log2FoldChange >= 1 and p-value adjust <= 0.05. The accession number for the

RNAseq data reported in this paper is GEO: GSE233657

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE233657).

The iModulon analysis consists of adding the differential expression data to each

iModulon and the order within the graphs was made based on the "system

category" field [23,36].

Acknowledgment

We thank Mishael Perez for initial RNAseq processing and Miguel Angel Vences-

Guzmán for technical support. LPBP was a recipient of the Postdoctoral

Scholarship Program in the UNAM-DGAPA. This work was supported by grants

from PAPIIT-UNAM IN214923 to JU and by grants from CONACyT (237713 and

425886) and PAPIIT UNAM (IN208116 and IN208319) to CS.

REFERENCES

(1)   Cordell, W. T.; Avolio, G.; Takors, R.; Pfleger, B. F. Milligrams to Kilograms: Making Microbes Work at Scale. *Trends Biotechnol* **2023**. https://doi.org/10.1016/J.TIBTECH.2023.05.002.

(2)   Calero, P.; Nikel, P. I. Chasing Bacterial *Chassis* for Metabolic Engineering: A Perspective Review from Classical to Non-Traditional Microorganisms. *Microb Biotechnol* **2019**, *12* (1), 98–124. https://doi.org/10.1111/1751-7915.13292.

(3)   Beites, T.; Mendes, M. V. Chassis Optimization as a Cornerstone for the Application of Synthetic Biology Based Strategies in Microbial Secondary Metabolism. *Front Microbiol* **2015**, *6*. https://doi.org/10.3389/fmicb.2015.00906.

(4)   Marquez-Zavala, E.; Utrilla, J. Engineering Resource Allocation in Artificially Minimized Cells: Is Genome Reduction the Best Strategy? *Microb Biotechnol* **2023**, *16* (5), 990–999. https://doi.org/10.1111/1751-7915.14233.

(5)   Chi, H.; Wang, X.; Shao, Y.; Qin, Y.; Deng, Z.; Wang, L.; Chen, S. Engineering and Modification of Microbial Chassis for Systems and Synthetic Biology. *Synthetic and Systems Biotechnology*. KeAi Communications Co. March 1, 2019, pp 25–33. https://doi.org/10.1016/j.synbio.2018.12.001.

(6)   Lastiri-Pancardo, G.; Mercado-Hernández, J. S.; Kim, J.; Jiménez, J. I.; Utrilla, J. A Quantitative Method for Proteome Reallocation Using Minimal Regulatory Interventions. *Nat Chem Biol* **2020**, *16* (9), 1026–1033. https://doi.org/10.1038/s41589-020-0593-y.

(7)   Marzan, L.; Shimizu, K. Metabolic Regulation of Escherichia Coli and Its PhoB and PhoR Genes Knockout Mutants under Phosphate and Nitrogen Limitations as Well as at Acidic Condition. *Microb Cell Fact* **2011**, *10* (1), 39. https://doi.org/10.1186/1475-2859-10-39.

(8)   Bachhawat, P.; Swapna, G. V. T.; Montelione, G. T.; Stock, A. M. Mechanism of Activation for Transcription Factor PhoB Suggested by Different Modes of Dimerization in the Inactive and Active States. *Structure* **2005**, *13* (9), 1353–1363. https://doi.org/10.1016/j.str.2005.06.006.

(9)   Sohlenkamp, C. Membrane Homeostasis in Bacteria upon PH Challenge. In *Biogenesis of Fatty Acids, Lipids and Membranes*; Springer International Publishing, 2017; pp 1–13. https://doi.org/10.1007/978-3-319-43676-0_57-1.

(10)   Sohlenkamp, C.; Geiger, O. Bacterial Membrane Lipids: Diversity in Structures and Pathways. *FEMS Microbiology Reviews*. Oxford University Press September 5, 2015, pp 133–159. https://doi.org/10.1093/femsre/fuv008.

(11)   Kim, S. K.; Park, S. J.; Li, X. H.; Choi, Y. S.; Im, D. S.; Lee, J. H. Bacterial Ornithine Lipid, a Surrogate Membrane Lipid under Phosphate-Limiting Conditions, Plays Important Roles in Bacterial Persistence and Interaction with Host. *Environ Microbiol* **2018**, *20* (11), 3992–4008. https://doi.org/10.1111/1462-2920.14430.

(12) Vences-Guzmán, M. Á.; Guan, Z.; Ormeño-Orrillo, E.; González-Silva, N.; López-Lara, I.
M.; Martínez-Romero, E.; Geiger, O.; Sohlenkamp, C. Hydroxylated Ornithine Lipids
Increase Stress Tolerance in Rhizobium Tropici CIAT899. *Mol Microbiol* **2011**, *79* (6),
1496–1514. https://doi.org/10.1111/j.1365-2958.2011.07535.x.

(13) Vences-Guzmán, M. Á.; Guan, Z.; Escobedo-Hinojosa, W. I.; Bermúdez-Barrientos, J. R.;
Geiger, O.; Sohlenkamp, C. Discovery of a Bifunctional Acyltransferase Responsible for
Ornithine Lipid Synthesis in Serratia Proteamaculans. *Environ Microbiol* **2015**, *17* (5),
1487–1496. https://doi.org/10.1111/1462-2920.12562.

(14) Córdoba-Castro, L. A.; Salgado-Morales, R.; Torres, M.; Martínez-Aguilar, L.; Lozano, L.;
Vences-Guzmán, M. Á.; Guan, Z.; Dantán-González, E.; Serrano, M.; Sohlenkamp, C.
Ornithine Lipids in Burkholderia Spp. Pathogenicity. *Front Mol Biosci* **2021**, *7*.
https://doi.org/10.3389/fmolb.2020.610932.

(15) Nikaido, H. Molecular Basis of Bacterial Outer Membrane Permeability Revisited.
Microbiology and Molecular Biology Reviews **2003**, *67* (4), 593–656.
https://doi.org/10.1128/MMBR.67.4.593-656.2003.

(16) Escobedo-Hinojosa, W. I.; Vences-Guzmán, M. Á.; Schubotz, F.; Sandoval-Calderón, M.;
Summons, R. E.; López-Lara, I. M.; Geiger, O.; Sohlenkamp, C. OlsG (Sinac_1600) Is an
Ornithine Lipid N-Methyltransferase from the Planctomycete Singulisphaera Acidiphila.
Journal of Biological Chemistry **2015**, *290* (24), 15102–15111.
https://doi.org/10.1074/jbc.M115.639575.

(17) Rojas-Jiménez, K.; Sohlenkamp, C.; Geiger, O.; Martínez-Romero, E.; Werner, D.;
Vinuesa, P. A ClC Chloride Channel Homolog and Ornithine-Containing Membrane
Lipids of *Rhizobium Tropici* CIAT899 Are Involved in Symbiotic Efficiency and Acid
Tolerance. *Molecular Plant-Microbe Interactions®* **2005**, *18* (11), 1175–1185.
https://doi.org/10.1094/MPMI-18-1175.

(18) Silva-Rocha, R.; Martínez-García, E.; Calles, B.; Chavarría, M.; Arce-Rodríguez, A.; de las
Heras, A.; Páez-Espino, A. D.; Durante-Rodríguez, G.; Kim, J.; Nikel, P. I.; Platero, R.; de
Lorenzo, V. The Standard European Vector Architecture (SEVA): A Coherent Platform for
the Analysis and Deployment of Complex Prokaryotic Phenotypes. *Nucleic Acids Res*
2013, *41* (D1), D666–D675. https://doi.org/10.1093/nar/gks1119.

(19) Baba, T.; Ara, T.; Hasegawa, M.; Takai, Y.; Okumura, Y.; Baba, M.; Datsenko, K. A.;
Tomita, M.; Wanner, B. L.; Mori, H. Construction of *Escherichia Coli* K-12 In-frame,
Single-gene Knockout Mutants: The Keio Collection. *Mol Syst Biol* **2006**, *2* (1).
https://doi.org/10.1038/msb4100050.

(20) Barbosa, L. C.; Goulart, C. L.; Avellar, M. M.; Bisch, P. M.; von Kruger, W. M. A.
Accumulation of Ornithine Lipids in Vibrio Cholerae under Phosphate Deprivation Is
Dependent on VC0489 (OlsF) and PhoBR System. *Microbiology (United Kingdom)* **2018**,
164 (3), 395–399. https://doi.org/10.1099/mic.0.000607.

(21) Foster, J. W. Escherichia Coli Acid Resistance: Tales of an Amateur Acidophile. *Nat Rev*
Microbiol **2004**, *2* (11), 898–907. https://doi.org/10.1038/nrmicro1021.

(22) Guan, N.; Liu, L. Microbial Response to Acid Stress: Mechanisms and Applications. *Appl*
Microbiol Biotechnol **2020**, *104* (1), 51–65. https://doi.org/10.1007/s00253-019-10226-
1.

(23) Rychel, K.; Decker, K.; Sastry, A. V; Phaneuf, P. V; Poudel, S.; Palsson, B. O. IModulonDB:
A Knowledgebase of Microbial Transcriptional Regulation Derived from Machine
Learning. *Nucleic Acids Res* **2021**, *49* (D1), D112–D120.
https://doi.org/10.1093/nar/gkaa810.

(24) Wang, M.; Chan, E. W. C.; Wan, Y.; Wong, M. H. yin; Chen, S. Active Maintenance of
Proton Motive Force Mediates Starvation-Induced Bacterial Antibiotic Tolerance in
Escherichia Coli. *Commun Biol* **2021**, *4* (1). https://doi.org/10.1038/s42003-021-02612-
1.

(25) Takano, S.; Takahashi, H.; Yama, Y.; Miyazaki, R.; Furusawa, C.; Tsuru, S. Inference of
Transcriptome Signatures of Escherichia Coli in Long-Term Stationary Phase. *Sci Rep*
2023, *13* (1), 5647. https://doi.org/10.1038/s41598-023-32525-4.

(26) Ramoneda, J.; Stallard-Olivera, E.; Hoffert, M.; Winfrey, C. C.; Stadler, M.; Niño-García,
J. P.; Fierer, N. Building a Genome-Based Understanding of Bacterial PH Preferences. *Sci*
Adv **2023**, *9* (17). https://doi.org/10.1126/sciadv.adf8998.

(27) Ren, C.; Zheng, Y.; Liu, C.; Mencius, J.; Wu, Z.; Quan, S. Molecular Characterization of an
Intrinsically Disordered Chaperone Reveals Net-Charge Regulation in Chaperone Action.
J Mol Biol **2022**, *434* (5), 167405. https://doi.org/10.1016/j.jmb.2021.167405.

(28) Iyer, R.; Williams, C.; Miller, C. Arginine-Agmatine Antiporter in Extreme Acid Resistance
in *Escherichia Coli*. *J Bacteriol* **2003**, *185* (22), 6556–6561.
https://doi.org/10.1128/JB.185.22.6556-6561.2003.

(29) Tsai, M.-F.; Fang, Y.; Miller, C. Sided Functions of an Arginine–Agmatine Antiporter
Oriented in Liposomes. *Biochemistry* **2012**, *51* (8), 1577–1585.
https://doi.org/10.1021/bi201897t.

(30) Durán, N.; Justo, G. Z.; Durán, M.; Brocchi, M.; Cordi, L.; Tasic, L.; Castro, G. R.;
Nakazato, G. Advances in Chromobacterium Violaceum and Properties of Violacein-Its
Main Secondary Metabolite: A Review. *Biotechnology Advances*. Elsevier Inc.
September 1, 2016, pp 1030–1045. https://doi.org/10.1016/j.biotechadv.2016.06.003.

(31) Wehrs, M.; Tanjore, D.; Eng, T.; Lievense, J.; Pray, T. R.; Mukhopadhyay, A. Engineering
Robust Production Microbes for Large-Scale Cultivation. *Trends Microbiol* **2019**, *27* (6),
524–537. https://doi.org/10.1016/j.tim.2019.01.006.

(32) Neidhardt, F. C.; Bloch, P. L.; Smith, D. F. Culture Medium for Enterobacteria. *J Bacteriol*
1974, *119* (3), 736–747. https://doi.org/10.1128/jb.119.3.736-747.1974.

(33) Swain, P. S.; Stevenson, K.; Leary, A.; Montano-Gutierrez, L. F.; Clark, I. B. N.; Vogel, J.;
Pilizota, T. Inferring Time Derivatives Including Cell Growth Rates Using Gaussian
Processes. *Nat Commun* **2016**, *7* (1), 13766. https://doi.org/10.1038/ncomms13766.

(34) Hidalgo, D.; Martínez-Ortiz, C. A.; Palsson, B. O.; Jiménez, J. I.; Utrilla, J. Regulatory
Perturbations of Ribosome Allocation in Bacteria Reshape the Growth Proteome with a

Trade-off in Adaptation Capacity. *iScience* **2022**, *25* (3).
https://doi.org/10.1016/j.isci.2022.103879.

(35) Guerrero-Castro, J.; Lozano, L.; Sohlenkamp, C. Dissecting the Acid Stress Response of
Rhizobium Tropici CIAT 899. *Front Microbiol* **2018**, *9*.
https://doi.org/10.3389/fmicb.2018.00846.

(36) Lamoureux, C. R.; Decker, K. T.; Sastry, A. V; Rychel, K.; Gao, Y.; Mcconn, J. L.; Zielinski,
D. C.; Palsson, B. O. A Multi-Scale Transcriptional Regulatory Network Knowledge Base
for Escherichia Coli. https://doi.org/10.1101/2021.04.08.439047.

Supplementary material

Figure S1. Synthesis of ornithine lipids (OLs) in *E. coli* increases biomass

production under phosphate-limiting conditions. (A) Maximum specific growth rate

and (B) dry weight concentration (mgDW/mL) are shown. Cells were grown in 96-well

plates for 24 hours at pH 7.4 in MOPS/glucose medium without phosphate limitation (2.6

mM). Each strain was grown without plasmid, harboring the empty control plasmid or the

plasmid containing *olsFC*. The data represent the average of three independent

experiments. The error bars indicate the standard deviation using a two-tailed unpaired

student's t-test.

Figure S2. *E. coli* strains forming OL and synthesizing violacein. Thin layer chromatography (TLC) analysis of total lipids from *E. coli* strains forming OLs and harboring the violacein synthesis pathway. The TLC plate was stained with ninhydrin to detect primary amine-containing lipids which are in this case PE, OL, and OLs-OH. All strains analyzed for their lipid composition harbored the plasmid pJAM336:*vioABCDE* in addition to a second plasmid. Strains PFC and PYC with pSEVA (empty) and p*olsFC* with pJAM336:*vioABCDE* grown in MOPS medium with 0.4 mM phosphate and pH 7.4 (left) and MES 2.6 mM and pH 5.8 (right). PE: phosphatidylethanolamine; OL: unmodified ornithine lipid; OLs-OH:hydroxylated ornithine lipid.

720 **Figure S3. iModulon classification of pertubations.** This classification consists in the

721 clustering of genes that share an independent modulated signal. IModulons coverage is

722 represented in dark green when is ≥80% of genes, light green <80% and beige when no

723 changes in expression were observed.

729   **Table S1. Strains and plasmids used in this study**

STRAINS	GENOTYPE OR DESCRIPTIONS	REFERENCE
PFC	*E. coli* K12BW25113 with mutations in *phoB, yedW, cusR*	Lastiri-Pancardo *et al.*, 2020
PYC	*E. coli* K12BW25113 with mutations in *phoB, fhlC y cueR*	Lastiri-Pancardo *et al.*, 2020
PhoB	*E. coli* K12BW25113 with mutation in *phoB*	KEIO collection
E. coli K12 BW25113	*E. coli* K12BW25113 wild-type strain	Laboratory stock
E. coli K12 BW25113pSEVA	*E. coli* K12BW25113 wild-type strain harboring pSEVA631 Gm^r	This study
phoBpSEVA	*phoB* strain harboring pSEVA631 Gm^r	This study
PYCpSEVA	PYC strain harboring pSEVA631 Gm^r	This study
PFCpSEVA	PFC strain harboring pSEVA631 Gm^r	This study
PFCp*olsFC*	PFC strain harboring pSEVA:*olsFC* Amp^r	This study
PYCp*olsFC*	PYC strain harboring pSEVA:*olsFC* Amp^r	This study
PhoBp*olsFC*	*pho* strain harboring pSEVA:*olsFC* Amp^r	This study
K12p*olsFC*	*E. coli* K12BW25113 wild-type strain harboring pSEVA:*olsFC* Amp^r	This study
PYCpSEVA:*vioABCDE*	PYC strain harboring pSEVA Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PFCpSEVA:*vioABCDE*	PFC strain harboring pSEVA Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PYCp*olsFC*:*vioABCDE*	PYC strain harboring pSEVA:*olsFC* Amp^r and pAJM.336*pvioABCDE* Km^r	This study
PFCp*olsFC*:*vioABCDE*	PFC strain harboring pSEVA:*olsFC* Amp^r and pAJM.336*pvioABCDE* Km^r	This study

734 **Table S2. Oligonucleotides used in this study.** All sequences are written in 5' to 3'
735 direction.

pSEVA*olsF*	
pluxpR-T7/pTU1-A-lacZ REV	AGAGAGTCCAAGCTAAACATCTAGTATTTCTCCTCTTTCTCTAGTATTTAT
pluxpR-T7/pTU1-A-lacZ FWD	AGGACACCTGTATAGAGTAATACTAGAGCCAGGCATCGCA
Spro_2569olsF FWD	AGAAAGAGGAGAAATACTAGATGTTTAGCTTGGACTCTCT
Spro_2569olsF REV	TGCGATGCCTGGCTCTAGTATTACTCTATACAGGTGTCCTGT
pSEVA*olsFC*	
OlsF/Cind.WizGA FWD (1)	GAAACCGCCCGAAGCCCTGACGCAAAAAACCCCGCTTCGG
OlsF/Cind.WizGA REV (1)	TCCGTCATTCCAAACCTCCTATGCCTGGCTCTAGTATTACTCTATAC
OlsF/Cind.WizGA FWD (2)	GTAATACTAGAGCCAGGCATAGGAGGTTTGGAATGACGGAGA
OlsF/Cind.WizGA REV (2)	CCGAAGCGGGGTTTTTTGCGTCAGGGCTTCGGGCGGTTC
pAJM.336:*vioABCDE*	
pAJM.336 FWD	TTTTTTCTGCCTCGTGATACGCCTACTCGGTACCAAATTCCAGAAAAGAG
pAJM.336 REV	TGCAGATATCGGAAGAATGCTTCATCTAGTATTTCCCCTCTTTCTCTAGTATTAAAC
pSEVA63-viogene FWD (1)	ACTAGAGAAAGAGGGGAAATACTAGATGAAGCATTCTTCCGATATCTGCATTGTCG
pSEVA63-viogene REV (1)	TAGTATTTCTCCTCTTTCTCTAGTATAGAGGATCCCCGGGTACCGAGCTC
pSEVA63-viogene FWD (2)	GAGCTCGGTACCCGGGGATCCTCTATACTAGAGAAAGAGGAGAAATACTAGATGAG
pSEVA63-viogene REV (2)	CTCTTTTCTGGAATTTGGTACCGAGTAGGCGTATCACGAGGCAGAAAAAA
revBvio(2)	TTTCTCTAGTAGGATCCTTAGGCCTCTCTAGA
fwCvio(2)	TACTAGAGAAAGAGGAGAAATACTAGATGAAAAGAGCAATC