Latent mutations in the ancestries of alleles under selection

Wai-Tong (Louis) Fan[a], John Wakeley[b]

[a]*Department of Mathematics, Indiana University, 831 East 3rd St, Bloomington, 47405, IN, USA*
[b]*Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Ave, Cambridge, 02138, MA, USA*

Abstract

We consider a single genetic locus with two alleles A_1 and A_2 in a large haploid population. The locus is subject to selection and two-way, or recurrent, mutation. Assuming the allele frequencies follow a Wright-Fisher diffusion and have reached stationarity, we describe the asymptotic behaviors of the conditional gene genealogy and the latent mutations of a sample with known allele counts, when the count n_1 of allele A_1 is fixed, and when either or both the sample size n and the selection strength $|\beta|$ tend to infinity. Our study extends previous work under neutrality to the case of non-neutral rare alleles, asserting that when selection is not too strong relative to the sample size, even if it is strongly positive or strongly negative in the usual sense ($\beta \to -\infty$ or $\beta \to +\infty$), the number of latent mutations of the n_1 copies of allele A_1 follows the same distribution as the number of alleles in the Ewens sampling formula. On the other hand, very strong positive selection relative to the sample size leads to neutral gene genealogies with a single ancient latent mutation. We also demonstrate robustness of our asymptotic results against changing population sizes, when one of $|\beta|$ or n is large.

Keywords: Recurrent mutation, selection, Ewens sampling formula, coalescent, Wright-Fisher diffusion

1. Introduction

The observed copies of a particular allele in a sample descend from an unknown number of distinct mutations. If k_1 is the number of these 'latent' mutations for allele A_1 when it is observed n_1 times in a sample, then $k_1 \in \{1, 2, \ldots, n_1\}$. Although latent mutations are not observed directly, they can be modeled as outcomes of the stochastic ancestral process of a sample and inferred from patterns of variation in DNA data (Harpak et al., 2016; Seplyarskiy et al., 2021; Johnson et al., 2022). Previously in Wakeley et al. (2023), we obtained results on the distribution and the timing of latent mutations for rare neutral alleles. Here we consider non-neutral alleles which may be under strong selection and which may or may not be rare. We take two different approaches to modeling latent mutations under selection and recurrent mutation. The first approach uses the idea of coalescence in a random background of allele frequencies in the population (Barton et al., 2004). The second uses the conditional ancestral selection graph (Slade, 2000a) and demonstrates results consistent with those from the first approach.

We assume there are two possible alleles, A_1 and A_2, at a single genetic locus in a large haploid population. We begin by assuming that the population size N is constant over time. In Section 3.4 we consider time-varying population size. One allele or the other is favored by directional selection. Mutation is recurrent and happens in both both directions. In the diffusion approximation, time is measured in proportion to N_e generations where N_e is the effective population size (Ewens, 2004). Under the Wright-Fisher model of reproduction, $N_e = N$. Under the Moran model of reproduction (Moran, 1958, 1962), $N_e = N/2$. With these assumptions, the frequency of A_1 alleles is well approximated by a process X that solves (1) below and has parameters θ_1, θ_2 and β as $N \to \infty$. For a haploid population, $\theta_i = 2N_e u_i$ and $\beta = 2N_e s$, in which u_i is the per-generation rate of $A_{3-i} \to A_i$ mutations and s is the selection coefficient. If there is no dominance, these results can be extended to diploids, in which case $\theta_i = 4N_e u_i$ and $\beta = 4N_e s$.

Thus, we assume that allele-frequency dynamics in the population obey the Wright-Fisher diffusion (Fisher, 1930; Wright, 1931; Ewens, 2004) with parameters θ_1 and θ_2 for the rates of $A_2 \to A_1$ and $A_1 \to A_2$ mutation, and β for the selective advantage (if $\beta > 0$) or disadvantage (if $\beta < 0$) of allele A_1. That is, we let $X(t)$ be the relative frequency of A_1 in the population at time t, and assume that its forward-time dynamics is described by the stochastic differential equation

$$dX(t) = \left[\frac{\theta_1}{2}(1 - X(t)) - \frac{\theta_2}{2}X(t) + \frac{\beta}{2}X(t)(1 - X(t)) \right] dt + \sqrt{X(t)(1 - X(t))}\, dW_t, \quad t > 0 \quad (1)$$

in which W_t is the Wiener process, also called the standard Brownian motion.

Both of the approaches (random background and ancestral selection graph) we take to modeling latent mutations rely on the assumption that the population has reached equilibrium, which occurs in the limit $t \to \infty$. The stationary probability density of X is

$$\phi_\beta(x) = Cx^{\theta_1 - 1}(1 - x)^{\theta_2 - 1}e^{\beta x}, \quad 0 < x < 1 \quad (2)$$

(Wright, 1931). We explicitly denote the dependence on β because this parameter plays a key role in what follows. The normalizing constant C guarantees that $\int_0^1 \phi_\beta(x)dx = 1$. It is given by

$$C = \frac{\Gamma(\theta_1 + \theta_2)}{\Gamma(\theta_1)\Gamma(\theta_2)\,_1F_1(\theta_1; \theta_1 + \theta_2; \beta)} \quad (3)$$

in which $\Gamma(a)$ is the gamma function and $_1F_1(a; b; z)$ is the confluent hypergeometric function, or Kummer's function; see Abramowitz and Stegun (1964) and Slater (1960).

By definition, latent mutations occur in the ancestry of a sample. When a sample of total size n is taken without replacement from a population with stationary density (2), it will contain a random number $\mathcal{N}_1$ of copies of allele A_1 and $\mathcal{N}_2 = n - \mathcal{N}_1$ copies of allele A_2. The probability that $\mathcal{N}_1$ is equal to n_1 is equal to

$$q(n_1, n_2) := \mathbb{P}(\mathcal{N}_1 = n_1; n, \beta, \theta_1, \theta_2)$$

$$= \int_0^1 \binom{n}{n_1} x^{n_1}(1 - x)^{n - n_1}\phi_\beta(x)dx$$

$$= C\binom{n_1 + n_2}{n_1}\frac{\Gamma(\theta_1 + n_1)\Gamma(\theta_2 + n_2)}{\Gamma(\theta_1 + \theta_2 + n_1 + n_2)}\,_1F_1(\theta_1 + n_1; \theta_1 + \theta_2 + n_1 + n_2; \beta) \quad (4)$$

for $n_1 \in \{0, 1, \ldots, n\}$ and $n_2 = n - n_1$, and with C again given by (3). The notation $q(\cdot)$ is from Slade (2000a,b) and is convenient for the ancestral selection graph.

Suppose now we are given the sample count, that is, we know that among the n uniformly sampled individuals, n_1 of them are of type 1 and the remaining $n_2 = n - n_1$ are of type 2. Then the posterior distribution of the population frequency of A_1 conditional on the sample is

$$\phi_\beta^{(n_1,n_2)}(x) = \frac{\binom{n_1+n_2}{n_1} x^{n_1}(1-x)^{n_2}\phi_\beta(x)}{q(n_1,n_2)} \tag{5}$$

$$= \frac{\Gamma(\theta_1+\theta_2+n_1+n_2)x^{\theta_1+n_1-1}(1-x)^{\theta_2+n_2-1}e^{\beta x}}{C\,\Gamma(\theta_1+n_1)\Gamma(\theta_2+n_2)\,_1F_1(\theta_1+n_1;\theta_1+\theta_2+n_1+n_2;\beta)} \tag{6}$$

with C as in (3). Equation (5) follows from Bayes' theorem with prior distribution ϕ_β.

The sampling probability $q(n_1,n_2)$ in (4) and the resulting posterior distribution $\phi_\beta^{(n_1,n_2)}$ play major roles in the two approaches we take to modeling latent mutations. Specifically, transition probabilities in the conditional ancestral selection graph depend on ratios of sampling probabilities (Slade, 2000b) and the allele frequency in the ancestral process of Barton et al. (2004) has initial distribution $\phi_\beta^{(n_1,n_2)}(x)$ when conditioned on the sample count.

We describe the occurrence of latent mutations in the ancestry of allele A_1 (and sometimes A_2) conditional on its sample count n_1 which is fixed and finite. We say that A_1 is *rare* when n_1 is fixed (and finite) and the sample size n is much larger than n_1. Likely this means that the population frequency x is small, but it is also possible that the sample is just very unlikely. This concept of rarity, which focuses on the counts of the alleles rather than their relative frequencies in the population, is the same as in Wakeley et al. (2023) and was previously considered by Joyce and Tavaré (1995) and Joyce (1995). Because rates of events in conditional ancestral processes depend on the numbers of each allele, we will enforce rarity by considering n_1 fixed and finite, as n_2 grows. Then, $n \sim n_2$ and both are large.

By *strong selection* we mean large $|\beta|$. We model rarity and strong selection together under the assumption that $\beta = \tilde{\beta}n_2$ for some constant $\tilde{\beta} \in \mathbb{R}$. We study latent mutations and the ancestral processes which generate them under three scenarios: (i) $|\beta|$ large with n_2 fixed, (ii) n_2 large with β fixed, and (iii) both $|\beta|$ and n_2 large with $\tilde{\beta} = \beta/n_2$ fixed. In making approximations for large n_2 and/or large $|\beta|$, we make extensive use of asymptotic results for ratios of gamma functions and for the confluent hypergeometric function which are presented in Appendix section Appendix A.

The parameters θ_1 and θ_2 are fixed constants throughout, with $\theta_1, \theta_2 > 0$. For single nucleotide sites, these population-scaled mutation rates have been estimated from pairwise sequence differences for many species and are typically about 0.01 with a range of about 0.0001 to 0.1 (Leffler et al., 2012). Values for humans are smaller but they vary almost as widely among sites in the genome, with a mean of about 0.0008 and a range of about 0.0001 to 0.02 (Seplyarskiy et al., 2021, 2022; Wakeley et al., 2023). In contrast, there is no reason to suppose that the selection parameter $|\beta|$ is small (Eyre-Walker and Keightley, 2007; Chen et al., 2020). Note that our introduction of a constant $\tilde{\beta} = \beta/n_2$ is simply a device to specify the relative importance of rarity as opposed strong selection, not a hypothesis about biology.

The case of a rare neutral allele was considered in Wakeley et al. (2023) where it was shown that the number of latent mutations in the ancestry of the n_1 copies of allele A_1 follows the same distribution as the number of alleles in the Ewens sampling formula (Ewens, 1972) with sample size n_1 and mutation parameter θ_1. Let K_1 be the random number of these latent mutations for allele A_1 in the ancestry of the sample. Then if ξ_j is a Bernoulli random variable with probability

of success

$$\mathbb{P}(\xi_j = 1) = \frac{\theta_1}{\theta_1 + j - 1} \quad , \quad j = 1, 2, \ldots \tag{7}$$

then under neutrality for large sample size and conditional on $\mathcal{N}_1 = n_1$,

$$K_1 \sim \xi_{n_1} + \xi_{n_1-1} + \cdots + \xi_2 + \xi_1 \tag{8}$$

which gives the stated Ewens sampling result (Arratia et al., 1992). Note that, because coalescence is among exchangeable lineages, the full Ewens sampling formula should apply if we were to keep track of the sizes of latent mutations; see Crane (2016) and Tavaré (2021) for recent reviews.

Here we apply the model of coalescence in a random background described by Barton et al. (2004) to prove these results (7) and (8) for rare alleles in large samples and especially to extend the analysis of latent mutations to scenarios which include selection. We investigate both the number of latent mutations and their timing in the ancestry of the sample, and we allow that selection may be strong. We also show how the same scenarios can be treated using the conditional ancestral selection graph (Slade, 2000a), giving consistently the same results.

Briefly, we find that positive selection does not in general lead to (7) and (8), that very strong positive selection (relative to the sample size) leads to neutral gene genealogies with a single ancient latent mutation for the favored allele. This is described in Section 3 for scenario (i) and for the case $\tilde{\beta} \in (1, \infty)$ in scenario (iii). On other hand, when selection is not too strong relative to the sample size, then extreme rarity of A_1 in the sample can effectively override strong positive selection and retrieve (7) and (8). This is described in Section 3 for scenario (ii) and for the case $\tilde{\beta} \in (-\infty, 1)$ in scenario (iii). Figures 1, 2 and 3 illustrate our results in the three scenarios.

2. Sample frequencies and posterior population frequencies

In this section, we present asymptotic results for the sampling probability $q(n_1, n_2)$ in (4) and the posterior distribution $\phi_\beta^{(n_1, n_2)}(x)$ in our three regimes of interest: (i) $|\beta|$ large with n_2 fixed, (ii) n_2 large with β fixed, and (iii) both $|\beta|$ and n_2 large with $\tilde{\beta} = \beta/n_2$ fixed.

2.1. Asymptotics for sampling probabilities

In the case of strong selection and moderate sample size, that is $|\beta|$ large with n_2 fixed, applying (A.4a) and (A.4b) to (4) gives

$$q(n_1, n_2) = \begin{cases} \binom{n}{n_1} \dfrac{\Gamma(\theta_1 + n_1)}{\Gamma(\theta_1)} |\beta|^{-n_1} \left(1 + O\left(|\beta|^{-1}\right)\right) & \text{if } \beta < 0, \tag{9a} \\[2em] \binom{n}{n_1} \dfrac{\Gamma(\theta_2 + n - n_1)}{\Gamma(\theta_2)} \beta^{n_1 - n} \left(1 + O\left(\beta^{-1}\right)\right) & \text{if } \beta > 0. \tag{9b} \end{cases}$$

Here we focus on the leading-order terms but note that the next-order terms are straightforward to obtain using (A.4a) and (A.4b) and additional terms could be computed using (4.1.2) and (4.1.6) in Slater (1960). In (9a), each additional copy of A_1 decreases the sampling probability by a factor of $1/|\beta|$ so the most likely sample is one which contains no copies of A_1. In (9b), each additional copy of A_1 increases the sampling probability by a factor of β so the most likely sample is monomorphic for A_1. However, these results are perfectly symmetric for the two alleles. Switching allelic labels

and swapping $|\beta|$ for β changes (9a) into (9b). That is, allele A_2 experiences the same effects of positive/negative selection in (9a)/(9b) as the focal allele A_1 does in (9b)/(9a).

In the case of large sample size and moderate selection, that is n_2 large with β fixed, applying (A.5) to (4) gives

$$q(n_1, n_2) = C \frac{\Gamma(\theta_1 + n_1)}{n_1!} n_2^{-\theta_1} \left(1 + O\left(n_2^{-1}\right)\right).$$ (10)

This has the same form as the neutral result, equation (22) in Wakeley et al. (2023), only with the additional factor $1/{}_1F_1(\theta_1; \theta_1 + \theta_2; \beta)$ in the constant C. With respect to the count of the focal allele A_1, the distribution is similar to a (degenerate) negative-binomial distribution, like the corresponding result in Theorem 2 of Watterson (1974) for neutral alleles which propagate by a linear birth-death process. The effect of selection is only to uniformly raise or lower the chances of seeing n_1 copies of A_1 in a very large sample. The additional factor $1/{}_1F_1(\theta_1; \theta_1 + \theta_2; \beta)$ is a decreasing function of β, which is equal to 1 when $\beta = 0$ and approaches 0 quickly from there as β increases. Greater selection against (respectively, for) A_1 increases (respectively, decreases) the chance of it being rare but does not affect the shape of the distribution of n_1, at least to leading order in $1/n_2$.

In the case of large sample size and strong selection, that is both $|\beta|$ and n_2 large with $\tilde{\beta} = \beta/n_2$ fixed, applying (A.4a), (A.4b), (A.6a), (A.6b) to (4) gives

$$q(n_1, n_2) = \begin{cases} \dfrac{\Gamma(\theta_1 + n_1)}{n_1! \Gamma(\theta_1)} \left(\dfrac{1}{1 + |\tilde{\beta}|}\right)^{n_1} \left(\dfrac{|\tilde{\beta}|}{1 + |\tilde{\beta}|}\right)^{\theta_1} \left(1 + O\left(n_2^{-1}\right)\right) & \text{if } \tilde{\beta} < 0 & \text{(11a)} \\[3ex] B_1 \dfrac{\Gamma(\theta_1 + n_1)}{n_1!} \left(\dfrac{1}{1 - \tilde{\beta}}\right)^{n_1} \left(1 + O\left(n_2^{-1}\right)\right) & \text{if } 0 < \tilde{\beta} < 1 & \text{(11b)} \\[3ex] B_2 \dfrac{1}{n_1!} \left(\dfrac{\tilde{\beta} - 1}{\tilde{\beta}} n_2\right)^{n_1} \left(1 + O\left(n_2^{-1}\right)\right) & \text{if } \tilde{\beta} > 1 & \text{(11c)} \end{cases}$$

with constants B_1 and B_2 which are unremarkable except in their dependence on n_2:

$$B_1 \propto n_2^{\theta_2 - \theta_1} e^{-\tilde{\beta} n_2}$$

$$B_2 \propto n_2^{\theta_2 - \frac{1}{2}} \left(\tilde{\beta} e\right)^{-n_2}$$

such that $q(n_1, n_2)$ becomes tiny as n_2 grows, especially in the second case. In (11b) and (11c), allele A_1 is favored by selection so it will be unlikely for its sample count to be very small.

To see how (11b) and (11c) compare to (11a), consider how these three sampling probabilities change as n_1 increases:

$$\frac{q(n_1 + 1, n_2)}{q(n_1, n_2)} \approx \begin{cases} \dfrac{\theta_1 + n_1}{(1 + |\tilde{\beta}|)(n_1 + 1)} & \text{if } \tilde{\beta} < 0 & \text{(12a)} \\[3ex] \dfrac{\theta_1 + n_1}{(1 - \tilde{\beta})(n_1 + 1)} & \text{if } 0 < \tilde{\beta} < 1 & \text{(12b)} \\[3ex] \dfrac{(\tilde{\beta} - 1)n_2}{\tilde{\beta}(n_1 + 1)} & \text{if } \tilde{\beta} > 1 & \text{(12c)} \end{cases}$$

where the approximation is for large n_2, i.e. omitting the $O\big(n_2^{-1}\big)$ parts of (11a), (11b) and (11c). The first two differ numerically, because $1 + |\tilde{\beta}| > 1$ in (12a) whereas $1 - \tilde{\beta} < 1$ in (12b), but otherwise depend identically on n_1. Further, they both differ from corresponding neutral result only by these factors $1 + |\tilde{\beta}|$ and $1 - \tilde{\beta}$. Note that the corresponding neutral result can be obtained using (10) because it, again, differs only by a constant factor from neutral case. Equation (12c) is quite different. With $\tilde{\beta} > 1$, each additional copy of A_1 increases the sampling probability a large factor, making this case similar to the case of strong positive selection in (9b). This is as expected. What is surprising is (12b), namely that some cases of strong selection in favor of A_1 can be made to resemble neutrality simply by increasing the sample size.

2.1.1. Comparison to discrete Moran and Wright-Fisher models

We emphasize that our analyses in this work are of the Wright-Fisher diffusion model, given here as the SDE (1) with stationary distribution (2). It is of interest to know how well our results hold for discrete, exact models such as the Moran model and the Wright-Fisher model, especially as $|\beta| \to \infty$ or $n \to \infty$ for finite n_1, in which cases we might expect the diffusion to be a relatively poor description of the dynamics. In this section, we focus on (11a) and show that it can be obtained in a different way from a discrete-time Moran model, without first passing to the diffusion limit, but that this cannot be done in general starting from the discrete-time Wright-Fisher model.

To leading order in $1/n_2$, (11a) is identical to the probability mass function of a negative binomial distribution with parameters $p = |\tilde{\beta}|/(1 + |\tilde{\beta}|)$ and $r = \theta_1$. Here selection *against* A_1 is so strong that it never reaches appreciable frequency in the population. In the limit, or ignoring the $O\big(n_2^{-1}\big)$ part in (11a), this distribution sums to one over all $n_1 \in \{0, 1, 2, \ldots\}$. The corresponding sum for the degenerate distribution in (10) diverges, because under neutrality there is a non-trivial chance that A_1 reaches appreciable frequency in the population.

Consider a discrete-time haploid Moran model with population size N, in which allele A_2 is favored by selection. Specifically, A_1 and A_2 have equal chances of being chosen to reproduce but different chances of being chosen to die: each A_1 has an increased chance $1 + s$ compared to each A_2. Upon reproduction, the offspring of an A_i, $i \in \{1, 2\}$, has type A_i with probability $1 - u_{3-i}$ and the other type A_{3-i} with probability u_{3-i}. If there are currently ℓ copies of A_1 and $N - \ell$ copies of A_2, then in the next time step there will $\ell + 1$ copies of A_1 with probability

$$\frac{N - \ell}{N - \ell + \ell(1 + s)} \frac{\ell}{N}(1 - u_2) + \frac{N - \ell}{N - \ell + \ell(1 + s)} \frac{N - \ell}{N} u_1 \tag{13}$$

and $\ell - 1$ copies of A_1 with probability

$$\frac{\ell(1 + s)}{N - \ell + \ell(1 + s)} \frac{N - \ell}{N}(1 - u_1) + \frac{\ell(1 + s)}{N - \ell + \ell(1 + s)} \frac{\ell}{N} u_2. \tag{14}$$

This converges to the Wright-Fisher diffusion process (1) as $N \to \infty$ if time is measured in units of $N^2/2$ discrete steps with $\ell = Nx$, $u_1 = \theta_1/N$, $u_2 = \theta_2/N$ and $s = -\beta/N$.

As another way of obtaining (11a), we assume that $s \gg u_1, u_2$. In particular, let $Nu_1 \to \theta_1$ and $Nu_2 \to \theta_2$ as $N \to \infty$ just as in the diffusion model, but let s be a constant. Then we may appeal to the analogous model and limit process (iii) of Karlin and McGregor (1964) which had no selection but instead assumed that $u_2 \gg u_1$. Similarly here we expect that allele A_1 will be held in negligible relative frequency in the population and instead be present in a finite number of copies as $N \to \infty$, only here due to strong selection rather than strong mutation.

In this case, it is natural to measure time in generations, or N time steps in this population. The with $dt = 1/N$, we can rewrite (13) and (14) as

$$\left(\ell \frac{N-\ell}{N-\ell+\ell(1+s)}(1-u_2) + \frac{N-\ell}{N-\ell+\ell(1+s)}(N-\ell)u_1 \right) dt,$$

and

$$\left(\ell(1+s)\frac{N-\ell}{N-\ell+\ell(1+s)}(1-u_1) + \frac{\ell(1+s)}{N-\ell+\ell(1+s)}\ell u_2 \right) dt.$$

In the limit $N \to \infty$ this converges to a continuous-time process for A_1 in which in a small time dt

$$j \to \begin{cases} j+1 & \text{with probability } (\ell+\theta_1)dt & \text{(15a)} \\ j-1 & \text{with probability } \ell(1+s)dt & \text{(15b)} \end{cases}$$

in other words to a birth-death process with immigration where the birth rate is $\lambda = 1$, the death rate is $\mu = 1+s$ and the immigration rate is $\kappa = \theta_1$. Then from (52) in Kendall (1949), the distribution of the number of copies of A_1 in the population at stationarity will be negative binomial with parameters $1 - \lambda/\mu = s/(1+s)$ and $\kappa/\lambda = \theta_1$, or

$$p(\ell) = \binom{\ell+\theta_1-1}{\ell}\left(\frac{1}{1+s}\right)^\ell \left(\frac{s}{1+s}\right)^{\theta_1}, \qquad \ell \in \mathbb{Z}_+. \tag{16}$$

In getting to (11a) in the main text, which we note is for $\beta < 0$, we first applied the diffusion limit then let the selection parameter $|\beta|$ be large, specifically proportional to the number n_2 of copies of A_2 in a sample of large size n for a given fixed number n_1 of copies of A_1 in the sample. In the current haploid Moran model with selection against A_1, $|\beta| = Ns$, and the scalar $|\tilde\beta| = |\beta|/n_2 = Ns/n_2$. Define $\alpha := n_2/N$. Then $|\tilde\beta| = s/\alpha$ and we may think of α as (close to) the proportion of the population sampled, because $n_2/N \sim n/N$.

If the number of copies of A_1 in the population is ℓ, then the probability there are n_1 copies in a sample of size n taken without replacement from the total population of size N is given by the hypergeometric distribution

$$p(n_1|\ell;N) = \frac{\binom{\ell}{n_1}\binom{N-\ell}{n-n_1}}{\binom{N}{n}}, \qquad n_1 = 0,1,2,\ldots,\ell. \tag{17}$$

Setting $n - n_1 = n_2 = \alpha N$ and taking $N \to \infty$, (17) converges to the binomial distribution

$$p(n_1|\ell) = \binom{\ell}{n_1}\alpha^{n_1}(1-\alpha)^{\ell-n_1}, \qquad n_1 = 0,1,2,\ldots,\ell. \tag{18}$$

This gives another route to (11a), namely using (16) and (18):

$$
\begin{aligned}
p(n_1) &= \sum_{\ell=n_1}^{\infty} p(n_1|\ell)p(\ell) \\
&= \sum_{\ell=n_1}^{\infty} \binom{\ell}{n_1} \alpha^{n_1}(1-\alpha)^{\ell-n_1} \binom{\ell+\theta_1-1}{\ell} \left(\frac{1}{1+s}\right)^{\ell} \left(\frac{s}{1+s}\right)^{\theta_1} \\
&= \frac{1}{n_1!} \left(\frac{\alpha}{1+s}\right)^{n_1} \left(\frac{s}{\alpha+s}\right)^{\theta_1} \sum_{y=0}^{\infty} \frac{\Gamma(y+n_1+\theta_1)}{\Gamma(y+\theta_1)} \binom{y+\theta_1-1}{y} \left(\frac{1-\alpha}{1+s}\right)^{y} \left(\frac{\alpha+s}{1+s}\right)^{\theta_1} \\
&= \frac{\Gamma(n_1+\theta_1)}{n_1!\Gamma(\theta_1)} \left(\frac{\alpha}{\alpha+s}\right)^{n_1} \left(\frac{s}{\alpha+s}\right)^{\theta_1}
\end{aligned}
\tag{19}
$$

the end result (19) being equal to the leading order part of (11a) since $|\tilde{\beta}| = s/\alpha$.

We can contrast this with a discrete-time haploid Wright-Fisher model with population size N, in which time is already measured in generations. Using equation (33) in Nagylaki (1990) for the corresponding model of (genic) viability selection, the count ℓ of A_1 evolves from one generation (g) to the next with transition probabilities given by

$$
\ell_{g+1}|\ell_g \sim \text{Poisson}(\theta_1 + \ell_g(1-s)),
\tag{20}
$$

that is by a Poisson branching process with Poisson immigration rather than by the birth-death process with immigration in (15a) and (15b). Although here too the number of copies of A_1 in the population will converge to a stationary distribution (Heathcote, 1965), it will not in general be a negative binomial distribution. Therefore, our results which are valid for the diffusion model may not capture important features of the Wright-Fisher process. On the other hand, it is not surprising that they may correspond well with the Moran process; consult for example Etheridge (2011, Definition 2.26 and Remark 2.27).

2.2. Asymptotics for population frequencies conditional on the sample

Next, we obtain asymptotics for the posterior distribution $\phi_\beta^{(n_1,n_2)}$ deduced in (6). Note that

$$
\phi_\beta^{(n_1,n_2)}(x) = \frac{\Gamma(\theta_1+n_1,\theta_2+n_2)\, x^{\theta_1+n_1-1}(1-x)^{\theta_2+n_2-1}e^{\beta x}}{\Gamma(\theta_1+n_1)\Gamma(\theta_2+n_2){}_1F_1\left(\theta_1+n_1;\theta_1+\theta_2+n;\beta\right)},
\tag{21}
$$

since $\phi_\beta^{(n_1,n_2)} \propto x^{\theta_1+n_1-1}(1-x)^{\theta_2+n_2-1}e^{\beta x}$ by (6) and it is a probability distribution.

Let $\mathcal{P}([0,1])$ be the space of probability measures on $[0,1]$ endowed with the weak converegnce topology (i.e with test functions in the space $C_b([0,1])$ of bounded continuous functions).

Lemma 1. *Let $n_1 \in \mathbb{N}$ be fixed. The following convergences in $\mathcal{P}([0,1])$ hold.*

(i) *Suppose n_2 is fixed and $\beta \to \infty$. Then $\phi_\beta^{(n_1,n_2)}(x)\,dx \to \delta_1$.*

(ii) *Suppose β is fixed and $n_2 \to \infty$. Then $\phi_\beta^{(n_1,n_2)}(x)\,dx \to \delta_0$.*

(iii) Suppose $\tilde{\beta} \in \mathbb{R}$ is fixed and $\beta = \tilde{\beta} n_2 \to \infty$. Then

$$\phi_\beta^{(n_1,n_2)}(x)\, dx \to \begin{cases} \delta_0 & \text{when} \quad \tilde{\beta} \in (-\infty, 1] \\ \delta_{1-1/\tilde{\beta}} & \text{when} \quad \tilde{\beta} \in (1, \infty) \end{cases} \tag{22}$$

where δ_x is the Dirac delta function.

Proof of Lemma 1. Part (ii) follows from part (iii) with $\tilde{\beta} = 0$. To prove part (iii), we let $a = \theta_1 + n_1$ for simplicity. The function $\phi_\beta^{(n_1,n_2)}$ is a constant multiple of the function

$$x^{a-1}(1-x)^{n_2+\theta_2-1}e^{\beta x} = \left[(1-x)e^{\tilde{\beta} x}\right]^{n_2} x^{a-1}(1-x)^{\theta_2-1}$$
$$= e^{n_2\, S(x)}\, x^{a-1}(1-x)^{\theta_2-1},$$

where the function $S : [0,1) \to \mathbb{R}$ defined by $S(x) := \tilde{\beta} x + \ln(1-x)$

$$\begin{cases} \text{is strictly decreasing} & \text{when} \quad \tilde{\beta} \in (-\infty, 1] \\ \text{has a global maximum at } x = 1 - 1/\tilde{\beta} \in (0,1) & \text{when} \quad \tilde{\beta} \in (1, \infty) \end{cases} \tag{23}$$

Part (iii) then follows from asymptotic expansion of integrals such as the Laplace method. For example, when $\tilde{\beta} \in (-\infty, 1]$, for any $\delta \in (0,1)$, the ratio

$$\int_\delta^1 e^{n_2\, S(x)}\, x^{a-1}(1-x)^{\theta_2-1}\, dx \Big/ \int_0^1 e^{n_2\, S(x)}\, x^{a-1}(1-x)^{\theta_2-1}\, dx \to 0 \quad \text{as } n_2 \to \infty. \tag{24}$$

For any $f \in C_b([0,1])$ and any $\epsilon > 0$, there exists $\delta \in (0,1)$ such that $\sup_{y:\, 0<y<\delta} |f(y) - f(0)| < \epsilon$ and so

$$\left| \int_0^1 f(x)\phi_\beta^{(n_1,n_2)}(x)\, dx - f(0) \right|$$
$$\leq \left| \int_\delta^1 f(x)\phi_\beta^{(n_1,n_2)}(x)\, dx \right| + \left| \int_0^\delta f(x)\phi_\beta^{(n_1,n_2)}(x)\, dx - f(0) \right|$$
$$\leq \|f\| \int_\delta^1 \phi_\beta^{(n_1,n_2)}(x)\, dx + \epsilon + |f(0)| \int_\delta^1 \phi_\beta^{(n_1,n_2)}(x)\, dx.$$

Hence by (24), $\limsup_{n_2\to\infty} \left| \int_0^1 f(x)\phi_\beta^{(n_1,n_2)}(x)\, dx - f(0) \right| \leq \epsilon$. Since $\epsilon > 0$ is arbitrary, we have shown that $\left| \int_0^1 f(x)\phi_\beta^{(n_1,n_2)}(x)\, dx - f(0) \right| \to 0$ as $n_2 \to \infty$.

Part (i) follows from the same argument. □

3. Conditional coalescence in a random background

In this section, we extend the approach of coalescence in a random background in Barton et al. (2004) to study the number and timing of latent mutations and other asymptotic properties of the conditional gene genealogy given the sample frequencies of A_1 and A_2. We also extend our results

to time-varying populations in Section 3.4. While the setting of Barton et al. (2004) covers the case of two linked loci (one selected, the other neutral) here we focus on one selected locus.

Suppose we are given a sample from the selected locus at the present time $t = 0$, and that we know the allelic types of the sample but we do not know how the sample was produced. What is the genealogy of the sample? This question was answered by Barton et al. (2004), where they modeled the ancestral process using the structured coalescent with allelic types as subpopulations. The structured coalescent can be a model of subdivision with migration between local populations (Takahata, 1988; Notohara, 1990; Herbots, 1997) or a model of selection with mutation between allelic types (Kaplan et al., 1988; Darden et al., 1989). For samples from a population at stationarity as in Section 1, Barton et al. (2004) proved that this could be done rigorously starting with a Moran model with finite N then passing to the diffusion limit. Barton and Etheridge (2004) explored some properties of gene genealogies under this model, and Etheridge et al. (2006) used the same idea to describe genetic ancestries following a selective sweep.

Even if the sample frequencies are known, the allele frequencies in the population are unknown. A key feature of this method is to model allele-frequency trajectories backward in time. As pointed out by Barton et al. (2004), the Moran model with finite N is *reversible*, meaning that under its stationary distribution, the time-reversed process is the same (in distribution) as the forward-time Moran process. This is not a property of the Wright-Fisher model with finite N but does hold for their shared diffusion limit (1) with stationary distribution (2); see for instance Millet et al. (1989) for why this holds. Figure 1 gives an illustration of a genealogy with mutations and allele frequencies varying backward in time.

Let $n_i^{(N)}(t)$ be the number of ancestral lineages of type i at time t backward, and let $p_t^{(N)}$ be the fraction of type 1 in the population at time t backward. By (Barton et al., 2004, Lemma 2.4), under the Moran model with stationary distribution, $(p_t^{(N)}, n_1^{(N)}(t), n_2^{(N)}(t))_{t \in \mathbb{R}_+}$ is a Markov process for each fixed N. Furthermore, (Barton et al., 2004, Lemma 3.1) describes the joint convergence of the processes $(p^{(N)}, n_1^{(N)}, n_2^{(N)})$ as $N \to \infty$.

Lemma 2 (Lemmas 2.4 and 3.1 of Barton et al. (2004)). *Let $p_t^{(N)}$ and $n_i^{(N)}(t)$ be the fraction of type 1 in the population and the number of ancestral lineages of type i respectively, at time t backward, under the stationary Moran model. Then $(p^{(N)}, n_1^{(N)}, n_2^{(N)})$ is a Markov process for each $N \in \mathbb{N}$. As $N \to \infty$, this process converges in distribution to a Markov process $(p_t, n_1(t), n_2(t))_{t \in \mathbb{R}_+}$ with state space $[0,1] \times \mathbb{Z}_+ \times \mathbb{Z}_+$. This limiting process is described as follows:*

(i) $(p_t)_{t \in \mathbb{R}_+}$ is a solution to equation (1) with stationary initial distribution ϕ_β. In particular, it does not depend on $(n_1(t), n_2(t))_{t \in \mathbb{R}_+}$.

(ii) Suppose the current state is (p, m_1, m_2). Then $(n_1(t), n_2(t))_{t \in \mathbb{R}_+}$ evolves as

$$
(m_1, m_2) \to \begin{cases} (m_1 - 1, m_2) & \text{at rate} \quad \frac{1}{p}\binom{m_1}{2} & \text{coalescence of type 1} \\ (m_1 - 1, m_2 + 1) & \text{at rate} \quad \frac{1-p}{p} m_1 \frac{\theta_1}{2} & \text{mutation of 1 to 2} \\ (m_1, m_2 - 1) & \text{at rate} \quad \frac{1}{1-p}\binom{m_2}{2} & \text{coalescence of type 2} \\ (m_1 + 1, m_2 - 1) & \text{at rate} \quad \frac{p}{1-p} m_2 \frac{\theta_2}{2} & \text{mutation of 2 to 1} \end{cases} \tag{25}
$$

We compare the notation here with that of Barton et al. (2004) and Etheridge (2011). Type 1 here is their type P, type 2 here is their type Q. So, $\frac{\theta_2}{2}$ (resp. $\frac{\theta_1}{2}$) here is μ_1 (resp. μ_2) in Barton

et al. (2004), and is v_1 (resp. v_2) in (Etheridge, 2011, eqn. (2.11)). The rescaled selection coefficient $\frac{\beta}{2}$ here is s in Barton et al. (2004). The Moran model in Barton et al. (2004) has population size $2N$ and each of the $(2N)(2N-1)/2$ unordered pairs is picked to interact (one dies and immediately the other reproduces) at rate $1/2$. Therefore, the diffusion limit in (Barton et al., 2004, Lemma 3.1) is slower by a factor of $1/2$ than the limit we consider here in this paper.

The proof of (Barton et al., 2004, Lemma 3.1) leads to more information for the limiting process. Let $n_i^{(N),obs}(t)$ be the number of type i lineages, among the $n_i^{(N)}(0)$ lineages that were present (thus observed) at $t = 0$, that remain at time t backward. Thus $n_i^{(N),obs}(t)$ is non-increasing in t, but $n_i^{(N)}(t)$ can increase as t increases due to mutations from type $3 - i$ to type i. Clearly, $n_i^{(N),obs}(0) = n_i^{(N)}(0)$. To keep track of the number of mutation events versus the number of coalescent events, we let $L_i^{(N)}(t)$ be the total number of latent mutations for type i during $[0,t]$ backward in time. We have the following generalization of Lemma 2.

Lemma 3 (Joint convergence). *Under the stationary Moran model, the backward process*

$$(p^{(N)};\, n_1^{(N)}, n_1^{(N),obs}, L_1^{(N)};\, n_2^{(N)}, n_2^{(N),obs}, L_2^{(N)})$$

is a Markov process for each fixed $N \in \mathbb{N}$. As $N \to \infty$, this process converges in distribution to a continuous-time Markov process $(p_t;\, n_1(t), n_1^{obs}(t), L_1(t);\, n_2(t), n_2^{obs}(t), L_2(t))_{t \in \mathbb{R}_+}$ with state space $[0,1] \times \mathbb{Z}_+^6$ such that

(i) *$(p_t)_{t \in \mathbb{R}_+}$ is a solution to equation (1) with stationary initial distribution ϕ_β. In particular, it does not depend on $(n_1(t), n_1^{obs}(t), L_1(t);\, n_2(t), n_2^{obs}(t), L_2(t))_{t \in \mathbb{R}_+}$.*

(ii) *At state $(p;\, m_1, a_1, \ell_1;\, m_2, a_2, \ell_2)$, the process $(n_1(t), n_1^{obs}(t), L_1(t);\, n_2(t), n_2^{obs}(t), L_2(t))_{t \in \mathbb{R}_+}$ evolves as $(m_1, a_1, \ell_1;\, m_2, a_2, \ell_2) \to$*

$$
\begin{cases}
(m_1 - 1, a_1 - 1, \ell_1;\, m_2, a_2, \ell_2) & \text{at rate} \quad \frac{1}{p}\binom{a_1}{2} & \text{coalescence of two type } 1^{obs} \\[4pt]
(m_1 - 1, a_1, \ell_1;\, m_2, a_2, \ell_2) & \text{at rate} \quad \frac{1}{p}\left[\binom{m_1}{2} - \binom{a_1}{2}\right] & \text{other coalescence of type 1} \\[4pt]
(m_1 - 1, a_1 - 1, \ell_1 + 1;\, m_2 + 1, a_2, \ell_2) & \text{at rate} \quad \frac{1-p}{p}\, a_1 \frac{\theta_1}{2} & \text{mutation of } 1^{obs} \\[4pt]
(m_1 - 1, a_1, \ell_1;\, m_2 + 1, a_2, \ell_2) & \text{at rate} \quad \frac{1-p}{p}\,(m_1 - a_1)\frac{\theta_1}{2} & \text{mutation of other type 1}
\end{cases}
$$

and, similarly,

$$
\begin{cases}
(m_1, a_1, \ell_1;\, m_2 - 1, a_2 - 1, \ell_2) & \text{at rate} \quad \frac{1}{1-p}\binom{a_2}{2} & \text{coalescence of two type } 2^{obs} \\[4pt]
(m_1, a_1, \ell_1;\, m_2 - 1, a_2, \ell_2) & \text{at rate} \quad \frac{1}{1-p}\left[\binom{m_2}{2} - \binom{a_2}{2}\right] & \text{other coalescence of type 2} \\[4pt]
(m_1 + 1, a_1, \ell_1;\, m_2 - 1, a_2 - 1, \ell_2 + 1) & \text{at rate} \quad \frac{p}{1-p}\, a_2 \frac{\theta_2}{2} & \text{mutation of } 2^{obs} \\[4pt]
(m_1 + 1, a_1, \ell_1;\, m_2 - 1, a_2, \ell_2) & \text{at rate} \quad \frac{p}{1-p}\,(m_2 - a_2)\frac{\theta_2}{2} & \text{mutation of other type 2}
\end{cases}
$$

Now suppose that in addition to knowing the sample counts n_1 and n_2, we also know that these are the outcome of uniform random sampling, as in (4). Let $\mathbb{P}_{\mathbf{n}}$ be the conditional probability measure given that a uniformly picked sample has allelic counts $\mathbf{n} = (n_1, n_2)$. Under $\mathbb{P}_{\mathbf{n}}$,

the limiting process in Lemma 3 has initial distribution $p_0 \sim \phi_\beta^{(n_1,n_2)}(x)\,dx$ given by (21), and $(n_1(0), n_1^{obs}(0), L_1(0); n_2(0), n_2^{obs}(0), L_2(0)) = (n_1, n_1, 0; n_2, n_2, 0)$. This follows from Bayes' theorem, because by part (i) of Lemmas 2 and 3, the initial frequency has prior distribution given by (2).

Focus on type 1 for now. We care about the sequence of events (coalescence and mutation) backward in time for type 1, and the timing of these events. At each of these events, the number of type 1 lineages decreases by 1, either by coalescence or by mutation from type 1 to type 2. Hence $n_1^{obs}(t)$ is non-increasing, but $n_1(t)$ can increase over time (backward) due to mutations from type 2 to type 1 (see Figure 1 for an illustration). Furthermore, the difference $n_1(t) - n_1^{obs}(t)$ is the number of type 1 at time t that came from lineages that are of type 2 in the sample (at $t = 0$). We do not care about these $n_1(t) - n_1^{obs}(t)$ lineages, nor the mutation events from type 2 to type 1. Analogous considerations hold for type 2.

From Lemma 3, we immediately obtain the following simplified description for the conditional ancestral process in the limit $N \to \infty$ for the two types. This description is the starting point of our analysis for constant population size; later in Proposition 2 we also obtain the analogous result for time-varying population size.

Proposition 1 (Conditional ancestral process). *The process $(p_t, n_1^{obs}(t), L_1(t), n_2^{obs}(t), L_2(t))_{t \in \mathbb{R}_+}$ under $\mathbb{P}_\mathbf{n}$ is a Markov process with state space $[0,1] \times \{0, 1, \cdots, n_1\}^2 \times \{0, 1, \cdots, n_2\}^2$ described as follows:*

(i) $(p_t)_{t \in \mathbb{R}_+}$ is a solution to (1) with initial distribution $\phi_\beta^{(n_1,n_2)}$. In particular, it does not depend on the process $(n_1^{obs}, L_1, n_2^{obs}, L_2)$.

(ii) The process $(n_1^{obs}, L_1, n_2^{obs}, L_2)$ starts at $(n_1, 0, n_2, 0)$. When the current state is $(p, a_1, \ell_1, a_2, \ell_2)$, this process evolves as

$$(a_1, \ell_1) \to \begin{cases} (a_1 - 1, \ell_1) & \text{at rate} \quad \frac{1}{p}\binom{a_1}{2} & \text{coalescence of type } 1^{obs} \\ (a_1 - 1, \ell_1 + 1) & \text{at rate} \quad \frac{1-p}{p} a_1 \frac{\theta_1}{2} & \text{mutation of } 1^{obs} \text{ to } 2 \end{cases} \tag{26}$$

and, independently,

$$(a_2, \ell_2) \to \begin{cases} (a_2 - 1, \ell_2) & \text{at rate} \quad \frac{1}{1-p}\binom{a_2}{2} & \text{coalescence of type } 2^{obs} \\ (a_2 - 1, \ell_2 + 1) & \text{at rate} \quad \frac{p}{1-p} a_2 \frac{\theta_2}{2} & \text{mutation of } 2^{obs} \text{ to } 1 \end{cases} \tag{27}$$

The total rate in (26), at which a_1 decreases by 1, is

$$\frac{a_1}{2p}\big((1-p)\theta_1 + a_1 - 1\big) := \lambda_{a_1}(p), \tag{28}$$

and the one-step transition probabilities are

$$(a_1, \ell_1) \to \begin{cases} (a_1 - 1, \ell_1) & \text{with probability} \quad \frac{a_1 - 1}{(1-p)\theta_1 + a_1 - 1} \\ (a_1 - 1, \ell_1 + 1) & \text{with probability} \quad \frac{(1-p)\theta_1}{(1-p)\theta_1 + a_1 - 1} =: h_{a_1}(p) \end{cases} \tag{29}$$

As t increases from 0 to ∞, the process $(n_1^{obs}(t))_{t\in\mathbb{R}_+}$ decreases from n_1 to 0, and the process $(L_1(t))_{t\in\mathbb{R}_+}$ increases from 0 to a random number $K_1 := \lim_{t\to\infty} L_1(t) \in \mathbb{N}$ which is the total number of latent mutations for type 1. Similarly, the total number of latent mutations for type 2 is defined by $K_2 := \lim_{t\to\infty} L_2(t) \in \mathbb{N}$.

We now give a more explicit description of (K_1, K_2) using the frequency process p and independent Bernoulli random variables. Note these are conditional on the sample counts, $(\mathcal{N}_1 = n_1, \mathcal{N}_2 = n_2)$ as in (8).

Let $\tau_1 < \tau_2 < \cdots < \tau_{n_1}$ be the jump times of the process n_1^{obs}. At time τ_1, the process n_1^{obs} decreases from n_1 to $n_1 - 1$, etc., until finally at τ_{n_1}, n_1^{obs} decreases from 1 to 0. It can be checked that $\tau_{n_1} < \infty$ almost surely under $\mathbb{P}_{\mathbf{n}}$, using the ergodicity of the process p and (28), so that $(n_1^{obs}(t))_{t\in\mathbb{R}_+}$ will indeed decrease to 0 eventually under $\mathbb{P}_{\mathbf{n}}$ almost surely. The allele frequencies at these random times are $p_{\tau_1}, p_{\tau_2}, \ldots, p_{\tau_{n_1}}$. By Proposition 1, under $\mathbb{P}_{\mathbf{n}}$, we have

$$K_1 \overset{d}{=} \xi_{n_1}(p_{\tau_1}) + \xi_{n_1-1}(p_{\tau_2}) + \xi_{n_1-2}(p_{\tau_3}) + \cdots + \xi_2(p_{\tau_{n_1-1}}) + 1, \tag{30}$$

where $\{\xi_k(\cdot)\}_{k=2}^{n_1}$ is a family of independent random processes such that, for a constant $p \in [0,1]$,

$$\xi_k(p) = \begin{cases} 0 & \text{with probability} \quad \frac{k-1}{(1-p)\theta_1+k-1} \\ 1 & \text{with probability} \quad \frac{(1-p)\theta_1}{(1-p)\theta_1+k-1} \end{cases} \tag{31}$$

Similarly, if we let $s_1 < s_2 < \cdots < s_{n_2}$ be the jump times of the process n_2^{obs}, then

$$K_2 \overset{d}{=} \zeta_{n_2}(p_{s_1}) + \zeta_{n_2-1}(p_{s_2}) + \zeta_{n_2-2}(p_{s_3}) + \cdots + \zeta_2(p_{s_{n_2-1}}) + 1, \tag{32}$$

where $\{\zeta_k(\cdot)\}_{k=2}^{n_2}$ is a family of independent random processes such that, for a constant $p \in [0,1]$,

$$\zeta_k(p) = \begin{cases} 0 & \text{with probability} \quad \frac{k-1}{p\theta_2+k-1} \\ 1 & \text{with probability} \quad \frac{p\theta_2}{p\theta_2+k-1} \end{cases} \tag{33}$$

Having described the ancestral process in Proposition 1 and the latent mutations in (30) and (32) under the conditional probability $\mathbb{P}_{\mathbf{n}}$, we study their asymptotic properties under 3 scenarios in the next 3 subsections. These 3 scenarios come from the asymptotic behaviors of the initial frequency p_0 described in Lemma 4.

Lemma 4 (Asymptotic initial frequency). *Let $n_1 \in \mathbb{N}$ be fixed. The initial frequency p_0 converges in probability under $\mathbb{P}_{\mathbf{n}}$ to a deterministic constant as follows.*

(i) Suppose n_2 is fixed and $\beta \to \infty$. Then $p_0 \to 1$.

(ii) Suppose β is fixed and $n_2 \to \infty$. Then $p_0 \to 0$.

(iii) Suppose $\tilde{\beta} \in \mathbb{R}$ is fixed and $\beta = \tilde{\beta} n_2 \to \infty$. Then

$$p_0 \to \begin{cases} 0 & \text{when} \quad \tilde{\beta} \in (-\infty, 1] \\ 1 - 1/\tilde{\beta} & \text{when} \quad \tilde{\beta} \in (1, \infty) \end{cases} \tag{34}$$

Furthermore, when $\tilde{\beta} \in (-\infty, 1)$, it holds that $n_2 p_0$ converges in distribution to a Gamma random variable $\mathrm{Gam}(n_1 + \theta_1, 1 - \tilde{\beta})$ with probability density function $\frac{(1-\tilde{\beta})^{(n_1+\theta_1)}}{\Gamma(n_1+\theta_1)} y^{n_1+\theta_1-1} e^{(\tilde{\beta}-1)y}$.

Proof of Lemma 4. Convergence in distribution to a constant is equivalent to convergence in probability. Hence this result, except the last statement, follows from Lemma 1.

When $\tilde{\beta} \in (-\infty, 1)$, we let $a = n_1 + \theta_1$ for simplicity. The probability density function of np_0 under $\mathbb{P}_{\mathbf{n}}$ is

$$
\begin{aligned}
\frac{1}{n} \phi_\beta^{(n_1,n_2)} \left(\frac{y}{n}\right) &= \frac{1}{n} \frac{1}{\mathrm{Beta}(a, n_2+\theta_2)_1 F_1\left(a; n+\theta_1+\theta_2; \beta\right)} \left(\frac{y}{n}\right)^{a-1} \left(1 - \left(\frac{y}{n}\right)\right)^{n_2+\theta_2-1} e^{\beta \frac{y}{n}} \\
&\approx \frac{1}{n} \frac{1}{\mathrm{Beta}(a, n_2+\theta_2)_1 F_1\left(a; n+\theta_1+\theta_2; \tilde{\beta} n_2\right)} \left(\frac{y}{n}\right)^{a-1} e^{-y} e^{\tilde{\beta} y} \\
&\approx \frac{1}{n^a} \frac{1}{\mathrm{Beta}(a, n_2+\theta_2) (1-\tilde{\beta})^{-a}} y^{a-1} e^{-y} e^{\tilde{\beta} y} \\
&\approx \frac{1}{\Gamma(a) (1-\tilde{\beta})^{-a}} y^{a-1} e^{-y} e^{\tilde{\beta} y},
\end{aligned}
$$

where we used (A.6a) and then (A.1) in the last two approximations above, as n and $\beta \to \infty$. Hence $n_2 p_0$ converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to the $\mathrm{Gam}(n_1 + \theta_1, 1 - \tilde{\beta})$ random variable. $\qquad\square$

3.1. Scenario (i): strong selection, arbitrary sample size

The first scenario is when $|\beta|$ large with n_2 fixed. We consider the case $\beta \to +\infty$ only, since the other case $\beta \to -\infty$ follows by switching the roles of type 1 and type 2.

The conditional genealogy of the $n_1 + n_2$ sampled individuals, under $\mathbb{P}_{\mathbf{n}}$, has three parts with different timescales. First, the n_2 type-2 individuals quickly evolve (coalesce and mutate) as in the Ewen sampling distribution, producing K_2 many type 1 alleles at a short time s_{n_2}. Thus, $K_2 \overset{d}{=} \sum_{k=1}^{n_1} \eta_k$ where $\{\eta_k\}$ are independent Bernoulli variables taking values in $\{0, 1\}$ and having means $\frac{\theta_2}{\theta_2+k-1}$. Next, the resulting $n_1 + K_2$ type 1 lineages will coalesce according to the Kingman coalescent without mutation until only one lineage remains. Finally, it takes a long time, $\tau_{n_1} \approx \frac{2\beta}{\theta_1\theta_2}$, for the single lineage to mutate. In particular, $K_1 \approx 1$.

This description is justified by Theorems 1 and 2. See Figure 1 for an illustration.

Figure 1: Conditional genealogy of a sample with observed frequencies $(n_1, n_2) = (3, 7)$ at the present time $t = 0$, when β is large. The fluctuating blue curve is the process $(p_t)_{t \in \mathbb{R}_+}$ of population frequency of type 1 backward in time. In this example, the 7 type-2 alleles quickly evolve (coalesce and mutate), producing $K_2 = 2$ type-1 alleles according to a Ewen sampling distribution. The $5 = 2 + 3$ type-1 alleles coalese according to the Kingman coalescent (with no mutation) until there is only one lineage, at τ_{n_1-1}. This lineage mutates after a long time, at $\tau_{n_1} \approx \frac{2\beta}{\theta_1 \theta_2}$. While the process $n_1(t)$ increases from 3 to 4, then to 5 before decreasing to 0, the process $n_1^{obs}(t)$ is monotonically decreasing.

Theorem 1. *Suppose $(n_1, n_2) \in \mathbb{N}^2$ is fixed. Then under $\mathbb{P_n}$, as $\beta \to \infty$,*

(i) $\sup_{t \in [0,T]} |1 - p_t| \to 0$ *in probability, for any $T \in (0, \infty)$; and*

(ii) the triplet (K_1, K_2, s_{n_2}) converges in distribution to $\left(1, \sum_{k=1}^{n_1} \eta_k, 0\right)$, where $\{\eta_k\}$ are independent Bernoulli variables taking values in $\{0, 1\}$ and having means $\frac{\theta_2}{\theta_2 + k - 1}$.

Proof. By (1), the process $q := 1 - p$ solves the equation

$$dq_t = \sqrt{q_t(1 - q_t)}\, dW_t - \frac{\theta_1}{2} q_t + \frac{\theta_2}{2}(1 - q_t) - \frac{\beta}{2} q_t(1 - q_t), \quad t \geq 0. \tag{35}$$

Fix any $\epsilon \in (0, 1)$. We shall to show that $\mathbb{P_n}(\sup_{t \in [0,T]} q_t > 2\epsilon) \to 0$ as $\beta \to \infty$.

By the comparison principle (Karatzas and Shreve, 1991, Proposition 2.18 in Chap. 5), we can replace the process q by another process $\widehat{q}$ that solves

$$d\widehat{q}_t = \sqrt{\widehat{q}_t(1 - \widehat{q}_t)}\, dW_t + \left[\frac{\theta_2}{2} - \frac{\beta}{2} \widehat{q}_t(1 - \widehat{q}_t) \right] dt \tag{36}$$

with an initial condition $\widehat{q}_0$ that has the same distribution as q_0. By Girsanov's theorem, we can further take away the constant drift $\frac{\theta_2}{2} dt$. That is, it suffices to show that there exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ on which $\mathbb{P}(\sup_{t \in [0,T]} \widehat{Q}_t > 2\epsilon) \to 0$ as $\beta \to \infty$, where the process $\widehat{Q}$ solves

$$d\widehat{Q}_t = \sqrt{\widehat{Q}_t(1 - \widehat{Q}_t)}\, dW_t - \frac{\beta}{2} \widehat{Q}_t(1 - \widehat{Q}_t)\, dt \tag{37}$$

with an initial condition $\widehat{Q}_0$ that has the same distribution as q_0. The initial frequency $q_0 \to 0$ in probability under $\mathbb{P_n}$, by Lemma 1(i). Hence it suffices to show that

$$\mathbb{P}\left(\sup_{t \in [0,T]} \left\{ \int_0^t \sqrt{\widehat{Q}_s(1 - \widehat{Q}_s)}\, dW_s - \frac{\beta}{2} \int_0^t \widehat{Q}_s(1 - \widehat{Q}_s)\, ds \right\} > \epsilon \right) \to 0 \quad \text{as } \beta \to 0. \tag{38}$$

This is true by the time-change representation of the martingale $M_t := \int_0^t \sqrt{\widehat{Q}_s(1 - \widehat{Q}_s)} \, dW_s$ (Karatzas and Shreve, 1991, Theorem 4.6 in Chap. 3) and the fact that

$$
\mathbb{P}\left(\sup_{t \in [0,T]} \left\{ B_{\langle M \rangle_t} - \frac{\beta}{2} \langle M \rangle_t \right\} > \epsilon \right)
$$

$$
= \mathbb{P}\left(\sup_{r \in [0, \langle M \rangle_T]} \left\{ B_r - \frac{\beta}{2} r \right\} > \epsilon \right)
$$

$$
\leq \mathbb{P}\left(\sup_{r \in [0, T/4]} \left\{ B_r - \frac{\beta}{2} r \right\} > \epsilon \right)
$$

$$
= \int_0^{T/4} \frac{1}{\sqrt{2\pi t^3}} \exp\left\{ \frac{-(\epsilon + \frac{\beta}{2} t)^2}{2t} \right\} dt \to 0 \quad \text{as} \quad \beta \to \infty,
$$

where in the inequality we used the fact that $\langle M \rangle_T \leq T/4$ almost surely (since $q_t(1 - q_t) \leq 1/4$ for all $t \in \mathbb{R}_+$). Convergence (i) is proved.

Note that the coalescence rate and the mutation rate in (26) converge to $a_1(a_1 - 1)/2$ and 0 respectively as $p \to 1$, and that in (27), both rates converge to infinity but their ratio converges in such a way that the limiting one-step transition probabilities are

$$
(a_2, \ell_2) \to \begin{cases} (a_2 - 1, \ell_2) & \text{with probability} \quad \frac{a_2 - 1}{\theta_2 + a_2 - 1} \\ (a_2 - 1, \ell_2 + 1) & \text{with probability} \quad \frac{\theta_2}{\theta_2 + a_2 - 1} \end{cases} \tag{39}
$$

Convergence (ii) then follows from (i) and the representations (30) and (32). $\qquad\square$

Next, we quantify the difference $q_t := 1 - p_t$ by rescaling. Let $(Q_t)_{t \in \mathbb{R}_+}$ be the $\mathbb{R}_+$-valued process that solves the stochastic differential equation

$$
dQ_t = \sqrt{Q_t} \, dW_t + \frac{1}{2}(\theta_2 - Q_t) \, dt, \qquad t \in \mathbb{R}_+, \tag{40}
$$

where W is the Wiener process.

Equation (40) (also (45) and (49) below) is equivalent to the Cox-Ingersoll-Ross (CIR) model for interest rates in financial mathematics. It has several other names including the Feller process and the square-root process (Dufresne, 2001). It has a unique strong solution. This equation is not explicitly solvable, but its transition density is explicitly known (Vanyolos et al., 2014) and its moments and distributions have been intensively studied.

Lemma 5. *Let $q_t = 1 - p_t$ for $t \in \mathbb{R}_+$. The process $(\beta q_{\frac{t}{\beta}})_{t \in \mathbb{R}_+}$ converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to the process Q solving* (40) *and starting at 0.*

Proof. Let $Y_t := \beta q_{\frac{t}{\beta}}$. Similar to (47),

$$
\begin{aligned}
Y_t - Y_0 &= \beta \left(q_{t/\beta} - p_0 \right) \\
&= \beta \int_0^{t/\beta} \sqrt{q_s(1-q_s)}\, dW_s + \frac{\beta}{2} \int_0^{t/\beta} \theta_2(1-q_s) - \theta_1 q_s - \beta p_s(1-p_s)\, ds \\
&\overset{d}{=} \frac{\beta}{\sqrt{\beta}} \int_0^t \sqrt{q_{r/\beta}(1-q_{r/\beta})}\, dW_r + \frac{\beta}{2} \int_0^t \theta_2(1-q_{r/\beta}) - \theta_1 q_{r/\beta} - \beta q_{r/\beta}(1-q_{r/\beta})\, \frac{dr}{\beta} \\
&= \sqrt{\beta} \int_0^t \sqrt{\frac{Y_r}{\beta}\left(1-\frac{Y_r}{\beta}\right)}\, dW_r + \frac{1}{2} \int_0^t \theta_2\left(1-\frac{Y_r}{\beta}\right) - \theta_1 \frac{Y_r}{\beta} - \beta \frac{Y_r}{\beta}\left(1-\frac{Y_r}{\beta}\right)\, dr \\
&= \int_0^t \sqrt{Y_r\left(1-\frac{Y_r}{\beta}\right)}\, dW_r + \frac{1}{2} \int_0^t \theta_2\left(1-\frac{Y_r}{\beta}\right) - \theta_1 \frac{Y_r}{\beta} - Y_r\left(1-\frac{Y_r}{\beta}\right)\, dr. \quad (41)
\end{aligned}
$$

By Lemma 1(i), $Y_0 \to 0$ in probability. The desired weak convergence in the canonical path space $C([0,\infty), \mathbb{R})$ can be proved by a standard compactness argument (Billingsley, 2013, Chap. 2) or (Ethier and Kurtz, 2009, Chap. 3). $\qquad\square$

In Theorem 2 below, we obtain that the mean of the age τ_{n_1} is about $\frac{2\beta}{\theta_1\theta_2}$.

Theorem 2 (Age of the oldest latent mutation of a favorable allele). *Suppose* $(n_1, n_2) \in \mathbb{N}^2$ *is fixed. Then under* $\mathbb{P}_{\mathbf{n}}$, *as* $\beta \to \infty$, $\frac{\tau_{n_1}}{\beta}$ *converges in distribution to an exponential random variable with mean* $\frac{2}{\theta_1\theta_2}$. *That is,*

$$
\frac{\tau_{n_1}}{\beta} \overset{d}{\longrightarrow} \mathrm{Exp}\left(\frac{\theta_1\theta_2}{2}\right) \qquad as \quad \beta \to \infty.
$$

Proof. Recall the rate of mutation in (26) with $a_1 = 1$. For any $t \in \mathbb{R}_+$, as $\beta \to \infty$,

$$
\mathbb{P}_{\mathbf{n}}\left(\frac{\tau_{n_1}}{\beta} > t\right) \approx \mathbb{E}\left[e^{\frac{-\theta_1}{2} \int_0^{\beta t} \frac{1-p_s}{p_s}\, ds} \right] \tag{42}
$$

$$
= \mathbb{E}\left[e^{\frac{-\theta_1}{2} \int_0^{\beta^2 t} \frac{q_{r/\beta}}{1-q_{r/\beta}}\, \frac{dr}{\beta}} \right]
$$

$$
= \mathbb{E}\left[e^{\frac{-\theta_1 t}{2\beta^2 t} \int_0^{\beta^2 t} \frac{\beta q_{r/\beta}}{1-q_{r/\beta}}\, dr} \right].
$$

Note that in the exponent, by Lemma 5,

$$
\lim_{\beta \to \infty} \frac{-\theta_1 t}{2\beta^2 t} \int_0^{\beta^2 t} \frac{\beta q_{r/\beta}}{1-q_{r/\beta}}\, dr \overset{d}{=} \frac{-\theta_1 t}{2} \lim_{T \to \infty} \frac{1}{T} \int_0^T Q(r)\, dr = \frac{-\theta_1 t}{2} \mathbb{E}_\infty[Q],
$$

where the last equality holds almost surely because the process Q is ergodic. Here $\mathbb{E}_\infty$ is the expectation with respect to the stationary distribution of the process Q to which it converges to as $t \to \infty$.

The limiting distribution of Q_t, as $t \to \infty$, is known to be the Gamma distribution $\mathrm{Gam}(\theta_2, 1)$ with probability density function $\frac{1}{\Gamma(\theta_2)} y^{\theta_2-1} e^{-y}$. Hence $\mathbb{E}_\infty[Q] = \theta_2$. $\qquad\square$

3.2. Scenario (ii): arbitrary selection, large sample size

The second scenario is when n_2 large with β fixed. This scenario is a special case of the next scenario (when $\tilde{\beta} = 0$).

The conditional genealogy of the n_1 sample type 1 individuals can be described as follows. The times of type 1 events are small under $\mathbb{P}_{\mathbf{n}}$ in the sense that τ_{n_1} is of order $O(1/n)$. However, if we measure time in proportion to $1/n$ coalescent time units and measure frequency on the scale of numbers of copies of alleles, then the n_1 type-1 individuals evolve (coalesce and mutate) as in the Ewen sampling distribution. In particular, $K_1 \approx 1 + \sum_{k=2}^{n_1} \eta_k$, where $\{\eta_k\}$ are independent Bernoulli variables taking values in $\{0, 1\}$ and having means $\frac{\theta_1}{\theta_1 + k - 1}$.

This description is justified by Theorems 3-4 and Corollary 1 below, with $\tilde{\beta} = 0$. In particular, the rescaled genealogy for type 1 can be described precisely under the rescaling above by the Feller diffusion with drift:

$$dZ_t = \sqrt{Z_t}\, dW_t + \frac{\theta_1}{2}\, dt, \qquad t \in \mathbb{R}_+, \tag{43}$$

with initial distribiution being the Gamma random variable $\mathrm{Gam}(n_1 + \theta_1, 1)$. See Figure 2 for an illustration.

3.3. Scenario (iii): strong selection, large sample size

The third scenario is when both $|\beta|$ and n_2 large with $\tilde{\beta} = \beta/n_2$ fixed. We shall further consider two cases: $\tilde{\beta} \in (-\infty, 1)$ and $\tilde{\beta} \in (1, \infty)$ in two subsections below. Lemma 4 implies that

$$\mathbb{E}_{\mathbf{n}}[p_0] \approx \begin{cases} \frac{\theta_1 + n_1}{1 - \tilde{\beta}} \frac{1}{n} & \text{when} \quad \tilde{\beta} \in (-\infty, 1) \\ 1 - 1/\tilde{\beta} & \text{when} \quad \tilde{\beta} \in (1, \infty) \end{cases} \qquad \text{as } n \to \infty. \tag{44}$$

Therefore, it makes sense to consider the two cases in scenario (iii).

3.3.1. Case $\tilde{\boldsymbol{\beta}} \in (-\infty, 1)$

In this case, under $\mathbb{P}_{\mathbf{n}}$ and as $n \to \infty$, we have that $p_0 \approx O(1/n)$ by Lemma 4. The genealogy of the n_1 sample type 1 individuals are the same as that in Senario (ii); see Figure 2. This description is precised and justified by Theorems 3-4 and Corollary 1 below.

Figure 2: Conditional genealogy of a sample with observed frequencies (n_1, n_2) at the present time $t = 0$, where $n_1 = 5$ and n_2 is large, and $\beta = \tilde{\beta} n_2$ for a constant $\tilde{\beta} \in (1, \infty)$. The n_2 samples are not shown. In this figure, $K_1 = 2$ and the two red bullets are mutation events from type 1 to type 2. In this scenario (iii), K_1 is distributed like the number of alleles in the Ewens sampling formula, and the timing of the type-1 events are small (of order $O(1/n)$ on the coalescent time scale). The rescaled process $\left(n p_{\frac{t}{n}} \right)_{t \in \mathbb{R}_+}$ is well approximated by the diffusion process (45) with initial distribution $\mathrm{Gam}(n_1 + \theta_1, 1 - \tilde{\beta})$.

Let $(Z_t)_{t \in \mathbb{R}_+}$ be the $\mathbb{R}_+$-valued process that has initial distribution $Z_0 \sim \mathrm{Gamma}(n_1 + \theta_1, 1 - \tilde{\beta})$ and solves the stochastic differential equation

$$dZ_t = \sqrt{Z_t}\, dW_t + \frac{1}{2}(\theta_1 + \tilde{\beta}\, Z_t)\, dt, \qquad t \in \mathbb{R}_+, \tag{45}$$

where W is the Wiener process.

Theorem 3 (Convergence of rescaled genealogy). *Suppose $\tilde{\beta} = \beta/n_2 \in (-\infty, 1)$ is fixed. As $n \to \infty$, the process $\left(n p_{\frac{t}{n}},\ n_1^{obs}(\frac{t}{n}),\ L_1(\frac{t}{n}) \right)_{t \in \mathbb{R}_+}$ converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to a Markov process $\left(Z_t,\ \widetilde{n}_1(t),\ \widetilde{L}_1(t) \right)_{t \in \mathbb{R}_+}$ with state space $\mathbb{R}_+ \times \{0, 1, \cdots, n_1\} \times \{0, 1, \cdots, n_1\}$ described as follows:*

(i) $(Z_t)_{t \in \mathbb{R}_+}$ is a solution to (45) with initial distribution $Z_0 \sim \mathrm{Gamma}(n_1 + \theta_1, 1 - \tilde{\beta})$. In particular, its transition kernel does not depend on $(\widetilde{n}_1, \widetilde{L}_1)$.

(ii) Suppose the current state is (z, a_1, ℓ_1). Then $(\widetilde{n}_1, \widetilde{L}_1)$ evolves as

$$(a_1, \ell_1) \to \begin{cases} (a_1 - 1, \ell_1) & \text{at rate} \quad \frac{1}{z}\binom{a_1}{2} & \text{coalescence of type 1} \\[2mm] (a_1 - 1, \ell_1 + 1) & \text{at rate} \quad \frac{1}{z}\, a_1 \frac{\theta_1}{2} & \text{mutation of 1 to 2} \end{cases} \tag{46}$$

Proof. Let $Y_t := np_{t/n}$. By Lemma 4, under $\mathbb{P}_\mathbf{n}$ we have $Y_0 = np_0 \to \mathrm{Gam}(n_1 + \theta_1, 1 - \tilde{\beta})$. By (1),

$$
\begin{aligned}
Y_t - Y_0 &= n\left(p_{t/n} - p_0\right) \\
&= n\int_0^{t/n} \sqrt{p_s(1-p_s)}\, dW_s + \frac{n}{2}\int_0^{t/n} \theta_1(1-p_s) - \theta_2 p_s + \beta p_s(1-p_s)\, ds \\
&\stackrel{d}{=} \frac{n}{\sqrt{n}} \int_0^t \sqrt{p_{r/n}(1-p_{r/n})}\, dW_r + \frac{n}{2}\int_0^t \theta_1(1-p_{r/n}) - \theta_2 p_{r/n} + \beta p_{r/n}(1-p_{r/n})\, \frac{dr}{n} \\
&= \sqrt{n}\int_0^t \sqrt{\frac{Y_r}{n}\left(1-\frac{Y_r}{n}\right)}\, dW_r + \frac{1}{2}\int_0^t \theta_1\left(1-\frac{Y_r}{n}\right) - \theta_2\frac{Y_r}{n} + \beta\frac{Y_r}{n}\left(1-\frac{Y_r}{n}\right)\, dr \\
&= \int_0^t \sqrt{Y_r\left(1-\frac{Y_r}{n}\right)}\, dW_r + \frac{1}{2}\int_0^t \theta_1\left(1-\frac{Y_r}{n}\right) - \theta_2\frac{Y_r}{n} + \frac{\beta}{n}Y_r\left(1-\frac{Y_r}{n}\right)\, dr, \qquad (47)
\end{aligned}
$$

where we used the fact $\int_0^{t/n} f(s)\, dW_s \stackrel{d}{=} \frac{1}{\sqrt{n}}\int_0^t f(r/n)dW_r$ as processes indexed by $t \in \mathbb{R}_+$.

Using (47), the fact $\sup_n \mathbb{E}[Y_0^2] < \infty$ and the assumption $\tilde{\beta} = \beta/n_2 \in \mathbb{R}$ is fixed, we can check by Gronwall's inequality that $\limsup_{n\to\infty} \mathbb{E}_\mathbf{n}[\sup_{t\in[0,T]} Y_t] < \infty$ for all $T > 0$.

Now, note that equation (45) is the same as (47) after we get rid of the terms $\frac{Y_r}{n}$ and replace $\frac{\beta}{n}$ by $\tilde{\beta}$. As $n \to \infty$, the process $(Y_t)_{t\in[0,T]}$ converges in distribution under $\mathbb{P}_\mathbf{n}$ to a process $(Z_t)_{t\in[0,T]}$ with initial state $Z_0 \sim \mathrm{Gam}(n_1 + \theta_1, 1 - \tilde{\beta})$ and solving (45). This weak convergence can be checked using a standard compactness argument (Billingsley, 2013, Chap. 2) or (Ethier and Kurtz, 2009, Chap. 3). $\qquad\square$

Let $\tilde{\tau}_1 < \tilde{\tau}_2 < \cdots < \tilde{\tau}_{n_1}$ be the jump times of the process $\tilde{n}_1$ in Theorem 3, at each of which the process decreases by 1.

Theorem 4 (Convergence of timing of events)**.** *Suppose $\tilde{\beta} = \beta/n_2 \in (-\infty, 1)$ fixed. As $n \to \infty$, the random vector $(n_2\, \tau_i,\ n_2\, p_{\tau_i})_{i=1}^{n_1}$ under $\mathbb{P}_\mathbf{n}$ converges in distribution to $(\tilde{\tau}_i, Z_{\tilde{\tau}_i})_{i=1}^{n_1}$.*

Proof. We first give a more explicit description of the jump times $\tilde{\tau}_1 < \tilde{\tau}_2 < \cdots < \tilde{\tau}_{n_1}$, in terms of the function

$$
\tilde{\lambda}_{a_1}(z) := \frac{a_1}{2z}(\theta_1 + a_1 - 1)
$$

that comes from (46) in Theorem 3. At the first jump time $\tilde{\tau}_1$, the process $\tilde{n}_1$ decreases from n_1 to $n_1 - 1$. Thus $\tilde{\tau}_1$ is the first jump time of a Poisson process with time inhomogeneous rate $\left(\tilde{\lambda}_{n_1}(Z_t)\right)_{t\in\mathbb{R}_+}$, given the trajectory $(Z_t)_{t\in\mathbb{R}_+}$. Hence,

$$
\mathbb{P}(\tilde{\tau}_1 > t) = \mathbb{E}\left[e^{-\int_0^t \tilde{\lambda}_{n_1}(Z_s)\, ds}\right], \qquad t \in \mathbb{R}_+. \qquad (48)
$$

Given $(\tilde{\tau}_1, Z_{\tilde{\tau}_1})$, the difference $\tilde{\tau}_2 - \tilde{\tau}_1$ is the first jump time of an independent Poisson process with time inhomogeneous rate $\left(\tilde{\lambda}_{n_1-1}(Z_{t+\tilde{\tau}_1})\right)_{t\in\mathbb{R}_+}$. Given $(\tilde{\tau}_2, Z_{\tilde{\tau}_2})$, the difference $\tilde{\tau}_3 - \tilde{\tau}_2$ is the first jump time of an independent Poisson process with time inhomogeneous rate $\left(\tilde{\lambda}_{n_1-2}(Z_{t+\tilde{\tau}_2})\right)_{t\in\mathbb{R}_+}$; and so on. Finally, given $(\tilde{\tau}_{n_1-1}, Z_{\tilde{\tau}_{n_1-1}})$, the difference $\tilde{\tau}_{n_1} - \tilde{\tau}_{n_1-1}$ is the first jump time of an independent Poisson process with time inhomogeneous rate $\left(\tilde{\lambda}_1(Z_{t+\tilde{\tau}_{n_1-1}})\right)_{t\in\mathbb{R}_+}$.

From (28) and Theorem 3, as $n \to \infty$ we have

$$\int_0^{t/n} \lambda_{a_1}(p_s)\, ds = \int_0^t \frac{\lambda_{a_1}(p_{s/n})}{n}\, ds = \int_0^t \frac{a_1}{n\, 2p_{s/n}}\big(a_1 - 1 + (1 - p_{s/n})\theta_1\big)\, ds$$

$$\to \int_0^t \frac{a_1}{2\, Z_s}(\theta_1 + a_1 - 1)\, ds = \int_0^t \widetilde{\lambda}_{a_1}(Z_s)\, ds.$$

Hence $\mathbb{P}_{\mathbf{n}}(n_2\tau_1 > t) \to \mathbb{P}(\widetilde{\tau}_1 > t)$ for all $t \geq 0$, by (48). Combining with Theorem 3, we have that $n_2\,(\tau_1,\; p_{\tau_1})$ under $\mathbb{P}_{\mathbf{n}}$ converges in distribution to $(\widetilde{\tau}_1,\; Z_{\widetilde{\tau}_1})$ as $n_2 \to \infty$.

Applying the strong Markov property of the process Z at $\widetilde{\tau}_1$ and that of the process p at τ_1, we can similarly show that $(\tau_1, \tau_2 - \tau_1, p_{\tau_1}, p_{\tau_2} - p_{\tau_1})\, n_2$ under $\mathbb{P}_{\mathbf{n}}$ converges to $(\widetilde{\tau}_1, \widetilde{\tau}_2 - \widetilde{\tau}_1,\; Z_{\widetilde{\tau}_1}, Z_{\widetilde{\tau}_2} - Z_{\widetilde{\tau}_1})$ in distribution. Continuing in the same way, we obtain that $n_2 \big((\tau_i - \tau_{i-1},\; p_{\tau_i} - p_{\tau_{i-1}})_{i=1}^{n_1}$ under $\mathbb{P}_{\mathbf{n}}$ converges in distribution to $(\widetilde{\tau}_i - \widetilde{\tau}_{i-1},\; Z_{\widetilde{\tau}_i} - Z_{\widetilde{\tau}_{i-1}})_{i=1}^{n_1}$, where $\tau_0 = \widetilde{\tau}_0 = 0$. The desired convergence then follows. $\qquad\square$

In Wakeley et al. (2023, Appendix), we showed that for the case $\beta = 0$ (no selection) and $n_1 > 1$, the jump times $\tau_1 < \tau_2 < \cdots < \tau_{n_1}$ of n_1^{obs} are of order $1/n$ and the re-scaled vector $(n\tau_i)_{i=1}^{n_1}$ converges in distribution. Theorem 4 therefore generalizes the latter convergence in the presence of selection, in scenario (ii) and in the case $\tilde{\beta} \in (-\infty, 1)$ within scenario (iii). This can further be generalized to time-varying populations, as we shall show below. By equation (18) in Wakeley et al. (2023), $\mathbb{E}_{\mathbf{n}}[\tau_1] \approx \frac{2\log n_2}{n_2}$ and $\mathbb{E}_{\mathbf{n}}[\widetilde{\tau}_1] = \infty$ when $n_1 = 1$ and $\beta = 0$.

Theorem 4 implies that when $\tilde{\beta} \in (-\infty, 1)$, the age τ_{n_1} of the oldest latent mutation is well approximated by $\frac{1}{n_2}\widetilde{\tau}_{n_1}$, and the total number of latent mutations for type 1 is predicted by a the Ewens sampling distribution, as $n \to \infty$.

Corollary 1. *Suppose $\tilde{\beta} = \beta/n_2 \in (-\infty, 1)$ fixed. Then as $n \to \infty$,*

(i) K_1 converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to $1 + \sum_{k=2}^{n_1} \eta_k$, where $\{\eta_k\}$ are independent Bernoulli variables taking values in $\{0, 1\}$ and having means $\frac{\theta_1}{\theta_1 + k - 1}$.

(ii) $n_2\,\tau_{n_1}$ converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to $\widetilde{\tau}_{n_1}$.

Proof. The vector $(p_{\tau_i})_{i=1}^{n_1-1}$ converges in probability to the zero vector in $\mathbb{R}^{n_1-1}$. This implies that

$$\Big(h_k(p_{\tau_{n_1-k+1}})\Big)_{k=2}^{n_1} \to \left(\frac{\theta_1}{\theta_1 + k - 1}\right)_{k=2}^{n_1} \in \mathbb{R}^{n_1-1},$$

where we recall $h_k(p) := \frac{(1-p)\theta_1}{(1-p)\theta_1 + k - 1}$ defined in (29). Hence the following weak convergence in $\mathbb{R}^{n_1-1}$ holds:

$$\Big(\xi_k(p_{\tau_{n_1-k+1}})\Big)_{k=2}^{n_1} \xrightarrow{d} (\eta_k)_{k=2}^{n_1}.$$

The proof is complete by (30). $\qquad\square$

3.3.2. Case $\tilde{\beta} \in (1, \infty)$

In this case, under $\mathbb{P}_{\mathbf{n}}$ and as $n \to \infty$, we have that $p_0 \to 1 - 1/\tilde{\beta} > 0$ by (34). The process $(p_t)_{t\in\mathbb{R}_+}$ increases very quickly and stays close to 1. The conditional ancestral process for the n_1 type 1 samples has two parts with different timescales. First it coalesces only as the Kingman coalescent (without mutation) until there is only one lineage. Then it takes a very long time (about $n\frac{2\tilde{\beta}}{\theta_1\theta_2} \approx \frac{2\beta}{\theta_1\theta_2}$) for the single latent mutation to occur. In particular, $K_1 \approx 1$.

This description is justified by Theorems 5-6. See Figure 3 for an illustration.

Figure 3: Conditional genealogy of a sample with observed frequencies (n_1, n_2) at the present time $t = 0$, where $n_1 = 4$ and n_2 is large, and $\beta = \tilde{\beta} n_2$ for a constant $\tilde{\beta} \in (1, \infty)$. The n_2 samples are not shown. This scenario is reminiscent of scenario (i) if we focus on the genealogy of only type 1.

Theorem 5. *Suppose $\tilde{\beta} = \beta/n_2 \in (1, \infty)$ is fixed. As $n \to \infty$, under $\mathbb{P}_{\mathbf{n}}$,*

(i) $\sup_{t \in [S,T]} |1 - p_t| \to 0$ *in probability, for any $0 < S < T < \infty$; and*

(ii) *For any $T > 0$, the process $(n_1^{obs}(t), L_1(t))_{t \in [0,T]}$ converges in distribution to a process $(\tilde{n}^{obs}(t), 0)_{t \in [0,T]}$ where $\tilde{n}^{obs}$ is a pure death process with jump rate $\binom{k}{2} = \frac{k(k-1)}{2}$ from k to $k - 1$.*

(iii) $K_1 \to 1$ *in probability.*

Proof. We first observe that the process p gets close to 1 quickly, when $\tilde{\beta} \in (1, \infty)$, in the sense that for any $\epsilon \in (0, 1)$,

$$\lim_{n \to \infty} \mathbb{P}_{\mathbf{n}} (\tau_{1-\epsilon} > S) = \lim_{n \to \infty} \mathbb{P}_{\mathbf{n}} \left(\sup_{t \in [0,S]} p_t < 1 - \epsilon \right) = 0,$$

where $\tau_{1-\epsilon} := \inf\{t \in \mathbb{R}_+ : p_t > 1 - \epsilon\}$ is the first time p hits above $1 - \epsilon$. This is true because $p_0 \to 1 - 1/\tilde{\beta} \in (0, 1)$ in probability, so that the growth term $\tilde{\beta} n_2 \, p_t (1 - p_t)$ is large at least when $t > 0$ is small. Next, suppose the process p starts at $1 - \epsilon$ (i.e. the process $q = 1 - p$ starts at ϵ), we show that the exit time of the process q out of the interval $[0, 2\epsilon)$ is longer than T with probability tending to 1, as $n \to \infty$. More precisely,

$$\mathbb{P}_{1-\epsilon} \left(\inf_{t \in [0, T]} p_t < 1 - 2\epsilon \right) = \mathbb{P}_\epsilon \left(\sup_{t \in [0,T]} q_t > 2\epsilon \right).$$

The last display tends to 0 as $n_2 \to \infty$, as in the proof of Theorem 1(i).

From these two estimates and the strong Markov property of the process p, we have that for

any $\epsilon \in (0,1)$,

$$
\mathbb{P}_{\mathbf{n}}\left(\sup_{t\in[S,T]}|1-p_t|>2\epsilon\right)
$$

$$
\leq \mathbb{P}_{\mathbf{n}}\left(\sup_{t\in[S,T]}|1-p_t|>2\epsilon,\ \tau_{1-\epsilon}\leq S\right)+\mathbb{P}_{\mathbf{n}}\left(\tau_{1-\epsilon}>S\right)
$$

$$
=\mathbb{E}_{\mathbf{n}}\left[1_{\{\tau_{1-\epsilon}\leq S\}}\,\mathbb{P}_{1-\epsilon}\left(\inf_{t\in[S-\tau_{1-\epsilon},\,T-\tau_{1-\epsilon}]}p_t<1-2\epsilon\right)\right]+\mathbb{P}_{\mathbf{n}}\left(\tau_{1-\epsilon}>S\right)
$$

$$
\leq \mathbb{P}_{1-\epsilon}\left(\inf_{t\in[0,T]}p_t<1-2\epsilon\right)+\mathbb{P}_{\mathbf{n}}\left(\tau_{1-\epsilon}>S\right)\to 0 \qquad \text{as } n\to\infty.
$$

The proof of part (i) is complete.

By Part (i) and (28), the times for the type 1 events are of order $O(1)$ and $h_k(p_t)\to 0$ where k_k is defined in (29). Hence Parts (ii) and (iii) follow. $\qquad\square$

Now we consider the second part of the genealogy, when there is a single lineage left (i.e. during τ_{n_1-1} and τ_{n_1}).

To estimate the age τ_{n_1} of the single latent mutation, we can ignore the n_1-1 jump times of the process $\widetilde{n}^{obs}$ (since they are of order 1 by Theorem 5). The frequency of type 1 is tightly regulated in the sense that it is close to 1 in the sense of Theorem 5(i). However, we need to know "how close it is to 1" in order to get an estimate of τ_{n_1}, because simply setting $p=1$ in $\frac{1-p}{2p}\theta_1$ will give us zero.

We quantify the difference $1-p_t$ by rescaling the process $q:=1-p$ as in Theorem 3. The emergent process is very similar to (45). Let $(\widetilde{q}_t)_{t\in\mathbb{R}_+}$ be the $\mathbb{R}_+$-valued process that solves the stochastic differential equation

$$
d\widetilde{q}_t=\sqrt{\widetilde{q}_t}\,dW_t+\frac{1}{2}(\theta_2-\tilde{\beta}\widetilde{q}_t)\,dt,\qquad t\in\mathbb{R}_+, \tag{49}
$$

where W is the Wiener process. This process $\widetilde{q}$ is a mean reverting Cox-Ingersoll-Ross (CIR) model. It will never touch zero if $\theta_2\geq 1$; otherwise it can occasionally touch the boundary zero point. The CIR model is an ergodic process with stationary distribution given by the Gamma distribution $\mathrm{Gam}(\theta_2,\tilde{\beta})$, which has probability density function $\frac{\tilde{\beta}^{\theta_2}}{\Gamma(\theta_2)}\,y^{\theta_2-1}\,e^{-\tilde{\beta}y}$.

Lemma 6. *Suppose $\tilde{\beta}=\beta/n_2\in\mathbb{R}$ is fixed. Let $q_t=1-p_t$ for $t\in\mathbb{R}_+$. Suppose $nq_0\to c_0$ as $n\to\infty$. Then the process $(nq_{\frac{t}{n}})_{t\in\mathbb{R}_+}$ converges in distibution to the solution process $\widetilde{q}$ to (49) starting at c_0.*

Proof. Following the proof of Lemma 5, we use (35), let $Y_t:=nq_{\frac{t}{n}}$ and suppose $Y_0\to c_0$. Then, similar to (41),

$$
Y_t-Y_0=n\left(q_{t/n}-p_0\right)
$$

$$
\overset{d}{=}\int_0^t\sqrt{Y_r\left(1-\frac{Y_r}{n}\right)}\,dW_r+\frac{1}{2}\int_0^t\theta_2\left(1-\frac{Y_r}{n}\right)-\theta_1\frac{Y_r}{n}-\frac{\beta}{n}Y_r\left(1-\frac{Y_r}{n}\right)\,dr. \tag{50}
$$

Using the assumption that $\tilde{\beta}=\beta/n_2\in\mathbb{R}$ is fixed, the desired weak convergence can be proved by a standard compactness argument (Billingsley, 2013, Chap. 2) or (Ethier and Kurtz, 2009, Chap. 3). $\qquad\square$

Theorem 6 below is analogous to Theorem 2. We obtain that the mean of the age τ_{n_1} is about $n\frac{2\tilde{\beta}}{\theta_1\theta_2} \approx \frac{2\beta}{\theta_1\theta_2}$ when it is larger than $\frac{2n}{\theta_1\theta_2}$ and n is large.

Theorem 6 (Age of the unique latent mutation)**.** *Suppose $\tilde{\beta} = \beta/n_2 \in (1,\infty)$ is fixed. As $n \to \infty$, $\frac{\tau_{n_1}}{n}$ converges in distribution under $\mathbb{P}_{\mathbf{n}}$ to an exponential random variable with mean $\frac{2\tilde{\beta}}{\theta_1\theta_2}$. That is,*

$$\frac{\tau_{n_1}}{n} \xrightarrow{d} \mathrm{Exp}\left(\frac{\theta_1\,\theta_2}{2\,\tilde{\beta}}\right) \qquad as \quad n \to \infty.$$

Proof. We argue as in the proof of Theorem 2. For any $t \in \mathbb{R}_+$, as $n \to \infty$, by Theorem 5,

$$\mathbb{P}_{\mathbf{n}}\left(\frac{A_1}{n} > t\right) \approx \mathbb{E}\left[\exp\left\{\frac{-\theta_1\,t}{2n^2t}\int_0^{n^2t}\frac{nq_{r/n}}{1-q_{r/n}}\,dr\right\}\right].$$

Note that in the exponent, by Lemma 6,

$$\lim_{n\to\infty}\frac{-\theta_1\,t}{2n^2t}\int_0^{n^2t}\frac{nq_{r/n}}{1-q_{r/n}}\,dr \stackrel{d}{=} \frac{-\theta_1\,t}{2}\lim_{T\to\infty}\frac{1}{T}\int_0^T\widetilde{q}(r)\,dr = \frac{-\theta_1\,t}{2}\mathbb{E}_\infty[\widetilde{q}],$$

where the last equality holds almost surely because the process $\widetilde{q}$ solving (49) is ergodic. Here $\mathbb{E}_\infty$ is the expectation with respect to the stationary distribution of the process $\widetilde{q}$ to which it converges as $t \to \infty$.

The limiting distribution of $\widetilde{q}_t$, as $t \to \infty$, is known to be the Gamma distribution $\mathrm{Gam}(\theta_2, \tilde{\beta})$. Hence $\mathbb{E}_\infty[\widetilde{q}] = \theta_2/\tilde{\beta}$. $\qquad\square$

3.4. Time-varying population size

For a nonconstant size population, namely one that has $\rho(t)N$ individuals at forward time t where ρ is a non-constant "demographic" function, neither the Moran process nor its diffusion approximation possess a stationary distribution. However, the random background approach of Barton et al. (2004) can be generalized to this setting by considering the time-reversed frequency process.

Our main message here is that the limiting pictures in scenarios (i) and (ii) are robust against changing population sizes and the initial distribution μ_0 of X_0. Let $\rho : \mathbb{R}_+ \to (0,\infty)$ be a continuous and strictly positive function. Under the conditional probability $\mathbb{P}_{\mathbf{n}}$, the same pictures for the conditional coalescence process for scenarios (i) and (ii) still hold in their respective limits. For scenario (iii), the limiting picture depends on the large deviation behavior of $\mathbb{P}_{\mu_0}(X_T \in dy)$ as $n \to \infty$. Roughly speaking, this robustness comes from the fact that by taking either β or $n_2 \sim n$ large, the events among finite-count alleles in the sample are so sped up that the population size will have hardly changed by the time all their coalescent and latent mutation events have occurred.

More precisely, in the diffusion limit of the Moran process, the relative frequency of A_1 in the population at time t satisfies the time-inhomogeneous stochastic differential equation

$$dX(t) = \left[\frac{\theta_1}{2}(1 - X(t)) - \frac{\theta_2}{2}X(t) + \frac{\beta}{2}X(t)(1 - X(t))\right]dt + \sqrt{\frac{X(t)(1 - X(t))}{\rho(t)}}\,dW_t, \qquad (51)$$

in which W_t is the Wiener process; see for instance Evans et al. (2007). Let $T > 0$ be the present time (forward) which is assumed to be large. Then the backward frequency process is

$$p_t := X(T - t) \qquad for \quad t \in [0, T], \qquad (52)$$

which is the time-reversal of the process X, therefore $(p_t)_{t \in [0,T]}$ is still Markovian (Millet et al., 1989). Note that the law of the present time frequency of type 1, $p_0 = X(T) = X_T$, now depends on the distribution μ_0 of the initial frequency $X(0)$. This law is denoted by $\mathbb{P}_{\mu_0}(X_T \in dy)$.

Suppose a sample of n individuals are picked uniformly at random at the present time $T > 0$, and we know that n_1 of them are of type 1 (and $n - n_1$ are of type 2). Let $\mathbb{P}_{\mathbf{n}} = \mathbb{P}_{\mathbf{n}, \mu_0}$ be the conditional probability measure given the sample count $\mathbf{n} = (n_1, n_2)$, and $\mathcal{L}^{\mathbf{n}} := \mathbb{P}_{\mathbf{n}}(X_T \in dy) = \mathbb{P}_{\mu_0}(X_T \in dy \,|\, \mathbf{n}) \in \mathcal{P}([0,1])$ be the conditional law of $p_0 = X(T)$ under $\mathbb{P}_{\mathbf{n}}$. Then

$$\mathcal{L}^{\mathbf{n}} = C_T \, y^{n_1} (1 - y)^{n_2} \, \mathbb{P}_{\mu_0}(X_T \in dy), \tag{53}$$

where $C_T = \left(\int_0^1 y^{n_1} (1 - y)^{n_2} \mathbb{P}_{\mu_0}(X_T \in dy) \right)^{-1}$ is a normalizing constant. This follows from Bayes' theorem, just like (5) did, but with prior distribution $\mathbb{P}_{\mu_0}(X_T \in dy)$.

Similar to Proposition 1, the conditional ancestral process in the diffusion limit can be described as follows.

Proposition 2 (Conditional ancestral process). *Let $T > 0$ and $\mu_0 \in \mathcal{P}([0,1])$ be fixed, and $\rho : [0,T] \to (0, \infty)$ be any bounded function. The process $(p_t, n_1^{obs}(t), L_1(t), n_2^{obs}(t), L_2(t))_{t \in [0,T]}$ under $\mathbb{P}_{\mathbf{n}}$ is a Markov process with state space $[0,1] \times \{0, 1, \cdots, n_1\}^2 \times \{0, 1, \cdots, n_2\}^2$ described as follows:*

(i) $(p_t)_{t \in [0,T]}$, defined by (52), has the law of $(X(T - t))_{t \in [0,T]}$ under $\mathbb{P}_{\mathbf{n}}$. In particular, it has initial distribution $\mathcal{L}^{\mathbf{n}}$ in (53) and it does not depend on the process $(n_1^{obs}, L_1, n_2^{obs}, L_2)$.

(ii) The process $(n_1^{obs}, L_1, n_2^{obs}, L_2)$ starts at $(n_1, 0, n_2, 0)$. When the current state is $(p, a_1, \ell_1, a_2, \ell_2)$, this process evolves as

$$(a_1, \ell_1) \to \begin{cases} (a_1 - 1, \ell_1) & \text{at rate} \quad \frac{1}{p} \binom{a_1}{2} & \text{coalescence of type } 1^{obs} \\ (a_1 - 1, \ell_1 + 1) & \text{at rate} \quad \frac{1-p}{p} a_1 \frac{\theta_1}{2} & \text{mutation of } 1^{obs} \text{ to } 2 \end{cases} \tag{54}$$

and, independently,

$$(a_2, \ell_2) \to \begin{cases} (a_2 - 1, \ell_2) & \text{at rate} \quad \frac{1}{1-p} \binom{a_2}{2} & \text{coalescence of type } 2^{obs} \\ (a_2 - 1, \ell_2 + 1) & \text{at rate} \quad \frac{p}{1-p} a_2 \frac{\theta_2}{2} & \text{mutation of } 2^{obs} \text{ to } 1 \end{cases} \tag{55}$$

Remark 1. A more explicit description for the process $(X_t)_{t \in [0,T]}$, hence also for its time-reversal $(p_t = X_{T-t})_{t \in [0,T]}$, can be obtained by Doob's h-transform (Doob, 1957, 2001). More precisely, we define the function

$$h(t, x) := \mathbb{P}(\mathbf{n} \,|\, X_t = x) = \mathbb{E}\left[\binom{n}{n_1} X_T^{n_1} (1 - X_T)^{n_2} \,\Big|\, X_t = x \right].$$

Then $(X_t)_{t \in [0,T]}$ under the conditional probability $\mathbb{P}_{\mathbf{n}}$ solves the following SDE that is different from (51).

$$dX_t = \left[b(X_t) + \frac{X_t(1 - X_t)}{\rho(t)} \, \partial_x \log h(t, X_t) \right] dt + \sqrt{\frac{X_t(1 - X_t)}{\rho(t)}} \, d\widetilde{W}_t,$$

where $b(x) := \frac{\theta_1}{2}(1-x) - \frac{\theta_2}{2}x + \frac{\beta}{2}x(1-x)$ is the drift term in (51), and $\widetilde{W}$ is a Brownian motion. Criteria for which the process $(p_t)_{t \in [0,T]}$ satisfies a stochastic differential equation can be found in Millet et al. (1989).

Next, we look at asymptotics. The following analogue of Lemma 1 holds for any initial distribution μ_0 of $X(0)$ and any demographic function ρ that is bounded and positive. Note that in Lemma 1, $\mu_0 = \phi_\beta$ depends on β, but here μ_0 is fixed.

Proposition 3. *Let $T > 0$ and $\mu_0 \in \mathcal{P}([0,1])$ be fixed, and $\rho : [0,T] \to (0,\infty)$ be any bounded function. The following convergences in $\mathcal{P}([0,1])$ hold.*

  (i) Suppose n_2 is fixed and $\beta \to \infty$. Then $\mathcal{L}^{\mathbf{n}} \to \delta_1$.

  (ii) Suppose β is fixed and $n_2 \to \infty$. Then $\mathcal{L}^{\mathbf{n}} \to \delta_0$.

  (iii) Suppose $\tilde{\beta} \in \mathbb{R}$ is fixed and $\beta = \tilde{\beta}n_2 \to \infty$. Suppose $\mathbb{P}_{\mu_0}(X_T \in dy)$ has a density $p(T, \mu_0, y)\, dy$ and there exists a large deviation rate function $\mathcal{I} : [0,1] \to [0,\infty]$ such that

$$\frac{\log p(T, \mu_0, y)}{n_2} \to \mathcal{I}(y) \qquad \text{as } n \to \infty.$$

Suppose also that $(1-y)\, e^{\mathcal{I}(y)}$ has a unique maximum at $y_* \in [0,1]$. Then $\mathcal{L}^{\mathbf{n}} \to \delta_{y_*}$, and $\mathcal{I}(y_*) = \frac{1}{1-y_*}$.

Remark 2. The assumptions in (iii) holds when $\rho(t) \equiv 1$ and $\mu_0 = \phi_\beta$ in (2), i.e. constant population size with stationary initial condition. In this case, $\mathcal{I}(y) = \tilde{\beta}\, y$ is the linear function, and $\mathcal{L}^{\mathbf{n}} = \phi_\beta^{(n_1,n_2)}$ in (21). The rate function has a phase transition at $\tilde{\beta} = 1$, as shown in Lemma 1. Namely, $y_* = 0$ when $\tilde{\beta} \in (-\infty, 1)$ and $y_* = 1 - 1/\tilde{\beta}$ when $\tilde{\beta} \in (1, \infty)$. It would be interesting to check whether a similar phase transition occur for general demographic function $\rho(t)$.

Remark 3. The large deviation principle for $\mathbb{P}_{\mu_0}(X_T \in dy)$ as $n_2 \to \infty$ can be checked using the Gärtner-Ellis theorem (Dembo, 2009, Theorem 2.3.6). When it holds, the rate function $\mathcal{I}$ is equal to the Legendre transform of the function

$$\Lambda(\gamma) := \lim_{n_2 \to \infty} \frac{1}{n_2} \log \mathbb{E}_{\mu_0}[e^{\gamma\, n_2\, X_T}].$$

Proof. A proof follows from that of Lemma 1. Let $f \in C_b([0,1])$, a bounded continuous function on $[0,1]$. Then

$$\mathbb{E}_{\mathbf{n}}[f(p_0)] = \int_0^1 f(y)\, \mathcal{L}^{\mathbf{n}}(dy) = \frac{\mathbb{E}_{\mu_0}[f(X_T)\, X_T^{n_1}(1-X_T)^{n_2}]}{\mathbb{E}_{\mu_0}[X_T^{n_1}(1-X_T)^{n_2}]}. \tag{56}$$

For part (i), note that if (n_1, n_2) is fixed and $\beta \to \infty$, then $\mathbb{P}_{\mu_0}(X(T) < 1 - \epsilon) \to 0$ for any $\epsilon > 0$ as in the proof of Theorem 5(i). Hence $\mathbb{E}_{\mu_0}[|f(X(T)) - f(1)|] \to 0$ for any $f \in C_b([0,1])$. In particular, $\mathbb{P}_{\mu_0}(X_T \in dy) \to \delta_1$. Hence the right hand side of (53) tends to δ_1 in $\mathcal{P}([0,1])$, as $\beta \to \infty$.

For part (ii), note that $(1-y)^{n_2}$ has maximum at $y = 0$, and $y^{n_1} \mathbb{P}_{\mu_0}(X_T \in dy)$ has nothing to do with n_2. Hence the right hand side of (53) tends to δ_0 in $\mathcal{P}([0,1])$, as $n_2 \to \infty$.

For part (iii), the numerator of (56) is

$$\mathbb{E}_{\mu_0}[f(X_T)\,X_T^{n_1}(1-X_T)^{n_2}] = \int_0^1 f(y)\,y^{n_1}(1-y)^{n_2}\,\mathbb{P}_{\mu_0}(X_T \in dy)$$

$$\approx \int_0^1 f(y)\,y^{n_1}\,[(1-y)\,e^{\mathcal{I}(y)}]^{n_2}\,\varphi(y)\,dy,$$

for some function φ such that $\frac{\ln \varphi(y)}{n_2} \to 0$, by assumptions of part (iii). Since $(1-y)\,e^{\mathcal{I}(y)}$ has a unique maximum at y_*, $\lim_{n \to \infty} \mathbb{E}_{\mathbf{n}}[f(p_0)] = f(y_*)$ by (56) and a standard argument as in the proof of Lemma 1. $\qquad\square$

By Proposition 2 and Proposition 3, the same limiting pictures for the conditional coalescence process for scenarios (i) and (ii) hold for arbitrary continuous function $\rho : \mathbb{R}_+ \to (0, \infty)$. Continuity of ρ ensures that ρ is bounded away from zero on any compact time interval $[0, T]$ and therefore analogous approximations for the frequency process $(p_t)_{t \in [0,T]}$ under $\mathbb{P}_{\mathbf{n}}$ (such as Lemma 5) still hold, where the new approximating functions now involve the h function in Remark 1.

More precisely, in scenario (i), Theorem 1 still holds, and Theorem 2 still holds but with a possibly different limiting random variable. Hence $s_{n_2} \sim O(1/\beta)$ is very small, $K_1 \to 1$ and the single latent mutation for type 1 is very old. The approximation in Lemma 5 will be replaced by an analogous statement that involves the h function in Remark 1. In scenario (ii), $\tau_{n_1} \sim O(1/n)$ is very small and $K_1 \approx 1 + \sum_{k=2}^{n_1} \eta_k$, where $\{\eta_k\}$ are independent Bernoulli variables taking values in $\{0,1\}$ and having means $\frac{\theta_1}{\theta_1+k-1}$. Theorem 3 with $\beta \in \mathbb{R}$ fixed still holds, but the statement need to be modified because the approximating process Z in (45) will be replaced by another one that involves the h function in Remark 1, as was needed in modifying Lemma 5.

Scenario (iii) is harder to analyze and we leave it for future work. We conjecture that if $y_* = 0$, then the conditional genealogy behaves like scenario (ii); and if $y_* \in (0,1]$, then the conditional genealogy behaves like scenario (i).

4. Conditional ancestral selection graph

Our aim in this section is to see how the results of the previous section can be obtained from a different model: the ancestral selection graph. We concentrate on the ancestry of focal allele A_1 and on constant population size, and proceed more heuristically than in the previous section.

The ancestral selection graph is an augmented coalescent model for the joint distribution of the gene genealogy and the allelic states of the sample (Krone and Neuhauser, 1997; Neuhauser and Krone, 1997). It includes the usual coalescent rate 1 per pair of lineages and mutation rate $\theta/2$ per lineage. Additionally, under the stationary model of Section 1, it includes a *branching* rate of $|\beta|/2$ per lineage. When a branching event occurs, the lineage splits into an *incoming* lineage and a *continuing* lineage. One of these is *real*, meaning it is included in the gene genealogy. The other is *virtual*, meaning it is there only to model the gene genealogy correctly with selection. Which is which could be resolved if their allelic states were known: the incoming lineage is real if its allelic type is the one favored by selection, otherwise the continuing lineage is real. But the allelic states are not known in the construction of the ancestral selection graph.

The conditional ancestral selection graph models gene genealogies given a sample with allelic states specified (Slade, 2000a,b; Fearnhead, 2001, 2002; Stephens and Donnelly, 2003; Baake and Bialowons, 2008). In this case it is known which lineages are real and which are virtual. This

allows a simplification in which there is a reduced rate of branching and only virtual lineages of the disfavored type are produced (Slade, 2000a). A second simplification is possible if mutation is parent-independent: then any lineage which mutates may be discarded (Fearnhead, 2002).

We assume parent-independent mutation, specifically $\theta_1 = \theta\pi_1$ and $\theta_2 = \theta\pi_2$, with $\pi_1 + \pi_2 = 1$. Any two-allele mutation model can be restated in this way, leaving the stationary frequency distribution (2) and the sampling probability (4) unchanged. But doing so introduces 'spurious mutations to one's own type' (Donnelly, 1986) or 'empty mutations' (Baake and Bialowons, 2008) which have no biological reality and in particular should not be counted as latent mutations. Adding them to our model allows us to discard real A_2 lineages and any virtual lineage once these mutate, but we must distinguish between actual mutations and empty mutations in the ancestry of A_1.

The resulting conditional process tracks the numbers real and virtual ancestral lineages from the present time $t = 0$ back into the past. Let $r_1(t)$, $r_2(t)$ and $v_i(t)$, where $i = 1$ if $\beta < 0$ or $i = 2$ if $\beta > 0$, be the numbers of real type-1, real type-2 and virtual type-i lineages at past time t. The process begins in state $r_1(0) = n_1$, $r_2(0) = n_2$, $v_i(0) = 0$ and stops the first time $r_1(t) + r_2(t) = 1$. We suppress t in what follows, and focus on the instantaneous rates of transition of the process.

The conditional ancestral process is obtained by considering rates of events in the unconditional process, which has total rate $(r_1 + r_2 + v_i)(\theta + |\beta| + r_1 + r_2 + v_i - 1)/2$, then weighting rates of events depending on how likely they are to produce the sample. Rates of some events are down-weighted to zero. For instance, the sample could not have been obtained if there were a coalescent event between lineages with different allelic types, whereas in the unconditional process these happen with rate $r_1 r_2$ plus either $r_2 v_1$ or $r_1 v_2$, depending on whether $\beta < 0$ or $\beta > 0$.

Rates of events for which the sample has a non-zero chance of being observed are up-weighted or down-weighted by ratios of sampling probabilities like (4). This method of conditioning a Markov process on its eventual outcome is stated simply in Kemeny and Snell (1960, p. 64), a familiar example being the Wright-Fisher diffusion conditioned on eventual fixation (Ewens, 2004, p. 89), and is characterized more generally by Doob's h-transform (Doob, 1957, 2001). In the conditional ancestral selection graph, the Markov process is the (unconditional) ancestral process of Krone and Neuhauser (1997) and the eventual outcome is the sample with allelic states specified.

In our formulation, the samples and their ancestral lineages all are distinguishable, which we denote with a subscript "o" for ordered as in Wakeley et al. (2023). The probability of any particular allelic configuration in the ancestry of the sample, in which there are r_1 lineages of type 1, r_2 lineages of type 2 and v_i lineages of type $i \in \{1, 2\}$, is

$$q_o(r_1, r_2, v_1) = \int_0^1 x^{r_1 + v_1}(1 - x)^{r_2}\phi_\beta(x)dx \quad \text{if } \beta < 0 \tag{57}$$

$$q_o(r_1, r_2, v_2) = \int_0^1 x^{r_1}(1 - x)^{r_2 + v_2}\phi_\beta(x)dx \quad \text{if } \beta > 0 \tag{58}$$

with $\phi_\beta(x)$ as in (2). Note, the additional binomial coefficient in the sampling probability (4) is the number of possible orderings of a sample containing n_1 and n_2 copies of A_1 and A_2.

The rate of any particular event with non-zero probability in the conditional process is the product of its rate in the unconditional process and a ratio of sampling probabilities from either (57) or (58). For event $(r_1, r_2, v_i) \to (r_1', r_2', v_i')$, the required ratio is $q_o(r_1', r_2', v_i')/q_o(r_1, r_2, v_i)$. The denominator $q_o(r_1, r_2, v_i)$ is the probability of the sample given all events so far in the conditional ancestral process, which have led to the current state (r_1, r_2, v_i), and the numerator $q_o(r_1', r_2', v_i')$ is the probability of the sample given these events and the event $(r_1, r_2, v_i) \to (r_1', r_2', v_i')$. Appendix

section Appendix B provides the details of how the minimal ancestral process we use here to model latent mutations in the ancestry of the sampled copies of allele A_1 is obtained from the full conditional ancestral process, using the simplifications of Slade (2000a) and Fearnhead (2002).

The resulting conditional ancestral process differs depending on whether $\beta < 0$ or $\beta > 0$ but in either case it includes five possible transitions from state (r_1, r_2, v_i). If $\beta < 0$,

$$(r_1, r_2, v_1) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_1) & \text{at rate} \quad r_1 \frac{\theta \pi_1}{2} \frac{q_o(r_1-1, r_2+1, v_1)}{q_o(r_1, r_2, v_1)} \\[2mm] (r_1 - 1, r_2, v_1) & \text{at rate} \quad \binom{r_1}{2} \frac{q_o(r_1-1, r_2, v_1)}{q_o(r_1, r_2, v_1)} \\[2mm] (r_1, r_2, v_1 + 1) & \text{at rate} \quad (r_1 + r_2 + v_1) \frac{|\beta|}{2} \frac{q_o(r_1, r_2, v_1+1)}{q_o(r_1, r_2, v_1)} \\[2mm] (r_1, r_2, v_1 - 1) & \text{at rate} \quad \left(v_1 \frac{\theta \pi_1}{2} + r_1 v_1 + \binom{v_1}{2} \right) \frac{q_o(r_1, r_2, v_1-1)}{q_o(r_1, r_2, v_1)} \\[2mm] (r_1, r_2 - 1, v_1) & \text{at rate} \quad \left(r_2 \frac{\theta \pi_2}{2} + \binom{r_2}{2} \right) \frac{q_o(r_1, r_2-1, v_1)}{q_o(r_1, r_2, v_1)} \end{cases} \tag{59}$$

whereas if $\beta > 0$,

$$(r_1, r_2, v_2) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_2) & \text{at rate} \quad \frac{r_1 \theta \pi_1}{2} \frac{q_o(r_1-1, r_2+1, v_2)}{q_o(r_1, r_2, v_2)} \\[2mm] (r_1 - 1, r_2, v_2) & \text{at rate} \quad \binom{r_1}{2} \frac{q_o(r_1-1, r_2, v_2)}{q_o(r_1, r_2, v_2)} \\[2mm] (r_1, r_2, v_2 + 1) & \text{at rate} \quad (r_1 + r_2 + v_2) \frac{\beta}{2} \frac{q_o(r_1, r_2, v_2+1)}{q_o(r_1, r_2, v_2)} \\[2mm] (r_1, r_2, v_2 - 1) & \text{at rate} \quad \left(v_2 \frac{\theta \pi_2}{2} + r_2 v_2 + \binom{v_2}{2} \right) \frac{q_o(r_1, r_2, v_2-1)}{q_o(r_1, r_2, v_2)} \\[2mm] (r_1, r_2 - 1, v_2) & \text{at rate} \quad \left(r_2 \frac{\theta \pi_2}{2} + \binom{r_2}{2} \right) \frac{q_o(r_1, r_2-1, v_2)}{q_o(r_1, r_2, v_2)} \end{cases} \tag{60}$$

which differ owing to the different resolutions of branching events when $\beta < 0$ versus $\beta > 0$. We may note that the total rates of events in (59) and (60) are less than in the unconditional ancestral process because the conditional process has a reduced rate of branching (Slade, 2000a) and because empty mutations do not change the number or types of ancestral lineages. If $\beta < 0$, the total rate is $r_1 \theta \pi_1 / 2 + r_2 |\beta| / 2$ less, whereas if $\beta > 0$, it is $r_1 \theta \pi_1 / 2 + r_1 \beta / 2$ less.

Asymptotic approximations for the ratios $q_o(r_1', r_2', v_i') / q_o(r_1, r_2, v_i)$ in these rates of events can be obtained using the results in Appendix section Appendix A. In the following three subsections we present approximations to the conditional ancestral process for our three scenarios of interest: (i) $|\beta|$ large with n_2 fixed, (ii) n_2 large with β fixed, and (iii) both $|\beta|$ and n_2 large with $\tilde{\beta} = \beta / n_2$ fixed. Because initially $r_2 = n_2$, we consider r_2 large in the scenarios with n_2 large. For each scenario, we compute the rates events up to leading order in $|\beta|$ or r_2, then consider how these conform to the corresponding results of Section 3.

4.1. Scenario (i): strong selection, arbitrary sample size

Here $|\beta|$ is large with n_2 fixed, along with n_1 and θ. In Section 3.1, Theorem 1, we treated the ancestries of A_1 and A_2 simultaneously as $\beta \rightarrow +\infty$, so that A_1 was favored and A_2 was disfavored. Here we cover these same two possibilities by modeling the ancestry of A_1, using (59) when A_1 is disfavored ($\beta < 0$) and (60) when A_1 is favored ($\beta > 0$). We disregard the ancestry of the non-focal allele A_2 except insofar as it is needed to model events in the ancestry of A_1.

When $\beta < 0$, using (A.4a) in (59) gives

$$(r_1, r_2, v_1) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_1) & \text{at rate} \quad r_1 \frac{|\beta|}{2} \frac{\theta \pi_1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\ (r_1 - 1, r_2, v_1) & \text{at rate} \quad r_1 \frac{|\beta|}{2} \frac{r_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\ (r_1, r_2, v_1 + 1) & \text{at rate} \quad \frac{(r_1 + r_2 + v_1)(\theta \pi_1 + r_1 + v_1)}{2} + O(|\beta|^{-1}) \\ (r_1, r_2, v_1 - 1) & \text{at rate} \quad v_1 \frac{|\beta|}{2} \frac{\theta \pi_1 + 2 r_1 + v_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\ (r_1, r_2 - 1, v_1) & \text{at rate} \quad r_2 \frac{\theta \pi_2}{2} + \binom{r_2}{2} + O(|\beta|^{-1}) \end{cases} \qquad (61)$$

for the ancestry of a disfavored allele under strong selection (as $\beta \to -\infty$). Latent mutations and coalescent events occur with rates proportional to $|\beta|$. Virtual lineages are removed similarly quickly but are produced at a much lower rate. So v_1 will stay zero during the $O(1/|\beta|)$ time it takes for the requisite n_1 latent mutations or coalescent events to occur. Then the analogous result to (39), namely (7) and (8), follows from the first two lines of (61). Coalescence and mutation among the copies of A_2 occur at the slower rate, so none of these should occur before all the type-1 lineages disappear. These results were first suggested in Wakeley (2008).

When $\beta > 0$, using (A.4b) in (60) gives

$$(r_1, r_2, v_2) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_2) & \text{at rate} \quad r_1 \frac{1}{\beta} \frac{\theta \pi_1 (\theta \pi_2 + r_2 + v_2)}{2} + O(\beta^{-2}) \\ (r_1 - 1, r_2, v_2) & \text{at rate} \quad \binom{r_1}{2} + O(\beta^{-1}) \\ (r_1, r_2, v_2 + 1) & \text{at rate} \quad \frac{(r_1 + r_2 + v_2)(\theta \pi_2 + r_2 + v_2)}{2} + O(\beta^{-1}) \\ (r_1, r_2, v_2 - 1) & \text{at rate} \quad v_2 \frac{\beta}{2} \frac{\theta \pi_2 + 2 r_2 + v_2 - 1}{\theta \pi_2 + r_2 + v_2 - 1} + O(1) \\ (r_1, r_2 - 1, v_2) & \text{at rate} \quad r_2 \frac{\beta}{2} \frac{\theta \pi_2 + r_2 - 1}{\theta \pi_2 + r_2 + v_2 - 1} + O(1) \end{cases} \qquad (62)$$

for the ancestry of a favored allele under strong selection (as $\beta \to +\infty$). Now A_2 is undergoing the fast process just described for A_1 in (61), so these lineages will disappear quickly. Again the rate of removal of virtual lineages greatly exceeds their rate of production. In $O(1/\beta)$ time, the ancestral state will become $(r_1, r_2, v_2) = (n_1, 0, 0)$. But now with A_1 favored, the rates of coalescence and latent mutation differ by a factor of β, so the first $n_1 - 1$ events will be coalescent events, followed by a long wait for a single latent mutation with mean $\theta^2 \pi_1 \pi_2 / (2\beta)$ as in Theorem 2.

4.2. Scenario (ii): arbitrary selection, large sample size

Here n_2 is large with β fixed, along with n_1 and θ. Because $r_2 = n_2$ at the start of the ancestral process, we present rates of events to leading order in $1/r_2$. In Section 3.2 we deferred this scenario to Section 3.3, because in the limit it is equivalent to $\tilde{\beta} = 0$. Of course, there are two ways for $\tilde{\beta}$ to approach zero, and the sign of $\tilde{\beta}$ matters in (45) for any $\tilde{\beta}$ not strictly equal to zero. Here we consider the two cases, $\beta < 0$ and $\beta > 0$, separately.

When $\beta < 0$, using (A.5) in (59) gives

$$
(r_1, r_2, v_1) \to
\begin{cases}
(r_1 - 1, r_2 + 1, v_1) & \text{at rate} \quad r_1 \frac{r_2}{2} \frac{\theta \pi_1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1 - 1, r_2, v_1) & \text{at rate} \quad r_1 \frac{r_2}{2} \frac{r_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1, r_2, v_1 + 1) & \text{at rate} \quad \frac{|\beta|(\theta \pi_1 + r_1 + v_1)}{2} + O(r_2^{-1}) \\[2mm]
(r_1, r_2, v_1 - 1) & \text{at rate} \quad v_1 \frac{r_2}{2} \frac{\theta \pi_1 + 2 r_1 + v_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1, r_2 - 1, v_1) & \text{at rate} \quad \frac{r_2^2}{2} + O(r_2)
\end{cases}
\tag{63}
$$

This differs from the neutral case (Wakeley et al., 2023) only by the possibility of virtual lineages, which again will be removed quickly if they are produced. With $v_2 = 0$, (7) and (8) follow, the process of latent mutation and coalescence happens in $O(1/r_2)$ time, and the argument in Wakeley et al. (2023, Appendix) guarantees that these conclusions hold despite the very fast decrease of r_2. Theorem 4 addresses the corresponding issues for the model of Section 3.

When $\beta > 0$, using (A.5) in (60) gives

$$
(r_1, r_2, v_2) \to
\begin{cases}
(r_1 - 1, r_2 + 1, v_2) & \text{at rate} \quad r_1 \frac{r_2}{2} \frac{\theta \pi_1}{\theta \pi_1 + r_1 - 1} + O(1) \\[2mm]
(r_1 - 1, r_2, v_2) & \text{at rate} \quad r_1 \frac{r_2}{2} \frac{r_1 - 1}{\theta \pi_1 + r_1 - 1} + O(1) \\[2mm]
(r_1, r_2, v_2 + 1) & \text{at rate} \quad \frac{r_2 |\beta|}{2} + O(1) \\[2mm]
(r_1, r_2, v_2 - 1) & \text{at rate} \quad v_2 r_2 + O(1) \\[2mm]
(r_1, r_2 - 1, v_2) & \text{at rate} \quad \frac{r_2^2}{2} + O(r_2)
\end{cases}
\tag{64}
$$

which differs from (63) in two ways. Now the rate of production of virtual lines is non-negligible. But here their presence does not affect the rates of latent mutation and coalescence. Again we have (7) and (8), and the process of latent mutation and coalescence happens in $O(1/r_2)$ time.

4.3. Scenario (iii): strong selection, large sample size

Here both $|\beta|$ and n_2 are large with $\tilde{\beta} = \beta / n_2$ fixed, along with n_1 and θ. Again since the process begins with $r_2 = n_2$, we present rates of events to leading order in $1/r_2$. Because the conditional ancestral process differs for $\beta < 0$ versus $\beta > 0$, i.e. with (59) and (60), and the asymptotic approximation we use for the hypergeometric function differs for $\tilde{\beta} < 1$ versus $\tilde{\beta} > 1$, i.e. with (A.6a) and (A.6b), here we have three cases. Note these are the same three cases in (11a), (11b) and (11c).

When $\tilde{\beta} < 0$, using (A.6a) in (59) gives

$$
(r_1, r_2, v_1) \to
\begin{cases}
(r_1 - 1, r_2 + 1, v_1) & \text{at rate} \quad r_1 \frac{r_2\left(1 + |\tilde{\beta}|\right)}{2} \frac{\theta \pi_1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1 - 1, r_2, v_1) & \text{at rate} \quad r_1 \frac{r_2\left(1 + |\tilde{\beta}|\right)}{2} \frac{r_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1, r_2, v_1 + 1) & \text{at rate} \quad \frac{|\tilde{\beta}|(\theta \pi_1 + r_1 + v_1)}{2\left(1 + |\tilde{\beta}|\right)} + O(r_2^{-1}) \\[2mm]
(r_1, r_2, v_1 - 1) & \text{at rate} \quad v_1 \frac{r_2\left(1 + |\tilde{\beta}|\right)}{2} \frac{\theta \pi_1 + 2 r_1 + v_1 - 1}{\theta \pi_1 + r_1 + v_1 - 1} + O(1) \\[2mm]
(r_1, r_2 - 1, v_1) & \text{at rate} \quad \frac{r_2^2}{2} + O(r_2)
\end{cases}
\tag{65}
$$

which is comparable to (61) and (63). Again we may effectively ignore virtual lineages. The rates of latent mutation and coalescence in (61) and (63) differ only by the interchange of r_2 for $|\beta|$. In (65), the factor $r_2\left(1+|\tilde{\beta}|\right)$ encompasses the effects of both. The larger $|\tilde{\beta}|$ is, the more quickly these events will occur, and again (7) and (8) describe the number of latent mutations.

When $0 < \tilde{\beta} < 1$, using (A.6a) in (60) gives

$$(r_1, r_2, v_2) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_2) & \text{at rate} \quad r_1 \frac{r_2\left(1-\tilde{\beta}\right)}{2} \frac{\theta\pi_1}{\theta\pi_1 + r_1 - 1} + O(1) \\[2mm] (r_1 - 1, r_2, v_2) & \text{at rate} \quad r_1 \frac{r_2\left(1-\tilde{\beta}\right)}{2} \frac{r_1 - 1}{\theta\pi_1 + r_1 - 1} + O(1) \\[2mm] (r_1, r_2, v_2 + 1) & \text{at rate} \quad \frac{r_2\tilde{\beta}}{2} + O(1) \\[2mm] (r_1, r_2, v_2 - 1) & \text{at rate} \quad v_2 r_2 + O(1) \\[2mm] (r_1, r_2 - 1, v_2) & \text{at rate} \quad \frac{r_2^2}{2} + O(r_2) \end{cases} \tag{66}$$

which is comparable to (64). In contrast to (65), now with A_1 favored, the larger $\tilde{\beta}$ is (i.e. the closer it is to 1) the smaller the rates of latent mutation and coalescence become. Otherwise, for any given $\tilde{\beta}$, the same conclusions regarding latent mutations and their timing follow from (66) as from (65), and these conform to what is stated in Corollary 1.

When $\tilde{\beta} > 1$, using (A.6b) in (60) gives

$$(r_1, r_2, v_2) \rightarrow \begin{cases} (r_1 - 1, r_2 + 1, v_2) & \text{at rate} \quad r_1 \frac{\theta\pi_1}{2} \frac{1}{\tilde{\beta} - 1} + O\left(r_2^{-1}\right) \\[2mm] (r_1 - 1, r_2, v_2) & \text{at rate} \quad \binom{r_1}{2} \frac{\tilde{\beta}}{\tilde{\beta} - 1} + O\left(r_2^{-1}\right) \\[2mm] (r_1, r_2, v_2 + 1) & \text{at rate} \quad \frac{r_2}{2} + O(1) \\[2mm] (r_1, r_2, v_2 - 1) & \text{at rate} \quad v_2 r_2 \tilde{\beta} + O(1) \\[2mm] (r_1, r_2 - 1, v_2) & \text{at rate} \quad \frac{r_2^2}{2}\tilde{\beta} + O(r_2) \end{cases} \tag{67}$$

which paints a very different picture. Whereas (61), (63), (64), (65) and (66) all give the Ewens sampling result described by (7) and (8) and have these events occurring quickly on the coalescent time scale, (67) is rather like (62) in that the rates of latent mutation and coalescence are too slow to register on the time scale of events involving the non-focal allele A_2. So, an instantaneous transition occurs. In (67), it is a transition from r_2 large to r_2 comparable to r_1. Then this case (67) will collapse quickly to the corresponding case (62) where coalescence without mutation will happen among the A_1 followed by a long wait for a single latent mutation. For the model in Section 3.3.2, this is described by Theorem 5 and Theorem 6. Finally we may note that initial the rates of latent mutation and coalescence in (67) are precisely those predicted for the model in Section 3.3.2 from (26) starting at $p_0 \rightarrow 1 - 1/\tilde{\beta}$ as specified for $\tilde{\beta} \in (1, \infty)$ in (34).

5. Discussion

In this paper, we have considered a two allele model at a single genetic locus in a large haploid population with possibly time-varying size. The locus is subject to mutation and selection. Suppose a sample of size n was drawn uniformly from the infinite population under the diffusion approximation.

By extending the framework of Barton et al. (2004), we described the asymptotic behaviors of the conditional genealogy and the number of latent mutations of the sample, given the sample frequencies of the two alleles. This moves beyond our previous work (Wakeley et al., 2023) by the inclusion of selection and by the use of an entirely different model, i.e. coalescence in a random background (Barton et al., 2004). This yields novel results, including for example, in the strong selection case in which the selection strength β is proportional to the sample size n and both go to infinity (our scenario (iii)), that under rescaling the genealogy of the rare allele can be described in terms of a Cox-Ingersoll-Ross (CIR) diffusion with an initial Gamma distribution.

The concept of rare allele in this paper and in Wakeley et al. (2023) is the same as the one considered by Joyce and Tavaré (1995) and Joyce (1995). It focuses on the counts of the alleles in a large sample rather than their relative frequencies in the population. In scenarios (ii) and (iii) in this paper we consider a fixed number n_1 of the rare type 1 when the sample size n tends to infinity. Joyce and Tavaré (1995) considered rare alleles in a large sample drawn from the stationary distribution of a d-dimensional Wright-Fisher diffusion with selection and mutation. They showed that the allele counts of rare alleles have approximately independent Poisson distributions with parameters that do *not* depend on the selection parameters. The case $d = 2$ with genic selection in Joyce and Tavaré (1995) corresponds to scenario (ii) in our work here when β is fixed. Our results for scenario (iii) therefore not only extend the Poisson convergence in Joyce and Tavaré (1995) to the case $\lim_{n\to\infty} \beta/n \in (-\infty, 1)$, but also contrast it with the case $\lim_{n\to\infty} \beta/n \in (1, \infty)$. In particular, the effects of selection can still be washed out in the limiting process, even if selection grows with the sample size n, in both the positive and the negative directions.

Joyce (1995) obtained a similar result for a Wright-Fisher diffusion with selection and infinite-alleles mutation. This diffusion process is a particular case of the Fleming-Viot process (see the survey Ethier and Kurtz (1993)) and it has a unique stationary distribution denoted ν_{selec}. Joyce (1995) considered a large sample of size n drawn from ν_{selec}. Let $C_b(n) \in \mathbb{Z}_+^b$ be the first b allele counts in a sample of size n drawn from the stationary distribution, and K_n be the total number of alleles in the sample. Joyce (1995) showed that for any fixed b, the distribution of $(C_b(n), K_n)$ under ν_{selec} is arbitrarily close to that under the neutral model. It would be interesting to know if analogous results for our scenario (iii) also holds for the infinite allele model. In particular, is there a threshold for the selection strength relative to n that control whether selection are washed out or not in the limit as $n \to \infty$?

Many open questions remain. For time-varying populations, little is known in scenario (iii). For example, will the assumptions in Proposition 3 hold for general demographic function? Will there be a phase transition for the value of y_* in terms of $\tilde{\beta}$ and if so, what will determine the phase transition? Also, both our results and those of Joyce and Tavaré (1995) and Joyce (1995) are for the infinite-population diffusion limit. Further consideration of the issues raised in Section 2.1.1 is needed to assess the relevance of these results to various kinds of finite populations.

Finally, we have ignored the possibility of spatial structure. Spatially heterogeneous populations in which reproduction rates, death rates, mutation rates and selection strength can depend both on spatial position and local population density present challenges. This is because the population dynamics now takes place in high or infinite dimension (Hallatschek and Nelson, 2008; Barton et al., 2010; Durrett and Fan, 2016; Louvet and Véber, 2022; Etheridge et al., 2023). For example, the spatial version of (1), the stochastic Fisher-Kolmogorov-Petrovsky-Piscunov (FKPP) equation introduced by Shiga (1988), is a stochastic partial differential equation that arises as the scaling limit of various discrete models under weak selection (Mueller and Tribe, 1995; Durrett and Fan,

2016; Fan, 2021). Under the stochastic FKPP, Hallatschek and Nelson (2008) and Durrett and Fan (2016) studied the backward-time lineage dynamics of a single sample individual, conditioned on knowing its type. It would be interesting to see if our results in this paper can be extended to spatial stochastic models with selection.

Acknowledgements

We thank Alison Etheridge for raising the question about the applicability of our limiting results to Wright-Fisher reproduction (cf. Section 2.1.1). We also thank Daniel Rickert and Kejia Geng for assistance in producing the figures. Research partially supported by National Science Foundation grants DMS-1855417 and DMS-2152103, and Office of Naval Research grant N00014-20-1-2411 to Wai-Tong (Louis) Fan.

Appendix A. Asymptotic approximations used in the text

From the series expansion for a ratio of gamma functions with a common large parameter, 6.1.47 in Abramowitz and Stegun (1964) or equation (1) in Tricomi and Erdélyi (1951), we have

$$\frac{\Gamma(a+n_2)}{\Gamma(b+n_2)} = n_2^{a-b}\left(1 + \frac{(a-b)(a+b-1)}{2n_2} + O\left(n_2^{-2}\right)\right) \tag{A.1}$$

for constants a and b which will depend on the application. For example, we can apply (A.1) twice in the sampling probability (4) when n_2 is large: once with $a = n_1 + 1$ and $b = 1$ (in the binomial coefficient) and once with $a = \theta_2$ and $b = \theta_1 + \theta_2 + n_1$.

The confluent hypergeometric function is commonly defined in terms of the integral

$$_1F_1\left(a; b; z\right) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 e^{zu} u^{a-1}(1-u)^{b-a-1} du \tag{A.2}$$

or in terms of the series

$$_1F_1\left(a; b; z\right) = \sum_{k=0}^{\infty} \frac{a^{(k)} z^k}{b^{(k)} k!} \tag{A.3}$$

which converge for all $z \in \mathbb{R}$ and $b > a > 0$, where $a^{(k)}$ is the rising factorial $a(a+1)\cdots(a+k-1)$ with $a^{(0)} = 1$. Again a and b depend on the context, e.g. as in (3) and (4). The parameter z corresponds to the selection parameter β.

For large $|\beta|$ and with constant a and b,

$$_1F_1\left(a; b; \beta\right) = \begin{cases} \dfrac{\Gamma(b)}{\Gamma(b-a)}|\beta|^{-a}\left(1 - \dfrac{a(b-a-1)}{|\beta|} + O\left(|\beta|^{-2}\right)\right) & \text{if } \beta < 0 \qquad \text{(A.4a)} \\[2ex] \dfrac{\Gamma(b)}{\Gamma(a)}e^{\beta}\beta^{a-b}\left(1 - \dfrac{(b-a)(a-1)}{\beta} + O\left(\beta^{-2}\right)\right) & \text{if } \beta > 0 \qquad \text{(A.4b)} \end{cases}$$

where the middle, neutral case is given only for completeness. Equation (A.4a) is from (4.1.2) in Slater (1960), and (A.4b) is from (4.1.6) in Slater (1960) or may be obtained from (A.4a) using Kummer's first theorem which appears as (1.4.1) in Slater (1960).

For large n_2, with constants a, b and z,

$$_1F_1\left(a; b+n_2; z\right) = 1 + \frac{az}{n_2} + O\left(n_2^{-2}\right) \tag{A.5}$$

directly from (A.3).

For large n_2 and $\beta = \tilde{\beta} n_2$, with constants a and b,

$$_1F_1\left(a; b+n_2; \tilde{\beta} n_2\right) \approx \begin{cases} \left(1 - \tilde{\beta}\right)^{-a} & \text{if } \tilde{\beta} < 1 \qquad \text{(A.6a)} \\[2ex] \dfrac{\sqrt{2\pi}}{\Gamma(a)}\left(1 - \dfrac{1}{\tilde{\beta}}\right)^{a-1}\left(\dfrac{1}{\tilde{\beta}}\right)^{b-a+n_2} n_2^{a-\frac{1}{2}} e^{n_2\left(\tilde{\beta}-1\right)} & \text{if } \tilde{\beta} > 1 \qquad \text{(A.6b)} \end{cases}$$

which we present only to leading order for simplicity. Equation (A.6a) follows from (A.3) and (A.6b) was obtained by applying Laplace's method to the integral in (A.2) for this case.

Appendix B. Events in the conditional ancestral selection graph

Here we show how the minimal conditional ancestral process in Section 4 is obtained from the full conditional ancestral process. To begin, we assume that at some time in the conditional ancestral process there were r_1, r_2, v_1 and v_2 real and virtual lineages of type 1 and type 2. The associated sampling probability is $q_o(r_1, r_2, v_1, v_2)$, the straightforward extension of (57) or (58) to include both type-1 and type-2 virtual lineages. How branching events are resolved depends on which allele is favored by selection. We begin here by assuming that A_2 is favored, or $\beta < 0$. Grouping events by the types of lineages involved (real or virtual of type 1 or type 2) then by whether it is mutation,

branching or coalescence gives fourteen possibilities which occur at the following rates.

$$r_1 \frac{\theta \pi_1}{2} \left(\frac{q_o(r_1, r_2, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1 - 1, r_2 + 1, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.1}$$

$$r_2 \frac{\theta \pi_2}{2} \left(\frac{q_o(r_1, r_2, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1 + 1, r_2 - 1, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.2}$$

$$v_1 \frac{\theta \pi_1}{2} \left(\frac{q_o(r_1, r_2, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1, r_2, v_1 - 1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.3}$$

$$v_2 \frac{\theta \pi_2}{2} \left(\frac{q_o(r_1, r_2, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1, r_2, v_1 + 1, v_2 - 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.4}$$

$$r_1 \frac{|\beta|}{2} \frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.5}$$

$$r_2 \frac{|\beta|}{2} \left(2 \frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.6}$$

$$v_1 \frac{|\beta|}{2} \frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.7}$$

$$v_2 \frac{|\beta|}{2} \left(2 \frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.8}$$

$$\binom{r_1}{2} \frac{q_o(r_1 - 1, r_2, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.9}$$

$$\binom{r_2}{2} \frac{q_o(r_1, r_2 - 1, v_1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.10}$$

$$r_1 v_1 \frac{q_o(r_1, r_2, v_1 - 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.11}$$

$$r_2 v_2 \frac{q_o(r_1, r_2, v_1, v_2 - 1)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.12}$$

$$\binom{v_1}{2} \frac{q_o(r_1, r_2, v_1 - 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.13}$$

$$\binom{v_2}{2} \frac{q_o(r_1, r_2, v_1, v_2 - 1)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.14}$$

The sum of (B.1) through (B.14) is equal to the total rate of events in the unconditional ancestral process, $(r_1 + r_2 + v_1 + v_2)(\theta + |\beta| + r_1 + r_2 + v_1 + v_2 - 1)/2$. Twenty-two distinct events $(r_1, r_2, v_1, v_2) \to (r'_1, r'_2, v'_1, v'_2)$ are represented, one for each of the ratios of sampling probabilities, $q_0(r'_1, r'_2, v'_1, v'_2)/q_0(r_1, r_2, v_1, v_2)$. Note that the assumption of parent-independent mutation leads to the four kinds of spurious or empty mutation events in (B.1) through (B.4) which do not change the ancestral state of the sample $(r'_1 = r_1, r'_2 = r_2, v'_1 = v_1, v'_2 = v_2)$. Also, only those events which have have non-zero probabilities of giving the data appear in (B.1) through (B.14); coalescent events

between lineages with different types and type-i mutation events on type $3 - i$ lineages would make the data impossible.

Recall that the resolution of branching events depends on which allele is favored by selection. The events and their probabilities in (B.5) through (B.8) are just for the case $\beta < 0$, where A_2 is the favored allele. Each branching event creates an incoming lineage and a continuing lineage, both of which may be of type 1 or type 2. Let (I, C) be the types of these lineages. In (B.5) and (B.7), only one of the four (I, C) pairs has non-zero probability of producing the data: $(I = 1, C = 1)$ corresponding to the event $(r_1, r_2, v_1, v_2) \to (r_1, r_2, v_1 + 1, v_2)$. In (B.6) and (B.8), the possibility $(I = 1, C = 1)$ is discarded as it would then be impossible for the descendant lineage to be of type 2. The other three possibilities have non-zero chances of producing the data, and associated events

$$(r_1, r_2, v_1, v_2) \to \begin{cases} (r_1, r_2, v_1 + 1, v_2) & \text{when } (I = 1, C = 2) & \text{(B.15a)} \\ (r_1, r_2, v_1 + 1, v_2) & \text{when } (I = 2, C = 1) & \text{(B.15b)} \\ (r_1, r_2, v_1, v_2 + 1) & \text{when } (I = 2, C = 2) \,. & \text{(B.15c)} \end{cases}$$

In contrast, if $\beta > 0$ then branching events on type-2 lineages are the ones for which only one of the four (I, C) pairs has non-zero probability of producing the data: $(I = 2, C = 2)$ corresponding to the event $(r_1, r_2, v_1, v_2) \to (r_1, r_2, v_1, v_2 + 1)$. When $\beta > 0$, if the branching event occurs on a type-1 lineage, then in place of (B.15a), (B.15b) and (B.15c) we have

$$(r_1, r_2, v_1, v_2) \to \begin{cases} (r_1, r_2, v_1 + 1, v_2) & \text{when } (I = 1, C = 1) & \text{(B.16a)} \\ (r_1, r_2, v_1, v_2 + 1) & \text{when } (I = 1, C = 2) & \text{(B.16b)} \\ (r_1, r_2, v_1, v_2 + 1) & \text{when } (I = 2, C = 1) \,. & \text{(B.16c)} \end{cases}$$

Therefore, when $\beta > 0$, (B.5) through (B.8) must be replaced with

$$r_1 \frac{\beta}{2} \left(\frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + 2 \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.17}$$

$$r_2 \frac{\beta}{2} \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.18}$$

$$v_1 \frac{\beta}{2} \left(\frac{q_o(r_1, r_2, v_1 + 1, v_2)}{q_o(r_1, r_2, v_1, v_2)} + 2 \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \right) \tag{B.19}$$

$$v_2 \frac{\beta}{2} \frac{q_o(r_1, r_2, v_1, v_2 + 1)}{q_o(r_1, r_2, v_1, v_2)} \tag{B.20}$$

Equations (B.1) through (B.7) and (B.9) through (B.14) are the same for $\beta > 0$ and $\beta < 0$.

The simplifications discovered by Slade (2000a) and Fearnhead (2002) follow from the simple fact that each sampled lineage is either of type 1 or type 2. Slade (2000a) noticed that when both the descendant lineage and the incoming lineage have the favored type, the type of the continuing lineage does not matter so there is no need to introduce a new virtual lineage. Instead, these two possibilities can be collapsed into a single null event which does not change the numbers and types of ancestral lineages. That is, we can use

$$q_o(r_1, r_2, v_1 + 1, v_2) + q_o(r_1, r_2, v_1, v_2 + 1) = q_o(r_1, r_2, v_1, v_2) \tag{B.21}$$

in (B.6), (B.8), (B.17) and (B.19). As a result, no type-2 virtual lineages will be created.

Along the same lines, Fearnhead (2002) noticed that when mutation is parent-independent there is no need to follow ancestral lineages once they have mutated, because the ancestral lineage could be of either type. Any such lineage can be removed from the ancestral process. Here we use

$$q_o(r_1, r_2, v_1, v_2) + q_o(r_1 + 1, r_2 - 1, v_1, v_2) = q_o(r_1, r_2 - 1, v_1, v_2) \qquad \text{(B.22)}$$

in (B.2), and other appropriate identities in (B.3) and (B.4). But we do not make use of this simplification in (B.1) because our specific goal is to model latent mutations in the ancestry of A_1. These are actual mutations, where the ancestral type was A_2. The remaining $A_1 \to A_1$ empty mutations are null events, which do not change the numbers and types of ancestral lineages.

The conditional ancestral processes for $\beta < 0$ and $\beta > 0$ given by (59) and (60) in the main text each include just five kinds of (non-null) events. We obtain these by applying the simplifications of Slade (2000a) and Fearnhead (2002) then grouping events by their outcomes. For example, the coalescent events in (B.10) have effect $r_2 \to r_2 - 1$, as do the combined mutations in (B.2) once the simplification of Fearnhead (2002) is applied. So these appear together as one kind of event, the fifth case in both (59) and (60).

We do not include null events in (59) and (60) since these by definition have no effect on the ancestral lineages. In the case $\beta < 0$, the null events are empty mutations on type-1 real lineages and branching events on type-2 real lineages where the incoming line is also of type 2. These occur with total rate $r_1 \theta \pi_1 / 2 + r_2 |\beta| / 2$. In the case $\beta > 0$, the null events are empty mutations on type-1 real lineages and branching events on type-1 real lineages where the incoming line is also of type 1. These occur with total rate $r_1 \theta \pi_1 / 2 + r_1 \beta / 2$.

References

Abramowitz, M., Stegun, I.A., 1964. Handbook of Mathematical Functions. Dover, New York.

Arratia, R., Barbour, A.D., Tavaré, S., 1992. Poisson process approximations for the Ewens sampling formula. The Annals of Applied Probability 2, 519 – 535. doi:`10.1214/aoap/1177005647`.

Baake, E., Bialowons, R., 2008. Ancestral processes with selection: branching and Moran models. Banach Center Publications 80, 33–52. doi:`10.4064/bc80-0-2`.

Barton, N.H., Etheridge, A.M., 2004. The effect of selection on genealogies. Genetics 166, 1115–1131. doi:`10.1093/genetics/166.2.1115`.

Barton, N.H., Etheridge, A.M., Sturm, A.K., 2004. Coalescence in a random background. The Annals of Applied Probability 14, 754 – 785. doi:`10.1214/105051604000000099`.

Barton, N.H., Etheridge, A.M., Véber, A., 2010. A new model for evolution in a spatial continuum. Electronic Journal of Probability 15, 162–216. doi:`10.1214/EJP.v15-741`.

Billingsley, P., 2013. Convergence of probability measures. John Wiley & Sons.

Chen, J., Glémin, S., Lascoux, M., 2020. From drift to draft: How much do beneficial mutations actually contribute to predictions of Ohta's slightly deleterious model of molecular evolution? Genetics 214, 1005–1018. doi:`10.1534/genetics.119.302869`.

Crane, H., 2016. The ubiquitous Ewens sampling formula. Statistical Science 31, 1–19. doi:`10.1214/15-STS529`.

Darden, T., Kaplan, N.L., Hudson, R.R., 1989. A numerical method for calculating moments of coalescent times in finite populations with selection. Journal of Mathematical Biology 27, 355–368. doi:`10.1007/BF00275818`.

Dembo, A., 2009. Large deviations techniques and applications. Springer.

Donnelly, P., 1986. Dual processes in population genetics, in: Tautu, P. (Ed.), Stochastic Spatial Processes, Springer Berlin Heidelberg, Berlin, Heidelberg. pp. 94–105. doi:`10.1007/BFb0076240`.

Doob, J.L., 1957. Conditional brownian motion and the boundary limits of harmonic functions. Bulletin de la Société mathématique de France 86, 431–458. URL: `http://www.numdam.org/article/BSMF_1957__85__431_0.pdf`.

Doob, J.L., 2001. Classical Potential Theory and Its Probabilistic Counterpart. Springer-Verlag Berlin, Heidelberg. doi:`10.1007/978-3-642-56573-1`.

Dufresne, D., 2001. The integrated square-root process. Centre for Actuarial Studies, Department of Economics, University of Melbourne.

Durrett, R., Fan, W.T.L., 2016. Genealogies in expanding populations. Ann. Appl. Probab. 26, 3456–3490. doi:`10.1214/16-AAP1181`.

Etheridge, A., 2011. Some Mathematical Models from Population Genetics: École D'Été de Probabilités de Saint-Flour XXXIX-2009. Springer Berlin, Heidelberg. doi:`10.1007/978-3-642-16632-7`.

Etheridge, A., Pfaffelhuber, P., Wakolbinger, A., 2006. An approximate sampling formula under genetic hitchhiking. The Annals of Applied Probability 16, 685–729. doi:`10.1214/105051606000000114`.

Etheridge, A.M., Kurtz, T.G., Letter, I., Ralph, P.L., Lung, T.T.H., 2023. Looking forwards and backwards: dynamics and genealogies of locally regulated populations. arXiv preprint arXiv:2305.14488 .

Ethier, S.N., Kurtz, T.G., 1993. Fleming–viot processes in population genetics. SIAM Journal on Control and Optimization 31, 345–386.

Ethier, S.N., Kurtz, T.G., 2009. Markov processes: characterization and convergence. John Wiley & Sons.

Evans, S.N., Shvets, Y., Slatkin, M., 2007. Non-equilibrium theory of the allele frequency spectrum. Theoretical population biology 71, 109–119.

Ewens, W.J., 1972. The sampling theory of selectively neutral alleles. Theoretical Population Biology 3, 87–112. doi:`10.1016/0040-5809(72)90035-4`.

Ewens, W.J., 2004. Mathematical Population Genetics, Volume I: Theoretical Foundations. Springer-Verlag, Berlin.

Eyre-Walker, A., Keightley, P.D., 2007. The distribution of fitness effects of new mutations. Nature Reviews Genetics 8, 610–618. doi:`10.1038/nrg2146`.

Fan, W.T.L., 2021. Stochastic PDEs on graphs as scaling limits of discrete interacting systems. Bernoulli 27, 1899–1941. doi:`10.3150/20-bej1296`.

Fearnhead, P., 2001. Perfect simulation from population genetic models with selection. Theoretical Population Biology 59, 263–279. doi:`https://doi.org/10.1006/tpbi.2001.1514`.

Fearnhead, P., 2002. The common ancestor at a nonneutral locus. Journal of Applied Probability 39, 38–54. doi:`10.1239/jap/1019737986`.

Fisher, R.A., 1930. The Genetical Theory of Natural Selection. Clarendon, Oxford.

Hallatschek, O., Nelson, D.R., 2008. Gene surfing in expanding populations. Theoretical population biology 73, 158–170.

Harpak, A., Bhaskar, A., Pritchard, J.K., 2016. Mutation rate variation is a primary determinant of the distribution of allele frequencies in humans. PLoS Genetics 12, e1006489. doi:`10.1371/journal.pgen.1006489`.

Heathcote, C.R., 1965. A branching process allowing immigration. Journal of the Royal Statistical Society. Series B (Methodological) 27, 138–143. URL: `http://www.jstor.org/stable/2984491`.

Herbots, H.M., 1997. The structured coalescent, in: Donnelly, P., Tavaré, S. (Eds.), Progress in Population Genetics and Human Evolution (IMA Volumes in Mathematics and its Applications, vol. 87. Springer-Verlag, New York, pp. 231–255.

Johnson, K.E., Adams, C.J., Voight, B.F., 2022. Identifying rare variants inconsistent with identity-by-descent in population-scale whole-genome sequencing data. Methods in Ecology and Evolution 13, 2429–2442. doi:`10.1111/2041-210X.13991`.

Joyce, P., 1995. Robustness of the Ewens sampling formula. Journal of Applied Probability 32, 609–622. doi:`10.2307/3215116`.

Joyce, P., Tavaré, S., 1995. The distribution of rare alleles. Journal of Mathematical Biology 33, 602–618. doi:`10.1007/BF00298645`.

Kaplan, N.L., Darden, T., Hudson, R.R., 1988. The coalescent process in models with selection. Genetics 120, 819–829. doi:`10.1093/genetics/120.3.819`.

Karatzas, I., Shreve, S.E., 1991. Brownian motion and stochastic calculus. volume 113. Springer Science & Business Media.

Karlin, S., McGregor, J., 1964. On some stochastic models in genetics, in: Gurland, J. (Ed.), Stochastic Models in Medicine and Biology. The University of Wisconsin Press, Madison, pp. 245–271.

Kemeny, J.G., Snell, J.L., 1960. Finite Markov Chains. D Van Nostrand Company, New York.

Kendall, D.G., 1949. Stochastic processes and population growth. Journal of the Royal Statistical Society. Series B (Methodological) 11, 230–282. URL: `http://www.jstor.org/stable/2984078`.

Krone, S.M., Neuhauser, C., 1997. Ancestral processes with selection. Theoretical Population Biology 51, 210–237. doi:`10.1006/tpbi.1997.1299`.

Leffler, E.M., Bullaughey, K., Matute, D.R., Meyer, W.K., Séurel, L., Venkat, A., Andolfatto, P., Przeworski, M., 2012. Revisiting an old riddle: What determines genetic diversity levels within species? PLOS Biology 10, 1–9. doi:`10.1371/journal.pbio.1001388`.

Louvet, A., Véber, A., 2022. Growth properties of the infinite-parent spatial lambda-fleming viot process. arXiv preprint arXiv:2205.03937 .

Millet, A., Nualart, D., Sanz, M., 1989. Integration by parts and time reversal for diffusion processes. The Annals of Probability , 208–238.

Moran, P.A.P., 1958. Random processes in genetics. Proc. Camb. Phil. Soc. 54, 60–71. doi:`10.1017/S0305004100033193`.

Moran, P.A.P., 1962. Statistical Processes of Evolutionary Theory. Clarendon Press, Oxford.

Mueller, C., Tribe, R., 1995. Stochastic pde's arising from the long range contact and long range voter processes. Probability theory and related fields 102, 519–545.

Nagylaki, T., 1990. Models and approximations for random genetic drift. Theoretical Population Biology 37, 192–212. doi:`10.1016/0040-5809(90)90035-T`.

Neuhauser, C., Krone, S.M., 1997. The genealogy of samples in models with selection. Genetics 145, 519–534. doi:`10.1093/genetics/145.2.519`.

Notohara, M., 1990. The coalescent and the genealogical process in geographically structured population. Journal of Mathematical Biology 29, 59–75. doi:`10.1007/BF00173909`.

Seplyarskiy, V., Lee, D.J., Koch, E.M., Lichtman, J.S., Luan, H.H., Sunyaev, S.R., 2022. A mutation rate model at the basepair resolution identifies the mutagenic effect of Polymerase III transcription. bioRxiv doi:`10.1101/2022.08.20.504670`. doi: 10.1101/2022.08.20.504670.

Seplyarskiy, V.B., Soldatov, R.A., Koch, E., McGinty, R.J., Goldmann, J.M., Hernandez, R.D., Barnes, K., Correa, A., Burchard, E.G., Ellinor, P.T., McGarvey, S.T., Mitchell, B.D., Vasan, R.S., Redline, S., Silverman, E., Weiss, S.T., Arnett, D.K., Blangero, J., Boerwinkle, E., He, J., Montgomery, C., Rao, D.C., Rotter, J.I., Taylor, K.D., Brody, J.A., Chen, Y.D.I., de las Fuentes, L., Hwu, C.M., Rich, S.S., Manichaikul, A.W., Mychaleckyj, J.C., Palmer, N.D., Smith, J.A., Kardia, S.L.R., Peyser, P.A., Bielak, L.F., O'Connor, T.D., Emery, L.S., NHLBI Trans-Omics for Precision Medicine (TOPMed) Consortium, TOPMed Population Genetics Working Group, Gilissen, C., Wong, W.S.W., Kharchenko, P.V., Sunyaev, S., 2021. Population sequencing data reveal a compendium of mutational processes in the human germ line. Science 373, 1030–1035. doi:`10.1126/science.aba7408`.

Shiga, T., 1988. Stepping stone models in population genetics and population dynamics, in: Albeverio, S., Blanchard, P., Hazewinkel, M., Streit, L. (Eds.), Stochastic processes in physics and engineering. Springer, pp. 345–355.

Slade, P.F., 2000a. Most recent common ancestor probability distributions in gene genealogies under selection. Theoretical Population Biology 58, 291–305. doi:`10.1006/tpbi.2000.1488`.

Slade, P.F., 2000b. Simulation of selected genealogies. Theoretical Population Biology 57, 35–49. doi:`10.1006/tpbi.1999.1438`.

Slater, L.J., 1960. Confluent Hypergeometric Functions. Cambridge University Press, Cambridge.

Stephens, M., Donnelly, P., 2003. Ancestral inference in population genetics models with selection (with Discussion). Australian & New Zealand Journal of Statistics 45, 395–430. doi:`10.1111/1467-842X.00295`.

Takahata, N., 1988. The coalescent in two partially isolated diffusion populations. Genetics Research 52, 213–222. doi:`10.1017/S0016672300027683`.

Tavaré, S., 2021. The magical Ewens sampling formula. Bulletin of the London Mathematical Society 53, 1563–1582. doi:`10.1112/blms.12537`.

Tricomi, F., Erdélyi, A., 1951. The asymptotic expansion of a ratio of gamma functions. Pacific Journal of Mathematics 1, 133–142. doi:`10.2140/pjm.1951.1.133`.

Vanyolos, A., Cho, M., Glasgow, S.A., 2014. Probability density of the cir model. Available at SSRN 2508699 .

Wakeley, J., 2008. Conditional gene genealogies under strong purifying selection. Molecular Biology and Evolution 25, 2615–2626. doi:`10.1093/molbev/msn209`.

Wakeley, J., Fan, W.T.L., Koch, E.M., Sunyaev, S.R., 2023. Recurrent mutation in the ancestry of a rare variant. Genetics, iyad049. doi:`10.1093/genetics/iyad049`.

Watterson, G.A., 1974. The sampling theory of selectively neutral alleles. Advances in Applied Probability 6, 463–488. doi:`10.2307/1426228`.

Wright, S., 1931. Evolution in Mendelian populations. Genetics 16, 97–159. URL: `https://www.genetics.org/content/16/2/97`.