Implicit motor adaptation patterns in a redundant motor task manipulating a
stick with both hands

Toshiki Kobayashi[1,2] & Daichi Nozaki[1]*

[1]Graduate School of Education, The University of Tokyo, Tokyo 113-0033, Japan

[2]Japan Society for the Promotion of Science, Tokyo 102-0083, Japan

*Corresponding author: Daichi Nozaki, Graduate School of Education, The University
of Tokyo, Tokyo 113-0033, Japan.

Email: nozaki@p.u-tokyo.ac.jp

Author Contributions: Conceptualization, Methodology, Investigation, Formal
Analysis, Writing: TK, DN, Supervision: DN.

Keywords: Reaching movement; Redundancy; Bimanual movement; Motor learning

Abstract

The remarkable ability of the motor system to adapt to novel environments has
traditionally been investigated using kinematically non-redundant tasks, such as
planar reaching movements. Due to this limitation, it remains poorly understood how
the motor system achieves adaptation by altering the movement patterns of our
redundant body. To address this issue, we developed a redundant motor task in
which participants reached for targets with the tip of a virtual stick held with both
hands. Despite the redundancy of the task, participants consistently employed a
stereotypical strategy of flexibly changing the tilt angle of the stick depending on the
direction of tip movement. Thus, this baseline relationship between tip-movement
direction and stick-tilt angle constrained both the physical and visual movement
patterns of the redundant system. Our task allowed us to systematically investigate
how the motor system implicitly changed both the tip-movement direction and the
stick-tilt angle in response to imposed visual perturbations. Both types of
perturbations, whether directly affecting the task (tip-movement direction) or not
(stick-tilt angle around the tip), drove adaptation, and the patterns of implicit
adaptation were guided by the baseline relationship. Consequently, tip-movement
adaptation was associated with changes in stick-tilt angle, and intriguingly, even
seemingly ignorable stick-tilt perturbations significantly influenced tip-movement
adaptation, leading to tip-movement direction errors. These findings provide a new
understanding that the baseline relationship plays a crucial role not only in how the
motor system controls movement of the redundant system, but also in how it implicitly
adapts to modify movement patterns.

Introduction

If a movement error occurs while reaching a target, the motor system automatically and implicitly updates the motor command for the next movement to reduce the error. This remarkable motor learning ability contributes not only to maintaining accurate movements in the presence of noise but also to flexibly adapting the movement to new environments. In the standard planar arm-reaching task, which has been widely used to study the mechanisms of motor learning, participants manipulate a visual cursor with their hands primarily by moving their shoulder and elbow joints (Flash & Hogan, 1985; Gordon et al., 1994) (Fig. 1a). The arm-reaching task has contributed to uncovering many important properties of motor adaptation when a novel environment is introduced (Thoroughman & Shadmehr, 1999; Mazzoni & Krakauer, 2006; Taylor & Ivry, 2011; Takiyama et al., 2015; Hayashi et al., 2016).

Notably, there is no redundancy between hand position and joint angles in this task. The joint angles are uniquely determined by the target to be reached (Fig. 1b). Because of this non-redundancy, when the motor system adapts the reaching movement to the visual rotation imposed on the cursor (Fig. 1c, left), the adapted pattern of joint angles is also uniquely determined (Fig. 1c, right). Naturally, the adapted pattern of joint angles should be identical to the pattern when reaching in the adapted direction (Fig. 1d). However, our body systems for performing practical motor tasks generally have more degrees of freedom than necessary to accomplish the tasks (Bernstein, 1996; Singh et al., 2016). For example, in the golf swing, there are countless patterns of whole-body configurations to achieve identical clubhead positions at ball contact.

The problem arising from such a redundant structure can be illustrated by considering the reaching movement of a hypothetical limb with three joints (Fig. 1e). In this case, there are an infinite number of possible joint angle patterns for the same hand position (Fig. 1f). However, given that the motor system solves task redundancy by minimizing motor cost (Collins, 1995; O'Sullivan et al., 2009) or movement error (White & Diedrichsen, 2010; Zhang et al., 2018), we should observe a stereotypical pattern of joint angles according to the targets (Fig. 1g). The problem is how the motor system changes the direction of cursor movement to implicitly adapt to the visual rotation imposed on the cursor (Fig. 1h). After adaptation, does the motor system move the joint angles in the same way that it voluntarily aims in the direction of the adapted hand movement? (Fig. 1i, left). Alternatively, does the motor system adopt a new joint movement pattern? (Fig. 1i, right). It has not been thoroughly investigated

71   how the motor system coordinates to change the redundant body system to implicitly
72   adapt to the perturbation of the end-effector (i.e., cursor or hand).

Figure 1. Kinematic redundancy and adaptation patterns. (**a**) In the ordinary arm-reaching movement task, there is no redundancy between the task goal (x_c, y_c) and the joint angles (θ_1, θ_2). (**b**) The movement pattern at each target is uniquely determined by the joint angles. (**c**) The adaptation patterns to a visual rotation are also uniquely determined. (**d**) This adapted joint angle pattern should be identical to that of voluntary reaching in the adapted direction. (**e**, **f**) In the case of planar reaching movement by a hypothetical limb with three joints (θ_1, θ_2, θ_3), the cursor can reach the target in infinite patterns. (**g**) Given that the motor system solves the task redundancy by optimization, we should observe a stereotypical pattern of joint angles according to the targets. (**h**, **i**) The question is whether, when a visual rotation is imposed on the cursor (end-effector relevant perturbation), the motor system adopts the same joint movement pattern as when voluntarily aiming in the adapted direction (*left*) or a new joint movement pattern (*right*). (**j**) The joint angles can be perturbed while the cursor position remains unchanged (end-effector irrelevant perturbation). (**k**) This perturbation can be ignored because it does not affect the cursor (*left*). Alternatively, the joint angles may be corrected (*right*).

Redundancy poses another type of problem for implicit movement adaptation.
If the joint angles are perturbed while the hand position remains unchanged (Fig. 1j),
how does the motor system respond to the perturbation? The motor system may
simply ignore the perturbation because it does not affect the cursor position (Fig. 1k,
left). Alternatively, the motor system may attempt to correct the joint movements (Fig.

1k, right). Conventionally, this type of problem has been investigated in the context of
how the motor system responds to the movement variability that affects or does not
affect task performance (i.e., task-relevant or task-irrelevant variability). Previous
studies have consistently shown that the motor system actively corrects for task-
relevant variability while allowing task-irrelevant variability (Mosier et al., 2005;
Latash, 2012; van Beers et al., 2013; Sternad, 2018; Scholtz & Schoner, 1999;
Scholtz et al., 2000). Optimal feedback control theory suggests that the strategies
based on such a "minimal intervention principle" are optimal in terms of achieving
task goals (e.g., movement accuracy) (Todorov & Jordan, 2002; Todorov, 2004; Scott,
2004; Diedrichsen, 2007; Valero-Cuevas et al., 2009; Ronsse et al., 2010; Sternad et
al., 2011; Selgrade & Chang, 2015). These studies predict that the motor system
should ignore the perturbations to joint angles that do not affect the task goal (Fig 1k,
left). However, these studies cannot directly test whether the redundant body
movement pattern is also unchanged when the end-effector movement remains
unchanged.

Conversely, it has also been reported that the motor system adapts to change
the movement in response to the task-irrelevant perturbation (Wolpert et al., 1995;
Flanagan & Rao, 1995; Morehead et al., 2017; Diedrichsen et al., 2010; Schaefer et
al., 2012; Kim et al., 2019). For example, when the cursor trajectory was curved only
in the middle of the reaching movement, the hand movement was adaptively modified
even though the perturbation did not cause a reaching error at the target (Wolpert et
al., 1995; Flanagan & Rao, 1995). These studies created the task-irrelevant condition
by applying the perturbation to the cursor (Wolpert et al., 1995; Flanagan & Rao,
1995; Morehead et al., 2017) or by combining the cursor perturbation with the target
shape manipulation (Diedrichsen et al., 2010; Schaefer et al., 2012; Kim et al., 2019).
Thus, the problem of motor adaptation to such task-irrelevant perturbations in the
previous studies is totally different from the problem of adaptation arising from body
redundancy (Figs. 1i and 1k). In light of this, in this study, we use the terms "end-
effector relevant perturbation" (e.g., Fig. 1h) and "end-effector irrelevant perturbation"
(e.g., Fig. 1j) when appropriate, instead of the more commonly used terms "task-
relevant perturbation" and "task-irrelevant perturbation".

To investigate the problem of how the motor system coordinates to adapt the
redundant body system to these perturbations, a new experimental approach is
required. This study introduces a novel bimanual stick manipulation task specially
designed to address this issue (Fig. 2a). Participants manipulated a virtual stick held
with both hands to reach visual targets by the tip of the stick. This is a redundant task

because the same tip position can be achieved with different stick angles (Fig. 2b).
Importantly, this task allowed us to visually perturb the tip position (i.e., end-effector
relevant perturbation) and the stick angle (i.e., end-effector irrelevant perturbation)
either separately or simultaneously. Using this approach, we can directly investigate
how the motor system changes the redundant body system when adapting to these
two types of perturbations.

Results

Movement strategy to manipulate the stick
A total of 72 healthy participants took part in the study (Experiments 1–3). They were
asked to manipulate a virtual stick with both hands (length = 40 cm; distance between
hands = 15 cm) to move the right tip of the stick towards the targets (Fig. 2a). This
task is redundant because the same target can be reached in countless movement
patterns. The participants could move both hands in parallel without tilting the stick
(Fig. 2b, top). Alternatively, they could perform the task by tilting the stick (Fig. 2b,
bottom). During the first part of the task (baseline phase: 360 trials) in all the
experiments (Experiments 1–3), the participants moved the tip from a starting point to
a target (movement distance = 10 cm) that appeared in one of nine directions (0°:
horizontal direction, ±10°, ±20°, ±30°, and ±40°) (Fig. 2c).

The trajectories of the tip and both hands appeared to be almost straight, but
both hands were not moved in parallel, except when aiming at the 0° target (Fig. 2d,
top). The velocity profiles of the hands and tip were bell-shaped (Fig. 2d, bottom),
similar to a normal unimanual reaching movement (Flash & Hogan, 1985; Gordon et
al., 1994). The time-course changes in the stick-tilt angle relative to the initial
horizontal tilt angle showed that the participants moved the tip while slightly tilting the
stick, except when aiming at the 0° target (Fig. 2e). This tendency was observed from
the beginning of the experiment (t-test compared with 0°, $|t(71)| > 6.4$, Bonferroni
corrected $P < 0.001$; Fig. 2f).

Figure 2. Bimanual stick manipulation task. (**a**) Participants held handles of a manipulandum with both hands and moved the right tip of a virtual stick on the monitor. (**b**) In this task, there are countless movement patterns to move the tip to the target. Two representative strategies are shown: parallel translation (*top*) and tilting the stick (*bottom*). (**c**) During the baseline phase, participants were instructed to move the tip from a starting point to a target that appeared in one of nine directions. (**d**) Movement trajectories (*top*) and velocity profiles (*bottom*) of the left hand, right hand, and tip averaged across participants. Each color corresponds to the target direction shown in **c**. The shaded areas represent the SEM across participants. (**e**) The time-course of the stick-tilt angle shows that participants moved the tip while tilting the stick. The shaded areas indicate the SEM. (**f**) Trial-to-trial changes in stick-tilt angle at the peak velocity of the tip. The strategy of tilting the stick was observed from the beginning of the experiment (*t*-test compared with zero, |*t*(71)| > 6.4, Bonferroni corrected *P* < 0.001). The shaded areas indicate the SEM.

Movement strategies to manipulate the stick for individuals are illustrated on
the plane expressed by two variables: tip-movement direction relative to a horizontal
direction (TMD) and the stick-tilt angle relative to the initial horizontal state (STA) (Fig.
3a). Although each participant could have performed the task differently (Fig. 2b), all
participants showed stereotypical movement patterns: The stick-tilt angle was
significantly modulated by the tip-movement direction (*t*-test compared with 0°, |*t*(71)|
> 5.5, Bonferroni corrected *P* < 0.01; Fig. 3a, inset). This monotonically increasing
curve, which will be referred to as the "baseline TMD–STA relationship", reflects a
constraint between the direction of tip movement and the angle of stick tilt angle. The
relationship showed that clockwise (CW) tip movement direction was accompanied
by CW stick-tilt, and counterclockwise (CCW) tip-movement direction was
accompanied by CCW stick-tilt. This constraint of movement patterns was likely
created by minimizing the movement distance of the hands. If both hands are moved

in parallel, the total distance between the initial and final positions of the hands
should be 20 cm (10 cm for each) regardless of the targets; however, the
experimentally observed distance was reduced by tilting the stick (Fig. 3b). The
average distance was a near-minimal value (versus parallel translation: Cohen's d =
1.914, versus theoretical minimum: Cohen's d = 0.334; Fig. 3c).

Figure 3. Relationship between the tip-movement direction and stick-tilt angle.
(**a**) A relationship between each individual's tip-movement direction and stick-tilt angle, when the tip speed peaked is represented as the thin gray lines drawn with colored plots. The averaged relationship is plotted as a bold gray line (baseline TMD–STA relationship) with the SD (crosses) and 95% confidence intervals (ellipses). The inset shows the stick-tilt angles at peak velocity and movement offset (error bars: SD, *: *t*-test compared with 0°, Bonferroni-corrected *P* < 0.01). The monotonically increasing curve indicates the stereotypical strategy of manipulating the stick (diagrams on the right side). (**b**) Sum of the distances between the initial and final positions of both hands (i.e., the sum of the lengths of pink arrows in Fig. 2b). The dotted and dashed gray lines indicate the distances when both hands are moved in parallel and the mathematically derived minimum distances by tilting the stick, respectively. The error bars denote the SD. (**c**) The distance averaged across target directions was closer to the minimum value (Cohen's d = 1.914) than the parallel translation distance (Cohen's d = 0.334). The error bar indicates the SD across participants.

Experiment 1: Motor adaptation pattern to the end-effector relevant perturbation

During the second part of the experiment (adaptation phase: 240 trials), the target appeared only in the 0° direction, and three types of visual perturbations were introduced. Most importantly, this task enabled the introduction of end-effector relevant perturbation (Experiment 1), or end-effector irrelevant perturbation (Experiment 2), or a combination of both types of perturbations (Experiment 3). In Experiment 1A (N = 19), the tip-movement direction was visually rotated in the CCW direction around the starting position (tip perturbation), whereas the stick-tilt angle remained unchanged (Fig. 4a). The amount of tip directional rotation was gradually increased by 1° per a trial up to 30° so that the participants did not notice the presence of perturbation.

Figure 4b illustrates how this visual perturbation changes the states of stick movement in visual and physical spaces. The movement in the visual space was the one visible to the participants, whereas the movement in the physical space was the actual hand movement, which was invisible to them. On the visual TMD–STA plane (Fig. 4b, left), the visual perturbation to the tip can be expressed as a shift of the state along the TMD axis. The error of the tip-movement direction in the visual space needs to be compensated by changing the movement of both hands. On the physical TMD–STA plane (Fig. 4b, right), this requirement can be expressed as a shift of the task goal in the direction opposite to the visual perturbation.

Owing to the redundant nature of this task, participants could adapt to change the visual tip movement using different strategies. For example, they could change the tip movement without changing the stick-tilt while keeping the strategy of moving both hands in parallel (Fig. 4c, top). Alternatively, they might adapt by changing the stick tilt as if they were aiming at a -30° target (Fig. 4c, bottom). These strategies can be expressed as a state shift towards the new task goal line along the TMD-axis or along the baseline TMD–STA relationship in the physical TMD–STA plane (Fig. 4d).

Figure 4. Experiment 1: End-effector relevant visual perturbations. (**a**) During the adaptation phase of Experiment 1, the tip-movement direction was rotated by 30° around the starting point (tip perturbation). The dotted and solid black arrows indicate the intended and actual tip trajectory, respectively. (**b**) The changes in the movement pattern are expressed as the state shift on the TMD–STA plane (*left*: visual space, *right*: physical space). The dashed red lines denote the task goal (i.e., a target direction). On the visual TMD–STA plane, the perturbation (a blue arrow) shifts the movement pattern. On the physical TMD–STA plane, the task goal is shifted. (**c**) The participants could correct the tip error without (pattern i, *top*) or by (pattern ii, *bottom*) altering the stick-tilt angle. (**d**) The adaptive strategies are expressed as the shift on the visual TMD–STA plane (*left*) and the physical TMD–STA plane (*right*). The visual error caused by the perturbation could be compensated by different adaptation patterns (blue arrows): Representative patterns are expressed as bold arrows (patterns i and ii).

The experimental results demonstrated that the participants adapted to the tip
perturbation by tilting the stick. On the visual TMD–STA plane (Fig. 5a), after the
adaptation, the tip-movement direction remained at the baseline level (paired *t*-test,
t(18) = -1.320, *P* = 0.203), while the stick was tilted significantly more in the CW
direction than when aiming at the 0° target in the baseline phase (paired *t*-test, *t*(18) =
10.003, *P* < 0.01). Thus, the tip perturbation was compensated by adaptation, but the
adaptation produced an abnormal visual stick-tilt. This result is likely to support the
idea of the minimal intervention principle because participants ignored this visual
stick-tilt which did not directly influence the task.

On the physical TMD–STA plane, adaptation appeared to progress along the
baseline TMD–STA relationship throughout the adaptation phase (Figs. 5b and 5c).
The implicitly adapted tip-movement direction and stick-tilt angle were similar to those
when voluntarily aiming at the -30° target in the baseline phase. Although the
difference was small but significant for the tip-movement direction (paired t-test, $t(18)$
= -10.053, $P < 0.01$), there was no significant difference in the stick-tilt angle (paired
t-test, $t(18)$ = -1.081, P = 0.294). Thus, the motor system implicitly achieved
adaptation to the tip perturbation as if it had voluntarily aimed at the -30° target (i.e.,
pattern ii in Figs. 4c and 4d) (Fig. 5d). This adaptation characteristic was also
supported by the observation that the participants who tilted their stick more when
aiming at the -30° target in the baseline phase exhibited a greater stick- tilt in the
adapted movement patterns (R = 0.601, P = 0.007; Fig. 5e). In Experiment 1B (N =
13), we imposed the 30° tip perturbation abruptly to allow participants to use explicit
strategy, and obtained nearly identical results (Fig. S1).

As described above, the adapted tip-movement direction on the physical
TMD–STA plane was significantly different from the tip-movement direction when
aiming at the -30° target in the baseline phase (Fig. 5b). In contrast, such significant
difference was not observed for Experiment 1B (Fig. S1). Why was such a difference
only in Experiment 1A? We speculated that this difference was caused by the implicit
adaptation to correct the abnormal visual stick-tilt angle. As shown in Fig. 5a, after
the adaptation, the stick was visually tilted in the CW direction. If the physical
movement pattern is constrained by the baseline TMD–STA relationship, the CCW
correction of the stick-tilt angle should be accompanied by the CCW change in tip-
movement, which would interfere with the adaptation to the tip perturbation.

Figure 5. Experiment 1: Adaptation patterns to gradually imposed end-effector relevant perturbations. (**a**) In Experiment1A, the adaptation trajectory (a thin blue line) on the visual TMD–STA plane gradually deviated from a baseline pattern (a gray circle) mainly along the STA-axis. The tip-movement direction remained at the baseline level (paired *t*-test, *t*(18) = -1.320, *P* = 0.203). In contrast, the stick-tilt angle after the adaptation (a blue circle) significantly differed from that in the baseline pattern when aiming at the 0° target (a gray circle) (paired *t*-tests, *t*(18) = 10.003, *P* < 0.001), indicating that the participants saw the abnormal visual stick-tilt. The plots indicate mean ± SD across participants. (**b**) On the physical TMD–STA plane, the adaptation trajectory followed the baseline TMD–STA relationship. Although the difference was small but significant for the tip-movement direction compared with the baseline pattern when aiming at the -30° target (paired *t*-test, *t*(18) = -10.053, *P* < 0.01), there was no significant difference in the stick-tilt angle (paired *t*-test, *t*(18) = -1.081, *P* = 0.294). (**c**) The area indicated by the squared outline in **b** is enlarged. The adaptive process can be divided into nine trial segments (trials 1–5, 6–10, 11–15, 16–20, 21–25, 26–30, 30–120, 121–210, and 211–240). (**d**) The motor system implicitly achieved adaptation to the tip perturbation as if it had voluntarily aimed at the -30° target. (**e**) Participants who showed a larger stick-tilt when aiming at the -30° target during the baseline phase also exhibited a larger stick-tilt in the adapted pattern (*R* = 0.599, *P* < 0.001).

Supplemental Figure 1. Adaptation patterns to abruptly imposed end-effector relevant perturbations. (**a**) In Experiment 1B, a 30° tip perturbation was applied from the first trial of the adaptation phase. On the visual TMD–STA plane, the stick-tilt angle after the adaptation (a blue circle) significantly differed from that in the baseline pattern when aiming at the 0° target (a gray circle) (paired *t*-tests, *t*(12) = 7.529, *P* < 0.001), whereas the tip-movement direction remained at the baseline level (paired *t*-test, *t*(12) = 0.430, *P* = 0.675). The formats are the same as Fig. 5a. (**b**) On the physical TMD–STA plane, neither tip-movement direction nor stick-tilt angle in the adapted pattern was significantly different from that in the baseline pattern when aiming at the -30° target (paired *t*-test, TMD: *t*(12) = -1.628, *P* = 0.129, STA: *t*(12) = -0.490, *P* = 0.633).

Experiment 2: Motor adaptation pattern to end-effector irrelevant perturbation

Thus, it is still possible that the motor system does not necessarily ignore, but

corrects for, the stick-tilt error that does not directly influence the tip position.

Experiment 2 (*N* = 20) was designed to directly test this possibility by imposing a

stick-tilt perturbation around the right tip (i.e., end-effector irrelevant perturbation) (Fig.

6a). As the tip moved from the starting point, the stick started to tilt around the tip up

to ±6° when the tip reached the target (the tilt amount was proportional to the tip's

distance from the starting point). The ±6° tilt perturbation was chosen according to

the stick tilt at the movement offset when reaching the ±30° target for the baseline

phase (see Fig. 3a). Ten participants participated in the CCW stick-tilt perturbation

condition (E2CCW group: 6°), and the remaining ten participants took part in the CW

stick-tilt perturbation condition (E2CW group: -6°). Because the amount was small,

the participants did not notice the presence of perturbation throughout the experiment.

Since this perturbation changed only the stick-tilt angle, it is expressed as a state

shift along the STA-axis on the visual TMD–STA plane (Fig. 6b). According to the

minimal intervention principle, the motor system does not need to change anything

because the stick-tilt perturbation does not affect the tip position (i.e., the task goal)

(Figs. 6c and 6d, pattern i). However, if the motor system is actively compensating
for the stick-tilt perturbation, it should change the stick-tilt angle (Figs. 6c and 6d,
pattern ii).

On the visual TMD–STA plane, the stick-tilt angles remained different from

Figure 6. Experiment 2: End-effector irrelevant visual perturbations. (**a**) During
the adaptation phase of Experiment 2, the stick-tilt angle was rotated around the tip.
The perturbation of ±6° was imposed from the first trial. Participants received either
the counterclockwise perturbation (green, E2CCW group) or the clockwise
perturbation (orange, E2CW group). (**b**) On the visual TMD–STA plane, the
perturbation is expressed as a vertical shift of the state along the task goal (dashed
red line). On the physical TMD–STA plane, the perturbation does not influence the
task goal. Note that, in both planes, the states are on the task goal lines. The formats
are the same as in Fig. 4b. (**c**) Participants may not change the movement pattern
because the perturbation does not affect the tip position (pattern i, *top*). Alternatively,
the perturbation can lead to a stick-tilt correction (pattern ii, *bottom*). (**d**) These
strategies are expressed as a shift on the TMD–STA plane. Note that pattern i means
that there is no change in the movement pattern.

those in the baseline phase (Fig. 7a) (two-way (direction of stick-tilt perturbation (i.e.,
groups), trial phase (first 30 trials or last 30 trials)) repeated-measures ANOVA, a
significant main effect of groups: $F(1,18) = 54.457$, $P < 0.001$; no effect of trial phase:
$F(1,18) = 2.554$, $P = 0.127$; no interaction: $F(1,18) = 1.613$, $P = 0.220$). Thus, the
adaptive changes did not fully compensate for the perturbation-induced stick-tilt
errors. However, the data on the physical TMD–STA plane clearly demonstrated that

participants implicitly changed the stick-tilt angle in the direction opposite to that of

the perturbation from the early adaptation phase (Fig. 7b): Without the correction, the

data of the E2CCW and E2CW groups should have largely overlapped. However,

with respect to stick-tilt angle, two-way repeated-measures ANOVAs revealed the

main effect of stick-tilt perturbation direction ($F(1, 18) = 12.483$, $P < 0.01$) but not the

main effect of trial phase ($F(1, 18) = 3.331$, $P = 0.085$). The interaction between these

factors was not significant ($F(1, 18) = 1.362$, $P = 0.258$).

Unexpectedly, this stick-tilt correction was accompanied by undesirable tip-

movement directional errors (Fig. 7c). With respect to the tip-movement direction,

two-way repeated-measures ANOVAs revealed the main effect of the direction of

stick-tilt perturbation ($F(1, 18) = 5.104$, $P = 0.037$) but not the main effect of trial

phase ($F(1, 18) = 1.288$, $P = 0.271$) (the interaction between these factors was not

significant: $F(1, 18) = 0.804$, $P = 0.382$). Notably, these seemingly irrational adaptation

patterns were consistent with the patterns predicted from the baseline physical TMD–

STA relationship: CCW (CW) stick-tilt adaptation to the CW (CCW) stick-tilt

perturbation slightly shifted the tip-movement direction in the CCW (CW) direction

(Fig. 7d).

We also examined how the amount of adaptation depended on the amount of

stick-tilt when voluntarily aiming at the ±30° target in the baseline phase. The 6° stick-

tilt perturbation should be insignificant for participants who tilt the stick at a larger

angle; thus, this perturbation tends to be ignored. Conversely, for participants who did

not tilt the stick, a 6° stick-tilt perturbation should be significant and must be corrected.

Figures 7e and 7f show how the amount of stick-tilt correction (Fig. 7e) and tip-

movement directional change (Fig. 7f) depended on the amount of stick-tilt angle

when aim at the ±30° target in the baseline phase. As predicted, we observed a

negative correlation (stick-tilt correction: $R = -0.572$, $P < 0.01$; tip-movement

directional change: $R = -0.440$, $P = 0.052$), although the tip-movement direction

change was not significant.

Taken together, the results of Experiment 1 and Experiment 2 suggest that

the implicit adaptation in the redundant stick-manipulating task is achieved by the

following two mechanisms: First, the motor system attempts to correct for any

perturbation in visual feedback, regardless of whether it is end-effector relevant or

end-effector irrelevant. Second, the physical pattern of movement correction is

constrained by the baseline TMD–STA relationship.

Figure 7. Experiment 2: Adaptation patterns to end-effector irrelevant visual perturbations. (**a**) Movement patterns are shown by green (E2CCW group) and orange (E2CW group) plots on the visual TMD–STA plane. The open and filled plots represent the movement patterns in the first and last 30 trials, respectively, during the adaptation phase (mean ± SD, across participants). (**b**) On the physical TMD–STA plane, the separated movement patterns between both groups showed that participants corrected the end-effector irrelevant stick-tilt errors. (**c**) The enlarged plot of the squared area in **b** shows that end-effector relevant tip-movement directional errors accompanied the adaptation. These effects persisted until the end of the adaptation phase (two-way repeated-measures ANOVAs). (**d**) Schematics of how the stick-tilt perturbation altered the movement patterns. (**e**, **f**) How participants tilted the stick during the baseline phase was negatively correlated with the stick-tilt angle and the tip-movement direction. To control for the individual variability, the baseline stick-tilt angle or tip-movement direction when aiming at the 0° target was subtracted from each value.

Experiment 3: Interaction between end-effector relevant and end-effector irrelevant perturbations

Experiment 3 (N = 20) was designed to further investigate the interaction between adaptations to end-effector relevant and end-effector irrelevant perturbations by applying both simultaneously (Fig. 8a). We hypothesized that adaptation to the tip perturbation would be influenced by the direction of the simultaneously applied stick-tilt perturbation. Specifically, the CCW tip perturbation was combined with the CCW stick-tilt perturbation (E3CCW group: N = 10) or with the CW stick-tilt perturbation (E3CW group: N = 10). These visual perturbation patterns can be expressed as state shifts in the visual TMD–STA plane (green and orange arrows for E3CCW and E3CW groups, respectively, in Fig. 8b). On the physical TMD–STA plane, the correction pattern required for the E3CCW group (Fig. 8c, top) is aligned with the baseline TMD–STA relationship (green arrow in Fig. 8d), from which we can predict that adaptation should proceed smoothly. In contrast, the correction pattern required for the E3CW group (Fig. 8c, bottom) is orthogonal to the baseline TMD–STA relationship (orange arrow in Fig. 8d), which may prevent smooth adaptation. We tested these predictions by simultaneously imposing the tip perturbation which was the same as the perturbation used in Experiment 1A and the stick-tilt perturbation which was the same as the perturbation used in Experiment 2. Again, the participants were unaware of the perturbations throughout the experiment.

Figure 8. Experiment 3: Interaction between end-effector relevant and end-effector irrelevant visual perturbations. (**a**) During the adaptation phase of Experiment 3, both the CCW tip perturbation and the CCW or CW stick-tilt perturbation were introduced. The participants were assigned to the CCW stick-tilt rotation condition (green, E3CCW group) or the CW stick-tilt rotation condition (orange, E3CW group). (**b**) These visual perturbations can be expressed as state shifts on the visual TMD–STA plane (green and orange arrows for E3CCW and E3CW groups, respectively). (**c**) The adaptation pattern to reduce visual errors (*top*: E3CCW group, *bottom*: E3CW group). (**d**) On the physical TMD–STA plane, the correction pattern required for the E3CCW group is aligned with the baseline TMD–STA relationship (green arrow), whereas the correction pattern required for the E3CW group is orthogonal to the baseline TMD–STA relationship (orange arrow).

Contrary to the predictions, the difference in stick-tilt perturbation directions
did not affect the final adapted movement patterns: As shown in the visual TMD–STA
plane (Fig. 9a), the difference in stick-tilt angle remained unchanged (paired *t*-test,
$t(18) = 6.920$, $P < 0.001$). Similarly, in the physical TMD–STA plane (Fig. 9b), there
were no significant differences in movement pattern between the two groups (tip-
movement direction: paired *t*-test, $t(18) = -0.479$, $P = 0.638$, stick-tilt angle: paired *t*-
test, $t(18) = -0.386$, $P = 0.704$). However, a trial-by-trial variability of the movement
pattern in the latter adaptation phase (trial# 121–240) showed a slight difference
between groups (*t*-test, tip-movement direction: $t(18) = -2.046$, $P = 0.056$, stick-tilt
angle, $t(18) = -3.235$, $P = 0.005$), suggesting that the adaptation processes were
implicitly affected by combination of perturbations (Fig. 9a, inset). In partial agreement

with our predictions, two groups showed different adaptation processes: trial-
dependent trajectories to the final adapted pattern were clearly separated for both
groups (Fig. 9c). Indeed, a repeated-measures multivariate analysis of variance
(MANOVA) showed a significant effect of group (Pillai's Trace $F(8, 162) = 0.155$, $P <$
0.001) and trial segment (Pillai's Trace $F(8, 162) = 0.719$, $P < 0.001$); however, no
significant interaction was observed (Pillai's Trace $F(8, 162) = 0.039$, $P = 0.982$). As a
result, the adaptation of the tip movement was more delayed in the E3CW group (trial
segment 2: t-test, $t(18) = -3.684$, Bonferroni-corrected $P < 0.05$; Fig. 9d).

Figure 9. Experiment 3: Adaptation patterns to end-effector relevant and end-effector irrelevant visual perturbations. (**a**) Adaptation patterns on the visual TMD–STA plane are shown (same formats as Fig. 5a). A green (orange) plot represents the adapted movement pattern in the E3CCW group (E3CW group). The tip-movement direction did not significantly differ (*t*-test, *t*(18) = -0.473, *P* = 0.642), while the stick-tilt angle remained a significant difference between groups (*t*-test, *t*(18) = 6.920, *P* < 0.001). A trial-by-trial variability of the movement pattern in the latter adaptation phase (trial# 121–240) showed a difference between groups (*t*-test, tip-movement direction: *t*(18) = -2.046, *P* = 0.056, stick-tilt angle, *t*(18) = -3.235, *P* = 0.005) (**b**) Similarly, the adapted movement patterns on the physical TMD–STA plane did not show a significant difference (*t*-test, tip-movement direction: *t*(18) = -0.479, *P* = 0.638, stick-tilt angle: *t*(18) = -0.386, *P* = 0.704). (**c**) However, the adaptive processes appeared to be different. The movement patterns of the two groups are compared using a repeated-measures MANOVA. A multivariate analysis revealed a significant effect of group (Pillai's trace *F*(8, 162) = 0.155, *P* < 0.001) and trial segment (Pillai's trace *F*(8, 162) = 0.719, *P* < 0.001); no significant interaction can be observed (Pillai's trace *F*(8, 162) = 0.039, *P* = 0.982). The definition of trial segments is the same as that shown in Fig. 5c. (**d**) The learning curves for the tip-movement. The adaptation in the E3CW group was always delayed but showed a significant difference only during the early phase (trial segment 2: *t*-test, *t*(18) = -3.684, Bonferroni-corrected *P* < 0.05).

Discussion

Our bodies have more degrees of freedom than necessary to achieve a relatively low-dimensional task goal. Understanding how the motor system resolves this redundancy remains a challenging problem (Bernstein et al., 1996). In this study, we investigated the problems of motor control and adaptation specific to kinematically redundant systems. The first problem is how the motor system coordinates the redundant body movements to control the end-effector (Fig. 1f). The second problem is how the motor system coordinates to modify redundant body movements when implicitly adapting to the perturbation imposed on the end-effector (Fig. 1i). The third problem is how the motor system responds to the perturbation to the redundant dimension that does not directly affect the end-effector (Fig. 1k). These two types of perturbations have been conventionally referred to as "task-relevant" and "task-irrelevant" perturbations. In previous studies, task-irrelevant perturbations were created by applying the perturbation to the end-effector (e.g., cursor) only at the midpoint of the movement (Wolpert et al., 1995; Flanagan & Rao, 1995) or by using a wide target (e.g., arc) (Schaefer et al., 2012; Kim et al., 2019). Because the task-irrelevant conditions realized in these studies are completely different from the scope of our study, we specifically used the terms "end-effector relevant perturbation" and "end-effector irrelevant perturbation".

To investigate these issues, we designed a novel stick manipulation reaching task. This redundant motor task allows the investigation of adaptation patterns in the redundant system following the introduction of end-effector relevant, end-effector irrelevant, or both types of perturbations. First, we showed that participants adopted a stereotyped strategy to reach the target with the stick tip, and that the movement pattern (stick tip movement and stick tilt angle) was constrained by a baseline TMD–STA relationship. Second, when only end-effector relevant perturbations were imposed on the tip, the motor system attempted to modify the physical movement patterns along the baseline TMD–STA relationship to achieve the task goal (Experiment 1). Third, even when perturbations were applied only to the end-effector irrelevant dimension (i.e., the stick-tilt angle), the motor system still adapted to the end-effector irrelevant perturbation and this adaptation was accompanied by an end-effector relevant error (Experiment 2). Finally, when both types of perturbations were imposed simultaneously, the adaptation processes were influenced depending on the combination of perturbations (Experiment 3).

Kinematic relationship between end-effector relevant and end-effector irrelevant dimensions

In our stick manipulation task, participants were asked to reach the tip of the stick to the targets. Participants can achieve the reaching task in different ways. For example, they can use a simple strategy by moving both hands in the same direction (Fig. 2b, top) or a more complicated strategy by tilting the stick (Fig. 2b, bottom). Nevertheless, they showed stereotypical patterns of moving the stick tip by tilting the stick from the beginning of the task (Fig. 2f): Reaching the CW (or CCW) targets was accompanied by the CW (or CCW) stick tilt (Fig. 3a). Notably, compared with the case where both hands moved in parallel, the movement distances of both hands to perform the task were significantly reduced to a near-optimal level (Figs. 3b and 3c), which contributed to the reduction of work required (Collins, 1995; O'Sullivan et al., 2009). As a result of this optimization process, tip-movement and stick-tilt were uniquely associated with each other by the baseline TMD–STA relationship (Fig. 3a). Physically, this relationship constrains how the motor system should move the stick. Visually, it tells the motor system how the stick movement should be seen.

The strength of the task to investigate the motor adaptation in redundant system

Our task has several advantages. First, the task has created redundancy in a natural way. Manipulating a stick with both hands is relatively common in everyday life (using a broom, golf or bat swing etc.). This familiarity with the task was reflected in Fig. 2f which shows that all participants started tilting the stick to reach the tip toward the targets from the beginning of the experiment. This characteristic is ideal for investigating how this inherent movement pattern influences motor adaptation. Second, this task allows us to investigate how the perturbation to the tip-movement direction leads to a change in both the tip-movement direction itself and the stick-tilt angle. It has not been investigated how the motor system implicitly achieves adaptation to the end-effector perturbation (i.e., the end-effector relevant perturbation) by changing the movement patterns of the redundant body system. Third, this task can create a novel condition in which the perturbations are imposed on the redundant system while the tip position remains unchanged (i.e., the end-effector irrelevant perturbation). No previous studies have investigated how the motor system implicitly responds to such a perturbation by altering (or not altering) the movement of the redundant body system.

Adaptation to end-effector relevant perturbation

When the end-effector relevant perturbation was implicitly imposed to the tip-
movement direction (visual rotation by 30° CCW), the participants adapted to change
the tip-movement direction in the opposite CW direction (Experiment 1), as reported
in many previous studies using an ordinary reaching movement (Cunningham, 1989;
Krakauer et al., 2000). Importantly, our task helped to investigate how such motor
adaptations in the end-effector relevant dimension (tip-movement direction) are
accompanied by the movement pattern changes in the end-effector irrelevant
dimension (stick-tilt angle). Participants implicitly corrected the tip-movement
direction by changing the stick tilt as if they were voluntarily aiming at the -30° target.
This adaptation can be visualized as a state shift along the baseline TMD–STA
relationship in the physical TMD–STA plane (Fig. 5b). The baseline TMD–STA
relationship originally contributes to dictate how the motor system voluntarily moves the
stick. It is likely that the motor system also uses this relationship as a scaffold to
implicitly guide how to correct the tip-movement direction. Hayashi et al. (2016) have
demonstrated that the adaptation of the standard reaching movement to a visual
perturbation was significantly influenced by the baseline relationship between
the movement and target directions (i.e., a visuomotor map). After distorting the
visuomotor map so that the movement direction was more (or less) sensitively
modulated by the target direction, the movement adaptation was enhanced (or
reduced). The present finding that the baseline movement pattern is used for the
implicit adaptation is consistent with this previous finding.

Notably, this adaptation pattern changed the visual feedback of the stick tilt.
When reaching the 0° target in the baseline condition, the stick-tilt angle was almost
0° (Fig. 3a). However, after adaptation, the amount of stick-tilt was equivalent to that
when aiming at the -30° target (Fig. 5b). Thus, the visually observed stick-tilt after
adaptation was different from that observed in the baseline condition (Fig. 5a).
Nevertheless, the adapted physical movement pattern persisted to follow the
baseline TMD–STA relationship, suggesting that the motor system mainly paid
attention to end-effector relevant tip-movement and ignored end-effector irrelevant
stick-tilt angle. This adaptation pattern seems to be consistent with the minimal
intervention principle.

However, a slight difference in the adaptation patterns between Experiment
1A and Experiment 1B implies a violation of the minimal intervention principle. In
Experiment 1A, in which a 30° visual rotation was implicitly introduced by gradually
increasing its amplitude, the final adapted tip-movement direction was slightly but
significantly different from that when voluntarily aiming at the adapted direction (i.e., -

30° target) (Fig. 5b). In contrast, no such residual difference was not observed in

Experiment 1B, where participants could use an explicit aiming strategy (Fig. S1b).

Visuomotor adaptation is often incomplete even after sufficient training, and residual

errors are observed (Krakauer et al., 2000; Ethier et al., 2008; Malone et al., 2011;

Vaswani et al., 2015). Although the presence of residual errors has been attributed to

the imbalance between learning and forgetting, a recent study demonstrated that only

the implicit learning system is responsible for producing residual errors by modulating

error sensitivity with the consistency of errors in previous trials (Albert et al., 2021).

We speculate that an alternative mechanism is involved in the generation of a

residual difference in Experiment 1A. If the motor system does not ignore the

abnormal stick-tilt but attempts to correct it, and this correction pattern is physically

constrained by the baseline TMD–STA relationship, the stick-tilt correction should

accompany the de-adaptation of the tip-movement direction and lead to the

generation of a residual error. As discussed below, the results of Experiment 2 and

Experiment 3 support this idea.

Adaptation to end-effector irrelevant perturbation

Our task can create a novel end-effector irrelevant stick-tilt perturbation and allows us

to study how the motor system responds to it. According to the minimal intervention

principle, the motor system should ignore stick-tilt perturbations because they do not

influence task goal achievement (i.e., so called "task-irrelevant" perturbation).

However, inconsistent with this idea, Experiment 2 clearly demonstrated that the

motor system responded to the stick-tilt perturbations (Figs. 7b and 7c). This result

indicates that the motor system predicts the visual feedback of an end-effector

irrelevant stick-tilt angle when performing the task and tries to eliminate the

dissociation between the observed and predicted stick-tilt angle (i.e., sensory

prediction error). This idea was corroborated by the fact that participants who

demonstrated a shallower TMD–STA relationship during the baseline phase showed

a greater amount of stick-tilt correction (Fig. 7e). The ±6° stick-tilt perturbation should

cause the larger prediction error for these participants.

One might consider that previous studies have already shown that the motor

system responds to the task-irrelevant perturbations (Wolpert et al., 1995; Flanagan

& Rao, 1995; Diedrichsen et al., 2010; Schaefer et al., 2012; Kim et al., 2019).

However, the task-irrelevant conditions implemented in these previous studies are

quite different from the end-effector irrelevant condition in our study. These studies

created the task-irrelevant condition by combining the perturbation of visual cursors
with the manipulation of movement trajectory (Wolpert et al., 1995; Flanagan & Rao,
1995; Morehead et al., 2017) or target/cursor shape (Diedrichsen et al., 2010;
Schaefer et al., 2012; Kim et al., 2019). Therefore, these perturbations are not "end-
effector irrelevant". Exceptionally, in a previous study, an end-effector irrelevant
perturbation was implemented in a bimanual postural task to investigate postural
reflex activity (Omrani et al., 2013). However, to our knowledge, such an end-effector
irrelevant perturbation has never been used to investigate the motor adaptation
pattern in a redundant system. Furthermore, as discussed in the next section, our
task design revealed a novel aspect of motor adaptation to the end-effector irrelevant
perturbation.

Adaptation to end-effector irrelevant perturbation influences end-effector
relevant performance
Unexpectedly, Experiment 2 shows that adaptation to the end-effector irrelevant
stick-tilt perturbation was accompanied by an increase in the end-effector relevant
tip-movement direction error (Fig. 7c). This counterintuitive result can be explained by
the physical constraint of the adaptation pattern along the baseline TMD–STA
relationship (Fig. 3a). According to this constraint, CCW (or CW) stick-tilt adaptation
should change the tip-movement direction in the CCW (or CW) direction (Fig. 7d).
The influence of stick-tilt perturbation on tip-movement direction error tended to be
more pronounced for participants with a smaller stick-tilt angle at baseline, i.e., those
with a shallower baseline TMD–STA relationship (Fig. 7f), further supporting this idea.
The results of Experiment 3 also support the existence of such an interaction
between tip-movement and stick-tilt adaptation. When the tip-movement and stick-tilt
perturbations were imposed simultaneously, the adaptation trajectories on the
physical TMD–STA plane were significantly influenced by the perturbation
combination patterns (Fig. 9c). If the motor system had simply attempted to correct
the stick-tilt to reduce the sensory prediction error, this spillover effect on the tip-
movement direction would not have been observed.

One remaining question is why the motor system needs to correct for end-
effector irrelevant stick-tilt errors in the first place? We speculate that there may be
two reasons. First, a possible simple explanation is that the motor system predicts all
sensory consequences generated by the action and attempts to correct any
differences between the predicted and actual sensory information, even when the

task is successfully achieved (Mazzoni & Krakauer, 2006; Schaefer et al., 2012;
Morehead et al., 2017; Kim et al., 2019). Second, given that tip-movement direction
and stick-tilt angle are tightly coupled (i.e., the presence of baseline TMD–STA
relationship), the presence of an end-effector irrelevant stick-tilt visual error implies
the presence of an end-effector relevant tip-movement direction error. In other words,
the motor system interprets end-effector irrelevant errors as evidence that end-
effector relevant errors have been also produced. When sensory information contains
noise and/or uncertainty, the reliability of sensory information estimation can be
improved by utilizing additional sensory information (Ernst & Banks, 2002). From this
viewpoint, utilizing stick-tilt information may be useful for improving the accuracy of
the estimation of the tip-movement direction.

Figure 10 illustrates the possible mechanism by which the motor system alters the
movement pattern of the redundant system when encountering visual errors. The
motor system develops a stereotyped movement pattern through an optimization
process that is expressed by a baseline TMD–STA relationship (gray lines in Figs.
10a and 10b). This relationship not only dictates how the motor system physically
controls the redundant system, but also informs the resulting visual feedback. When
visual perturbations are applied to the tip-movement direction or the stick-tilt angle
(bold arrows in Fig. 10a), the motor system tries to eliminate the dissociation of the
perturbed state from the original state (thin arrows in Fig. 10a). It even responds to
the end-effector irrelevant perturbation (red and blue arrows in Fig. 10a), because the
error in the end-effector irrelevant dimension implies the presence of an end-effector
relevant error (black dashed arrows in Fig. 10a) under the condition that the stick-tilt
angle is tightly coupled with the tip-movement direction (i.e., the baseline TMD–STA
relationship). However, these correction patterns are not physically allowed. Since
the tip-movement direction and the stick-tilt angle are physically constrained by the
baseline TMD–STA relationship, these correction patterns are biased toward the
relationship (bold arrows in Fig. 10b).

In summary, this study shows that the baseline TMD–STA relationship, originally
developed to control the redundant system, also plays a crucial role for the motor
system in how it should implicitly respond to the visual errors and how it should
correct the physical movement. This idea provides a novel framework for
understanding how the motor system leans to control the redundant system when
encountering visual errors.

Figure 10. A hypothesis for how the motor system alters the movement pattern of redundant system. The motor system develops a stereotypical relationship in redundant space, possibly through an optimization process (a gray line). (**a**) In the visual space, this relationship plays a crucial role for the motor system in how it should implicitly respond to the various visual perturbations (bold arrows) in the redundant space. The motor system tries to eliminate the dissociation of the perturbed state from the original state (thin arrows). According to the relationship, the motor system interprets end-effector irrelevant errors (red or blue) as evidence that end-effector relevant errors have been also produced (dashed arrows). (**b**) However, the correction patterns are constrained by the relationship in the physical space. The relationship determined which movement pattern the motor system should perform. Thus, the correction patterns gravitate toward the relationship (bold arrows).

Materials and Methods

Participants

A total of 72 healthy adults (25 females and 47 males, aged 22.1 ± 2.5 years old; mean ± SD) participated in the study. Participants were recruited using an online recruitment system (https://www.jikken-baito.com). All the participants were right-handed according to their laterality score (83.3 ± 13.7; mean ± SD) obtained using the Edinburgh Handedness Inventory (Oldfield, 1971). They provided written informed consent, and the experimental protocol was approved by the ethics committee of The University of Tokyo (#19-225).

Experimental apparatus

The participants performed reaching movements using a virtual stick displayed on a horizontal screen with both handles of a manipulandum (KINARM End-Point Lab; Kinarm, Kingston, Canada) (Scott, 1999) (Fig. 2a). This screen prevented the participants from directly seeing their arms and handles, and displayed a start position, a target (1.4 cm in diameter), and a virtual stick (length: 40 cm, width: 0.5 cm). A white circle (1 cm diameter) was displayed on the right tip of the stick. The participants held the left tip of the stick with their left hand (left handle) and the stick 15 cm away from the left hand with their right hand (right handle). The distance between the handles was fixed using a strong elastic force produced by the manipulandum (spring constant = 2000 N/m). The positions and velocities of the hands were analogue/digitally converted at 1.129 kHz and then recorded at 1 kHz for offline analysis.

Baseline phase

All the experiments (Experiments 1–3) comprised a baseline phase (360 trials) followed by an adaptation phase (240 trials). The baseline phase, which was common to all experiments, was performed to investigate the basic strategy of how the participants moved the stick to reach the targets with the right tip. Before each reaching movement, the participants were asked to move the stick to the home position with the help of a manipulandum (Fig. 2c). After maintaining the home position for 0.5−1 s, a gray target was pseudo-randomly chosen from nine possible targets (0°: horizontal direction, ±10°, ±20°, ±30°, ±40°) appearing 10 cm away from the initial right tip position of the stick. After waiting for another 0.5−1 s, the color of the target changed to pink, indicating that they should immediately start moving the tip to the target. The participants were instructed to maintain the peak velocities of the tip as constant as possible across the trials. When the tip reached

the target, a feedback message, "fast" or "slow," was presented on the monitor if the
movement speed was higher or lower than the range of 300−450 mm/s, respectively.
After maintaining the tip at the target for 0.5 s, the visual feedback of the stick
disappeared. Then, the handles automatically returned to their home position, and the
next trial began. During the subsequent 240 trials in the adaptation phase, the target
appeared only in the 0° direction, and distinct visual perturbations were introduced in
Experiments 1–3 as described below. The home positions, size of the visual feedback
and the target, inter-trial interval, and criteria for the tip velocity feedback for the
adaptation phase were the same as those in the baseline phase.

Experiment 1

Experiment 1A (N = 19) and Experiment 1B (N = 13) were performed to investigate
how participants moved the stick when adapting to end-effector relevant perturbations
(tip perturbations). In Experiment 1A, the tip-movement direction on the monitor was
gradually rotated CCW direction around the starting position by 1° per trial up to 30°
(Fig. 4a). The stick-tilt angle remained unchanged (i.e., the stick-tilt angle was
identical to the angle of the line connecting both hands). In Experiment 1A,
participants did not perceive the presence of rotation and the adaptation was
considered to be dominated by the implicit process. On the other hand, in Experiment
1B, the 30° visual rotation was abruptly applied to the tip-movement direction, which
allowed them to use an explicit strategy.

Experiment 2

Experiment 2 (N = 20) examined whether the motor system was indifferent to
end-effector irrelevant perturbations (stick-tilt perturbations). The stick-tilt angle on
the monitor was rotated around the right tip in the CW direction (E2CW group, N =
10) or in the CCW direction (E2CCW group, N = 10). Notably, these perturbations did
not affect the tip-movement direction. During each reaching trial, the stick-tilt angle
increased from 0° to 6° in proportion to the distance of the tip position from the
starting point (0.6°/cm) (Fig. 6a). Participants did not notice any perturbations during
the experiment.

Experiment 3

In Experiment 3 (N = 20), to examine how end-effector irrelevant stick-tilt
perturbations influenced adaptation to end-effector relevant tip perturbations, both tip-
movement and stick-tilt perturbations were simultaneously imposed (Fig. 8a). Ten
participants participated in a task in which CCW tip and CCW stick-tilt rotations were
imposed (E3CCW group). The remaining ten participants participated in a task in

which CCW tip and CW stick-tilt rotations were imposed (E3CW group). The
procedure for applying tip and stick-tilt perturbations was the same as in Experiment
1A and Experiment 2, respectively. The participants did not notice perturbations
during the experiment.

Data analysis
Data analyses were performed using MATLAB R2023a (MathWorks, Natick,
Massachusetts, USA). The behavioral data were low-pass filtered (10 Hz, fourth-
order Butterworth filter). The tip-movement direction was calculated as the movement
direction of the tip relative to the horizontal direction (Fig. 3a). The stick-tilt angle was
defined as the tilting angle of the stick from the horizontal line (Fig. 3a). Positive
values for both parameters were defined as the degree to the CCW direction. Unless
otherwise noted, the tip-movement direction and the stick-tilt angle at the tip's peak
velocity were used to evaluate the feedforward motor commands.

Minimum distance of hand movement
The total movement distance of both hands in Figs. 3b and 3c was estimated by
summing the Euclidean distances between the initial and final positions of the hands.
The distance is represented as a function of the stick-tilt angle at the movement
offset (θ).

$$f(\phi, \theta) = \sqrt{(d\cos\phi - L\cos\theta + L)^2 + (d\sin\phi - L\sin\theta)}$$
$$+ \sqrt{(d\cos\phi - R\cos\theta + R)^2 + (d\sin\phi - R\sin\theta)} \qquad (1)$$

The first term on the right side represents the distance of the left-hand movement,
whereas the second term represents the distance of the right-hand movement. ϕ, d,
L, and R represent a target direction (0°, ±10°, ±20°, ±30°, and ±40°), a target
distance (10 cm), a distance between the left hand and the tip (40 cm), and a
distance between the right hand and the tip (25 cm), respectively. The minimum
distance of hand movement was derived by solving the following equation:

$$\frac{\partial f}{\partial \theta} = 0 \qquad (2)$$

635   **Statistical analysis**
636   To statistically evaluate the behavioral data, we conducted t-tests, correlation
637   analyses, two-way repeated-measures ANOVA, and MANOVA using MATLAB,
638   the software package JASP (http://jasp-stats.org), and R studio.

Data Availability

Data collected in Experiments 1–3 and the code are uploaded in zenodo and are available at https://zenodo.org/record/8033232.

References

Flash T, Hogan N. 1985. The Coordination of Arm Movements: An Experimentally Confirmed Mathematical Model. *Journal of Neuroscience* **5**:1688−1703.

Gordon J, Ghilardi MF, Ghez C. 1994. Accuracy of planar reaching movements I. Independence of direction and extent variability. *Experimental Brain Research* **99**:97−111.

Thoroughman KA, Shadmehr R. 1999. Electromyographic Correlates of Learning an Internal Model of Reaching Movements. *Journal of Neuroscience* **19**:8573−8588.

Mazzoni P, Krakauer JW. 2006. An Implicit Plan Overrides an Explicit Strategy during Visuomotor Adaptation. *Journal of Neuroscience* **26**:3642−3645.

Taylor JA, Ivry RB. 2011. Flexible Cognitive Strategies during Motor Learning. *PLoS Computational Biology* **7**:e1001096.

Takiyama K, Hirashima M, Nozaki D. 2015. Prospective errors determine motor learning. *Nature Communications* **6**:5925.

Hayashi T, Yokoi A, Hirashima M, Nozaki D. 2016. Visuomotor Map Determines How Visually Guided Reaching Movements are Corrected Within and Across Trials. *eNeuro* **3**:e0032-16.2016.

Bernstein NA, Latash MA, Turvey MT. 1996. Dexterity and Its Development. Psychology Press.

Singh P, Jana S, Ghosal A, Murthy A. 2016. Exploration of joint redundancy but not task space variability facilitates supervised motor learning. *Proceedings of the National Academy of Sciences of the United States of America* **113**:14414−14419.

Collins JJ. 1995. The Redundant Nature of Locomotor Optimization Laws. *Journal of Biomechanics* **28**:251−267.

O'Sullivan I, Burdet E, Diedrichsen J. 2009. Dissociating Variability and Effort as Determinants of Coordination. *PLoS Computational Biology* **5**:e1000345.

White O, Diedrichsen J. 2010. Responsibility assignment in redundant systems. *Current Biology* **20**:1290−1295.

Zhang Z, Guo D, Huber ME, Park S-W, Sternad D. 2018. Exploiting the geometry of
the solution space to reduce sensitivity to neuromotor noise. *PLoS Computational*
Biology **14**:e1006013.

Mosier KM, Scheidt RA, Acosta S, Mussa-Ivaldi FA. 2005. Remapping Hand
Movements in a Novel Geometrical Environment. *Journal of Neurophysiology*
94:4362−4372.

Latash ML. 2012. The bliss (not the problem) of motor abundance (not redundancy).
Experimental Brain Research **217**:1−5.

van Beers RJ, Brenner E, Smeets JBJ. 2013. Random walk of motor planning in
task-irrelevant dimensions. *Journal of Neurophysiology* **109**:969−977.

Sternad D. 2018. It's not (only) the mean that matters: variability, noise and
exploration in skill learning. *Current Opinion in Behavioral Sciences* **20**:183−195.

Scholz JP, Schöner G. 1999. The uncontrolled manifold concept: identifying control
variables for a functional task. *Experimental Brain Research* **126**:289−306.

Scholz JP, Kang N, Patterson D, Latash ML. 2000. Uncontrolled manifold analysis of
single trials during multi-finger force production by persons with and without Down
syndrome. *Experimental Brain Research* **153**:45−58.

Todorov E, Jordan MI. 2002. Optimal feedback control as a theory of motor
coordination. *Nature Neuroscience* **5**:1226−1235.

Todorov E. 2004. Optimality principles in sensorimotor control. *Nature Neuroscience*
7:907−915.

Scott SH. 2004. Optimal feedback control and the neural basis of volitional motor
control. *Nature Reviews Neuroscience* **5**:534−546.

Diedrichsen J. 2007. Optimal Task-Dependent Changes of Bimanual Feedback
Control and Adaptation. *Current Biology* **17**:1675−1679.

Valero-Cuevas FJ, Venkadesan M, Todorov E. 2009. Structured Variability of
Muscle Activations Supports the Minimal Intervention Principle of Motor
Control. *Journal of Neurophysiology* **102**:59−68.

Ronsse R, Wei K, Sternad D, 2010. Optimal Control of a Hybrid Rhythmic-Discrete
Task: The Bouncing Ball Revisited. *Journal of Neurophysiology* **103**:2482−2493.

Sternad D, Abe MO, Hu X, Müller H. 2011. Neuromotor Noise, Error Tolerance and
Velocity-Dependent Costs in Skilled Performance. *PLoS Computational Biology*
7:e1002159.

Selgrade BP, Chang Y-H. 2015. Locomotor control of limb force switches from
minimal intervention principle in early adaptation to noise reduction in late
adaptation. *Journal of Neurophysiology* **113**:1451−1461.

Wolpert DM, Ghahramani Z, Jordan MI. 1995. Are arm trajectories planned in
kinematic or dynamic coordinates? An adaptation study. *Experimental Brain*
Research **103**:460−470.

Flanagan JR, Rao AK. 1995. Trajectory Adaptation to a Nonlinear Visuomotor
Transformation: Evidence of Motion Planning in Visually Perceived Space.
Journal of Neurophysiology **74**:2174−2178.

Morehead JR, Taylor JA, Parvin DE, Ivry RB. 2017. Characteristics of Implicit
Sensorimotor Adaptation Revealed by Task-irrelevant Clamped Feedback.
Journal of Cognitive Neuroscience **29**:1061−1074.

Diedrichsen J, White O, Newman D, Lally N. 2010. Use-Dependent and Error-Based
Learning of Motor Behaviors. *Journal of Neuroscience* **30**:5159−5166.

Schaefer SY, Shelly IL, Thoroughman KA. 2012. Beside the point: motor
adaptation without feedback-based error correction in task-irrelevant conditions.
Journal of Neurophysiology **107**:1247−1256.

Kim HE, Parvin DE, Ivry RB. 2019. The influence of task outcome on implicit
motor learning. *eLife* **8**:e39882.

Cunningham HA. 1989. Aiming Error Under Transformed Spatial Mappings
Suggests a Structure for Visual-Motor Maps. *Journal of Experimental Psychology*
15:493−506.

Krakauer JW, Pine ZM, Ghilardi M-F, Ghez C. 2000. Learning of Visuomotor
Transformations for Vectorial Planning of Reaching Trajectories. *Journal of*
Neuroscience **20**:8916−8924.

Ethier V, Zee DS, Shadmehr R. 2008. Spontaneous Recovery of Motor
Memory During Saccade Adaptation. *Journal of Neurophysiology* **99**:2577−2583.

Malone LA, Vasudevan EVL, Bastian AJ. 2011. Motor Adaptation Training for
Faster Relearning. *Journal of Neuroscience* **31**:15136−15143.

Vaswani PA, Shmuelof L, Haith AM, Delnicki RJ, Huang VS, Mazzoni P, Shadmehr R,
Krakauer JW. 2015. Persistent Residual Errors in Motor Adaptation Tasks:
Reversion to Baseline and Exploratory Escape. *Journal of Neuroscience*
35:6969−6977.

Albert ST, Jang J, Sheahan HR, Teunissen L, Vandevoorde K, Herzfeld DJ,
Shadmehr R. 2021. An implicit memory of errors limits human sensorimotor
adaptation. *Nature Human Behaviour* **5**:920−934.

Omrani M, Diedrichsen J, Scott SH. 2013. Rapid feedback corrections during a
bimanual postural task. *Journal of Neurophysiology* **109**:147−161.

Ernst MO, Banks MS. 2002. Humans integrate visual and haptic information in a
statistically optimal fashion. *Nature* **415**:429−433.

Oldfield RC. 1971. The Assessment and Analysis of Handedness: The Edinburgh
Inventory. *Neuropsychologia* **9**:97−113.
Scott SH. 1999. Apparatus for measuring and perturbing shoulder and elbow joint
positions and torques during reaching. *Journal of Neuroscience Methods*
89:119−127.

Acknowledgments
We thank members of the Nozaki laboratory for their helpful comments and
suggestions, Jinnosuke Ueno, Junnosuke Sakuragi, and Ryosuke Miyazawa for
performing preliminary experiments, Asako Munakata and Mayumi Yoda for
coordinating experiments. This study was supported by a grant from the Japan
Society for the Promotion of Science Research Fellowships for Young Scientists to
T.K. (20J13734) and a KAKENHI to D.N. (17H00874, 21H04860) and to T.K.
(23K10739).

Competing Interest Statement
Authors declare that they have no competing interests.