TITLE

Consideration of sex as a biological variable in diabetes research across twenty years

AUTHORS

Celena Cherian[1*], Hayley Reeves[1,2*], Duneesha De Silva[1,3&], Serena Tsao[1,4&], Katie E.

Marshall[5], Elizabeth J. Rideout[1§]

AFFILIATIONS

[1] Department of Cellular and Physiological Sciences, Life Sciences Institute, The University of

British Columbia, Vancouver, Canada.

[2] Present address: School of Molecular Biosciences, University of Glasgow, Glasgow, UK.

[3] Present address: Department of Orthopaedics, The University of British Columbia,

Vancouver, Canada.

[4] Department of Anatomy and Cell Biology, Faculty of Medicine and Health Sciences, McGill

University, Montréal, Canada.

[5] Department of Zoology, The University of British Columbia, Vancouver, Canada.

[*] These authors contributed equally to this work
[&] These authors contributed equally to this work
[§] Corresponding author
20
21 **CORRESPONDING AUTHOR DETAILS**

22 Elizabeth J. Rideout, Life Sciences Center, 2350 Health Sciences Mall (RM3308), Vancouver,

23 BC, Canada V6T 1Z3. Email: elizabeth.rideout@ubc.ca

24
25 **KEYWORDS**

26 Biological sex, sex-based analysis, sex difference, diabetes

ABSTRACT

Sex differences exist in the risk of developing both type 1 and type 2 diabetes, and in the risk of developing diabetes-associated complications. Sex differences in glucose homeostasis, islet and β cell biology, and peripheral insulin sensitivity have also been reported in multiple animals. To determine the degree to which biological sex has been addressed in published literature related to diabetes and insulin biology, we developed a scoring system to assess the inclusion of biological sex in papers related to these topics. We scored manuscripts published in *Diabetes*, published by the American Diabetes Association, as this journal focuses on diabetes and diabetes-related research. We scored papers published across three years within a 20-year period (1999, 2009, 2019), a timeframe that spans the introduction of funding agency and journal policies to improve the consideration of biological sex as a variable. Our analysis shows fewer than 15% of papers used sex-based analysis in even one figure across all study years, a trend that was reproduced across journal-defined categories of diabetes research (*e.g.*, islet studies, signal transduction). Single-sex studies accounted for approximately 40% of all manuscripts, of which >87% used male subjects only. While we observed a modest increase in the overall inclusion of sex as a biological variable during our study period, our results highlight significant opportunities to improve consideration of sex as a biological variable in diabetes research. In particular, we show that journal policies represent one way to promote better consideration of biological sex as a variable. In the long term, improved practices will reveal sex-specific mechanisms underlying diabetes risk and complications, generating insights to support the development of sex-informed prevention and treatment strategies.

INTRODUCTION

Diabetes mellitus is a disease that affects millions of individuals globally. According to the

International Diabetes Federation, approximately 537 million individuals (20-79 years)

worldwide live with diabetes and it has caused 6.7 million deaths[1]. There are also rare

monogenic forms of diabetes that affect many people[2]. While diverse factors influence the risk

of developing diabetes[3–9], there is growing recognition that biological sex impacts the risk of

developing diabetes and diabetes-related complications[8–13]. In several populations, the risk of

developing type 2 diabetes (T2D) is approximately 40% higher in males than in

females[8,9,11,12,14,15]. In adults aged 15-40, males also have a 60% higher risk of developing

type 1 diabetes (T1D)[16]. This male bias in T1D is particularly notable given that females are at

a higher risk of nearly all other autoimmune-related diseases[17–19]. After diabetes onset,

females are at higher risk of diabetic heart disease, stroke mortality, depression, and anxiety,

whereas males show faster progression of nephropathy and require lower extremity

amputation more often[9,20–25].

Clues into factors that contribute to sex differences in the risk of developing diabetes

and diabetes-related complications include a male-female difference in peripheral insulin

sensitivity. In humans and many animal models[26–29], females typically show higher insulin

sensitivity[30–34]. Given the established links between reduced peripheral insulin sensitivity and

T2D[27,34–41], and between insulin sensitivity and diabetes-associated complications[9,21–25,42],

biological sex is an important variable to consider when studying insulin sensitivity across

physiological and pathological contexts. Sex differences also exist across many animals,

including humans, in the biology of insulin-producing β cells[30,43–47]. Male-female differences

have been reported in the number of pancreatic β cells[47], and studies in humans and rodents

report profound sex differences in β cell gene expression, function, and stress responses in

both normal and T2D contexts[30–34,48,49]. Female β cells also show higher glucose-stimulated

insulin secretion under normal physiological conditions and in T2D[9,30–32], differences that

cannot be solely attributed to sex differences in peripheral insulin sensitivity[30]. Because there

are established links between β cell dysfunction in rodent models and humans in T2D[49–55],

and emerging evidence suggests β cell dysfunction may also contribute to T1D[56,57], biological

sex is an important variable to consider in studies on islets and β cells in multiple

physiological and pathological contexts. Yet, the degree to which biological sex has been

considered in the literature with respect to islet and β cells is incompletely known.

Broad-based text searches of diabetes articles, and detailed assessments of

randomized controlled trials and human observational studies in top diabetes journals,

suggest that biological sex is rarely considered as a variable in diabetes research[58–61].

Indeed, one study showed that only 6.5% of randomized controlled trials and human

observational studies reported all outcomes separately according to biological sex[58]. While

these studies were critical in highlighting the overall poor consideration of both sexes in

diabetes research, several questions remain. For example, the prior focus on randomized

controlled trials and observational studies means that we lack detailed information on whether

biological sex is included in studies that focus on cell, tissue, and animal models of diabetes.

We similarly lack information on whether there are differences in practices related to

biological sex between different diabetes subject areas (*e.g.* islet biology, transplantation),

and whether funding agency policies mandating the inclusion of both sexes[59,62–65] lead to

better consideration of biological sex in diabetes research. Answering these questions will

highlight opportunities for improvement in research design, methods, and analysis across all

areas of diabetes research, and establish whether policies introduced by major funding

agencies, journals, and governments have produced sufficient change in research

practices[62–66].

We performed a detailed analysis of all manuscripts published in the journal *Diabetes*,

published by the American Diabetes Association, in 1999, 2009, and 2019 (819 publications)

(Fig. 1A). Studying the inclusion of sex as a biological variable in one journal rather than

between journals provides advantages due to uniform journal policies and similar manuscript

quality. Because *Diabetes* publishes papers in multiple subject areas, and in both clinical and

biomedical disciplines, we captured and compared differences in common practices across

diverse fields related to diabetes mechanisms and risk. Overall, our findings suggest that

fewer than 15% of studies published in *Diabetes* during our study period analyzed data using

biological sex as a variable. Approximately 34% of studies included both sexes in data

collection, but did not analyze their data using biological sex as a variable, and 40% of studies

included only a single sex. While we observed a significant increase in the inclusion of both

sexes in diabetes research over time, we observed a smaller increase in the number of

studies that used sex-based analysis over time. These trends were broadly similar across

many subfields of diabetes research, and funding status had no significant effect on overall

consideration of biological sex as a variable in diabetes research. Together, our data

suggests progress has been made in diabetes-related research with respect to the inclusion

of both sexes and uptake of sex-based analysis; however, significant opportunities for

improvement remain. This is especially true for single-sex studies, the vast majority of which

used male and not female animals. Greater knowledge of how biological sex affects diabetes

mechanisms and complications will ultimately support the development of sex-informed

prevention and treatment strategies to improve equity in health outcomes[61,66].

RESULTS

Assessing the inclusion of biological sex as a variable in diabetes research over time

The start of the twenty-year period we considered in our study corresponded with the introduction of major policy initiatives aimed at improving the consideration of biological sex as a variable in Canada, Europe, and the United States[62–66]. To determine whether the inclusion of biological sex as a variable in diabetes research changed over time, we monitored the average score of the papers in our study in 1999, 2009, and 2019. The average score of papers published in 1999 was 1.47 ± 0.07, whereas the average scores of manuscripts published in 2009 and 2019 were 1.91 ± 0.08 and 2.28 ± 0.1, respectively (Fig. 1B). While this represents a significant increase in average score between 1999 and 2019 (year effect $p<0.001$; ANOVA in a generalized linear model with Poisson distribution), an increase in average score of only 0.8 over 20 years indicates that significant room for improvement remains in the inclusion of sex as a biological variable in diabetes research.

To describe the dynamics underlying the change in average score over time in more detail, we examined the proportion of studies classified into each of our six categories. The proportion of studies assigned a score of 0 significantly declined over time, from 27% in 1999, to 19% in 2009, and 7.3% in 2019 (year effect $p<0.001$; ANOVA in a generalized linear model with binomial distribution) (Fig. 1C). In contrast, there was a significant increase in the proportion of papers over time assigned a score of 3, 4, or 6 (year effect $p=0.002$, $p=0.004$, $p=0.029$ respectively; ANOVA in a generalized linear model with binomial distribution) (Fig. 1C). Papers assigned a score of 3 comprised 25.5% of papers in 1999, 36.5% in 2009, and 37.6% in 2019 (Fig. 1C). Papers assigned a score of 4 made up 4.2% of papers in 1999, 6% papers in 2009, and 10.6% papers in 2019, whereas papers assigned a score of 6 represented only 0.91% of studies in 1999, 1.6% in 2009, and 3.9% in 2019 (Fig. 1C).

No significant change was observed in the proportion of papers scored as a 1, 2, or 5

between 1999 and 2019 (year effect $p=0.618$, $p=0.500$, $p=0.991$ respectively; ANOVA in a

generalized linear model with binomial distribution) (Fig. 1C). Based on this data, the

categories with the most pronounced changes between 1999 and 2019 were 0 (decrease)

and 3 (increase) (Fig. 1C). This suggests the higher average score in 2019 may be partly

attributed to a drop in studies that did not discuss biological sex at all and an increase in

mixed-sex studies with no sex-based analysis. Increased inclusion of both sexes represents a

promising step toward better consideration of biological sex as a variable in diabetes

research; however, important next steps for improvement include greater use of sex-based

analysis.

Male animals are used more often in single-sex studies

Previous studies on the inclusion of biological sex as a variable in biomedical research reveal

that male animals are used more frequently than female animals[59,60,68]. For example, a study

from 2010 analyzed almost 2,000 animal studies across 10 biological fields in several journals

and found a male bias in eight fields, with the most pronounced bias in neuroscience,

pharmacology, and physiology[60]. To determine whether one sex was used more often in

diabetes research, for all papers assigned a score of 1 or 2 (single-sex studies) we recorded

the sex that was used. Studies assigned a score of 1-2 represented 42% of studies in 1999,

40% of studies in 2009, and 40% of studies in 2019 (Fig. 1D), suggesting single-sex studies

represented a large proportion of total studies published during this period. In 1999, 71.7% of

single-sex studies used only male animals (Fig. 1D). In 2009 and 2019, 82.4% and 87.3% of

single-sex studies used only male animals, respectively (Fig. 1D). When we examined the

number of male-only studies over time, we observed a significant increase between 1999 and

2019 (year effect $p=0.034$; ANOVA in a generalized linear model with binomial distribution)

(Fig. 1D). This suggests that in diabetes research, as in other fields of biomedical

science[59,60,68], male animals are used more frequently than female animals.

Evaluating the inclusion of biological sex as a variable in diabetes research across

clinical and biomedical disciplines

Diabetes research includes both clinical and biomedical disciplines. Given that policies

governing the inclusion of biological sex as a variable were introduced at an earlier date for

clinical research compared with biomedical research[59,60,69,70], we compared the average score

assigned to studies in each of these disciplines in 1999, 2009, and 2019. Across all three

study years, the average score for papers in the clinical discipline (2.78 ± 0.08) was

significantly higher than the average score for papers in the biomedical discipline ($1.48 \pm$

0.05) (discipline effect $p<0.001$; ANOVA in a generalized linear model with Poisson

distribution) (Fig. 2A). We observed a significant increase in the average scores across both

biomedical and clinical disciplines between 1999 and 2019 (year effect $p<0.001$; ANOVA in a

generalized linear model with Poisson distribution) (Fig. 2A), with a trend toward a larger

increase over time in the biomedical discipline.

To identify factors that contribute to the score difference between the biomedical and

clinical disciplines, we examined the probability that a study in each category would be

assigned a specific score (Fig. 2B,C). While there was a significant decrease in the proportion

of both biomedical and clinical studies assigned a score of 0 over time (year:discipline

interaction $p<0.001$; ANOVA in a generalized linear model with binomial distribution), the

probability that a biomedical study would be assigned a score of 0 was higher than the

probability of a clinical study being assigned this score across all study years (discipline effect

$p<0.001$; generalized linear model with binomial distribution) (Fig. 2B,C). The probability of a

biomedical study receiving a score of 1 was also significantly higher than the probability of a

clinical study being assigned this score in all three study years (discipline effect $p<0.001$;

generalized linear model with binomial distribution) (Fig. 2B).

In the clinical discipline, the score that was assigned to the highest proportion of

studies across all years was 3 (61.2% of studies; Fig. 2C). Indeed, the probability of a clinical

study being assigned a score of 3, 4 or 5 was significantly higher than the probability of a

biomedical study being assigned these scores in all study years (discipline effect $p<0.001$,

$p=0.002$ and $p=0.03$, respectively; generalized linear model with binomial distribution) (Fig.

2B,C). While our data indicate that more authors indicated the sex of the model system they

used in 2019 than in 1999, significant room for improvement remains. For example, in 2019

the majority of biomedical studies still used only a single sex (~50%) and most failed to use

sex-based analysis (86.6%). In the clinical discipline, fewer than 15% of published studies

used sex-based analysis during our study period.

Comparing the inclusion of biological sex as a variable between diabetes subject areas

Beyond the diverse approaches to diabetes research between clinical and biomedical

disciplines, diabetes research spans a broad range of topics, from genetic and genomic

approaches, to immunology, to pharmacology. We therefore wanted to compare the inclusion

of biological sex as a variable across different diabetes subject areas. From 2009 onwards, all

studies published in *Diabetes* correspond to one of 10 primary topics/subjects (Table 2). Note

that not all areas maintained identical labels in 2009 and 2019. For example, the topic

'Genetics' in 2009 was compared to 'Genetics/Genomes/Proteomics/Metabolomics' in 2019

(hereafter called Genetics/Omics). Similarly, the 'New Methodologies and Databases' was

compared to 'Technological Advances' in 2019 (hereafter called New Methodology and

Technology). To ensure adequate sample sizes within each topic, we excluded subject areas

with fewer than ten papers in either 2009 or 2019 (Signal Transduction, New Methodology

and Technology, Pharmacology & Therapeutics).

In 2009, the average score was highest in the Genetics/Omics and Obesity subject

areas (2.5 ± 0.17 and 2.62 ± 0.39, respectively), and the lowest in Immunology and

Transplantation and Islet topics (1.62 ± 0.27 and 1.35 ± 0.23, respectively; Fig. 3A). In 2019,

the average scores were highest in the Genetics/Omics, Obesity, and Immunology and

Transplantation topics (3.04 ± 0.20, 2.67 ± 0.34, and 2.72 ± 0.38, respectively) (Fig. 3A). In

both 2009 and 2019, the higher score for studies under the Genetics/Omics subject area may

be attributed to the higher probability of a study in that subject area being assigned a score of

3 compared with other subject areas (Fig. 3B-H). Despite the different average scores

between subject areas, our model suggests that all subject areas showed a similar change in

average score over time (year:subject interaction $p=0.347$; ANOVA in a generalized linear

model with Poisson distribution). Thus, our analysis did not reveal widespread differences in

practices related to the inclusion of biological sex as a variable between different diabetes

subject areas.

Evaluating the effect of funding source on the inclusion of biological sex as a variable

in diabetes research

Funding agencies across many countries have introduced policies aimed at increasing the

inclusion of biological sex as a variable in research[59,62–66]. The US National Institutes of

Health (NIH) represents a key source of funds for health research, with significant

investments in diabetes research[63]. The NIH was among the first funding agencies worldwide

to introduce policies related to the inclusion of both sexes in both clinical and biomedical

research[59,62,64–66]. To evaluate whether NIH funding affected the inclusion of biological sex as

a variable in diabetes research, we compared the average score assigned to studies by

authors that declared NIH funding with scores of studies by authors that did not declare NIH

funding (Fig. 4A). We found that the average score of NIH-funded studies was not

significantly higher than studies that declared funding from other sources (funding effect

p=0.063; year:funding interaction p=0.974; ANOVA in a generalized linear model with Poisson

distribution) (Fig. 4A). We next compared the proportion of studies within each score category

between NIH-funded studies and non-NIH-funded studies (Fig 4B,C). For the majority of

score categories (0-5), NIH funding had no significant effect on the proportion of studies

assigned a specific score (funding effect p=0.942, p=0.236, p=0.897, p=0.545, p=0.771,

p=0.277, respectively; ANOVA in a generalized linear model with binomial distribution) (Fig.

4B,C). In contrast, NIH-funded studies showed a significantly higher proportion of studies

assigned a score of 6 across all study years compared with non-NIH-funded studies (funding

effect p<0.001; funding:year interaction p=0.049; ANOVA in a generalized linear model with

binomial distribution) (Fig. 4B,C). This data indicates that NIH funding was not associated with

overall shifts in practices related to the consideration of biological sex as a variable in

diabetes research. Indeed, despite the fact that policies related to the inclusion of biological

sex as a variable in clinical research[16] were introduced earlier than similar policies related to

biomedical research[62,64,65], there was no significant interaction between year, funding source

and type of study when we asked whether NIH funding affected the average scores assigned

to clinical and biomedical studies (Fig. 4D). This reinforces our conclusion that NIH funding

was not associated with increased consideration of biological sex as a variable in diabetes

research.

Evaluating the effect of journal policies on inclusion of sex as a biological variable

In 2019, a new policy was introduced at *Diabetes* mandating that authors of studies that use

human islets provide detailed information on donor characteristics[71–73]. The goal of including

this detailed donor information was to enhance transparency and to enable better

comparisons among data generated using human islets. One of the required donor

characteristics included in the Human Islet checklist is biological sex[73]. Because disclosing

the sex of experimental models represents one way that authors can improve consideration of

biological sex in their research, we assessed the effect of this policy change on the scores

assigned to papers in the 'Islet studies' subject area before and after the policy came into

effect in April 2019. We scored all manuscripts published in this subject area in 2018 (36

papers) and 2020 (32 papers), as this was the topic most likely to be affected by a policy

regarding human islet documentation. We did not score papers published in 2019 as these

papers were likely submitted and/or reviewed prior to the policy change.

We found the average score of papers in the Islet subject area in 2018 was $1.66 \pm$

0.64, whereas the average score of studies in this area was 2.5 ± 0.19 in 2020 (Fig. 5A). This

represented an increase of 0.8 in the average score of studies in the Islet subject area over

only a two-year period (Fig. 5A). While this increase was not statistically significant

(year:subject interaction $p=0.248$; ANOVA in a generalized linear model with Poisson

distribution) it is worth noting that the magnitude of this increase is equivalent to the increased

we observed in all studies across our twenty-year study period. Potential explanations for the

increase in average score for studies in the Islet subject area include a trend toward more

studies assigned a score of 4-6, and a trend toward fewer studies assigned a score of 0-1

(Fig. 5B). To determine whether this increase was reproduced in another subject area less

likely to be affected by the policy change regarding human islets, we repeated our analysis of

manuscripts published in 2018 and 2020 in the 'Metabolism' subject area. The average score

for studies in the 'Metabolism' subject area increased by 0.03 over the span of the policy

change and was not significantly different in 2018 compared with 2020 (year:subject

interaction p=0.248; ANOVA in a generalized linear model with Poisson distribution) (Fig. 5A).

Taken together our analysis suggests that the editorial policy change at *Diabetes* may have

contributed to a significant improvement in the consideration of biological sex as a variable in

diabetes research.

DISCUSSION

The overall goal of our study was to assess the degree to which biological sex was included

as a variable in diabetes research across a twenty-year period. We chose to assess studies in

a journal that is dedicated to publishing high-quality studies in the area of diabetes research,

and across a timeframe that captures widespread changes to journal and funding agency

policies regarding sex- and gender-based analysis[62,64,65,71–73]. While our analysis revealed

significant increases in the inclusion of biological sex as a variable over time, in alignment

with the reported increase in the word "sex" in text-based searches of a similar body of

literature[58], our detailed evaluation of methods demonstrates that many studies do not

adequately address biological sex as a variable in diabetes research. Given that biological

sex affects diabetes risk[1,9,16,56], treatment efficacy[3,8,72], and risk of developing diabetes

complications[9,20,21,25,42], this knowledge gap must be closed in order to develop sex-informed

prevention and treatment strategies for people living with diabetes.

One major barrier to the full consideration of biological sex as a variable in diabetes

research was related to inclusion. In all three study years across a twenty-year period,

between 46 and 69% of studies did not include both sexes (score 0-2). This low inclusion of

both sexes was particularly notable in biomedical studies (between 59% and 81% of studies

across the 3 years were assigned a score of 0-2) and in diabetes subject areas that contained

a large proportion of biomedical studies (*e.g.,* islets and metabolism). A lack of inclusion of

both sexes has been described in studies of literature related to neuroscience, psychiatry and

psychology, pharmacology, and physiology[60,68,69,74,75], and our data aligns with findings from

others[58,69,70,74,76,77] suggesting similar trends exist in diabetes research. Indeed, the inclusion

of both sexes did not show a meaningful increase over a twenty-year period in many subject

areas of diabetes research. Addressing this overall lack of inclusion is an important task going

forward for the diabetes community as funding agencies and journals worldwide aim to

improve transparency and reproducibility in research[59,66]. Because our study showed the

majority of single-sex studies use only male subjects, the major change that is needed in

biomedical studies is the inclusion of female subjects. Indeed, our analysis suggests that the

gap in knowledge of female cells and animals will not be closed if current trends in single-sex

studies continue. Given that female subjects are protected from dysregulation of glucose

homeostasis[27,30–32,34,37,38], but are at a higher risk of developing complications from

diabetes[9,21,42], mechanistic insights gained from studying females will provide valuable

insights into diabetes prevention and treatment.

Another major barrier to the full consideration of biological sex as a variable in diabetes

research was a lack of sex-based analysis. Studies assigned a score of 3 comprised between

17% and 28% of those published in the biomedical discipline and between 51% and 77% of

clinical studies between 1999 and 2019. In the biomedical discipline, the most common

reason studies were assigned a score of 3 was that they pooled samples (*e.g.*, islets)

collected from male and female experimental models prior to collecting measurements. This

suggests that in biomedical studies, the major changes that are needed to support sex-based

analysis are to 1) collect samples and data from male and female experimental models

separately, and 2) to use appropriate statistical methods to analyze the data from each sex

separately. In clinical studies, the main reason studies were assigned a score of 3 was that

biological sex was not treated as a variable. Instead, most studies in the clinical discipline

considered biological sex as a covariate. Given that biological sex can interact with treatment

to drive outcomes, including biological sex only as a covariate can mask sex-dependent

effects. Biological sex should therefore be included either as an interactor, or analyses should

be stratified by sex[75]. The main change that is needed in the clinical discipline is therefore to

adjust study analysis methods to fully consider biological sex as a variable. Because many

studies show sex differences in islet and β cell function[41,47,48,78], the genetic architecture of

disease[9,41,42,55], and metabolic differences[27,30,32,33,38], shifts in practice across the biomedical

and clinical disciplines, and in different diabetes subject areas, will allow researchers to

develop evidence-based models for these topics in each sex. These models will form the

basis for the development of sex-informed prevention and treatment efforts for individuals at

risk for, or living with, diabetes.

Despite the growing number of studies showing that biological sex is not given

adequate consideration in biomedical and clinical research[60,69,70,79], the most effective

strategies to promote better uptake of biological sex as a variable are not fully understood.

Our data, in alignment with similar studies[74,76,77], indicate that funding by agencies with

clearly-stated requirements related to the consideration of biological sex does not significantly

influence author practices (Fig. 4). One potential reason for this lack of influence on author

practices is that monitoring compliance with sex- and gender-related policies poses multiple

challenges. Due to a policy change during the course of our study, we were able to assess

how journal policies affect the consideration of biological sex as a variable. Our data suggests

that journal policies may represent an effective tool in promoting the uptake and integration of

biological sex as a variable in diabetes research. In the 'Islet studies' subject area, the

increase in average score of 0.8 between 2018 and 2020 was similar in magnitude to the 0.91

increase in average score across the biomedical discipline over a twenty-year period. This

suggests the implementation of robust journal policies related to integration of biological sex

as a variable will promote better practices in the future. Applying these policies will require a

culture shift across editors, reviewers, and authors, however, proper consideration of

biological sex as a variable will provide significant benefits in terms of data reproducibility,

transparency, and scientific rigor[70,74,76,79].

Based on our analysis, we suggest several practices authors can adopt to improve

consideration of biological sex as a variable in their research. First, we discuss changes that

can be made in the short term. For example, one way to immediately improve transparency

regarding integration of biological sex as a variable is to ensure the sex of experimental

subjects are explicitly stated in the methods section and in figure legends. If only a single sex

is used, justification should be provided for the use of a single-sex experimental group.

Authors should also identify the use of a single sex as a limitation at some point in the paper,

and state that future studies will be needed to determine whether findings made in one sex

apply to the other. If both sexes are used, but not all phenotypes are recorded in each sex,

the figure legends should include this information to ensure the reader is aware of which sex

was used to generate specific data points.

It is also important to ensure data from each sex is collected and analyzed separately,

and not pooled. Pooling male and female cells or tissues during analysis risks overlooking

important effects if a sex-specific outcome does not achieve statistical significance within a

mixed-sex group. Similarly, potential interactions between biological sex and other variables

should be explicitly tested for, or the analysis should be stratified by sex, to allow for detection

of sex-specific effects. Treatment of biological sex as a covariate potentially masks those

effects. Because these suggestions are all related to changes in text, and how data are

displayed, described, and analyzed, they can be applied to studies that are either complete or

in progress. Longer-term changes include using male and female cell lines and animal

models, collecting and analyzing data from males and females separately, and applying

statistical methods to reveal how biological sex affects diverse phenotypes. The solid

foundation of knowledge that will emerge from these studies will help overcome the current

lack of information on male and female cells and animal models. It will also serve as a first

step in embracing diversity in sex chromosome complement and sex hormone levels, in line

with emerging recognition that biological sex is not a binary variable[13,61,70,75,77,79].

In conclusion, better knowledge of normal physiology in males and females will allow

the development of sex-informed prevention efforts, and better knowledge of pathophysiology

will support the discovery of more effective treatment options that reduce the burden of

complications. Broader consideration of biological sex as a variable in diabetes research will

therefore provide significant benefits to all individuals at risk for, and living with, diabetes.

METHODS

Article selection criteria. Our analysis included original research articles for which the full

text was available on the *Diabetes* online archive. We did not include reviews, editorials, or

commentaries in our analysis. We also excluded case studies, as reports on individuals

preclude the meaningful analysis of both sexes (Fig. 1A).

Scoring the inclusion of sex as a biological variable. Each paper was scored based on

the degree to which biological sex was integrated into study design and data analysis. A full

breakdown of the scoring system is included as Table 1. Briefly, a score of 0 was assigned to

studies in which the sex of animal models or participants was not stated at any point in the

text including the methods section. A score of 1 was assigned to studies that used a single

sex (male or female) but did not provide any rationale for this choice. A score of 2 was

assigned to studies that used a single sex but provided a reason for this decision. A score of

3 was given to mixed-sex studies. These studies included both sexes, but did not 1) indicate

the number of male and female subjects used and 2) analyze their data by sex. Clinical

studies that used biological sex as a covariate were also assigned a score of 3 because this

type of analysis does not properly consider biological sex effects as a variable. A score of 4

was given to studies that included biological sex as a variable in the analysis for as little as

one piece of data (*e.g.*, one panel of a figure reported male and female phenotypes

separately). A score of 5 was assigned to studies that included biological sex as a variable in

>50% of the data presented in the manuscript. A score of 6 was assigned for studies that

included both males and females across all experiments, and in which all data was analyzed

and reported by sex. The full data set is available in Supplemental Files 1 and 2.

Scoring reproducibility. After the full data set was collected, a subset of articles was re-

scored by another person to assess inter-individual scoring variability. Of the 86 papers that

were selected at random for re-scoring, 96.5% (83/86) were assigned the same score by

different individuals. This indicates consistency between individuals in the application of our

scoring system.

Recording the sex of experimental subject. For papers with a score of '1' or '2' the sex of

the experimental subjects, as reported by the authors, was recorded in our spreadsheet.

Recording diabetes discipline and subfields. Each paper was classified as either 'clinical'

or 'biomedical' to allow a comparison between these two disciplines regarding the inclusion of

biological sex as a variable. Clinical studies included randomized controlled trials and studies

in which subjected human participants were subjected to treatment protocols. Observational

studies involving human subjects were also classed as clinical when material or recordings

beyond a single blood sample, superficial tissue, or DNA sample was taken for analysis. All

other papers, including those which used animal models or cell lines, were classed as

biomedical. For papers with both clinical and biomedical data, the classification was based on

the type of data from which the majority (>50%) of the results were generated. For papers

published in 2009 and 2019 the research subject area of each study was also recorded based

on the classification system used by *Diabetes* (*e.g.*, 'Islet biology', 'Obesity'; see Table 2).

Because papers published in 1999 were not categorized according to subject area, this

information was only collected for 2009 and 2019.

Funding status. We recorded the funding source for all papers in which this information was

reported by the authors.

Statistical analysis. All statistical analyses were performed using R[67]. For all analyses,

generalized linear models were used. For mean score changes, a Poisson error distribution

was assumed as scores are integers > -1. To calculate changes in proportions of scores, and

the percentage of male- and female-biased studies, the scores were converted to a binary

format and a binary (logistic) distribution was assumed. All analysis was started with a

maximal model and *p*-values from the resulting ANOVA table are reported. p values are

available in Supplemental File 3.

REFERENCES

1.   Sun H, Saeedi P, Karuranga S, et al. IDF Diabetes Atlas: Global, regional and country-level
diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract*.
2022;183:109119. doi:10.1016/j.diabres.2021.109119
2.   Bonnefond A, Unnikrishnan R, Doria A, et al. Monogenic diabetes. *Nat Rev Dis Primer*.
2023;9(1):12. doi:10.1038/s41572-023-00421-w
3.   American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus. *Diabetes
Care*. 2010;33(Supplement_1):S62-S69. doi:10.2337/dc10-S062
4.   Inzucchi SE. Management of Hyperglycemia in the Hospital Setting. *N Engl J Med*.
2006;355(18):1903-1911. doi:10.1056/NEJMcp060094
5.   Lin X, Xu Y, Pan X, et al. Global, regional, and national burden and trend of diabetes in 195
countries and territories: an analysis from 1990 to 2025. *Sci Rep*. 2020;10(1):14790.
doi:10.1038/s41598-020-71908-9
6.   Gregory GA, Robinson TIG, Linklater SE, et al. Global incidence, prevalence, and mortality of
type 1 diabetes in 2021 with projection to 2040: a modelling study. *Lancet Diabetes Endocrinol*.
2022;10(10):741-760. doi:10.1016/S2213-8587(22)00218-2
7.   Saeedi P, Petersohn I, Salpea P, et al. Global and regional diabetes prevalence estimates for
2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes
Atlas, 9th edition. *Diabetes Res Clin Pract*. 2019;157:107843. doi:10.1016/j.diabres.2019.107843
8.   Kautzky-Willer A, Harreiter J. Sex and gender differences in therapy of type 2 diabetes.
Diabetes Res Clin Pract. 2017;131:230-241. doi:10.1016/j.diabres.2017.07.012
9.   Kautzky-Willer A, Harreiter J, Pacini G. Sex and Gender Differences in Risk, Pathophysiology
and Complications of Type 2 Diabetes Mellitus. *Endocr Rev*. 2016;37(3):278-316. doi:10.1210/er.2015-
1137
10.   Oster RT, Johnson JA, Hemmelgarn BR, et al. Recent epidemiologic trends of diabetes mellitus
among status Aboriginal adults. *Can Med Assoc J*. 2011;183(12):E803-E808. doi:10.1503/cmaj.101882
11.   Zhou B, Lu Y, Hajifathalian K, et al. Worldwide trends in diabetes since 1980: a pooled analysis
of 751 population-based studies with 4·4 million participants. *The Lancet*. 2016;387(10027):1513-
1530. doi:10.1016/S0140-6736(16)00618-8
12.   Dyck R, Osgood N, Lin TH, Gao A, Stang MR. Epidemiology of diabetes mellitus among First
Nations and non-First Nations adults. *Can Med Assoc J*. 2010;182(3):249-256.
doi:10.1503/cmaj.090846
13.   Mauvais-Jarvis F, Bairey Merz N, Barnes PJ, et al. Sex and gender: modifiers of health, disease,
and medicine. *Lancet Lond Engl*. 2020;396(10250):565-582. doi:10.1016/S0140-6736(20)31561-0
14.   Heise L, Greene ME, Opper N, et al. Gender inequality and restrictive gender norms: framing
the challenges to health. *The Lancet*. 2019;393(10189):2440-2454. doi:10.1016/S0140-
6736(19)30652-X
15.   Rodgers JL, Jones J, Bolleddu SI, et al. Cardiovascular Risks Associated with Gender and Aging. *J
Cardiovasc Dev Dis*. 2019;6(2):19. doi:10.3390/jcdd6020019
16.   Gale EAM, Gillespie KM. Diabetes and gender. *Diabetologia*. 2001;44(1):3-15.
doi:10.1007/s001250051573
17.   Ferretti MT, Iulita MF, Cavedo E, et al. Sex differences in Alzheimer disease - the gateway to
precision medicine. *Nat Rev Neurol*. 2018;14(8):457-469. doi:10.1038/s41582-018-0032-9
18.   Whitacre CC. Sex differences in autoimmune disease. *Nat Immunol*. 2001;2(9):777-780.
doi:10.1038/ni0901-777

19.   DeMeo D, Ramagopalan S, Kavati A, et al. Women manifest more severe COPD symptoms
across the life course. *Int J Chron Obstruct Pulmon Dis*. 2018;Volume 13:3021-3029.
doi:10.2147/COPD.S160270
20.   Anand SS, Islam S, Rosengren A, et al. Risk factors for myocardial infarction in women and
men: insights from the INTERHEART study. *Eur Heart J*. 2008;29(7):932-940.
doi:10.1093/eurheartj/ehn018
21.   Chatterjee S, Peters SAE, Woodward M, et al. Type 2 Diabetes as a Risk Factor for Dementia in
Women Compared With Men: A Pooled Analysis of 2.3 Million People Comprising More Than 100,000
Cases of Dementia. *Diabetes Care*. 2016;39(2):300-307. doi:10.2337/dc15-1588
22.   Icks A, Trautner C, Haastert B, Berger M, Giani G. Blindness Due to Diabetes: Population-based
Age- and Sex-specific Incidence Rates. *Diabet Med*. 1997;14(7):571-575. doi:10.1002/(SICI)1096-
9136(199707)14:7<571::AID-DIA384>3.0.CO;2-Y
23.   Ohkuma T, Peters SAE, Woodward M. Sex differences in the association between diabetes and
cancer: a systematic review and meta-analysis of 121 cohorts including 20 million individuals and one
million events. *Diabetologia*. 2018;61(10):2140-2154. doi:10.1007/s00125-018-4664-5
24.   Huebschmann AG, Huxley RR, Kohrt WM, Zeitler P, Regensteiner JG, Reusch JEB. Sex
differences in the burden of type 2 diabetes and cardiovascular risk across the life course.
Diabetologia. 2019;62(10):1761-1772. doi:10.1007/s00125-019-4939-5
25.   Barrett-Connor EL. Why Is Diabetes Mellitus a Stronger Risk Factor for Fatal Ischemic Heart
Disease in Women Than in Men?: The Rancho Bernardo Study. *JAMA*. 1991;265(5):627.
doi:10.1001/jama.1991.03460050081025
26.   Amengual-Cladera E, Lladó I, Gianotti M, Proenza AM. Sex differences in the effect of high-fat
diet feeding on rat white adipose tissue mitochondrial function and insulin sensitivity. *Metabolism*.
2012;61(8):1108-1117. doi:10.1016/j.metabol.2011.12.016
27.   Macotela Y, Boucher J, Tran TT, Kahn CR. Sex and Depot Differences in Adipocyte Insulin
Sensitivity and Glucose Metabolism. *Diabetes*. 2009;58(4):803-812. doi:10.2337/db08-1054
28.   Tan SY, Lei X, Little HC, et al. CTRP12 ablation differentially affects energy expenditure, body
weight, and insulin sensitivity in male and female mice. *Am J Physiol-Endocrinol Metab*.
2020;319(1):E146-E162. doi:10.1152/ajpendo.00533.2019
29.   Vital P, Larrieta E, Hiriart M. Sexual dimorphism in insulin sensitivity and susceptibility to
develop diabetes in rats. *J Endocrinol*. 2006;190(2):425-432. doi:10.1677/joe.1.06596
30.   Horie I, Abiru N, Eto M, et al. Sex differences in insulin and glucagon responses for glucose
homeostasis in young healthy Japanese adults. *J Diabetes Investig*. 2018;9(6):1283-1287.
doi:10.1111/jdi.12829
31.   Li T, Jiao W, Li W, Li H. Sex effect on insulin secretion and mitochondrial function in pancreatic
beta cells of elderly Wistar rats. *Endocr Res*. 2016;41(3):167-179.
doi:10.3109/07435800.2015.1124437
32.   Tura A, Pacini G, Moro E, Vrbíková J, Bendlová B, Kautzky-Willer A. Sex- and age-related
differences of metabolic parameters in impaired glucose metabolism and type 2 diabetes compared
to normal glucose tolerance. *Diabetes Res Clin Pract*. 2018;146:67-75.
doi:10.1016/j.diabres.2018.09.019
33.   Basu R, Dalla Man C, Campioni M, et al. Effects of Age and Sex on Postprandial Glucose
Metabolism. *Diabetes*. 2006;55(7):2001-2014. doi:10.2337/db05-1692
34.   Kautzky-Willer A, Brazzale AR, Moro E, et al. Influence of Increasing BMI on Insulin Sensitivity
and Secretion in Normotolerant Men and Women of a Wide Age Span. *Obesity*. 2012;20(10):1966-
1973. doi:10.1038/oby.2011.384

35.   Mank JE, Rideout EJ. Developmental mechanisms of sex differences: from cells to organisms.
Development. 2021;148(19):dev199750. doi:10.1242/dev.199750
36.   Millington JW, Biswas P, Chao C, et al. A low-sugar diet enhances *Drosophila* body size in males
and females via sex-specific mechanisms. *Development*. 2022;149(6):dev200491.
doi:10.1242/dev.200491
37.   Nuutila P, Knuuti MJ, Mäki M, et al. Gender and Insulin Sensitivity in the Heart and in Skeletal
Muscles: Studies Using Positron Emission Tomography. *Diabetes*. 1995;44(1):31-36.
doi:10.2337/diab.44.1.31
38.   Færch K, Borch-Johnsen K, Vaag A, Jørgensen T, Witte DR. Sex differences in glucose levels: a
consequence of physiology or methodological convenience? The Inter99 study. *Diabetologia*.
2010;53(5):858-865. doi:10.1007/s00125-010-1673-4
39.   Borissova AM, Tankova T, Kirilov G, Koev D. Gender-dependent effect of ageing on peripheral
insulin action: Effect of Ageing on Peripheral Insulin Action. *Int J Clin Pract*. 2005;59(4):422-426.
doi:10.1111/j.1368-5031.2005.00209.x
40.   Geer EB, Shen W. Gender differences in insulin resistance, body composition, and energy
balance. *Gend Med*. 2009;6:60-75. doi:10.1016/j.genm.2009.02.002
41.   Gannon M, Kulkarni RN, Tse HM, Mauvais-Jarvis F. Sex differences underlying pancreatic islet
biology and its dysfunction. *Mol Metab*. 2018;15:82-91. doi:10.1016/j.molmet.2018.05.017
42.   Peters SAE, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart
disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including
858,507 individuals and 28,203 coronary events. *Diabetologia*. 2014;57(8):1542-1551.
doi:10.1007/s00125-014-3260-6
43.   Rideout EJ, Narsaiya MS, Grewal SS. The Sex Determination Gene transformer Regulates Male-
Female Differences in Drosophila Body Size. Kopp A, ed. *PLOS Genet*. 2015;11(12):e1005683.
doi:10.1371/journal.pgen.1005683
44.   Millington JW, Brownrigg GP, Chao C, et al. Female-biased upregulation of insulin pathway
activity mediates the sex difference in Drosophila body size plasticity. *eLife*. 2021;10:e58341.
doi:10.7554/eLife.58341
45.   Millington JW, Brownrigg GP, Basner-Collins PJ, Sun Z, Rideout EJ. Genetic manipulation of
insulin/insulin-like growth factor signaling pathway activity has sex-biased effects on *Drosophila* body
size. Lipshitz HDa, ed. *G3 GenesGenomesGenetics*. 2021;11(3):jkaa067. doi:10.1093/g3journal/jkaa067
46.   Stancill JS, Osipovich AB, Cartailler JP, Magnuson MA. Transgene-associated human growth
hormone expression in pancreatic β-cells impairs identification of sex-based gene expression
differences. *Am J Physiol-Endocrinol Metab*. 2019;316(2):E196-E209.
doi:10.1152/ajpendo.00229.2018
47.   Marchese E, Rodeghier C, Monson RS, et al. Enumerating β-Cells in Whole Human Islets: Sex
Differences and Associations With Clinical Outcomes After Islet Transplantation. *Diabetes Care*.
2015;38(11):e176-e177. doi:10.2337/dc15-0723
48.   Arrojo E Drigo R, Erikson G, Tyagi S, et al. *Aging of Human Endocrine Pancreatic Cell Types Is
Heterogeneous and Sex-Specific*. Physiology; 2019. doi:10.1101/729541
49.   Saisho Y. β-cell dysfunction: Its critical role in prevention and management of type 2 diabetes.
World J Diabetes. 2015;6(1):109. doi:10.4239/wjd.v6.i1.109
50.   Segerstolpe Å, Palasantza A, Eliasson P, et al. Single-Cell Transcriptome Profiling of Human
Pancreatic Islets in Health and Type 2 Diabetes. *Cell Metab*. 2016;24(4):593-607.
doi:10.1016/j.cmet.2016.08.020

51.   Xin Y, Kim J, Okamoto H, et al. RNA Sequencing of Single Human Islet Cells Reveals Type 2
Diabetes Genes. *Cell Metab*. 2016;24(4):608-615. doi:10.1016/j.cmet.2016.08.018
52.   Lawlor N, George J, Bolisetty M, et al. Single-cell transcriptomes identify human islet cell
signatures and reveal cell-type–specific expression changes in type 2 diabetes. *Genome Res*.
2017;27(2):208-222. doi:10.1101/gr.212720.116
53.   Baron M, Veres A, Wolock SL, et al. A Single-Cell Transcriptomic Map of the Human and Mouse
Pancreas Reveals Inter- and Intra-cell Population Structure. *Cell Syst*. 2016;3(4):346-360.e4.
doi:10.1016/j.cels.2016.08.011
54.   Muraro MJ, Dharmadhikari G, Grün D, et al. A Single-Cell Transcriptome Atlas of the Human
Pancreas. *Cell Syst*. 2016;3(4):385-394.e3. doi:10.1016/j.cels.2016.09.002
55.   Deng S, Vatamaniuk M, Huang X, et al. Structural and Functional Abnormalities in the Islets
Isolated From Type 2 Diabetic Subjects. *Diabetes*. 2004;53(3):624-632. doi:10.2337/diabetes.53.3.624
56.   Roep BO, Thomaidou S, Van Tienhoven R, Zaldumbide A. Type 1 diabetes mellitus as a disease
of the β-cell (do not blame the immune system?). *Nat Rev Endocrinol*. 2021;17(3):150-161.
doi:10.1038/s41574-020-00443-4
57.   Sims EK, Mirmira RG, Evans-Molina C. The role of beta-cell dysfunction in early type 1
diabetes. *Curr Opin Endocrinol Diabetes Obes*. 2020;27(4):215-224.
doi:10.1097/MED.0000000000000548
58.   Day S, Wu W, Mason R, Rochon PA. Measuring the data gap: inclusion of sex and gender
reporting in diabetes research. *Res Integr Peer Rev*. 2019;4(1):9. doi:10.1186/s41073-019-0068-4
59.   Mazure CM, Jones DP. Twenty years and still counting: including women as participants and
studying sex and gender in biomedical research. *BMC Womens Health*. 2015;15(1):94.
doi:10.1186/s12905-015-0251-9
60.   Beery AK, Zucker I. Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev*.
2011;35(3):565-572. doi:10.1016/j.neubiorev.2010.07.002
61.   Arnold AP. Promoting the understanding of sex differences to enhance equity and excellence
in biomedical science. *Biol Sex Differ*. 2010;1(1):1. doi:10.1186/2042-6410-1-1
62.   Clayton JA, Collins FS. Policy: NIH to balance sex in cell and animal studies. *Nature*.
2014;509(7500):282-283. doi:10.1038/509282a
63.   Viergever RF, Hendriks TCC. The 10 largest public and philanthropic funders of health research
in the world: what they fund and how they distribute their funds. *Health Res Policy Syst*.
2016;14(1):12. doi:10.1186/s12961-015-0074-z
64.   Collins FS, Tabak LA. Policy: NIH plans to enhance reproducibility. *Nature*. 2014;505(7485):612-
613. doi:10.1038/505612a
65.   NIH Policy. Consideration of Sex as a Biological Variable in NIH-funded Research.
66.   *Women and Health Research: Ethical and Legal Issues of Including Women in Clinical Studies,*
Volume 1. National Academies Press; 1994:2304. doi:10.17226/2304
67.   R Core Team (2023). _R: A Language and Environment for Statistical   Computing.
https://www.R-project.org
68.   Zucker I, Beery AK. Males still dominate animal studies. *Nature*. 2010;465(7299):690-690.
doi:10.1038/465690a
69.   Karp NA, Reavey N. Sex bias in preclinical research and an exploration of how to change the
status quo. *Br J Pharmacol*. 2019;176(21):4107-4118. doi:10.1111/bph.14539
70.   Tannenbaum C, Schwarz JM, Clayton JA, De Vries GJ, Sullivan C. Evaluating sex as a biological
variable in preclinical research: the devil in the details. *Biol Sex Differ*. 2016;7(1):13, s13293-016-0066-
x. doi:liss

71.   Poitout V, Satin LS, Kahn SE, et al. A Call for Improved Reporting of Human Islet Characteristics
in Research Articles. *Diabetes*. 2019;68(2):239-240. doi:10.2337/dbi18-0055
72.   Hart NJ, Powers AC. Use of human islets to understand islet biology and diabetes: progress,
challenges and suggestions. *Diabetologia*. 2019;62(2):212-222. doi:10.1007/s00125-018-4772-2
73.   Policy: Reporting on Human Islets. Published online April 2019.
https://diabetesjournals.org/journals/pages/human-islets
74.   Rechlin RK, Splinter TFL, Hodges TE, Albert AY, Galea LAM. An analysis of neuroscience and
psychiatry papers published from 2009 and 2019 outlines opportunities for increasing discovery of sex
differences. *Nat Commun*. 2022;13(1):2137. doi:10.1038/s41467-022-29903-3
75.   Stewart AN, MacLean SM, Stromberg AJ, et al. Considerations for Studying Sex as a Biological
Variable in Spinal Cord Injury. *Front Neurol*. 2020;11:802. doi:10.3389/fneur.2020.00802
76.   Choleris E, Galea LAM, Sohrabji F, Frick KM. Sex differences in the brain: Implications for
behavioral and biomedical research. *Neurosci Biobehav Rev*. 2018;85:126-145.
doi:10.1016/j.neubiorev.2017.07.005
77.   Duffy KA, Epperson CN. Evaluating the evidence for sex differences: a scoping review of human
neuroimaging in psychopharmacology research. *Neuropsychopharmacology*. 2022;47(2):430-443.
doi:10.1038/s41386-021-01162-8
78.   Liu G, Li Y, Zhang T, et al. Single-cell RNA Sequencing Reveals Sexually Dimorphic
Transcriptome and Type 2 Diabetes Genes in Mouse Islet β Cells. *Genomics Proteomics Bioinformatics*.
2021;19(3):408-422. doi:10.1016/j.gpb.2021.07.004
79.   Tannenbaum C, Ellis RP, Eyssel F, Zou J, Schiebinger L. Sex and gender analysis improves
science and engineering. *Nature*. 2019;575(7781):137-146. doi:10.1038/s41586-019-1657-6

FIGURE LEGENDS

Figure 1 – Assessing the inclusion of biological sex as a variable in *Diabetes*

manuscripts published over time. (A) Schematic representation of the workflow for

manuscript selection in our study. (B) Average score of studies published in 1999, 2009, and

2019. (*** $p<0.001$; generalized linear model with Poisson distribution; error bars indicate

SEM) (C) Proportion of studies assigned scores between 0-6 in 1999, 2009, and 2019.

Dashed lines represent a statistically insignificant change and solid lines represent a

statistically significant change (generalized linear model with binomial distribution). (D)

Proportion of single-sex studies that used male animals in 1999, 2009, and 2019 (** $p<0.01$;

generalized linear model with binomial distribution).

Figure 2 – Evaluating the inclusion of biological sex as a variable in *Diabetes*

manuscripts across clinical and biomedical disciplines. (A) Average scores assigned to

studies published in biomedical and clinical disciplines in 1999, 2009, and 2019 (*** $p<0.001$;

generalized linear model with Poisson distribution; error bars indicate SEM). (B) Proportion of

biomedical studies assigned scores between 0-6 in 1999, 2009, and 2019. (C) Proportion of

clinical studies assigned scores between 0-6 in 1999, 2009, and 2019.

Figure 3 – Comparing the inclusion of biological sex as a variable between *Diabetes*

manuscripts published across diabetes subject areas. (A) Average scores assigned to

studies published in different diabetes subject areas in 2009 and 2019. (ns not significant;

generalized linear model with Poisson distribution; error bars indicate SEM) (B) Proportion of

studies assigned scores between 0-6 in 2009 and 2019 in the 'Complications' subject area.

(C) Proportion of studies assigned scores between 0-6 in 2009 and 2019 in the

'Genetics/Omics' subject area. No studies in either year were assigned a score of 2 or 5. (D)

Proportion of studies assigned scores between 0-6 in 2009 and 2019 in the 'Immunology and

Transplantation' subject area. No studies in either year were assigned a score of 5 or 6. (E)

Proportion of studies assigned scores between 0-6 in 2009 and 2019 in the 'Islet' subject

area. No studies in either year were assigned a score of 5 or 6. (F) Proportion of studies

assigned scores between 0-6 in 2009 and 2019 in the Obesity subject area. (G) Proportion of

studies assigned scores between 0-6 in 2009 and 2019 in the Metabolism subject area. (H)

Proportion of studies assigned scores between 0-6 in 2009 and 2019 in the Pathophysiology

subject area. No studies in either year were assigned a score of 5 or 6.

Figure 4 – Evaluating the effect of funding source on the inclusion of biological sex as

a variable in diabetes research (A) Average scores assigned to studies that declared NIH

funding compared with studies that did not declare NIH funding in 1999, 2009, and 2019 (ns

not significant; generalized linear model with Poisson distribution; error bars indicate SEM).

(B) Proportion of NIH-funded studies assigned scores between 0-6 in 1999, 2009, and 2019.

(C) Proportion of non-NIH-funded studies assigned scores between 0-6 in 1999, 2009, and

2019. (D) Scores assigned to studies that declared NIH funding in the biomedical and clinical

disciplines in 1999, 2009, and 2019 (ns not significant; generalized linear model with Poisson

distribution; error bars indicate SEM).

Figure 5 – Evaluating the effect of journal policies on inclusion of sex as a biological

variable in *Diabetes* manuscripts. (A) Average scores assigned to studies in the 'Islet' and

'Metabolism' subject areas between 2018 and 2020 (before and after the 2019 policy change,

respectively) (** p<0.01; ns not significant; generalized linear model with Poisson distribution;

error bars indicate SEM). (B) Proportion of studies in the 'Islet' subject area assigned scores

between 0-6 in 2018 and 2020.

744 **TABLE 1 – Scoring system to assess the consideration of biological sex as a variable**
745

Score	Meaning
0	No mention of sex
1	One sex used, without explicit justification
2	One sex used, with explicit justification
3	Both sexes included; data not analyzed separately by sex or data not provided
4	Both sexes included; fewer than half of results analyzed by sex (<50%)
5	Both sexes included; more than half of results analyzed by sex (>50%)
6	Both sexes included; all data analyzed and reported by sex

746
747

748 **TABLE 2 – Number of papers in each diabetes subject area**

Diabetes subject area	Number of papers in 2009	Number of papers in 2019
Complications	41	37
Genetics / Omics (Genetics - 2009) (Genetics/Genomes/Proteomics/Metabolomics - 2019)	54	25
Immunology and Transplantation	24	11
Islet studies	37	25
Obesity studies	21	18
Metabolism	81	37
Pathophysiology	27	12
New Methodology / Technology (New Methodologies and Databases - 2009) (Technological Advances - 2019)	3	2
Pharmacology	8	9
Signal Transduction	16	2

749
750

751
752

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

A Average Islet study score before and after policy change

Islet studies   Metabolism

B Proportion of Islet studies scores before and after policy change

0   2   4   6
1   3   5