Full title: High-fat diet-induced obesity increases susceptibility to endogenous and ethyl

carbamate-induced somatic mutagenesis

Short title: High-fat diet and mutagenesis

Rebecca Lichtler[1,#a,*], Mark J. Wilson[1#b], Jeffrey K. Wickliffe[1,#c]

[1]Department of Environmental Health Sciences, School of Public Health and Tropical

Medicine, Tulane University, New Orleans, Louisiana, United States of America

[#a] Current Address: Department of Biological Sciences, College of Sciences, North

Carolina State University, Raleigh, North Carolina, United States of America

[#b] Current Address: Chemical Insights Research Institute, Underwriters Laboratories

Research Institutes, Marietta, Georgia, United States of America

[#c] Current Address: Department of Environmental Health Sciences, School of Public

Health, University of Alabama at Birmingham, Birmingham, Alabama, United States of

America

*Corresponding author, email: rclichtl@ncsu.edu (RL)

Author contributions: RL, MJW, and JKW all contributed to the conceptualization,

investigation, methodology, and data analysis of the described experiments. RL drafted

and MJW and JKW reviewed and edited the manuscript.

Competing interests: The authors declare they have no actual or potential competing

financial interests.

Abstract

Obesity and dietary fat consumption are associated with increased risk for certain

cancers. We conducted a set of experiments designed to investigate both the inherent

mutagenicity of high-fat diet (HFD) and the effect of HFD exposure on sensitivity to ethyl

carbamate, a known mutagen. Greater body mass in females was associated with

higher frequency of Pig-a mutant erythrocytes and HFD-induced obese female mice

exhibited increased sensitivity to ethyl carbamate-induced mutagenesis compared to

control mice. Together, these findings indicate that HFD exposure induces both

endogenously generated changes related to mutagenesis and interacts with the

mutagenic effects of ethyl carbamate.

1. Introduction

Obesity and overweight are highly prevalent in the United States and are

associated with a number of conditions, including certain cancers. Of all new cancer

cases diagnosed in 2015, 55% of those diagnosed in women and 24% of those

diagnosed in men can be linked to obesity[1]. In fact, incidence of obesity-related

cancers rose 7% between 2005 and 2014, while non-obesity related cancers declined

by 13%, and the fastest growing obesity-related cancer type during that period was liver

cancer, which rose by 29% [1,2]. Women, in particular, appear to be at increasing risk

for obesity related cancers. Among women, rates of obesity-related cancers rose

between 2001 and 2018, while cancers due to other risk factors declined or remained

steady during the same time period [3]. Women with higher body mass index (BMI)

were less likely to survive liver cancer following treatment than those with BMI

considered to be within the normal range [4]. Along with evidence that higher BMI or

adiposity increases cancer risk, dietary composition appears to play a role as well. High

dietary fat consumption was associated with a 50% increase in the likelihood of having

breast cancer in women [5]. Among men and women, higher consumption of dietary fat

was associated with a 60% increase in colorectal cancer[6]. A meta-analysis of

prospective cohort studies suggest consuming a diet high in fat, and saturated fat in

particular, was associated with a 7-14% increase in lung cancer risk [7]. Another meta-

analysis of prospective cohort studies that tracked liver cancer incidence found a 34%

increased risk of liver cancer among those with the highest saturated fat consumption

[8]. A recent study of patients with clonal hematopoiesis of indeterminate potential

(CHIP), a phenomenon linked to hematologic cancers where blood cells accumulate

mutations, uncovered obesity as risk factor for this condition[9].

The pathway from a genotoxic exposure, either dietary or chemical, to cancer

development involves several stages, the first of which is initiation, or the establishment

of a somatic gene mutation [10]. Despite the abundance of population-based

association studies that link dietary fat consumption and cancer risk, very few controlled

experimental studies have been conducted to evaluate the mutagenicity of HFD alone.

Additionally, there is a dearth of studies that investigate the effects of co-exposure to

HFD and environmental mutagens[11]. We have previously shown that high-fat diet

(HFD) exposure in male C57BL/6J mice leads to higher frequency of spontaneous

somatic mutations using the Pig-a gene mutation assay [12]. The Pig-a gene mutation

assay was developed to measure the frequency of somatic gene mutations in peripheral

red blood cells, by exploiting the glycosylphosphatidylinositol (GPI) cell surface anchor

protein, found ubiquitously on red blood cells[13–16]. GPI anchor biosynthesis is

dependent on a functional *Pig-a* gene, located on the X chromosome, meaning a single

mutation can result in a GPI-deficient cell phenotype, and the frequency of GPI-deficient

red blood cells can be enumerated using flow cytometric analysis[17].

Here we present the results from a set of experiments investigating the

mutagenicity of HFD-induced obesity including both male and female mice for the first

time. We hypothesized that HFD-induced obesity will increase the risk for both

endogenous, spontaneous somatic mutagenesis and exogenous mutagen-induced

mutagenesis. To test this hypothesis, we conducted two experiments, one designed to

determine the effect of HFD alone (60% calories from fat) over time in both males and

females on frequency of spontaneous somatic mutations using the Pig-a gene mutation

assay. We also designed a two-factorial experiment whereby mice who had undergone

long-term HFD exposure were challenged with oral ethyl carbamate and assessed for

Pig-a gene mutation frequency after a sufficient mutation manifestation period. Ethyl

carbamate is a known mutagen and carcinogen that is produced in food and beverage

fermentation and has been shown to induce somatic gene mutations using the Pig-a

gene mutation assay [16,18]. While the general public is not at high risk, those that

regularly consume stone fruit brandy or fermented soy products may have a lifetime

tumor risk of 0.01%, 100-times the risk that is generally considered acceptable [19–21].

Ethyl carbamate is converted into more toxic epoxide-containing compounds by

esterases and the cytochrome P450 superfamily of phase I metabolic enzymes, several

members of which are known to be perturbed in the context of obesity [22]. Together,

this set of experiments explores both the endogenous and exogenous risk of somatic

mutagenesis, one of the key components of the pathway to carcinogenicity, in the

context of diet-induced obesity.

2. Methods

2.1 General animal care and analysis

Mice were reared at the Tulane University Downtown Vivarium on a 12-hour

light/dark cycle. Research staff supplied C57BL/6J mice with HFD containing 60%

calories from fat from Research Diets, Inc. (D12492) or with a low-fat diet (LFD)

containing 10% calories from fat (D12450J), *ad libitum*. The LFD formulation serves as

the most appropriate control to the HFD formulation because it is source-matched and

contains equivalent sucrose content. Water and veterinary care were provided by

vivarium staff. Unless otherwise stated, Prism software (GraphPad; San Diego, CA,

USA) was used to conduct statistical analysis and to generate figures. Unless otherwise

indicated, data in figures is presented as the arithmetic mean $\pm$ standard error with

individual data points. Statistical significance was determined using a threshold of

$\alpha=0.05$, and results of statistical mean comparison tests are presented as the t statistic

with degrees of freedom in subscript, followed by the corresponding p value.

2.2 Biochemical assays

To ensure that long-term HFD exposure had generated a physiologically obese

phenotype in our model, plasma leptin, alanine amino transferase (ALT), and aspartate

aminotransferase (AST), and total liver triglycerides were quantified using commercially

available kits within the vehicle-control (deionized water) treated mice from the ethyl

carbamate challenge experiment. Enzymatic activity of AST and ALT was measured as

the synthesis of glutamate or pyruvate, respectively, that occurred per minute of

incubation with each substrate at 37°C by Aspartate Aminotransferase Activity Assay Kit

(Abcam ab105135) and the Alanine Transaminase Activity Assay Kit (Abcam

ab105134). Plasma leptin was measured with the Mouse/Rat Leptin Quantikine ELISA

Kit (R&D Systems, MOB00). Liver triglycerides were measured with the Triglyceride

Quantification Assay kit (Abcam, ab65336), following tissue lysis and isolation of lipids

with homogenization and boiling in a solution of 5% NP-40 in deionized water. All

colorimetric and fluorescent results were analyzed using a Tecan Infinite 200 Pro and i-

control software (V1.9, Mannedorf, Switzerland).

2.3 HFD-only experiment

2.3.1 Animal care and sample collection

This experiment was conducted under the approval of Tulane University IACUC

protocol 546. Mice were maintained on HFD or LFD conditions from 8 weeks old until 52

weeks old. Each dietary group consisted of 12 males and 12 females. Blood was taken

at three time points, when mice were 17, 33 and 52 weeks of age. This sampling

method was accomplished through repeated submandibular venipuncture under

isoflurane anesthetization [23].

2.3.2 Erythrocyte Pig-a mutant frequency analysis

Each blood sample was immediately transferred to a K_2EDTA-coated

microtainers (BD, 365974) and kept on ice. Samples were packed into Whirl-Pak® bags

and shipped to Litron Laboratories (Rochester, NY). Pig-a mutant frequency analysis

was carried out by Litron research staff following *In Vivo* MutaFlow® kit protocols

[12,16,24].

2.3.4 Statistical analysis

Body mass and mutant frequency were analyzed using mixed effects analysis of variance (ANOVA) for repeated measures with Bonferroni's post hoc multiple comparisons test performed at each time point between LFD and HFD subjects. Simple linear regression analysis of the relationship between body mass and Pig-a mutant frequency was also conducted at each time point, and p value for slope deviation from zero and R^2 correlation coefficient were reported.

2.4 Ethyl carbamate challenge experiment

2.4.1 Animal care and sample collection

This experiment was conducted under approved Tulane University IACUC protocol 36-105. 48 males and 48 females were reared on HFD or LFD from 8 weeks old until they reached 42 weeks old, when the experiment was terminated. Nine weeks prior to termination, at 33 weeks, mice were divided into groups of 4-6, and exposed to 0, 100, 200, or 400 mg/kg/day of ethyl carbamate (CAS # 51-79-6; Sigma, U2500, Lot # WXBC3505V) in deionized water via oral gavage for 3 consecutive days following the procedure outlined by Labash et al. [16]. At 42 weeks, to maximize yield and quality of blood samples, mice were weighed and then anesthetized with no more than 5% isoflurane (v/v) and 100% oxygen saturation at a 1 L/min flow rate. The sternal cavity was opened for full visualization of the heart, and blood was collected from the left ventricle with a heparin-coated 22G needle and 1 mL syringe. The blood was then transferred to K_2EDTA-coated microtainers (BD, 365974) to prevent coagulation. Immediately after collection, 60 µL of each sample was transferred to a tube containing

an anticoagulant solution provided by Litron Laboratories and kept on ice for further

processing following the Mouse MutaFlow[PLUS] Kit manual (Litron Laboratories,

v170925).

2.4.2 Pig-a mutant frequency analysis

Pig-a mutant frequency analysis was conducted at Tulane University using a

modified MutaFlow method. In summary, white blood cells and platelets were removed

with Lympholyte®-Mammal (CedarLane, CL5110) and remaining reticulocytes,

erythrocytes, and platelets were labeled with Anti-CD24-PE (Miltenyi, 130-110-688) and

Anti-CD61-PE (Miltenyi, 130-102-628) antibodies and anti-PE magnetic microbeads

(Miltenyi, 130-048-801). A small portion of each sample was retained for cell density

determination and the remainder was enriched for mutant cells via immunomagnetic

separation on LS columns (Miltenyi, 130-042-401) within a magnetic field. SYTO 13

nucleic acid dye (Invitrogen, S7575) was added to discriminate mature erythrocytes

from immature reticulocytes and remaining lymphocytes. Total mature erythrocyte and

Pig-a mutant erythrocyte counts were enumerated via the Miltenyi MACSQuant

Analyzer 10 flow cytometer. Pig-a mutant mature erythrocytes were classified as cells

exhibiting absence of the PE fluorescence signal and intracellular SYTO 13, indicating a

lack of both CD24 surface markers and nucleic acids. Mutant frequency was calculated

as the number of mutant mature erythrocytes per 10^6 total mature erythrocytes.

2.4.3 Statistical analysis

Pig-a mutant frequency was analyzed using two-way ANOVA with diet and ethyl

carbamate dose as the two independent factors, with Bonferroni's post hoc comparison

184 test between LFD and HFD groups within each dose group. A dose-response curve was

185 plotted to compare the effect of ethyl carbamate dose on Pig-a mutant frequency

186 between LFD and HFD groups, with slope comparison and R^2 calculation as well as an

187 F test to compare dose-response curve slopes between HFD and LFD groups. Males

188 and females were analyzed separately.

190 **3. Results**

191  **3.1 Biochemical markers of obesity**

192  HFD exposure in the untreated group of the ethyl carbamate challenge

193 experiment (vehicle controls) led to elevated liver triglycerides in females (t_8=2.409,

194 p=0.0425) and males (t_8=2.942, p=0.0186) and elevated plasma leptin in females

195 (t_{10}=7.187, p<0.0001) and males (t_{10}=4.717, p=0.00082) (Fig 1A-B). Activity of liver

196 enzymes AST (females: t_6=2.857, p=0.0289; males: t_6=3.841, p=0.0086) and ALT

197 (females: t_6=3.354, p=0.0153; males: t_6=5.574, p=0.0014) were elevated in plasma, as

198 well (Fig 1C-D). These biochemical findings indicate HFD exposure until 42 weeks of

199 age was sufficient to generate increased body mass, fatty liver, and elevated circulating

200 levels of liver enzymes and a circulating adipokine, all characteristic features of HFD-

201 induced obesity.

Fig 1. Effect of HFD-exposure on biochemical markers of obesity at 42 weeks-of-age. (A) Total liver triglycerides, (B) plasma leptin, (C) plasma AST activity, and (D) plasma ALT activity. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001 per two-tailed Student's t test between LFD and HFD groups, conducted within males and females separately (n=4-6).

3.2 HFD-only experiment

To assess the mutagenicity of long-term HFD exposure, the erythrocyte Pig-a mutation assay was conducted with sequential blood sampling at 17, 33 and 52 weeks old. By week 52, one LFD female was euthanized due to injury, and two HFD males died due to undetermined causes. Both HFD males and HFD females were significantly heavier than their LFD counterparts at each sampling timepoint. On average, HFD

females were 6.98 ($t_{17.11}$=6.334, p<0.0001), 19.99 ($t_{15.12}$=8.281, p<0.0001), and 32.10 ($t_{14.92}$=13.27, p<0.0001) grams heavier than LFD females at 17, 33, and 52 weeks, respectively (Fig 2A). On average, HFD males were 10.62 ($t_{14.24}$=8.151, p<0.0001), 14.83 ($t_{11.34}$=11.18, p<0.0001), and 15.90 ($t_{11.07}$=9.494, p<0.0001) grams heavier than LFD males at 17, 33, and 52 weeks, respectively (Fig 2B).

Fig 2. Effect of diet on body mass at each sampling timepoint. Male (A) and female (B) body mass in LFD and HFD groups at 17, 33 and 52 weeks of age.

**** p<0.0001 per mixed effects analysis (for repeated measures) with Bonferroni's multiple comparisons test, between LFD and HFD groups at each timepoint, conducted in males and females independently (n=10-12).

One extreme outlier LFD female was excluded from all Pig-a mutant frequency analyses due to very high frequency of Pig-a mutant erythrocytes detected at 33 and 52 weeks (>4 standard deviations). Blood samples that did not pass quality control parameters determined by the operators at Litron Laboratories were not reported. Both raw and fold-difference erythrocyte mutant frequency values were reported to normalize

interexperiment variability. Despite a trend of an approximately 1.5-2-fold elevation,

there was no significant difference in mutant frequency at any sampling timepoint

between LFD and HFD females (Fig 3A-B). Because of the variability in body mass

observed in response to HFD and the stochastic nature of somatic mutagenesis, simple

linear regression was performed to determine a dose-response curve for the association

between body mass and mutant frequency at each sampling time point. In females,

simple linear regression for the association between body mass and erythrocyte mutant

frequency per million total erythrocytes indicated the slope was not significantly different

from zero at 17 weeks, indicating the was no association between body mass and

mutation frequency at the first sampling timepoint (Fig 3C, Table 1). By 33 weeks, there

was significant positive association between body mass and mutant frequency, with an

R^2 of 0.2004 and significantly non-zero slope (p=0.0322), indicating mice with a higher

body mass were more likely to experience more frequent somatic mutations (Fig 3D,

Table 1). By 52 weeks the association between body mass and mutant frequency

remained positive, with an R^2 of 0.2518 and a significant slope (p=0.0286) (Fig 3E,

Table 1), indicating higher body mass was associated with higher incidence of somatic

mutations, and this association grew stronger with age in females. Among males, while

there was a similar trend towards higher (~2-fold) mutant frequency among the HFD

group, there was no significant differences between the LFD and HFD groups at any

time point (Fig 4A-B), and there was no significant correlation between body mass and

mutant frequency at any sampling timepoint (Fig 4C-E, Table 2).

Figure 3. Effects of diet and body mass on spontaneous mutation frequency in erythrocytes at 17, 33 and 52 weeks of age in females. (A) Mutant frequency per million erythrocytes, and (B) Mutant frequency fold-difference between LFD and HFD

groups at each time point, per mixed effects analysis (for repeated measures) with

Bonferroni's multiple comparisons test, between LFD and HFD group sat each timepoint

(n=10-12). (C) Correlation between body mass and erythrocyte mutant frequency per

million cells at 17 weeks. (D) Correlation between body mass and erythrocyte mutant

frequency per million cells at 33 weeks. (E) Correlation between body mass and

erythrocyte mutant frequency per million cells at 52 weeks.

Table 1. Simple linear regression correlating body mass and Pig-a mutant

frequency in females after 17, 33 and 52 weeks of dietary exposure.

Week	Equation	R^2	Deviation from zero p value
17	y=0.01262x+0.01830	0.02438	0.5232
33	y=0.008555x-0.01226	0.2004	**0.0322**
52	y=0.007024x+0.06099	0.2518	**0.0286**

265 Bol

266 d

267 typ

268 e

269 indicates the slope deviates significantly from zero (α=0.05).

Fig 4. Effects of diet and body mass on spontaneous mutation frequency in erythrocytes at 17, 33 and 52 weeks of age in males. (A) Mutant frequency per million erythrocytes, and (B) Mutant frequency fold-difference between LFD and HFD groups at each time point, per mixed effects analysis (for repeated measures) with

Bonferroni's multiple comparisons test, between LFD and HFD group sat each timepoint (n=10-12). (C) Correlation between body mass and erythrocyte mutant frequency per million cells at 17 weeks in males. (D) Correlation between body mass and erythrocyte mutant frequency per million cells at 33 weeks in males. (E) Correlation between body mass and erythrocyte mutant frequency per million cells at 52 weeks in males.

Table 2. Simple linear regression correlating body mass and Pig-a mutant frequency in males after 17, 33 and 52 weeks of dietary exposure.

Age (weeks)	Equation	R^2	Deviation from zero p value
17	y=0.01836x-0.4354	0.1339	0.0939
33	y=0.05517x-1.743	0.1504	0.1118
52	y=0.01093x-0.1897	0.02333	0.5086

Ethyl carbamate challenge experiment

To determine the effect of HFD on sensitivity to a known mutagen, mice were reared on HFD or LFD starting at 7 weeks of age, exposed to ethyl carbamate at 33 weeks, and the experiment was terminated following a 9-week mutation manifestation period. Dietary conditions were maintained for the entirety of the experiment, and 93 of 96 total mice survived until termination at 42 weeks old. One female HFD mouse was sacrificed at 17 weeks due to excessive pruritus with unhealed dermal lesions and lack of weight gain. One male HFD mouse was sacrificed at 22 weeks due to sudden rapid weight loss, several large obstructive growths were observed in the colon during necropsy. One female LFD mouse was cannibalized at 29 weeks. All HFD groups were significantly heavier than their corresponding LFD controls and there was no significant

effect of ethyl carbamate dose group on final body mass, (p<0.0001 for all LFD to HFD

comparisons) (Fig 5).

Fig 5. The final body mass among all dietary and ethyl carbamate treatment

groups. (A) final body mass in males, and (B) final body mass in males. **** p <0.0001

per two-way ANOVA with Bonferroni-corrected comparison between all groups (only

those comparisons within dose groups are shown here).

HFD exposure in females led to increased susceptibility to somatic mutations

induced by exposure to 400 mg/kg/day ethyl carbamate (t_{38}=4.67, p=0.0001; Fig 6A).

The difference was on average 3.8-fold higher in the HFD group than the LFD group,

within those who received 400 mg/kg/day ethyl carbamate (t_{38}=4.67, p=0.0001; Fig 6B).

To determine if HFD females were more sensitive than LFD females to the same ethyl

carbamate dose regimen, dose response curves were plotted with regression analysis

between ethyl carbamate dose and Pig-a mutant frequency response. Simple linear

regression indicated the dose-response slope does not significantly differ from zero in

the LFD group (p=0.3872) and there was no correlation between ethyl carbamate dose

and mutant frequency response (R^2=0.0376; Fig 6C, Table 3). The HFD regression

curve is highly significantly different from zero (p<0.0001), and there is a strong

correlation between ethyl carbamate dose and mutant frequency (R^2 =0.5924; Fig 6C,

Table 3). The slope of the HFD curve is significantly higher than the LFD curve,

indicating HFD exposure significantly increases the susceptibility of female mice to ethyl

carbamate induced somatic mutagenesis, and that the two dietary treatment groups

display different dose-response curves in response to ethyl carbamate (p=0.0003; Table

3).

Fig 6. Effects of diet and body mass on mutant frequency in erythrocytes of female mice exposed to 0, 100, 200, and 400 mg/kg/day ethyl carbamate. (A) Mutant frequency per million erythrocytes in each ethyl carbamate dose group. (B) Mutant frequency fold-difference between LFD and HFD groups in each ethyl carbamate dose group. (C) Dose-response curves for ethyl carbamate dose and erythrocyte mutant frequency with simple linear regression analysis, points represent

330　mean ±standard error.　　*** p<0.001 per two-way ANOVA with Bonferroni's multiple

331　comparison test between LFD and HFD groups (n=4-6).

333　**Table 3. Simple linear regression dose-response analysis correlating ethyl**

334　**carbamate dose and Pig-a mutant frequency response in females.**

Diet	Equation	R^2	Deviation from zero p value	Slope comparison p value
LFD	y=0.0006029x + 0.4032	0.0376	0.3872	**0.0003**
HFD	y=0.005166x - 0.02898	0.5924	**<0.0001**	

Bold type indicates the slope or slope difference deviates significantly from zero

(α=0.05).

Among males there was no significant difference between the response of the HFD and

LFD groups to any level of ethyl carbamate treatment (Fig 7A). Though there was a

trend suggesting HFD males exhibit an approximately 2-fold increase in Pig-a mutant

frequency, regardless of ethyl carbamate treatment, these differences were not

statistically significant (Fig 7B). Linear regression analysis of the correlation between

ethyl carbamate dose and mutant frequency with slope comparison between the HFD

and LFD groups suggest a positive trend, although the slopes were not significant and

the R^2 values were low (LFD: p=0.1032, R^2=0.1486; HFD: p=0.0951, R^2=0.1471; Fig

7C). There was ultimately no difference between the slopes for the LFD and HFD

curves, suggesting both dietary groups are equally sensitive to ethyl carbamate-induced

mutagenesis (p=0.5654, Table 4).

Fig 7. Effects of diet and body mass on mutant frequency in erythrocytes of male mice exposed to 0, 100, 200, and 400 mg/kg/day ethyl carbamate. (A) Mutant frequency per million erythrocytes in each ethyl carbamate dose group. (B) Mutant frequency fold-difference between LFD and HFD groups in each ethyl carbamate dose group. (C) Dose-response curves for ethyl carbamate and erythrocyte mutant frequency with simple linear regression analysis, points represent mean ±standard error (n=4-5).

Table 4: Simple linear regression dose response analysis between ethyl

Diet	Equation	R^2	Deviation from zero p value	Slope comparison p value
LFD	y=0.002403x+0.1553	0.1486	0.1032	
HFD	y=0.003947x+0.5569	0.1471	0.0951	0.5654

carbamate dose and Pig-a mutant frequency response in males.

4. Discussion

The purpose of this set of experiments was to evaluate the mutagenicity of HFD both alone and with a chemical mutagen challenge using the Pig-a gene mutation assay. The findings presented here describe the first time that the effect of HFD on spontaneous somatic gene mutation frequency was tested in both males and females with repeated sampling over the lifespan. Interestingly, females in this long-term experiment were more responsive to HFD-induced weight gain as they aged than males, as indicated by the fact that HFD females were twice as heavy as LFD females, while HFD males were less than 50% heavier than LFD males after they reached 1 year of age (Fig 2). This was likely due to the fact that even males on the LFD become overweight over time while females do not, a phenomenon well-documented in the C57BL/6J strain[25]. Because of this tendency of males to gain weight over time, regardless of diet, we postulate that both LFD and HFD males represent some degree of biological overweight/obesity.

In females, linear regression analysis demonstrates that there is a significant dose-responsive relationship between body mass and Pig-a mutant frequency at 33 and 52 weeks old. This association between body mass and Pig-a mutant frequency

strengthens with age and continued HFD exposure (Fig 3C-E, Table 1), suggesting that

the length of the duration of HFD exposure over the lifespan is linked to degree of

spontaneous gene mutation risk. These findings suggest that HFD-induced weight gain

in females is sufficient to increase the risk of the initiation phase of carcinogenesis.

HFD females developed, on average, 3.8-times more somatic mutations in

response to high-dose ethyl carbamate than LFD females (Fig 6B). In fact, obesity

appears to potentiate the mutagenicity of ethyl carbamate in females, as indicated by

the significance of the ethyl carbamate-mutant frequency dose-response slope among

HFD, but not LFD, females (Fig 6C, Table 3). Among HFD males, mutagenicity of ethyl

carbamate appears to saturate between 100 and 400 mg/kg/day, with a peak at 200

mg/kg/day (Fig 7A-C). Our previous findings suggested that HFD males exhibited a 2-

fold elevation in mutation frequency, and that trend was recapitulated in this experiment,

though the difference did not reach the threshold of statistical significance[12]. HFD

males tended to exhibit non-significantly higher mutant frequencies than LFD males in

each ethyl carbamate dose group, however the dose-response curves for LFD and HFD

males shared a common slope (Fig 7C, Table 4). These results suggests that males on

both diets were similarly sensitive to ethyl carbamate. We attribute the overall lack of

association between diet and mutant frequency with the underlying weight gain

observed in LFD males, and this supports our assertion that studies that only include

male C57Bl6/J mice are not sufficient to explore the interactive effects of high-fat diet-

induced obesity and environmental toxicants.

Further investigation is needed to elucidate the mechanism behind the increased

ethyl carbamate sensitivity observed in HFD-exposed female mice. Several human

studies have indicated that individuals with a higher body mass index metabolize
xenobiotics differently. For example, the cytochrome P450 3A4 isoform tends to be less
active in the context of obesity, while the 2e1 isoform tends to be more active. This
group of xenobiotic metabolizing enzymes, part of the phase I class of chemical
metabolism, can be responsible for both detoxifying and increasing the toxicity of drugs
and other chemicals, depending on the substrate and efficiency of downstream phase II
metabolizers. Our findings emphasize the importance of investigating the mechanisms
underlying the HFD-induced mutagen sensitivity phenotype.

Interpretation of sex differences observed in this set of experiments is
complicated by the different trends in weight gain between male and female mice. That
is, males on the LFD gained weight consistently throughout the duration of the study,
while females on the LFD maintain proportionally lower body mass. Increased weight
gain by LFD males maintained for longer periods of time may mask differences that
might be evident a study that included in truly lean (non-overweight, non-obese) males
as a control group. Additional calorie restriction or reduced dietary fat may be required
to generate a male equivalent of the female non-obese phenotype within this model
system.

5. Conclusions

The findings presented here emphasize the importance of studying obesity as a
mutagen and carcinogen and the necessity of including both sexes in diet-induced
obesity research. For chemical risk assessment to properly protect the most vulnerable
groups, obesity must be factored in. In our female model, obesity appears to act to both

directly induce *de novo* mutations and to increase sensitivity to xenobiotic-induced

mutagenesis. We demonstrated that obesity is likely involved in the initiation

(mutagenesis) stage of cancer progression, however, further studies are needed to

investigate the differences in male and female response to HFD exposure. The

underlying the mechanism behind the association between HFD and ethyl carbamate

exposure and somatic mutagenesis is also important to investigate further.

Acknowledgements

We thank Stephen Dertinger, Dorothea Torous, and Svetlana Avlasevich of Litron

Laboratories for conducting the Pig-a mutation assay for the long term HFD experiment.

All funding for these projects was from intramural sources at Tulane University.

References

1. Steele CB, Thomas CC, Henley SJ, Massetti GM, Galuska DA, Agurs-Collins T, et
al. *Vital Signs*☐: Trends in Incidence of Cancers Associated with Overweight and
Obesity — United States, 2005–2014. MMWR Morb Mortal Wkly Rep. 2017;66:
1052–1058. doi:10.15585/mmwr.mm6639e1
2. Howlader N, Noone AM, Krapcho M, Miller D, Brest A, Yu M, et al. SEER Cancer
Statistics Review (CSR) 1975-2018. Bethesda, MD; 2021 Apr.
3. Cotangco KR, Liao C-I, Eakin CM, Chan A, Cohen J, Kapp DS, et al. Trends in
Incidence of Cancers Associated With Obesity and Other Modifiable Risk Factors
Among Women, 2001–2018. Prev Chronic Dis. 2023;20: 220211.
doi:10.5888/pcd20.220211
4. Cha B, Yu JH, Jin Y-J, Suh YJ, Lee J-W. Survival Outcomes According to Body
Mass Index in Hepatocellular Carcinoma Patient: Analysis of Nationwide Cancer
Registry Database. Sci Rep. 2020;10: 8347. doi:10.1038/s41598-020-65460-9
5. Mozafarinia M, Sasanfar B, Toorang F, Salehi-Abargouei A, Zendehdel K.
Association between dietary fat and fat subtypes with the risk of breast cancer in an
Iranian population: a case-control study. Lipids Health Dis. 2021;20: 138.
doi:10.1186/s12944-021-01557-y
6. Seyyedsalehi MS, Collatuzzo G, Huybrechts I, Hadji M, Rashidian H, Safari-
Faramani R, et al. Association between dietary fat intake and colorectal cancer: A

multicenter case-control study in Iran. Front Nutr. 2022;9.
doi:10.3389/fnut.2022.1017720

7. 458 Yang JJ, Yu D, Takata Y, Smith-Warner SA, Blot W, White E, et al. Dietary Fat
459 Intake and Lung Cancer Risk: A Pooled Analysis. Journal of Clinical Oncology.
460 2017;35: 3055–3064. doi:10.1200/JCO.2017.73.3329

8. 461 Zhao L, Deng C, Lin Z, Giovannucci E, Zhang X. Dietary Fats, Serum Cholesterol
462 and Liver Cancer Risk: A Systematic Review and Meta-Analysis of Prospective
463 Studies. Cancers (Basel). 2021;13: 1580. doi:10.3390/cancers13071580

9. 464 Pasupuleti SK, Ramdas B, Burns SS, Palam LR, Kanumuri R, Kumar R, et al.
465 Obesity-induced inflammation exacerbates clonal hematopoiesis. Journal of
466 Clinical Investigation. 2023;133. doi:10.1172/JCI163968

10. 467 Siddiqui IA, Sanna V, Ahmad N, Sechi M, Mukhtar H. Resveratrol
468 nanoformulation for cancer prevention and therapy. Ann N Y Acad Sci. 2015;1348:
469 20–31. doi:10.1111/nyas.12811

11. 470 Fu H, Tang B, Lang J, Du Y, Cao B, Jin L, et al. High-Fat Diet Promotes
471 Macrophage-Mediated Hepatic Inflammation and Aggravates Diethylnitrosamine-
472 Induced Hepatocarcinogenesis in Mice. Front Nutr. 2020;7.
473 doi:10.3389/fnut.2020.585306

12. 474 Wickliffe JK, Dertinger SD, Torous DK, Avlasevich SL, Simon-Friedt BR, Wilson
475 MJ. Diet-induced obesity increases the frequency of *Pig-a* mutant erythrocytes in
476 male C57BL/6J mice. Environ Mol Mutagen. 2016;57: 668–677.
477 doi:10.1002/em.22058

13. 478 Kimoto T, Suzuki K, Kobayashi X mei, Dobrovolsky VN, Heflich RH, Miura D, et
479 al. Manifestation of Pig-a mutant bone marrow erythroids and peripheral blood
480 erythrocytes in mice treated with N-ethyl-N-nitrosourea: Direct sequencing of Pig-a
481 cDNA from bone marrow cells negative for GPI-anchored protein expression.
482 Mutation Research/Genetic Toxicology and Environmental Mutagenesis. 2011;723:
483 36–42. doi:10.1016/j.mrgentox.2011.03.016

14. 484 Bemis JC, Hall NE, Dertinger SD. Erythrocyte-Based Pig-a Gene Mutation Assay.
485 2019. pp. 29–57. doi:10.1007/978-1-4939-9646-9_2

15. 486 Olsen A-K, Dertinger SD, Krüger CT, Eide DM, Instanes C, Brunborg G, et al. The
487 *Pig-a* Gene Mutation Assay in Mice and Human Cells: A Review. Basic Clin
488 Pharmacol Toxicol. 2017;121: 78–92. doi:10.1111/bcpt.12806

16. 489 Labash C, Avlasevich SL, Carlson K, Berg A, Torous DK, Bryce SM, et al. Mouse
490 *Pig-a* and micronucleus assays respond to *N* -ethyl- *N* -nitrosourea, benzo[*a*
491]pyrene, and ethyl carbamate, but not pyrene or methyl carbamate. Environ Mol
492 Mutagen. 2016;57: 28–40. doi:10.1002/em.21965

17. 493 Miura D, Dobrovolsky VN, Kasahara Y, Katsuura Y, Heflich RH. Development of
494 an in vivo gene mutation assay using the endogenous *Pig-A* gene: I. Flow
495 cytometric detection of CD59-negative peripheral red blood cells and CD48-
496 negative spleen T-cells from the rat. Environ Mol Mutagen. 2008;49: 614–621.
497 doi:10.1002/em.20414

18. 498 IARC. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to
499 Man: Some Anti-thyroid and Related Substances, Nitrofurans and Industrial
500 Chemicals. Lyon; 1974.

501   19.   Zhou K, Siroli L, Patrignani F, Sun Y, Lanciotti R, Xu Z. Formation of Ethyl
502   Carbamate during the Production Process of Cantonese Soy Sauce. Molecules.
503   2019;24: 1474. doi:10.3390/molecules24081474
504   20.   Hasnip S, Crews C, Potter N, Christy J, Chan D, Bondu T, et al. Survey of Ethyl
505   Carbamate in Fermented Foods Sold in the United Kingdom in 2004. J Agric Food
506   Chem. 2007;55: 2755–2759. doi:10.1021/jf063121c
507   21.   Schlatter J, Lutz WK. The carcinogenic potential of ethyl carbamate (urethane):
508   Risk assessment at human dietary exposure levels. Food and Chemical Toxicology.
509   1990;28: 205–211. doi:10.1016/0278-6915(90)90008-B
510   22.   Brill MJE, Diepstraten J, van Rongen A, van Kralingen S, van den Anker JN,
511   Knibbe CAJ. Impact of Obesity on Drug Metabolism and Elimination in Adults and
512   Children. Clin Pharmacokinet. 2012;51: 277–304. doi:10.2165/11599410-
513   000000000-00000
514   23.   Golde WT, Gollobin P, Rodriguez LL. A rapid, simple, and humane method for
515   submandibular bleeding of mice using a lancet. Lab Anim (NY). 2005;34: 39–43.
516   doi:10.1038/laban1005-39
517   24.   Dertinger SD, Bryce SM, Phonethepswath S, Avlasevich SL. When pigs fly:
518   Immunomagnetic separation facilitates rapid determination of Pig-a mutant
519   frequency by flow cytometric analysis. Mutation Research/Genetic Toxicology and
520   Environmental Mutagenesis. 2011;721: 163–170.
521   doi:10.1016/j.mrgentox.2011.01.009
522   25.   Reynolds TH, Dalton A, Calzini L, Tuluca A, Hoyte D, Ives SJ. The impact of age
523   and sex on body composition and glucose sensitivity in C57BL/6J mice. Physiol
524   Rep. 2019;7: e13995. doi:https://doi.org/10.14814/phy2.13995
525