Decoding Phase Separation of Prion-Like Domains through Data-Driven Scaling Laws

M. Julia Maristany[1]**, Anne Aguirre Gonzalez**[2]**, Jorge R. Espinosa**[3]**, Jan Huertas**[2]**,**
Rosana Collepardo-Guevara[2,4]**, Jerelle A. Joseph**[5,6]*****

*For correspondence:
jerellejoseph@princeton.edu (JAJ)

[1]Department of Physics, University of Cambridge, Cambridge, UK; [2]Yusuf Hamied
Department of Chemistry, University of Cambridge, Cambridge, UK; [3]Universidad
Complutense de Madrid, Madrid, Spain; [4]Department of Genetics, University of
Cambridge, Cambridge, UK; [5]Department of Chemical and Biological Engineering,
Princeton University, USA; [6]Omenn–Darling Bioengineering Institute, Princeton
University, USA

Abstract Proteins containing prion-like low complexity domains (PLDs) are common drivers of
the formation of biomolecular condensates and are prone to misregulation due to amino acid
mutations. Here, we exploit the accuracy of our residue-resolution coarse-grained model, Mpipi,
to quantify the impact of amino acid mutations on the stability of 140 PLD mutants from six
proteins (hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1). Our simulations reveal the existence
of scaling laws that quantify the range of change in the critical solution temperature of PLDs as a
function of the number and type of amino acid sequence mutations. These rules are consistent
with the physicochemical properties of the mutations and extend across the entire family tested,
suggesting that scaling laws can be used as tools to predict changes in the stability of PLD
condensates. Our work offers a quantitative lens into how the emergent behavior of PLD
solutions varies in response to physicochemical changes of single PLD molecules.

Introduction

Biomolecular condensates are highly multicomponent systems with remarkable specificity: they
exhibit non-random molecular compositions and, concomitantly, contain distinct physicochemi-
cal environments defined by the collective behavior of their components [1, 2]. Phase separation
has emerged as a leading mechanism to account for the formation of biomolecular condensates.
In this thermodynamic process, segregative (density-driven) and associative (connectivity-driven)
transitions are coupled to produce protein- and nucleic acid-rich phases suspended in the cellu-
lar cytoplasm or nucleoplasm [3]. Within this framework, intrinsically disordered proteins/regions
(IDP/IDRs) have emerged as strong contributors to the multivalency and phase separation capac-
ity of many naturally occurring proteins [1, 4–12]. Moreover, IDRs characterized by amino acid se-
quences of low complexity and enriched in aromatic residues, such as the prion-like low-complexity
domains (PLDs) of RNA–binding proteins [8, 13–16], are prone to form condensates that behave
like viscous fluids. Furthermore, the PLDs of various naturally occurring proteins, such as Fused
in Sarcoma (FUS), Trans-activation response DNA-binding protein 43 (TDP-43), and the heteroge-
neous nuclear ribonucleoprotein A1 (hnRNPA1) have been reported to transition from liquid-like
condensates into dynamically arrested solids and glasses over time [17, 18]. Such arrested states

have been postulated as precursors of pathological protein aggregates and amyloids [19]. Understanding the molecular factors that regulate the ability of PLDs to phase separate is of crucial importance, as these are conjectured to also facilitate their aggregation [20].

Biomolecular condensates are networked fluids sustained by multivalent macromolecules that interconnect with more than two partners simultaneously [1, 2, 21–25]. Thus, the factors that increase the molecular valency, give rise to more densely connected liquid networks, and more stable condensates condensates [1, 23, 26–29]. For PLDs, multivalency has been shown to be strongly promoted by the presence of aromatic residues, which act as "stickers" that enable associative π–π interactions, and also cation–π contacts with arginine and lysine [8, 9, 30–36]. Indeed, the range of coexistence of PLD condensates has been shown to change most significantly in response to mutations involving aromatic residues [9, 37]. Quantitative experimental phase diagrams for a set of hnRNPA1-IDR variants in combination with mean-field simulations demonstrated that tyrosine drives stronger associative interactions than phenylalanine, arginine is very context-specific, and lysine destabilizes PLD–PLD interactions [33]. More importantly, such study demonstrates that, at least for hnRNPA1, there exist quantitative rules that explain the changes in thermodynamic parameters of condensates in response to precise amino-acid sequence mutations.

In this work, we investigate whether there are scaling laws that can predict how the critical parameters of a broader collection of PLDs change in response to amino acid sequence mutations. Scaling laws are mathematical relationships that describe how the properties of a system change for a given variable, such as its size or composition [38]. In the context of polymer-like systems such as IDPs, scaling laws have offered invaluable insight into vastly different aspects of their physical behavior: for example, they were instrumental in determining that the relationship between viscosity and radius of gyration in polymer solutions is described by a power law between the two quantities [39]; demonstrating that aging scaling during polymer collapse is a universal property in homopolymers [40]; determining that in single polymer systems, their end-to-end distance is directly proportional to their length [41]; showing that foldable sequences with more compact unfolded states are recent products of protein evolution [42]. In the context of protein condensates, we hypothesized that scaling laws, if found, may allow us to quantify how condensates respond to different intrinsic perturbations—such as amino acid mutations or presence/absence of different post-translational modifications, changes in protein length, increased/decreased flexibility—or external stimuli—such as changes to the pH of the medium or the ionic environment. Thus, we focused on on determining if such laws exist for PLDs by exploring six different RNA-binding proteins (i.e., hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1; Fig. 1 A,B), linked to the etiology of neurodegenerative disorders through pathology and genetics [43–53].

Leveraging the accuracy and efficiency of a modern sequence-dependent coarse-grained model [32], we quantify the phase diagrams (temperature-vs-density plane) of large set of mutants spanning all six PLDs mentioned above. We performed Direct Coexistence molecular dynamics (MD) simulations (Fig. 1 C), exploring simultaneously the protein-rich and protein-depleted phases of the condensate for each mutant, in the temperature–concentration phase space. Our massive set of simulations encompasses 140 PLD mutants, each sampled at 8 to 12 different temperatures. Accounting for independent replicas to ensure proper sampling and convergence, the dataset presented here comprises the results of 2000 independent Direct Coexistence simulations (Fig. 1 D), totaling 1.2 milliseconds of accumulated sampling.

Our work puts forward a data-driven scaling-law framework for the study of PLD phase behavior. We reveal that the changes in the critical solution temperatures of PLDs, as a function of the number and type of amino acid sequence mutations (e.g., aromatic, positively charged, etc), can indeed be quantified by scaling laws, which are highly conserved across the family of PLDs we investigate. The addition of aromatic residues yields a positive linear correlation between mutation fraction and critical temperature changes, consistent with the dominant role of aromatic residues in PLD phase separation. We find that protein length dictates scaling effects when aromatic residues are mutated into uncharged, non-aromatic amino acids. We also quantify the effects of arginine

mutations on condensate stability, where removing arginines decreases the critical temperature of the condensate—revealing an inverse relationship between the fraction of arginine deletions and critical temperatures. Finally, polar amino acids emerge as subtle modulators of the phase behavior of PLDs. By examining the impact of different amino acid substitutions on the critical solution temperature, our results provide insights into the underlying molecular mechanisms of protein condensation. The scaling laws presented in this study serve two purposes: firstly, to verify the generality of the molecular grammar that governs the phase behavior across various PLD proteins, and secondly, to quantify the extent to which the critical solution temperature of a PLD protein solution will change in response to specific amino acid sequence changes. We anticipate that scaling laws for other families of proteins with very different compositional biases (e.g., transcription factors, nucleoporins, or molecular chaperones) may exist, but could be distinct from those we report here for PLDs. Overall, our work suggests that scaling laws represent useful predictors of the phase behavior of protein families.

Results

Generating and Analyzing an Extensive Dataset of Critical Solution Temperatures for Prion-like Domain Variants

The phase behavior of PLDs and their modulation with amino acid sequence mutations has been extensively investigated with a variety of experimental techniques and with theory and simulation methods for systems including the hnNRPA1 PLD [33, 54], the FUS PLD [8, 49, 51, 55–63], and the TDP-43 PLD [45, 46, 50, 64, 65], among many others. Here, we perform residue-resolution coarse-grained Molecular Dynamics (MD) simulations (see Fig. 1 C, with further details in Methods) with our Mpipi model [32] to quantitatively compare the critical parameters of a total of 140 mutants of the PLDs of TDP-43, FUS, EWSR1, hnRNPA1, RBM14, and TIA1 (Fig. 1 A) under consistent solution conditions. From our simulations, we compute temperature-vs-density phase diagrams and estimate the critical solution temperature for each system. The PLDs of the proteins we investigate have similar composition biases—i.e., are highly rich in neutral, polar, and aromatic residues—and present less than 5% of total charged amino acids in their sequence (Fig. 1 B). Thus, our simulation results allow us to assess if there exists a code—i.e., a simple set of scaling laws—that can predict how the critical parameters of the family of PLDs will scale in response to amino acid sequence mutations.

Our residue-resolution model Mpipi (see Methods) is ideally suited to investigate the phase behavior of PLDs because it achieves the correct balance of the dominant π–π and cation–π interactions, which drive the phase behavior of PLDs. Indeed, the Mpipi model recapitulates the quantitative experimental phase diagrams of hnRNPA1 variants by Bremer and colleagues [33] with remarkable accuracy [32]. It is worth noting that the Mpipi model was not parameterized based on the experimental dataset of Bremer and colleagues, but rather the model parameters were developed independently by combining atomistic umbrella sampling MD simulations of amino acid pairs with the bioinformatics data of Vernon and colleagues for π-driven contacts [35]. Using Flory–Huggins–Staverman theory, critical temperatures for phase separation were extracted from the experimental data of Bremer et al., which were then used to validate the Mpipi model [32].

The amino acid sequences of all the proteins we investigated (Supporting Information) were designed based on five main considerations. (1) The set of mutants should serve to probe the effects of both the strong modulators of PLD phase separation—namely, aromatic and arginine interactions—and the subtle modulators—that is, polar and neutral amino acids, previously identified experimentally [8, 9, 33]. (2) To ensure that the observed changes in the critical solution temperatures can be attributed unequivocally to specific amino-acid-type changes, we mutated only one amino acid type at a time. However, multiple mutations of the same amino acid type were allowed. (3) Mutations were distributed homogeneously across the entire protein. (4) Deleted amino acids (e.g. aromatic deletions) were replaced by weakly interacting amino acids (i.e. serines,

Figure 1. Data generation and representation of dataset for probing scaling laws of PLD phase behavior (A) Representation of the prion-like domains (grey-shaded regions) of each considered protein, alongside the associated number of variants simulated, totaling a set of 140 sequences. **(B)** Composition of the wild-type sequence of the PLDs simulated, in terms of the percentage of neutral amino acids (A, C, I, L, M, P, S, T, and V; no net charge at pH 7 and no π electrons in the side chain), glycine (G), negative (D, E), positive (K, H, R), neutral-π (N and Q; no net charge at pH 7 with π electrons in the side chain) and aromatic (F, W, Y) residues. **(C)** Top panel: Representation of the simulation model: each amino acid of the protein is represented by a single bead, bonded interactions are modeled with springs, and non-bounded interactions are modeled with a combination of the Wang–Frenkel potential and a Coulomb term with Debye-Hückel screening. Middle panel: The Direct Coexistence simulation method, in the slab geometry, is employed, where both the protein-rich and protein-depleted phases are simulated simultaneously. Bottom panel: Finally, to obtain the data point in the concentration–temperature phase space, at a given temperature the density profile is computed and the concentrations of the rich and depleted phases are obtained via a suitable fitting (Methods). **(D)** Representation of the entire computational data set. Orange data points represent variants where charged or aromatic residues were mutated (i.e., stickers), while cyan data points represent all other types of mutations studied (i.e., spacers).

Figure 2. Mpipi model quantitatively captures phase behavior for a large set of hnRNPA1 PLD mutants. Mutations are divided into **A** aromatic, **B** Arg/Lys, **C** charged, and **D** Gly/Ser sequence variants of the PLD of the protein hnRNPA1, and comparison with experimental results. The hnRNPA1 PLD was used to validate the accuracy of the model, by extracting critical temperatures from experimental phase diagrams of a large set of these variants [33]. To assess how well the Mpipi model performed for PLDs, we compared critical temperatures obtained from simulations with the corresponding ones extracted from the experimental data for those variants that were available (depicted in the top right of each binodal, where *r* is the Pearson coefficient and *D* is the root mean square deviation between simulated and experimental values). For the variants where the experimental critical temperature could not be determined, we make a comparison between the critical temperature with the saturation concentration, since for these systems we expect them to be inversely correlated. The naming system for the variants is discussed in detail in the text.

141 threonines, glycines, and alanines). For each PLD protein, the identity of the weakly interacting
142 residues used as replacements were chosen to preserve the percentages of the various types of
143 weakly interacting amino acids present originally in the unmodified PLD. (5) Achieve a data set that
144 spans a wide array of critical solution temperatures, as shown in Fig. 1 D.

Validation of the Mpipi Model for Phase behavior of Prion-like Domains

The availability of quantitative experimental binodals for the hnRNPA1-IDR and various amino acid
sequence variants [33] has allowed us to test the accuracy of the Mpipi model to describe the
phase diagrams of PLD solutions [32]. Here, we make such a test more stringent by extending
the hnRNPA1-IDR validation set from 9 PLD variants to 21, selecting mutations spanning changes
in aromatic, charged, polar, and neutral residues. The amino acid sequences of all the hnRNPA1-
IDR variants considered are described in the Supporting Information and are denoted using the
intuitive nomenclature established by Bremer and colleagues: variants that mutate n residues of
type X into m residues of type Z are labeled as $-nX + mZ$, and so forth. For more details on the
nomenclature, see Methods.

For each hnRNPA1-IDR variant, we perform Direct Coexistence MD simulations, compute criti-
cal temperatures from the simulations, and compare them with the experimental critical temper-
atures (Fig. 2 A–D). We use the fitting methodology outlined in Methods to estimate the critical
solution temperatures from the experimental data.

When comparing with experimental critical temperatures, our model achieves a Pearson correlation coefficient higher than 0.93 for all types of mutations analyzed. Additionally, for all cases, the root-mean-square deviation (D) between the experimental and simulated critical temperatures is ≤ 14 Kelvin. Our validation set, which expands the range of proteins variants originally tested [32], highlights that the Mpipi potential can effectively capture the thermodynamic behavior of a wide range of hnRNPA1-PLD variants, and suggests that Mpipi is adequate for proteins with similar sequence compositions, as in the set of proteins analyzed in this study. In recent work by others [66], Mpipi was tested against experimental radius of gyration data for 137 disordered proteins and the model produced highly accurate results, which further suggests the applicability of the approach to a broad range of sequences.

Large-Scale Data Generation on Critical Parameters of PLD Phase Separation

The next step of the study consists of generating a large amount of data, needed to report the critical temperatures of the PLD condensates. In the Direct Coexistence method, the diluted phase and the condensate are simulated in the same rectangular box separated by an interface. The phase diagrams we compute explore the plane of temperature (vertical axis) versus density (horizontal axis). For each phase diagram, we run a set of about 20 Direct Coexistence simulations–each simulation explores the phase behavior of the system at different constant temperatures. For a given value of the temperature, if two phases are detected in our simulations, we measure a "density profile": the density of the proteins as a function of the long side of the simulation box (Bottom panel in Fig. 1 C). We then plot the two coexistence densities for each temperature we simulate and build the coexistence curve (also known as binodal), Fig. 3. The coexistence curve shows the values of temperature and protein density for which phase separation will occur. The region above the coexistence curve is the "one-phase region" where the temperature is too high for the enthalpic gain coming from the associative protein–protein interactions to overcome the entropic loss upon phase separation. The region below the coexistence curve, the "two-phase coexistence region" represents lower temperatures for which the enthalpic gain due to protein–protein interactions is now sufficiently high to favor phase separation into a condensed (protein-enriched) and a diluted (protein-depleted) liquid phase. The maximum in the coexistence curve is known as the critical point: the upper critical solution temperature (herein, critical solution temperature) and corresponding critical density. For temperatures higher than the critical solution temperature, phase separation is no longer observed.

The protein solutions that we investigate with our coarse-grained simulations must be sufficiently large, in terms of the number of copies of the same protein, to minimize finite size effects—for instance, to adequately sample bulk phenomena without dominating interfacial effects. At the same time, our simulations must be small enough to ensure computational convergence in a feasible timescale. Following extensive testing of computational convergence for systems of different sizes and MD trajectories of different time lengths, we determined that a system composed of 64 protein replicas, with trajectories of 200 ns ensured rigorous convergence and reproducible results, while taking into careful consideration the trade-off between accuracy and computational demands. The size of the simulation boxes were also optimized, again to ensure finite size effects did not affect the predictions (See Methods and Materials, and Supporting Information).

As a result, we generated a dataset of critical parameters of over an order of magnitude higher than previously reported studies [9, 33, 67–70]. This extensive dataset (Fig. 3) not only offers new insights into phase separation phenomena but also sets a new benchmark for data-driven research in this field.

A Data-Driven Scaling Law Approach for Studying Phase Behavior of Prion-like Domains

Scaling laws can reveal key relationships between system variables, such as composition and critical parameters, providing a useful theory to predict and interpret the regulation of complex bio-

Figure 3. Complete set of 140 binodals, in the temperature versus density phase space, generated via largscale molecular dynamics simulations. Binodals are grouped according to the mutation type (Aromatic, Arg/Lys, Gln/Asn (Polar), Ala/Ser and Thr/Gly/Ser (Neutral)). Each set is further grouped based on the PLD in question.

logical phenomena such as phase separation. In the context of PLD phase behavior, scaling laws would provide a quantitative theory to analyze how alterations in amino acid composition affect the critical solution parameters—a crucial aspect for understanding protein behavior in biological systems. Here, our objective is to discern which scaling laws can effectively capture the variations of PLD critical parameters as a function of sequence mutations. To accomplish this, we employ a systematic approach of varying the amino acid sequence of the proteins considered, to determine how the phase behavior of PLDs changes with amino acid compositional perturbations.

For each type of mutation we analyze, we define and test three types of scaling laws. These scaling laws illustrate a different relationship between critical solution temperature changes and the number of amino acid mutations. The first and simplest scaling law is a linear relationship between these two quantities,

$$\Delta T_C = S_{X_1 \to X_2} N_{X_1 \to X_2}, \tag{1}$$

where ΔT_c is the variation on the critical temperature between the mutated protein and the corresponding unmodified reference sequence, X_i is an amino acid (e.g., Y, F, R, and so forth) and $N_{X_1 \to X_2}$ is the number of X_1 residues that have been mutated to X_2. Thus, $S_{X_1 \to X_2}$ in this case quantifies the variation in the critical solution temperature of the protein solution for a mutation of one X_1 residue to X_2. If the data is well represented by this first scaling law, it would imply that the effects of amino acid mutations on PLD phase behavior depend exclusively on the total number of mutations, but are not affected by other PLD parameters, such as the protein length. Thus, the first scaling law (Eq. 1) is the simplest possible model we test here for how incremental mutations

may collectively influence the phase behavior of PLDs.

Next, we define a second type of scaling law, which hypothesizes that the changes to the critical solution temperature upon mutation is proportional to the fraction of residues that are mutated, rather than their raw number. That is,

$$\Delta T_c = S_{X_1 \to X_2} \frac{N_{X_1 \to X_2}}{L} = S_{X_1 \to X_2} f_{X_1 \to X_2}, \tag{2}$$

where, L represents the number of total amino acids in the PLD, or protein length, and $f_{X_1 \to X_2} = \frac{N_{X_1 \to X_2}}{L}$ is the fraction of X_1 residues that have been mutated to X_2. Thus, this second scaling law hypothesizes that the effect of mutations on the critical solution temperature is more pronounced in shorter proteins (where $N_{X_1 \to X_2}$ mutations constitute a larger fraction of the protein) and less so in longer proteins (where $N_{X_1 \to X_2}$ mutations represent a smaller fraction of the total number of amino acids). The expectation is that because longer PLDs have higher valencies than shorter ones, the density of inter-molecular connections they form inside condensates is more robust towards perturbations than that of shorter PLDs [71].

Finally, we define a third scaling law,

$$\Delta T_c = S_{X_1 \to X_2} \frac{N_{X_1 \to X_2}}{\sqrt{L}} \tag{3}$$

where the change in critical solution temperature is hypothesized to be directly proportional to the number of mutations ($N_{X_1 \to X_2}$) but inversely linear to $\sqrt{L}$. Like our second scaling law, this third relationship also presumes that longer PLDs form condensates with critical parameters that are less affected by mutations than shorter PLDs. However, by considering $\sqrt{L}$, instead of L as in Eq. 2, the expectation is that the buffering effects of protein length are more modest than those stated by Eq. 2. Notably, the amino acid composition of PLDs can be understood by the stickers-and-spacers framework [72, 73]. Aromatic and arginine residues have been proposed as stickers, contributing to the stability of the liquid network via associative interactions, yet some groups argue that all residues matter equally. Our data suggests that while a strict black-and-white delineation might not be appropriate, certain residues such as Tyr or Arg demonstrably have a larger impact on T_c when mutated. Conversely, other residues exhibit smaller effects, acting more like spacers interspersed among the stickers. Eq. 3 suggests that as a protein lengthens, the number of residues that can establish sufficiently strong inter-protein interactions grows too; yet, it grows much more conservatively than L because a fraction of L is made of spacers.

With these scaling laws as ansatz, we set to uncover trends in the dataset we built from our MD simulations of 140 variants of different PLDs. Adopting these laws as foundational hypotheses, we explore the relationship between alterations in composition and changes in critical solution temperatures, offering insights into the molecular drivers of PLD phase separation.

Aromatic Mutations Exhibit a Linear Correlation Between Mutation Fraction and Critical Solution Temperatures

Aromatic amino acids, namely tyrosine (Tyr, Y), phenylalanine (Phe, F), and tryptophan (Trp, W), are widely regarded as essential contributors to phase separation of PLDs, due to their ability to form π–π and cation–π contacts [9, 30, 31, 33, 35, 36, 74–76]. Among aromatics, experiments and simulations agree that Tyr is a stronger contributor to the stability of condensates than Phe [8, 31, 33, 60], and that both the composition and the patterning of aromatic residues in PLDs have an impact on the condensate critical parameters and its material state [18, 68, 77, 78].

Aromatic residues are highly abundant in PLDs, with the fraction of aromatic residues in the PLD variants analyzed in this study ranging from 0.19 to 0.43 (Fig. 1B). Furthermore, because different aromatics are unequal contributors to PLD phase behavior, we analyzed the impact of mutations for Tyr and Phe and Trp separately.

Figure 4. Mutations in aromatic amino acids have strong effects on the critical solution temperature of PLDs. A. Tyr to Phe mutations. The number of Tyr mutated to Phe (x-axis) versus the change in critical temperature, (y-axis), computed as the critical temperature of the variant minus that of the corresponding PLD wild-type sequence. The trendline defines the simplest, dominating scaling law for this mutation. The R^2 value of 0.88 shows the agreement of our data to the scaling law defined by Equation 4. **B.** Tyr or Phe to Trp mutations. In this case, the overall trendline, defined by equation 5 has an R^2 of 0.99. **C.** Analysis of the variants involving mutations of aromatic residues to uncharged, non-aromatic amino acids, a.k.a. aromatic deletions. In the y-axis, the change in the critical temperature of the variant to that of the wild type sequence, and in the x-axis, a renormalized measure of mutations: the fraction of aromatic residues mutated times the $\sqrt{L}$, to account for the competing physical interactions between PLDs. Error bars indicate the standard error associated with replicas of the simulation.

To quantify the impact of mutating Tyr to Phe in shifting the critical solution temperatures of our set of PLDs—i.e., determining which of our proposed scaling laws best describes the data—we designed 17 protein mutants that varied the Tyr to Phe ratio in the family of PLDs and computed their phase diagrams (Fig. 4A). Our simulations reveal that Tyr is stronger than Phe as a driving force for phase separation for the entire PLD family considered, a finding that replicates the results of Bremer *et al.* [33] for hnRNPA1 and confirms that it extends to other PLD proteins. By analyzing the change of critical solution temperature with respect to the unmodified PLD (or wild type, WT) against the number of mutations, our results reveal a linear relationship between the number of Tyr to Phe mutations and the change in the critical temperature of the system; our first scaling law, $S_{Y \to F}$, Eq. 4. Here, ΔT_c represents the temperature difference between the mutated protein and its wild-type sequence, and $N_{Y \to F}$ represents the number of Tyr residues that were mutated to Phe in the sequence.

$$S_{Y \to F} = \frac{\Delta T_c}{N_{Y \to F}} = -0.40 \pm 0.04 \tag{4}$$

Note that, by definition $S_{Y \to F} = -S_{F \to Y}$, since $N_{Y \to F}$ is defined as positive for Tyr to Phe mutations and negative for Phe to Tyr mutations, or, in other words, $N_{Y \to F} = -N_{F \to Y}$. For our data set, we find $S_{Y \to F}$ to be -0.40 ± 0.04, where the error given is the standard error. The way to interpret this result is that a single mutation of a Tyr residue to a Phe residue in a given PLD protein would result in a -0.4 K variation in the critical solution temperature for phase separation. Such a modest variation due to a single-point mutation is smaller than the typical error associated with estimating the absolute value of the critical solution temperature itself using residue-resolution coarse-grained Direct Coexistence simulations (approximately 3 K). Unambiguously measuring the value of the critical solution temperature with sub-Kelvin accuracy poses a considerable challenge also in experiments [79, 80]. However, we can uncover statistically significant trends by analyzing the behavior of the critical solution temperature in a large set of PLDs with diverse numbers of mutations rather than a single case. In particular, our analysis reveals that the impact of Tyr to Phe mutations is modest and the effect of multiple mutations can be described adequately as cumulative.

In addition to these analyses, the role of Tyr/Phe to Trp mutations, albeit Trp being a rare residue in prion domains, was investigated in TIA1 due to the presence of Trp in the wild type of this PLD, Fig. 4B. The simplest scaling law that describes this mutation (Equation 5) is fundamentally equal to the law describing Tyr to Phe mutations. However, a notable distinction is that the $S_{F/Y \to W}$ scaling constant for our data is approximately 4.3 ± 0.1, which is one order of magnitude greater than $S_{Y \to F}$, consistent with the much stronger π–π contacts established by Trp. It is important to note that, in this case, similarly to Try to Phe mutations, multiple Ansatzs align closely with the observed data, which underscores the fact that, in the context of these mutations, the size of the side chains does not significantly alter phase behavior, as the predominance of aromatic perturbations exerts a dominating effect.

$$S_{F/Y \to W} = \frac{\Delta T_c}{N_{F/Y \to W}} = 4.3 \pm 0.1 \tag{5}$$

Across the whole family of PLDs analyzed, our data reveals that changes in aromatic amino acid identity impact the critical solution temperatures modestly (< 1 K per mutations involving F and Y) to strongly (4 K for mutations involving Trp) and independently of the protein length. Such a result is consistent with the mutations in aromatic identity preserving the total number of associative interactions that the PLDs can establish, and hence, the density of molecular connections in the condensed liquid network. That is, aromatic mutations are expected to influence the energetics and lifetimes of the multivalent interactions among PLDs, but not the number of intermolecular contacts they form.

Strong Critical Temperature Reduction Due To Aromatic Deletions is Buffered By Protein Length

We next examine the role of replacing aromatic residues (Tyr and Phe) with uncharged, non-aromatic residues—such as Gly, Ser, Gln, Ala, and Asn—which are prevalent in PLDs. The mutations were designed to preserve amino acid sequence compositional biases by replacing Tyr and Phe with uncharged, non-aromatic residues in the same proportions as they appear in the wild type of each sequence, i.e., maintaining the relative amounts of each type of residue found in the sequence. All amino acid sequences are available in the Supporting Information.

Our simulations show that aromatic deletions strongly reduce the critical solution temperature of the PLD solutions, Fig. 4C. The best scaling law from our ansätz to describe the data is Eq. 3, which involves the square root of the PLD length. That is, $S_{Y/F \to X}$, where X represents any uncharged, non-aromatic residue.

$$S_{Y/F \to X} = \frac{\Delta T_c \sqrt{L}}{N_{Y/F \to X}} = -56 \pm 5 \qquad (6)$$

Here, once again, $N_{Y/F \to X}$ is the number of mutations, and L is the length of the PLD. For our data set, the value of the scaling constant $S_{Y/F \to X}$ is -56 ± 5. To interpret this result, let us consider a model PLD of two hundred amino acids ($L = 200$). If we mutate one Tyr to an uncharged, non-aromatic amino acid, the change in the critical temperature—computed via the formula $S_{Y/F \to X} \times \frac{N_{Y \to X}}{\sqrt{L}}$—is approximately -4 K. Such a value is consistent with the dominance of π–π and cation–π contacts in PLD phase separation reported experimentally [9, 30, 31, 33, 35, 36, 74–76]. As stated above, aromatic residues behave as associate hubs within PLDs condensates. When these aromatic residues are replaced with uncharged non-aromatic amino acids the density of connections network of molecular connections in the liquid is strongly perturbed, as we have shown earlier [81]. To observe this in more detail, we computed select contact maps (see Fig. 2) – notably, mutations that removed aromatic amino acids strongly altered the primary sites of interactions.

Our results reveal that the length of the PLDs strongly influences how the critical parameters of PLD solutions change in response to aromatic deletions. The effect of mutations is stronger in shorter proteins because each aromatic deletion represents a higher percentage of the total number of associative residues in the PLD, than in longer proteins. Longer proteins have higher valencies and redundant "stickers" that facilitate a network rearrangement.

Our results highlight the delicate balance between amino acid composition and protein length in determining the phase behavior of PLDs, providing a deeper understanding of the molecular basis of protein phase transitions.

Arginine Mutations Drastically Alter Condensate Stability

The distribution and balance of charged amino acids–i.e., aspartic acid (Asp, D), glutamic acid (Glu, E), histidine (His, H), arginine (Arg, R), and lysine (Lys, K)—within an IDR affects its solubility, conformational ensemble, and the range of stability of the ensuing condensates it might form. PLDs are not particularly rich in charged residues: in the wild-type PLDs of our data set, the fraction of charged residue ranges from 0.02 to 0.1 (Fig. 1B), and the fraction of positively charged amino acids is narrowly distributed around 0.015, except for hnRNPA1, which has a value of 0.09 (Fig. 1B). However, despite their low prevalence in PLDs, charged residues, in particular Arg, have been shown to be important contributors to the stability of a wide-range of biomolecular condensates [30, 33, 82–84].

While both Arg and Lys are positively charged amino acids at physiological pH, it has been demonstrated that they have unequal roles in the stability of biomolecular condensates [30, 33, 60, 83, 85]. Both residues can form charge–charge and cation–π associative interactions, but the case of Arg is unique: the sp^2-hybridized planar guanidinium group of Arg confers it a special capacity to form strong hybrid π–π/cation–π contacts with the π-electrons of aromatic rings [31]. Thus, to

Figure 5. Critical effects of Arg mutations in the phase transition temperature of the condensates (A) Analysis of the critical solution temperature of PLDs with Arg to Lys mutations. The number of Arg to Lys mutations divided by PLD length (x-axis) versus the change in the critical solution temperature, that is, $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendline, representing Equation 7, fits the data with an $R^2 = 0.94$, and defines the stability measure of this perturbation. **(B)** Analysis of the critical solution temperature of PLDs with Arg mutations to uncharged, non-aromatic amino acids that maintain the WT compositional percentages (mainly Gly, Ser, and Ala. For more details see Supporting Information). Arg deletion fraction (x-axis) versus $T_c(\text{variant}) - T_c(\text{WT})$ (y-axis). The trendline, which fits the data with $R^2 = 0.96$, defines the stability measure, represented in Equation 8.

---

quantify the impact of Arg in the range of stability of PLD condensates, we designed mutants that replaced Arg with Lys and vice-versa (Fig. 5).

The best fitting scaling law associated with mutating Arg to Lys is inversely dependent on the fraction of mutations, rather than the number of residues mutated (Eq. 7). The resulting scaling constant for this particular mutation, i.e., $S_{R \to K}$, has a value of -1300 ± 100, indicating that for a typical PLD of $L = 200$, the mutation of a single Arg into Lys results in a reduction of the critical temperature of the system by $6.5\,\text{K}$. Hence, replacing Arg with Lys is a destabilizing mutation of PLD condensates in agreement with the wide-body of experimental evidence [30, 33, 60, 83, 85].

$$S_{R \to K} = \frac{\Delta T_c}{f_{R \to K}} = -1300 \pm 100 \tag{7}$$

The length dependency of this particular scaling law suggests that the impact of Arg to Lys mutations on the phase behavior of PLDs primarily depends on the relative concentration of these mutations (mutation fraction). Additionally, it proposes that longer sequences may exhibit increased resilience against Arg to Lys mutations. This resilience is likely attributed to the greater separation of charge within the longer sequences.

To further analyze the effect of the charge of the PLDs in this set of mutants, in Fig. 5 we include a scalebar showing the net charge per residue (NCPR). Mutants that present higher NCPR present the biggest decrease in their phase-separating propensities, which is consistent with the expectation that condensates with a neutral overall charge are more energetically stable [33, 86, 87].

When we replace Arg with uncharged, non-aromatic residues, the resulting scaling law also depends inversely on the fraction of mutations, with a value of -640 ± 50, amounting to a reduction of $3.2\,\text{K}$ in T_c per Arg deletion for a model PLD of length $L = 200$.

$$S_{R \to X} = \frac{\Delta T_c}{f_{R \to X}} = -640 \pm 50 \tag{8}$$

Therefore, our results reveal that it is almost twice more destabilizing to mutate Arg to Lys than to replace Arg with any uncharged, non-aromatic amino acid, namely, Gly, Ser or Ala. Mutating Arg

to an uncharged non-aromatic amino acid is less destabilizing than mutating it to Lys likely because the charge-reduced variants decrease the cation–cation repulsion of the PLDs, with respect to the Arg-rich or Lys-rich systems. Furthermore, mutating Arg to Lys decreases the enthalpic gain for condensate formation not only via the replacement of stronger Arg-based associative interactions for weaker Lys-based ones but also by increasing the overall solubility of the Lys-rich PLD [36, 88], given that the hydration free energy of Arg is less favorable than that of Lys [31, 36, 82].

Polar Amino Acids Subtly Modulate PLD Phase behavior

Uncharged, non-aromatic amino acids, including Gly, Ala, proline (P, Pro), Ser, and Thr, are commonly found in PLDs. Because they generally form weaker residue–residue interactions and present smaller excluded volumes than aromatics, they are classified as spacers that contribute to maintaining the solubility of proteins and material states of condensates [68, 72]. These amino acids can also have other functions in proteins, and the role they play as spacers depends on their chemical make-up, but also on the specific context in which they are found [33, 37, 89].

Here, we investigate the role of glutamine (Gln, Q) and asparagine (Asp, N). Both Gln and Asn are polar amino acids, hence their charge distribution is more asymmetric, presenting a significant dipole moment. Consequently, they can establish transient ion–dipole, dipole–dipole and π–dipole interactions with other residues. While the Mpipi model does not account explicitly for polarity, it is worth noting that it was parametrized based on bioinformatics data, which implicitly incorporates this aspect. Thus, our study allows for the observation of emergent behaviors that stem from this inherent characteristic.

We designed mutations that vary the relative amounts of Gln and Asn, and examined trends in their phase-separating propensity. Increasing the number of Gln per unit length by mutating Asn to Gln increases the ability of the system to phase separate, likely due to Gln having a higher dipole moment than Asn, which is approximated in our model via marginally stronger non-ionic interactions formed by Gln over Asn (Fig. 6 A).

$$S_{N \to Q} = \frac{\Delta T_c}{f_{N \to Q}} = 50 \pm 5 \tag{9}$$

The value of the scaling constant $S_{N \to Q}$ is 50±5, which is significantly lower than that of mutations of residues with a net charge or aromatic deletions (rounding 1300, for example, for arginine to lysine mutations). This result indicates that mutations of Asn to Gln or vice versa have a smaller impact in the critical parameters than aromatic deletions or mutations of charged residues.

Finally, we also explored the role of Ser, Gly, Thr, and Ala on the critical parameters of the condensate. While Ser and Thr are also polar amino acids, their dipole moment is significantly smaller than that of Gln and Asp. Consistently, the scaling constants involving mutations Ser and Thr mutations—i.e., $S_{S \to T}$— are considerably smaller (Fig. 6 B and C) overall. A comparable pattern is noted for $S_{G \to T}$, with G expected to exhibit significantly lower polarity compared to Gln and Asp.

$$S_{S \to T} = \frac{\Delta T_c}{N_{S \to T}} = -0.09 \pm 0.03 \tag{10}$$

$$S_{G \to T} = \frac{\Delta T_c}{N_{G \to T}} = -0.08 \pm 0.02 \tag{11}$$

Regarding Ala, we also tested its relative role with respect to Ser.

$$S_{A \to S} = \frac{\Delta T_c}{N_{A \to S}} = 0.16 \pm 0.05 \tag{12}$$

Amino acids with larger side chains, such as Thr, have a larger excluded volume than those with smaller side chains, such as Gly or Ser. The corresponding value of the scaling constants $S_{S \to T}$ and $S_{G \to T}$ suggests that in the case of Thr mutations, mutations of neutral, non-aromatic amino acids to those with larger excluded volumes inhibit phase separation of the PLD by lowering the corresponding critical temperature, while mutations to those amino acids with smaller

Figure 6. Polar and neutral amino acid mutations show subtle modulation of the critical solution temperatures of PLDsA) Analysis of the critical solution temperature of PLDs with Asn to Gln mutations. The number of Gln to Asn mutations renormalized by dividing by PLD length (x-axis) versus the temperature change T_c(variant) − T_c(WT) (y-axis). The trendline, which fits the data with an $R^2 = 0.84$, fits the scaling law Equation 9. **(B)** Analysis of the critical solution temperature of PLDs with Gly/Ser to Thr mutations (x-axis) versus the temperature change T_c(variant) − T_c(WT) (y-axis). The trendlines, representing Equations 10 and 11 fit the data with an $R^2 = 0.7$. **(C)** The number of Ala to Ser mutations (x-axis) versus the temperature change T_c(variant) − T_c(WT). The trendline, Equation 12, fits the data with an $R^2 = 0.77$

excluded volumes have an opposite effect. This is likely due to increased flexibility of the protein backbone, allowing the chain to adopt multiple conformations that promote multivalent contacts. Importantly, these trends cannot be anticipated by inspecting the parameters of the Mpipi model; instead, they emerge from examining the behavior of the collective system.

On the other hand, for mutations involving Ser, Gly, and Ala–all small amino acids with minor side chain size variations–the impact on phase separation seems to stem predominantly from changes in solvation. Solvation plays a key role in phase separation since it can help balance the driving forces of this process [90, 91]. While hydrophobic regions of the protein, with low solvation free energies, tend to form aggregates to reduce interactions with the solvent environment; hydrophilic regions promote interactions with water. Whether these regions inhibit or promote phase separation is likely due to a balance of opposing effects. On the one hand, hydrophilic regions can form energetically favorable contacts with solvent, promoting dispersion and suppressing phase separation. However, when these regions are recruited to condensates the release of water molecules can help to off-set the entropic cost of sequestering hydrophobic regions within a protein droplet, thereby facilitating phase separation.

Specifically, mutations from Gly to Ser (not shown) increase the size of the side chain while also enhancing solvation, while mutations from Ala to Ser do not significantly change the size of the side chain but significantly increase the protein solvation. In the first case, the critical parameters of phase separation remain virtually unchanged, only presenting a very minor decrease. In the case of Ala to Ser mutations, phase separation is slightly enhanced. Both scaling laws $S_{G \to S}$ and $S_{A \to S}$ reflect such an effect, and underlie the context-dependent nature of these mutations.

Overall, mutations of unchanged, non-aromatic amino acids seem to have an impact on phase separation due to two competing effects: a change in side chain size and a change in solvation. However, it is important to note that the magnitude of the scaling laws that govern these mutations is minimal, and thus, likely extremely sensitive to small fluctuations such as changes in the system and solution conditions. Notably, these findings are consistent with experiments that report small changes in phase separation propensity for mutations involving neutral, non-aromatic residues [8, 33].

Beyond composition: patterning considerations

Beyond composition, the relative positioning of amino acids in the sequence, i.e. sequence pattern, also plays a role in protein folding, function and, in this case, phase behavior [9, 92, 93]. Due to the sequence heterogeneity of PLDs, and, more generally, IDPs, it is particularly complex to define adequate patterning measures [87, 94–96].

In this work, we have taken advantage of our large data set to analyze the impact of amino acid sequence patterning by adopting a patterning order parameter [9, 86, 97], σ_{aro}, that has been designed to describe patterning of aromatic residues in such IDPs.

To compute σ_{aro}, we divide the amino acid sequence into peptide segments of equal length, compute a measure of sequence aromatic asymmetry for each segment, and finally sum the values of such measure over all the segments. Consistent with literature [9, 86, 97], we take l to be 5 and 6. σ_{aro} is then given by:

$$\sigma_{\text{aro}} = \left(\sum_{\text{segments}} \frac{\sigma_i}{\sigma_{\text{max},i}} \right) \frac{N_{\text{stickers}}}{L}, \tag{13}$$

where L is the total number of amino acids in the protein domain, N_{stickers} the number of sticker residues—i.e. charged or aromatic residues—and σ_i has the functional form

$$\sigma_i = \left(\left(2\frac{N_{\text{aro},i}}{l_i} - 1 \right)^2 - \sigma_{\text{T}} \right)^2 . \quad (14)$$

Here, $N_{\text{aro},i}$ is the number of aromatic residues in the i-th segment, l_i the segment length, and σ_{T} is a function of the total number of aromatic residues in the sequence N_{aro}:

$$\sigma_{\text{T}} = \left(2\frac{N_{\text{aro}}}{L} - 1 \right)^2 \quad (15)$$

In turn, $\sigma_{\text{max},i}$ is the measure of aromatic amino acid distribution for an artificial sequence with all the aromatic amino acids concentrated in part of the chain.

Notably, our results, presented in Fig. 7, indicate that the sequences with the most dispersed aromatic residues (i.e., low values of σ_{aro}) phase separate more readily. Dispersed residues result in greater effective valency of a PLD, leading to a higher probability of forming a percolated network structure to sustain the condensate. Additionally, this result demonstrates that the patterning order parameter, σ_{aro}, can be used to predict the phase-separating propensity of a PLDs as the function of aromatic residue distribution. This result is consistent with previous studies that demonstrate the importance of aromatic residue patterning on the phase behaviour of PLDs [9, 98].

Figure 7. Prion-like domains featuring sequences with dispersed aromatic residues show greater propensities for phase separation. A Representation of the aromatic amino acids present in the wild type sequences of each PLD considered **B** Location of the positively charged (red) and negatively charged (blue) amino acids present in the wild type sequences of each PLD considered **C** Analysis of the critical solution temperature versus the σ order parameter, defined in Equation 13. Those groups of variants with the same 13 value show their average critical temperature highlighted, and softer values for other data points. For lower values of σ, the critical temperature present in the condensates is higher, and vice versa, indicating that those variants with a more homogeneous distribution of aromatic amino acids have condensates more stable at higher temperatures.

Discussion

Computer simulations have advanced our ability to map emergent properties of protein solutions to the chemical make-up of their individual biomolecules. For instance, modern residue-resolution protein coarse-grained models [32, 74, 99–103] can now be used to quantify the modulation of protein phase behavior as a result of amino acid sequence mutations of their native proteins with near-quantitative agreement with experiments [32, 101, 102]. In this study, we exploit the accuracy and efficiency of our protein model, Mpipi [32], to compute and analyze the temperature-vs-density phase diagrams of a large set of PLD proteins—140 variants of the proteins hnRNPA1, TDP43, FUS, EWSR1, RBM14, and TIA1. Collectively, our set of phase diagrams reveal conserved scaling laws

that quantitatively predict the change in the critical solution temperatures of single-component PLD solutions as a function of the number and type of amino acid mutations of the PLDs.

To facilitate the interpretation of the scaling laws derived in our study, we have organized the data in a table that illustrates the effect of single point mutations on the critical solution temperatures of the family of PLDs, as a function of the PLD length.

L	Y $\rightarrow$ F	F/Y $\rightarrow$ W	F/Y $\rightarrow$ X	R $\rightarrow$ K	R $\rightarrow$ X
100	-0.4 $\pm$ 0.04	4.3 $\pm$ 0.1	-5.6 $\pm$ 0.5	-13 $\pm$ 1	-6.4 $\pm$ 0.5
200	-0.4 $\pm$ 0.04	4.3 $\pm$ 0.1	-3.9 $\pm$ 0.4	-6.5 $\pm$ 0.5	-3.2 $\pm$ 0.3
300	-0.4 $\pm$ 0.04	4.3 $\pm$ 0.1	-3.2 $\pm$ 0.3	-4.3 $\pm$ 0.3	-2.1 $\pm$ 0.2

L	N $\rightarrow$ Q	S $\rightarrow$ T	G $\rightarrow$ T	A $\rightarrow$ S
100	0.5 $\pm$ 0.05	-0.09 $\pm$ 0.03	-0.08 $\pm$ 0.02	0.16 $\pm$ 0.05
200	0.25 $\pm$ 0.03	-0.09 $\pm$ 0.03	-0.08 $\pm$ 0.02	0.16 $\pm$ 0.05
300	0.17 $\pm$ 0.02	-0.09 $\pm$ 0.03	-0.08 $\pm$ 0.02	0.16 $\pm$ 0.05

Table 1. Predicted effect of single point mutations on the critical temperature, in Kelvin, of PLDs, as a function of their length, L

Consistent with experimental evidence and computational predictions, interactions involving aromatic residues—namely, π–π and cation–π—are dominant in stabilizing condensates formed by PLDs [8, 32, 33]. Notably, our simulations reveal the largest decrease in the critical solution temperature of a PLD condensate occurs when an aromatic residue is replaced by an uncharged, non-aromatic amino acid. Our scaling law for such aromatic deletions ($56\,\mathrm{K}/\sqrt{L}$) predicts that the destabilizing effect of replacing an aromatic with a non-aromatic residue is linearly amplified by the number of such mutations and screened by the square root of the length of the protein ($\sqrt{L}$), indicating a disruption in the percolated network of interactions sustaining the condensate. Condensates made of longer PLDs are more robust against aromatic deletions because, by construction, each aromatic residue represents a smaller percentage of the total valency of the PLD in question. In other words, longer PLDs are higher valency molecules than their shorter counterparts and can, thus, form a densely connected percolating condensed-liquid network, even when a few of their aromatic stickers have been deleted.

Our findings also highlight the crucial role of Arg in the critical parameters of PLD condensates. Specifically, our scaling law for Arg deletions ($640\,\mathrm{K}/L$) predicts that the destabilizing effect of an Arg mutation to an uncharged, non-aromatic residue is inversely proportional to the length of the protein (L), as well as linear with respect to the number of mutations. More specifically, it indicates that, for a PLD of the length of 100 amino acids, the change in the critical temperature will be of the order of 6.4 K. While formulaically similar to the scaling law for aromatic deletions, a notable difference is a change in the dependence on the PLD length. In the case of Arg deletions, the effect of mutations is screened by L rather than $\sqrt{L}$, indicating a stronger screening dependence on the PLD length. This indicates that for long PLDs, the destabilizing effects due to Arg deletions are diminished when compared to aromatic deletions, while in shorter PLDs this trend is reversed with each Arg deletion exerting a more significant destabilizing effect than the aromatic deletions.

These results underscore once again how for PLDs cation–π and π–π interactions are the main drivers of phase separation [9, 30, 31, 33, 35, 36, 74–76]. While it may seem counterintuitive for Arg deletions to have a lesser effect in the critical parameters of longer PLDs than their aromatic counterparts (as Arg residues are known to form strong cation–π interactions [104, 105]) the explanation lies in the type of systems in question, which are particularly rich in aromatic residues. Longer PLDs have, as a result of their higher valency, a larger network of possible pairwise interactions, which can mitigate the destabilizing effect of Arg deletions. In the case of shorter PLDs, each Arg residue represents a larger proportion of the total valency, and thus its deletion has a proportionally larger impact on the critical parameters. In contrast, aromatic residues contribute

to the stability of the condensate through both cation–π and π–π interactions, which are predicted to be weaker than their cation–π analogues [31, 32], but more numerous in PLDs. In a longer PLD, the deletion of an aromatic residue results in the loss of more potential interactions, creating a larger destabilizing effect than the deletion of an Arg residue. However, in shorter PLDs, due to their lower total valency, the loss of potential associative interactions that result from the deletion of an aromatic residue is lower. Hence, the destabilizing effect resulting from this mutation is comparatively less than for the Arg deletion.

Our scaling laws have also allowed us to quantify the destabilizing effect of Arg to Lys mutations in the phase separation of PLDs. Concretely, mutations of Arg residues to Lys result in an overall decrease in the critical temperature of phase separation, two-fold in comparison to the impact of mutating Arg into an alternate uncharged non-aromatic residue. Consequently, the scaling law considerably intensifies for mutations from Arg to Lys, nearly doubling the effects of each mutation in the critical temperature of the condensate (See Table 1). This implies that each mutation of Arg to Lys exerts a substantially more destabilizing effect than mutations of Arg to other uncharged, non-aromatic residues, consistent with experimental evidence [30, 33, 83, 106] and highlighting the significant, modulatory role Lys plays in the phase behavior of PLDs.

Uncharged, non-aromatic residues such as Ser, Gly, Asn, Gln, and Thr are found to play a marginal role in PLD condensate stability [33, 37, 68, 72]. Notably, PLD condensates are highly stable in the face of Gly, Ser, Thr and Ala mutations. This behavior is quantified by their respective scaling laws, with values for the scaling constants for mutations from Gly to Thr, Ser to Thr and Ala to Ser at $-0.09\,\mathrm{K}$, $-0.08\,\mathrm{K}$ and $0.16\,\mathrm{K}$ respectively. These numerical values indicate that, for a change of one degree in the critical temperature, approximately 10 amino acids of this set would need to undergo mutation. Notably, these scaling laws do not display a significant length dependency. This observation aligns with the idea that these mutations primarily affect the condensate properties through modulation in excluded volume and solvation, rather than by interrupting specific interaction networks.

In contrast, the condensates display heightened sensitivity towards Asn and Gln mutations, underscored by the Asn to Gln scaling of $50\,\mathrm{K}/L$. This law is also inversely length dependent, indicating that mutations on these amino acids affect interaction networks and that mutations Asn to Gln result in a heightened ability of the system to phase separate. We hypothesize that, in addition to differences in solvation volume, the polarity of each amino acid plays a role by forming charge–dipole and dipole–dipole interactions. Both Gln and Asn are polar amino acids, with Gln having a longer side chain than Asn, which could potentially interact with a larger number of nearby atoms. Furthermore, the additional methylene group (-CH2-) may slightly alter the electron distribution around the amide group, which can lead to the formation of more stable dipole–dipole contacts. It is important to note that our computational model, Mpipi, is a charge-fixed model, and hence does not directly capture the effects of polarization. However, the lack of physical details in the Mpipi model is compensated by the parameters having been derived from bioinformatics data.

While the Mpipi model has proven to be a powerful tool in predicting phase behaviour of PLDs, it is vitally important to acknowledge its limitations. Notably, the model does not account for temperature-dependent dependencies related to solvent effects, such as hydrophobic interactions and charge-mediated interactions facilitated by desolvation penalties, which implies that the model might not accurately predict the effects of mutations in particular sequences, such as sequences high in proline or aliphatic residues, where the interaction landscape is temperature sensitive. Additionally, the Mpipi model is not predictive of folding, or secondary structure. An assumption we made was that the mutations introduced do not significantly alter the secondary structure of each PLD, which is a necessary simplification for the scope of this study.

Collectively, the definition of a set of scaling laws acts as a unifying framework for the study of the critical parameters of PLD phase separation, by allowing for direct comparison of individual sequence effects. Such laws are cumulative, meaning that the effects of successive mutations can be predicted by the addition of the appropriate laws, as long as the resulting mutation keeps within

PLD composition and patterning biases.

Our work explores a new framework to describe, quantitatively, the phase behavior of a large set of protein variants via the identification of scaling laws. Our scaling laws can predict changes in the stability of condensates based on amino acid sequence mutations. This code not only allows for testing the effects of composition bias on PLD phase separation but also serves as a predictive tool, allowing for a better selection of mutations to achieve desired changes in different condensates. This study is focused on PLDs, but we hypothesize similar scaling behavior in other IDPs, with the laws having a dependence on the protein length that reflects the nature of the underlying interaction networks of the system. Our findings have important implications for understanding PLD condensation and for designing modifications to counter their aberrant aggregation, by setting the basis for characterizing and quantifying the molecular determinants of PLD phase behavior.

Methods and Materials

Design of the dataset

To design sequence variants to test the phase behavior of prion-like domains, we use in-silico targeted mutagenesis to introduce specific changes to the amino acid sequence of the PLD of interest. These changes involve substituting different amino acids, while carefully maintaining patterning and compositional biases, to test the role of given amino acids in the phase-separating propensities of the respective PLDs.

The nomenclature employed is consistent with that of Bremer and colleagues [33]. Namely, if n amino acids of type X are removed from the sequence and replaced with n amino acids of type Y, the variant is labeled $-nX + nY$. If instead, n amino acids of type X are removed from the sequence and replaced with amino acids (often different types) in proportion to the compositional bias of the PLD, the variant is simply labeled $-nX$.

To begin, we select a family of PLDs composed of hNRNPA1, RBM14, FUS, TDP43, TIA1 and EWSR1 and perform mutations on each sequence to generate a set of variants. Accordingly, we assess whether the effects of sequence mutations in PLD phase behaviors hold for the whole family of PLDs. Furthermore, our data set allows us to develop a set of rules that can aid in understanding and predicting PLD phase behavior. The amino acid sequences of the PLDs, and their corresponding variants, investigated herein are provided in the Supporting Information.

In total, we compute phase diagrams of 140 PLD variants via molecular dynamics (MD). This task required a large amount of CPU hours, over a 1.25 million in total. Given these demands, efficient utilization of high-performance computing (HPC) resources and proper parallelization of the computational tasks are paramount to ensuring optimal performance. To this end, we compute MD trajectories using the LAMMPS (Large-scale Atomic/Molecular Massively Parallel Simulator) software [107], exploiting its efficient parallelization algorithms. Another significant computational challenge involves the large size of data generated. Each single run produced a significant amount of trajectory, backup, and analysis data, and a large number of files. On average, each run produced over 50 files and a minimum of 5 GB of data. To efficiently handle this, we implement a robust, tiered, data storage and retrieval system, ensuring that the vast amounts of data is organized through a combination of indexing and partitioning, enhancing search efficiency and accessibility. We leveraged the Lustre filesystem for its high performance and scalability, making it the ideal choice for managing and accessing large volumes of active data required by our computational jobs. Custom scripts are employed for automated data processing, enabling us to extract meaningful insights from the complex datasets. All scripts are available in our GitHub repository.

Critical Parameter Determination Through MD

We used MD simulations to determine the full binodals (i.e., the boundaries that delimit the two-phase region from the one-phase region) for the PLDs. Specifically, we perform MD simulations in the canonical ensemble – i.e., fixed volume (V), temperature (T), and number of particles (N).

For each binodal, a series of independent NVT simulations are performed at different tempera-
tures and the density of each of the coexisting phases are determined. Thus, this approach yields
binodals in the temperature–density phase space.

Direct Coexistence Simulations

Specifically, we use Direct Coexistence simulations, which enables the simultaneous simulation of
both high- and low-density phases of a mixture, separated by an interface, within a single simula-
tion box. Additionally, these simulations are performed with periodic boundary conditions in each
dimension.

To initiate the NVT Direct Coexistence simulations, we start with one copy of a PLD in a cu-
bic simulation cell and replicate it to the target number of chains (64 in total). We then iteratively
scaled the simulation box to compress the chains into a high density direct coexistence simulation
(approx. 0.8–1 g/cm³) [100]. Here, we use a Langevin thermostat [108] (at the target temperature,
with a damping time of 40 ps) for temperature control. Keeping the length of the box in the X and
Y dimensions fixed, we then elongate the Z dimension of the box. We then simulate each system
at the target temperature for over 200 ns (see Convergence test in the SI). From each simulation
we compute the density profile in the Z direction and determine the density of the dilute and con-
densed phases. These data are then used to produce the binodals.

Mpipi model

All the simulations are run using the Mpipi coarse-grained force-field [32] with an implicit solvent.
In the Mpipi model electrostatic, cation–π and π–π interactions are carefully balanced to capture
the phase behavior of PLDs. Notably, Mpipi achieves quantitative agreement with experiments in
terms of prediction of critical solution temperatures of such systems. In Mpipi, each amino acid is
represented by a single unique bead – i.e., of corresponding mass, molecular diameter (σ), charge
(q), and an energy scale that reflects the relative frequency of planar π–π contacts (ϵ) [32].

The potential energy in the Mpipi model is computed as follows:

$$E_{\text{mpipi}} = E_{\text{bond}} + E_{\text{elec}} + E_{\text{pair}}. \tag{16}$$

Beads are connected via harmonic springs

$$E_{\text{bond}} = \sum_i \frac{1}{2} k(r_i - r_i^0)^2. \tag{17}$$

Here k is the bond force constant (19.1 kcal mol^{-1} Å^{-2}) and r_i^0 is the equilibrium bond length,
which is set to 3.81 Å.

The electrostatic contribution is modeled via a Coulomb interaction term, with Debye–Hückel
screening.

$$E_{\text{elec}} = \sum_{i,j} \frac{q_i q_j}{4\pi \epsilon_r \epsilon_0 r_{ij}} \exp{(-\kappa r_{ij})}, \tag{18}$$

where q_i and q_j are the charges of the bead i and j, respectively. ϵ_0 is the electric constant and
$\epsilon_r = 80$ is the relative dielectric constant of water. $\kappa = 0.126$ Å^{-1} is the inverse Debye screening
length, which corresponds to monovalent salt concentration of 150 mM.

Finally, short-ranged non-bonded interactions are modeled with the Wang–Frenkel potential

$$E_{\text{pair}} = \sum_{i,j} \phi_{i,j} = \sum_{i,j} \epsilon_{i,j} \alpha_{i,j} \left[\left(\frac{\sigma_{i,j}}{r} \right)^{2\mu_{i,j}} - 1 \right] \left[\left(\frac{\sigma_{i,j}}{r} \right)^{2\mu_{i,j}} - 1 \right]^{2\nu_{i,j}} \tag{19}$$

where E_{pair} is the potential energy between all pairs of amino acids, $\phi_{i,j}$ is the potential energy
between the pairs of amino acids i and j, ϵ_{ij} is an energy parameter that dictates the strength of
interaction of that particular pair, $\sigma_{i,j}$ is average the Van der Waals radii of the pair that dictate the
position of the energy minima, r is the distance between center of masses and $\nu_{i,j}$ and $\mu_{i,j}$ are fitted

exponential parameters that determine the shape of the energy curve. Full details of the Mpipi model are described in [32].

Data fitting

To estimate the critical temperatures of the binodals computed in Section , we use the law of coexistence densities

$$\left(\rho_{\text{high}}(T) - \rho_{\text{low}}(T)\right)^{3.06} = d\left(1 - \frac{T}{T_{\text{c}}}\right) \tag{20}$$

The constant 3.06 in the equation is a dimensionless empirical factor that was derived from simulations of the 3D Ising model. The critical densities are computed by assuming that the law of rectilinear diameters holds, namely

$$\rho_{\text{high}}(T) + \rho_{\text{low}}(T) = 2\rho_c + 2A(T - T_{\text{c}}) \tag{21}$$

where $\rho_{\text{high}}(T)$, $\rho_{\text{low}}(T)$ and ρ_{c} are the densities of the high-density and low-density phases and the critical density, respectively, T_{c} is the critical temperature, and d and A are fitting parameters.

Finite-size scaling

In direct coexistence simulations in the slab geometry, finite-size effects can arise from either the cross-sectional area (i.e., size of the interface) or from the long axis (i.e., fraction bulk condensate). To ensure our calculations are robust with respect to system size, we perform finite-size scaling at fixed system number density: (1) varying the size of the cross section and (2) varying the length of the long axis. Using these tests, we confirm that the phase diagrams computed for systems of varying sizes (27, 64, 125 and 216 chains) exhibit no significant difference in predicted densities (see Figure 2 in Supporting Information). Our scaling analysis revealed that 64 chains in a box of dimensions 125x125x1000 is the minimum size that ensures proteins do not interact with their periodic images and that the condensed phase is large enough to represent a true bulk phase.

Acknowledgments

We thank Pin Yu Chew for useful comments on the manuscript. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 803326 to R.C.G.). MJM would like to acknowledge the Winton Programme for Physics of Sustainability for doctoral funding. AAG is funded by the ERC (grant agreement No 803326). JRE acknowledges funding from the Ramon y Cajal fellowship (RYC2021-030937-I) and the Spanish National Agency for Research under the grant PID2022-136919NA-C33. This work was supported by the Engineering and Physical Sciences Research Council (EPSRC) [grant number EP/X02332X/1 to JH] under the UK Research and Innovation (UKRI) Post-doctoral Fellowships Guarantee Scheme [project TF-CHROM-LLPS]. JAJ acknowledges research support from departmental start-up funds provided by the Department of Chemical and Biological Engineering and the Omenn–Darling Bioengineering Institute at Princeton University. This project made use of time on HPC granted via the UK High-End Computing Consortium for Biomolecular Simulation, HECBioSim (http://hecbiosim.ac.uk), supported by EPSRC (grant no. EP/R029407/1).

Code and Data Availability

The source code for running the Mpipi model in LAMMPS, example input scripts, as well as instructions for reproducing the data through a demo can be found at the GitHub repository: https://github.com/JuliMaristany/PLD_Scaling_Laws. The data supporting the findings of this manuscript will be available via a figshare repository.

References

[1] Salman F. Banani, Hyun O. Lee, Anthony A. Hyman, and Michael K. Rosen. Biomolecular condensates: Organizers of cellular biochemistry. *Nature Reviews Molecular Cell Biology*, 18(5):285–298, 5 2017.

[2] Pilong Li, Sudeep Banjade, Hui-Chun Cheng, Soyeon Kim, Baoyu Chen, Liang Guo, Marc Llaguno, Javoris V Hollingsworth, David S King, Salman F Banani, et al. Phase transitions in the assembly of multivalent signalling proteins. *Nature*, 483(7389):336–340, 2012.

[3] Tanja Mittag and Rohit V Pappu. A conceptual framework for understanding phase separation and addressing open questions and challenges. *Molecular cell*, 82(12):2201–2214, 2022.

[4] Yongdae Shin and Clifford P. Brangwynne. Liquid phase condensation in cell physiology and disease. *Science*, 357(6357):eaaf4382, 9 2017.

[5] Edward Gomes and James Shorter. The molecular language of membraneless organelles. *Journal of Biological Chemistry*, 294(18):7115–7127, 2019.

[6] Timothy J Nott, Evangelia Petsalaki, Patrick Farber, Dylan Jervis, Eden Fussner, Anne Plochowietz, Timothy D Craggs, David P Bazett-Jones, Tony Pawson, Julie D Forman-Kay, et al. Phase transition of a disordered nuage protein generates environmentally responsive membraneless organelles. *Molecular cell*, 57(5):936–947, 2015.

[7] Jarrett Smith, Deepika Calidas, Helen Schmidt, Tu Lu, Dominique Rasoloson, and Geraldine Seydoux. Spatial patterning of p granules by rna-induced phase separation of the intrinsically-disordered protein meg-3. *Elife*, 5:e21337, 2016.

[8] Jie Wang, Jeong Mo Choi, Alex S. Holehouse, Hyun O. Lee, Xiaojie Zhang, Marcus Jahnel, Shovamayee Maharana, Régis Lemaitre, Andrei Pozniakovsky, David Drechsel, Ina Poser, Rohit V. Pappu, Simon Alberti, and Anthony A. Hyman. A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins. *Cell*, 174(3):688–699, 7 2018.

[9] Erik W Martin, Alex S Holehouse, Ivan Peran, Mina Farag, J Jeremias Incicco, Anne Bremer, Christy R Grace, Andrea Soranno, Rohit V Pappu, and Tanja Mittag. Valence and patterning of aromatic residues determine the phase behavior of prion-like domains. *Science*, 367(6478):694–699, 2020.

[10] Erik W Martin and Alex S Holehouse. Intrinsically disordered protein regions and phase separation: sequence determinants of assembly or lack thereof. *Emerging topics in life sciences*, 4(3):307—329, December 2020.

[11] Ammon E Posey, Alex S Holehouse, and Rohit V Pappu. Phase separation of intrinsically disordered proteins. 611:1–30, 2018.

[12] Shana Elbaum-Garfinkle, Younghoon Kim, Krzysztof Szczepaniak, Carlos Chih-Hsiung Chen, Christian R Eckmann, Sua Myong, and Clifford P Brangwynne. The disordered p granule protein laf-1 drives phase separation into droplets with tunable viscosity and dynamics. *Proceedings of the National Academy of Sciences*, 112(23):7189–7194, 2015.

[13] Titus M Franzmann, Marcus Jahnel, Andrei Pozniakovsky, Julia Mahamid, Alex S Holehouse, Elisabeth Nüske, Doris Richter, Wolfgang Baumeister, Stephan W Grill, Rohit V Pappu, et al. Phase separation of a yeast prion protein promotes cellular fitness. *Science*, 359(6371):eaao5654, 2018.

[14] Shovamayee Maharana, Jie Wang, Dimitrios K Papadopoulos, Doris Richter, Andrey Pozniakovsky, Ina Poser, Marc Bickle, Sandra Rizk, Jordina Guillén-Boixet, Titus M Franzmann, et al. Rna buffers the phase separation behavior of prion-like rna binding proteins. *Science*, 360(6391):918–921, 2018.

[15] Nieves Lorenzo Gotor, Alexandros Armaos, Giulia Calloni, Marc Torrent Burgas, R Martin Vabulas, Natalia Sanchez De Groot, and Gian Gaetano Tartaglia. Rna-binding and prion domains: the yin and yang of phase separation. *Nucleic acids research*, 48(17):9491–9504, 2020.

[16] Amandine Molliex, Jamshid Temirov, Jihun Lee, Maura Coughlin, Anderson P. Kanagaraj, Hong Joo Kim, Tanja Mittag, and J. Paul Taylor. Phase Separation by Low Complexity Domains Promotes Stress Granule Assembly and Drives Pathological Fibrillization. *Cell*, 163(1):123–133, 9 2015.

[17] Louise Jawerth, Elisabeth Fischer-Friedrich, Suropriya Saha, Jie Wang, Titus Franzmann, Xiaojie Zhang, Jenny Sachweh, Martine Ruer, Mahdiye Ijavi, Shambaditya Saha, et al. Protein condensates as aging maxwell fluids. *Science*, 370(6522):1317–1323, 2020.

[18] Ibraheem Alshareedah, Wade M Borcherds, Samuel R Cohen, Mina Farag, Anurag Singh, Anne Bremer, Rohit V Pappu, Tanja Mittag, and Priya R Banerjee. Sequence-encoded grammars determine material properties and physical aging of protein condensates. *bioRxiv*, pages 2023–04, 2023.

[19] Miriam Linsenmeier, Maria Hondele, Fulvio Grigolato, Eleonora Secchi, Karsten Weis, and Paolo Arosio. Dynamic arrest and aging of biomolecular condensates are modulated by low-complexity domains, rna and biochemical activity. *Nature Communications*, 13(1):3030, 2022.

[20] Macy L Sprunger and Meredith E Jackrel. Prion-like proteins in phase separation and their link to disease. *Biomolecules*, 11(7):1014, July 2021.

[21] SF Banani. Compositional control of phase-separated cell. bodies. *Cell*, 166:651–663, 2016.

[22] Jie Wang, Jeong Mo Choi, Alex S. Holehouse, Hyun O. Lee, Xiaojie Zhang, Marcus Jahnel, Shovamayee Maharana, Régis Lemaitre, Andrei Pozniakovsky, David Drechsel, Ina Poser, Rohit V. Pappu, Simon Alberti, and Anthony A. Hyman. A Molecular Grammar Governing the Driving Forces for Phase Separation of Prion-like RNA Binding Proteins. *Cell*, 174(3):688–699, 7 2018.

[23] Jeong Mo Choi, Alex S. Holehouse, and Rohit V. Pappu. Physical Principles Underlying the Complex Biology of Intracellular Phase Transitions. *Annual Review of Biophysics*, 49:107–133, 5 2020.

[24] Jeong-Mo Choi, Anthony A Hyman, and Rohit V Pappu. Generalized models for bond percolation transitions of associative polymers. *Physical Review E*, 102(4):042403, 2020.

[25] Tyler S Harmon, Alex S Holehouse, Michael K Rosen, and Rohit V Pappu. Intrinsically disordered linkers determine the interplay between phase separation and gelation in multivalent proteins. *elife*, 6:e30294, 2017.

[26] Christiane Iserman, Christine Desroches Altamirano, Ceciel Jegers, Ulrike Friedrich, Taraneh Zarin, Anatol W Fritsch, Matthäus Mittasch, Antonio Domingues, Lena Hersemann, Marcus Jahnel, et al. Condensation of ded1p promotes a translational switch from housekeeping to stress protein production. *Cell*, 181(4):818–831, 2020.

[27] Omar Adame-Arana, Christoph A Weber, Vasily Zaburdaev, Jacques Prost, and Frank Jülicher. Liquid phase separation controlled by ph. *Biophysical Journal*, 119(8):1590–1605, 2020.

[28] Matthias Christoph Munder, Daniel Midtvedt, Titus Franzmann, Elisabeth Nuske, Oliver Otto, Maik Herbig, Elke Ulbricht, Paul Müller, Anna Taubenberger, Shovamayee Maharana, Liliana Malinovska, Doris Richter, Jochen Guck, Vasily Zaburdaev, and Simon Alberti. A ph-driven transition of the cytoplasm from a fluid- to a solid-like state promotes entry into dormancy. *eLife*, 5:e09347, mar 2016.

[29] Hasan Cinar, Zamira Fetahaj, Süleyman Cinar, Robert M. Vernon, Hue Sun Chan, and Roland H. A. Winter. Temperature, hydrostatic pressure, and osmolyte effects on liquid–liquid phase separation in protein condensates: Physical chemistry and biological implications. *Chemistry – A European Journal*, 25(57):13049–13069, 2019.

[30] Rachel S Fisher and Shana Elbaum-Garfinkle. Tunable multiphase dynamics of arginine and lysine liquid condensates. *Nature communications*, 11(1):4628, 2020.

[31] Georg Krainer, Timothy J. Welsh, Jerelle A. Joseph, Jorge R. Espinosa, Sina Wittmann, Ella de Csilléry, Akshay Sridhar, Zenon Toprakcioglu, Giedre Gudiškytė, Magdalena A. Czekalska, William E. Arter, Jordina Guillén-Boixet, Titus M. Franzmann, Seema Qamar, Peter St George-Hyslop, Anthony A. Hyman, Rosana Collepardo-Guevara, Simon Alberti, and Tuomas P.J. Knowles. Reentrant liquid condensate phase of proteins is stabilized by hydrophobic and non-ionic interactions. *Nature Communications*, 12(1):1–14, 12 2021.

[32] Jerelle A. Joseph, Aleks Reinhardt, Anne Aguirre, Pin Yu Chew, Kieran O. Russell, Jorge R. Espinosa, Adiran Garaizar, and Rosana Collepardo-Guevara. Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy. *Nature Computational Science*, 1(11):732–743, 2021.

[33] Anne Bremer, Mina Farag, Wade M. Borcherds, Ivan Peran, Erik W. Martin, Rohit V. Pappu, and Tanja Mittag. Deciphering how naturally occurring sequence features impact the phase behaviors of disordered prion-like domains. *Nature chemistry*, 14(2):196, 2 2022.

[34] Jacob P Brady, Patrick J Farber, Ashok Sekhar, Yi-Hsuan Lin, Rui Huang, Alaji Bah, Timothy J Nott, Hue Sun Chan, Andrew J Baldwin, Julie D Forman-Kay, et al. Structural and hydrodynamic properties of an intrinsically disordered region of a germ cell-specific protein on phase separation. *Proceedings of the National Academy of Sciences*, 114(39):E8194–E8203, 2017.

[35] RM Vernon. Pi–Pi contacts are an overlooked protein feature relevant to phase separation. *eLife*, 7, 2018.

[36] Martin J Fossat, Xiangze Zeng, and Rohit V Pappu. Uncovering differences in hydration free energies and structures for model compound mimics of charged side chains of amino acids. *The Journal of Physical Chemistry B*, 125(16):4148–4161, 2021.

[37] Erik W Martin and Tanja Mittag. Relationship of sequence and phase separation in protein low-complexity regions. *Biochemistry*, 57(17):2478–2487, 2018.

[38] R. C. Ball. *Scaling in Polymer Physics*, pages 483–490. Springer US, Boston, MA, 1991.

[39] Dave E. Dunstan. The viscosity-radius relationship for concentrated polymer solutions. *Scientific Reports*, 9(1), January 2019.

[40] Suman Majumder, Henrik Christiansen, and Wolfhard Janke. Understanding nonequilibrium scaling laws governing collapse of a polymer. *The European Physical Journal B*, 93:1–19, 2020.

[41] Michael Rubinstein, Ralph H Colby, et al. *Polymer physics*, volume 23. Oxford university press New York, 2003.

[42] Hagen Hofmann, Andrea Soranno, Alessandro Borgia, Klaus Gast, Daniel Nettels, and Benjamin Schuler. Polymer scaling laws of unfolded and intrinsically disordered proteins quantified with single-molecule spectroscopy. *Proceedings of the National Academy of Sciences*, 109(40):16155–16160, 2012.

[43] Joseph P Clarke, Patricia A Thibault, Hannah E Salapa, and Michael C Levin. A comprehensive analysis of the role of hnrnp a1 function and dysfunction in the pathogenesis of neurodegenerative disease. *Frontiers in Molecular Biosciences*, 8:659610, 2021.

[44] Hong Joo Kim, Nam Chul Kim, Yong-Dong Wang, Emily A Scarborough, Jennifer Moore, Zamia Diaz, Kyle S MacLea, Brian Freibaum, Songqing Li, Amandine Molliex, et al. Mutations in prion-like domains in hnrnpa2b1 and hnrnpa1 cause multisystem proteinopathy and als. *Nature*, 495(7442):467–473, 2013.

[45] I-Fan Wang, Lien-Szu Wu, and CK James Shen. Tdp-43: an emerging new player in neurodegenerative diseases. *Trends in molecular medicine*, 14(11):479–485, 2008.

[46] Myungjin Jo, Shinrye Lee, Yu-Mi Jeon, Seyeon Kim, Younghwi Kwon, and Hyung-Jun Kim. The role of tdp-43 propagation in neurodegenerative diseases: integrating insights from clinical and experimental studies. *Experimental & molecular medicine*, 52(10):1652–1662, 2020.

[47] Peter EA Ash, Shuwen Lei, Jenifer Shattuck, Samantha Boudeau, Yari Carlomagno, Maria Medalla, Bryce L Mashimo, Guillermo Socorro, Louloua FA Al-Mohanna, Lulu Jiang, et al. Tia1 potentiates tau phase separation and promotes generation of toxic oligomeric tau. *Proceedings of the National Academy of Sciences*, 118(9):e2014188118, 2021.

[48] Daniel J Apicco, Peter EA Ash, Brandon Maziuk, Chelsey LeBlang, Maria Medalla, Ali Al Abdullatif, Antonio Ferragud, Emily Botelho, Heather I Ballance, Uma Dhawan, et al. Reducing the rna binding protein tia1 protects against tau-mediated neurodegeneration in vivo. *Nature neuroscience*, 21:72–80, 2018.

[49] Hao Deng, Kai Gao, and Joseph Jankovic. The role of fus gene variants in neurodegenerative diseases. *Nature Reviews Neurology*, 10(6):337–348, 2014.

[50] Clotilde Lagier-Tourenne and Don W Cleveland. Rethinking als: The fus about tdp-43. *Cell*, 136(6):1001–1004, 2009.

[51] Julien Couthouis, Michael P Hart, Renske Erion, Oliver D King, Zamia Diaz, Tadashi Nakaya, Fadia Ibrahim, Hyung-Jun Kim, Jelena Mojsilovic-Petrovic, Saarene Panossian, et al. Evaluating the role of the fus/tls-related gene ewsr1 in amyotrophic lateral sclerosis. *Human molecular genetics*, 21(13):2899–2911, 2012.

[52] Junghee Lee, Phuong T Nguyen, Hyun Soo Shim, Seung Jae Hyeon, Hyeonjoo Im, Mi-Hyun Choi, Sooyoung Chung, Neil W Kowall, Sean Bong Lee, and Hoon Ryu. Ewsr1, a multifunctional protein, regulates cellular function and aging via genetic and epigenetic pathways. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 1865(7):1938–1945, 2019.

[53] Yun R Li, Oliver D King, James Shorter, and Aaron D Gitler. Stress granules as crucibles of als pathogenesis. *Journal of cell biology*, 201(3):361–372, 2013.

[54] Mina Farag, Wade M. Borcherds, Anne Bremer, Tanja Mittag, and Rohit V. Pappu. Phase separation in mixtures of prion-like low complexity domains is driven by the interplay of homotypic and heterotypic interactions. *bioRxiv*, 2023.

[55] Stefan Reber, Daniel Jutzi, Helen Lindsay, Anny Devoy, Jonas Mechtersheimer, Brunno Rocha Levone, Michal Domanski, Eva Bentmann, Dorothee Dormann, Oliver Mühlemann, Silvia M L Barabino, and Marc-David Ruepp. The phase separation-dependent FUS interactome reveals nuclear and cytoplasmic function of liquid–liquid phase separation. *Nucleic Acids Research*, 49(13):7713–7731, July 2021.

[56] Zhe Feng, Xudong Chen, Xiandeng Wu, and Mingjie Zhang. Formation of biological condensates via phase separation: Characteristics, analytical methods, and physiological implications. *Journal of Biological Chemistry*, 294(40):14823–14835, 2019.

[57] Timothy J Welsh, Georg Krainer, Jorge R Espinosa, Jerelle A Joseph, Akshay Sridhar, Marcus Jahnel, William E Arter, Kadi L Saar, Simon Alberti, Rosana Collepardo-Guevara, et al. Surface electrostatics govern the emulsion stability of biomolecular condensates. *Nano letters*, 22(2):612–621, 2022.

[58] Feng Luo, Xinrui Gui, Heng Zhou, Jinge Gu, Yichen Li, Xiangyu Liu, Minglei Zhao, Dan Li, Xueming Li, and Cong Liu. Atomic structures of FUS LC domain segments reveal bases for reversible amyloid fibril formation. *Nat. Struct. Mol. Biol.*, 25(4):341–346, April 2018.

[59] Dylan T Murray, Masato Kato, Yi Lin, Kent R Thurber, Ivan Hung, Steven L McKnight, and Robert Tycko. Structure of FUS protein fibrils and its relevance to self-assembly and phase separation of low-complexity domains. *Cell*, 171(3):615–627.e16, October 2017.

[60] Seema Qamar, GuoZhen Wang, Suzanne J Randle, Francesco Simone Ruggeri, Juan A Varela, Julie Qiaojin Lin, Emma C Phillips, Akinori Miyashita, Declan Williams, Florian Ströhl, et al. Fus phase separation is modulated by a molecular chaperone and methylation of arginine cation-π interactions. *Cell*, 173(3):720–734, 2018.

[61] Tetsuro Murakami, Seema Qamar, Julie Qiaojin Lin, Gabriele S Kaminski Schierle, Eric Rees, Akinori Miyashita, Ana R Costa, Roger B Dodd, Fiona TS Chan, Claire H Michel, et al. Als/ftd mutation-induced phase transition of fus liquid droplets and reversible hydrogels into irreversible hydrogels impairs rnp granule function. *Neuron*, 88(4):678–690, 2015.

[62] Avinash Patel, Hyun O Lee, Louise Jawerth, Shovamayee Maharana, Marcus Jahnel, Marco Y Hein, Stoyno Stoynov, Julia Mahamid, Shambaditya Saha, Titus M Franzmann, et al. A liquid-to-solid phase transition of the als protein fus accelerated by disease mutation. *Cell*, 162(5):1066–1077, 2015.

[63] Anastasia C Murthy, Gregory L Dignon, Yelena Kan, Gül H Zerze, Sapun H Parekh, Jeetain Mittal, and Nicolas L Fawzi. Molecular interactions underlying liquid- liquid phase separation of the fus low-complexity domain. *Nature structural & molecular biology*, 26(7):637–648, 2019.

[64] Auré Lie Zbinden, Manuela Pé Rez-Berlanga, Pierre De Rossi, and Magdalini Polymenidou. Phase Separation and Neurodegenerative Diseases: A Disturbance in the Force. *Developmental Cell*, 55:45–68, 2020.

[65] Elizabeth L Guenther, Qin Cao, Hamilton Trinh, Jiahui Lu, Michael R Sawaya, Duilio Cascio, David R Boyer, Jose A Rodriguez, Michael P Hughes, and David S Eisenberg. Atomic structures of tdp-43 lcd segments and insights into reversible or pathogenic aggregation. *Nature structural & molecular biology*, 25(6):463–471, 2018.

[66] Jeffrey M Lotthammer, Garrett M Ginell, Daniel Griffith, Ryan Emenecker, and Alex S Holehouse. Direct prediction of intrinsically disordered protein conformational properties from sequence. *Biophysical Journal*, 123(3):43a, 2024.

[67] Sven Hennig, Geraldine Kong, Taro Mannen, Agata Sadowska, Simon Kobelke, Amanda Blythe, Gavin J. Knott, K. Swaminathan Iyer, Diwei Ho, Estella A. Newcombe, Kana Hosoki, Naoki Goshima, Tetsuya Kawaguchi, Danny Hatters, Laura Trinkle-Mulcahy, Tetsuro Hirose, Charles S. Bond, and Archa H. Fox. Prion-like domains in RNA binding proteins are essential for building subnuclear paraspeckles. *Journal of Cell Biology*, 210(4):529–539, August 2015.

[68] Alex S. Holehouse, Garrett M. Ginell, Daniel Griffith, and Elvan Böke. Clustering of aromatic residues in prion-like domains can tune the formation, state, and organization of biomolecular condensates. *Biochemistry*, 60(47):3566–3581, 2021.

[69] Yuan Lin, David S.W. Protter, Michael K. Rosen, and Roy Parker. Formation and Maturation of Phase-Separated Liquid Droplets by RNA-Binding Proteins. *Molecular Cell*, 60(2):208–219, 10 2015.

[70] Veronica H Ryan, Gregory L Dignon, Gül H Zerze, Charlene V Chabata, Rute Silva, Alexander E Conicella, Joshua Amaya, Kathleen A Burke, Jeetain Mittal, and Nicolas L Fawzi. Mechanistic view of hnrnpa2 low-complexity domain structure, interactions, and phase separation altered by mutation and arginine methylation. *Molecular cell*, 69(3):465–479, 2018.

[71] Ignacio Sanchez-Burgos, Jorge R Espinosa, Jerelle A Joseph, and Rosana Collepardo-Guevara. Valency and binding affinity variations can regulate the multilayered organization of protein condensates with many components. *Biomolecules*, 11(2):278, 2021.

[72] Jeong-Mo Choi and Rohit V Pappu. The stickers and spacers framework for describing phase behavior of multivalent intrinsically disordered proteins. *Biophysical Journal*, 118(3):492a, 2020.

[73] Garrett M Ginell and Alex S Holehouse. An introduction to the stickers-and-spacers framework as applied to biomolecular condensates. In *Phase-Separated Biomolecular Condensates: Methods and Protocols*, pages 95–116. Springer, 2022.

[74] Suman Das, Yi Hsuan Lin, Robert M. Vernon, Julie D. Forman-Kay, and Hue Sun Chan. Comparative roles of charge, π, and hydrophobic interactions in sequence-dependent phase separation of intrinsically disordered proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 117(46):28795–28805, 11 2020.

[75] H Jane Dyson, Peter E Wright, and Harold A Scheraga. The role of hydrophobic interactions in initiation and propagation of protein folding. *Proceedings of the National Academy of Sciences*, 103(35):13057–13061, 2006.

[76] Charles D Andrew, Samita Bhattacharjee, Nicoleta Kokkoni, Jonathan D Hirst, Gareth R Jones, and Andrew J Doig. Stabilizing interactions between aromatic and basic side chains in α-helical peptides and proteins. tyrosine effects on helix circular dichroism. *Journal of the American Chemical Society*, 124(43):12706–12714, 2002.

[77] Wei Li and Hao Jiang. Nuclear protein condensates and their properties in regulation of gene expression. *Journal of Molecular Biology*, 434(1):167151, 2022. Protein Interactions in Liquid-Liquid Phase Separation.

[78] Andrés R Tejedor, Rosana Collepardo-Guevara, Jorge Ramirez, and Jorge R Espinosa. Time-dependent material properties of aging biomolecular condensates from different viscoelasticity measurements in molecular dynamics simulations. *The Journal of Physical Chemistry B*, 2023.

[79] Simon Alberti, Amy Gladfelter, and Tanja Mittag. Considerations and Challenges in Studying Liquid-Liquid Phase Separation and Biomolecular Condensates. *Cell*, 176(3):419–434, 1 2019.

[80] Soumik Ray and Alexander K Buell. Emerging experimental methods to study the thermodynamics of biomolecular condensate formation. *The Journal of Chemical Physics*, 160(9), 2024.

[81] Jorge R. Espinosa, Jerelle A. Joseph, Ignacio Sanchez-Burgos, Adiran Garaizar, Daan Frenkel, and Rosana Collepardo-Guevara. Liquid network connectivity regulates the stability and composition of biomolecular condensates with many components. *Proceedings of the National Academy of Sciences*, 117(24):13238–13247, 6 2020.

[82] Yuri Hong, Saeed Najafi, Thomas Casey, Joan-Emma Shea, Song-I Han, and Dong Soo Hwang. Hydrophobicity of arginine leads to reentrant liquid-liquid phase separation behaviors of arginine-rich proteins. *Nature Communications*, 13(1):7326, 2022.

[83] Jamie A Greig, Tu Anh Nguyen, Michelle Lee, Alex S Holehouse, Ammon E Posey, Rohit V Pappu, and Gregory Jedd. Arginine-enriched mixed-charge domains provide cohesion for nuclear speckle condensation. *Molecular cell*, 77(6):1237–1250, 2020.

[84] Amy R Strom, Alexander V Emelyanov, Mustafa Mir, Dmitry V Fyodorov, Xavier Darzacq, and Gary H Karpen. Phase separation drives heterochromatin domain formation. *Nature*, 547(7662):241–245, 2017.

[85] Dennis J. Mitchell, Lawrence Steinman, D T Kim, C G Fathman, and Jonathan B. Rothbard. Polyarginine enters cells more efficiently than other polycationic homopolymers. *Journal of Peptide Research*, 56(5):318–325, 2000.

[86] Rahul K. Das and Rohit V. Pappu. Conformations of intrinsically disordered proteins are influenced by linear sequence distributions of oppositely charged residues. *Proceedings of the National Academy of Sciences*, 110(33):13392–13397, 2013.

[87] Megan C. Cohan, Min Kyung Shinn, Jared M. Lalmansingh, and Rohit V. Pappu. Uncovering non-random binary patterns within sequences of intrinsically disordered proteins. *Journal of Molecular Biology*, 434(2):167373, 2022.

[88] Xiangze Zeng, Kiersten M. Ruff, and Rohit V. Pappu. Competing interactions give rise to two-state behavior and switch-like transitions in charge-rich intrinsically disordered proteins. *Proceedings of the National Academy of Sciences*, 119(19), May 2022.

[89] David De Sancho. Phase separation in amino acid mixtures is governed by composition. *Biophysical Journal*, 121(21):4119–4127, 2022.

[90] Jonas Ahlers, Ellen M. Adams, Verian Bader, Simone Pezzotti, Konstanze F. Winklhofer, Jörg Tatzelt, and Martina Havenith. The key role of solvent in condensation: Mapping water in liquid-liquid phase-separated fus. *Biophysical Journal*, 120(7):1266–1275, 2021.

[91] Akira Nomoto, Suguru Nishinami, and Kentaro Shiraki. Solubility parameters of amino acids on liquid–liquid phase separation and aggregation of proteins. *Frontiers in Cell and Developmental Biology*, 9:691052, 2021.

[92] Wenwei Zheng, Gregory L. Dignon, Matthew L. Brown, Young C. Kim, and Jeetain Mittal. Hydropathy patterning complements charge patterning to describe conformational preferences of disordered proteins. *Journal of Physical Chemistry Letters*, 11(9):3408–3415, 2020.

[93] Benjamin G Weiner, Andrew GT Pyo, Yigal Meir, and Ned S Wingreen. Motif-pattern dependence of biomolecular phase separation driven by specific interactions. *PLoS computational biology*, 17(12):e1009748, 2021.

[94] Jonathan Huihui and Kingshuk Ghosh. Intrachain interaction topology can identify functionally similar intrinsically disordered proteins. *Biophysical Journal*, 120(10):1860–1868, 2021.

[95] Kingshuk Ghosh, Jonathan Huihui, Michael Phillips, and Austin Haider. Rules of physical mathematics govern intrinsically disordered proteins. *Annual Review of Biophysics*, 51(1):355–376, May 2022.

[96] Ushnish Rana, Clifford P. Brangwynne, and Athanassios Z. Panagiotopoulos. Phase separation vs aggregation behavior for model disordered proteins. *The Journal of Chemical Physics*, 155(12):125101, September 2021.

[97] Erik W Martin, Alex S Holehouse, Christy R Grace, Alex Hughes, Rohit V Pappu, and Tanja Mittag. Sequence determinants of the conformational properties of an intrinsically disordered protein prior to and upon multisite phosphorylation. *Journal of the American Chemical Society*, 138(47):15323–15335, 2016.

[98] Alex S Holehouse, Garrett M Ginell, Daniel Griffith, and Elvan Boke. Clustering of aromatic residues in prion-like domains can tune the formation, state, and organization of biomolecular condensates: published as part of the biochemistry virtual special issue "protein condensates". *Biochemistry*, 60(47):3566–3581, 2021.

[99] Roshan Mammen Regy, Jacob Thompson, Young C. Kim, and Jeetain Mittal. Improved coarse-grained model for studying sequence dependent phase separation of disordered proteins. *Protein Science*, 30(7):1371–1379, May 2021.

[100] Gregory L. Dignon, Wenwei Zheng, Young C. Kim, Robert B. Best, and Jeetain Mittal. Sequence determinants of protein phase behavior from a coarse-grained model. *PLoS Computational Biology*, 14(1), 1 2018.

[101] Giulio Tesei, Thea K. Schulze, Ramon Crehuet, and Kresten Lindorff-Larsen. Accurate model of liquid–liquid phase behavior of intrinsically disordered proteins from optimization of single-chain properties. *Proceedings of the National Academy of Sciences*, 118(44), October 2021.

[102] Roshan Mammen Regy, Wenwei Zheng, and Jeetain Mittal. Using a sequence-specific coarse-grained model for studying protein liquid–liquid phase separation. pages 1–17, 2021.

[103] Andrew P. Latham and Bin Zhang. Maximum entropy optimized force field for intrinsically disordered proteins. *Journal of Chemical Theory and Computation*, 16(1):773–781, November 2019.

[104] Sohee Park, Yongjin Lee, Yongseok Jho, and Dong Soo Hwang. Molecular hydration tunes the cation–π interaction strength in aqueous solution. *Advanced Materials Interfaces*, 10(4):2201732, December 2022.

[105] Justin P. Gallivan and Dennis A. Dougherty. A computational study of cation-π interactions vs salt bridges in aqueous media: implications for protein engineering. *Journal of the American Chemical Society*, 122(5):870–874, January 2000.

[106] Tina Ukmar-Godec, Saskia Hutten, Matthew P. Grieshop, Nasrollah Rezaei-Ghaleh, Maria-Sol Cima-Omori, Jacek Biernat, Eckhard Mandelkow, Johannes Söding, Dorothee Dormann, and Markus Zweckstetter. Lysine/RNA-interactions drive and regulate biomolecular condensation. *Nature Communications*, 10(1), July 2019.

[107] A. P. Thompson, H. M. Aktulga, R. Berger, D. S. Bolintineanu, W. M. Brown, P. S. Crozier, P. J. in 't Veld, A. Kohlmeyer, S. G. Moore, T. D. Nguyen, R. Shan, M. J. Stevens, J. Tranchida, C. Trott, and S. J. Plimpton. LAMMPS - a flexible simulation tool for particle-based materials modeling at the atomic, meso, and continuum scales. *Comp. Phys. Comm.*, 271:108171, 2022.

[108] Niels Grønbech-Jensen and Oded Farago. A simple and effective verlet-type algorithm for simulating langevin dynamics. *Molecular Physics*, 111(8):983–991, 2013.

Full amino acid sequences

RBM14

```
WT(Q96PK6)   1 PMTAQAASYRAQPSVSLGAPYRGQLASPSSQSAAASSLGPYGGAQPSASALSSYGGQAAAASSLNSYGAQGSSLASYG
-20Y         1 .......G...........S....................Q...........G.....................S.
-10Y         1 .......G...........S....................Q.................................
-5Y          1 ...................S.....................................................
-30Y+30F     1 .......F...........F....................F...........F.............F.........F.
-20Y+20F     1 .......F...........F....................F...........F.....................
-10Y+10F     1 .......F................................F..................................
-5Y+5F       1 .......F.................................................................
-10T+10S     1 ..S......................................................................
-10T+10G     1 ..G......................................................................
-5T+5S       1 ..S......................................................................
-5T+5G       1 ..G......................................................................
-20Q+20N     1 ....N...............N...................N...................N.......
-10Q+20N     1 ....N...............N...................N...................N.......
-5Q+5N       1 ....N....................................................................
-4N+4Q       1 ..................................................Q......................
```

```
WT(Q96PK6)  79 NQPSSYGAQAASSYGVRAAASSYNTQGAASSLGSYGAQAASYGAQSAASSLAYGAQAASYNAQPSASYNAQSAPYAAQ
-20Y        79 .....G.......Q........Q...........G......S................G.......Q......S...
-10Y        79 ............Q...............S.....................G......................
-5Y         79 ............G...........................................................
-30Y+30F    79 ....F......F.........F.......F......F.......F.....F.......F......F...
-20Y+20F    79 ....F......F.................F......F.............F......F...........
-10Y+10F    79 ....F.......................F....................F...................
-5Y+5F      79 ............F.......................................................F...
-10T+10S    79 ...................S....................................................
-10T+10G    79 ...................G....................................................
-5T+5S      79 ......................................................................
-5T+5G      79 ......................................................................
-20Q+20N    79 .N.......................N...............N............N.............N
-10Q+20N    79 .......................N...............N........N................N
-5Q+5N      79 .N.....................................................N......
-4N+4Q      79 ....................Q...........................Q........
```

```
WT(Q96PK6) 157 QAASYSSQPAAYVAQPATAAAYASQPAAYAAQATTPMAGSYGAQPVVQTQLNSYGAQASMGLSGSYGAQSAAAATGSY
-20Y       157 ....Q................S......S.........Q..........S.................
-10Y       157 .....................S..........Q.................................
-5Y        157 ....................S...................Q..........................
-30Y+30F   157 ....F......F.......F......F........F.......F.......F..........F
-20Y+20F   157 ....F......F...............F........F.......F........F
-10Y+10F   157 ....F.......................F.......................F
-5Y+5F     157 .................................................F
-10T+10S   157 ..................S.............SS.........S...............S...
-10T+10G   157 ..................G.............GG.........G...............G...
-5T+5S     157 ..................S.............S..........................S...
-5T+5G     157 ..................G.............G..........................G...
-20Q+20N   157 N...........N........N...........N...N.......N
-10Q+20N   157 N.........................N.......N
-5Q+5N     157 ...........................N
-4N+4Q     157 ......................................................
```

```
WT(Q96PK6) 235 GAAAAYGAQPSATLAAPYRTQSSASLAASYAAQQHPQAAASYRGQPGNAYDGAGQPSAAYLSMSQGAVANANSTPPPY
-20Y       235 .....S............Q..........Q...........S......S..........Q..............S
-10Y       235 ..................S..........................S.................
-5Y        235 ..............................S.......................
-30Y+30F   235 .....F...........F..........F.......F.....F........F..............F
-20Y+20F   235 .................F..........F.......F.....F........F
-10Y+10F   235 .................F..........F.....F........
-5Y+5F     235 .................................F
-10T+10S   235 ...........S......S.............................S....
-10T+10G   235 ...........G......G.............................G....
-5T+5S     235 ...........S.........................................
-5T+5G     235 ...........G.........................................
-20Q+20N   235 .......N...............NN.........N.......N.......N
-10Q+20N   235 ......................N.............................
-5Q+5N     235 ...........................N.........................
-4N+4Q     235 ...........................................Q.................Q.....
```

TDP43

```
WT(Q13148)    1 R G G G A G L G N N Q G S N M G G G M N F G A F S I N P A M M A A A Q A A L Q S S W G M M G M L A S Q Q N Q S G P S G N N
-Y+F          1 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5F+5Y        1 . . . . . . . . . . . . . . . . . . . . . . . . Y . . Y . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5F           1 . . . . . . . . . . . . . . . . . . . . . . . . G . . S . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-20Q+20N      1 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-10Q+10N      1 . . . . . . . . . N . . . . . . . . . . . . . . . . . . . . . . . . . N . . . N . . . . . . . . . . . . . N N . N . . . . . . .
-5Q+5N        1 . . . . . . . . . N . . . . . . . . . . . . . . . . . . . . . . . . . . . N . . . . . . . . . . . . . . . N . . . . . . . . .
-15N+15Q      1 . . . . . . . . Q . . . Q . . . . Q . . . . . Q . . . . . . . . . . . . . . . . . . . . . . . . . . . . . Q . . . . . . Q Q
-10N+10Q      1 . . . . . . . . Q . . . Q . . . . Q . . . . . Q . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . Q Q
-5N+5Q        1 . . . . . . . . Q . . . . . . . . Q . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . Q .

WT(Q13148)   62 Q N Q G N M Q R E P N Q A F G S G N N S Y S G S N S G A A I G W G S A S N A G S G S G F N G G F G S S M D S K S S G W G M
-Y+F         62 . . . . . . . . . . . . . . . . . . . . . . . . . F . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5F+5Y       62 . . . . . . . . . . . . . . Y . . . . . . . . . . . . . . . . . . . . . . . . . . . Y . . . Y . . . . . . . . . . . . . .
-5F          62 . . . . . . . . . . . . . . G . . . . . . . . . . . . . . . . . . . . . . . . . . . S . . . G . . . . . . . . . . . . . .
-20Q+20N     62 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-10Q+10N     62 N . N . . . N . . . . N . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5Q+5N       62 N . . . . . N . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-15N+15Q     62 . Q . . Q . . . . . Q . . . . . . Q Q . . . . . Q . . . . . . . . . . Q . . . . . . . . Q . . . . . . . . . . . . . . . . .
-10N+10Q     62 . Q . . . . . . . . Q . . . . . Q Q . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5N+5Q       62 . . . . Q . . . . . . . . . . . . Q . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
```

TIA1

```
WT(P31483)    1 M I N P V Q Q Q N Q I G Y P Q P Y G Q W G Q W Y G N A Q Q I G Q Y M P N G W Q V P A Y G M Y G Q A W N Q Q G F N Q T Q
-12Y          1 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-6Y           1 . . . . . . . . . . . Q . . . G . . . . . S . . . . . . . . . . . . . . . . . . . . . Q . . . . . . . . . . . . . . .
-9Y+9F        1 . . . . . . . . . . . . . F . . . F . . . . . F . . . . . . . . . . F . . . . . . . . . F . . F . . . . . . . . . . .
-4Y+4F        1 . . . . . . . . . . . . . F . . . . . . . . . F . . . . . . . . . . F . . . . . . . . . . . . . . . . . . . . . . . .
-9Y+9W        1 . . . . . . . . . . . . . W . . . W . . . . . W . . . . . . . W . . . . . . . . . . . W . . W . . . . . . . . . . . .
-4Y+4W        1 . . . . . . . . . . . . . W . . . . . . . . . W . . . . . . . . . . W . . . . . . . . . . . . . . . . . . . . . . . .
-5W+5F        1 . . . . . . . . . . . . . . . . . . F . F . . . . . . . . . . . . . F . . . . . . . . . . . . F . . . . . . . . . . .
-5W+5Y        1 . . . . . . . . . . . . . . . . . . Y . Y . . . . . . . . . . . . . Y . . . . . . . . . . . . Y . . . . . . . . . . .
-20Q+20N      1 . . . . . N N N . N . . . . N . . . N . . N . . . . . N N . . N . . . . . . . N . . . . . . . . N . . . N N . . . N . N
-10Q+10N      1 . . . . . N N N . . . . . . . . . . . . . . . . . . . . . N N . . . . . . . . . . . . . . . . . . . . . N . . . . . . .
-9N+9Q        1 . . Q . . . . Q . . . . . . . . . . . . . . . . . Q . . . . . . . . . . Q . . . . . . . . . . Q . . . . Q . . .

WT(P31483)   60 S S A P W M G P N Y G V Q P P Q G Q N G S M L P N Q P S G Y R V A G Y E T N
-12Y         60 . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-6Y          60 . . . . . . . . . G . . . . . . . . . . . . . . . . . . . . . . . . . . .
-9Y+9F       60 . . . . . . . . . F . . . . . . . . . . . . . . . . . . F . . . F . . .
-4Y+4F       60 . . . . . . . . . F . . . . . . . . . . . . . . . . . . . . . . . . . . .
-9Y+9W       60 . . . . . . . . W . . . . . . . . . . . . . . . . . W . . . . W . . .
-4Y+4W       60 . . . . . . . . W . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5W+5F       60 . . . . F . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-5W+5Y       60 . . . Y . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . . .
-20Q+20N     60 . . . . . . . . N . . N . N . . . . . . N . . . . . . . . . . . . . . .
-10Q+10N     60 . . . . . . . . N . . . . N . . . . . N . . . . . . . . . . . . . . .
-9N+9Q       60 . . . . . . . . Q . . . . . . . . . . Q . . . . . Q . . . . . . . . . -
```

1033 **FUS**

```
(P35637)    1 MASNDYTQQATQSYGAYPTQPGQGYSQQSSQPYGQQSYSGYSQSTDTSGYGQSSYSSYGQS
-28Y+28F    1 .....F.......F..F......F.......F....F.F.......F...F..F...
-14Y+14F    1 .....F.......F.......F.......F.......F.......F.......F...
-7Y+7F      1 .....F.......F.......F.......F.......F.....
-14Y        1 ............Q.........G.........S.........Q......G...
-7Y         1 ............Q.........G.........G...
-10T+10S    1 .....S...S......S.........S.S...
-10T+10G    1 .....G...G......G.........G.G...
-5T+5S      1 .....S......S.........S...
-5T+5G      1 .....G......G.........G...
-10G+10T    1 ............T.........T...
-10S+10T    1 ..T.......T.......T...
-20Q+20N    1 .......N...N......N....N......N......N.........N.
-10Q+10N    1 .......N......N......N......N.
-5Q+5N      1 .......N......N.
-4N+4Q      1 .

(P35637)   62 QNTGYGTQSTPQGYGSTGGYGSSQSSQSSYGQQSSYPGYGQQPAPSSTSGSYGSSSQSSSY
-28Y+28F   62 ....F.......F....F.......F....F..F.......F.......F
-14Y+14F   62 ....F.......F.......F.......F.....
-7Y+7F     62 ....F.......F.......F.....
-14Y       62 ..........S.........Q.......G.........S
-7Y        62 ..........S.........Q...
-10T+10S   62 ..S...S..S......S.........S...
-10T+10G   62 ..G...G..G......G.........G...
-5T+5S     62 .....S......S...
-5T+5G     62 .....G......G...
-10G+10T   62 .........T.........T...
-10S+10T   62 .........T.......T.......T..T...
-20Q+20N   62 .......N.......N....N......N......N....
-10Q+10N   62 .......N......N......N...
-5Q+5N     62 .......N...
-4N+4Q     62 .Q...

(P35637)  123 GQPQSGSYSQQPSYGGQQQSYGQQQSYNPPQGYGQQNQYNSSSGGGGGGGGGGGGNYGQDQSS
-28Y+28F  123 .......F....F......F.....F.....F.....F.......F.....
-14Y+14F  123 .......F.......F.......F.......F.....
-7Y+7F    123 ................F.............F...
-14Y      123 ...........Q.........S.........G...
-7Y       123 ...........S.........G...
-10T+10S  123 .
-10T+10G  123 .
-5T+5S    123 .
-5T+5G    123 .
-10G+10T  123 .........T.........T...T....T...
-10S+10T  123 ..........T.......T...
-20Q+20N  123 ...N......N......N....N.N......N..N.........N..
-10Q+10N  123 ...N......N......N......N...
-5Q+5N    123 .........N.........N..
-4N+4Q    123 .........Q..Q.........Q.....

(P35637)  184 MSSGGGSGGGYGNQDQSGGGGSGGYGQQDRGGRGRGGSGGGGGGGGGGGGYNRSSG
-28Y+28F  184 .......F.......F.........F...
-14Y+14F  184 .......F.....
-7Y+7F    184 .
-14Y      184 .........G.........G...
-7Y       184 .........G...
-10T+10S  184 .
-10T+10G  184 .
-5T+5S    184 .
-5T+5G    184 .
-10G+10T  184 .......T.......T.......T...
-10S+10T  184 ..T.....
-20Q+20N  184 .
-10Q+10N  184 .
-5Q+5N    184 .
-4N+4Q    184 .........Q...
```

1034 **EWSR1**

Dataset representation

```
WT(Q01844)    1 MASTDYSTYSQAAAQQGYSAYTAQPTQGYAQTTQAYGQQSYGTYGQPTDVSYTQAQTTATYGQTAYATSYGQP
-20Y          1 ....T..Q........S..G.......................T.........Q............S......
-10Y          1 ....T..Q.......S..G.......................................T.........
-5Y           1 ....T..............................................Q...
-30Y+30F      1 ....F..F.......F.......F.......F...F.........F.......F...F...F.
-20Y+20F      1 ....F.........F.......F.......F.......F...F.
-10Y+10F      1 ....F.........F.......F.......F...F.
-5Y+5F        1 ....F.
-30T+30S      1 ..S...........S...S...S.......S...S...S....S.S..S.
-30T+30G      1 ..G...........G...G...G.......G...G...G.G..G.
-20T+20S      1 ..S...........S...S.......S...S...S..S.
-10T+10S      1 ....S..........................S...........S....S.
-20T+20G      1 ..G...................G.......G.......G...G...G.
-10T+10G      1 ..G...................G.......G...G.
-30S+30T      1 ..T...T..T.................T...T...T.............T.
-20G+20T      1 ...............T.......T.......T..T...T.............T.
-10G+10T      1 ...............T.......T.......T.............T.
-20S+20T      1 ..T...T...T...............T.............T.
-10S+10T      1 ..T..........T.............................T.
-20Q+20N      1 .......N...N.......N.......N.......N.......N.........N.
-10Q+10N      1 .......N.......N.......N.
-5Q+5N        1 .......N.......N.
-4N+4Q        1

WT(Q01844)   74 PTGYTTPTAPQAYSQPVQGYGTGAYDTTTATVTTTQASYAAQSAYGTQPAYPAYGQQPAATAPTRPQDGNKPT
-20Y         74 ...T.......G........Q...................T..S.
-10Y         74 ...Q...........S...............G.
-5Y          74 ...............S...............G.
-30Y+30F     74 ...........F....F...F.........F...F.......F.
-20Y+20F     74 ..F...........F...F.........F...F.......F.
-10Y+10F     74 ..F...........F...........F...F.
-5Y+5F       74 ...............F.
-30T+30S     74 .S..SS.S...............SSS...SSS...................S...S
-30T+30G     74 ....GG.G..............GG..G.GG.......G.......G.......G
-20T+20S     74 .S...S............S...S..S..S.......S.......S....
-10T+10S     74 ..S...S............S...S...S.......S
-20T+20G     74 .G...G..............G...G..G..G.......G.......G.
-10T+10G     74 .....G.............G...G.......G.
-30S+30T     74 ...................T...T.
-20G+20T     74 ..T............T..T.......T.......T......T.
-10G+10T     74 ...............T...T.......T.......T.
-20S+20T     74 ...................T.
-10S+10T     74
-20Q+20N     74 .............N.........N...........N.......N.
-10Q+10N     74 .............N...............N.
-5Q+5N       74 ...............................N.
-4N+4Q       74 ...........................................Q...

WT(Q01844)  147 ETSQPQSSTGGYNQPSLGYGQSNYSYPQVPGSYPMQPVTAPPSYPPTSYSSTQPTSYDQSSYSQQNTYGQPSS
-20Y        147 ...................Q.G.....T...............S.........Q.....
-10Y        147 ........T...................S.
-5Y         147 ........T.
-30Y+30F    147 ........F....F...F.......F.......F...F.......F...F...F.
-20Y+20F    147 ........F...F.......F.......F...F.
-10Y+10F    147 ........F.......F.......F.
-5Y+5F      147 ........F.
-30T+30S    147 .S.......................S...S..S..S.......S.
-30T+30G    147 .G.......G...................G...G..G..G.......G.
-20T+20S    147 .S.......................S...S.......S.
-10T+10S    147 ...........................S...S.
-20T+20G    147 .G...................G...G...G.
-10T+10G    147 .....................G...G.
-30S+30T    147 .T....T.......T...T...T.........T...T..T...T...TT.T........TT
-20G+20T    147 .......T....T.T.......T...............T.
-10G+10T    147 .......T...........T.
-20S+20T    147 .T....T.......T.......T.......T..T.......T..T.......T
-10S+10T    147 ..T...........T...................T...........T.
-20Q+20N    147 .....N.........N.............N.......N.
-10Q+10N    147 .........N.....................N.
-5Q+5N      147 ...................N.
-4N+4Q      147 .........Q.......Q.................................Q......

WT(Q01844)  220 YGQQSSYGQQSSYGQQPPTSYPPQTGSYSQAPSQYSQQSSSYGQQSSFRQDHPSSMGVYGQ
-20Y        220 T...............G..............S.
-10Y        220
-5Y         220
-30Y+30F    220 F....F....F.......F.....F.
-20Y+20F    220 ....F...............F.......F.
-10Y+10F    220 F.
-5Y+5F      220
-30T+30S    220 .............S....S.
-30T+30G    220 .............G....G.
-20T+20S    220 .............S.
-10T+10S    220 ..........S.
-20T+20G    220 ..................G.
-10T+10G    220
-30S+30T    220 .....T....TT.........T....T..T..T...T.
-20G+20T    220 .T....T.............T.............T...........T.T.
-10G+10T    220 .....T.......................T.
-20S+20T    220 .....T....T.........T...T...T.T.
-10S+10T    220 .........T.......T...T.
-20Q+20N    220 ..N.....N......N..........N.....N.
-10Q+10N    220 ..N...........N..............N.
-5Q+5N      220 ...............N.
-4N+4Q      220
```

Appendix 0—figure 8. Representation of the entire computational data set, divided on mutations on 'sticker' or 'spacer' residues. Orange data points, on the left, represent variants where charged or aromatic residues were mutated (i.e., stickers), while cyan data points, on the right, represent all other types of mutations studied (i.e., spacers)

 Appendix 1

 ## Convergence tests

 Molecular Dynamic simulations are inherently sensitive to both the length of simulated time
 and system scaling factor such as the number of particles present in the system and the size
 of the simulation box. In this study, we systematically tested all these aspects to ensure
 proper convergence of the results.

 The system was placed in the middle of the simulation box, and the length of X and Y
 was the same as the length of the condensate, while the box was elongated in Z, to about
 ten times the size of X, Y(see 1). Importantly, the cross-section is large enough to avoid
 protein self-interactions through the periodic boundary conditions (e.g., cross-section sizes
 are at least twice the protein radius of gyration in condensate bulk conditions).

 We tested for time convergence by plotting the obtained density of the protein wild types
 after increasing simulation times, from 50 ns to 350 ns. On simulated timesteps above 100
 ns, the density of the condensate exhibited no significant differences. We thus concluded
 that, to optimize speed-up, 200 ns provided sufficient sampling for each point of the binodal.

 Finally, we decided to keep the number of particles at a constant value of 64, which we
 found to be a good balance between minimizing finite-size effects and optimizing simulation
 speed (see Figure 2). We tested this by varying the corresponding number of protein chains
 of the FUS WT as a model protein from 27 to 216 and found that in these systems there was
 not a significant difference in the simulated density. Additionally, with 64 protein chains, we
 could ensure a bulk large enough to sustain the condition that the cross-section of the box
 should be higher than twice the radii of gyration of the protein chain.

Appendix 1—figure 1. Convergence tests for the WT sequence of the PLD of FUS. The density profiles of different and independent simulations are plotted across the perpendicular axis to the condensate interfaces. Each simulation was performed for different timescales (as specified in the legend) to check for proper convergence.

Appendix 1—figure 2. Convergence tests for the WT sequence of the PLD of FUS. The density profiles of different and independent simulations are plotted across the perpendicular axis to the condensate interfaces. Each simulation was performed for different system sizes (aka, number of chains and corresponding box size, as specified in the legend) to check for proper convergence. To the right, a representation of the systems of increasing number of chains we employed to check for convergence. In each case, the overall density of the box is kept constant, while the cross-section and lenght is varied in order to mantain the density constraint. For 27 chains, the box dimensions are 109x109x765 Å; for 64 chains 145x145x1021 Å; for 125 chains 182x182x1276 Å and for the final case of 216 218x218x1531 Å.

 Appendix 2

 ## Normalised and comparative contact maps

Further study of the differences between variants at the chemical level was carried out by computing the inter-molecular contacts per residue inside a phase-separated droplet. For each variant, the contact matrix was computed at a temperature around ten percent below the T_c. The cutoff to define a pairwise distance as a contact was dynamically calculated for each pair, as 1.2 times the average Van der Waals radius of both pair residues. Given residues i and j belong to different molecules, the contact maps were computed as shown below:

$$c_{i,j} = \sum_{i,j} \chi_{ij} \tag{22}$$

$$\chi_{ij} = \begin{cases} 1 & \text{if } \left| r_i - r_j \right| \leq r_{ij}^{\text{cutoff}} \\ 0 & \text{otherwise} \end{cases} \tag{23}$$

To get a more accurate picture of the main interactions driving phase separation for each variant, we normalised the contact maps to account for the occurrence of each residue in the sequence. This allowed us to account for the disparities in residue distributions in the sequences, and extract the relevant residue–residue interactions. For each residue–residue pair, the normalisation was performed as follows:

$$C = \begin{vmatrix} \alpha_{M,M} & \alpha_{M,G} & \cdots & \alpha_{M,I} \\ \alpha_{G,M} & \alpha_{G,G} & \cdots & \alpha_{G,I} \\ \cdots & \cdots & \cdots & \cdots \\ \alpha_{I,M} & \alpha_{I,G} & \cdots & \alpha_{I,I} \end{vmatrix} \tag{24}$$

$$\alpha_{i,j} = \frac{c_{i,j}}{n_i \cdot n_j} \tag{25}$$

where n_i is the number of i residues in the IDP sequence. The resulting contact matrix is min–max normalised:

$$\beta_{i,j} = \frac{\alpha_{i,j} - \min C}{\max C - \min C} \tag{26}$$

$$C_{\text{norm}} = \begin{vmatrix} \beta_{M,M} & \beta_{M,G} & \cdots & \beta_{M,I} \\ \beta_{G,M} & \beta_{G,G} & \cdots & \beta_{G,I} \\ \cdots & \cdots & \cdots & \cdots \\ \beta_{I,M} & \beta_{I,G} & \cdots & \beta_{I,I} \end{vmatrix} \tag{27}$$

The comparative contact maps (Fig. 1 and 2), on the other hand, are computed as the difference in contacts of a protein variant respective to its wild type, as:

$$C_{\text{comparative}} = C_{\text{variant}} - C_{\text{WT}} \tag{28}$$

which is finally mean-normalised.

$$\gamma_{i,j} = \frac{\alpha_{i,j} - \overline{C}_{\text{comparative}}}{\max C_{\text{comparative}} - \min C_{\text{comparative}}} \tag{29}$$

$$C_{\text{comparative,normed}} = \begin{vmatrix} \gamma_{M,M} & \gamma_{M,G} & \cdots & \gamma_{M,I} \\ \gamma_{G,M} & \gamma_{G,G} & \cdots & {}_v2Prions_v2gamma_{G,I} \\ \cdots & \cdots & \cdots & \cdots \\ \gamma_{I,M} & \gamma_{I,G} & \cdots & \gamma_{I,I} \end{vmatrix} \tag{30}$$

Appendix 2—figure 1. Normalised contact maps of different *R to K* variants of PLDs: A) TDP B) FUS, C) hnRNPA1

Appendix 2—figure 2. Normalised contact maps of different aromatic variants of PLDs: A) FUS B) hnRNPA1.