1 **Main Manuscript for**
2
3 # The Rapidly Evolving X-linked *miR-506* Family
4 # Finetunes Spermatogenesis to Enhance Sperm
5 # Competition
6

7 Zhuqing Wang[1,2], Yue Wang[1,2], Tong Zhou[1], Sheng Chen[1,2], Dayton Morris[2],
8 Rubens Daniel Miserani Magalhães[2], Musheng Li[1], Shawn Wang[1], Hetan Wang[1,2],
9 Yeming Xie[1], Hayden McSwiggin[1,2], Daniel Oliver[1], Shuiqiao Yuan[1], Huili Zheng[1,2],
10 Jaaved Mohammed[3], Eric C. Lai[3], John R. McCarrey[4], Wei Yan[1,2,5]*
11
12
[1]Department of Physiology and Cell Biology, University of Nevada, Reno School of
Medicine, Reno, NV 89557, USA.
[2]The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center,
Torrance, CA 90502, USA.
[3]Department of Developmental Biology, Memorial Sloan-Kettering Institute, 1275
York Ave, Box 252, New York, NY 10065, USA.
[4]Department of Biology, University of Texas at San Antonio, San Antonio, TX, USA.
[5]Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles,
CA 90095, USA.
*Corresponding author: Wei Yan   **Email:** wei.yan@lundquist.org

23
24
25

26 **Author Contributions:** Z.W. and W.Y. designed the research. Z.W., Y. W., S. C.,
27 D.M., S. W., H. W., H.M., J. R. M., and H.Z., performed bench experiments. Z.W.,
28 T.Z., R.D.M.M., M.L., Y.X., J.M., and E. C. L. performed bioinformatics analysis.
29 Z.W., and W.Y. wrote the manuscript.

31 **Competing Interest Statement:** The authors declare no conflict of interest.
32
33 **Classification:** Biological Sciences; Genetics; Physiology.
34
Keywords: *reproductive fitness; sperm competition; evolution; selective pressure;*
microRNAs; spermatogenesis; male fertility

This PDF file includes:
Main Text
Figures 1 to 6

Abstract
Despite rapid evolution across eutherian mammals, the X-linked *miR-506* family
miRNAs are located in a region flanked by two highly conserved protein-coding
genes (*Slitrk2* and *Fmr1*) on the X chromosome. Intriguingly, these miRNAs are
predominantly expressed in the testis, suggesting a potential role in
spermatogenesis and male fertility. Here, we report that the X-linked *miR-506*
family miRNAs were derived from the MER91C DNA transposons and their
sequences diverged *via* LINE1-mediated retrotransposition during evolution.
Selective inactivation of individual miRNAs or clusters caused no discernable
defects, but simultaneous ablation of five clusters containing nineteen members of
the *miR-506* family led to reduced male fertility in mice. Despite normal sperm
counts, motility and morphology, the KO sperm were less competitive than wild-
type sperm when subjected to a polyandrous mating scheme. Transcriptomic and
bioinformatic analyses revealed that these X-linked *miR-506* family miRNAs, in
addition to targeting a set of conserved genes, have gained more targets that are
critical for spermatogenesis and embryonic development during evolution. Our
data suggest that the *miR-506* family miRNAs function to enhance sperm
competitiveness and reproductive fitness of the male by finetuning gene
expression during spermatogenesis.

Significance Statement
The X-linked *miR-506* family has rapidly evolved in mammals, but their
physiological significance remains elusive. Given their abundant and preferential
expression in the testis and sperm, these X-linked miRNAs likely play a functional
role in spermatogenesis and/or early embryonic development. However, the
deletion of either individual miRNA genes or all of the five miRNA clusters
encoding 38 mature miRNAs did not cause major fertility defects in mice. When
these mutant males were subjected to conditions resembling polyandrous mating,
the mutant sperm were much less competitive than the wild-type sperm, rendering
the mutant males "functionally infertile". Our data suggest that the *miR-506* family
of miRNAs regulates sperm competition and the reproductive fitness of the male.
72
73

Main Text

Introduction

Spermatogenesis is highly conserved among all vertebrates. Although it generally consists of three phases (mitotic, meiotic, and haploid), many characteristics appear to be species-specific, *e.g.,* the duration of each of the three phases, the seminiferous epithelial organization, and the shape and length of spermatozoa, likely reflecting the adaptive changes during evolution (1-3). Several cellular events are unique to the male germ cells, *e.g.,* meiosis, genetic recombination, and haploid male germ cell differentiation (also called spermiogenesis) (4). Unique cellular processes are often accompanied by a more complex yet unique transcriptome, which may explain why the testis expresses more genes than any other organs in the body, with the possible exception of the brain (5). Regulation of gene expression during spermatogenesis occurs at both transcriptional and post-transcriptional levels (6). As a post-transcriptional regulator, miRNAs are abundantly expressed in the testis and are required for spermatogenesis (7-11). miRNAs typically function at post-transcriptional levels by binding the complementary sequences in the untranslated regions (UTRs) of mRNAs – particularly in the 3'UTRs through the "seed sequence" (2^{nd}-7^{th} nucleotides)(12). Numerous miRNAs are subject to rapid evolution, probably in response to the accelerated rate of divergence of UTRs compared to the exonic sequences (13). Divergence of genomic sequences can be mediated by transposable elements (TEs), which are known as building blocks of the genome and mostly map to UTRs and intronic regions of protein-encoding genes (14). Each miRNA can bind numerous target mRNAs, and one mRNA can be targeted by multiple different miRNAs. This "one-to-multiple" relationship between miRNAs and mRNAs amplifies their potential to coordinate gene expression in the cell (12). Moreover, miRNA genes often exist in clusters, which are transcribed as a unit followed by nuclear and cytoplasmic cleavage events to generate individual miRNAs (12).

Multiple clusters of miRNA genes containing the same seed sequences are categorized as a miRNA family, and miRNAs within the same family likely evolved from a common ancestor sequence (15). Of great interest, many of the testis-enriched miRNA clusters map to the X chromosome (16). Sex-linked genes are generally subject to the male germline-specific phenomenon called meiotic sex chromosome inactivation (MSCI), which silences transcription during most, if not all, of meiosis (16). Indeed, prior to 2009, there were no confirmed reports of any sex-linked genes escaping the repressive effects of MSCI. Surprisingly, however, we found that many X-linked miRNA genes do escape MSCI, suggesting that they may contribute to particularly important functions during spermatogenesis (16). This notion has since been tested by the generation of knockouts (KOs) of individual miRNA genes or individual clusters of miRNA genes normally expressed during spermatogenesis. However, these KOs resulted in minimal, if any, phenotypic effects and did not appear to impede normal spermatogenesis or male fertility (15). This left the field facing multiple unanswered questions, including 1) why do so many X-linked miRNAs express uniquely or preferentially during

spermatogenesis and escape MSCI, 2) what's their origin, and 3) how and why did
they evolve rapidly?
To address these questions and better understand the functional role played
by these X-linked miRNAs, we investigated the evolutionary history of this unique
miRNA family and also generated KOs of individual, paired, triple, quadruple, or
quintuple sets of miRNA clusters within this family and tested the effects on male
fertility, initially by standard monandrous mating assays. Consistent with previous
efforts to inactivate miRNA genes in *C. elegans* and mice (15, 17-19), KOs of
either individual members or individual clusters of the *miR-506* family induced no
discernable phenotypes and did not impact male fertility. This may reflect the level
of functional redundancy inherent within the members and clusters of this miRNA
family. It was only when four or more clusters of the *miR-506* family were ablated
that relevant phenotype became detectable, which was manifested in the form of
reduced litter size despite normal sperm counts, motility, and morphology.
Interestingly, the most common male fertility testing for lab rodents is based on a
monandrous mating scheme, *i.e.,* one fertility-proven female mated with one male.
However, there are additional aspects of male fertility in many mammalian species,
particularly those that are normally litter-bearing. In the wild, litter-bearing females
often mate with multiple different males such that a single litter may include pups
sired by more than one male (20, 21). This polyandrous mating introduces the
potential for additional aspects of male reproductive fitness to accrue, one of which
involves sperm competition (20, 21). Sperm competition can occur when sperm
from more than one male are present in the female reproductive tract
simultaneously, such that they then compete to fertilize each oocyte (22). Sperm
competition has been now recognized as a significant evolutionary force directly
impacting male reproductive success (22). Using experiments that mimic
polyandrous mating, we found that the quinKO male mice indeed displayed
compromised sperm competition. Hence, the X-linked *miR-506* family miRNAs
appear to function to finetune spermatogenesis to enhance sperm competition and,
consequently, male reproductive fitness.
150
151 **Results**
152
X-linked *miR-506* family miRNAs flanked by two highly conserved protein-
coding genes *Slitrk2* and *Fmr1* rapidly evolved across species. X-linked
genes are generally more divergent between species than autosomal ones, a
phenomenon known as the "faster-X effect" (23). However, despite a high degree
of conservation of two protein-coding genes, *Slitrk2* and *Fmr1*, on the X
chromosome across species, the miRNA genes located between these two loci are
divergent among clades across the eutherian mammals (15, 24). Through tracing
the evolution of this genomic region, we found that *Slitrk2* and *Fmr1* were
syntenically linked to autosomes in zebrafish and birds (Fig. 1A), but had migrated
onto the X chromosome in most mammals, with divergence and multiplication of
numerous miRNA genes that belong to the *miR-506* family in between (Fig. 1A and
Table S1). By mapping these miRNAs of various species using the UCSC genome
browser (25), we found that all members of the *miR-506* family are located in a

region flanked by *Slitrk2* and *Fmr1* (Fig. 1A). Consistent with previous reports (15,
24), *Slitrk2* and *Fmr1* are usually on the positive strand, whereas the *miR-506*
family miRNAs are in the reverse orientation (Fig. 1A). Based on the location of
these miRNAs, we named the miRNAs proximal to *Slitrk2* (*miR-892c ~ miR-891a*
in humans) and *Fmr1* (*miR-513c ~ miR-514a3* in humans) SmiRs (*Slitrk2*-proximal
miRNAs) and FmiRs (*Fmr1*-proximal miRNAs), respectively.
To evaluate the sequence conservation of these miRNAs across species,
we adopted the Multiz Alignment and Conservation pipeline, which utilizes
PhastCons and PhyloP algorithms (25), to search miRNA datasets from 100
different species using the human genome as a reference (Fig. 1B and Fig. S1).
The mean values of phyloP and phastCons of the *Fmr1* and *Slitrk2* coding
sequences (CDS) were ~4.9 and ~0.97, respectively (Fig. 1C and D), indicating
that these regions are highly conserved. In contrast, the mean values of phyloP
and phastCons of SmiRs were ~1.0 and ~0.002, respectively, and those of FmiRs
are ~0.03 and ~0.11, respectively (Fig. 1 C and D). The phyloP and phastCons
values of the CDS, FmiRs, and SmiRs are significantly different from each other
(adjusted p-value < 0.05, Kruskal-Wallis test) (Fig. 1 C and D), indicating that
FmiRs and SmiRs are highly divergent, and SmiRs are more divergent than FmiRs.
We then assessed the genomic sequence similarity among various species
using D-GENIES-based dot plot analyses (26). Although the *Slitrk2-Fmr1* genomic
regions were highly variable among different species, the sequences within some
clades shared a high degree of similarities, e.g., primates (rhesus monkeys,
chimpanzees, and humans), *cetartiodactyla* (sheep and cows), *rodentia* (e.g., mice
and rats) and *carnivora* (e.g., dogs and cats) (Fig. S2A). Although the *miR-506*
family miRNAs were highly divergent, some orthologs displayed a higher degree of
sequence conservation (Fig. S2 B and C), e.g., the SmiRs within primates were
similar (Fig. S2B); *miR-891a* and *miR-891b* in rhesus monkeys were similar to
miR-891b and *miR-891a* in humans and chimpanzees, respectively, and *miR-892b*
in chimpanzees was homologous to *miR-892c* in humans and rhesus monkeys in
terms of sequence and location (Figs. S2B and S3). The FmiRs, including *miR-201*
(assigned as *Mir-506-P1* (paralogue 1) in MirGeneDB (27)), *miR-547* (*Mir-506-P2*),
and *miR-509* (*Mir-506-P7*) in mice and rats, are orthologues of *miR-506*, *miR-507*
and *miR-509* in humans, respectively (Fig. S2C, and Table S1). Of interest,
although the *miR-506* family miRNAs are highly divergent, the seed sequences of
some miRNAs, such as *Mir-506-P6* and *Mir-506-P7*, remain conserved (Fig. S2C),
and these miRNAs represent the dominant mature miRNAs (28). It is noteworthy
that the majority of the substitutions among the *miR-506* family are U→C and
A→G (Fig. S2 B and C). Furthermore, we analyzed the conservation of the *miR-*
506 family in modern humans using data from the 1000 Genomes Project (1kGP)
(Fig. 1 E and F) (29). We compared SmiRs, FmiRs, and all miRNAs with
pachytene piRNAs, which are known to be highly divergent in modern humans but
barely exert any biological functions (30). Of interest, the derived allele frequency
(DAF) and mean nucleotide diversity (MND) of FmiRs and all miRNAs were
significantly smaller than that of the pachytene piRNAs, whereas SmiRs were
significantly smaller than all miRNAs (Fig. 1 E and F) (adjusted p-value < 0.05,
Kruskal-Wallis test), suggesting that the *miR-506* family miRNAs are more

conserved than pachytene piRNAs in modern humans. Taken together, these data
indicate that the X-linked *miR-506* family, although rapidly evolving as a whole,
contains both divergent and conserved miRNAs, suggesting both conserved and
novel functions across species.
216
X-linked *miR-506* family miRNAs are derived from MER91C DNA transposons
and expanded *via* LINE1 retrotransposition. To visualize the family history of
these miRNAs, we built a phylogram for the *miR-506* family (Fig. S4). The
phylogram suggests that these miRNAs shared a common ancestor and that the
FmiRs emerged earlier than the SmiRs, which is also supported by the fact that
some FmiRs exist in green sea turtles (Figs. S1 and S4). These data suggest that
the *miR-506* family miRNAs arose much earlier than previously thought (25). The
two subfamilies, FmiRs and SmiRs, despite their common ancestors, may have
evolved at different paces and thus, might be functionally divergent.
Studies have shown that transposable elements (TEs) drive evolution
through transpositions (31). CRISPR-Cas9/Cas12a genome editing can induce
irreversible small indels at the cutting sites (also called "scars")(32), which have
been used for lineage tracing (33). Inspired by this strategy, we attempted to trace
the evolution of the *miR-506* family miRNAs by searching the transposon database
for the transpositional "scars" (partial TE sequences) after transposition. To search
the potential TE sources of the *miR-506* family miRNAs, we downloaded all
transposons in the human, horse, dog, and guinea pig genomes and aligned them
to their corresponding *miR-506* family miRNAs using BLAST (Basic Local
Alignment Search Tool) (34). The MER91C DNA transposon (~100-150 million
years) (35) was the only transposon that aligned to FmiRs of the *miR-506* family (>
94% identical matches) in all four species (Table S2). The remaining miRNAs of
the *miR-506* family, mainly SmiRs, are flanked by LINEs (long interspersed
elements) or SINEs (short interspersed element, including Alu) (Fig. S3)
retrotransposons (Fig. S3 and Table S2). LINE1 is the most abundant and the only
active TE in the human genome (36). Although SINEs cannot mobilize by
themselves, they can hijack the LINE1 mechanism to achieve transpositions (36).
Given that the FmiRs (e.g., *miRs-506~509*) emerged much earlier than the
SmiRs (fig. S1 and fig. S4) and that *miR-513* (belonging to FmiRs) and SmiRs
(including *miR-891a* and *miR-891b*) share a common ancestor (Fig. S4), we
reasoned that the X-linked *miR-506* family might be derived from the *MER91C*
DNA transposon and further expanded through LINE1-mediated retrotransposition.
To test this hypothesis, we first aligned the X-linked *miR-506* family miRNAs from
several species to a human MER91C DNA transposon. Indeed, numerous FmiRs
of almost all species analyzed aligned to the MER91C DNA transposon despite
few mismatches (Fig. 2A), phylogenetic tree further confirmed that the MER91C is
the sister group of the *miR-506* family miRNAs (Fig. 2B and Fig. S5), suggesting
that the MER91C DNA transposon is the likely source of the older *miR-506* family
miRNAs. Further supporting this notion, the MER91C DNA transposons could form
hairpin structures, which is a prerequisite for miRNA biogenesis (Fig. 2C and Fig.
S6A). Moreover, analyses of the testis small RNA datasets from humans,
marmosets, dogs, and horses revealed the peaks corresponding to these miRNAs

(Fig. 2D and Fig. S6A). Finally, by overexpressing several MER91C DNA regions
randomly selected from humans, dogs, and horses in HEK293T cells, we found
that these DNA regions were indeed capable of producing miRNAs (Fig. 2 E and F,
and Fig. S6 B and C). Co-expression of MER91C DNA regions and AGO2
significantly increased the abundance of human MER91C miRNAs, as compared
with overexpression of MER91C DNA regions alone (Fig. 2 E and F), suggesting
that these miRNAs could be loaded onto and protected by AGO2. Together, these
data indicate that the MER91C DNA transposon serves as a source for the *miR-*
506 family miRNAs. As to the rest of the *miR-506* family miRNAs, we first
compared the proportion of TEs in the miRNA-flanking regions (length of different
classes of TEs/length of the genomic region) among humans, chimpanzees,
horses, mice, and rats (Fig. 2 G and Fig. S6D). LINE1 retrotransposons were much
more abundant in the *FmiR* genomic region when compared to either all the
chromosomes or the *FmiR* proximal gene *Fmr1* (adjusted p-value < 0.05, one-way
ANOVA). Similarly, LINE1s were more enriched in the *SmiR* region than in the
SmiR proximal gene *Slitrk2* (adjusted p-value < 0.05, one-way ANOVA) (Fig. 2 G).
Since the length of LINE1 retrotransposons (~6kb) is intrinsically longer than that
of the SINEs (100-600bp), which may bias the results, we further calculated
proportions of the nearest TE classes for the *miR-506* family and all miRNAs (Fig.
2 H and Fig. S6E and Table S2). In humans, chimpanzees, horses, mice, and rats,
the *miR-506* family miRNAs displayed a significantly higher proportion of LINE1
when compared to all miRNAs (baseline) (p < 0.05, paired t-test) (Fig. 2 H and
Table S2), suggesting that LINE1 may involve in *miR-506* family expansion. Taken
together, these results suggest that the X-linked *miR-506* miRNAs were originally
derived from the MER91C DNA transposon and further expanded through LINE1-
mediated retrotransposition.
284
X-linked *miR-506* family miRNAs are predominantly expressed in
spermatogenic cells and sperm. Several previous studies have shown that the
X-linked *miR-506* family miRNAs are predominantly expressed in the testis of
multiple species (15, 16, 24, 37-39). By analyzing the publicly available small RNA
sequencing (sRNA-seq) datasets from multiple species, including humans, rhesus
monkeys, mice, rats, rabbits, dogs, and cows (27, 40, 41), we further confirmed
that *miR-506* family miRNAs were indeed highly abundant in the testis, but barely
expressed in other organs (Fig. S7, and Table S3). To further determine whether
these miRNAs are expressed in male germ cells in rodent testes, we conducted
sRNA-seq using pachytene spermatocytes, round spermatids, and sperm purified
from adult mice (Fig. S8A and Table S3). Consistent with previous data (15, 16,
24), these miRNAs were abundantly expressed in spermatogenic cells in murine
testes (Fig. 3A). ~80% of these miRNAs were significantly upregulated (FDR <
0.05) when pachytene spermatocytes developed into round spermatids, and ~83.3%
were significantly upregulated (FDR < 0.05) when developed into cauda sperm
(Fig. 3A). By analyzing the publicly available sRNA-seq datasets from humans (42),
marmosets (37) and horses (43), we determined the expression patterns of the
miR-506 family miRNAs in the testes of these species (Fig. 3 B, C and D, Table
S3). The significantly increasing abundance of the SmiRs from immature to mature

testes in horses (43) supports the elevated expression in haploid male germ cells
(round, elongating/elongated spermatids, and sperm) compared to meiotic male
germ cells (spermatocytes) (Fig. 3D). More interestingly, the SmiRs and FmiRs
appear to be differentially expressed in the testes of various species, e.g., relative
levels of the SmiRs were greater than those of the FmiRs in mice (Fig. 3 A and
Table S3). Both the SmiRs and FmiRs were highly expressed in horses (Fig. 3D
and Table S3). Levels of the FmiRs in marmoset (37) and human testes (42) were
much greater than in those of the SmiRs (Fig. 3 B and C and Table S3). Overall,
the *miR-506* family miRNAs are abundant in the testis and predominantly
expressed in haploid male germ cells, *i.e.,* spermatids and spermatozoa.
314
Ablation of X-linked *miR-506* family miRNAs compromises male fertility due
to reduced sperm competitiveness. To define the physiological role of the *miR-*
506 family, we sequentially deleted these miRNA genes using CRISPR-Cas9-
based genome editing (Fig. 4A) (15). We first generated the KO mice lacking either
the *miR-883* single cluster (*miR-883* sKO) or the *miR-465* single cluster (*miR-465*
sKO) (Fig. 4A), as these two clusters are the most abundantly expressed in the
mouse testes (Fig. 3 A and Fig. S7). No discernable defects were observed, and
these KO males developed normally and were fertile (Fig. 4 B and C). On the *miR-*
883 sKO background, we further deleted the *miR-741* cluster, which we termed
double KO (dKO) (Fig. 4 A), but no discernable abnormalities were observed in the
dKO males either (Fig. S8 B-E). On the dKO background, we next deleted either
the *miR-465*, termed triple KO (tKO), or the *miR-471* and *miR-470* clusters, termed
quadruple KO (quadKO) (Fig. 4 A). Lastly, we ablated the *miR-465* cluster on the
quadKO background, named quintuple KO (quinKO) (Fig. 4 A). To reduce potential
off-target effects due to multiple rounds of CRISPR-Cas9 targeting, we also
generated a KO mouse line with only 4 guide RNAs (gRNAs) flanking the SmiR
region, named X-linked SmiR KO (XS) (Fig. 4 A). The XS mice were genetically
equivalent to the quinKO mice, and phenotypically identical to quinKOs (Fig. 4 B
and C), suggesting the phenotype observed was not due to the accumulating off-
target effects. To further exclude the potential off-target effect, all KO mouse
strains were backcrossed with WT C57BL/6J mice for at least 5 generations before
data collection. In addition, T7 endonuclease I (T7EI) assays showed no
discernable off-target effects in the quinKO mice (Fig. S8F).
While the litter size was still comparable between the tKO and WT control
mice, the quadKO, quinKO, and XS males produced significantly smaller litters (~5
vs. ~8 pups/litter) (adjusted p-value < 0.05, One-Way ANOVA) (Fig. 4 B). Of
interest, no significant changes were detected in litter interval, testis weight or
histology in any of the four types of KOs, as compared to WT controls (Fig. 4 C, D
and G). Computer-assisted sperm analyses (CASA) revealed no significant
differences in sperm counts and motility parameters among the four types of KOs
(Fig. 4 E and F and Fig. S8H). Overall, there appears to be an inverse correlation
(R^2 = 0.9139, p < 0.05, F-test) between the number of miRNAs inactivated and the
litter size (Fig. S8G). Interestingly, several human studies have correlated the
dysregulated *miR-506* family miRNAs with impaired male fertility due to maturation
arrest and oligo-asthenozoospermia (Table S4)(44-48). These data suggest that

the *miR-506* family may play an important role in spermatogenesis and male
fertility.
352   Most of the protein-coding genes that are exclusively or preferentially
expressed in the testis with an essential role in spermatogenesis are highly
conserved across species (49). Despite their male germ cell-predominant
expression, the *miR-506* family miRNAs appear to have evolved rapidly to diverge
their sequences, suggesting that these miRNAs might control certain "non-
conserved" aspects of spermatogenesis, leading to enhanced sperm
competitiveness for male reproductive success. Supporting this hypothesis,
previous reports have documented that females of most species throughout the
animal kingdoms mate with multiple males before pregnancy, suggesting that
sperm competition may serve as a selection mechanism to bias the birth of
offspring sired by the males with more competitive sperm (20, 21). Studies have
also shown female rodents in the wild mate with multiple males and produce litters
of mixed paternity, and that pups born to the females following such polyandrous
mating display greater survival rates than those produced from females following
monandrous mating (50). Given that CASA detected no difference in swimming
patterns between quinKO and WT sperm (Fig. S8H), we next carried out sperm
competition experiments that mimic polyandrous mating in the wild. Since the *miR-
506* family miRNAs are X-linked, the Y sperm from the quinKO mice are genetically
indistinguishable from those of WT controls. We, therefore, adopted the mTmG
male mice (51) for sperm competition experiments because the embryos or
offspring fathered by the mTmG males can be easily identified based on the
constitutively expressed membrane-tagged tomato red (mT) fluorescence and/or
PCR genotyping.
375   We first conducted sequential mating with two mating events ~6-8 hours
apart. Interestingly, all of the pups born were fathered by mTmG males (n=8) when
the WT females were mated first with mTmG males and subsequently with the
quinKO males. In contrast, when the WT females were mated first with quinKO
males and subsequently with mTmG males, ~89% of the pups born were fathered
by quinKO males, and the remaining ~11% of pups were from mTmG males (n=28)
(Fig. 4 H-I). It is noteworthy that in the sequential mating experiments, the two
coituses occurred ~6-8 hours apart due to practical reasons, whereas in the wild,
polyandrous mating may take place much faster. To better mimic polyandrous
mating *in vitro*, we mixed the WT and quinKO sperm in different ratios and used
the mixed sperm to perform *in vitro* fertilization (IVF). MII oocytes fertilized by
mTmG, quinKO, or a mixture of two types of sperm at three ratios (mTmG: quinKO
= 1:1, 4:1 and 1:4) all displayed comparable rates at which fertilized oocytes
developed into blastocysts (Fig. S8I). Interestingly, when a 1:1 ratio (mTmG sperm:
quinKO sperm) was used, ~73% of the resulting blastocysts were derived from
mTmG sperm, whereas the remaining ~27% were from quinKO sperm (n=179) (p
< 0.0001, Chi-squared test) (Fig. 4 J). When a 4:1 sperm ratio (mTmG: quinKO)
was used, ~92% of the blastocysts were from mTmG sperm and only 8% were
from quinKO sperm (n=170) (p < 0.05, Chi-squared test) (Fig. 4 J). In contrast,
when a 1:4 sperm ratio (mTmG: quinKO) was used, blastocysts derived from
mTmG and quinKO sperm represented ~28% and ~72% of the total, respectively

(n=135) (Fig. 4 J). We also performed co-artificial insemination (AI) using mTmG
and quinKO sperm. When a 1:1 sperm ratio (mTmG: quinKO) was used, ~62.5%
of the embryos (n=96) were derived from mTmG sperm (p< 0.05, Chi-squared test)
(Fig. 4 K). When a 1:4 ratio was used, ~35.3% of the embryos (n=17) were from
the mTmG mice (Fig. 4K and Fig. S8J). Together, these results indicate that the
quinKO sperm are less competitive than the control mTmG sperm both *in vivo* and
in vitro. Previous studies suggest that sperm aggregation and midpiece size might
be involved in sperm competitiveness (52, 53), but no changes in these two
parameters were observed in the quinKO sperm (Fig. S8 K and L). Although the
blastocyst rate (out of 2-cell embryos) of quinKO was comparable to that of the
mTmG mice, the 2-cell rates (out of zygotes) were significantly reduced (p< 0.05,
paired t-test) in the quinKO mice (~39%, n=67) when compared to the mTmG mice
(~88%, n=58) (Fig. S8M), implying that the quinKO sperm are indeed less efficient
in fertilizing eggs and/or supporting early embryonic development, especially the
first cleavage of the zygotes.
411
X-linked *miR-506* family miRNAs mostly target the genes involved in
spermatogenesis and embryonic development and compensate for each
other. To identify the target genes of these X-linked *miR-506* family miRNAs, we
performed RNA-seq analyses using testis samples from the five types of KOs
(*miR-465* sKO, dKO, tKO, quadKO, and quinKO) (Fig. S9A and Table S5).
Comparisons between the KO and WT testes revealed thousands of differentially
expressed genes (DEGs) (fold change ≥ 2, FDR< 0.05, Fig. S9A and Table S5).
The DEGs identified were then compared with the predicted *miR-506* target genes
using four different databases, including TargetScan (54), microrna.org (55),
miRWalk (56) and mirDB (57), to predict the differentially expressed targets (DETs)
of the *miR-506* family miRNAs (Fig. S9A and Table S5). We obtained 2,692, 2,028,
1,973, 3,405, and 1,106 DETs from *miR-465* sKO, dKO, tKO, quadKO and quinKO
testes, respectively. GO terms of DETs from each KO testis revealed that the
DETs were mostly involved in embryonic development, response to stimulus,
centrosome cycle, epithelium morphogenesis, organelle organization, cell
projection, RNA metabolic process, and DNA repair (Fig. S9B). The 431 DETs
identified to be shared across all five KO testes were also enriched in similar
pathways (Fig. 5 An and B). Several genes, including *Crisp1*, *Egr1*, and *Trpv4*,
were selected for validation using qPCR, Western blots and luciferase-based
reporter assays. Consistent with the RNA-seq data, qPCR showed that *Crisp1*,
Egr1, and *Trpv4* were significantly downregulated in the quinKO testes (Fig. S9C).
Western blots also confirmed that *CRISP1* is downregulated in the quinKO testis
when compared to the WT testis (Fig. S9D). Luciferase assays further confirmed
that *Egr1* and *Crisp1* are targets of the *miR-506* family members (Fig. S9 E and F).
Egr1 3'UTR luciferase activity was upregulated by *miR-465c*, while downregulated
by *miR-743b* (fig. S9E). *miR-465a*, *miR-465c*, *miR-470*, *miR-741*, and *miR-743a*
upregulated *Crisp1* 3'UTR luciferase activity, while *miR-743b* exerted the opposite
effect (Fig. S9F). Of interest, KO of *Crisp1* in mice or inhibition of *CRISP1* in
human sperm appears to phenocopy the quinKO mice (58, 59), e.g., sperm motility
in the *Crisp1* KO mice is comparable to that in WT mice, but their ability to

penetrate the eggs was reduced in the *Crisp1* KO mice (58); a similar effect was
also observed in human sperm treated with anti-hCRISP1 antibody (59).
The inverse correlation between the number of miRNAs inactivated and the
severity of the phenotype strongly hints that these miRNAs compensate for each
other (Fig. 4B and Fig. S8G). To test this hypothesis, we performed sRNA-seq on
four KO (*miR-465* sKO, tKO, quadKO and quinKO) testes. The sRNA-seq data
showed that these miRNAs were no longer expressed in the corresponding KOs,
confirming the successful deletion of these miRNAs in these KOs (Fig. 5C and
Table S6). Interestingly, in *miR-465* sKO testes, *miR-201, miR-547, miR-470*, *miR-
471*, *miR-742, miR-871*, *miR-881, miR-883a,* and *miR-883b* were all significantly
upregulated (FDR < 0.05). Similarly, *miR-201, miR-547, miR-470*, *miR-471*, *miR-
871,* and *miR-883b* were all significantly upregulated in the tKO testes (FDR <
0.05); *miR-201* and *miR-547* were all significantly upregulated in the quadKO and
the quinKO testes (FDR < 0.05) (Fig. 5 C and Table S6). These results support the
notion that genetic compensation exists among the X-linked *miR-506* family
miRNAs.
458
Rapid evolution of the *miR-506* family is not driven by the increased
complexity of 3'UTRs of the conserved targets but rather adaptive to
targeting more genes. To minimize false positives, the DETs in mice were
selected using the following criteria: (1) dysregulated by fold change ≥ 2 and FDR
< 0.05. (2) Falling within the predicted targets. (3) Intersected with at least two
different KO mouse samples. Using the 3,043 DETs identified in mice as a
reference, we searched the predicted targets of the *miR-506* family miRNAs in rats
and humans to determine if these target genes were shared across species (Fig.
5A). While 2,098 (~69%) target genes were shared among all three species, 2,510
(~82%) were common to both humans and mice, and 2,202 (~72%) were shared
between mice and rats (Fig. 6 A and Table S7). To test the accuracy of the
predicted targets, we selected several genes in humans and performed luciferase
assays using human *miR-506* family miRNAs and their corresponding target genes
(Fig. S10 A and B). Among these targets, *Crisp1* and *Fmr1* were shared among
humans, mice, and rats, and confirmed to be targeted by the *miR-506* family
miRNAs in mice (fig. S9C, S9D, S9F, and ref 11 and 15)(11, 15). Luciferase
assays also confirmed that human *CRISP1* (*hCRISP1*) and *FMR1*(*hFMR1*) were
targets of the *miR-506* family, and *miR-510* and *miR-513b* both could activate
hCRISP1 3'UTR luciferase activity (Fig. S10A), whereas *miR-509-1, miR-509-2*,
miR-509-3, *miR-513b*, *miR-514a*, and *miR-514b* could enhance *hFMR1* 3'UTR
luciferase activity (Fig. S10B). These results confirmed the accuracy of our
predicted targets in humans.
We considered two likely explanations for the paradox where the majority of
their target genes were shared across species despite the rapid evolution of the
miR-506 family miRNAs: 1) The 3'UTR sequences in extant target genes became
increasingly divergent during evolution such that the *miR-506* family miRNAs had
to adapt to maintain their ability to bind these 3'UTRs; or 2) that the *miR-506* family
miRNAs evolved rapidly in a manner that allowed them to target mRNAs encoded
by additional genes involved in spermatogenesis. To distinguish the two

possibilities, we first compared the extent of similarities among the 3'UTR
sequences of the 2,510 shared target genes between humans and mice (Fig. 6 A
and Table S7). We adopted the PhyloP scores to measure the evolutionary
conservation at individual nucleotide sites in the 3'UTRs of the shared target genes.
The overall conservation appeared to be greater in the regions targeted by *miR-506* family miRNAs than in the non-target regions in the 3'UTRs of the shared
target genes in both mice and humans (Fig. 6 B and C) with few exceptions (Fig. 6
D and E) ($p < 0.05$, t-test). These data suggest that the regions targeted by the X-
linked *miR-506* family miRNAs are under relatively stronger purifying, rather than
adaptive, selection. We then tested the second hypothesis that the rapid evolution
of the *miR-506* family resulted in more extant mRNAs being targeted by these
miRNAs. We first compared the average target numbers of each *miR-506* family
miRNA between humans and mice using the 2,510 shared targets between
predicted targets in humans and the dysregulated targets in mice (Fig. 6 A and F).
Among these shared targets, the human *miR-506* family members could target
~1,268 targets for each miRNA, whereas the miRNAs in mice could only target
~1,068 ($p < 0.05$, t-test) (Fig. 6 F), indicating that the *miR-506* family miRNAs had
gained more targets in humans when compared to those in mice. Furthermore, we
analyzed the number of all potential targets of the *miR-506* family miRNAs
predicted by the aforementioned four algorithms among humans, mice and rats.
The total number of targets for all the X-linked *miR-506* family miRNAs among
different species did not show significant enrichment in humans (Fig. S10C),
suggesting the sheer number of target genes does not increase in humans. We
then compared the number of target genes per miRNAs. When comparing the
number of target genes per miRNA for all the miRNAs (baseline) between humans
and mice, we found that human miRNAs have more target genes than murine
miRNAs ($p<0.05$, t-test) (Fig. S10D), implicating a higher biological complexity in
humans. This became even more obvious for the X-linked *miR-506* family ($p<0.05$,
t-test) (fig. S10D). In humans, the X-linked *miR-506* family targets significantly
more genes than the average of all miRNAs combined ($p<0.05$, t-test) (Fig. S10D).
In contrast, in mice, we did not observe a significant difference in target mRNA
number between X-linked miRNAs and all of the mouse miRNAs (mouse baseline)
(Fig. S10D). These results imply that human's X-linked *miR-506* family has gained
additional target genes relative to mice during recent mammalian evolution. We
also investigated the number of *miR-506* family miRNA targeting sites within the
individual target genes in both humans and mice, but no significant differences
were found between humans and mice (Fig. 6 G). Taken together, these results
suggest the human X-linked *miR-506* family has undergone additional selective
pressure causing them to gain additional regulatory functions after the evolutionary
divergence between primates and rodents, and that this has been manifested as
X-linked miRNAs targeting additional, novel mRNAs expressed during
spermatogenesis, rather than by these miRNAs simply targeting additional sites
within mRNAs that already targeted (Fig. 6 H).
531
532
533 **Discussion**

534
Successful reproduction is pivotal for the perpetuation of species, and sperm are
constantly facing selective pressures (60). To enhance their chance of fertilizing
the eggs, sperm need to adapt accordingly, and miRNAs-mediated regulation of
gene expression in spermatogenesis provides a rapidly adaptable mechanism for
this purpose. Although miRNAs were initially believed to be evolutionarily
conserved, the number of non-conserved miRNAs has been steadily increasing
(61). Among the non-conserved miRNAs, many are derived from TEs, suggesting
that TEs serve as a major source of miRNA sequences (61). The delayed
recognition of TE-derived miRNAs, in part, results from the fact that repetitive
sequences were usually excluded during the computational annotation of miRNAs.
In theory, TEs can serve as a good donor of miRNA sequences for the following
reasons: 1) TEs are ubiquitous and abundant in the genome and are known to
contribute to the regulatory elements of the coding genes, e.g., UTRs (62). As one
of the regulatory factors that target mainly the 3'UTRs, TE-derived miRNAs can
regulate a larger number of mRNAs with multiple miRNA-targeting sites. 2) TEs
are among the most rapidly evolving sequences in the genome (63) and thus, can
continuously produce species/lineage-specific miRNA genes to diversify their
regulatory effects. Consistent with these notions, the present study provides
evidence supporting that the *miR-506* family miRNAs originated from the MER91C
DNA transposons and subsequently underwent sequence divergence *via* LINE1-
mediated retrotransposition during evolution. Of more interest, the *miR-506* family
miRNAs, despite their rapid evolution, are all expressed in spermatogenic cells in
the testis, supporting a lineage-specific functional diversification of TE-derived
miRNAs. In fact, the *miR-506* family miRNAs were among the first reported TE-
derived miRNAs because of their location in a small region of the X chromosome
and their confined, abundant expression in the testis (16).
It has been demonstrated that under some circumstances, genes that
evolve under sexual conflicts should preferentially move to the X chromosome,
especially when they are male-beneficial, female-deleterious, and act recessively
(64, 65). The X chromosome is enriched with genes associated with male
reproductions (15, 16, 66). The rapid evolution of the X-linked *miR-506* family
strongly suggests that these miRNA genes were under selection to expand and
diversify their regulatory effects on spermatogenesis. Indeed, our data strongly
suggest that targeting new mRNAs was likely the driving force for the rapid
evolution of the *miR-506* family of miRNAs. However, the expansion and sequence
divergence of the X-linked *miR-506* family may simply reflect natural drifting
without functional significance, similar to some of the pachytene piRNA clusters
(30). We argue that the neutral drifting theory may not be true to the *miR-506*
family for the following reasons: 1) Despite highly divergent overall sequences of
the *miR-506* family, some miRNAs share the same seed regions across multiple
species, suggesting that these regions may undergo strong selections. 2) The *miR-*
506 family miRNAs, especially the FmiRs, are highly conserved in modern humans,
implying a strong selection of these miRNAs. 3) Knockout the *miR-506* family
(either quinKO or XS) results in male subfertility, reflecting a biological function. 4)
The quinKO sperm are less competitive than the WT sperm both in *vivo* and *in*

vitro. 5) Several human studies have linked the dysregulation of the *miR-506* family
with male infertility/subfertility (44-48). Similarly, one study in *Drosophila* also
showed that the rapidly evolving testis-restricted miRNAs underwent adaptive
evolution rather than neutral drifting (67).
Since TEs are abundant in UTRs, TE-derived miRNAs can target a much
greater number of mRNAs than those derived from distinct non-repetitive genomic
loci (61). Indeed, thousands of the dysregulated genes detected in the *miR-465*
sKO, dKO, tKO, quadKO and quinKO testes are involved in multiple pathways of
spermatogenesis. By analyzing the sequences divergence, we noticed that the
most common sequence substitutions among all of the *miR-506* miRNA sequences
were U-to-C and A-to-G, which were likely mediated by ADARs (adenosine
deaminases acting on RNA) that can change A to I (which is functionally equivalent
to G) (68). Interestingly, ~90% of the A-to-I editing appears to have occurred in Alu
elements (belonging to the SINE family), and some of the edits occurred in
miRNAs (68). Since G and U can form the so-called G-U wobble base pair (69),
those U-to-C or A-to-G substitutions can, in theory, target similar sequences and
exert regulatory functions (70), suggesting that the evolving miRNA sequences
could target not only the original sequences but also new sites with similar
sequences. Consistent with this notion, the predicted target genes of the *miR-506*
family in mice can also be found in rats and humans, suggesting the target genes
are shared across species despite the quick divergence of the miRNA sequences
across species. This is also supported by our data showing that the binding sites
for the *miR-506* family of miRNAs are more conserved than the surrounding, non-
targeting regions in the 3'UTRs of the predicted target mRNAs. Furthermore, seed
sequences among some *miR-506* family miRNAs remain the same despite the
high divergence of these miRNAs, and these conserved seed sequences appear to
be present in the dominant mature miRNAs. Thus, the seed region of these
miRNAs appears to have undergone strong selection. Supporting this notion,
previous studies have shown correlations between miRNA expression and the
evolution of miRNAs and target sites (71, 72). Of interest, miRNAs with the same
seed sequences may exert divergent functions. For example, the mature *miR-465a*,
miR-465b, and *miR-465c* only have a few mismatches outside of the seed region,
but only *miR-465c* exerts functional activation of *Egr1*. A similar effect has also
been reported in the *miR-465* cluster on the *Alkbh1* 3'UTR activity (18). Similarly,
despite the same seed sequences in the *miR-465* or *miR-743* cluster, *miR-465a*
and *miR-465c* have differential activating effects on the 3'UTR of *Crisp1*, and *miR-*
743a and *miR-743b* exert opposite effects on the *Crisp1* 3'UTR , further confirming
their functional divergence. Therefore, the sequences outside of the miRNA seed
region may play an important role in their functions, which have also been
observed in *C. elegans* and human HEK293 cells (18, 73, 74).
To unequivocally demonstrate the physiological role of miRNAs*,* it would be
ideal to delete not only the miRNAs but also their binding sites in their target
transcripts *in vivo.* A few previous studies have established the miRNA:target
relationship by deleting the miRNA-binding sites in target transcripts in *C. elegans*,
Drosophila, and cell lines (75-77), but similar studies have not been reported in
mice or humans. The strategy may work in mRNAs with 3'UTRs containing only

one or two miRNA-binding sites, but for more complex 3'UTRs of mRNAs in mice
and humans that often contain multiple binding sites for the same or different
miRNAs, deletion of one miRNA binding site may not cause any discernable
effects as the loss of function can easily be compensated by other miRNAs.
Nevertheless, by deleting all five clusters of the *miR-506* family one by one, we
were able to overcome the compensatory effects among the family
members/clusters and successfully revealed the physiological role of this miRNA
family.
It is well-known that mice in the wild are promiscuous, and one female often
mates with multiple males sequentially, giving rise to polyandrous litters derived
from sperm from more than one sire (20, 21). Polyandrous mating establishes a
situation where sperm from multiple males co-exist in the female reproductive tract,
with the most competitive ones fertilizing eggs and producing offspring (22, 78).
Therefore, a male that may be fertile in the monandrous mating scheme, may
rarely sire offspring in a polyandrous mating scenario, rendering this male
functionally "infertile". Therefore, sperm competitiveness reflects the general
reproductive fitness of the male (78). Although the quinKO males tend to produce
smaller litters under the monandrous mating scheme, their sperm counts, sperm
motility and morphology are indistinguishable from those of WT sperm. This is not
surprising given that miRNAs of the *miR-506* family most likely function to control
certain non-essential aspects of spermatogenesis. Sperm can be subject to
competition at multiple steps during fertilization, including their migration through
the female reproductive tract (cervix, uterine cavity and oviduct), binding the
cumulus-oocyte complexes, penetration of zona pellucida, etc. Therefore, IVF may
not be ideal for evaluating sperm competition in the real world as it bypasses
several key sites where sperm competition likely takes place. Artificial insemination
(AI) may represent a better way to assess sperm competition than IVF, but it is
probably less desirable than polyandrous mating for the following reasons: First, in
the wild, sperm from two males rarely, if not never, enter the female reproductive
tract simultaneously. We had tried to place two males into the cage with one
female, but the two males ended up fighting, and the submissive one never mated.
Second, sperm are delivered directly into the uterus or oviduct during AI (79, 80),
thus bypassing the potential sites for sperm competition (e.g., cervix and uterine
cavity). Although our breeding scheme also involves sperm competition, by
shortening the time between the two mating events in a laboratory setting, the
sequential mating method reported here may be further improved to better mimic
the natural polyandrous mating in the future. Moreover, future analyses of the
quinKO sperm may help identify biochemical or molecular biomarkers for sperm
competitiveness.
In summary, our data suggest that the *miR-506* family miRNAs are derived
initially from the MER91C DNA transposon and their rapid evolution results from
LINE1-meditated transposition to diversify their regulatory effects on both
conserved and non-conserved genes involved in spermatogenesis. These miRNAs
share many of their targets and can compensate for each other's absence, and
they work jointly through regulating their target genes in spermatogenesis to
ensure sperm competitiveness and reproductive fitness of the male.

672
673
Materials and Methods

KO mice were generated using CRISPR-Cas9-based genome editing, as
described (15). F1 KO mice were backcrossed for at least five generations before
data collection. Proportions of TEs were quantified based on the genomes
retrieved from the UCSC genome browser. PhyloP and phastCons were applied to
compare the conservation among miRNAs, and phyloP was used to evaluate the
conservation of miRNA target genes. RNA-seq and sRNA-seq were performed to
identify differential expressed mRNAs and miRNAs, respectively. qPCR, WB, and
luciferase assays were carried out to validate miRNA targets. Detailed methods
can be found in supplementary methods.
684
685

Acknowledgments

We would like to thank Dr. Kevin J. Peterson, Dartmouth College, Hanover, NH, for
his help with phylogenetic analyses of the *miR-506* miRNA genes. This work was
supported by grants from the NIH (HD071736, HD085506, HD098593, HD099924,
and P30GM110767 to WY), NIH/National Center for Advancing Translational
Science (NCATS) UCLA CTSI (UL1TR001881-01 to ZW and WY), and the
Templeton Foundation (PID: 61174 to WY). Work in the Lai lab was supported by
the NIH (R01-GM083300 and R01-HD108914) and Memorial Sloan-Kettering
Institute Grant P30-CA008748.

References

1. E. F. Oakberg, Duration of spermatogenesis in the mouse. *Nature* **180**, 1137-1138 (1957).
2. B. Muciaccia *et al.*, Novel stage classification of human spermatogenesis based on acrosome development. *Biol Reprod* **89**, 60 (2013).
3. N. H. Gu, W. L. Zhao, G. S. Wang, F. Sun, Comparative analysis of mammalian sperm ultrastructure reveals relationships between sperm morphology, mitochondrial functions and motility. *Reprod Biol Endocrinol* **17**, 66 (2019).
4. L. Hermo, R. M. Pelletier, D. G. Cyr, C. E. Smith, Surfing the wave, cycle, life history, and genes/proteins expressed by testicular germ cells. Part 1: background to spermatogenesis, spermatogonia, and spermatocytes. *Microsc Res Tech* **73**, 241-278 (2010).
5. P. Khaitovich, W. Enard, M. Lachmann, S. Paabo, Evolution of primate gene expression. *Nat Rev Genet* **7**, 693-702 (2006).
6. R. K. Idler, W. Yan, Control of messenger RNA fate by RNA-binding proteins: an emphasis on mammalian spermatogenesis. *J Androl* **33**, 309-337 (2012).
7. S. Ro, C. Park, K. M. Sanders, J. R. McCarrey, W. Yan, Cloning and expression profiling of testis-expressed microRNAs. *Dev Biol* **311**, 592-602 (2007).
8. Q. Wu *et al.*, The RNase III enzyme DROSHA is essential for microRNA production and spermatogenesis. *J Biol Chem* **287**, 25173-25190 (2012).
9. M. D. Papaioannou *et al.*, Sertoli cell Dicer is essential for spermatogenesis in mice. *Dev Biol* **326**, 250-259 (2009).
10. Y. Zhang *et al.*, MicroRNAs control mRNA fate by compartmentalization based on 3' UTR length in male germ cells. *Genome Biol* **18**, 105 (2017).
11. M. Guo *et al.*, Uncoupling transcription and translation through miRNA-dependent poly(A) length control in haploid male germ cells. *Development* **149** (2022).
12. D. P. Bartel, MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215-233 (2009).
13. C. Mayr, Evolution and Biological Roles of Alternative 3'UTRs. *Trends Cell Biol* **26**, 227-237 (2016).
14. P. J. Thompson, T. S. Macfarlan, M. C. Lorincz, Long Terminal Repeats: From Parasitic Elements to Building Blocks of the Transcriptional Regulatory Repertoire. *Mol Cell* **62**, 766-776 (2016).
15. Z. Wang *et al.*, X-linked miR-506 family miRNAs promote FMRP expression in mouse spermatogonia. *EMBO Rep* **21**, e49024 (2020).
16. R. Song *et al.*, Many X-linked microRNAs escape meiotic sex chromosome inactivation. *Nat Genet* **41**, 488-493 (2009).
17. J. Bao *et al.*, MicroRNA-449 and microRNA-34b/c function redundantly in murine testes by targeting E2F transcription factor-retinoblastoma protein (E2F-pRb) pathway. *J Biol Chem* **287**, 21686-21698 (2012).

18. Z. Wang *et al.*, Ablation of the miR-465 Cluster Causes a Skewed Sex Ratio
in Mice. *Front Endocrinol (Lausanne)* **13**, 893854 (2022).
19. E. A. Miska *et al.*, Most Caenorhabditis elegans microRNAs are individually
not essential for development or viability. *PLoS Genet* **3**, e215 (2007).
20. M. D. Dean, K. G. Ardlie, M. W. Nachman, The frequency of multiple
paternity suggests that sperm competition is common in house mice (Mus
domesticus). *Mol Ecol* **15**, 4141-4151 (2006).
21. R. C. Firman, L. W. Simmons, Polyandry facilitates postcopulatory
inbreeding avoidance in house mice. *Evolution* **62**, 603-611 (2008).
22. G. A. Parker, Sperm competition and its evolutionary consequences in the
insects. *Biological reviews* **45**, 525-567 (1970).
23. R. P. Meisel, T. Connallon, The faster-X effect: integrating theory and data.
Trends Genet **29**, 537-544 (2013).
24. F. Zhang *et al.*, Evolution of an X-Linked miRNA Family Predominantly
Expressed in Mammalian Male Germ Cells. *Mol Biol Evol* **36**, 663-678
(2019).
25. J. Casper *et al.*, The UCSC Genome Browser database: 2018 update.
Nucleic Acids Res **46**, D762-D769 (2018).
26. F. Cabanettes, C. Klopp, D-GENIES: dot plot large genomes in an
interactive, efficient and simple way. *PeerJ* **6**, e4958 (2018).
27. B. Fromm *et al.*, MirGeneDB 2.0: the metazoan microRNA complement.
Nucleic Acids Res **48**, D1172 (2020).
28. A. Kozomara, M. Birgaoanu, S. Griffiths-Jones, miRBase: from microRNA
sequences to function. *Nucleic Acids Res* **47**, D155-D162 (2019).
29. M. Byrska-Bishop *et al.*, High-coverage whole-genome sequencing of the
expanded 1000 Genomes Project cohort including 602 trios. *Cell* **185**, 3426-
3440 e3419 (2022).
30. D. M. Ozata *et al.*, Evolutionarily conserved pachytene piRNA loci are highly
divergent among modern humans. *Nat Ecol Evol* **4**, 156-168 (2020).
31. N. V. Fedoroff, Presidential address. Transposable elements, epigenetics,
and genome evolution. *Science* **338**, 758-767 (2012).
32. Z. Wang *et al.*, Efficient genome editing by CRISPR-Mb3Cas12a in mice. *J
Cell Sci* **133** (2020).
33. A. McKenna *et al.*, Whole-organism lineage tracing by combinatorial and
cumulative genome editing. *Science* **353**, aaf7907 (2016).
34. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local
alignment search tool. *J Mol Biol* **215**, 403-410 (1990).
35. J. Giordano *et al.*, Evolutionary history of mammalian transposons
determined by genome-wide defragmentation. *PLoS Comput Biol* **3**, e137
(2007).
36. Z. Wang *et al.*, Insertion of a chimeric retrotransposon sequence in mouse
Axin1 locus causes metastable kinky tail phenotype. *Mob DNA* **10**, 17
(2019).
37. T. Hirano *et al.*, Small RNA profiling and characterization of piRNA clusters
in the adult testes of the common marmoset, a model primate. *RNA* **20**,
1223-1237 (2014).

38.   R. Zhang, Y. Peng, W. Wang, B. Su, Rapid evolution of an X-linked
microRNA cluster in primates. *Genome Res* **17**, 612-617 (2007).
39.   E. M. Koenig *et al.*, The beagle dog MicroRNA tissue atlas: identifying
translatable biomarkers of organ toxicity. *BMC Genomics* **17**, 649 (2016).
40.   A. Keller *et al.*, miRNATissueAtlas2: an update to the human miRNA tissue
atlas. *Nucleic Acids Res* **50**, D211-D221 (2022).
41.   P. R. Bushel *et al.*, RATEmiRs: the rat atlas of tissue-specific and enriched
miRNAs for discerning baseline expression exclusivity of candidate
biomarkers. *RNA Biol* **17**, 630-636 (2020).
42.   I. V. Gainetdinov *et al.*, Assessment of piRNA biogenesis and function in
testicular germ cell tumors and their precursor germ cell neoplasia in situ.
BMC Cancer **18**, 20 (2018).
43.   B. Li *et al.*, Identification of piRNAs and piRNA clusters in the testes of the
Mongolian horse. *Sci Rep* **9**, 5022 (2019).
44.   M. Abu-Halima *et al.*, Altered microRNA expression profiles of human
spermatozoa in patients with different spermatogenic impairments. *Fertil*
Steril **99**, 1249-1255 e1216 (2013).
45.   Z. Heidary, M. Zaki-Dizaji, K. Saliminejad, H. R. Khorram Khorshid,
MicroRNA profiling in spermatozoa of men with unexplained
asthenozoospermia. *Andrologia* **51**, e13284 (2019).
46.   H. Tian *et al.*, Semen-specific miRNAs: Suitable for the distinction of infertile
semen in the body fluid identification? *Forensic Sci Int Genet* **33**, 161-167
(2018).
47.   X. Qing *et al.*, Dysregulation of an X-linked primate-specific epididymal
microRNA cluster in unexplained asthenozoospermia. *Oncotarget* **8**, 56839-
56849 (2017).
48.   C. Wang *et al.*, Altered profile of seminal plasma microRNAs in the
molecular diagnosis of male infertility. *Clin Chem* **57**, 1722-1731 (2011).
49.   B. Xia *et al.*, Widespread Transcriptional Scanning in the Testis Modulates
Gene Evolution Rates. *Cell* **180**, 248-262 e221 (2020).
50.   R. C. Firman, Polyandrous females benefit by producing sons that achieve
high reproductive success in a competitive environment. *Proc Biol Sci* **278**,
2823-2831 (2011).
51.   M. D. Muzumdar, B. Tasic, K. Miyamichi, L. Li, L. Luo, A global double-
fluorescent Cre reporter mouse. *Genesis* **45**, 593-605 (2007).
52.   H. S. Fisher, H. E. Hoekstra, Competition drives cooperation among closely
related sperm of deer mice. *Nature* **463**, 801-803 (2010).
53.   H. S. Fisher, E. Jacobs-Palmer, J. M. Lassance, H. E. Hoekstra, The
genetic basis and fitness consequences of sperm midpiece size in deer
mice. *Nat Commun* **7**, 13652 (2016).
54.   V. Agarwal, G. W. Bell, J. W. Nam, D. P. Bartel, Predicting effective
microRNA target sites in mammalian mRNAs. *Elife* **4** (2015).
55.   D. Betel, A. Koppal, P. Agius, C. Sander, C. Leslie, Comprehensive
modeling of microRNA targets predicts functional non-conserved and non-
canonical sites. *Genome Biol* **11**, R90 (2010).

56. H. Dweep, N. Gretz, miRWalk2.0: a comprehensive atlas of microRNA-
target interactions. *Nat Methods* **12**, 697 (2015).
57. Y. Chen, X. Wang, miRDB: an online database for prediction of functional
microRNA targets. *Nucleic Acids Res* **48**, D127-D131 (2020).
58. V. G. Da Ros *et al.*, Impaired sperm fertilizing ability in mice lacking
Cysteine-RIch Secretory Protein 1 (CRISP1). *Dev Biol* **320**, 12-18 (2008).
59. J. A. Maldera *et al.*, Human fertilization: epididymal hCRISP1 mediates
sperm-zona pellucida binding through its interaction with ZP3. *Mol Hum*
Reprod **20**, 341-349 (2014).
60. E. H. Morrow, How the sperm lost its tail: the evolution of aflagellate sperm.
Biol Rev Camb Philos Soc **79**, 795-814 (2004).
61. J. Piriyapongsa, L. Marino-Ramirez, I. K. Jordan, Origin and evolution of
human microRNAs from transposable elements. *Genetics* **176**, 1323-1337
(2007).
62. I. K. Jordan, I. B. Rogozin, G. V. Glazko, E. V. Koonin, Origin of a
substantial fraction of human regulatory sequences from transposable
elements. *Trends Genet* **19**, 68-72 (2003).
63. J. Gonzalez, D. A. Petrov, Evolution of genome content: population
dynamics of transposable elements in flies and humans. *Methods Mol Biol*
855, 361-383 (2012).
64. W. R. Rice, Sex Chromosomes and the Evolution of Sexual Dimorphism.
Evolution **38**, 735-742 (1984).
65. J. R. Gibson, A. K. Chippindale, W. R. Rice, The X chromosome is a hot
spot for sexually antagonistic fitness variation. *Proc Biol Sci* **269**, 499-505
(2002).
66. P. J. Wang, J. R. McCarrey, F. Yang, D. C. Page, An abundance of X-linked
genes expressed in spermatogonia. *Nat Genet* **27**, 422-426 (2001).
67. J. Mohammed *et al.*, Adaptive evolution of testis-specific, recently evolved,
clustered miRNAs in Drosophila. *RNA* **20**, 1195-1209 (2014).
68. K. Nishikura, A-to-I editing of coding and non-coding RNAs by ADARs. *Nat*
Rev Mol Cell Biol **17**, 83-96 (2016).
69. G. Varani, W. H. McClain, The G x U wobble base pair. A fundamental
building block of RNA structure crucial to RNA function in diverse biological
systems. *EMBO Rep* **1**, 18-23 (2000).
70. J. G. Doench, P. A. Sharp, Specificity of microRNA target selection in
translational repression. *Genes Dev* **18**, 504-511 (2004).
71. A. Simkin, R. Geissler, A. B. R. McIntyre, A. Grimson, Evolutionary
dynamics of microRNA target sites across vertebrate evolution. *PLoS Genet*
16, e1008285 (2020).
72. J. Meunier *et al.*, Birth and expression evolution of mammalian microRNA
genes. *Genome Res* **23**, 34-45 (2013).
73. J. P. Broughton, M. T. Lovci, J. L. Huang, G. W. Yeo, A. E. Pasquinelli,
Pairing beyond the Seed Supports MicroRNA Targeting Specificity. *Mol Cell*
64, 320-333 (2016).

74.   A. Helwak, G. Kudla, T. Dudnakova, D. Tollervey, Mapping the human
miRNA interactome by CLASH reveals frequent noncanonical binding. *Cell*
153, 654-665 (2013).
75.   N. Pinzon *et al.*, microRNA target prediction programs predict many false
positives. *Genome Res* **27**, 234-245 (2017).
76.   M. Ecsedi, M. Rausch, H. Grosshans, The let-7 microRNA directs vulval
development through a single target. *Dev Cell* **32**, 335-344 (2015).
77.   D. L. Garaulet, B. Zhang, L. Wei, E. Li, E. C. Lai, miRNAs and Neural
Alternative Polyadenylation Specify the Virgin Behavioral State. *Dev Cell* **54**,
410-423 e414 (2020).
78.   M. Gomendio, J. Martin-Coello, C. Crespo, C. Magana, E. R. Roldan,
Sperm competition enhances functional capacity of mammalian
spermatozoa. *Proc Natl Acad Sci U S A* **103**, 15113-15117 (2006).
79.   A. Nagy, M. Gertsenstein, K. Vintersten, R. Behringer, *Manipulating the*
mouse embryo: a laboratory manual (Cold Spring Harbor Laboratory Press
Cold Spring Harbor, NY:, 2003), vol. 10.
80.   B. J. Stone, K. H. Steele, A. Fath-Goodin, A rapid and effective nonsurgical
artificial insemination protocol using the NSET device for sperm transfer in
mice without anesthesia. *Transgenic Res* **24**, 775-781 (2015).
896

897   **Figures and Tables**
898
Figure 1. Genomic location, sequence alignment and evolution conservation
of the X-linked *miR-506* family.
A. Genomic location of the X-linked *miR-506* family miRNAs (black bars) and the
two flanking coding genes, *Slitrk2* (green blocks) and *Fmr1* (red blocks). The
number of miRNAs within each cluster is indicated underneath the miRNA clusters.
B. Evolution conservation of X-linked *miR-506* family based on Multiz Alignment &
Conservation using the human genome as a reference. Positive PhyloP scores
indicate conservation and vice versa. PhastCons has a score between 0~1, and
the higher the score, the more conserved the DNA region is. C-D. Comparison of
mean phyloP (C) phastCons (D) scores among CDS of *Slitrk2* and *Fmr1*,
intergenic region (IGR), pachytene piRNAs, SmiR, FmiR, and all miRNAs. **, ***,
**** indicate adjusted p-value < 0.01, 0.001, and 0.0001, respectively. ns, not
significant. Kruskal-Wallis test was used for statistical analyses. E-F. Comparison
of derived allele (E) and mean nucleotide (F) frequencies among pachytene
piRNAs, SmiR, FmiR, and all miRNAs. *, ** and **** indicate adjusted p-value <
0.05, 0.01 and 0.0001, respectively. ns, not significant. Kruskal-Wallis test was
used for statistical analyses.
916
Figure 2. Evolutionary history of the X-linked *miR-506* family.
A. Sequences alignment of FmiRs from various species using human MER91C
DNA transposon as the reference. The first line is the human MER91C DNA
transposon, and below are the miRNAs of various species. Mismatched
nucleotides are highlighted with various colors. B. A phylogenetic tree of the
MER91C DNA transposons and the X-linked *miR-506* family miRNAs. The
MER91C DNA transposons are labeled in purple. C. RNA structure of the MER91C
DNA transposon-derived miRNA (hsa-miR-513c). D. sRNA-seq reads (lower panel)
of the MER91C DNA transposon-derived miRNA, hsa-miR-513c. E.
Representative gel images showing expression levels of the MER91C DNA
transposon-derived miRNA (hsa-miR-513a1) in HEK293T cells. The asterisk (*)
indicates the expected miRNA size. U6 was used as the loading control. F. qPCR
analyses of expression levels of MER91C DNA transposon-derived miRNA (hsa-
miR-513a1) in HEK293T cells. *, **, **** indicate adjusted p-value < 0.05, 0.01,
0.0001, respectively. One-way ANOVA was used for statistical analyses. G. The
proportions of LINE1 in the genomic regions of all chromosomes, *Fmr1*, *FmiR*,
Slitrk2, and *SmiR* in humans, chimpanzees, horses, mice, and rats. * and ***
indicate adjusted p-value < 0.05 and p < 0.001, respectively. One-way ANOVA
was used for statistical analyses. H. The proportion of LINE1 between the closest
TEs to all miRNAs and to *miR-506* family miRNAs. ** indicates p < 0.01. Paired t-
test was used for statistical analyses.
938
939
940   **Figure 3. Expression profiles of X-linked *miR-506* family in mammalian testes**
941   **and male germ cells.**

A. Heatmaps showing the *miR-506* family expression in the testis, pachytene spermatocytes, round spermatids and sperm in mice. Biological triplicates of the testis samples (n=3) and duplicates of pachytene spermatocytes, round spermatids and sperm samples isolated from 2-4 mice were used for sRNA-seq. *, **, ***, and **** indicate FDR < 0.05, 0.01, 0.001, and 0.0001, respectively, when comparing round spermatids to pachytene spermatocytes. #, ##, ###, and #### indicate FDR < 0.05, 0.01, 0.001, and 0.0001, respectively, when comparing cauda sperm to pachytene spermatocytes. ns and na indicate not significantly and not not applicable, respectively.B-C. LogCPM bar graphs showing the *miR-506* family expression in the testis of humans (B) and marmosets (C). D. LogCPM bar graph showing the *miR-506* family expression in sexually immature and mature horse testes. n=3. *, **, ***, and **** indicate FDR < 0.05, 0.01, 0.001, and 0.0001, respectively.

Figure 4. Ablation of X-linked *miR-506* family miRNAs compromised sperm competitiveness and reproductive fitness in male mice.
A. Schematics showing the strategy used to generate six lines of KO mice lacking individual or combined miRNA clusters within the *miR-506* family using CRISPR-Cas9. B-C. Litter size (B) and litter interval (C) of six *miR-506* KO lines, at least 10 litters from 3 different breeding pairs for each KO line were counted. Dunnett's multiple comparisons test as the posthoc test following one-way ANOVA was used for the statistical analysis. ns, not significant. * and **indicate adjusted p-value <0.05 and 0.01, respectively. D-F. Analyses of testis weight (D), sperm counts (E) and sperm motility (F) in four *miR-506* KO lines. n ≥ 3 and Dunnett's multiple comparisons test as the posthoc test following one-way ANOVA was used for the statistical analysis. G. Testicular histology of WT and four *miR-506* KO lines showing largely normal spermatogenesis. Scale bars = 50 μm. H. Representative genotyping results of the sequential mating experiments. I. Sequential mating of WT female mice with mTmG and quinKO males. Upper panel, an overview of the polyandrous mating scheme. "mTmG V.S. quinKO": mTmG male mice mated first; "quinKO V.S. mTmG": quinKO male mice mated first. J. Percentage of mTmG blastocysts obtained from IVF using WT MII oocytes and mixed sperm from mTmG (control) and quinKO males at different ratios. Data were based on three independent IVF experiments. The expected ratio was indicated as the blue line. Chi-squared test was used for statistical analyses. * and **** indicate p<0.05 and 0.0001, respectively. K. Percentage of mTmG embryos obtained from co-artificial insemination using different ratios of mTmG and quinKO sperm. Data were based on 9 and 3 independent AI experiments for the 1:1 and 1:4 sperm ratio (mTmG: quinKO), respectively. The expected ratio was indicated as the blue line. Chi-squared test was used for statistical analyses. *, p<0.05.

Fig. 5. Target genes and genetic compensation of the X-linked *miR-506* family miRNAs. A. Intersections of the DETs among different KO testes. B. GO term enrichment analyses of the 431 DETs shared among the four different miR-

988 506 KO testes. C. MA plots showing the expression levels of the miR-506 family
989 miRNAs in WT, sKO, tKO, quadKO, and quinKO testes. Three biological replicates
990 (n=3) were used for sRNA-seq analyses.
991
992
Figure 6. Rapid evolution of the X-linked *miR-506* family miRNAs correlates
with increased complexity of genetic networks that regulate
spermatogenesis across mammalian species. A. Overlap between the
dysregulated targets in mice and the predicted targets in humans and rats. B.
Comparison of the cumulative distribution between the *miR-506* family targeting
sites and the other regions in humans. *: p< 0.05; t-test was used for statistical
analyses. C. Comparison of the cumulative distribution between the *miR-506*
family targeting sites and the other regions in mice. *: p< 0.05; t-test was used for
statistical analyses. D. Paired comparison of the PhyloP score between the *miR-*
506 family targeting sites and the other regions in humans. E. Paired comparison
of the PhyloP score between the *miR-506* family targeting sites and the other
regions in mice. F. Comparison of the number of the targets per miRNA for the X-
linked *miR-506* family in mice and humans. *: p< 0.05; t-test was used for
statistical analyses. G. The number of target sites within individual target mRNAs
in both humans and mice. H. Schematics show that human *miR-506* family
miRNAs gained more targets relative to those of mice during evolution.
1009