Main Manuscript for

Confinement by liquid-liquid interface replicates *in vivo* neutrophil deformations and elicits bleb based migration

Jonathan H. Schrope[1,2,3,4], Adam Horn[2], Kaitlyn Lazorchak[2,4] , Clyde W. Tinnen[3], Jack J Stevens[1,2], Mehtab Farooqui[3,5],Tanner Robertson[2], Jiayi Li[1], David Bennin[5], Terry Juang[1,3], Adeel Ahmed[5], Chao Li[5,7], Anna Huttenlocher[2,6,7,*], David J Beebe[1,3,5,7,*]

[1]Department of Biomedical Engineering, University of Wisconsin-Madison, Madison, WI, USA.
[2]Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI, USA.
[3]Department of Pathology and Laboratory Medicine, University of Wisconsin-Madison, Madison, WI, USA.
[4]Medical Scientist Training Program, University of Wisconsin-Madison, Madison, WI, USA.
[5]Carbone Cancer Center, University of Wisconsin-Madison, Madison, WI, USA.
[6]Department of Pediatrics, University of Wisconsin-Madison, Madison, WI, USA.
[7]These authors contributed equally.

* Co-corresponding authors: Anna Huttenlocher, David J Beebe

Email(s): huttenlocher@wisc.edu, djbeebe@wisc.edu

This PDF file includes:

Main Text
Figures 1 to 6

Abstract

Leukocytes navigate through interstitial spaces resulting in deformation of both the motile leukocytes and surrounding cells. Creating an *in vitro* system that models the deformable cellular environment encountered *in vivo* has been challenging. Here, we engineer microchannels with a liquid-liquid interface that exerts confining pressures (200-3000 Pa) similar to cells in tissues, and, thus, is deformable by cell generated forces. Consequently, the balance between migratory cell-generated and interfacial pressures determines the degree of confinement. Pioneer cells that first contact the interfacial barrier require greater deformation forces to forge a path for migration, and as a result migrate slower than trailing cells. Critically, resistive pressures are tunable by controlling the curvature of the liquid interface, which regulates motility. By granting cells autonomy in determining their confinement, and tuning environmental resistance, interfacial deformations are made to match those of surrounding cells *in vivo* during interstitial neutrophil migration in a larval zebrafish model. We discover that, in this context, neutrophils employ a bleb-based mechanism of force generation to deform a barrier exerting cell-scale confining pressures.

Significance Statement

Immune cells sense physical forces provided by surrounding cellular tissues to regulate their motility. Here, we introduce the use of liquid-liquid interfaces to model forces exerted by surrounding cells during interstitial motility *in vivo*. Neutrophils interacting with the interface employ a bleb-based mechanism of force generation to induce interfacial deformation. This work furthers our understanding of the mechanisms employed by immune cells to traverse through deformable barriers akin to cells in the body, and introduces a pioneering technology enabling the study of cell interaction with soft materials.

MAIN TEXT

Introduction

Precise regulation of leukocyte migration to sites of infection, malignancy, and injury is required for successful mounting and resolution of the immune response (1–4). Leukocyte motility is tightly regulated *in vivo* by a combination of chemical gradients and local mechanical cues arising from the surrounding physical environment (3–6). After exiting circulation, leukocytes migrate through tissue interstitial spaces and generate forces to overcome resistive forces applied by surrounding cells (3, 4, 6, 7). The balance between these leukocyte-generated and resisting forces regulates cell morphology and motility (4, 6). Existing *in vitro* migration platforms have uncovered that the degree of mechanical confinement and resulting variations in local cell shape regulate leukocyte speed, directionality, polarity, and migration mode (5, 8–12), thus implicating a role of local mechanical pressures in regulating the immune response. In the absence of confinement, leukocyte migration is largely mediated by actin polymerization-driven formation of a pseudopod at the leading edge (3, 4, 13, 14). However, under confinement, leukocytes can elect an alternative migration strategy driven by rear acto-myosin contractility driving membrane protrusions (blebs) through rapid fluctuations in intracellular hydrostatic pressure (12, 15–18).

It is well established that confining forces provided by the surrounding extracellular matrix (ECM) influence cell motility and effector function. However, our understanding of how interactions with surrounding cells themselves regulate motility is limited. The reasons for this are twofold; first, limitations in optical access of most *in vivo* models has thus far restricted capture of these single cell-cell physical interactions. Second, existing *in vitro* systems to study confined cell migration largely focus on cell-ECM interactions, or utilize materials orders of magnitude stiffer than cells themselves (5, 19–21) (**Fig. S1**). The development of ECM-mimics with tunable mechanical properties (22–24), and the ability to fabricate channels made of ECM-mimics (25–27), has yielded critical insights into how cell-matrix interactions modulate migration speed and directionality (28–31). However, such approaches neglect to model interactions with surrounding cells (stiffness ~ 200-3000 Pa on single cell length scales (20) with very different mechanical properties than matrix fibers (~300 MPa) (20). Alternative approaches utilize polymeric materials such as polyacrylamide, agarose, or polydimethylsiloxane (PDMS) based microchannels to study how controlled confinement of individual cells modulates cell traction (32, 33), polarity (34, 35), migration direction (10, 27, 36, 37), deformations (38), migration mode (18, 39), and speed (9, 40). However, the rigidity of these materials force cells into unnatural morphologies determined by pre-defined channel geometries. This has limited the ability to explore mechanisms of force generation employed by leukocytes to deform surrounding cells and forge paths through host tissues. Thus, there remains an unmet need for engineering an *in vitro* material interface that confines cells with physiologically relevant pressures to model physical interactions with neighboring cells and enable new insights into how interaction with soft materials regulates cell motility.

Here, we utilize differential surface patterning of a glass substrate to engineer microchannels on single cell length scales (i.e., channel height < 10 um) bound by a liquid-liquid interface. We examine interactions of primary neutrophils, the most abundant cells in the immune system and an exemplary model of amoeboid cell migration, with the liquid interface during chemotaxis. We find that the liquid-liquid interface exerts pressures comparable to cells themselves; sufficiently rigid for confinement, yet deformable in response to single cell generated forces. The result is greater autonomy granted to motile cells themselves to control the degree of confinement and thus morphology. Migrating neutrophils that reach the interface first (pioneer cells) require greater interfacial deformation than trailing cells and as a result migrate slower. Pioneer cells interacting with the interfacial barrier exhibit distinct morphological polarity characterized by frontward protrusions and rear-ward nuclear positioning. By careful tuning of the interfacial pressures resisting cell-generated forces, deformations of the interface are made to match those of epithelial cells during interstitial migration within a larval zebrafish model. While the majority of

observations of bleb-based migration have occurred under rigid confinement, here we find that under *in vivo*-like confinement, neutrophils elect a bleb-based mechanism of force generation driven by rear contractility to deform the interface. Altogether, this work introduces a tunable material interface that replicates single cell-scale confining forces within a system accessible to a broad range of researchers to study how cells sense and respond to interactions with soft barriers to regulate motility.

Results

Exclusive Liquid Repellency (ELR) enables liquid microchannels of heights less than single cells

In this section, the surface engineering principles enabling construction of liquid channels of heights less than a single cell is described. Recently, groups have engineered liquid-walled microfluidic channels by either printing aqueous media onto a chemically homogeneous solid substrate under oil overlay (41–43), or by depositing media onto superhydrophilic regions of a chemically patterned (i.e., heterogeneous) substrate (44). In either case, interfacial forces within the three phase (solid-aqueous-oil) system resist aqueous fluid expansion and thus pin the media to user-controlled geometries. In such strategies, however, obtaining channel heights on single cell length scales (< 10 μm leukocyte diameter) is challenging, as the forces pinning the aqueous phase to the desired geometry do not reach the theoretical maximum possible within a three-phase liquid-liquid-solid system. Here, we leverage our recent discovery of exclusive liquid repellency (ELR) (45–47) to control interfacial geometry with the theoretical maximum of aqueous-confining forces, enabling engineering of conduit heights comparable to that of single cells (~1-10 μm).

Engineering such channels is rooted in foundational principles of interfacial physics. The degree to which a solid surface repels a liquid within a three phase solid-liquid-liquid system is reflected by the Young's contact angle (θ) that a liquid droplet (Liquid 1) makes on the surface (**Fig. 1A, top**). Young's contact angle is determined by the balance between all interfacial energies (γ) present within the system. Meticulous engineering of this energy balance results in a surface exhibiting exclusive liquid repellency (ELR), that is, complete repulsion of a liquid, resulting in a contact angle of 180° (45, 46). Here, we differentially pattern a glass substrate to exhibit regions of ELR to aqueous media (Liquid 1) in the presence of silicon oil (Liquid 2) (i.e., Aqueous Repellent Surface (ARS)) (**Fig. 1A, middle**), or regions of ELR to silicon oil (Liquid 1) in the presence of media (Liquid 2) (i.e., Oil Repellent Surface (ORS)) (**Fig. 1A, bottom**). Controlling surface patterning of ARS and ORS regions to specific geometries (**Fig. S2**) results in a double-ELR system that provides the theoretical maximum of interfacial forces to pin both media and oil onto their preferred surface. Importantly, this enables the construction of channels bound by a liquid-liquid interface with heights less than single cell diameter (< 10 μm) (**Fig. 1B**). Media is deposited onto the pre-patterned surface by "sweeping" a hanging droplet of media across the substrate with a pipette to construct an array of > 50 channels on a standard microscope slide within seconds (**Fig. 1C**). Here, we implement a simple pattern that results in two hemispheric droplets connected by a single, linear, liquid channel (**Fig. 1D-E**). However the method allows for creation of an endless number of configurations (**Fig. S3**).

While collagen is the most abundant extracellular matrix (ECM) protein in the body and provides physical and chemical (48) cues to migrating cells, its high viscosity and rapid polymerization rate limits incorporation into traditional closed microchannels on sub-100 μm length scales. Existing methods often require external pumping of collagen into closed channels (49), decreasing throughput and increasing adoption barrier. Here, channels are constructed in open fluid, allowing for direct physical access so that performing the sweep technique with neutralized collagen yields microchannels on single cell length scales coated by fibrillar collagen (**Fig. 1F-G, Fig. S4**).

The liquid-liquid interface confines, yet deforms in response to, immobile cells

In microchannel systems composed of rigid walls, the degree of confinement, and thus cell shape is determined entirely by the channel geometry. However during interstitial migration *in vivo*,

leukocyte shape is determined by an equilibrium state between cell-generated and resistive pressures provided by the surrounding cellular environment. Thus, we sought to investigate whether the liquid-liquid interface 1) possesses sufficient rigidity to confine cells and 2) is "soft" enough to deform in response to single cell-scale forces (i.e. single cell stiffness). Primary human neutrophils were incorporated into the hanging droplet during the sweep and thus placed directly into channels of varying width (200 μm to 30 μm) (**Fig. 2A**). In the absence of cells, channel height is determined by an equilibrium state reflecting the pressure exerted by the aqueous phase itself; channel height (red line) increases with channel width (**Fig. 2C, Fig. S5A**). Cells placed into progressively smaller channels exhibited decreased cell height (blue line), implicating sufficient force applied by the interface to mechanically confine single cells (**Fig. 2B-C, Fig. S5B-F**). This demonstrates the ability to alter cell confinement by varying channel geometry. Notably, when channel height is significantly less than cell diameter (30 and 100 μm width channels), then cell height exceeds that of the channel alone (in the absence of cells). This implies that cells exert passive forces to deform the interface when initial channel height is sufficiently small. Indeed, confocal imaging along the length of a 30 μm width channel reveals that cells deform the interface from an initial equilibrium height ($h_{0,equilib}$) to an increased equilibrium height ($h_{cell,equilib}$) (**Fig. 2D**). Thus the interface is not a rigid barrier such as plastic or PDMS elastomer but is deformable by the presence of individual cells.

       In contrast to rigid substrates with fixed stiffness dependent on material properties alone, the pressure exerted by a liquid-liquid interface is dependent both on interfacial components (i.e. oil and media) and the curvature of the interface. Thus we next sought to quantify the confining pressures exerted on cells by the interface. According to the Law of Laplace, the pressure exerted by any curved liquid-liquid interface is a function of the interfacial tension (material dependent constant; γ_{OM} ~ 41.8 mN/m for water and silicone oil (46)) and the radii of interfacial curvature across all curved planes (side yz and cross-sectional xz in this case). Within this double ELR system that completely immobilizes the aqueous phase to the predefined channel width, deformations in channel height alter interfacial curvature (κ) and thus the reciprocal, radius of curvature ($R = 1/\kappa$) (**Fig. 2E**). The result is an equilibrium confining pressure that reflects the pressures exerted by cells themselves. Critically, interfacial curvature (and thus pressure) is controllable by altering channel width (**Fig. 2F**); interfacial pressures in the presence (blue) or absence (red) of cells decreases with increased channel width (**Fig. 2G**). This demonstrates the ability to control confining pressures by altering channel geometry. This mathematical framework would predict that the cell nucleus, the stiffest part of the cell, exerts the greatest pressure against the interface thus generating the greatest interfacial deformation. Indeed, the interfacial height is always greatest over the cell nucleus (**Fig. 2H**). Thus, in contrast to rigid-walled channels where cell confinement is purely a function of predefined geometrical constraints, these data support a model whereby the channels adopt a confining pressure reflecting an equilibrium defined by the physical properties of cells themselves (**Fig. 2I**).

Neutrophil motility is dependent on required deformation of the interface
Given that the interface exerts confining pressures comparable to cells themselves, we sought to examine how interaction with the deformable barrier regulates cell motility. Creation of channels followed by sequential addition of neutrophils to the inlet droplet opposite from chemoattractant in the outlet droplet establishes a chemotaxis assay under confinement (**Fig. 3A, Fig S6A-E, Movie S1**). In this assay, cells occupy a layer of culture media between the collagen layer and oil-media interface, thus chemotaxis would require active deformation of the interface to allow space for migration (**Fig. 3B**). The initial cells to enter the channel, denoted as "pioneer" cells, migrate slowly, followed by "trailing" cells that migrate rapidly (**Fig. 3C-D**). Confocal imaging reveals differential interfacial heights before and after pioneer cell passage (**Fig. 3E-F**), establishing that pioneer cells actively deform the interface to allow space for migration. Therefore, while both populations are confined by the interface in the transverse (z) direction (**Fig. S6F**), pioneer cells must also interact with the interface in the axial (y) direction at the cell front. It is reasonable that this mechanical resistance is responsible for decreased pioneer cell speed. However, it is also possible that pioneer cell migration alters the chemical gradient in a manner to increase motility of trailing cells. To investigate this, we added a combination of fMLP and FITC dye of similar

molecular weight (FITC 389.382 g/mol vs fMLP 437.56 g/mol) to visualize an approximate chemokine gradient during the course of migration (**Fig. S7A**). We found that pioneer cell passage did not significantly alter the chemical gradient (**Fig. S7B**). To further decouple positioning within the chemical gradient from required interaction with the interface, we analyzed a subset of trailing cells that migrate fast enough to catch up to and thus become pioneer cells. As these cells move through the stable global gradient, they significantly decrease in speed once they reach the interface (**Fig. 3G**). Taken together, these data indicate a purely mechanical mechanism whereby the interaction with (and deformation of) the interface restricts cell motility. Pioneer cells actively deform the channel interface, leaving decreased mechanical resistance for trailing cells which in turn migrate faster. From a technical standpoint, these results demonstrate the ability to decouple chemical cues such as position within a stable gradient, from mechanical cues of interaction with the interfacial barrier to isolate a role of physical cues on cell motility. Given the ability to tune pressures exerted by the interface, we next studied how alterations in the physical properties of the surrounding environment regulate cell motility.

Liquid interface generates a tunable gradient of confinement that regulates motility
In contrast to immobile cells, migratory cells typically exhibit morphological polarity, largely dependent on nuclear positioning as they extend cytoplasmic protrusions at the leading edge to deform the surrounding environment. Rigid microchannels impose fixed environmental stiffness that is not responsive to cell-generated forces, so that all cell components (nucleus and cytoplasm) typically assume the entirety of available space within the channel. Here, we find that pioneer neutrophils exhibit rear-ward nuclear positioning and front-ward cytoplasmic protrusions (**Fig. 4A, Fig. S8A, Movie S2**). These regions deform the interface to different degrees; the nucleus provokes the maximal interfacial deformation, characterized by increased interfacial curvature (**Fig. 4B**) and thus maximal confining pressures, compared to frontward protrusions (**Fig. 4C**). Thus in contrast to immobile cells directly placed into channels, which exhibit symmetric deformations in the xz and yz planes, pioneer cells interacting with the interface generate spatially-variant radii of curvature across both planes. The result is a gradient of confining pressures across the length of the cell. This gradient is tunable by altering channel width (**Fig. S8B**). Increased width generates a more shallow gradient across cell length (yz plane), with less significant differences in confining pressure between the forward protrusion and nucleus-containing rear (**Fig. 4D**). Therefore tuning interface curvature (xz plane) controls cell shape (yz plane) during pioneer cell migration. Bulk analysis confirms that increased channel width results in decreased confining pressures over both the cell nucleus (**Fig. S8D**) and frontward protrusions (**Fig. S8E**). Thus akin to interstitial migration *in vivo*, the greater the force generated by the cell, the higher resistive pressure exerted by the surrounding elastic environment.

    To explore how altering confining pressure gradients regulate motility, we engineered channels of varying width (30 vs 100 µm) and length (300 vs 500 µm, respectively) that exhibit different confining pressures yet similar chemical gradients (**Fig. S8B-C**). While the chemical gradient within 30 µm width channels was slightly steeper than that in 100 µm channels (typically increasing motility), cells migrated significantly slower, suggesting that increased confining pressures negatively impacts motility (**Fig. 4E**). Ultimately, these data indicate that the liquid-liquid interface is sensitive to differential pressures generated across a single migratory cell, and that cell morphology and motility is tunable by controlling channel geometry. These data would predict rearward nuclear positioning eliciting maximal surrounding cell deformation during passage between surrounding cells *in vivo*, which we demonstrate in a larval zebrafish model.

Tuning interface pressures recapitulates surrounding cell deformations during interstitial migration *in vivo*
Migratory leukocytes sense membrane deformations to regulate motility and effector function (50–53). Membrane shape *in vivo* is determined by a balance between outward cell generated forces and resistive pressures arising from surrounding cells. Thus, the magnitude of surrounding cell deformations is a reflection of the relative force relationship between migratory and surrounding cells. We next explored whether tuning the resistive pressures applied by the

interface could recapitulate relevant leukocyte morphology and surrounding cell deformations during interstitial migration *in vivo*.

Limitations in optical access of most *in vivo* models has hindered the ability to capture leukocyte physical interactions with surrounding cells during migration. Here, we leverage the genetic tractability and optical transparency of the larval zebrafish to capture deformations of neutrophils and surrounding keratinocytes during interstitial migration within the zebrafish epidermis (**Fig. 5A**). We performed high resolution imaging of spontaneous neutrophil motility within the epidermis of a dual transgenic larval zebrafish model expressing BFP in the neutrophil cytoplasm, mCherry in the nucleus and GFP in surrounding basal keratinocytes (*Tg(LyzC:TagBFP/LyzC:H2B-mCherry) x Tg(Krtt1c19e:acGFP)*). Such imaging reveals that neutrophils interact with and deform surrounding basal keratinocytes during interstitial migration (**Movie S3**). Neutrophils first extend a cytoplasmic protrusion to exert outward forces on surrounding keratinocytes, followed by a period of maximal keratinocyte deformation during passage of the cell body (**Fig. 5B**). As predicted within liquid channels, the nucleus is rear-ward oriented, and its passage induces maximal deformations of surrounding cells as the case of the liquid interface (**Fig. 5C**). Tuning the interfacial shape by controlling channel width (30 µm vs 100 µm) reproduces neutrophil protrusion width (**Fig. 5D**) and surrounding cell deformations during passage of the cell body (**Fig. 5E**) observed *in vivo*. These data indicate that tuning of baseline channel pressures reproduces relevant transient force balances during leukocyte interstitial migration *in vivo*, validating the relevance of this system to model physical cell-cell interactions during migration.

Neutrophils employ a bleb-based mechanism of force generation to deform an interface exerting cell-scale confining pressures

We next sought to explore the mechanism of force generation that cells employ at the leading edge to deform the interface and generate space for migration. On two-dimensional substrates, leukocytes largely rely on polymerization of branched actin networks stabilized by actin-related protein 2/3 (Arp2/3) to extend protrusions at the leading edge. Recent studies have demonstrated that under rigid confinement, neutrophils can employ a bleb-based motility mechanism independent of actin polymerization at the leading edge (18), however it is not known how cells generate forces to deform a surrounding elastic environment exerting cell scale confining forces. Blebbing is driven by Rho-kinase dependent contraction at the rear driving increased intracellular hydraulic pressure. This drives detachment and rapid projection of the cell membrane through the F-actin cortex at the cell front (**Fig. 6A**). Blebs form on significantly faster timescales (< 2 µm/sec and < 10 sec lifespans) than lamellipodia (12). We investigated the hypothesis that interaction with the interface provoked a bleb-based mechanism of migration to generate outward deforming forces.

We first examined migration of primary neutrophils under inhibition of the Arp-2/3 complex using the small molecule CK666. While migration was inhibited within an established two-dimensional migration system (**Fig. S9A-B)**, it was unaffected under confinement within liquid channels (**Fig. 6B**). This suggests a migration mode independent of Arp-2/3 mediated actin polymerization at the leading edge. Intriguingly, rapid imaging (1 sec intervals) revealed sheet-like pseudopodial protrusions in trailing cells and formation of rapid, rounded protrusions that mimic the shape and kinetics of blebs in pioneer cells (**Fig 6C, Fig S9C, Movie S4**). These bleb-like structures are disproportionately formed by pioneer cells interacting with the interface compared to trailing cells (**Fig S9D**). Inhibition of rear contractility under treatment with Rho-Kinase inhibitor Y-27632 hinders the ability of pioneer cells to migrate through the interfacial barrier (**Fig. 6D**). Under such treatment, pioneer cell speed is sufficiently slow such that trailing cell motility is limited by that of pioneers and thus were excluded from analysis. Attempts at confirming blebbing in primary neutrophils by visualization of F-actin was limited by photobleaching of fluorescent actin probes and resistance to transfection. Thus, we imaged migration of neutrophil-like PLB-985 cells expressing LifeAct-mRuby. Pioneer neutrophil-like cells exhibited prominent blebbing at the leading edge compared to trailing cells that favored pseudopodial-based protrusions (**Fig. 6E-F, Fig. S9E, Movie S5-6**). Furthermore, trailing cells that eventually reach the interface to become pioneers transition from utilizing pseudopodia to bleb-based protrusions (**Fig. 6G, Movie S7**).

323 Taken together, these data indicate a mechanism of bleb-based force generation at the leading
324 edge to deform an interface exerting cell-scale confining pressures.
325

Discussion

It is well established that cells sense physical properties of the surrounding tissue environment to regulate motility, oftentimes by sensing changes in cell membrane shape. This realization has predominantly relied on development of synthetic biomaterials in attempts to model physical properties of *in vivo* tissues. Existing systems to study confined cell migration largely rely on either ECM-based or rigid materials to provide confinement. The former offer limited ability to isolate variables of matrix stiffness without altering pore size and chemical composition, and the latter fail to mimic situations whereby a cell must generate forces to forge their own path through surrounding tissues (54). Here, we engineer a deformable liquid barrier lending authority to cells themselves in determining the degree of confinement and thus cell morphology. While cell-generated forces control the degree of confinement within a given channel geometry, resistance provided by the environment is tunable by controlling interfacial shape. By allowing such cellular autonomy and tuning environmental confining pressures, interfacial deformations match those of surrounding cells *in vivo*. To date, characterization of leukocyte interaction with surrounding cells *in vivo* is largely absent from existing literature due to limitations in optical access of mammalian models. Here, we validate the physiological relevance of confinement within this system by comparing deformations of the interface to those of surrounding cells during neutrophil interstitial motility within a larval zebrafish model. The ability to capture such interactions *in vivo* introduces an appealing model to study how cellular architecture and other physical properties of surrounding cells themselves regulate the immune response.

Given the temporal changes in tissue architecture that occur throughout aging and many disease states (i.e. tissue wounding, infection, malignancy) (55, 56), understanding how altered tissue mechanics regulates host immune cell motility holds particular relevance to human disease. Within liquid microchannels, pioneer cells that interact with the interfacial barrier require greater interfacial deformations to allow space for migration and as a result migrate slower. Increasing confining pressures by tuning channel geometries negatively impacts motility. These results would predict impaired leukocyte recruitment through tissues composed of stiffer cells, or more tightly packed cellular architectures. Future work might employ genetic engineering tactics to study how altered tissue mechanical properties impact leukocyte motility.

Seminal studies have demonstrated that purely mechanical cues can elicit transitions in cell migration mode. Srivastava et al., and Zatulovskiy et al. found that increasing applied transverse load ($-z$ direction) or stiffness of an agarose overlay induced a transition from pseudopodial to bleb-based migration in Dictyostelium cells (12, 57). In such studies, blebs were more often formed at areas of negative membrane curvature (57, 58). In addition, Goudarzi et al., found that blebs formed by zebrafish primordial germ cells occur at regions of local membrane invaginations which act as a source of membrane during bleb expansion (59). In liquid channels, both pioneer and trailing cells are under similar transverse confinement (z direction). Yet pioneers interact with the interface providing additional axial resistance (y direction). The ability of the interface to respond to cell-generated pressures results in distinct morphological polarity of pioneer cells. It is possible that alterations in membrane curvature induced by pioneer cell interaction with the interface drives transitioning to bleb-based motility. Cells use a number of mechanisms to sense membrane curvature, including BAR-domain proteins and PIEZO channels, both of which have been implicated in regulating bleb formation (57, 59). Our system represents an opportunity to decouple different chemical and mechanical cues (i.e. axial vs. transverse confining pressures) to further investigate the mechanisms by which cells sense their external environment to transition between migration modes.

In addition to applying physiologically-relevant confinement onto migrating cells, the ability to establish and control the shape of a liquid-liquid interface on single-cell length scales represents an intriguing potential to measure spatial and temporal variations in cell generated forces during the course of migration. Where traction force microscopy enables probing forces at the cell-substrate interface, observing the curvature of the liquid interface during cell migration

offers a simple readout of transverse forces (exerted in $+z$ direction). Distinct from
hydrogel-based systems where mechanical properties are complicated by measurement length
scale (60), equilibrium pressures of the liquid interface are independent of length scale. Using a
read out of interfacial curvature could provide a mathematically straightforward way to measure
cell-generated pressures in the transverse direction during motility that is independent of active
external intervention required in other methods (i.e. atomic force microscopy).
Thus far, advances in biomaterials engineering have largely been confined to modeling
cell interactions with the extracellular matrix. The ability to model physical interactions with
surrounding cells provokes an intriguing conceptual shift in the engineering of synthetic materials
to model *in vivo* tissues. From both a cell biology and technical engineering perspective, this work
advances our understanding of how leukocytes forge paths through dense cellular architectures,
while introducing a pioneering technology with implications for further development of soft
material-based systems to model an array of *in vivo* tissue properties. Further elucidation of how
immune cells sense and respond to physiologically relevant physical cues might facilitate
development of therapeutic approaches targeting immune cell migration machinery in the context
of human disease.
394
395
396
397 **Materials and Methods**
398
Construction of liquid-liquid microchannels
A chambered coverglass (no. 1.5 borosilicate glass, 0.13 to 0.17 mm thick; Thermo Fisher
Scientific, 155360) was treated first with O_2 plasma (Diener Electronic Femto, Plasma
Surface Technology) at 60 W for 3 min and then moved to an oven for grafting of
PDMS-Silane (1,3-dichlorotetramethylsiloxane; Gelest, SID3372.0; about 10 µl per device)
onto the glass surface by chemical vapor deposition (CVD) for 30 mins. The
PDMS-grafted slide was masked by a PDMS stamp (stamp construction described in *Li et*
al. (47)) and treated with O_2 plasma at 60 W for 3 min. After surface patterning, the PDMS
stamp was removed by tweezers and stored in a clean space for reuse. The chambered
coverglass slides were overlaid with oil (silicone oil, 5 cSt; Sigma-Aldrich, 317667) to
generate a surface exhibiting ELR to aqueous media. Liquid-liquid channels were created
by sweeping a hanging droplet across the patterned surface with a wide-orifice pipette.
Channels were constructed with collagen by sweeping a 1:1:1 solution of rat tail collagen I
(Corning, 10 mg/mL)) neutralized with one part neutrophil culture media (Roswell Park
Memorial Institute (RPMI) 1640 Medium, Thermo Fisher Scientific,11875093) + 2% fetal
bovine serum (FBS; Thermo Fisher Scientific, 10437010) and 1% Penicillin/Streptomycin))
and one part 2X HEPES buffer, yielding a final concentration 3.33 mg/mL of rat tail
collagen at pH 7.2. For experiments requiring visualization of the collagen layer, one in 15
parts RFP-labeled rat tail collagen was added to allow for confocal imaging of fiber
architecture. The collagen solution was incubated at 37 °C, 21% O2, 5% CO2, and 95%
RH for 15 minutes to allow for polymerization before transfer to an on-stage incubator
(Bold Line, Okolab) for imaging.
421
Whole blood collection and neutrophil isolation
All blood samples were drawn according to Institutional Review Boards (IRB) approved
protocols per the Declaration of Helsinki at the University of Wisconsin-Madison in the
Beebe Lab (IRB# 2020 − 1623) and in the Huttenlocher Lab (IRB#, 2017-003). Informed
consent was obtained from all subjects in the study. Whole blood was collected with
standard ethylenediaminetetraacetic acid (EDTA) tubes (BD, #36643, EDTA [K2]
STERILE, 1054688) and then stored at RT (∼ 22 °C) or 37 °C in stationary storage before
isolation.
Neutrophils were isolated from whole blood using magnetic bead-based negative
selection per protocol using either the EasySep Direct Human Neutrophil Isolation Kit
(STEMCELL, 19666) or the MACSxpress Neutrophil Isolation Kit (Miltenyi Biotech,

130-104-434) purified with the MACSxpress Erythrocyte Depletion Kit (Miltenyi Biotech,
130-098-196). After isolation, the neutrophil pellet was directly resuspended in RPMI 1640
culture media with 2% FBS and 1% Penicillin/Streptomycin (as above). Following
resuspension, liquid cultures were stored at 37° in a standard CO_2 incubator (Thermo
Scientific, HERACELL 240i) for no more than 1 hour before seeding into the device and
transfer to an onstage incubator for imaging.
439

Quantification of cell and channel heights following passive sweep
Primary neutrophils were placed directly into channels by incorporation into the neutralized
collagen mixture at concentration of 80k/µL before forming channels by sweep. Cells were
labeled immediately after resuspension by incubation at 37 °C and 5% CO_2 for 15 minutes
with with the cytoplasmic dye Calcein-AM green (1 µm; Invitrogen, C3100MP) and the
nuclear dye Hoescht 33342 (10µg/mL; Invitrogen, H1399) in RPMI 1640 media with 2%
FBS and 1% Penicillin/Streptomycin before combination with the neutralized collagen. The
collagen plus cell suspension was swept across the patterned surface as described above
to place cells directly into channels. Confocal imaging was performed on a Nikon AR1
spinning disk microscope courtesy of the UW Optical Core to gather z-stacks with step
size of 200 nm. Cell and channel heights, in addition to interfacial curvatures, were
measured from cross-sectional orthogonal views at regions where a cell was, or was not
present (both within the same channel). A single representative cell at the center of the
channel was chosen for analysis per unique channel replicate, with a total of three channel
replicates for each independent human donor (n = 3).
455

Quantification of cell morphology and motility during chemotaxis
To study cell chemotaxis under confinement by the liquid-liquid interface (Fig. 3), channels
were constructed by sweeping with neutralized collagen and incubated at 37° to allow
polymerization, as described above. Immediately after sweep, excess volume was
manually removed by pipette to ensure equal volumes in both inlet and outlet droplets.
Following isolation from whole blood, neutrophils were re-suspended in RPMI 1640 media
containing 2% FBS and 1% Penicillin/Streptomycin at a concentration of 40,000 cells per
µL. In experiments where interfacial height or curvature was measured from volumetric
confocal stacks (Fig. 3 and Fig. 4), media was visualized by addition of fluorescein (FITC)
sodium salt (40 µM; Sigma Aldrich, F6377). Vybrant DiD (Invitrogen, V22887) was used to
label the cell membrane; 5µL dye was added to 1mL of cell suspension in above media
immediately following isolation from whole blood (per the Invitrogen protocol) and
incubated at 37°C and 5% CO_2 for 15 minutes prior to washing and cell resuspension to
the desired concentration of 40k cells per µL for seeding into the liquid inlet droplets.
Following resuspension, 1.5 µL of cell suspension was added to the inlet droplet
by pipette. Immediately after, 1.5 µL of 100 nM *N*-Formylmethionyl-leucyl-phenylalanine
(fMLP; Millipore Sigma F3506) in the same media was added to the outlet droplet. After
such addition, the cell suspension was mixed by pipetting to ensure homogeneity and cell
positioning near the channel entrance. Addition of equal amounts of volume to both the
inlet and outlet droplets is essential to restrict generation of convective flow throughout the
channel, so that chemokine gradients are driven by diffusion alone. Cell motility was
visualized on a Nikon epi-fluorescent microscope (Nikon Eclipse TE3000 with a Nikon
Plan Fluor 20x/0.50 objective, motorized stage from Ludl Electronic Products and Prime
BSI Express camera from Teledyne Photometrics or a Nikon TI Eclipse inverted
microscope with Nikon Plan Fluor 20x/0.50 objective and Hamamatsu camera).
Quantification of cell speed was performed in ImageJ/FIJI using the manual tracking
plug-in of time lapse movies taken at time intervals of 30 seconds. Volumetric imaging of
cells interacting with the interface was performed by labeling the media with FITC dye and
imaging with the aforementioned Nikon AR1 spinning disk confocal microscope with step
size of 200 nm. Through such volumetric imaging, interfacial heights were measured by
the height of the FITC labeled media layer. Similarly, measurements of interfacial
pressures during pioneer cell migration (Fig. 4) were made from measuring interfacial

curvature (defined by the FITC labeled media) from cross-sectional (xz plane) and side (yz
plane) volumetric images.
490
Zebrafish maintenance and handling
Animal care and use was approved by the Institutional Animal Care and Use Committee of
University of Wisconsin and strictly followed guidelines set by the federal Health Research
Extension Act and the Public Health Service Policy on the Humane Care and Use of
Laboratory Animal, administered by the National Institutes of Health Office of Laboratory
Animal Welfare. All protocols using zebrafish in this study were approved by the University
of Wisconsin-Madison Research Animals Resource Center (protocol M005405-A02). To
generate the larval zebrafish model used to visualize neutrophil morphology and
keratinocyte deformations, previously published neutrophil-cytoplasm labeled
Tg(LyzC:TagBFP) fish were crossed with nuclear labeled *Tg(LyzC:H2B-mCherry)* fish to
generate a stable line (*Tg(LyzC:TagBFP/LyzC:H2B-mCherry).* These fish were bred with a
stable line expressing cytoplasmic acGFP in basal keratinocytes *Tg(Krtt1c19e:acGFP)*)
(Gift of Dr. Alvaro Sagasti) to generate larvae used in each experiment. All fish were
maintained on the AB background. Following breeding, fertilized embryos were transferred
to E3 medium (5 mM NaCl, 0.17 mM KCl, 0.44 mM $CaCl_2$, 0.33 mM $MgSO_4$, 0.025 mM
NaOH, and 0.0003% Methylene Blue) and maintained at 29°C. Larval zebrafish were
anesthetized using 0.2 mg/ml tricaine (ethyl 3-aminobenzoate; Sigma-Aldrich) before any
experimentation or live imaging.
509
Tracking *in vivo* keratinocyte deformations
Dechorionated, tricaine-anesthetized three days-post fertilization (dpf) larvae were
mounted in 2% low-melting point agarose (Sigma-Aldrich) on a 35 mm glass bottom dish
(#1.5H Glass, Cellvis, CA, USA). Time-lapse imaging of spontaneous neutrophil migration
through basal keratinocytes in the yolk sac and neck region was performed on an
spinning-disk confocal (CSU-X; Yokogawa) on a Zeiss Observer Z.1 inverted microscope
and an electron-multiplying charge-coupled device Evolve 512 camera (Photometrics),
with a EC Plan-Neofluar 40×/NA 0.75 air objective (1-2μm optical sections, 2,355 × 512
resolution) using ZenPro 2012 software (Zeiss). Images were captured at 10 second
intervals with z-step intervals of 3 μm. Quantification of deformations was performed on
resulting maximal intensity projections; background subtraction was manually performed in
FIJI/ImageJ in images displayed in the main text (Fig. 5).
522
Quantification of bleb formation in primary neutrophils and neutrophil-like PLB-985
cells
Both primary neutrophils and PLB-985 neutrophil-like cells were used to determine the
migration mode utilized by neutrophils during interaction with the confining interface (Fig.
6). Immediately following isolation from whole blood, primary neutrophils were incubated
with 10 μM CK666 (Millipore Sigma, SML0006) or LY-27632 (Millipore Sigma, Y0503) to
inhibit CK-666 or Rho-Kinase at 37 °C and 5% CO_2 for 30 mins. Bleb-like structures in
primary neutrophils were quantified from brightfield time lapse images taken at 1 sec
intervals.
PLB-985 cells expressing LifeAct-mRuby were generated by lentiviral transfection.
HEK293T cells were grown to 70% confluency in a 10-cm tissue culture dish for each
lentiviral target and transfected with pLV-EF1a-IRES-Neo (Addgene #85139) containing
LifeAct-mRuby, VSV-G, and CMV 8.9.1. The 72-h viral supernatant was collected and
concentrated using a lentivirus concentrator (Lenti-X; Takara Bio Inc.) following the
manufacturer's instructions. PLB-985 cells (1 × 10^6 total number) were infected with viral
supernatant for 3 days in the presence of 15 μg/ml polybrene. Stable cell lines were
generated with 1 μg/ml neomycin selection. Resulting PLB-985 cells expressing
LifeAct-mRuby were differentiated into neutrophil-like cells by treatment with 1.25% DMSO
(Sigma-Aldrich, D2650) for 6 days at 37 °C and 5% CO_2 in RPMI 1640 with 2% FBS and
1% Penicillin/Streptomycin. Cell motility was assayed in an equivalent assay as that of

primary cells, with imaging performed on the previously described Zeiss Observer Z.1 inverted spinning disc confocal microscope, at 0.5-1 sec intervals. Background subtraction was manually performed in FIJI/ImageJ in images displayed in the main text (Fig. 6).

Statistical Analysis
Data were analyzed (Prism 9.0; GraphPad Software). Statistical significance was assessed using Student's *t* tests when comparing two conditions/groups, and one-way analysis of variance (ANOVA) corrected using the Tukey's test when comparing multiple groups.

Funding

Biotechnology Training Program
University of Wisconsin-Madison
T23GM135066
PI: Brian Fox

Ruth L. Kirschstein National Research Service Award (NRSA) Fellowship
National Heart, Lung, and Blood Institute
1F30HL174128
PI: Jon Schrope

Carbone Cancer Center Support Grant
University of Wisconsin
NIH P30CA014520

Microscale models of inflammation and its resolution
NIH R01AI34749
PI: Beebe and Huttenlocher

Cell Migration and Wound Repair
NIH R35GM118027-08
PI: Huttenlocher

A multiplexed micro scale assay for real time analysis of pediatric immune cell function
NIH U24AI152177
PI: Beebe

Under-oil open microfluidic system (UOMS) for studying systemic fungal infection
NIH R01AI54940
PI: Beebe

Competing interests: David J. Beebe holds equity in Bellbrook Labs LLC, Tasso Inc., Salus Discovery LLC, Lynx Biosciences Inc., Stacks to the Future LLC, Flambeau Diagnostics LLC, and Onexio Biosystems LLC.

Data and materials availability: All data are available in the main text or the supplementary materials, or upon request.

Acknowledgments
The authors would like to thank the University of Wisconsin Imaging Core for the use of microscopy facilities as well as Biorender and Imaris for use of software to make figure schematics.

References
1. A. D. Luster, R. Alon, U. H. von Andrian, Immune cell migration in inflammation: present and future therapeutic targets. *Nat Immunol* **6**, 1182–1190 (2005).
2. S. de Oliveira, E. E. Rosowski, A. Huttenlocher, Neutrophil migration in infection and wound repair: going forward in reverse. *Nat Rev Immunol* **16**, 378–391 (2016).
3. H. D. Moreau, M. Piel, R. Voituriez, A.-M. Lennon-Duménil, Integrating Physical and Molecular Insights on Immune Cell Migration. *Trends in Immunology* **39**, 632–643 (2018).
4. P. Friedl, B. Weigelin, Interstitial leukocyte migration and immune function. *Nat Immunol* **9**, 960–969 (2008).
5. M. L. Heuzé, *et al.*, Migration of dendritic cells: physical principles, molecular mechanisms, and functional implications. *Immunol Rev* **256**, 240–254 (2013).
6. S. van Helvert, C. Storm, P. Friedl, Mechanoreciprocity in cell migration. *Nat Cell Biol* **20**, 8–20 (2018).
7. W. Weninger, M. Biro, R. Jain, Leukocyte migration in the interstitial space of non-lymphoid organs. *Nat Rev Immunol* **14**, 232–246 (2014).
8. E. Sitarska, *et al.*, Sensing their plasma membrane curvature allows migrating cells to circumvent obstacles. *Nat Commun* **14**, 5644 (2023).
9. L. Boneschansker, J. Yan, E. Wong, D. M. Briscoe, D. Irimia, Microfluidic platform for the quantitative analysis of leukocyte migration signatures. *Nat Commun* **5**, 4787 (2014).
10. H. V. Prentice-Mott, *et al.*, Biased migration of confined neutrophil-like cells in asymmetric hydraulic environments. *Proc Natl Acad Sci U S A* **110**, 21006–21011 (2013).
11. A. E. Ekpenyong, *et al.*, Mechanical deformation induces depolarization of neutrophils. *Science Advances* **3**, e1602536 (2017).
12. E. Zatulovskiy, R. Tyson, T. Bretschneider, R. R. Kay, Bleb-driven chemotaxis of Dictyostelium cells. *Journal of Cell Biology* **204**, 1027–1044 (2014).
13. L. K. Fritz-Laylin, *et al.*, Actin-based protrusions of migrating neutrophils are intrinsically lamellar and facilitate direction changes. *eLife* **6**, e26990.
14. R. H. Insall, L. M. Machesky, Actin Dynamics at the Leading Edge: From Simple Machinery to Complex Networks. *Developmental Cell* **17**, 310–322 (2009).
15. T. Lämmermann, R. N. Germain, The multiple faces of leukocyte interstitial migration. *Semin Immunopathol* **36**, 227–251 (2014).
16. E. K. Paluch, E. Raz, The role and regulation of blebs in cell migration. *Curr Opin Cell Biol* **25**, 582–590 (2013).
17. J. Schick, E. Raz, Blebs—Formation, Regulation, Positioning, and Role in Amoeboid Cell Migration. *Front. Cell Dev. Biol.* **10** (2022).
18. B. R. Graziano, *et al.*, Cell confinement reveals a branched-actin independent circuit for neutrophil polarity. *PLoS Biology* **17** (2019).
19. C. D. Paul, W.-C. Hung, D. Wirtz, K. Konstantopoulos, Engineered Models of Confined Cell Migration. *Annu Rev Biomed Eng* **18**, 159–180 (2016).
20. C. F. Guimarães, L. Gasperini, A. P. Marques, R. L. Reis, The stiffness of living tissues and its implications for tissue engineering. *Nature Reviews Materials* **5**, 351–370 (2020).
21. M.-E. Brett, A. L. Crampton, D. K. Wood, Rapid generation of collagen-based microtissues to study cell–matrix interactions. *Technology* **04**, 80–87 (2016).
22. K. Siemsen, *et al.*, Tunable 3D Hydrogel Microchannel Networks to Study Confined Mammalian Cell Migration. *Adv Healthc Mater* **10**, e2100625 (2021).

23. X. Li, Q. Sun, Q. Li, N. Kawazoe, G. Chen, Functional Hydrogels With Tunable
Structures and Properties for Tissue Engineering Applications. *Frontiers in*
Chemistry **6** (2018).
24. A. M. Kloxin, M. W. Tibbitt, A. M. Kasko, J. A. Fairbairn, K. S. Anseth, Tunable
Hydrogels for External Manipulation of Cellular Microenvironments through
Controlled Photodegradation. *Adv Mater* **22**, 61–66 (2010).
25. C. M. Kraning-Rush, S. P. Carey, M. C. Lampi, C. A. Reinhart-King, Microfabricated
collagen tracks facilitate single cell metastatic invasion in 3D. *Integr Biol (Camb)* **5**,
606–616 (2013).
26. J. A. Mosier, *et al.*, Extent of Cell Confinement in Microtracks Affects Speed and
Results in Differential Matrix Strains. *Biophys J* **117**, 1692–1701 (2019).
27. M. R. Zanotelli, *et al.*, Energetic costs regulated by cell mechanics and confinement
are predictive of migration path during decision-making. *Nat Commun* **10**, 4185
(2019).
28. S.-Y. Cheng, *et al.*, A hydrogel-based microfluidic device for the studies of directed
cell migration. *Lab Chip* **7**, 763–769 (2007).
29. J. François, *et al.*, The interplay between matrix deformation and the coordination of
turning events governs directed neutrophil migration in 3D matrices. *Sci Adv* **7**,
eabf3882 (2021).
30. M. Ehrbar, *et al.*, Elucidating the Role of Matrix Stiffness in 3D Cell Migration and
Remodeling. *Biophys J* **100**, 284–293 (2011).
31. C. A. Reinhart-King, M. Dembo, D. A. Hammer, Cell-Cell Mechanical
Communication through Compliant Substrates. *Biophysical Journal* **95**, 6044–6051
(2008).
32. N. Wang, E. Ostuni, G. M. Whitesides, D. E. Ingber, Micropatterning tractional forces
in living cells. *Cell Motil Cytoskeleton* **52**, 97–106 (2002).
33. A. Afthinos, *et al.*, Migration and 3D Traction Force Measurements inside Compliant
Microchannels. *Nano Lett.* **22**, 7318–7327 (2022).
34. F. Wang, *et al.*, Lipid products of PI(3)Ks maintain persistent cell polarity and
directed motility in neutrophils. *Nature Cell Biology* **4**, 513–518 (2002).
35. T. Y.-C. Tsai, *et al.*, Efficient Front-Rear Coupling in Neutrophil Chemotaxis by
Dynamic Myosin II Localization. *Dev Cell* **49**, 189-205.e6 (2019).
36. J. Renkawitz, *et al.*, Nuclear positioning facilitates amoeboid migration along the
path of least resistance. *Nature* **568**, 546–550 (2019).
37. V. Ambravaneswaran, I. Y. Wong, A. J. Aranyosi, M. Toner, D. Irimia, Directional
decisions during neutrophil chemotaxis inside bifurcating channels. *Integr Biol*
(Camb) **2**, 639–647 (2010).
38. D. C. Williams, D. K. Wood, High-throughput quantification of red blood cell
deformability and oxygen saturation to probe mechanisms of sickle cell disease.
Proceedings of the National Academy of Sciences **120**, e2313755120 (2023).
39. K. M. Stroka, *et al.*, Water Permeation Drives Tumor Cell Migration in Confined
Microenvironments. *Cell* **157**, 611–623 (2014).
40. B. Hamza, *et al.*, Retrotaxis of human neutrophils during mechanical confinement
inside microfluidic channels. *Integr Biol (Camb)* **6**, 175–183 (2014).
41. C. Deroy, *et al.*, Predicting flows through microfluidic circuits with fluid walls.
Microsyst Nanoeng **7**, 1–9 (2021).
42. C. Soitu, *et al.*, Microfluidic chambers using fluid walls for cell biology. *Proceedings*
of the National Academy of Sciences **115**, E5926–E5933 (2018).
43. C. Deroy, *et al.*, Assaying Macrophage Chemotaxis Using Fluid-Walled Microfluidics.
Advanced Materials Technologies **7**, 2200279 (2022).
44. W. Feng, *et al.*, Harnessing liquid-in-liquid printing and micropatterned substrates to
fabricate 3-dimensional all-liquid fluidic devices. *Nat Commun* **10**, 1095 (2019).
45. C. Li, *et al.*, Double-exclusive liquid repellency (double-ELR): an enabling technology
for rare phenotype analysis. *Lab Chip* **18**, 2710–2719 (2018).
46. C. Li, *et al.*, Exclusive Liquid Repellency: An Open Multi-Liquid-Phase Technology

for Rare Cell Culture and Single-Cell Processing. *ACS Appl. Mater. Interfaces* **10**,
17065–17070 (2018).
47. C. Li, *et al.*, Under oil open-channel microfluidics empowered by exclusive liquid
repellency. *Science Advances* **6**, eaay9919 (2020).
48. Z. Zhou, F. Cui, Q. Wen, H. Susan Zhou, Effect of vimentin on cell migration in
collagen-coated microchannels: A mimetic physiological confined environment.
Biomicrofluidics **15**, 034105 (2021).
49. B. Burkel, *et al.*, Preparation of 3D Collagen Gels and Microchannels for the Study of
3D Interactions In Vivo. *J Vis Exp* 53989 (2016). https://doi.org/10.3791/53989.
50. H. Du, *et al.*, Tuning immunity through tissue mechanotransduction. *Nat Rev*
Immunol **23**, 174–188 (2023).
51. J. Geng, *et al.*, TLR4 signalling via Piezo1 engages and enhances the macrophage
mediated host response during bacterial infection. *Nat Commun* **12**, 3519 (2021).
52. V. S. Meli, *et al.*, YAP-mediated mechanotransduction tunes the macrophage
inflammatory response. *Sci Adv* **6**, eabb8471 (2020).
53. R. G. Scheraga, B. D. Southern, L. M. Grove, M. A. Olman, The Role of TRPV4 in
Regulating Innate Immune Cell Function in Lung Inflammation. *Front Immunol* **11**,
1211 (2020).
54. G. Charras, E. Sahai, Physical influences of the extracellular environment on cell
migration. *Nat Rev Mol Cell Biol* **15**, 813–824 (2014).
55. M. C. Lampi, C. A. Reinhart-King, Targeting extracellular matrix stiffness to attenuate
disease: From molecular mechanisms to clinical trials. *Science Translational*
Medicine **10**, eaao0475 (2018).
56. J. Huynh, *et al.*, Age-Related Intimal Stiffening Enhances Endothelial Permeability
and Leukocyte Transmigration. *Sci Transl Med* **3**, 112ra122 (2011).
57. N. Srivastava, D. Traynor, M. Piel, A. J. Kabla, R. R. Kay, Pressure sensing through
Piezo channels controls whether cells migrate with blebs or pseudopods.
Proceedings of the National Academy of Sciences **117**, 2506–2512 (2020).
58. R. A. Tyson, E. Zatulovskiy, R. R. Kay, T. Bretschneider, How blebs and pseudopods
cooperate during chemotaxis. *Proceedings of the National Academy of Sciences*
111, 11703–11708 (2014).
59. M. Goudarzi, *et al.*, Bleb Expansion in Migrating Cells Depends on Supply of
Membrane from Cell Surface Invaginations. *Dev Cell* **43**, 577-587.e5 (2017).
60. D. Lee, H. Zhang, S. Ryu, "Elastic Modulus Measurement of Hydrogels" in
Cellulose-Based Superabsorbent Hydrogels, Md. I. H. Mondal, Ed. (Springer
International Publishing, 2018), pp. 1–21.

 Figures and Tables

Figure 1. ELR-enabled construction of liquid microchannels on single cell length scales.
A) Schematic depicting Young's contact angle which describes the degree of liquid repulsion by a solid surface in a three phase (Solid-Liquid-Liquid) system. Controlling surface chemistry enables generation of surfaces exhibiting exclusive liquid repellency (ELR), characterized by complete liquid repulsion yielding a contact angle of 180°. Such surfaces can be Aqueous Repellent or Oil Repellent Surfaces (ARS, ORS respectively). **B)** Schematic depicting differential surface patterning of a glass substrate to exhibit ARS and ORS regions to create a double-ELR system. This enables creation of liquid channels with the maximum possible interfacial forces pinning both aqueous liquid and oil onto their preferred surface, enabling channel heights < 10 µm. **C)** Schematic of channel construction by "sweeping" a hanging droplet of aqueous media across ORS regions to form two hemispheric droplets and a connecting channel (> 50 on standard microscope slide). **D)** Macroscale side view of a media channel after sweep and addition of 1 µL of culture media to each droplet to improve visualization. **E)** Confocal image of a culture media channel with height < 1 µm (cross-sectional scale bars represent 3 µm). The channel height tapers at the entrance to a relatively consistent height (points III-IV) until a gradual, symmetric height increase towards the channel exit (right side). **F)** Schematic depicting incorporation of fibrous collagen into microchannels by performing the sweep with neutralized collagen then allowing polymerization. **G)** Representative confocal image of collagen-coated channel (cross-sectional scale bars represent 3 µm).

A placing cells directly into channel

cells in collagen hanging droplet

sweep → sweep →

inlet droplet | outlet droplet

B varying channel widths

cs (xz)

calcein

top (xy)

hoechst

30 μm | 100 μm | 150 μm | 200 μm

C

- channel alone
- cell

chan > cell
.0417

chan < cell
< .0001 | .0230 | .8489

height (μm)

channel width (μm)

D side (yz)

$h_{0,equilib}$ | $h_{cell,equilib}$

$\kappa_{yz,\,equilib}$

E

$\kappa_{yz,equilib}$ | $\kappa_{xz,equilib}$

cross-section (xz)

$P_{equilib,interface}$

$\kappa_{xz,\,cell}$

$\kappa_{xz,\,no\,cell}$

$$P_{equilib,interface} = \gamma_{OM}\left(\frac{1}{R_{xz,local}} + \frac{1}{R_{yz,local}}\right)$$

F channel cross-section (xz)

$P_{equilib,interface,\,xz}$

h

w_0 | Δw | $\kappa_{xz,\,w_0+\Delta w}$

$R_{xz,\,w_0}$ | $\kappa_{xz,\,w_0}$

$R_{xz,\,w_0+\Delta w}$

G

- cell
- no cell

equilibrium interfacial pressure (Pa)

channel width (μm)

H

0.0460

mean interfacial height (μm)

nucleus | cytoplasm

I side (yz)

oil

media

$P_{equilib,interface}$

$P_{0,interface}$

P_{media}

$\kappa_{xz,local}$

nucleus

cytoplasm

$h_{max,\,cell} = h_{max,interface}$

Figure 2. Tension of liquid-liquid interface exerts confining pressures on the scale of single cell stiffness A) Schematic depicting direct addition of cells embedded in neutralized collagen into liquid microchannels by incorporation into the hanging droplet during the sweep process. **B)** Primary human neutrophils placed within channels of varying sizes (30-200 μm widths) exhibit differential degrees of cell confinement. **C)** Quantification of cell height as a function of channel width. Each point represents the mean cell (blue) or initial channel (red) height over three channel replicates for three different human donors. P-values represent differences between cell and channel height (blue vs red) determined by t-test. In channels of sufficiently low height (30 and 100 μm width), cell height exceeds that of channel height; on 200 μm width channels of height greater than cell diameter, cells exhibit a resting diameter less than that of channel height. **D)** Channel side view revealing that cells are confined by, yet deform the channel interface from $h_{0,equilib}$ to $h_{cell,equilib}$ to generate curvature (κ) in yz the plane. **E)** Confocal surface rendering depicting interfacial curvatures in the xz and yz planes resulting from cell-generated deformation. Central nuclear positioning in such immobile cells results in equivalent curvature in the xz and yz planes. **F)** Schematic defining the channel interfacial pressure ($P_{equilib,interface}$), curvature (κ), and radius of curvature (R), in the xz plane. Increases in channel width alter interfacial pressure. **G)** Quantification of interfacial pressure ($P_{equilib,interface}$) in the presence or absence of a cell as a function of channel width. Each point represents the mean of three independent channel replicates. **H)** The interfacial height is greatest over regions of the cell containing the nucleus compared to those containing only cytoplasm. Each point represents a cell measured on an independent channel replicate (n = 3), shade represents donor (n = 3). Statistical significance was determined by an unpaired, two-sample t-test assuming equal variance. **I)** Model schematic depicting equilibrium interfacial pressure defined by cell physical properties. In locations devoid of a cell, $P_{equilib,interface}$ reflects the pressure exerted by the media itself. In the presence of a cell, $P_{equilib,interface}$ is determined by the pressure exerted by the cell nucleus.

Figure 3. Migration speed is dependent on required deformation of interface. A) Schematic of migration experiment whereby collagen channels are created by the sweep technique followed by sequential addition of cells in the inlet droplet across from chemoattractant in the outlet droplet. **B)** Representative (confocal) side view of channel entrance. Cells are confined to the media layer between the collagen coating and oil-media interface. **C)** Timelapse images and **D)** corresponding quantification of migration speed for pioneer and trailing cells (n = 5 cells depicted by circles over 3 channel replicates and 3 independent human donors denoted by shade). **E)** Confocal image of pioneer cells deforming the interface (dashed white line). Media was visualized by addition of FITC dye. **F)** Quantification of interfacial heights before and after passage of pioneer cells. Each point represents an independent channel replicate (n = 10). **G)** Model schematic and corresponding quantification of migration speed for neutrophils that start as trailing cells but catch up to and become pioneer cells over time. Each data point represents the mean speed of a cell before and after the transition to becoming a pioneer (n = 3 cells observed on independent channel replicates over n = 3 independent human donors, represented by shade). Statistical significance was determined by either an unpaired (D,E) or paired (G) two-sample t-test assuming equal variance. All data gathered on 30 μm width channels.

Figure 4. Liquid-liquid interface exerts gradient of confining pressure on migratory cells exhibiting morphological polarity A) Representative side view of neutrophil migration within a 100 µm width channel exhibiting rear-ward nuclear positioning and a front-ward protrusion. This results in differential interfacial deformation and thus equilibrium confining pressure across the length of the cell (yz plane). **B)** Three-dimensional confocal reconstruction depicting differential interfacial curvature across cell length. **C)** Quantification of equilibrium confining pressure over the cell rear and front. Statistical significance determined by an unpaired, two-sample t-test assuming equal variance. **D)** Representative confining pressure gradients over the length of pioneer cells within channels of differential width (30 µm vs 100 µm). **E)** Cell speed as a function of channel width. Each point represents a single cell, data pooled over three independent channels and two independent donors (denoted by shade). Statistical significance determined by a one-way ANOVA with multiple comparisons between every group.

817 Figure 5. Tuning interfacial geometry recapitulates surrounding cell deformations during
818 neutrophil migration *in vivo*. A) Schematic of skin layers in larval zebrafish three days
post-fertilization. At this developmental stage, the epidermis consists of a single layer of both
superficial (tan) and basal (green) keratinocytes. **B)** Schematic and corresponding time-lapse
images depicting distinct phases of interstitial migration *in vivo* whereby neutrophils extend
protrusions to deform surrounding basal keratinocytes (green) followed by passage of the cell
body resulting in maximal deformation of surrounding cells. **C)** Schematic depicting interfacial
deformation during protrusion extension and passage of the cell body *in vitro*. **D-E)** Tuning
interfacial geometries recapitulates neutrophil protrusion width (D) and maximal surrounding cell
deformations induced by the neutrophil cell body (E). Circles represent single cell passage
events, shade represents independent biological replicates (n = 3), either a different fish clutch or
human donor. Statistical significance was determined by a one-way ANOVA assuming equal
variances with multiple comparisons to the *in vivo* group.

Figure 6. Neutrophils employ a bleb-based mechanism of force generation to deform the confining interface. A) Schematic of leukocyte migration modes, including actin polymerization-dependent pseudopod formation and bleb formation through rear contractility driving membrane extension. **B)** Treatment with Arp2/3 inhibitor CK666 does not impact migration speed within liquid channels. Circles represent mean track speed of individual cells pooled over three technical channel replicates; shade represents individual donor replicates (n = 3). **C)** Image of bleb-like protrusions formed by primary neutrophils compared to pseudopodia of trailing cells; scale bar represents 10 µm. **D)** Inhibition of rear contractility with Rho Kinase inhibitor Y-27632 decreases pioneer migration speed in primary neutrophils. White arrow denotes migration direction. Circles represent cell speeds pooled over three channel replicates; shade represents human donor replicate. **E)** Timelapse images of neutrophil-like PLB-985 cells expressing LifeAct-mRuby depicting bleb formation within pioneer cells and pseudopodia characteristic of trailing cells. White arrow denotes migration direction. Scale bar represents 5 µm. **F)** Quantification of blebs per minute for pioneer and trailing neutrophil-like PLB-985 cells. Each point represents a cell, data pooled over three independent biological replicates represented by shade. **G)** Trailing cells that reach the interface transition from pseudopodia to bleb-based protrusions (denoted by white arrows); scale bar represents 10 µm.