A transcriptomics-based drug repositioning approach to identify drugs with similar

activities for the treatment of muscle pathologies in spinal muscular atrophy (SMA) models

Joseph M Hoolachan [1], Eve McCallion [1], Emma R Sutton [1], Özge Çetin [1], Paloma Pacheco-Torres

[2], Maria Dimitriadi [2], Magnus Okoh [1], Lisa M Walter [3, 4], Peter Claus [3, 4], Matthew JA Wood [5],

Daniel P Tonge [6], & Melissa Bowerman [1, 7].

[1] School of Medicine, Keele University, Staffordshire, UK.

[2] School of Life and Medical Sciences, University of Hertfordshire, Hertfordshire, UK.

[3] SMATHERIA gGmbH – Non-Profit Biomedical Research Institute, Hannover, Germany.

[4] Centre of Systems Neuroscience (ZSN), Hannover Medical School, Hannover, Germany.

[5] Department of Paediatrics, University of Oxford, Oxford, UK.

[6] School of Life Sciences, Keele University, UK.

[7] Wolfson Centre for Inherited Neuromuscular Disease, RJAH Orthopaedic Hospital, Oswestry,

UK.

Corresponding Author: m.bowerman@keele.ac.uk

ABSTRACT

Spinal muscular atrophy (SMA) is a genetic neuromuscular disorder caused by the reduction of

survival of motor neuron (SMN) protein levels. Although three SMN-augmentation therapies are

clinically approved that significantly slow down disease progression, they are unfortunately not

cures. Thus, complementary SMN-independent therapies that can target key SMA pathologies and

that can support the clinically approved SMN-dependent drugs are the forefront of therapeutic

development. We have previously demonstrated that prednisolone, a synthetic glucocorticoid (GC)

improved muscle health and survival in severe $Smn^{-/-};SMN2$ and intermediate $Smn^{2B/-}$ SMA mice.

However, long-term administration of prednisolone can promote myopathy. We thus wanted to

identify genes and pathways targeted by prednisolone in skeletal muscle to discover clinically

approved drugs that are predicted to emulate prednisolone's activities. Using an RNA-sequencing,

bioinformatics and drug repositioning pipeline on skeletal muscle from symptomatic prednisolone-

treated and untreated $Smn^{-/-};SMN2$ SMA and $Smn^{+/-};SMN2$ healthy mice, we identified molecular

targets linked to prednisolone's ameliorative effects and a list of 580 drug candidates with similar

predicted activities. Two of these candidates, metformin and oxandrolone, were further

investigated in SMA cellular and animal models, which highlighted that these compounds do not

have the same ameliorative effects on SMA phenotypes as prednisolone; however, a number of

other important drug targets remain. Overall, our work further supports the usefulness of

prednisolone's potential as a second-generation therapy for SMA, identifies a list of potential SMA

drug treatments and highlights improvements for future transcriptomic-based drug repositioning

studies in SMA.

INTRODUCTION

Spinal muscular atrophy (SMA) is a heterogenous autosomal recessive neuromuscular disorder (NMD) characterized by motor neuron degeneration alongside progressive muscle atrophy and weakness [1]. Being the leading monogenic cause of infant mortality [2], around 96% of SMA cases are mapped to homozygous loss-of-function and deletion mutations in the *survival of motor neuron 1* (*SMN1*) gene [3,4], which ubiquitously expresses SMN, a protein that current and ongoing research has linked to diverse housekeeping and tissue-specific cellular functions [5–7]. Although complete SMN loss is embryonic lethal in most organisms[8], humans can overcome the complete loss of the *SMN1* gene due to incomplete rescue by the homologous *SMN2* gene [9,10]. In essence, the presence of a single nucleotide mutation in *SMN2* promotes exon 7 alternative splicing that limits full length SMN (FL-SMN) expression in this gene to 10% [11]. Consequently, the limited FL-SMN expression makes *SMN2* gene copy number an important disease modifier, impacting SMA type and severity [12].

In recent years, novel SMN restorative SMA treatments have emerged that either increase FL-*SMN2* expression by an anti-sense oligonucleotide (ASO) (Nusinersen marketed as (Spinraza®)) [13,14] or a small molecule (Evrysdi®) [15,16] or promote exogenous FL-*SMN1* expression by an adeno-associated virus 9 (AAV-9) delivery system (Zolgensma®) [17,18]. Despite the significant increased life expectancy and improved quality of life associated with these therapies [14,16,18,19], they are not cures and their efficacy is dependent upon early intervention [20].

Thus, additional SMN-independent therapies that target affected tissues such as muscle are needed to further enhance and support the benefits of SMN-dependent treatments [21]. Indeed, pre-clinical studies and primary patient data have reported innate muscular defects in SMA, which include myogenesis [22], regeneration [23], contraction [24,25], regulation [25], growth [26], and metabolism [27],

highlighting skeletal muscle as a primary therapeutic target. Although two novel skeletal muscle-

specific SMA therapies, Apitegromab[TM] [28] (ClinicalTrials.gov ID: NCT03921528) and

Reldesemtiv[TM] [29] (ClinicalTrials.gov ID: NCT02644668), are showing progress in clinical trials,

the high expenses involved in novel drug research and development (R&D) [30] alongside the costs

of the three clinically approved SMN-dependent therapies may lead to elevated prices for

combinatorial treatments [31], thus further widening the accessibility gap for SMA patients.

A useful alternative for ensuring accessibility of SMN-independent treatments for all SMA

patients would be the identification of cost-effective generic drugs via drug repositioning, a

strategy aimed at finding new therapeutic activities for existing pharmacological compounds [32].

One such example is prednisolone, a synthetic glucocorticoid (GC) administered to relieve muscle

inflammation in Duchenne muscular dystrophy (DMD) patients [33,34]. Interestingly, evidence also

emerged that short-term prednisolone treatment additionally conferred ergogenic muscle benefits

in DMD patients [35,36], which was also validated in the *mdx* mouse model of DMD [37,38].

Although GCs are not used for the treatment of SMA patients, prednisolone is administered for a

short period (~30 days, 1 mg/kg) to alleviate the immunological adverse effects of Zolgensma®

[18]. However, given prednisolone's potential muscle benefits, we have previously investigated and

demonstrated that prednisolone treatment (5 mg/kg, every other day) in severe $Smn^{-/-};SMN2$ and

intermediate $Smn^{2B/-}$ SMA mice improved survival, weight and neuromuscular phenotype [27].

Although the study was aimed at investigating prednisolone's activity on the GC-Krüppel-Like-

Factor-15 (KLF15) pathway in SMA skeletal muscle, synergistic muscle improvement was also

observed in prednisolone-treated $Smn^{-/-};SMN2$ SMA mice overexpressing Klf15 specifically in

skeletal muscle [27], suggesting that prednisolone may act on SMA skeletal muscle via numerous

effectors and pathways.

Despite the benefits observed, our study did not evaluate prednisolone's long-term effects [18,27].

Given that chronic GC usage increases myopathy [35,39], it is unclear whether long-term prednisolone

treatments would be similarly detrimental in SMA muscle. Furthermore, the rapid onset and

progression of disease in SMA mouse models (1-3 weeks on average) [9,40,41] does not allow

sufficient comparison of intermittent vs chronic studies.

Thus, in this study, we used a transcriptomics and drug repositioning pipeline based on our

previously published experimental paradigm [42] to uncover the genes and pathways restored by

prednisolone in skeletal muscle of SMA mice and identify existing non-GC drugs predicted to

have similar activities. Our study uncovered that prednisolone restored pathways linked to growth,

metabolism, and regulation in SMA skeletal muscle and identified 20 leading commercially

available non-GC drugs predicted to emulate its action. Based on oral bioavailability and evidence

of safety treatment in children, we selected and validated metformin and oxandrolone in SMA

cellular and animal models. Although, both metformin and oxandrolone improved neuromuscular

activity in the *Caenorhabditis elegans (C. elegans)* model for severe SMA, we found that higher

metformin doses reduced survival in the intermediate $Smn^{2B/-}$ SMA mouse model. On the other

hand, oxandrolone treatment partially improved survival in $Smn^{2B/-}$ SMA mice, albeit not to the

same extent as prednisolone [27].

Nevertheless, our study computationally uncovered new mechanisms behind prednisolone's

beneficial activity in SMA muscle, identified numerous potential SMA muscle-specific

therapeutic candidates and highlighted the importance of transcriptomic-based drug repositioning

for SMN-independent drug discovery.

METHODS

Animal Procedures

Experiments with the severe (Taiwanese) $Smn^{-/-};SMN2$ (FVB/N background, FVB.Cg-$Smn1tm1HungTg(SMN2)2Hung/J$) SMA mice [40] and $Smn^{+/-};SMN2$ healthy control littermates were carried out in the University of Oxford Biomedical Sciences Unit (BSU), in accordance with the UK Home Office authorisation (Animals Scientific Procedures Act (1986), UK Home Office Project Licence PDFEDC6F0).

The $Smn^{2B/-}$ SMA mouse model [43] was generated by breeding $Smn^{2B/2B}$ mice (generously provided by Dr Rashmi Kothary (University of Ottawa), Dr Lyndsay Murray (University of Edinburgh) and Professor Matthew Wood (University of Oxford) before being sent to Charles River for rederivation) with $Smn^{+/-}$ mice (B6.Cg-$Smn1/J$, stock #007963, Jackson Labs). All live procedures on wild type (WT) (C57BL/6 background), $Smn^{2B/-}$ SMA and $Smn^{2B/+}$ healthy littermates were performed in the Keele University BSU, in accordance with the UK Home Office authorisation (Animals Scientific Procedures Act (1986), UK Home Office Project Licence P99AB3B95).

For all behavioural experiments, body weights and righting reflex [44] (up to 30 seconds) were assessed daily from birth until animals reached their humane endpoint, defined in our UK Home Office Project Licence (P99AB3B95) as the time at which the animal displays hindlimb paralysis, immobility, inability to right (greater than 30 seconds), 4 consecutive days of weight loss and/or up to 20% body weight loss.

As previously described [27], prednisolone (5 mg tablet, Almus) (5 mg/kg dissolved in water) was administered every second day by gavage from post-natal day (P)0 to P7 in $Smn^{-/-};SMN2$ SMA and $Smn^{+/-};SMN2$ healthy mice and from P0 to P20 in $Smn^{2B/-}$ SMA and $Smn^{2B/+}$ healthy mice. Metformin hydrochloride (#PHR1084, Sigma-Aldrich) was dissolved in 0.9% saline physiological

solution (tablet dissolved in sterile water) (#07982, Sigma) and administered daily (200 or 400

mg/kg) by gavage from P5 to humane endpoint in $Smn^{2B/-}$ SMA, $Smn^{2B/+}$ healthy and WT mice.

Oxandrolone (#SML0437, Sigma-Aldrich) was prepared in 0.5% carboxymethylcellulose (CMC)

solution (powder dissolved in 0.9% saline solution) (#C5678, SLS) by sonication at 37 kHz for 3

minutes and administered (4 mg/kg) daily by gavage from P8 to P21 in $Smn^{2B/-}$ SMA, $Smn^{2B/+}$

healthy and WT mice.

Blood-glucose measurement

Blood was collected from non-fasted $Smn^{2B/-}$ SMA and $Smn^{2B/+}$ healthy mice and glucose levels

were immediately measured (mmol/L) via True Metrix® GO Self-Monitoring Blood Glucose

Meter (Trividia Health™) and True Metrix® Test Strips (Trividia Health™).

RNA-Sequencing (RNA-Seq)

Total RNA was extracted from triceps of symptomatic P7 untreated and prednisolone-treated

Taiwanese $Smn^{-/-};SMN2$ SMA and $Smn^{+/-};SMN2$ healthy mice (Table S1). The triceps were

immediately placed in RNALater (#AM7030, ThermoFisher) following dissection and stored at -

20°C under further analysis. For mRNA isolation, 500 ng of total RNA was used as input for the

NEBNext® Poly(A) mRNA Magnetic Isolation Module' (#E7490L, New England Biolabs (NEB))

in accordance with the manufacturer's standard instruction. Library preparation was carried out

using the NEBNext® Ultra Directional RNA Library Prep Kit for Illumina (#E7420L, NEB).

Barcoded libraries from each experimental sample were combined in equimolar concentrations of

1.5 pM prior to sequencing at 75bp x 1 (single-end) read metric on a NextSeq 550 (Illumina)

system.

Differential gene expression analysis

For the RNA-Seq data from the triceps of P7 untreated and prednisolone-treated Taiwanese *Smn⁻/⁻;SMN2* SMA and *Smn⁺/⁻;SMN2* healthy mice (Table S1), differential gene expression (DGE) analysis was performed in Galaxy (usegalaxy.org) [45]. After initial quality control assessments via FastQC v0.72+galaxy1 [46], we trimmed reads based on SLIDINGWINDOW of 4 bp at average quality read of 32 in Trimmomatic v0.36.5 [47] and trimmed the first 12 abnormal bases in Trim sequences v1.0.2 [48]. After quality control confirmation with FastQC v0.72+galaxy1 [46] the processed 63 bp single-reads were aligned to an in-built UCSC *Mus musculus* mm10 genome via HISAT2 v2.1.0 [45,49] under a reverse (or antisense) strand setting. Count quantification of aligned single-reads to mapped coding genes was performed by FeatureCounts v1.6.3+galaxy2 [45,50] using an in-built Entrez *Mus musculus* mm10 gene transfer file (GTF) with known gene identifier set at *-exon level*. Mapping and count quantification was visualized through MultiQC v1.6 [51]. For DGE analysis of our raw transcript counts, we used DESeq2 v2.11.40.2 [52] under the design formula of "Condition" (SMA vs Healthy) and "Treatment" (Untreated vs Prednisolone) after removal of 1 outlier (prednisolone-treated *Smn⁻/⁻;SMN2* sample N0603). We set differentially expressed gene (DEG) significance at log2 fold change (FC) > 0.6 and false discovery rate (FDR) < 0.05. The Entrez Gene IDs were translated to official *Mus musculus* gene symbols by AnnotatemyID v3.7.0 [45]. The normalized count files generated by DESeq2 of prednisolone-treated Taiwanese *Smn⁻/⁻;SMN2* SMA mice vs untreated *Smn⁻/⁻;SMN2* SMA and *Smn⁺/⁻;SMN2* healthy mice were generated into heatmaps by Heatmap2 v2.2.1+galaxy1 [45] under *-data log2 transformed* and scaled by *-row scale genes*.

Pathway analysis and drug repositioning

Pathway analysis of the prednisolone-treated vs untreated Taiwanese *Smn-/-;SMN2* SMA mice was performed with iPathwayGuide [53] (Advaita) with default criteria of log2FC > 0.6 and FDR < 0.05 for DEGs. The impact analysis performed by iPathwayGuide incorporated our DEGs expression (log2FC) and its topological position in the KEGG pathway database [54] to calculate significantly impacted pathways $p < 0.05$ (KEGG v1910 Release 90.0+/05-29, May 19, GODb v1910 2019-Apr 26). Furthermore, overrepresentation analysis [55] (ORA) with elimination pruning [56] was performed for gene ontology (GO) pathways [57] and predicted upstream regulators (STRING v11.0, Jan 19th, 2019). Drug candidate identification was performed through an in-built KEGG drugs database [54] aligned to KEGG pathways in iPathwayGuide and by drug-gene interactions for upstream regulators in the Drug Gene Interaction Database [58] (DGIdb) v3.0.

C2C12 cell culture

The immortalised murine C2C12 myoblast-like cell line [59] (#CRL-1772, ATCC, USA) was maintained in growth media comprised of high glucose (4.5 g/L) and L-glutamine (0.6 g/L) Dulbecco's Modified Eagle's Media (DMEM) (#BE12-741F, Lonza), 10% foetal bovine serum (FBS) (#10438026, Gibco), and 1% penicillin-streptomycin (10,000 U/ml) (#15140122, Gibco). C2C12 myoblasts were differentiated into myotubes in a differentiation media comprised of high glucose (4.5 g/L) and L-glutamine (0.6 g/L) DMEM (#BE12-741F, Lonza), 2% horse serum (HS) (#26050070, Gibco), 1% penicillin-streptomycin (10,000 U/ml) (#15140122, Gibco), and 0.1% insulin (1 µg/ml) (#I6634, Sigma) for 2–8 days with media replacement every 48 hours. All cultured cells were incubated in humid 37°C and 5% CO_2 conditions (Heracell 150i CO_2 incubator, ThermoScientific).

***In vitro* drug treatment**

Proliferating C2C12 myoblasts were seeded in 6-well plates (x4 wells per group). *In vitro* drug

treatments began at 50-60% confluence for C2C12 myoblasts and D7 stage for C2C12 myotubes.

For metformin groups, they were treated with metformin hydrochloride (#PHR1084, Sigma-

Aldrich) dissolved in phosphate buffered saline (PBS), pH 7.4 (#10010023, ThermoFisher) at

concentrations of 0.3, 0.6, 1 and 2 mM for 24 hours against a PBS control (0.1% v/v). For

oxandrolone groups, they were treated with oxandrolone (#SML0437, Sigma-Aldrich) dissolved

in ethanol absolute > 99.8% (#20821.296, VWR) at concentrations of 1, 10 and 100 μM for 24

hours against an ethanol absolute > 99.8% vehicle control (0.1% v/v).

Lactate dehydrogenase (LDH) assay

Drug cytotoxicity was measured by the lactate dehydrogenase (LDH)-Glo™ Cytotoxicity Assay

Kit (#J2380, Promega) following manufacturer's instructions. Luminescence was measured at 400

nm using the GloMax® Explorer Multimode Microplate Reader (#GM3500, Promega).

Bromodeoxyuridine (BrDU) cell proliferation assay

Cell proliferation was measured by the Bromodeoxyuridine (BrDU) Cell Proliferation Assay Kit

(#QIA58, Sigma-Aldrich) following manufacturer's instructions. Absorbance was measured at

450 – 540 nm using the GloMax® Explorer Multimode Microplate Reader (#GM3500, Promega).

Small interfering RNA-mediated *Smn* knockdown in C2C12 cells.

A 10 μM *Smn* small interfering RNA (siRNA) (Duplex name: mm.RiSmn1.13.1) (Integrated DNA

Technologies (IDT)) was used to knock down *Smn* levels against a 10 μM scrambled siRNA

(scrambled DsiRNA, #51-01-19-08) (IDT) negative control. The *Smn* and scrambled siRNAs were

aliquoted separately into an siRNA-lipofectamine complex containing Lipofectamine®

RNAiMAX Reagent (#13778075, Invitrogen) and Opti-MEM (#31985062, Gibco) following

manufacturer's instructions. C2C12 myoblasts were transfected for 48 hours with *Smn* depletion,

whilst C2C12 myotubes were freshly transfected at differentiation (D) stages D0, D3 and D6 for

48 hours with *Smn* depletion confirmed via quantitative polymerase chain reaction (qPCR) (Table

S2).

Serum starvation-induced canonical muscle atrophy in differentiated C2C12 myotubes

Differentiated C2C12 myotubes were incubated in serum-free glucose (4.5 g/L) and L-glutamine

(0.6 g/L) DMEM (#BE12-741F, Lonza) with 1% Penicillin-Streptomycin (10,000 U/ml)

(#15140122, Gibco) for 24 hours. Atrophy was confirmed by upregulation of atrogenes *Atrogin-1*

and *MuRF1* via qPCR (Table S2) and morphology via microscopy (Motic AE31E).

RNA extraction and quantitative polymerase chain reaction (qPCR)

RNA extraction for C2C12 cells was performed with the ISOLATE II RNA Mini Kit (#BIO-

52073, Meridian BIOSCIENCE) as per manufacturer's instructions. Skeletal muscle (triceps and

Tibialis anterior (TA)) and spinal cord tissue samples underwent homogenization with 7 mm

stainless steel beads (#69990, Qiagen) in a Tissue Lyser LT (#85600, Qiagen) set at 60

oscillations/second for 2 minutes followed by microcentrifugation at >10,000 RCF (MSE Sanyo

Hawk 15/05) for 1 minute. RNA extractions from harvested skeletal muscle was performed with

the RNeasy Fibrous Tissue Kit (#74704, Qiagen) and all other harvested tissues with ISOLATE II

RNA Mini Kit (#BIO-52073, Meridian BIOSCIENCE) as per manufacturer's instructions. RNA

concentrations (ng/µl) were measured with the NanoDrop 1000 spectrophotometer

(ThermoScientific) before reverse transcription was performed using the qPCRBIO cDNA

Synthesis Kit (#PB30.11-10, PCR Biosystems) as per manufacturer's instructions.

The cDNA was then diluted by 1:5 in nuclease-free water (#10526945, ThermoFisher). qPCR was

performed using 2x PCRBIO Sygreen Blue Mix Hi-ROX (#PB20.16-20, PCR Biosystems),

nuclease-free water (#10526945, ThermoFisher) and 10 µM forward and reverse primers obtained

from IDT (Table S2). The qPCR reactions were performed in the StepOnePlus™ Real-Time PCR

System (ThermoScientific) with the following program: initial denaturation (95°C for 2 minutes),

40 cycles of 95 °C for 5 seconds and 60 °C for 30 seconds and ending with melt curve stage (95 °C

for 15 seconds, 60 °C for 1 minutes and 95 °C for 15 seconds). Relative gene expression was

quantified using the Pfaffl method [60] and referenced against the validated *RNA polymerase II*

polypeptide J (*PolJ*) housekeeping gene [27,61] (Table S2). Primer efficiency was calculated using

LinRegPCR V11.0 [62].

***Caenorhabditis elegans* drug treatments**

The *Caenorhabditis elegans* (*C. elegans*) SMA strains used in this study was LM99 *smn-*

1(ok355)I/hT2(I;III), which was segregated into homozygotes *smn-1(ok355)*, lethal homozygotes

hT2/hT2 and control heterozygotes *smn-1(ok355)I/hT2* [63]. These animals were maintained at 20°C

on Nematode Growth Medium (NGM) plates seeded with *Escherichia coli* OP50 bacteria [64]. *C.*

elegans were treated by mixing metformin hydrochloride (#PHR1084, Sigma-Aldrich) dissolved

in water at concentrations of 0, 1, 10 and 50 mM and oxandrolone (#SML0437, Sigma-Aldrich)

dissolved in DMSO at concentrations of 0, 1, 10 and 50 µM with NGM agar solution.

Caenorhabditis elegans **neuromuscular assays**

Neuromuscular assays were performed on day 3 animals raised on plates containing the pertinent

solvent or drug. The pharyngeal pumping assay was performed as previously described [65]. Briefly,

animals were filmed with a 150x objective using a AxioCam ICc5 camera at 175 frames/10sec on

a Discovery.V8 SteREO microscope. Pumps were manually counted using the Zen Pro software

v2.3. A pumping event was defined as a grinder movement in any axis. For locomotion assays

animals were filmed with a 63x objective using a AxioCam ICc5 camara at 15 frames/sec on a

Discovery.V8 SteREO microscope. Reversals, and paralysis time for 5 minutes ($\pm$SEM) were

quantified using WormLab 1.1 software (MBF Bioscience). Final data represents three

independent trials (n $\geq$ 25 animals in total per genotype).

Statistical Analyses

Statistical analyses were carried out using the most up to date GraphPad PRISM software. Prior to

any analyses, outliers were identified via Grubb's test (GraphPad) and subsequently removed.

Appropriate statistical tests include unpaired t-test, one-way analysis of variance (ANOVA), and

two-way ANOVA. Each post-hoc analyses used is noted in the respective figure legend. Kaplan-

Meier survival curves were analysed with a log-rank test. Statistical significance was considered

at $p < 0.05$, described in graphs as *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ and ****$p < 0.0001$.

RESULTS

Prednisolone restores the expression of a large subset of genes involved in canonical skeletal muscle pathways in SMA mice.

As described in an earlier study, we have previously demonstrated that treating SMA mice with prednisolone significantly improved several disease phenotypes, including survival, weight, and muscle health [27]. To have a more in depth understanding of the impact of prednisolone on SMA skeletal muscle at a molecular level, we performed bulk RNA-Seq on skeletal muscle of untreated and prednisolone-treated *Smn⁻ᐟ⁻;SMN2* SMA and *Smn⁺ᐟ⁻;SMN2* healthy mice. Specifically, we administered prednisolone (5 mg/kg, gavage, every 2 days) starting from P0 until P7 to *Smn⁻ᐟ⁻;SMN2* SMA and *Smn⁺ᐟ⁻;SMN2* healthy mice [27]. Triceps were harvested from P7 prednisolone-treated and untreated mice for RNA-Seq via Illumina NextSeq550 and a HISAT2-FeatureCounts-DESeq2 pipeline against a Mus Musculus mm10 genome for parameters of "condition" and "treatment" (Figures S1).

Initially, our principal component analysis (PCA) revealed distinct clusters of untreated *Smn⁻ᐟ⁻;SMN2* SMA and untreated *Smn⁺ᐟ⁻;SMN2* healthy littermates, with prednisolone-treated *Smn⁻ᐟ⁻;SMN2* SMA mice falling between the aforementioned groups (Figure 1.a). Importantly, we found that prednisolone treatment restored the expression of 1361 genes in *Smn⁻ᐟ⁻;SMN2* SMA mice to levels similarly observed in untreated *Smn⁺ᐟ⁻;SMN2* healthy mice (Figure 1.b; Tables S3-5).

Next, we determined the biological pathways associated with DEGs in prednisolone-treated *Smn⁻ᐟ⁻;SMN2* SMA mice compared to untreated *Smn⁻ᐟ⁻;SMN2* SMA mice. Using iPathwayGuide, we identified that 3056 significant DEGs (Log2FC > 0.6, FDR < 0.05) (Table S4) were targeted by prednisolone in the skeletal muscle of *Smn⁻ᐟ⁻;SMN2* SMA mice when compared to untreated *Smn⁻ᐟ⁻;SMN2* SMA mice and associated with 28 significant KEGG pathways (p <0.05) (Table 1).

Interestingly, these prednisolone-targeted pathways are closely associated with important skeletal

muscle processes such as metabolism, atrophy and regulatory function, alongside previous

associations with SMA-related pathways such as FoxO signalling [66], p53 signalling [67], AMPK

signalling [68], mitophagy [69], circadian rhythm [70], PPAR signalling [71] and autophagy [72] (Table 1).

An additional GO analysis also revealed similar skeletal muscle biological processes associated

with the DEGs in prednisolone-treated *Smn$^{-/-}$;SMN2* SMA mice such as myotube differentiation,

fatty acid oxidation, protein ubiquitination, sarcomere regulation, gluconeogenesis, and circadian

rhythm (Table 2; Tables S6-8).

Combined, our transcriptomics and pathway analyses suggest that prednisolone treatment

attenuated muscle pathologies in SMA mice [27] by targeting key muscle metabolism, atrophy and

regulatory pathways.

Drug repositioning algorithms identify novel pharmacological compounds predicted to

emulate prednisolone's activity in skeletal muscle of SMA mice.

As mentioned above, while prednisolone treatment significantly improves muscle health and

overall disease progression in SMA mice, chronic use of prednisolone can negatively impact

skeletal muscle [35,39]. As such, we used the DEGs and associated KEGG pathways identified in

prednisolone-treated *Smn$^{-/-}$;SMN2* SMA mice to discover alternative drugs predicted to mimic

prednisolone's molecular effects in SMA skeletal muscle. Initially, we utilized the in-built

integration of KEGG drugs database in iPathwayGuide [53] and the DGIdb v3.0 [58] database to

initially reveal a total of 580 compounds (Tables S9-12). To filter down our list, we focused on

the drug compounds 1) that targeted > 5 prednisolone-targeted pathways or linked to upstream

regulators, 2) were clinically approved and 3) were not associated with promotion of muscle-

wasting (e.g., primary anti-cancer drugs [73]), leaving a total of 20 potential candidates (Tables 3-4).

Interestingly, our combined *in silico* drug repositioning approach revealed a subset of candidates

previously investigated in SMA such as celecoxib [74] (ClinicalTrials.gov ID: NCT02876094) and

colforsin [75]. To further validate our bioinformatics strategy, we chose to continue our study with

drugs not yet assessed for SMA, focusing on those previously used safely in young patients and

orally bioavailable. With these criteria, we narrowed down our selection to metformin, a generic

asymmetric dimethyl-biguanide type 2 diabetes mellitus (T2DM) drug [76] and oxandrolone, a

synthetic anabolic steroid with a higher ratio of anabolic: androgynous effects for further study [77].

Thus, using a transcriptomics-based *in silico* drug repositioning platform, we were able to generate

a list of clinically approved pharmacological compounds that are predicted to emulate

prednisolone's activity in skeletal muscle.

Metformin's primary predicted target gene *Prkag3* is dysregulated in skeletal muscle of both

severe *Smn$^{-/-}$;SMN2* and intermediate *Smn$^{2B/-}$* SMA mice.

As previously mentioned, metformin is an orally administered T2DM drug that we selected as one

of the candidates to validate our bioinformatics-based drug repositioning approach. Importantly,

metformin has over 60 years of clinical use with a well-known safety profile [76] and recorded

administration in younger patients [78]. Furthermore, it has been previously repositioned and

conferred ergogenic activities in muscular disorders such as DMD [79] and congenital muscular

dystrophy type 1 A (CMDT1A) [80], highlighting its potential as a skeletal muscle therapy.

Our iPathwayGuide analysis predicted that metformin could emulate prednisolone's targeting of

the KEGG: 04068 FoxO signalling pathway (Figure S2.a). In particular, metformin was predicted

to mimic prednisolone's upregulation of *Prkag3*, which encodes for the AMPK-γ3 subunit of the

predominant skeletal muscle AMPK-α2β2γ3 isoform complex [81] (Figure S2). Furthermore,

Prkag3 upregulation was predicted to coherently downregulate the expression of *FoxO1*, *FoxO3*

and *Foxo4* isoforms, whilst upregulating *FoxO6* (Figure S2) supporting previous literature

associating these FoxO isoforms with promotion of muscle atrophy [66,82]. Importantly, the

expression pattern of these genes in the prednisolone-treated *Smn$^{-/-}$;SMN2* SMA mice were

normalized to healthy *Smn$^{+/-}$;SMN2* levels (Figure S2.c), supporting the usefulness of investigating

metformin and these targets in SMA skeletal muscle.

We thus measured the mRNA expression levels of *Prkag3* and *FoxO* isoforms in the triceps of

both symptomatic P7 severe *Smn$^{-/-}$;SMN2* and P19 intermediate *Smn$^{2B/-}$* SMA mice alongside their

respective healthy controls. We indeed observed that *Prkag3* levels were significantly

downregulated in skeletal muscle of both SMA mouse models (Figure 2.a), supporting the

bioinformatics data. However, none of the *FoxO* isoforms were significantly different between

SMA mice and their healthy controls (Figures 2.b-e). Previous research also reported no significant

upregulation of *FoxO* isoforms in P7 severe *Smn$^{-/-}$;SMN2* SMA mice via qPCR [66]. However, the

fact that our qPCR data did not reflect the bioinformatics predictions may be due to variability in

our experimental cohorts, the sequencing depth coverage not being sufficiently conservative

and/or intrinsic differences between RNA-Seq and primer-based qPCR approaches [83].

Overall, our qPCR experiments revealed that the primary metformin target *Prkag3* matched its

bioinformatics prediction of being downregulated in both severe *Smn$^{-/-}$;SMN2* and intermediate

Smn$^{2B/-}$ SMA mice, suggesting that this gene may be involved in both severe and milder SMA

pathologies and an appropriate therapeutic molecular target in SMA muscle.

The predicted target genes for metformin are mostly Smn-independent in an SMA muscle cellular model.

We next wanted to better understand if the aberrant expression of the metformin target genes was dependent on SMN expression and/or muscle atrophy. Thus, we firstly generated siRNA-mediated Smn-depleted C2C12 myoblast-like cells, a useful and previously successful *in vitro* model [84]. We confirmed by qPCR that *Smn* mRNA levels were significantly reduced by up to 90% in C2C12 myoblasts and D8 C2C12 myotubes compared to scrambled siRNA and untreated controls (Figure S3). We next investigated the effects of Smn knockdown on the expression of the predicted metformin target genes. In C2C12 myoblasts, we identified a significant upregulation of only the *FoxO3* gene in Smn-depleted C2C12 myoblasts compared to controls (Figure 3.a), which reflects previous microarray analyses of specific *FoxO* isoforms upregulated in quadriceps femoralis muscle biopsies from type 1 SMA patients [85]. However, in C2C12 myotubes we found that Smn knockdown (KD) had no effect on the expression of predicted metformin target genes (Figure 3.b), suggesting that for the most part, the expression of the predicted metformin genes is Smn-independent, thus representing ideal targets for SMN-independent therapies.

We next investigated if the expression of the predicted metformin target genes is affected *in vitro* by muscle atrophy. However, one difficulty in mimicking SMA muscle atrophy *in vitro* is establishing denervation. Thus, based on evidence of shared pathway similarities from different pro-atrophy factors such as starvation and denervation [86], we used a validated method of 24-hour serum-starvation in C2C12 myotubes to induce canonical atrophy, as confirmed by myotube loss and upregulation of pro-atrophic *atrogin-1* levels (Figure 3.c). Next, we evaluated the expression of the predicted metformin target genes and observed a significant upregulation of *FoxO3* and

FoxO4 isoforms (Figure 3.d), reflecting their established roles in atrophy-dependent ubiquitin-proteasome pathways [66].

We then evaluated whether metformin could attenuate muscle atrophy in C2C12 myotubes. Based on initial gene-dose response experiments in both control C2C12 myoblasts and D8 myotubes, we treated our cells with physiological (60 µM) and supraphysiological (2 mM) metformin concentrations for 24 hours (Figures S4) [88]. The 30 µM physiological metformin concentration for 24 hours did not attenuate muscle atrophy or impact the expression of the target genes in the serum starved C2C12 myotubes (Figure 3.e-f). However, for the supraphysiological 2 mM metformin concentration [88], we observed an upregulation of *Atrogin-1* levels (Figure 3.e), suggesting an exacerbation of muscle atrophy. Further analysis of the predicted metformin target genes revealed no significant impact on their expression patterns either (Figure 3.f), suggesting that exacerbation of atrophy in C2C12 myotubes by supraphysiological metformin concentrations involve factors mostly outside of our predicted targets. However, it should be noted that metformin may have different effects in SMA muscle as there are still differences between distinct pro-atrophic factors [86].

Overall, our *in vitro* studies revealed that although most of our predicted metformin target genes are SMN-independent with some linked to muscle atrophy, they were mostly not linked to metformin's influence on canonical atrophy in C2C12 myotubes.

Dose-dependent effect of metformin on disease progression and survival in *Smn²ᴮ/⁻* SMA mice.

Next, we assessed metformin in the intermediate *Smn²ᴮ/⁻* SMA mouse model [43]. The rationale for conducting our *in vivo* pharmacological studies in the *Smn²ᴮ/⁻* SMA mice was based on their longer

lifespan [43], responsiveness to SMN-independent therapies [42,70], established metabolic and

myopathy defects [43,66,70,89–91], and later symptomatic onset [43] making them a clinically relevant

model for starting treatment regimens >P5 time-points [42,70].

We initially administered a 200 mg/kg daily dose (diluted in 0.9% saline) starting from P5 until

humane endpoint in $Smn^{2B/-}$ SMA mice and $Smn^{2B/+}$ healthy control littermates, based on this

concentration's previous success in the muscle disorder CMDT1A [80]. We observed no significant

improvement in survival of $Smn^{2B/-}$ SMA mice treated with 200 mg/kg/day compared to untreated

(Figure 4.a) and vehicle-treated animals (Figure S5). We also found a significant reduction in the

body weight of 200 mg/kg/day metformin-treated $Smn^{2B/-}$ SMA mice compared to untreated SMA

animals, beginning 4 days after initial treatment at P9 (Figure 4.b.) However, we did not observe

any effects on weight in the 200 mg/kg/day metformin treated $Smn^{2B/+}$ healthy control mice (Figure

S6.b), indicating a disease specific effect of metformin. In terms of motor function, there was no

significant difference in the righting reflex between untreated and 200 mg/kg/day metformin-

treated $Smn^{2B/-}$ SMA mice (Figure 4.c) and $Smn^{2B/+}$ healthy control animals (Figure S6.a-c).

Since our initial 200 mg/kg/day metformin dosage did not improve disease onset or disease

progression in $Smn^{2B/-}$ SMA mice, we conducted pilot studies with a later treatment start point (P8)

and a lower dose (100 mg/kg/day), which demonstrated similar effects to our initial dosing regimen

(data not shown). We therefore tried a higher daily dosage of 400 mg/kg/day, starting at P5.

Surprisingly, the higher 400 mg/kg/day dose significantly reduced survival in $Smn^{2B/-}$ SMA pups

by 5 days (Figure 4.d), while having no significant impact on weight or righting reflex (Figures

4.e-f). Interestingly, 400 mg/kg/day metformin had no adverse effects in the healthy $Smn^{2B/+}$

control littermates (Figures S6.d-f), suggesting that the adverse effects of the higher dose of

metformin is disease specific.

Thus, our *in vivo* experiments demonstrated that metformin did not emulate prednisolone's

beneficial effects on SMA disease progression in SMA mice and is in fact not an ideal therapy

candidate, due to dose- and disease-dependent adverse effects in SMA.

Higher 400 mg/kg/day metformin dosage is associated with hypoglycaemia in non-fasted

$Smn^{2B/-}$ SMA mice.

We next investigated the potential causes behind metformin's adverse effects in SMA mice. With

metformin being a glucose lowering agent for T2DM, we initially assessed blood glucose levels

in P14 non-fasted, untreated, 200- and 400-mg/kg/day metformin treated $Smn^{2B/-}$ SMA and $Smn^{2B/+}$

healthy mice 2 hours after the final treatment. This time point was chosen to account for the

reduced median survival in the higher dose SMA cohort. We observed that neither 200- or 400-

mg/kg/day metformin treatments lowered blood glucose levels in $Smn^{2B/+}$ healthy mice (Figure

5.a). However, we reported a significant reduction in blood glucose levels in 400 mg/kg/day

metformin-treated $Smn^{2B/-}$ SMA mice compared to untreated SMA animals (Figure 5.a). Our

results suggest that hypoglycaemic shock could have been one of the possible causes behind the

premature death in the 400 mg/kg/day SMA cohort, further exacerbating the previously reported

hypoglycaemia in SMA models [89] and patients [92,93].

Metformin did impact the expression of the predicted target genes, but not muscle pathology

markers in $Smn^{2B/-}$ SMA mice.

We next evaluated the effect of metformin on the expression of the predicted target genes in TAs

from P14 untreated, 200- and 400-mg/kg/day metformin-treated $Smn^{2B/-}$ SMA and $Smn^{2B/+}$ healthy

mice 2 hours after final treatment. Contrary to the drug repositioning prediction that metformin

could reverse *Prkag3* downregulation in SMA muscle (Figure S3), we instead discovered that both

200- and 400-mg/kg/day metformin doses exacerbated *Prkag3* downregulation in *Smn*$^{2B/-}$ SMA

muscle (Figure 5.b). Furthermore, although metformin had no impact on *FoxO1*, *FoxO3* and

FoxO4 isoforms in both *Smn*$^{2B/+}$ healthy and *Smn*$^{2B/-}$ SMA muscle (Figures 5.c-e), for the 200

mg/kg/day *Smn*$^{2B/-}$ SMA cohort, metformin significantly reduced *FoxO6* expression (Figure 5.f),

which again contrasted our bioinformatics prediction that metformin would upregulate the

expression of this isoform in SMA muscle. Thus, the drug-gene response for metformin-treated

Smn$^{2B/-}$ SMA skeletal muscle reveals a contrasting pattern that does not match the bioinformatic

predictions.

We next investigated metformin's effects on the expression of dysregulated molecular markers

associated with muscle atrophy (*Atrogin-1* and *MuRF-1*) and glucose metabolism (*Glut4* and *Hk2*)

[94,95]. We however observed that neither atrophy (Figures 5.g-h) or glucose metabolism markers

(Figures 5.i-j) were affected by 200- and 400-mg/kg/day metformin treatments in the *Smn*$^{2B/-}$ SMA

mice when compared to untreated animals.

We also investigated markers associated with mitochondrial biogenesis and function in muscle

(*Pgc1-α, Tfam*, *Ndufs1*, and *Nrf1*), as previous research has established that these features are

impaired in SMA skeletal muscle [69] and a common mechanism of action for metformin is mild

inhibition of mitochondrial electron transport complex 1 (or NADH:ubiquinone oxidoreductase)

[96]. We found that that neither the 200- or 400-mg/kg/day dose of metformin impacted the

expression of mitochondrial genes in the skeletal muscle of *Smn*$^{2B/-}$ SMA mice (Figures 5.k-n),

although we do observe a significant downregulation of *Pgc1-α, Tfam*, and *Ndufs1* in the skeletal

muscle of 200 mg/kg/day metformin-treated *Smn*$^{2B/+}$ healthy counterparts (Figures 5.k-n).

Overall, our data highlights that metformin did not have a direct impact on the predicted target

genes in skeletal muscle of SMA mice. Furthermore, the absence of direct impact on muscle

atrophy, glucose metabolism, and mitochondrial function markers following metformin treatment

in $Smn^{2B/-}$ SMA muscle, suggests that the adverse effects associated with the 400 mg/kg/day dosage

may not have been linked to muscle-intrinsic effects.

A higher dose of metformin is associated with dysregulation of mitochondrial regulatory

genes in the spinal cord of $Smn^{2B/-}$ SMA mice.

We next investigated the effects of metformin on the spinal cord given that metformin is

systematically distributed [97], has the ability to cross the blood-brain-barrier (BBB) [98] and can impact

the mitochondria in the spinal cord [99]. We thus evaluated whether metformin altered the expression

of mitochondrial markers (*Pgc1-α*, *Tfam*, *Nrf1* and *Ndufs1*) in the spinal cord of P14 untreated,

200- and 400-mg/kg/day metformin treated $Smn^{2B/-}$ SMA mice compared to $Smn^{2B/+}$ healthy mice,

2 hours after final treatment.

For *Pgc1-α*, a master regulator of mitochondrial biogenesis and function, we observed that

although both 200- and 400-mg/kg/day metformin concentrations significantly reduced its

expression levels in $Smn^{2B/+}$ healthy spinal cords (Figure 6.a), it was only the higher concentration

that significantly reduced *Pgc1-α* expression in $Smn^{2B/-}$ SMA spinal cords (Figure 6.a). Similarly,

400 mg/kg/day metformin significantly reduced *Ndufs1* levels in both $Smn^{2B/+}$ healthy and $Smn^{2B/-}$

SMA spinal cords (Figure 6.b), suggesting that for these mitochondrial health markers the higher

metformin dose negatively affected their expression independent of disease status. On the other

hand, although not affected by metformin in $Smn^{2B/-}$ SMA spinal cords, *Nrf1* gene expression was

significantly reduced by both metformin doses in the spinal cord of $Smn^{2B/+}$ healthy mice (Figure

6.c), whilst *Tfam* was not affected by metformin in either cohort (Figure 6.d).

Our results demonstrating that the higher dose of metformin (400 mg/kg/day) appears to

specifically dysregulate certain mitochondrial genes in the spinal cord of $Smn^{2B/-}$ SMA mice is

supported by recent evidence of tissue-dependent differences in conserved cellular processes

between SMA motor neurons and skeletal muscle [72]. Thus, although further in-depth investigations

would be needed, our results on mitochondrial health markers suggest that metformin's adverse

effects in SMA mice could be linked to the exacerbation of neuronal mitochondrial dysfunction.

Oxandrolone's predicted target gene, *Ddit4*, is dysregulated in the skeletal muscle of severe

***Smn$^{-/-}$;SMN2* SMA mice.**

Our second drug candidate that we selected to mimic prednisolone activities was oxandrolone, a

synthetic orally bioavailable anabolic steroid that confers minimal androgynous effects [77].

Importantly for SMA, oxandrolone has been successful in the promotion of muscle growth for

DMD [100] and mixed gender burn injury patients [101].

Oxandrolone was predicted to upregulate the *Ar* gene in SMA muscle (Figure S7). The

upregulation of *Ar* was predicted to directly upregulate downstream effectors *Igfbp5* and *myogenin*

(or *MyoG*) (Figure S7), which both regulate muscle differentiation, regeneration and myofiber

growth [102,103]. Furthermore, *Ar* was predicted to indirectly upregulate *Dok5*, a signalling protein

linked to insulin and IGF-1 activity [104] and *Akap6*, which is involved in the modulation of muscle

differentiation and regeneration [105] (Figure S7). In addition to these factors, we also decided to

investigate *Ddit4* as an oxandrolone target based on its direct relation with *Ar* [106] and being one of

the top 20 downregulated DEG targets of prednisolone in *Smn⁻/⁻;SMN2* SMA skeletal muscle

(Table S4; Figure S7).

Similar to our metformin strategy above, we initially wanted to evaluate the mRNA expression

levels of these target genes in the triceps of both symptomatic P7 severe *Smn⁻/⁻;SMN2* and P19

intermediate *Smn²ᴮ/⁻* SMA mice alongside their respective healthy controls. Overall, we identified

no significant dysregulated expression of the target genes *Ar*, *Akap6*, *Igfbp5*, *Dok5* and *MyoG* in

both severe *Smn⁻/⁻;SMN2* and intermediate *Smn²ᴮ/⁻* SMA mice (Figures 7.a-e). However, for *Ddit4,*

we did identify a significant upregulation only in *Smn⁻/⁻;SMN2* SMA mice (Figures 7.f), supporting

both our bioinformatics data for this gene and its known pro-atrophic role [106], indicating that it

may play an important role in SMA muscle pathologies. In summary, the majority of the predicted

oxandrolone target genes did not significantly reflect their bioinformatic predictions.

***In vitro* oxandrolone treatment prevents canonical atrophy in C2C12 myotubes**

independently of the predicted Smn-independent targets.

Similar to our metformin *in vitro* studies, we wanted to evaluate whether reduced SMN levels or

atrophy influenced the expression of the predicted oxandrolone target genes in SMA skeletal

muscle. Although none of the target genes were affected in the Smn-depleted C2C12 myoblasts

(Figure 8.a), we found that Smn KD triggered a significant upregulation of *Dok5* only in C2C12

myotubes (Figure 8.b), suggesting that the expression of this gene may be Smn-dependent.

Nevertheless, the expression of the majority of the predicted oxandrolone target genes was Smn-

independent.

We next wanted to evaluate the ability of oxandrolone to attenuate canonical atrophy in serum-

deprived C2C12 myotubes [87]. However, in this case we performed the treatments in D5 C2C12

myotubes instead of D8, as although oxandrolone was non-toxic (Figures S8), it elicited a greater

androgen *Ar* response at the earlier differentiation stage (Figure S9). Following confirmation of

muscle atrophy in D5 C2C12 myotubes via significant *Atrogin-1* and *MuRF1* upregulation (Figure

8.c), we observed that the expression of the predicted oxandrolone target genes *Akap6*, *Igfbp5*,

MyoG and *Ddit4* was significantly downregulated in serum-deprived D5 C2C12 myotubes (Figure

8.d).

Interestingly, we found that 24-hour treatment with 1 μM oxandrolone attenuated canonical

muscle atrophy in these serum-starved D5 C2C12 myotubes as shown by significant

downregulation of *Atrogin-1* (Figure 8.e). However, we observed that *Igfbp5*, *MyoG* and *Ddit4*

were further downregulated by the 1 μM oxandrolone treatment (Figure 8.f), suggesting that

oxandrolone's effects on atrophy are linked to effectors independent of the predicted target genes.

Overall, our *in vitro* studies have shown that although the expression of the predicted oxandrolone

target genes is Smn-independent, they are not involved in oxandrolone's ameliorative effects on

canonical atrophy in C2C12 myotubes.

Oxandrolone treatment improves survival in *Smn$^{2B/-}$* SMA mice.

We next assessed the impact of oxandrolone in SMA mice. We initially tested preliminary

treatment regimens of 1 – 8 mg/kg/day starting from P5 or P8 in *Smn$^{2B/-}$* SMA and *Smn$^{2B/+}$* healthy

mice (data not shown), based on previous studies in models of spinal cord injury (SCI) [107] and burn

injury [108]. We also stopped oxandrolone treatments at P21 as previous research has shown that

shorter oxandrolone treatments are more effective [107]. These pilot studies allowed us to identify

the optimal dosing regimen of 4 mg/kg/day oxandrolone treatment from P8 to P21, which

significantly improved the median survival of *Smn$^{2B/-}$* SMA mice (Figure 9.a).

However, we found that the body weight of 4 mg/kg/day oxandrolone-treated $Smn^{2B/-}$ SMA mice

was significantly lower compared to their untreated counterparts (Figure 9.b), which is most likely

due to the intrinsic smaller sizes of the randomly assigned litters, as demonstrated by the difference

in weight starting 4 days prior to initial treatment (Figure 9.b). In terms of motor function, we

observed no significant difference in righting reflex between untreated and oxandrolone-treated

SMA animals (Figure 9.c). Furthermore, we identified no impact of vehicle treatment on survival,

weight, and righting reflex in $Smn^{2B/-}$ SMA mice (Figures S10).

In the $Smn^{2B/+}$ healthy mice, although 4 mg/kg/day oxandrolone had no effect on survival or motor

function in treated animals (Figures S11.a-b), we did observe a significant decrease in bodyweight

starting from P9, one day after initial treatment (Figure S11.c), suggesting that oxandrolone may

have impacted growth.

Nevertheless, our results demonstrated that although 4 mg/kg/day oxandrolone treatment

improved survival in $Smn^{2B/-}$ SMA mice, its effect on survival was still minor compared to

prednisolone [27], suggesting that oxandrolone is not a suitable substitute as an SMA skeletal muscle

therapy.

Oxandrolone did not impact the expression of the predicted target genes or muscle pathology

markers.

The improved 3-day survival in 4 mg/kg/day oxandrolone-treated $Smn^{2B/-}$ SMA mice led us to

evaluate whether this beneficial impact was related to targeting muscle pathologies. Thus, we

evaluated oxandrolone's effects on the expression of dysregulated molecular markers associated

with the SMA hallmark pathology of muscle atrophy (*Atrogin-1* and *MuRF-1*) in the triceps from

P19 late symptomatic, untreated and 4 mg/kg/day oxandrolone-treated $Smn^{2B/-}$ SMA and $Smn^{2B/+}$

healthy mice, 2 hours after final treatment. We observed no significant reduction in elevated

Atrogin-1 or *MuRF-1* gene expression levels by oxandrolone in the *Smn²ᴮ/⁻* SMA cohort (Figures

9.d-e), suggesting that oxandrolone did not attenuate muscle atrophy.

We next evaluated the effect of oxandrolone on the expression of the predicted target genes in the

same P19 untreated and 4 mg/kg/day oxandrolone-treated *Smn²ᴮ/⁻* SMA and *Smn²ᴮ/⁺* healthy mice.

We found that oxandrolone did not significantly impact the predicated target genes in the triceps

from the *Smn²ᴮ/⁻* SMA mice (Figures 9.f-k). However, we did observe that 4 mg/kg/day

oxandrolone treatment significantly upregulated *Dok5* expression in the *Smn²ᴮ/⁺* healthy mice

(Figure 9.i). Nevertheless, the pattern observed suggests that oxandrolone did not impact any of

the predicted genes in the muscle from *Smn²ᴮ/⁻* SMA mice.

Overall, our data shows that oxandrolone did not have an efficient effect on the predicted target

genes. Furthermore, its inability to ameliorate muscle atrophy marker dysregulation in SMA

skeletal muscle, suggests that improved survival in the *Smn²ᴮ/⁻* SMA mice by oxandrolone may be

independent of targeting skeletal muscle pathologies.

Both metformin and oxandrolone drug candidates attenuate neuromuscular phenotypes in

the *C. elegans* severe SMA model.

We next wanted to investigate our drug candidates in a separate SMA model to assess whether

they could attenuate neuromuscular dysfunctions in a distinct species. For this purpose, we used

the *C. elegans smn-1(ok355)* invertebrate model [63], based on shared conservation of the SMN

protein with vertebrate species [109] and the well described phenotypic defects of larval lethality

(reduced survival) and impaired neuromuscular function in pharyngeal pumping for feeding and

locomotion [63,65]. For metformin, administration at higher doses of 50 mM partially ameliorated

multiple phenotypes only in *C. elegans smn-1(ok355)* including pharyngeal pumping defects and

the locomotory defect of number of reversals (Figures 10.a-b), however only the lower dose of 1

mM metformin significantly ameliorated paralysis times (Figures 10.c). On the other hand,

oxandrolone across 1-50 µM doses significantly ameliorated pharyngeal pumping defects only

(Figure 10.d), with no significant effect on locomotory defects of reversal and paralysis times

(Figures 10.e-f), suggesting an improvement in muscular activity in the pharynx region. Overall,

given SMN conservation across species, oxandrolone could improve neuromuscular defects across

vertebrate and invertebrate models of SMA.

DISCUSSION

The goal of this study was to use a transcriptomics-based drug repositioning strategy to identify clinically approved drug candidates that could emulate prednisolone's beneficial effects in SMA skeletal muscle and life expectancy [27], without the risks associated with long-term GC exposure [39].

Our major finding was the observation that prednisolone treatment restored specific gene sets associated with key pathological SMA pathways such as FoxO signalling [66], p53 signalling [67], AMPK signalling [68], mitophagy [69], circadian rhythm [70], PPAR signalling [71], and autophagy [72] in $Smn^{-/-};SMN2$ SMA skeletal muscle. Although these pathway results highlight prednisolone's efficacy in improving skeletal muscle health, it should be noted that our transcriptomic data cannot distinguish whether these restorations are targeted directly by prednisolone or a consequence of improved muscle health. Nevertheless, our transcriptomic data complemented the prior phenotypic data of prednisolone's potential as a second-generation SMA therapy [27].

Despite a multitude of promising compounds identified that could be investigated in future studies, one of the findings of our study was that neither of our chosen orally bioavailable drug candidates, metformin and oxandrolone, reproduced prednisolone's previously reported effect on muscle growth and survival in SMA mice [27]. For metformin, we observed that both 200- and 400-mg/kg/day doses counterintuitively exacerbated $Prkag3$ downregulation in $Smn^{2B/-}$ SMA muscle instead of the predicted upregulation, which could have negative consequences since $Prkag3^{-/-}$ null mice presented metabolic and mitochondrial dysregulation [110,111] similar to those reported in SMA patients [69,93] and models [89,90]. More surprising was the reduced survival of $Smn^{2B/-}$ SMA mice treated with 400 mg/kg/day metformin, which we potentially linked to possible hypoglycaemic shock and/or dysregulation of neuronal mitochondrial components. On the other hand, although

various metformin doses ameliorated neuromuscular dysfunction in our SMA *C. elegans* model,

the negative effects observed in $Smn^{2B/-}$ SMA mice could be to due vertebrate vs invertebrate

systems. Curiously, previous research into AMPK agonism via 500 mg/kg AICAR treatment in

the *SmnΔ7* SMA mice contrasts our metformin data, as they showed improved skeletal muscle

health and no exacerbations in neuronal dysfunction [68]. One explanation for metformin and

AICAR's conflicting results in SMA mice could be related to differential pharmacological

activities. Indeed, AICAR is an AMP analogy with a low BBB penetrability [112] that directly binds

to the γ-AMPK isoform [113,114], whilst metformin can rapidly penetrate the BBB [98] and has been

associated with direct and indirect AMPK activation [97], an example of the latter involving

mitochondrial respiratory complex 1 inhibition [96]. With emerging evidence of naturally low

mitochondrial respiratory complex 1 activity in SMA motor neurons [115] and SMA pathway profiles

being tissue-dependent [72], one theory could be that the 400 mg/kg/day metformin dose exacerbated

mitochondrial respiratory complex 1 inhibition in SMA motor neurons. However, future studies

would be needed to verify this proposed model. Nonetheless, our findings could be important for

clinical drug safety, as with reported co-morbidities of diabetes in certain SMA patients [116,117], our

pre-clinical data suggests lower metformin doses or non-biguanide drugs may be important to

manage diabetes and not risk primary pathologies in SMA patients.

For oxandrolone, our mouse data showed that 4 mg/kg/day treatment from P8 partially improved

survival in $Smn^{2B/-}$ SMA mice, although not to the same extent observed with prednisolone [27]. In

addition, we identified that the lower dose of 1 μM oxandrolone *in vitro* attenuated canonical

atrophy in C2C12 myotubes, whilst *in vivo* oxandrolone attenuated neuromuscular dysfunction in

severe SMA *C. elegans* model, suggesting that in both our SMA vertebrate and invertebrate

species oxandrolone may be a beneficial SMN-independent treatment option. However, one factor

that we did not account for in our oxandrolone investigations was gender specificity. Although the

clinical literature has shown oxandrolone has minimal androgenous effects, studies with *Ar* KO

rodent models have demonstrated that Ar absence does not have the same impact on female muscle

size compared to males [118]. Furthermore, it is suggested that intramuscular Ar content may have a

stronger influence on hypertrophy than peripheral androgen levels [119]. Thus, we cannot conclude

whether oxandrolone's treatment efficacy compared to prednisolone was due to gender-specific

differences.

Despite our study's limitations, it highlighted refinements for future *in silico* SMA drug

repositioning studies. Compared to a previous study that successfully identified and validated

harmine's therapeutic potential in SMA muscle [42], ours did not include proteomics. The absence

of proteomics can be a caveat for drug studies as transcript levels alone do not proportionally

reflect drug-protein interactions, abundance and translational modifications [120,121]. However, a

limitation of both transcriptomics and proteomics approaches is that they cannot bridge drug-

pathway interactions with disease phenotypes, as demonstrated by a recent proteomics analysis of

Spinraza® treated type 2 and 3 SMA patients that could not correlate protein profiles with

functional improvements [122]. Thus, implementation of metabolomics may be beneficial for linking

pathway perturbations with metabolites associated with disease and stages of muscle atrophy [123].

To the best of our knowledge, this three-pronged multi-omics approach has not previously been

used in SMA drug repositioning, however it has been successful in the identification of 200

biomarker candidates for SMA [124].

Another consideration is systemic effects of the drugs as seen with the enhanced lethality of the

400 mg/kg/day metformin's dose in *Smn$^{2B/-}$* SMA mice being linked to hypoglycaemia and

mitochondrial dysfunction in neuronal tissue. Indeed, adverse systemic risks were also found with

a previous multi-omics drug repositioning study for SMA muscle that identified the development

of tremors in harmine-treated $Smn^{2B/-}$ SMA mice [42]. With tissue-dependent differences in

conserved pathways in SMA [72], future omics studies could integrate data from both neuronal and

skeletal muscle to minimize systemic adverse risks.

Even with these refinements, future drug repositioning studies for SMA skeletal muscle may need

to consider replacing bulk RNA-Seq. Skeletal muscle fibers are comprised of a myriad of different

muscle fiber and cell types, alongside non-muscular interconnecting tissues such as neurons,

tendons, adipose, immune cells and capillaries [125,126], which altogether does not truly represent the

transcriptomic profiles for distinct skeletal muscle cells. Indeed, our significant KEGG pathway

results included those associated exclusively with neuronal, immune, and capillary cells such as

glioma, atherosclerosis and Th17 cell differentiation. With alterations in fiber type composition

[91,127] and satellite cell dysregulation evidenced in SMA muscle [23], an emerging alternative to

predict drug candidates that target dysregulated SMA pathways in these muscle types would be

single-cell (scRNA-Seq) and/or single nuclear RNA-Seq (snRNA-Seq) [126], which have already

been useful tools in other muscular disorders such as DMD [128,129]. In addition, snRNA-Seq could

have further benefits to narrow in on muscle regions such as nuclei located near the NMJ, since

myonuclei display spatial and temporal expression pattern heterogeneity in multi-nucleated

myofibers [126].

Although the drug candidate's metformin and oxandrolone did not emulate prednisolone's

beneficial effects in SMA to the extent previously reported, our transcriptomics-drug repositioning

approach did better define prednisolone's activity in SMA muscle and provided a list of potential

candidates for future pre-clinical SMA drug repositioning studies. Furthermore, our study

highlights important refinements for future SMA drug repositioning studies.

Data Availability Statement

The datasets presented in this study can be found in the following online repository: NCBI BioProject, accession ID: PRJNA972323.

AUTHOR CONTRIBUTIONS

Conceptualization: J.M.H, M.B; Methodology: J.M.H, P.P.T, M.D, L.M.W, P.C, D.P.T & M.B; Formal Analysis: J.M.H, P.P.T, M.D, D.P.T & M.B; Investigation: J.M.H, E.M, E.R.S, O.C, P.P.T, M.D, M.O, L.M.W, P.C, & M.B; Software: J.M.H & D.P.T; Visualization: J.M.H; Resources: M.D, P.C, S.D, M.J.A.W, D.P.T & M.B; Writing – Original Draft: J.M.H & M.B; Writing – Review & Editing: J.M.H, E.M, E.R.S, O.C, P.P.T, M.D, M.O, L.M.W, P.C, S.D, M.J.A.W, D.P.T, & M.B; Supervision: D.P.T & M.B; Project Administration: M.B; Funding Acquisition: M.B

FUNDING

J.M.H. was funded by a Ph.D. studentship from the Keele University of School of Medicine. E.M. is supported by an Academy of Medical Sciences grant (SBF006/1162). E.R.S. was funded by a MDUK Ph.D. studentship (18GRO-PS48-0114). O.C. is supported by a Ph.D. studentship from the Republic of Turkey Ministry of National Education. M.B. was funded by SMA Angels Charity. P.C. is supported by the Deutsche Muskelstiftung and the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No 956185.

ACKNOWLEDGEMENTS

We would like to thank the personnel at the Biomedical Sciences Unit at both Keele University and University of Oxford. Many thanks go to members of the Combined Inherited Neuromuscular Disorders meetings for helpful discussions. For the RNA-Sequencing we are grateful for the aid and resources of Research Core Unit Transcriptomics at Hannover Medical School. *C. elegans* strains were provided by the *Caenorhabditis elegans* consortium (CGC), which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). Research in the M.D. lab has been funded by the Wellcome Trust and Royal Society. A special thanks goes out to Advaita for their technical support of iPathwayGuide and to CureSMA, SMA Europe and the Biochemical Society, UK for providing podium presentation opportunities for preliminary data of this project.

REFERENCES

1. Kolb, S. J. & Kissel, J. T. Spinal Muscular Atrophy. *Neurol. Clin.* **33**, 831–846 (2015).

2. Wojcik, M. H. *et al.* Infant Mortality: the Contribution of Genetic Disorders. *J. Perinatol. Off. J. Calif. Perinat. Assoc.* **39**, 1611–1619 (2019).

3. Lefebvre, S. *et al.* Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* **80**, 155–165 (1995).

4. Rodrigues, N. R. *et al.* Deletions in the survival motor neuron gene on 5q13 in autosomal recessive spinal muscular atrophy. *Hum. Mol. Genet.* **4**, 631–634 (1995).

5. Coovert, D. D. *et al.* The survival motor neuron protein in spinal muscular atrophy. *Hum. Mol. Genet.* **6**, 1205–1214 (1997).

6. Groen, E. J. N. *et al.* Temporal and tissue-specific variability of SMN protein levels in mouse models of spinal muscular atrophy. *Hum. Mol. Genet.* **27**, 2851–2862 (2018).

7. Singh, R. N., Howell, M. D., Ottesen, E. W. & Singh, N. N. Diverse role of Survival Motor Neuron Protein. *Biochim. Biophys. Acta* **1860**, 299–315 (2017).

8. Schrank, B. *et al.* Inactivation of the survival motor neuron gene, a candidate gene for human spinal muscular atrophy, leads to massive cell death in early mouse embryos. *Proc. Natl. Acad. Sci.* **94**, 9920–9925 (1997).

9. Monani, U. R. *et al.* The human centromeric survival motor neuron gene (SMN2) rescues embryonic lethality in Smn(-/-) mice and results in a mouse with spinal muscular atrophy. *Hum. Mol. Genet.* **9**, 333–339 (2000).

10. Rochette, C. F., Gilbert, N. & Simard, L. R. SMN gene duplication and the emergence of the SMN2 gene occurred in distinct hominids: SMN2 is unique to Homo sapiens. *Hum. Genet.* **108**, 255–266 (2001).

11.   Lorson, C. L., Hahnen, E., Androphy, E. J. & Wirth, B. A single nucleotide in the SMN gene

regulates splicing and is responsible for spinal muscular atrophy. *Proc. Natl. Acad. Sci. U. S. A.*

96, 6307–6311 (1999).

12.   Lefebvre, S. *et al.* Correlation between severity and SMN protein level in spinal muscular

atrophy. *Nat. Genet.* **16**, 265–269 (1997).

13.   Hua, Y., Vickers, T. A., Okunola, H. L., Bennett, C. F. & Krainer, A. R. Antisense Masking of

an hnRNP A1/A2 Intronic Splicing Silencer Corrects SMN2 Splicing in Transgenic Mice. *Am. J.*

Hum. Genet. **82**, 834–848 (2008).

14.   Finkel, R. S. *et al.* Nusinersen versus Sham Control in Infantile-Onset Spinal Muscular Atrophy.

N. Engl. J. Med. **377**, 1723–1732 (2017).

15.   Poirier, A. *et al.* Risdiplam distributes and increases SMN protein in both the central nervous

system and peripheral organs. *Pharmacol. Res. Perspect.* **6**, e00447 (2018).

16.   Servais, L. *et al.* FIREFISH Part 2: Efficacy and Safety of Risdiplam (RG7916) in Infants with

Type 1 Spinal Muscular Atrophy (SMA) (1302). *Neurology* **94**, (2020).

17.   Al-Zaidy, S. A. *et al.* AVXS-101 (Onasemnogene Abeparvovec) for SMA1: Comparative Study

with a Prospective Natural History Cohort. *J. Neuromuscul. Dis.* **6**, 307–317 (2019).

18.   Mendell, J. R. *et al.* Single-Dose Gene-Replacement Therapy for Spinal Muscular Atrophy. *N.*

Engl. J. Med. **377**, 1713–1722 (2017).

19.   Mercuri, E. *et al.* Nusinersen versus Sham Control in Later-Onset Spinal Muscular Atrophy. *N.*

Engl. J. Med. **378**, 625–635 (2018).

20.   Dangouloff, T. & Servais, L. Clinical Evidence Supporting Early Treatment Of Patients With

Spinal Muscular Atrophy: Current Perspectives. *Ther. Clin. Risk Manag.* **15**, 1153–1161 (2019).

21. Bowerman, M. *et al.* Therapeutic strategies for spinal muscular atrophy: SMN and beyond. *Dis.*

Model. Mech. **10**, 943–954 (2017).

22. Martínez-Hernández, R., Bernal, S., Alias, L. & Tizzano, E. F. Abnormalities in early markers

of muscle involvement support a delay in myogenesis in spinal muscular atrophy. *J. Neuropathol.*

Exp. Neurol. **73**, 559–567 (2014).

23. Hayhurst, M., Wagner, A. K., Cerletti, M., Wagers, A. J. & Rubin, L. L. A cell-autonomous

defect in skeletal muscle satellite cells expressing low levels of survival of motor neuron protein.

Dev. Biol. **368**, 323–334 (2012).

24. Boyer, J. G. *et al.* Early onset muscle weakness and disruption of muscle proteins in mouse

models of spinal muscular atrophy. *Skelet. Muscle* **3**, 24 (2013).

25. Stevens, L. *et al.* Childhood spinal muscular atrophy induces alterations in contractile and

regulatory protein isoform expressions. *Neuropathol. Appl. Neurobiol.* **34**, 659–670 (2008).

26. Martínez-Hernández, R. *et al.* The developmental pattern of myotubes in spinal muscular

atrophy indicates prenatal delay of muscle maturation. *J. Neuropathol. Exp. Neurol.* **68**, 474–481

(2009).

27. Walter, L. M. *et al.* Interventions Targeting Glucocorticoid-Krüppel-like Factor 15-Branched-

Chain Amino Acid Signaling Improve Disease Phenotypes in Spinal Muscular Atrophy Mice.

EBioMedicine **31**, 226–242 (2018).

28. Place, A. A Phase 2 Study to Evaluate the Efficacy and Safety of SRK-015 in Patients with

Later-Onset Spinal Muscular Atrophy (TOPAZ): An Introduction (4534). *Neurology* **94**, (2020).

29. Rudnicki, S. A. *et al.* Reldesemtiv in Patients with Spinal Muscular Atrophy: a Phase 2

Hypothesis-Generating Study. *Neurother. J. Am. Soc. Exp. Neurother.* **18**, 1127–1136 (2021).

30.   Dickson, M. & Gagnon, J. P. The cost of new drug discovery and development. *Discov. Med.*

4, 172–179 (2004).

31.   Dangouloff, T., Botty, C., Beaudart, C., Servais, L. & Hiligsmann, M. Systematic literature

review of the economic burden of spinal muscular atrophy and economic evaluations of treatments.

Orphanet J. Rare Dis. **16**, 47 (2021).

32.   Talevi, A. & Bellera, C. L. Challenges and opportunities with drug repurposing: finding

strategies to find alternative uses of therapeutics. *Expert Opin. Drug Discov.* **15**, 397–401 (2020).

33.   Cruz-Topete, D. & Cidlowski, J. A. One hormone, two actions: anti- and pro-inflammatory

effects of glucocorticoids. *Neuroimmunomodulation* **22**, 20–32 (2015).

34.   Matthews, E., Brassington, R., Kuntzer, T., Jichi, F. & Manzur, A. Y. Corticosteroids for the

treatment of Duchenne muscular dystrophy. *Cochrane Database Syst. Rev.* CD003725 (2016)

doi:10.1002/14651858.CD003725.pub4.

35.   Ricotti, V. *et al.* Long-term benefits and adverse effects of intermittent versus daily

glucocorticoids in boys with Duchenne muscular dystrophy. *J. Neurol. Neurosurg. Psychiatry* **84**,

698–705 (2013).

36.   McMillan, H. J. Intermittent glucocorticoid regimes for younger boys with duchenne muscular

dystrophy: Balancing efficacy with side effects. *Muscle Nerve* **59**, 638–639 (2019).

37.   Morrison-Nozik, A. *et al.* Glucocorticoids enhance muscle endurance and ameliorate Duchenne

muscular dystrophy through a defined metabolic program. *Proc. Natl. Acad. Sci.* **112**, E6780–

E6789 (2015).

38.   Bulfield, G., Siller, W. G., Wight, P. A. & Moore, K. J. X chromosome-linked muscular

dystrophy (mdx) in the mouse. *Proc. Natl. Acad. Sci. U. S. A.* **81**, 1189–1192 (1984).

39. Oray, M., Abu Samra, K., Ebrahimiadib, N., Meese, H. & Foster, C. S. Long-term side effects

of glucocorticoids. *Expert Opin. Drug Saf.* **15**, 457–465 (2016).

40. Hsieh-Li, H. M. *et al.* A mouse model for spinal muscular atrophy. *Nat. Genet.* **24**, 66–70

(2000).

41. Le, T. T. *et al.* SMNDelta7, the major product of the centromeric survival motor neuron (SMN2)

gene, extends survival in mice with spinal muscular atrophy and associates with full-length SMN.

Hum. Mol. Genet. **14**, 845–857 (2005).

42. Meijboom, K. E. *et al.* Combining multiomics and drug perturbation profiles to identify muscle-

specific treatments for spinal muscular atrophy. *JCI Insight* **6**, 149446 (2021).

43. Bowerman, M., Murray, L. M., Beauvais, A., Pinheiro, B. & Kothary, R. A critical smn

threshold in mice dictates onset of an intermediate spinal muscular atrophy phenotype associated

with a distinct neuromuscular junction pathology. *Neuromuscul. Disord. NMD* **22**, 263–276

(2012).

44. Nguyen, A. T., Armstrong, E. A. & Yager, J. Y. Neurodevelopmental Reflex Testing in

Neonatal Rat Pups. *J. Vis. Exp. JoVE* 55261 (2017) doi:10.3791/55261.

45. Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and collaborative biomedical

analyses: 2018 update. *Nucleic Acids Res.* **46**, W537–W544 (2018).

46. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence

Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

47. Trimmomatic: a flexible trimmer for Illumina sequence data | Bioinformatics | Oxford

Academic. https://academic.oup.com/bioinformatics/article/30/15/2114/2390096.

48. FASTX-Toolkit. http://hannonlab.cshl.edu/fastx_toolkit/.

49.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019).

50.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinforma. Oxf. Engl.* **30**, 923–930 (2014).

51.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for

multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

52.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

53.   Ahsan, S. & Drăghici, S. Identifying Significantly Impacted Pathways and Putative

Mechanisms with iPathwayGuide. *Curr. Protoc. Bioinforma.* **57**, 7.15.1-7.15.30 (2017).

54.   Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives

on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**, D353–D361 (2017).

55.   Draghici, S., Khatri, P., Martins, R. P., Ostermeier, G. C. & Krawetz, S. A. Global functional

profiling of gene expression. *Genomics* **81**, 98–104 (2003).

56.   Jantzen, S. G., Sutherland, B. J., Minkley, D. R. & Koop, B. F. GO Trimming: Systematically

reducing redundancy in large Gene Ontology datasets. *BMC Res. Notes* **4**, 267 (2011).

57.   The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing

strong. *Nucleic Acids Res.* **47**, D330–D338 (2019).

58.   Cotto, K. C. *et al.* DGIdb 3.0: a redesign and expansion of the drug–gene interaction database.

Nucleic Acids Res. **46**, D1068–D1073 (2018).

59.   Yaffe, D. & Saxel, O. Serial passaging and differentiation of myogenic cells isolated from

dystrophic mouse muscle. *Nature* **270**, 725–727 (1977).

60. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT–PCR.

Nucleic Acids Res. **29**, e45 (2001).

61. Radonić, A. *et al.* Guideline to reference gene selection for quantitative real-time PCR.

Biochem. Biophys. Res. Commun. **313**, 856–862 (2004).

62. Untergasser, A., Ruijter, J. M., Benes, V. & van den Hoff, M. J. B. Web-based LinRegPCR:

application for the visualization and analysis of (RT)-qPCR amplification and melting data. *BMC*

Bioinformatics **22**, 398 (2021).

63. Briese, M. *et al.* Deletion of smn-1, the Caenorhabditis elegans ortholog of the spinal muscular

atrophy gene, results in locomotor dysfunction and reduced lifespan. *Hum. Mol. Genet.* **18**, 97–

104 (2009).

64. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

65. Dimitriadi, M. *et al.* Decreased function of survival motor neuron protein impairs endocytic

pathways. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E4377-4386 (2016).

66. Deguise, M.-O. *et al.* Differential induction of muscle atrophy pathways in two mouse models

of spinal muscular atrophy. *Sci. Rep.* **6**, 28846 (2016).

67. Simon, C. M. *et al.* Converging Mechanisms of p53 Activation Drive Motor Neuron

Degeneration in Spinal Muscular Atrophy. *Cell Rep.* **21**, 3767–3780 (2017).

68. Cerveró, C. *et al.* Chronic Treatment with the AMPK Agonist AICAR Prevents Skeletal Muscle

Pathology but Fails to Improve Clinical Outcome in a Mouse Model of Severe Spinal Muscular

Atrophy. *Neurother. J. Am. Soc. Exp. Neurother.* **13**, 198–216 (2016).

69. Ripolone, M. *et al.* Impaired Muscle Mitochondrial Biogenesis and Myogenesis in Spinal

Muscular Atrophy. *JAMA Neurol.* **72**, 666–675 (2015).

70.   Walter, L. M. *et al.* Light modulation ameliorates expression of circadian genes and disease

progression in spinal muscular atrophy mice. *Hum. Mol. Genet.* **27**, 3582–3597 (2018).

71.   Ng, S. Y., Mikhail, A. & Ljubicic, V. Mechanisms of exercise-induced survival motor neuron

expression in the skeletal muscle of spinal muscular atrophy-like mice. *J. Physiol.* **597**, 4757–4778

(2019).

72.   Sansa, A. *et al.* Spinal Muscular Atrophy autophagy profile is tissue-dependent: differential

regulation between muscle and motoneurons. *Acta Neuropathol. Commun.* **9**, 122 (2021).

73.   Pin, F., Couch, M. E. & Bonetto, A. Preservation of muscle mass as a strategy to reduce the

toxic effects of cancer chemotherapy on body composition. *Curr. Opin. Support. Palliat. Care* **12**,

420–426 (2018).

74.   Farooq, F. *et al.* Celecoxib increases SMN and survival in a severe spinal muscular atrophy

mouse model via p38 pathway activation. *Hum. Mol. Genet.* **22**, 3415–3424 (2013).

75.   Mack, S. G., Cook, D. J., Dhurjati, P. & Butchbach, M. E. R. Systems biology investigation of

cAMP modulation to increase SMN levels for the treatment of spinal muscular atrophy. *PloS One*

9, e115473 (2014).

76.   Rojas, L. B. A. & Gomes, M. B. Metformin: an old but still the best treatment for type 2 diabetes.

Diabetol. Metab. Syndr. **5**, 6 (2013).

77.   Orr, R. & Fiatarone Singh, M. The anabolic androgenic steroid oxandrolone in the treatment of

wasting and catabolic disorders: review of efficacy and safety. *Drugs* **64**, 725–750 (2004).

78.   Soliman, A., De Sanctis, V., Alaaraj, N. & Hamed, N. The clinical application of metformin in

children and adolescents: A short update. *Acta Bio Medica Atenei Parm.* **91**, e2020086 (2020).

79.   Hafner, P. *et al.* Effect of Combination l-Citrulline and Metformin Treatment on Motor Function

in Patients With Duchenne Muscular Dystrophy: A Randomized Clinical Trial. *JAMA Netw. Open*

2, e1914171 (2019).

80.   Fontes-Oliveira, C. C., M. Soares Oliveira, B., Körner, Z., M. Harandi, V. & Durbeej, M. Effects

of metformin on congenital muscular dystrophy type 1A disease progression in mice: a gender

impact study. *Sci. Rep.* **8**, 16302 (2018).

81.   Birk, J. B. & Wojtaszewski, J. F. P. Predominant α2/β2/γ3 AMPK activation during exercise in

human skeletal muscle. *J. Physiol.* **577**, 1021–1032 (2006).

82.   Zhang, L. *et al.* Role and mechanism underlying FoxO6 in skeletal muscle in vitro and in vivo.

Int. J. Mol. Med. **48**, 143 (2021).

83.   Everaert, C. *et al.* Benchmarking of RNA-sequencing analysis workflows using whole-

transcriptome RT-qPCR expression data. *Sci. Rep.* **7**, 1559 (2017).

84.   Shafey, D., Côté, P. D. & Kothary, R. Hypomorphic Smn knockdown C2C12 myoblasts reveal

intrinsic defects in myoblast fusion and myotube morphology. *Exp. Cell Res.* **311**, 49–61 (2005).

85.   Millino, C. *et al.* Different atrophy-hypertrophy transcription pathways in muscles affected by

severe and mild spinal muscular atrophy. *BMC Med.* **7**, 14 (2009).

86.   Calura, E. *et al.* Meta-analysis of expression signatures of muscle atrophy: gene interaction

networks in early and late stages. *BMC Genomics* **9**, 630 (2008).

87.   Mirza, K. A., Pereira, S. L., Edens, N. K. & Tisdale, M. J. Attenuation of muscle wasting in

murine C2C 12 myotubes by epigallocatechin-3-gallate. *J. Cachexia Sarcopenia Muscle* **5**, 339–

345 (2014).

88.   Rivera, M. E., Lyon, E. S. & Vaughan, R. A. Effect of metformin on myotube BCAA

catabolism. *J. Cell. Biochem.* **121**, 816–827 (2020).

89. Bowerman, M. *et al.* Glucose Metabolism and Pancreatic Defects in Spinal Muscular Atrophy.

Ann. Neurol. **72**, 256–268 (2012).

90. Deguise, M.-O. *et al.* Abnormal fatty acid metabolism is a core component of spinal muscular

atrophy. *Ann. Clin. Transl. Neurol.* **6**, 1519–1532 (2019).

91. Boyer, J. G. *et al.* Myogenic program dysregulation is contributory to disease pathogenesis in

spinal muscular atrophy. *Hum. Mol. Genet.* **23**, 4249–4259 (2014).

92. Bruce, A. K., Jacobsen, E., Dossing, H. & Kondrup, J. Hypoglycaemia in spinal muscular

atrophy. *Lancet Lond. Engl.* **346**, 609–610 (1995).

93. Djordjevic, S. A. *et al.* Glucose and lipid metabolism disorders in children and adolescents with

spinal muscular atrophy types 2 and 3. *Neuromuscul. Disord. NMD* **31**, 291–299 (2021).

94. Birnbaum, M. J. Identification of a novel gene encoding an insulin-responsive glucose

transporter protein. *Cell* **57**, 305–315 (1989).

95. DeWaal, D. *et al.* Hexokinase-2 depletion inhibits glycolysis and induces oxidative

phosphorylation in hepatocellular carcinoma and sensitizes to metformin. *Nat. Commun.* **9**, 446

(2018).

96. Owen, M. R., Doran, E. & Halestrap, A. P. Evidence that metformin exerts its anti-diabetic

effects through inhibition of complex 1 of the mitochondrial respiratory chain. *Biochem. J.* **348 Pt**

3, 607–614 (2000).

97. Graham, G. G. *et al.* Clinical pharmacokinetics of metformin. *Clin. Pharmacokinet.* **50**, 81–98

(2011).

98. Łabuzek, K. *et al.* Quantification of metformin by the HPLC method in brain regions,

cerebrospinal fluid and plasma of rats treated with lipopolysaccharide. *Pharmacol. Rep. PR* **62**,

956–965 (2010).

99. Singh, J., Olle, B., Suhail, H., Felicella, M. M. & Giri, S. Metformin-induced mitochondrial

function and ABCD2 up-regulation in X-linked adrenoleukodystrophy involves AMP-activated

protein kinase. *J. Neurochem.* **138**, 86–100 (2016).

100. Fenichel, G., Pestronk, A., Florence, J., Robison, V. & Hemelt, V. A beneficial effect of

oxandrolone in the treatment of Duchenne muscular dystrophy: a pilot study. *Neurology* **48**, 1225–

1226 (1997).

101. Hart, D. W. *et al.* Anabolic Effects of Oxandrolone After Severe Burn. *Ann. Surg.* **233**, 556–

564 (2001).

102. Ren, H., Yin, P. & Duan, C. IGFBP-5 regulates muscle cell differentiation by binding to IGF-

II and switching on the IGF-II auto-regulation loop. *J. Cell Biol.* **182**, 979–991 (2008).

103. Ganassi, M., Badodi, S., Wanders, K., Zammit, P. S. & Hughes, S. M. Myogenin is an essential

regulator of adult myofibre growth and muscle stem cell homeostasis. *eLife* **9**, e60445 (2020).

104. Cai, D., Dhe-Paganon, S., Melendez, P. A., Lee, J. & Shoelson, S. E. Two new substrates in

insulin signaling, IRS5/DOK4 and IRS6/DOK5. *J. Biol. Chem.* **278**, 25323–25330 (2003).

105. Lee, S.-W. *et al.* AKAP6 inhibition impairs myoblast differentiation and muscle regeneration:

Positive loop between AKAP6 and myogenin. *Sci. Rep.* **5**, 16523 (2015).

106. Wu, Y. *et al.* REDD1 is a major target of testosterone action in preventing dexamethasone-

induced muscle loss. *Endocrinology* **151**, 1050–1059 (2010).

107. Zeman, R. J. *et al.* Improved functional recovery with oxandrolone after spinal cord injury in

rats. *NeuroReport* **20**, 864–868 (2009).

108. Ahmad, A., Herndon, D. N. & Szabo, C. Oxandrolone protects against the development of

multiorgan failure, modulates the systemic inflammatory response and promotes wound healing

during burn injury. *Burns J. Int. Soc. Burn Inj.* **45**, 671–681 (2019).

109. Bertrandy, S. *et al.* The RNA-Binding Properties of SMN: Deletion Analysis of the Zebrafish

Orthologue Defines Domains Conserved in Evolution. *Hum. Mol. Genet.* **8**, 775–782 (1999).

110. Nilsson, E. C. *et al.* Opposite transcriptional regulation in skeletal muscle of AMP-activated

protein kinase gamma3 R225Q transgenic versus knock-out mice. *J. Biol. Chem.* **281**, 7244–7252

(2006).

111. Barnes, B. R. *et al.* Changes in exercise-induced gene expression in 5'-AMP-activated protein

kinase gamma3-null and gamma3 R225Q transgenic mice. *Diabetes* **54**, 3484–3489 (2005).

112. Marangos, P. J. *et al.* Adenosinergic Modulation of Homocysteine-Induced Seizures in Mice.

Epilepsia **31**, 239–246 (1990).

113. Corton, J. M., Gillespie, J. G., Hawley, S. A. & Hardie, D. G. 5-aminoimidazole-4-carboxamide

ribonucleoside. A specific method for activating AMP-activated protein kinase in intact cells? *Eur.*

J. Biochem. **229**, 558–565 (1995).

114. Xiao, B. *et al.* Structural basis for AMP binding to mammalian AMP-activated protein kinase.

Nature **449**, 496–500 (2007).

115. Thelen, M. P., Wirth, B. & Kye, M. J. Mitochondrial defects in the respiratory complex I

contribute to impaired translational initiation via ROS and energy homeostasis in SMA motor

neurons. *Acta Neuropathol. Commun.* **8**, 223 (2020).

116. Borkowska, A. *et al.* Coexistence of type 1 diabetes mellitus and spinal muscular atrophy in an

8-year-old girl: a case report. *Acta Biochim. Pol.* **62**, 167–168 (2015).

117. Hossain, S. & Chao, J. MON-154 Abnormal Glucose Homeostasis in Spinal Muscular Atrophy

(SMA) Leading to a Transient Episode of Diabetic Ketoacidosis (DKA). *J. Endocr. Soc.* **3**, MON-

154 (2019).

118. MacLean, H. E. *et al.* Impaired skeletal muscle development and function in male, but not

female, genomic androgen receptor knockout mice. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.*

22, 2676–2689 (2008).

119. Morton, R. W. *et al.* Muscle Androgen Receptor Content but Not Systemic Hormones Is

Associated With Resistance Training-Induced Skeletal Muscle Hypertrophy in Healthy, Young

Men. *Front. Physiol.* **9**, (2018).

120. Liu, Y., Beyer, A. & Aebersold, R. On the Dependency of Cellular Protein Levels on mRNA

Abundance. *Cell* **165**, 535–550 (2016).

121. Knorre, D. G., Kudryashova, N. V. & Godovikova, T. S. Chemical and functional aspects of

posttranslational modification of proteins. *Acta Naturae* **1**, 29–51 (2009).

122. Kessler, T. *et al.* Cerebrospinal fluid proteomic profiling in nusinersen-treated patients with

spinal muscular atrophy. *J. Neurochem.* **153**, 650–661 (2020).

123. Gardell, S. J., Zhang, X., Kapoor, N., Petucci, C. & Coen, P. M. Metabolomics Analyses of

Muscle Atrophy Induced by Hind Limb Unloading. *Methods Mol. Biol. Clifton NJ* **1996**, 297–309

(2019).

124. Finkel, R. S. *et al.* Candidate proteins, metabolites and transcripts in the Biomarkers for Spinal

Muscular Atrophy (BforSMA) clinical study. *PloS One* **7**, e35462 (2012).

125. De Micheli, A. J., Spector, J. A., Elemento, O. & Cosgrove, B. D. A reference single-cell

transcriptomic atlas of human skeletal muscle tissue reveals bifurcated muscle stem cell

populations. *Skelet. Muscle* **10**, 19 (2020).

126. Williams, K., Yokomori, K. & Mortazavi, A. Heterogeneous Skeletal Muscle Cell and Nucleus

Populations Identified by Single-Cell and Single-Nucleus Resolution Transcriptome Assays.

Front. Genet. **13**, (2022).

127. Habets, L. E. *et al.* Magnetic resonance reveals mitochondrial dysfunction and muscle

remodelling in spinal muscular atrophy. *Brain* **145**, 1422–1435 (2022).

128. Saleh, K. K. *et al.* Single cell sequencing maps skeletal muscle cellular diversity as disease

severity increases in dystrophic mouse models. *iScience* **25**, 105415 (2022).

129. Chemello, F. *et al.* Degenerative and regenerative pathways underlying Duchenne muscular

dystrophy revealed by single-nucleus RNA sequencing. *Proc. Natl. Acad. Sci.* **117**, 29691–29701

(2020).

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

 FIGURE LEGENDS

Figure 1. Prednisolone treatment normalizes a subset of genes in severe *Smn$^{-/-}$;SMN2* SMA

mice to healthy levels observed in untreated *Smn$^{+/-}$;SMN2* mice.

Smn$^{-/-}$;SMN2 SMA and *Smn$^{+/-}$;SMN2* healthy mice received prednisolone treatment (5 mg/kg

gavage every 2 days) from P0. The triceps was harvested from P7 untreated and prednisolone-

treated *Smn$^{-/-}$;SMN2* SMA and *Smn$^{+/-}$;SMN2* healthy mice for RNA isolation and library

preparation for RNA-Sequencing. Differential gene expression analysis was performed by

DESeq2 v2.11.40.2 with study design set to "condition and "treatment". (**a**) Principal component

analysis based on transcriptomic profiles between P7 untreated *Smn$^{+/-}$;SMN2* (red, n=3),

prednisolone-treated *Smn$^{-/-}$;SMN2* (green, n=2) and untreated *Smn$^{-/-}$;SMN2* (blue, n=3) mice. (**b**)

Heatmap of the transcriptomic expression profiles (Log2FC >0.6; FDR <0.05) between P7

untreated *Smn$^{+/-}$;SMN2* (n=3, left), untreated *Smn$^{-/-}$;SMN2* (n=3, centre) and prednisolone-treated

Smn$^{-/-}$;SMN2 (n=3, right) mice. Upregulated genes highlighted in red and downregulated genes

highlighted in blue.

1115

Figure 2. The metformin target gene, *Prkag3,* is significantly downregulated in the skeletal

muscle of both symptomatic severe *Smn$^{-/-}$;SMN2* and intermediate *Smn$^{2B/-}$* SMA mice.

qPCR analysis of mRNA levels for predicted metformin target genes **a.** *Prkag3*, **b.** *Foxo1*, **c.**

Foxo3, **d.** *Foxo4* and **e.** *Foxo6* in the harvested triceps of symptomatic untreated P7 Taiwanese

Smn$^{-/-}$;SMN2 SMA mice (violet) and healthy *Smn$^{+/-}$;SMN2* controls (black) (left panel) and

symptomatic untreated P19 intermediate *Smn$^{2B/-}$* SMA mice (blue) and wild type (C57BL/6J

background) controls (black) (right panel). Data are shown as scatter plot represent as mean ±

SEM error bars; n = 4-7 animals per experimental group, unpaired t-test, *p <0.05. *Smn$^{-/-}$;SMN2*

Prkag3: $p = 0.04$; *Smn$^{-/-}$;SMN2 Foxo1*: $p = 0.82$; *Smn$^{-/-}$;SMN2 Foxo3*: $p = 0.54$; *Smn$^{-/-}$;SMN2*

Foxo4: $p = 0.37$; *Smn$^{-/-}$;SMN2 Foxo6*: $p = 0.34$; *Smn$^{2B/-}$ Prkag3*: $p = 0.02$; *Smn$^{2B/-}$ Foxo1*: $p = 0.39$;

Smn$^{2B/-}$ Foxo3: $p = 0.27$; *Smn$^{2B/-}$ Foxo4*: $p = 0.48$; *Smn$^{2B/-}$ Foxo6*: $p = 0.33$.

1127

Figure 3. Metformin target genes are pre-dominantly SMN-independent in an SMA muscle

C2C12 cellular model.

Smn siRNA knockdown (red) was performed for **a.** 48 hours in C2C12 myoblasts and **b.** every 48

hours throughout differentiation in D8 C2C12 myotubes. mRNA expression of metformin target

genes *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4* and *Foxo6* was measured by qPCR and compared to non-

transfected (black) and scrambled siRNA transfected controls (blue). D8 C2C12 myotubes were

serum-starved for 24 hours to induce canonical atrophy (red). mRNA expression of **c.** atrogenes

Atrogin-1 and *MuRF-1* and **d.** Metformin target genes *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4* and *Foxo6*

was measured by qPCR and compared against non-starved myotubes (black). Serum-starved D8

C2C12 myotubes were treated with either physiological 30 µM metformin (red) or

supraphysiological 2 mM metformin (blue) for 24 hours to evaluate mRNA expression via qPCR

of **e.** atrogenes *Atrogin-1* and *MuRF-1* and **f.** Metformin target genes *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4*

and *Foxo6* compared to serum-starved PBS vehicle treated control (black). Data are shown as

scatter plot that represent mean ± SEM error bars; n = 4 samples per group across two independent

experiments. Two-way ANOVA followed by uncorrected Fisher's least significant difference

(LSD). F = 3.543 (**a.**); F = 2.332 (**b.**); F = 4.9 (**c.**); F = 3.493 (**d.**); F = 0.057 (**e.**); F = 0.235 (**f.**); F

= 0.401, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 4. 200 mg/kg/day metformin does not improve disease phenotype, while 400

mg/kg/day metformin reduces survival.

All treated animals received a daily dose of metformin (either 200 or 400 mg/kg/day, diluted in

0.9% saline) by gavage starting at P5. **a.** Survival curves of untreated (n = 13, 21 days median

survival, black) and 200 mg/kg/day metformin-treated (n = 11, 21 days median survival, red)

$Smn^{2B/-}$ SMA mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test, ns = not

significant, $p = 0.237$. **b.** Daily weights of untreated (n = 13, black) and 200 mg/kg/day metformin-

treated (n = 11, red) $Smn^{2B/-}$ SMA mice. Data represented as mean ± SEM error bars; Two-way

ANOVA followed by a Sidak's multiple comparison test, F = 402.1, df = 455, *$p < 0.05$, **$p <$

0.01, ***$p < 0.001$, ****$p < 0.0001$. **c.** Daily righting reflex test for motor function activity up to

a 30 second maximum time point in untreated (n = 13, black) and 200 mg/kg/day metformin-

treated (n = 11, red) $Smn^{2B/-}$ SMA mice. Data are shown as bar chart with mean ± SEM error bars;

unpaired T-test, ns = not significant, $p = 0.833$. **d.** Survival curves of untreated (n = 13, 21 days

median survival, black) and 400 mg/kg/day metformin-treated (n = 4, 16 days median survival,

blue) $Smn^{2B/-}$ SMA mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test,

****$p < 0.0001$. **e.** Daily weights of untreated (n = 13, black) and 400 mg/kg/day metformin-

treated (n = 9, blue) $Smn^{2B/-}$ SMA mice. Data represented as mean ± SEM error bars; Two-way

ANOVA followed by a Sidak's multiple comparison test, F = 184.9.1, df = 300. **f.** Daily righting

reflex test for motor function activity up to a 30 second maximum time point in untreated (n = 13,

black) and 400 mg/kg/day metformin-treated (n = 9, blue) $Smn^{2B/-}$ SMA mice. Data are shown as

bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, $p = 0.733$.

1166

1167 **Figure 5. 400 mg/kg/day metformin significantly lowers blood-glucose levels in $Smn^{2B/-}$ SMA**

1168 **mice, with no impact on markers of atrophy, glucose metabolism and mitochondrial**

1169 **regulation in skeletal muscle.**

a. Blood-glucose concentrations (mmol/L) were measured 2 hours after final treatment from untreated (black) and 200 (red) or 400 mg/kg/day metformin-treated (blue), non-fasted P14 $Smn^{2B/+}$ healthy and $Smn^{2B/-}$ SMA mice. Data represented as bar chart with scatter graph represented as mean ± SEM error bars, n = 4 animals per group; two-way ANOVA with Tukey's multiple comparisons test, F = 25.49, * p <0.05. qPCR analysis of predicted metformin target genes **b.** *Prkag3*, **c.** *Foxo1*, **d.** *Foxo3*, **e.** *Foxo4*, **f.** *Foxo6*; atrogenes **g.** *Atrogin-1*, **h.** *MuRF1*; and glucose uptake and metabolism genes **i.** *Glut4*, **j.** *Hk2*; and mitochondrial regulatory genes **k.** *Pgc1-α,* **l.** *Tfam,* **m.** *Ndufs1* and **n.** *Nrf1* in the TA muscle from untreated and 200 or 400 mg/kg/day metformin-treated, P14 $Smn^{2B/+}$ healthy and $Smn^{2B/-}$ SMA mice. Data are shown as scatter graph represented as mean ± SEM error bars, n = 4 animals per group; two way ANOVA with Tukey's multiple comparisons test, *Prkag3* (F = 59.92), *Foxo1* (F = 1.507), *Foxo3* (F = 0.343), *Foxo4* (F = 6.475), *Foxo6* (F = 0.024), *Atrogin-1* (F = 0.381), *MuRF1* (F = 7.838), *Glut4* (F = 9.9), *Hk2* (F = 17.78), *Pgc1-α* (F = 29.84), *Tfam* (F = 0.423) *Ndufs1* (F = 22.66), and *Nrf1* (F = 0.164).

Figure 6. 400 mg/kg/day metformin dysregulates mitochondrial regulatory genes exclusively in spinal cord tissue from Smn$^{2B/-}$ SMA mice.

qPCR analysis of mitochondrial regulatory genes **a.** *Pgc1-α,* **b.** *Ndufs1,* **c.** *Nrf1* and **d.** *Tfam* in the spinal cord from untreated (black) and 200 (red) or 400 mg/kg/day metformin-treated (blue), P14 $Smn^{2B/+}$ healthy and $Smn^{2B/-}$ SMA mice. Data are shown as scatter graph represented as mean ± SEM error bars, n = 4 animals per group; two-way ANOVA with Tukey's multiple comparisons test. **a.** *Pgc1-α* (F = 1.526), **b.** *Ndufs1* (F = 1.135), **c.** *Nrf1* (F = 0.362) and **d.** *Tfam* (F = 0.614), *p < 0.05, **p < 0.01.

Figure 7. The oxandrolone target gene, *Ddit4*, is significantly upregulated in the skeletal

muscle of severe *Smn$^{-/-}$;SMN2* SMA mice.

qPCR analysis of mRNA levels for predicted oxandrolone target genes **a.** *Ar*, **b.** *Akap6*, **c.** *Igfbp5*,

d. *Dok5,* **e.** *MyoG* and **f.** *Ddit4* in the harvested triceps of untreated P7 Taiwanese *Smn$^{-/-}$;SMN2*

SMA mice (violet) and healthy *Smn$^{+/-}$;SMN2* controls (black) (left panel) and symptomatic

untreated P19 intermediate *Smn$^{2B/-}$* SMA mice (blue) and wild type (C57BL/6J background)

controls (black) (right panel). Data are shown as scatter plot that represents mean ± SEM error

bars; n = 4-7 animals per experimental group, unpaired t-test, **p* <0.05. *Smn$^{-/-}$;SMN2 Ar*: *p* = 0.38;

Smn$^{-/-}$;SMN2 Akap6: *p* = 0.68; *Smn$^{-/-}$;SMN2 Igfbp5*: *p* = 0.49; *Smn$^{-/-}$;SMN2 Dok5*: *p* = 0.79; *Smn$^{-/-}$*

;SMN2 MyoG: *p* = 0.64; *Smn$^{-/-}$;SMN2 Ddit4*: *p* = 0.02; *Smn$^{2B/-}$ Ar*: *p* = 0. 75; *Smn$^{2B/-}$ Akap6*: *p* = 0.

29; *Smn$^{2B/-}$ Igfbp5*: *p* = 0.52; *Smn$^{2B/-}$ Dok5*: *p* = 0.19; *Smn$^{2B/-}$ MyoG*: *p* = 0.15, *Smn$^{2B/-}$ Ddit4*: *p* =

0.16.

1205

Figure 8. Oxandrolone target genes are pre-dominantly SMN-independent in SMA muscle

C2C12 cellular model.

Smn siRNA knockdown (red) was performed for **a.** 48 hours in C2C12 myoblasts and **b.** every 48

hours throughout differentiation in D8 C2C12 myotubes. mRNA expression of oxandrolone target

genes *Ar*, *Akap6*, *Igfbp5*, *Dok5, MyoG* and *Ddit4* was measured by qPCR and compared to non-

transfected (black) and scrambled siRNA transfected (blue) controls. D5 C2C12 myotubes were

serum-starved for 24 hours to induce canonical atrophy (red). mRNA expression of **c.** atrogenes

Atrogin-1 and *MuRF-1* and **d.** oxandrolone target genes *Ar*, *Akap6*, *Igfbp5*, *Dok5, MyoG* and *Ddit4*

was measured by qPCR and compared to non-starved myotubes (black). Serum-starved D5 C2C12

myotubes were treated with1 µM oxandrolone for 24 hours (blue) to evaluate mRNA expression

via qPCR of **e.** atrogenes *Atrogin-1* and *MuRF-1* and **f.** oxandrolone target genes *Ar, Akap6, Igfbp5,*

Dok5, MyoG and *Ddit4* compared to serum-starved absolute ethanol vehicle treated control

(black). Data are shown as scatter graphs that represent mean ± SEM error bars; n = 4 samples per

group across two independent experiments. Two-way ANOVA followed by uncorrected Fisher's

least significant difference (LSD). **a.** F = 5.45; **b.** F = 6.87; **c.** F = 1.1; **d.** F = 2.03; **e.** F = 0.72; **f.**

F = 0.36, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

1222

Figure 9. 4 mg/kg/day oxandrolone treatment partially improves survival in *Smn$^{2B/-}$* SMA

mice.

All treated animals received a daily dose of oxandrolone (4 mg/kg/day, suspended in 0.5% CMC)

by gavage starting at P8. **a.** Survival curves of untreated (n = 15, 21 days median survival, black)

and 4 mg/kg/day oxandrolone-treated (n = 12, 24 days median survival, orange) *Smn$^{2B/-}$* SMA

mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test, , ***$p = 0.0006$. **b.**

Daily weights of untreated (n = 15, black) and 4 mg/kg/day oxandrolone-treated (n = 12, orange)

Smn$^{2B/-}$ SMA mice. Data represented as mean ± SEM error bars; Two-way ANOVA followed by

a Sidak's multiple comparison test, F = 610.8, df = 519, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

****$p < 0.0001$. **c.** Daily righting reflex test for motor function activity up to a 30 second maximum

time point in untreated (n = 15, black) and 4 mg/kg/day oxandrolone-treated (n = 12, orange)

Smn$^{2B/-}$ SMA mice. Data are shown as bar chart with mean ± SEM error bars; unpaired T-test, ns

= not significant, *$p = 0.775$*. qPCR analysis of mRNA levels for atrogenes **d.** *Atrogin-1* and **e.**

MuRF1 and predicted target genes **f.** *Ar,* **g.** *Akap6,* **h.** *Igfbp5,* **i.** *Dok5,* **j.** *MyoG,* and **k.** *Ddit4* in the

triceps muscle from untreated (black) and 4 mg/kg/day oxandrolone-treated (orange), P19 *Smn$^{2B/+}$*

healthy and *Smn$^{2B/-}$* SMA mice. Data are shown as bar chart with scatter graph represented as mean

± SEM error bars, n = 4 animals per group; two way ANOVA with Tukey's multiple comparisons

test, *Atrogin-1* (F = 0.1914), *MuRF1* (F = 1.214) , *Ar* (F = 0.003), *Akap6* (F = 2.40), *Igfbp5* (F =

0.06), *Dok5* (F = 1.72), *MyoG* (F = 0.29) and *Ddit4* (F = 0.64), *$p < 0.05$, **$p < 0.01$.

1242

Figure 10. Metformin and oxandrolone partially ameliorates neuromuscular defects in the

severe SMA *C. elegans* model.

Day 3 *C. elegans smn-1(ok355)* SMA homozygotes and *smn-1(ok355)I/hT2* control heterozygotes

were maintained at 20 °C on nematode growth medium (NGM) plates seeded with Escherichia coli

OP50 bacteria. In metformin conditions, the NGM contained metformin concentrations of 0 (black

square), 1 (blue circle), 10 (red triangle) and 50 mM (orange diamond) respectively. **a.** Pharyngeal

pumping rates (pumps/minute) defined as grinder movement in any axis at 175 frames/10 seconds.

Locomotion assays were filmed at 15 frames/sec and quantified for 5 minutes for **b.** reversals and

c. paralysis times in the *C. elegans* groups. In oxandrolone conditions, the NGM contained

oxandrolone concentrations of 0 (black square), 1 (blue circle), 10 (red triangle) and 50 μM (orange

diamond) respectively. **d.** Pharyngeal pumping rates (pumps/minute) defined as grinder movement

in any axis at 175 frames/10 seconds. Locomotion assays were filmed at 15 frames/sec and were

quantified for 5 minutes for **e.** reversals and **f.** paralysis times in the *C. elegans* groups. Data are

shown as scatter plot that represents mean ± SEM error bars; n > 25 animals per experimental

group across three independent trials; Two-way ANOVA with Tukey's multiple comparisons test,

a. F = 459.2, **b.** F = 108.6, **c.** F = 52.59, **d.** F = 501.8, **e.** F = 121.1, **f.** F = 57.06, ns = not significant

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

1260

1261

1262 **TABLES**

1263 **Table 1. KEGG pathways targeted in the skeletal muscle of symptomatic prednisolone-**

1264 **treated *Smn⁻/⁻;SMN2* SMA mice compared with untreated *Smn⁻/⁻;SMN2* SMA mice.**

1265

1266 **Table 2. Top Gene Ontology Biological Process pathways targeted in the skeletal muscle of**

1267 **symptomatic prednisolone-treated *Smn⁻/⁻;SMN2* SMA mice compared with untreated *Smn⁻/⁻***

1268 **;*SMN2* SMA mice.**

1269

1270 **Table 3. Top 10 clinically approved drugs identified by KEGG database based on**

1271 **prednisolone-targeted KEGG pathways in symptomatic prednisolone-treated *Smn⁻/⁻;SMN2***

1272 **SMA mice.**

1273

1274 **Table 4. Top 10 clinically approved drugs identified by DGIdb database based on**

1275 **prednisolone-targeted KEGG pathways in symptomatic prednisolone-treated *Smn⁻/⁻;SMN2***

1276 **SMA mice.**

1277

1278

1279

1280

1281

1282

1283

1284

SUPPLEMENTARY FIGURE LEGENDS

Figure S1. "Condition" and "Treatment" groups share distinct transcriptomic patterns for a specific sub-set of genes in prednisolone-treated *Smn$^{-/-}$;SMN2* SMA mice and untreated *Smn$^{-/-}$;SMN2* SMA and *Smn$^{+/-}$;SMN2* healthy mice.

Smn$^{-/-}$;SMN2 SMA and *Smn$^{+/-}$;SMN2* healthy mice received prednisolone treatment (5 mg/kg gavage every 2 days) from P0. The triceps was harvested from P7 untreated and prednisolone-treated *Smn$^{-/-}$;SMN2* SMA and *Smn$^{+/-}$;SMN2* healthy mice for RNA isolation and library preparation for RNA-Sequencing. Differential gene expression analysis was performed by DESeq2 v2.11.40.2 with study design set to "condition and "treatment". Principal component analysis based on transcriptomic profiles between **a.** P7 untreated *Smn$^{+/-}$;SMN2* (red, n=3) and untreated *Smn$^{-/-}$;SMN2* (blue, n=3) mice. **b.** P7 untreated *Smn$^{-/-}$;SMN2* (green, n=3) and prednisolone-treated *Smn$^{-/-}$;SMN2* (blue, n=2) SMA mice**.**

Figure S2. Bioinformatic identification of shared target genes between metformin and prednisolone in SMA skeletal muscle.

a. The pathway diagram contains proteins within the FOXO signalling pathway (KEGG: 04068) encoded by the predicted differentially expressed genes from prednisolone vs untreated *Smn$^{-/-}$;SMN2* SMA skeletal muscle. The AMPK protein (yellow circle) represents the AMPK-γ3 isoform gene *Prkag3*. Its activity on the iPathwayGuide interactive server highlights that it can be targeted by metformin via a built-in KEGG Drugs database. The red lines represent coherent cascades, which supports the consistency of the transcriptomic data and published pathway activity. In the FOXO signalling pathway, activation of AMPK represented by *Prkag3* coherently downregulates the FOXO protein (green circle), which represents *Foxo1*, *Foxo3*, *Foxo4*, and *Foxo6* isoforms. The

highest logFC patterns are shown in dark red and lowest in dark blue as indicated in the legend

value box. Graph generated in iPathwayGuide (Advaita). **b.** Differential gene expression pattern

by logFC (Y-axis) of predicted metformin target genes *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4* and *Foxo6*

based on transcriptomic data from prednisolone vs untreated *Smn$^{-/-}$;SMN2* SMA skeletal muscle.

Upregulated genes above X-axis are highlighted in red and downregulated genes below X-axis are

highlighted in blue. The box and whisker plot on Y-axis represents 1st quartile, median and 3rd

quartile. Graph generated in iPathwayGuide (Advaita). **c.** Heatmap visualization for predicted

metformin targets *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4* and *Foxo6* (Log2FC > 0.6; FDR < 0.05) between

untreated *Smn$^{+/-}$;SMN2* healthy mice (left), untreated *Smn$^{-/-}$;SMN2* SMA mice (centre) and

prednisolone-treated *Smn$^{-/-}$;SMN2* SMA mice (right). Colour key represents log2FC for

upregulated (red) and downregulated (blue) genes. Heatmap was generated by Heatmap

v2.1.1+galaxy1.

1320

Figure S3. Smn knockdown in C2C12 myoblasts and myotubes via *Smn* siRNA transfection.

Smn siRNA knockdown (red) was performed for **a.** 48 hours in C2C12 myoblasts and **b.** every 48

hours throughout differentiation in D8 C2C12 myotubes. *Smn* knockdown of mRNA levels was

confirmed by qPCR and compared to non-transfected (black) and scrambled siRNA control (blue)

groups. Data are shown as scatter graph that represent mean ± SEM error bars; n = 4 samples per

group across two independent experiments. One-way ANOVA followed by Tukey's multiple

comparisons test. C2C12 myoblasts F = 22.01; D8 C2C12 myotubes F = 115.4. ****p <0.0001.

1328

Figure S4. Both physiological and supraphysiological metformin concentrations affect the

expression of specific predicted target genes in C2C12 myoblasts and myotubes.

a. C2C12 myoblasts and **b.** D8 C2C12 myotubes were treated with a range of physiological (30

µM (red) and 60 µM (green)) and supraphysiological (1 mM (brown) and 2 mM (blue)) metformin

concentrations for 24 hours against a PBS vehicle control (black) to evaluate the mRNA expression

via qPCR of predicted target genes *Prkag3*, *Foxo1*, *Foxo3*, *Foxo4* and *Foxo6*. Data are shown as

bar charts with scatter graph that represent mean ± SEM error bars; n = 4 samples per group across

two independent experiments. Two-way ANOVA followed by uncorrected Fisher's least

significant difference (LSD). C2C12 myoblasts F = 7.822; D8 C2C12 myotubes F = 12.17. *p

<0.05, **p <0.01, ***p <0.001, ****p <0.0001.

1339

Figure S5. 0.9% saline vehicle treatment had no effect on survival or phenotype in *Smn$^{2B/-}$*

SMA and Smn$^{2B/+}$ healthy mice.

All treated animals received a daily dose of 0.9% physiological saline vehicle by gavage starting

at P5**. a.** Survival curves of untreated (n = 13, black) and 0.9% saline vehicle-treated (n = 14,

green) *Smn$^{2B/-}$* SMA mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test,

ns = not significant, *p* = 0.0653. **b.** Daily weights of untreated (n = 13, black) and 0.9% saline

vehicle-treated (n = 14, green) *Smn$^{2B/-}$* SMA mice. Data represented as mean ± SEM error bars;

Two-way ANOVA followed by a Sidak's multiple comparison test, F = 454.6, df = 501 **c.** Daily

righting reflex test for motor function activity up to a 30 second maximum time point in untreated

(n = 13, black) and 0.9% saline vehicle-treated (n = 14, green) *Smn$^{2B/-}$* SMA mice. Data are shown

as bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, *p* = 0.6626**. d.**

Survival curves of untreated (n = 16, black) and 0.9% saline vehicle-treated (n = 22, green) *Smn$^{2B/+}$*

healthy mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test, ns = not

significant, *p* > 0.9999. **e.** Daily weights of untreated (n = 16, black) and 0.9% saline vehicle-

treated (n = 22, green) $Smn^{2B/+}$ healthy mice. Data represented as mean ± SEM error bars; Two-

way ANOVA followed by a Sidak's multiple comparison test, F =328, df = 757. **f.** Daily righting

reflex test for motor function activity up to a 30 second maximum time point in untreated (n = 16,

black) and 0.9% saline vehicle-treated (n = 22, green) $Smn^{2B/+}$ healthy mice. Data are shown as

bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, $p = 0.9555$.

1359

Figure S6. 200 and 400 mg/kg/day metformin had no negative effect on survival or phenotype

in healthy $Smn^{2B/+}$ mice.

All treated animals received a daily dose of metformin (either 200 or 400 mg/kg/day, diluted in

0.9% saline) by gavage starting at P5. **a.** Survival curves of untreated (n = 16, black) and 200

mg/kg/day metformin-treated (n = 20, red) $Smn^{2B/+}$ healthy mice. Kaplan-Meier survival curve

shown with log rank (Mantel-Cox) test, ns = not significant, $p > 0.9999$. **b.** Daily weights of

untreated (n = 16, black) and 200 mg/kg/day metformin-treated (n = 20, red) $Smn^{2B/+}$ healthy mice.

Data represented as mean ± SEM error bars; Two-way ANOVA followed by a Sidak's multiple

comparison test, F = 549.5, df = 680. **c.** Daily righting reflex test for motor function activity up to

a 30 second maximum time point in untreated (n = 16, black) and 200 mg/kg/day metformin-

treated (n = 20, red) $Smn^{2B/+}$ healthy mice. Data are shown as bar chart with mean ± SEM error

bars; unpaired T-test, ns = not significant, $p = 0.9183$. **d.** Survival curves of untreated (n = 16,

black) and 400 mg/kg/day metformin-treated (n = 15, blue) $Smn^{2B/+}$ healthy mice. Kaplan-Meier

survival curve shown with log rank (Mantel-Cox) test, ns = not significant, $p > 0.9999$. **e.** Daily

weights of untreated (n = 16, black) and 400 mg/kg/day metformin-treated (n = 15, blue) $Smn^{2B/+}$

healthy mice. Data represented as mean ± SEM error bars; Two-way ANOVA followed by a

Sidak's multiple comparison test, F = 261.9, df = 435. **f.** Daily righting reflex test for motor

function activity up to a 30 second maximum time point in untreated (n = 16, black) and 400

mg/kg/day metformin-treated (n = 15, blue) $Smn^{2B/+}$ healthy mice. Data are shown as bar chart

with mean ± SEM error bars; unpaired T-test, ns = not significant, $p = 0.9966$.

1380

Figure S7. Oxandrolone is predicted to emulate the target patterns of prednisolone in the

skeletal muscle (Triceps) of $Smn^{-/-};SMN2$ SMA mice.

a. Predicted model based on upstream regulator patterns predicted by iPathwayGuide in

prednisolone vs untreated $Smn^{-/-};SMN2$ SMA skeletal muscle. *Ar* upregulates downstream targets

Igfbp5 and *MyoG*, whilst it downregulates *Ddit4*. *Igfbp5* upregulates *Dok5* and *MyoG* upregulates

Akap6. Upregulated genes shaded in red, downregulated genes shaded in blue. Downregulation of

Ddit4 based on previous published literature. **b.** Differential gene expression pattern by logFC (Y-

axis) of predicted oxandrolone targets *Ddit4*, *Igfbp5*, *Ar*, *MyoG*, *Akap6* and *Dok5* based on

transcriptomic data from prednisolone vs untreated $Smn^{-/-};SMN2$ SMA skeletal muscle.

Upregulated genes above X-axis are highlighted in red and downregulated genes below X-axis are

highlighted in blue. The box and whisker plot on Y-axis represents 1st quartile, median and 3rd

quartile. Graph generated in iPathwayGuide (Advaita). **c.** Heatmap visualization for predicted

oxandrolone targets *Ar*, *MyoG*, *Igfbp5*, *Dok5*, *Akap6* and *Ddit4* (log2FC > 0.6; FDR < 0.05)

between untreated $Smn^{+/-};SMN2$ healthy mice (left), untreated $Smn^{-/-};SMN2$ SMA mice (centre)

and prednisolone-treated $Smn^{-/-};SMN2$ SMA mice (right). Colour key represents log2FC for

upregulated (red) and downregulated (blue) genes. Heatmap was generated by Heatmap

v2.1.1+galaxy1.

1398

Figure S8. Low 1 μM oxandrolone treatment is non-toxic and does not impact proliferation in C2C12 myoblasts and myotubes.

a. C2C12 myoblasts and **b.** D5 C2C12 myotubes were treated with 1 μM oxandrolone for 24 (purple) and 72 hours (orange) and compared to an absolute ethanol vehicle (24 hours red, 72 hours blue) in addition to untreated (black) and 1% Triton-X max lactate dehydrogenase (LDH) control (green). LDH levels in cell culture supernatant were measured by proportional fluorescence absorption readings (nm). C2C12 myoblasts were treated with 1 μM oxandrolone for **c.** 24 (orange) and **d.** 72 hours (purple) against an absolute ethanol vehicle (24 hours red, 72 hours blue) in addition to blank media (brown), blank cells (cyan), and untreated cells (black) controls. Absorption readings (nm) were measured from anti-BrDU antibody immunostained samples. Data are shown as bar charts that represent mean ± SEM error bars; n = 6 samples per group across one independent experiments; one-way ANOVA followed by Dunnett's multiple comparisons test, **a.** F = 44.25, **b.** F = 3.092, **c.** F = 67.51, **d.** F = 64.71, $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$, ns = not significant.

Figure S9. D5 stage in C2C12 myotubes for oxandrolone to elicit an *Ar* gene response.

a. C2C12 myoblasts were treated with a range of oxandrolone concentrations of 1 (blue), 10 (red) and 100 μM (green) for 24 hours and compared to an absolute ethanol (vehicle, black) to evaluate the mRNA expression via qPCR of predicted target genes *Ar*, *Dok5*, *Igfbp5*, *Akap6*, *MyoG* and *Ddit4*. Data are shown as a bar chart and scatter graph that represent mean ± SEM error bars; n = 4 samples per group across two independent experiments. Two-way ANOVA followed by uncorrected Fisher's least significant difference (LSD), F = 1.693. $*p <0.05$, $**p <0.01$, $***p <0.001$, $****p <0.0001$. **b.** D8 C2C12 myotubes were treated with a range of oxandrolone

concentrations of 1 (blue), 10 (red) and 100 μM (green) for 24 hours and compared to an absolute

ethanol (vehicle, black) to evaluate the mRNA via qPCR of predicted target gene *Ar* and

downstream targets *Igfbp5* and *MyoG*. Data are shown as bar chart and scatter graph that represent

mean ± SEM error bars; n = 4 samples per group across two independent experiments. Two-way

ANOVA followed by uncorrected Fisher's least significant difference (LSD), F = 0.8708. **c.** D3

and **d.** D5 C2C12 myotubes were treated with 1 μM oxandrolone for 24 hours (blue) against an

absolute ethanol (vehicle, black) to evaluate the mRNA levels via qPCR of *Ar*. Data are shown as

scatter graphs that represent mean ± SEM error bars; n = 4 samples per group across one

independent experiment; unpaired T-test D3 C2C12 myotubes $p = 0.0203$, D5 C2C12 myotubes p

$= 0.0480$, *p <0.05. **e.** D5 C2C12 myotubes were treated with 1 μM oxandrolone (blue) for 24

hours against an absolute ethanol (vehicle, black) to evaluate the mRNA levels via qPCR of target

genes *Dok5*, *Igfbp5*, *Akap6*, *MyoG* and *Ddit4*. Data are shown as bar charts with scatter graphs that

represent mean ± SEM error bars; n = 4 samples per group across two independent experiments.

Two-way ANOVA followed by uncorrected Fisher's least significant difference (LSD), F =

0.7013.

1437

1438

Figure S10. 0.5% carboxymethyl cellulose vehicle treatment had no effect on survival or

phenotype in *Smn²ᴮ/⁻* SMA and *Smn²ᴮ/⁺* healthy mice.

All treated animals received a daily dose of 0.5% carboxymethyl cellulose (CMC) vehicle by

gavage starting at P8**. a.** Survival curves of untreated (n = 15, black) and 0.5% CMC vehicle-

treated (n = 13, violet) *Smn²ᴮ/⁻* SMA mice. Kaplan-Meier survival curve shown with log rank

(Mantel-Cox) test, ns = not significant, $p = 0.4222$. **b.** Daily weights of untreated (n = 15, black)

and 0.5% CMC vehicle-treated (n = 13, violet) $Smn^{2B-/-}$ SMA mice. Data represented as mean ±

SEM error bars; Two-way ANOVA followed by a Sidak's multiple comparison test, F = 335 df =

511 **c.** Daily righting reflex test for motor function activity up to a 30 second maximum time point

in untreated (n = 15, black) and 0.5% CMC vehicle-treated (n = 13, violet) $Smn^{2B/-}$ SMA mice.

Data are shown as bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, p

= 0.5602**. d.** Survival curves of untreated (n = 21, black) and 0.5% CMC vehicle-treated (n = 16,

violet) $Smn^{2B/+}$ healthy mice. Kaplan-Meier survival curve shown with log rank (Mantel-Cox) test,

ns = not significant, p > 0.9999. **e.** Daily weights of untreated (n = 21, black) and 0.5% CMC

vehicle-treated (n = 16, violet) $Smn^{2B/+}$ healthy mice. Data represented as mean ± SEM error bars;

Two-way ANOVA followed by a Sidak's multiple comparison test, F = 377.3, df = 724. **f.** Daily

righting reflex test for motor function activity up to a 30 second maximum time point untreated (n

= 21, black) and 0.5% CMC vehicle-treated (n = 16, violet) $Smn^{2B/+}$ healthy mice. Data are shown

as bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, p = 0.9638**.**

1458

1459

1460

Figure S11. 4 mg/kg/day oxandrolone decreased bodyweight in healthy $Smn^{2B/+}$ mice.

All treated animals received a daily dose of oxandrolone (4 mg/kg/day, suspended in 0.5% CMC)

by gavage starting at P8. **a.** Survival curves of untreated (n = 21, black) and 4 mg/kg/day

oxandrolone-treated (n = 10, orange) $Smn^{2B/+}$ healthy mice. Kaplan-Meier survival curve shown

with log rank (Mantel-Cox) test, ns = not significant, p > 0.9999. **b.** Daily weights of untreated (n

= 21, black) and 4 mg/kg/day oxandrolone-treated (n = 10, orange) $Smn^{2B/+}$ healthy mice. Data

represented as mean ± SEM error bars; Two-way ANOVA followed by a Sidak's multiple

comparison test, F = 395.5, df = 598, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. **c.** Daily righting reflex test for motor function activity up to a 30 second maximum time point in untreated (n = 21, black) and 4 mg/kg/day oxandrolone-treated (n = 10, orange) $Smn^{2B/+}$ healthy mice. Data are shown as bar chart with mean ± SEM error bars; unpaired T-test, ns = not significant, $p = 0.7865$.

1491 **SUPPLEMENTARY TABLES**

1492 **Table S1. RNA sequencing sample groups.**

1493

1494 **Table S2. Murine primers for qPCR.**

1495

1496 **Table S3. Significant differentially expressed genes (Log2FC > 0.6; FDR < 0.05) in skeletal**

1497 **muscle (triceps) between P7 untreated *Smn[-/-];SMN2* SMA and *Smn[+/-];SMN2* healthy mice.**

1498

1499 **Table S4. Significant differentially expressed genes (Log2FC > 0.6; FDR < 0.05) in skeletal**

1500 **muscle (triceps) between P7 prednisolone-treated vs untreated *Smn[-/-];SMN2* SMA mice.**

1501

1502 **Table S5. Raw Counts for significant differentially expressed genes in skeletal muscle**

1503 **(triceps) between P7 prednisolone-treated vs untreated *Smn[-/-];SMN2* SMA and untreated**

1504 ***Smn[+/-];SMN2* healthy mice.**

1505

1506 **Table S6. Complete list of significant Gene Ontology (GO) Term (Biological Processes) for**

1507 **P7 prednisolone-treated vs untreated *Smn[-/-];SMN2* SMA mice.**

1508

1509 **Table S7. Complete list of significant Gene Ontology (GO) Term (Molecular Functions) for**

1510 **P7 prednisolone-treated vs untreated *Smn[-/-];SMN2* SMA mice.**

1511

1512 **Table S8. Complete list of significant Gene Ontology (GO) Term (Cell Components) for P7**

1513 **prednisolone-treated vs untreated *Smn[-/-];SMN2* SMA mice.**

1514

1515   **Table S9. List of drugs identified in iPathwayGuide from KEGGS drug database that target**

1516   **significantly impacted prednisolone pathways.**

1517

1518   **Table S10. List of predicted upstream regulators from P7 prednisolone-treated vs untreated**

1519   *Smn$^{-/-}$;SMN2* **SMA mice used in DGIdb v.3. gene-drug target search.**

1520

1521   **Table S11. List of predicted agonist drugs from DGIdb v.3. for targeting upregulated**

1522   **upstream regulators.**

1523

1524   **Table S12. List of predicted antagonist drugs from DGIdb v.3. for targeting downregulated**

1525   **upstream regulators.**

1526

1527

1528

1529

Taiwanese *Smn⁻/⁻;SMN2*   Intermediate *Smn²ᴮ/⁻*

Taiwanese *Smn⁻/⁻;SMN2*

Intermediate *Smn²ᴮ/⁻*

Metformin

Oxandrolone

a.

Pharyngeal Pump

d.

Pharyngeal Pump

b.

Reversals

e.

Reversals

c.

Paralysis Time

f.

Paralysis Time

Table 1. KEGG pathways targeted in the skeletal muscle of symptomatic prednisolone-treated *Smn^{-/-};SMN2* SMA mice compared with untreated *Smn^{-/-};SMN2* SMA mice.

KEGG ID	Pathway name	#Genes (DE/All)	*p* value
04066	HIF-1 signaling pathway	30/91	0.001
05214	Glioma	25/71	0.002
04213	Longevity regulating pathway - multiple species	24/57	0.003
04068	FoxO signaling pathway	41/116	0.003
04713	Circadian entrainment	23/73	0.004
04211	Longevity regulating pathway	30/83	0.004
04115	p53 signaling pathway	25/67	0.006
00040	Pentose and glucuronate interconversions*	8/15	0.007
05010	Alzheimer disease	19/158	0.008
04152	AMPK signaling pathway	37/110	0.008
04080	Neuroactive ligand-receptor interaction	23/118	0.011
05418	Fluid shear stress and atherosclerosis	38/125	0.013
05223	Non-small cell lung cancer	22/64	0.014
05034	Alcoholism	25/104	0.014
04744	Phototransduction	4/9	0.015
05215	Prostate cancer*	28/88	0.016
04137	Mitophagy - animal	22/61	0.018
04659	Th17 cell differentiation	18/72	0.024
03030	DNA replication*	13/35	0.026
05031	Amphetamine addiction	13/49	0.028
05200	Pathways in cancer*	110/434	0.031
04710	Circadian rhythm	12/29	0.033
01210	2-Oxocarboxylic acid metabolism*	7/16	0.039
04140	Autophagy - animal	36/123	0.041
04372	Apelin signaling pathway	34/121	0.045
05226	Gastric cancer	36/123	0.048
00515	Mannose type O-glycan biosynthesis*	8/20	0.048
03320	PPAR signaling pathway	20/56	0.048

*Over-representation only

Table 2. Top Gene Ontology Biological Process pathways targeted in the skeletal muscle of symptomatic prednisolone-treated *Smn*[-/-]*;SMN2* SMA mice compared with untreated *Smn*[-/-]*;SMN2* SMA mice.

GO ID	GO Name	#Genes (DE/All)	*p value* (Weight)
GO:0000055	ribosomal large subunit export from nucleus	8/8	0.0000054
GO:0010830	regulation of myotube differentiation	25/53	0.000088
GO:0034504	protein localization to nucleus	81/247	0.00011
GO:0046320	regulation of fatty acid oxidation	17/32	0.00028
GO:0048662	negative regulation of smooth muscle cell proliferation	20/44	0.00045
GO:0031062	positive regulation of histone methylation	17/35	0.00046
GO:0046854	phosphatidylinositol phosphorylation	17/36	0.0007
GO:0016239	positive regulation of macroautophagy	26/64	0.00113
GO:0000083	regulation of transcription involved in G1/S transition of mitotic cell cycle	8/12	0.00113
GO:0042594	response to starvation	49/148	0.00114
GO:0031398	positive regulation of protein ubiquitination	36/101	0.00115
GO:0090073	positive regulation of protein homodimerization activity	7/10	0.00157
GO:0048671	negative regulation of collateral sprouting	6/8	0.00208
GO:0007050	cell cycle arrest	40/121	0.00319
GO:1990830	cellular response to leukaemia inhibitory factor	37/110	0.00321
GO:1901215	negative regulation of neuron death	60/198	0.00378
GO:0007623	circadian rhythm	50/160	0.00378
GO:0061635	regulation of protein complex stability	6/9	0.00508
GO:0033137	negative regulation of peptidyl-serine phosphorylation	12/26	0.00533
GO:0032088	negative regulation of NF-kappaB transcription factor activity	22/59	0.00537
GO:0006094	gluconeogenesis	23/63	0.00604
GO:0001937	negative regulation of endothelial cell proliferation	15/36	0.00629
GO:0010715	regulation of extracellular matrix disassembly	7/12	0.00681
GO:0030240	skeletal muscle thin filament assembly	7/12	0.00681
GO:0035358	regulation of peroxisome proliferator activated receptor signalling pathway	7/12	0.00681

Table 3. Top 10 clinically approved drugs identified by KEGG database based on prednisolone-targeted KEGG pathways in symptomatic prednisolone-treated *Smn-/-;SMN2* SMA mice.

Pathways Targeted	KEGG ID	Compound	Bioavailability	Gene Targets
10	D03297	Mecasermin (genetical recombination) (JAN)	Subcutaneous	*Igf1r*
10	D04870	Mecasermin rinfabate (USAN/INN)	Intravenous	*Igf1r*
9	D09680	Teprotumumab (USAN/INN)	Intravenous	*Igf1r*
8	D04966	Metformin (USAN/INN)	Oral	*Prkag3*
8	D00944	Metformin hydrochloride (JP16/USP)	Oral	*Prkag3*
7	D01697	Colforsin daropate hydrochloride (JAN)	Oral	*Adcy1, Adcy6 Adcy7*
6	D01146	Iguratimod (JAN/INN)	Oral	*Nfkb1*
5	D07058	Acamprosate (INN)	Oral	*Grin1, Grin2a, Grin2b, Grin2c, Grin2d*
5	D02754	Acitretin (USP/INN); Soriatane (TN)	Oral	*Rarb, Rxrb, Rxrg*
5	D00085	Insulin (JAN/USP)	Intravenous Subcutaneous	*Insr*

Table 4. Top 10 clinically approved drugs identified by DGIdb database based on prednisolone-targeted KEGG pathways in symptomatic prednisolone-treated *Smn⁻/⁻;SMN2* SMA mice.

Compound	Bioavailability	Gene Target
Testosterone	Busal Nasal Oral Topical Transdermal Subcutaneous	*Ar*
Oxandrolone	Oral	*Ar*
Nandrolone phenpropionate	Oral	*Ar*
Progesterone	Oral Vaginal Intramuscular	*Ers1*
Tibolone	Oral	*Ers1*
Cannabidiol	Oral Inhale	*Cnr1*
Insulin, neutral	Intravenous Intramuscular Subcutaneous	*Insr*
Celecoxib	Oral	*Pdpk1*
Tocilizumab	Intravenous Subcutaneous	*Il-6ra*
Sarilumab	Subcutaneous	*Il-6ra*