Pulsed ultrasound promotes secretion of anti-inflammatory extracellular vesicles from skeletal myotubes via elevation of intracellular calcium level

Atomu Yamaguchi[1], Noriaki Maeshige[1], Hikari Noguchi[1], Jiawei Yan[2], Xiaoqi Ma[1], Mikiko Uemura[1], Hiroyo Kondo[3], Kristopher Sarosiek[4], Hidemi Fujino[1]

[1]Department of Rehabilitation Science, Kobe University Graduate School of Health Sciences, Kobe, Japan

[2]School of Life Sciences and Technology, ShanghaiTech University, Shanghai, China.

[3]Department of Food Science and Nutrition, Nagoya Women's University, Nagoya, Japan

[4]John B. Little Center for Radiation Sciences, Harvard University T.H Chan School of Public Health, Boston, MA, USA

Correspondence:

Noriaki Maeshige

Department of Rehabilitation Science, Kobe University Graduate School of Health Sciences

654-0142, 7-10-2 Tomogaoka, Kobe, Hyogo, Japan

Tel: +81 78 796 4582

E-mail address: nmaeshige@pearl.kobe-u.ac.jp

Abstract

The regulation of inflammatory responses is an important intervention in biological function and macrophages play an essential role during inflammation. Skeletal muscle is the largest organ in the human body and releases various factors which mediate anti-inflammatory/immune modulatory effects. Recently, the roles of extracellular vesicles (EVs) from a large variety of cells are reported. In particular, EVs released from skeletal muscle are attracting attention due to their therapeutic effects on dysfunctional organs and tissues. Also, ultrasound (US) promotes release of EVs from skeletal muscle. In this study, we investigated the output parameters and mechanisms of US-induced EV release enhancement and the potential of US-treated skeletal muscle-derived EVs in the regulation of inflammatory responses in macrophages. High-intensity US (3.0 W/cm^2) irradiation increased EV secretion via elevating intracellular Ca^{2+} level without negative effects. Moreover, US-induced EVs suppressed expression levels of pro-inflammatory factors in macrophages. miRNA sequencing analysis revealed that miR-206-3p and miR-378a-3p were especially abundant in skeletal myotube-derived EVs. In this study we demonstrated that high-intensity US promotes the release of anti-inflammatory EVs from skeletal myotubes and exert anti-inflammatory effects on macrophages.

Keywords: Extracellular vesicles, Skeletal muscles, Myotubes, Macrophages, Ultrasound

Introduction

Inflammation is a crucial response to defend the body from infection. However, excess and prolonged inflammation can also be harmful and needs to be tightly regulated (Funes et al., 2018). Macrophages play a leading role in the innate immune system, causing inflammation during infection (Hirayama et al., 2017). Therefore, regulating the condition of macrophages is a major therapeutic strategy for the control of excess inflammation. Various organs including skeletal muscle are known to release anti-inflammatory factors. Skeletal muscle is the largest organ in the human body accounting for 40% of the body weight (Trovato et al., 2019) and responsible for whole-body metabolism, energy homeostasis, and locomotion. Recently, skeletal muscle is attracting attention as a secretory organ of anti-inflammatory factors (Trovato et al., 2019), and extracellular vesicles (EVs) are responsible for transporting various factors from skeletal muscle to target organs or cells. EVs are nano-sized vesicles secreted by most types of cells and facilitate cell-to-cell communication (Trovato et al., 2019; Raposo et al., 2013; Andaloussi et al., 2013) through transportation of proteins, mRNAs, and miRNAs (Zomer et al., 2010), leading to regulation of immune response (Andaloussi et al., 2013), tissue regeneration (Bittel et al., 2019), and cell proliferation/differentiation (Matsuzaka et al., 2016). EVs from skeletal muscle have also been reported to exert therapeutic effects in various dysfunctional organs (Rome et al., 2019; Bei et al., 2017; Madison et al., 2014) and our previous study revealed that skeletal myotube-derived EVs attenuate inflammatory responses of macrophages (Yamaguchi et al., 2023). EVs from mesenchymal stem cells (MSC) also have been reported to suppress inflammatory responses in lipopolysaccharide (LPS)-induced macrophages (Harrell et al., 2019; Xin et al., 2020; Shao et al., 2020) and Kim et al.

reported the anti-inflammatory action of MSC EVs depended on the concentration of

EVs (Kim et al., 2019). Therefore, an enhancement of EV release may enhance the

anti-inflammatory effects of muscle EVs.

Physical exercise is one of the main means to promote EV release (Bei et al., 2017;

Whitham et al., 2018). Fruhbeis et al. reported that total amount of EVs increased by 2.7

times after high-intensity exercise (Fruhbeis et al., 2015). However, high-intensity

exercise can be difficult and not feasible for people with underlying diseases. Therefore,

development of means to promote EV secretion from skeletal muscle without physical

burden is expected. Ultrasound (US) irradiation is used as a non-invasive therapy and

has physiological effects including cell proliferation, suppression of inflammatory

signaling (Ueno et al., 2021), and increase cell membrane permeability (Ma et al., 2022).

Our previous study revealed that high-intensity US can promote EV release (Maeshige

et al., 2021). An increase in intracellular Ca^{2+} concentration is one of the effects of US

irradiation, and elevation of intracellular Ca^{2+} is a key factor for EV secretion (Savina et

al., 2003). However, the role of Ca^{2+} in US-induced EV release and its

intensity-dependency are still unclear and anti-inflammatory effects of US-induced EVs

have not been elucidated. This study aimed to clarify the mechanism and

intensity-dependency of EV release enhancement by US and anti-inflammatory effects

of US-induced skeletal muscle-derived EVs on macrophages.

Results

US irradiation has no negative effect on myotube viability, protein content, and

energy metabolism

To measure the viability of US-irradiated myotubes, MTT assay and Zombie Red[TM]

immunofluorescence staining were performed. Our results showed no significant decrease in all US groups (Figure 1A, B). To assess the potential effect of US irradiation on cultured myotubes, the total protein content in myotubes was measured and no significant difference was observed among all groups (Figure 1C). Furthermore, CS activity was measured to determine the effect of US on energy metabolism. CS activity in each group did not decrease in the US-treated myotubes (Figure 1D).

US irradiation enhances release of EVs from myotubes

The concentration of EVs in the 3W group was 2 times higher than that in the control group. The 1W and 2W groups also showed an increase in EVs concentration, which was 1.64 and 1.68 times higher than that in the Control group (Figure 2A), but no significant difference. Regarding size distribution, the majority of released EVs were in the range of 50-150 nm and the size of EVs did not change in the US groups (Figure 2B).

Intracellular Ca^{2+} upregulation mediates US-induced enhancement of EV release from myotubes

To investigate the mechanism of the effect of US on EV release enhancement, we measured intracellular Ca^{2+} levels in myotubes after US irradiation. Compared to the control group, Ca^{2+} level in the 3W group was significantly increased immediately after US irradiation. The 2W group only showed a tendency of increased intracellular Ca^{2+} level, but no significant difference (Figure 3A).

Next, we investigated the calcium dependency of the promotive effect of US on EV release using Ca^{2+}-free medium. Here we used the US intensity of 3.0 W/cm^2, which

showed the greatest EV release enhancement effect. The groups with Ca^{2+}-free medium

showed significantly lower levels of Ca^{2+} compared with the control and US groups

(Figure 3B). The concentration of EVs in US group was significantly increased

compared to the control group while this increase was not observed in absence of Ca^{2+}

(Figure 3C). Most of the extracted EVs were 50-150 nm in diameter (Figure 3D).

Myotube viability did not decrease by Ca^{2+}-free culture (Figure 3E).

US-induced EVs exert anti-inflammatory effects on macrophages

To investigate the anti-inflammatory effect of myotube EVs on macrophages, qPCR

analysis was performed. While C2C12 myotube conditioned medium (CM) only

showed a tendency to suppress the expression of pro-inflammatory factors, US-treated

C2C12 myotube conditioned medium (US-CM) significantly suppressed the expression

of pro-inflammatory *Il-1b* and *Il-6*, compared to LPS-stimulation alone (LPS), and

expression of *Il-1b* in US-CM significantly decreased compared to non-treated C2C12

myotube conditioned medium. Furthermore, the expression of *Il-1b* and *Il-6* were not

suppressed when the EVs were eliminated from the conditioned media (Figure 4A).

When EV concentrations were equated, the upregulation of the anti-inflammatory effect

of myotube EVs by US was not observed (Figure 4B). Myotube EVs did not decrease

macrophage viability (Figure 4C). EV concentration in each group is shown in Figure

4D.

miRNA profile change in myotube EVs by US

To investigate the effect of US on miRNA profile in myotube-derived EVs, miRNA

sequencing analysis on EVs was performed. A total of 524 miRNAs were identified by

proteomic quantitative analysis. Twenty-nine miRNAs were expressed specifically in the control group, and 81 miRNAs were expressed specifically in the US group (Figure 5A). Lists of miRNAs specific to each group are shown in Supplemental material 1. Although miRNAs specific to each group were identified, they accounted for only 0.014% of the total miRNAs, and 99.99% of total miRNAs were common to the control and US groups (Figure 5B). According to the standard of a fold change of fold change $\geq$ 2 or $\leq$ 0.5 as well as an FDR < 0.05, we screened 13 up-regulated miRNAs (Table 1) and 14 down-regulated miRNAs (Table 2) in the EV group versus the control group. Differentially expressed miRNAs are displayed as a volcano plot (Figure 5C). The top 10 abundant miRNAs and their proportion to the total miRNAs are shown (Figure 5D). In both the control and US groups, the two most abundant miRNAs, miR-206-3p and miR-378a-3p, accounted over 60% of the total.

Discussion

This study demonstrated the facilitatory effect of US irradiation on EV release from cultured skeletal myotubes and the enhanced anti-inflammatory effect of EVs from US-irradiated myotubes. These findings contribute to the development of non-invasive complementary therapy using skeletal muscle-derived EVs for various inflammatory diseases.

Skeletal muscle-derived EVs have been reported to exert therapeutic effects in various dysfunctional organs (Rome et al., 2019), including prevention of cardiac damage by ischemia-reperfusion injury (Bei et al., 2017), enhancement of neurite outgrowth in motor neurons (Madison et al., 2014), and enhancement of angiogenesis (Ma et al., 2018). Furthermore, since skeletal muscle EVs mediate cross-talk between

skeletal muscle and other tissues (Whitham et al., 2018), methods to enhance the release

of skeletal muscle-derived EVs are potentially beneficial in the treatment of various

diseases and health management.

Our results showed that the intracellular Ca^{2+} level significantly increased in the

US-irradiated group, indicating that US irradiation promoted the uptake of Ca^{2+} to

myotubes. This result is supported by previous reports investigating Ca^{2+} uptake by US

in various types of cells (Ambattu et al., 2020; Scheffer et al., 2010; Fan et al., 2010;

Lentacker et al., 2014; Honda et al., 2004). In the present study, the effect of US on Ca^{2+}

uptake was observed in an intensity-dependent manner. US irradiation has been reported

to exert a physiological action to promote transient membrane permeabilization, which

can promote the Ca^{2+} influx pathways (Fan et al., 2010; Cao et al., 2008; Zhou et al.,

2008), and the efficiency of permeabilization depends on the intensity of irradiation

(Kumon et al., 2009; Zeghimi et al., 2015). Thus, the change of cell membrane structure

is supposed to be the mechanism of the enhancement of Ca^{2+} influx. In this study, EV

release was promoted by US of 3.0 W/cm^2, and this effect was cancelled by inhibition

of Ca^{2+}. Since intracellular Ca^{2+} level is reported to be a factor that promotes EVs

release (Savina et al., 2003; Taylor et al., 2020), US irradiation was shown to enhance

EV release from myotubes via promotion of influx of Ca^{2+}.

We found that C2C12 myotube conditioned medium was capable of suppressing

LPS- induced inflammatory responses in macrophages, and this effect was enhanced by

US irradiation to myotubes. In addition, the anti-inflammatory effects were cancelled by

eliminating EVs from the culture medium. This indicates that EVs in the culture

medium were responsible for this effect, which is consistent with our previous study

(Yamaguchi et al., 2023).

IL-1β is essential for an adequate acute inflammatory response to a variety of

pathogens and injuries, but its excessive overexpression leads to sepsis, septic shock, or

chronic inflammation (Dinarello, 2009; Piccioli et al., 2013). IL-6 also contributes to

host defense, however, dysregulated continual synthesis of it leads to prolonged chronic

inflammation (Tanaka et al., 2014) and can drive tumorigenesis (Chang et al., 2014).

Since blockade of these factors is reported to alleviate several inflammatory diseases

(Chevalier et al, 2005; Emsley et al., 2005; Choy et al., 2002; Nishimoto et al., 2004; Ito

et al., 2004; Yokota et al., 2005), eliciting the anti-inflammatory effects of skeletal

muscle-derived EVs by US and suppressing the production of IL-1β and IL-6 suggests a

new therapeutic strategy against inflammatory diseases utilizing US.

Meanwhile, it has been reported that the contents of EVs released from cells vary

depending on the cellular microenvironment and that certain factors are selectively

internalized in response to specific stimuli (van Niel et al., 2018). The results of our

analysis on miRNAs in EVs revealed that the EVs from US-exposed myotubes contain

several unique miRNAs. However, only about 0.01% of the total miRNAs were altered

by US irradiation, suggesting that the change was very small. Furthermore, no

upregulation of the anti-inflammatory effect of myotube EVs by US was observed when

the EV concentrations were equalized. Thus, the enhancement of the anti-inflammatory

effect of myotube conditioned medium by US irradiation is assumed to be due to

changes in the amount of EVs, not to changes in the content of EVs. Consistent with our

results, a previous study reported that MSC-EVs regulated inflammatory responses in

macrophages concentration-dependently (Kim et al., 2019).

Analysis on miRNA profiles in the EVs revealed that the two most abundant

miRNAs, miR-206-3p and miR-378a-3p, were common to both control and US EVs,

and these two types accounted for more than 60% of the total miRNA contents in the both conditions. The most abundant miR-206 is a member of the skeletal muscle-specific myo-miR family of miRNAs (Carpi et al., 2020). It is reported that miR-206 targets IL-17A and REG3A and suppresses macrophage inflammation (Huang et al., 2020). Furthermore, Lin et al. reported that transfection of miR-mimic-206-3p into macrophages suppressed macrophage inflammation by targeting PPP3CA and transfection of miR-inhibitor-206-3p increased the level of inflammatory factors in macrophages (Lin et al., 2020). miR-378a is highly expressed in skeletal muscle and is involved in metabolism and mitochondrial energy homeostasis (Krist et al., 2015). Rückerl et al. identified miR-378a-3p as a factor contributing to the induction of anti-inflammatory macrophage reprogramming through IL-4-induced gene transcription by targeting Akt (Rückerl et al., 2012). In addition, Kris et al. reported that miR-378a has anti-inflammatory effects on macrophages and its deficiency enhances severity of inflammation (Kris et al., 2020). Based on these previous studies, it is assumed that myotube-derived EVs elicited the anti-inflammatory effects in macrophages by delivering these miRNAs. Additionally, miR-206 has been reported to modulate fat metabolism in diabetes (Wu et al., 2017) and control the tumorigenesis process of cancer (Ding et al., 2017), and miR-378a has been shown to inhibit cardiac hypertrophy (Ganesan, 2013) and cancer growth (Zeng et al., 2017) and alleviate spinal cord injury (Zhang et al., 2021), indicating that these miRNAs are expected to have therapeutic effects against various diseases. Therefore, promoting the release of EVs containing these factors by US stimulation to skeletal muscle is potentially effective in the prevention and treatment of various diseases, and further investigations of its effects on other pathological conditions are expected.

In summary, this study showed that US irradiation promoted the secretion of myotube-derived EVs which have anti-inflammatory effects and suggested that US irradiation to skeletal muscles is a potent candidate as a novel treatment for various inflammatory disorders.

Materials and methods

Cell culture

C2C12 myoblasts were purchased from the American Type Culture Collection (ATCC, USA). Myoblasts were cultured on 35 mm dishes (Iwaki) in Dulbecco's modified Eagle medium (DMEM, Wako Junyaku Co., Ltd., Japan) supplemented with 10% fetal bovine serum (FBS) at 37 °C under 5% CO_2. At 90% confluence, differentiation into myotubes was initiated by changing the growth medium to differentiation medium (DMEM supplemented with 2% horse serum). Differentiation was continued for 6–7 days, refreshing the medium every 2 days. Efficiency of differentiation was confirmed by observing the contractile ability using electrical stimulation (Supplemental material 2). US was applied to the myotubes for 5 min after changing the differentiation medium to serum-free medium. After incubation for 12 h, the culture medium was collected to extract EVs.

To generate bone marrow-derived macrophages (BMDMs), bone marrow cells from femur and tibias of 7-week-old male C57BL/6 mice were harvested and cultured in Petri dishes with RPMI 1640 (DMEM, Wako Junyaku Co., Ltd., Japan) supplemented with 10% FBS, 1% Penicillin/Streptomycin, 1% L-Glutamine and 25% L929 cell supernatant for 8 days. Differentiated macrophages were harvested and plated in 12-well tissue culture plates with macrophage culture media (RPMI 1640 supplemented with 10%

FBS, 1% Penicillin/Streptomycin, 1% L-Glutamine and 10% L929 cell supernatant) for

subsequent experiments.

US irradiation

To expose the myotubes to US, the culture dish was placed on the probe of an US

transducer (SONICTIZER SZ-100, Minato Medical Science Co., Ltd., Japan). Coupling

gel was applied to fill the space between the probe surface and the bottom of the dish

and sterilized silicone was inserted into the culture medium and fixed to the dish with a

distance of 2 mm from the bottom of the dish (Figure 6). The US parameters used were

as follows: duty cycle of 20%; acoustic frequency of 1 MHz; duration of 5 min;

repetition frequency of 100 Hz; beam nonuniformity ratio of 2.0; and effective radiation

area of 8.0 cm^2 (Maeshige et al., 2021). In this study, three exposure intensities of 1.0

W/cm^2, 2.0 W/cm^2, and 3.0 W/cm^2 were investigated. After US irradiation, the

myotubes were incubated at 37 °C for 12 h. We adopted the intensities of 1.0-3.0 W/cm^2

because output intensities of 0.1-2.5 W/cm^2 are typically applied for therapeutic

purposes in clinical practice (Draper, 2014). After US irradiation, we monitored the

temperature of the culture medium with a thermometer (TM-947SDJ, SATO-SHOJI,

Japan) and confirmed it below 37 °C to distinguish the advantage of US therapy from

its thermal effect.

To investigate the involvement of intracellular Ca^{2+} in the mechanism of EV release

from myotubes, Ca^{2+}-free medium (Nacalai Tesque Inc., Japan) was used. The culture

medium for Ca^{2+}-free groups were changed from normal medium (serum free DMEM

with calcium) to Ca^{2+}-free medium (serum free DMEM without calcium) 1 h before US

irradiation.

EV isolation

EVs were isolated from the conditioned medium of cultured myotubes with polymer precipitation as previously reported (Ter-Ovanesyan et al., 2021). The conditioned medium was collected and centrifuged for 15 min at 3,000 × g to remove cell debris. The supernatant was transferred to a sterile vessel and added Exo Quick-TC (System Biosciences, Palo Alto, CA) (supernatant: Exo Quick reagent = 5 : 1). The tubes were stored overnight at 4 °C and centrifuged at 1,500 × g for 30 min at 4°C. All traces of fluid were removed and the pellet was resuspended with 100 μL of PBS.

EV characterization

A tunable resistive pulse sensing technology (qNano, IZON system; Izon Science Ltd., Christchurch, New Zealand) was used to measure the concentration, size distribution, and diameter of extracted EVs. The system was calibrated for voltage, stretch, pressure, and baseline current using standard beads: CPC100 (concentration; 1.0 × 10^{10} beads/mL). An NP100 nanopore (for 50-200 nm size range) was used and data analysis was performed by a qNano IZON software.

Macrophage treatment with conditioned medium of C2C12 myotubes

At 3 h after US irradiation, the conditioned media were collected and centrifuged (1500 × g, 10 min). Obtained supernatant was supplemented with 10% FBS, 1% Penicillin/Streptomycin, 1% L-Glutamine, and 10% L929 cell supernatant and applied to BMDMs. After 1.5 h-treatment by myotube conditioned medium, BMDMs were incubated with macrophage culture media overnight (12 h). Subsequently, BMDMs

were treated with 100 ng/mL LPS for 1.5 h for pro-inflammatory marker quantification.

To clarify the involvement of EVs in the anti-inflammatory effects on macrophages,

EV-depleted culture medium was added to macrophages. EV depletion was performed

using Exo Quick-TC reagent as previously reported (Mathew et al., 2019).

MTT assay

MTT assay was performed to evaluate cell viability 24 h after US irradiation to

myotubes. BMDM viability was also assessed 24 h after EV treatment to examine the

effect of the treatment with C2C12 conditioned medium on macrophages. The cells

were incubated for 3 h at 37 °C with MTT solution (MTT, Wako Junyaku Co., Ltd.,

Japan) dissolved in culture medium at 0.5 mg/mL, then dissolved in DMSO. The

absorbance at 595 nm was measured using a microplate reader.

Zombie Red^{TM} immunostaining

C2C12 myotube viability was assessed at 24 h after US irradiation or treatment with

1% povidone-iodine (positive control) by Zombie Red^{TM} immunostaining. This reagent

is an amine-reactive fluorescent dye that is non-permeant to live cells but permeant to

cells with a compromised plasma membrane. Myotubes were washed twice with PBS,

stained with Zombie Red^{TM} (1 : 1000) for 15 min at 37 °C, and fixed with 4%

paraformaldehyde for 30 min. Nuclei were stained with DAPI for 5 min at 37°C. The

images were captured with a 10 × objective on a BX50 (OLYMPUS, Japan).

Total protein assay

Total protein content in myotubes was measured at 24 h after US irradiation. The

cells were washed with PBS and collected into tubes using a cell scraper. After

centrifugation at 13,000 rpm for 20 sec, the pellet was processed with 100 μL of PRO-

PREP reagent (iNtRON Biotechnology Co., Ltd. Japan), followed by incubation on ice

for 20 min. After centrifugation at 13,000 rpm for 5 min, the protein concentration was

analyzed using the Bradford method.

Citrate synthase activity

Energy metabolism was evaluated 24 h after US irradiation to the myotube by citrate

synthase (CS) assay as previously reported (Srere, 1963). Briefly, the samples of total

protein extraction were mixed with the following solution: distilled water (DW) +

3mMAcetylCoA+1mMDTNB (1mM DTNB + 1 M Tris-HCL). Oxaloacetate was added

to the mixture and the absorbance at 415 nm was measured using a microplate reader

every 2 min for 10 min at 37°C.

Intracellular Ca^{2+} levels

The concentration of intracellular Ca^{2+} in the medium of C2C12 myotubes was

measured using a Metallo assay calcium kit LS (CPZIII) (Metallogenics LTD., Japan).

This method determines intracellular Ca^{2+} levels by observing the coloration in the

visible region caused by the chelate complex formation between Chlrophosphonazo-III

(CPZIII) and Ca^{2+}. Briefly, the cells were collected and centrifuged at 13,000 rpm for 20

sec to remove the supernatant. The pellet was suspended with a mixture of 200 μL of

RIPA buffer (RIPA Lysis Buffer, SCB Ltd., Japan) and 1 μL of hydrochloric acid (2 M).

After incubation for 30 min, the samples were centrifuged at 10,000 rpm for 10 min and

the Ca^{2+} concentration in the supernatant was analyzed. Ca^{2+} was dissociated by low pH

and collected as the supernatant. Finally, the absorbance was converted to concentration

(mM) according to the manufacturer's protocol.

Quantitative real-time PCR

To measure gene expression in BMDMs, mRNA was isolated with TRIzol RNA

Isolation protocol and used to make cDNA using iScript^{TM} cDNA Synthesis Kit

(BioRad). The StepOne^{TM} Real-Time PCR System was used to analyze the samples

under the following conditions: 95°C (3 min), 40 cycles of 95°C (10 sec) and 60°C (30

s). The reaction mixture consisted of 8 µL cDNA, 1.5 µL 10× buffer, 0.3 µL 10 mM

dTNPs, 1.5 µL 5 µM primers for each gene used in the study (F+R), 3.58µL H2O,

0.075µL Go Taq DNA polymerase and 0.045µL 2× SYBR green (Invitrogen). Target

genes were the pro-inflammatory markers IL-1β and IL-6 (Moore et al., 2020). Relative

expression values for target genes were calculated by normalization to the expression of

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH). Data was analyzed using the

ΔΔCT method. Sequences for qPCR primers are shown in Supplemental material 3.

miRNA sequencing

miRNA was extracted from myotube-derived EVs using TRIzol reagent (Takara

Biotechnology, Japan) according to the manufacturer's instructions. Raw miRNA

sequence data were obtained using an Illumina NovaSeq™ 6000 machine. After

acquiring the raw data, the fold change (mean of each miRNA in the US group / mean

of each RNA in the control group) and P-values were calculated for each miRNA. These

P-values were used to calculate the false discovery rate (FDR) for each miRNA, which

was further used as a filter to identify significant miRNAs with a fold change ≥ 2 or ≤

0.5 and an FDR < 0.05. The volcano plots were generated using the R 3.5.3 software.

Statistical analysis

Statistical analysis was conducted using Statistical 4 (OMS, Tokyo, Japan). For

two-group comparison, Student's *t*-test was used and for multiple comparisons, ANOVA

(Tukey's multiple comparison test as post-hoc) was used. Power analysis using G Power

software (Kang et al., 2021) was conducted to determine the sample size for each

experiment to provide a power of at least 0.8 at a significance level of 0.05 ($\alpha = 0.05$, β

$= 0.2$).

Data availability

We have submitted all relevant data of our experiments to the EV-TRACK

knowledgebase (EV-TRACK ID: EV220432) (Van Deun et al., 2017).

Acknowledgements

This study was supported by JSPS KAKENHI (grant number 17H04747 and

21H03852). We are grateful to Dr. P Kent Langston (Department of Immunology,

Harvard Medical School and Evergrande Center for Immunologic Diseases, Harvard

Medical School, Boston, MA, USA) for providing insightful advice.

References

Ambattu LA, Ramesan S, Dekiwadia C, Hanssen E, Li H, Yeo LY (2020) High

frequency acoustic cell stimulation promotes exosome generation regulated by a

calcium-dependent mechanism. Commun Biol 5;3(1):553.

Bei Y, Xu T, Lv D, Yu P, Xu J, Che L, Das A, Tigges J, Toxavidis V, Ghiran I, Shah R, Li Y, Zhang Y, Das S, Xiao J (2017) Exercise-induced circulating extracellular vesicles protect against cardiac ischemia-reperfusion injury. Basic Res Cardiol. 112(4):38.

Bittel DC, Jaiswal JK (2019) Contribution of Extracellular Vesicles in Rebuilding Injured Muscles. Front Physiol. 10:828.

Cao Z, Zhang T, Sun X, Liu M, Shen Z, Li B, Zhao X, Jin H, Zhang Z, Tian Y (2019) Membrane-permeabilized sonodynamic therapy enhances drug delivery into macrophages. PLoS One. 14(6):e0217511.

Chang Q, Daly L, Bromberg J (2014) The IL-6 feed-forward loop: a driver of tumorigenesis. Semin Immunol. 26(1):48-53.

Carpi S, Polini B, Nieri D, Dubbini N, Celi A, Nieri P, Neri T (2020) Expression Analysis of Muscle-Specific miRNAs in Plasma-Derived Extracellular Vesicles from Patients with Chronic Obstructive Pulmonary Disease. Diagnostics (Basel). 10(7):502.

Chevalier X, Giraudeau B, Conrozier T, Marliere J, Kiefer P, Goupille P (2005) Safety study of intraarticular injection of interleukin 1 receptor antagonist in patients with painful knee osteoarthritis: a multicenter study. J Rheumatol. 32(7):1317-23.

Choy EH, Isenberg DA, Garrood T, Farrow S, Ioannou Y, Bird H, Cheung N, Williams

B, Hazleman B, Price R, Yoshizaki K, Nishimoto N, Kishimoto T, Panayi GS (2002)

Therapeutic benefit of blocking interleukin-6 activity with an anti-interleukin-6 receptor

monoclonal antibody in rheumatoid arthritis: a randomized, double-blind,

placebo-controlled, dose-escalation trial. Arthritis Rheum. 46(12):3143-50.

Cron L, Watt KI, Kuchel RP, Jayasooriah N, Estevez E, Petzold T, Suter CM,

Gregorevic P, Kiens B, Richter EA, James DE, Wojtaszewski JFP, Febbraio MA (2018)

Extracellular Vesicles Provide a Means for Tissue Crosstalk during Exercise. Cell

Metab. 27(1):237-251.e4.

Funes SC, Rios M, Escobar-Vera J, Kalergis AM (2018) Implications of macrophage

polarization in autoimmunity. Immunology. 154(2):186-195.

Dinarello CA (2009) Immunological and inflammatory functions of the interleukin-1

family. Annu Rev Immunol. 27:519-50.

Ding W, Ren J, Ren H, Wang D (2017) Long Noncoding RNA HOTAIR Modulates

MiR-206-mediated Bcl-w Signaling to Facilitate Cell Proliferation in Breast Cancer. Sci

Rep. 7(1):17261.

Draper DO (2014) Facts and misfits in ultrasound therapy: steps to improve your

treatment outcomes. Eur J Phys Rehabil Med. 50(2):209-16.

EL Andaloussi S, Mäger I, Breakefield XO, Wood MJ (2013) Extracellular vesicles:

biology and emerging therapeutic opportunities. Nat Rev Drug Discov. 12(5):347-57.

Emsley HC, Smith CJ, Georgiou RF, Vail A, Hopkins SJ, Rothwell NJ, Tyrrell PJ (2005) Acute Stroke Investigators. A randomised phase II study of interleukin-1 receptor antagonist in acute stroke patients. J Neurol Neurosurg Psychiatry. 76(10):1366-72.

Fan Z, Kumon RE, Park J, Deng CX (2010) Intracellular delivery and calcium transients generated in sonoporation facilitated by microbubbles. J Control Release. 142(1):31-9.

Frühbeis C, Helmig S, Tug S, Simon P, Krämer-Albers EM (2015) Physical exercise induces rapid release of small extracellular vesicles into the circulation. J Extracell Vesicles. 4:28239.

Ganesan J, Ramanujam D, Sassi Y, Ahles A, Jentzsch C, Werfel S, Leierseder S, Loyer X, Giacca M, Zentilin L, Thum T, Laggerbauer B, Engelhardt S (2013) MiR-378 controls cardiac hypertrophy by combined repression of mitogen-activated protein kinase pathway factors. Circulation. 127(21):2097-106.

Harrell CR, Jovicic N, Djonov V, Arsenijevic N, Volarevic V (2019) Mesenchymal Stem Cell-Derived Exosomes and Other Extracellular Vesicles as New Remedies in the Therapy of Inflammatory Diseases. Cells. 8(12):1605.

Hirayama   D,   Iida   T,   Nakase   H   (2017)   The   Phagocytic   Function   of

Macrophage-Enforcing Innate Immunity and Tissue Homeostasis. Int J Mol Sci.

19(1):92.

Honda H, Kondo T, Zhao QL, Feril LB Jr, Kitagawa H (2004) Role of intracellular

calcium ions and reactive oxygen species in apoptosis induced by ultrasound.

Ultrasound Med Biol. 30(5):683-92.

Huang X, Gong F, Lu Z, Zhu J, Yu Q (2020) Downregulated MiR-206 expression

promotes the proliferation and migration of macrophages by regulating IL-17A/REG3A

pathway. Eur. J. Inflamm. 18:1-8.

Ito H, Takazoe M, Fukuda Y, Hibi T, Kusugami K, Andoh A, Matsumoto T, Yamamura

T, Azuma J, Nishimoto N, Yoshizaki K, Shimoyama T, Kishimoto T (2004) A pilot

randomized trial of a human anti-interleukin-6 receptor monoclonal antibody in active

Crohn's disease. Gastroenterology. 126(4):989-96; discussion 947.

Kang H (2021) Sample size determination and power analysis using the G*Power

software. J Educ Eval Health Prof. 18:17.

Kim H, Wang SY, Kwak G, Yang Y, Kwon IC, Kim SH (2019) Exosome-Guided

Phenotypic Switch of M1 to M2 Macrophages for Cutaneous Wound Healing. Adv Sci

(Weinh). 6(20):1900513.

Krist B, Florczyk U, Pietraszek-Gremplewicz K, Józkowicz A, Dulak J (2015) The Role

of miR-378a in Metabolism, Angiogenesis, and Muscle Biology. Int J Endocrinol. 2015:281756.

Krist B, Podkalicka P, Mucha O, Mendel M, Sępioł A, Rusiecka OM, Józefczuk E, Bukowska-Strakova K, Grochot-Przęczek A, Tomczyk M, Klóska D, Giacca M, Maga P, Niżankowski R, Józkowicz A, Łoboda A, Dulak J, Florczyk-Soluch U (2020) miR-378a influences vascularization in skeletal muscles. Cardiovasc Res. 116(7):1386-1397.

Kumon RE, Aehle M, Sabens D, Parikh P, Han YW, Kourennyi D, Deng CX (2009) Spatiotemporal effects of sonoporation measured by real-time calcium imaging. Ultrasound Med Biol. 35(3):494-506.

Lentacker I, De Cock I, Deckers R, De Smedt SC, Moonen CT (2014) Understanding ultrasound induced sonoporation: definitions and underlying mechanisms. Adv Drug Deliv Rev. 72:49-64.

Lin CC, Law BF, Hettick JM (2020) Acute 4,4'-Methylene Diphenyl Diisocyanate Exposure-Mediated Downregulation of miR-206-3p and miR-381-3p Activates Inducible Nitric Oxide Synthase Transcription by Targeting Calcineurin/NFAT Signaling in Macrophages. Toxicol Sci. 173(1):100-113.

Ma C, Wang J, Liu H, Chen Y, Ma X, Chen S, Chen Y, Bihl JI, Yang YI (2018) Moderate Exercise Enhances Endothelial Progenitor Cell Exosomes Release and Function. Med Sci Sports Exerc. 50(10):2024-2032.

Madison RD, McGee C, Rawson R, Robinson GA (2014) Extracellular vesicles from a muscle cell line (C2C12) enhance cell survival and neurite outgrowth of a motor neuron cell line (NSC-34). J Extracell Vesicles. 3.

Maeshige N, Langston PK, Yuan ZM, Kondo H, Fujino H (2021) High-intensity ultrasound irradiation promotes the release of extracellular vesicles from C2C12 myotubes. Ultrasonics. 110:106243.

Mathew B, Ravindran S, Liu X, Torres L, Chennakesavalu M, Huang CC, Feng L, Zelka R, Lopez J, Sharma M, Roth S (2019) Mesenchymal stem cell-derived extracellular vesicles and retinal ischemia-reperfusion. Biomaterials. 197:146-160.

Matsuzaka Y, Tanihata J, Komaki H, Ishiyama A, Oya Y, Rüegg U, Takeda SI, Hashido K (2016) Characterization and Functional Analysis of Extracellular Vesicles and Muscle-Abundant miRNAs (miR-1, miR-133a, and miR-206) in C2C12 Myocytes and mdx Mice. PLoS One. 11(12):e0167811.

Ma X, Yamaguchi A, Maeshige N, Uemura M, Noguchi H, Kondo H, Fujino H (2022) Enhancement of astaxanthin incorporation by pulsed high-intensity ultrasound in LPS-stimulated macrophages. J Med Ultrason (2001). 49(2):125-132.

Moore KJ, Sheedy FJ, Fisher EA (2013) Macrophages in atherosclerosis: a dynamic balance. Nat Rev Immunol. 13(10):709-21.

Nishimoto N, Yoshizaki K, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, Hashimoto J, Azuma J, Kishimoto T (2004) Treatment of rheumatoid arthritis with humanized anti-interleukin-6 receptor antibody: a multicenter, double-blind, placebo-controlled trial. Arthritis Rheum. 50(6):1761-9.

Piccioli P, Rubartelli A (2013) The secretion of IL-1β and options for release. Semin Immunol. 25(6):425-9.

Raposo G, Stoorvogel W (2013) Extracellular vesicles: exosomes, microvesicles, and friends. J Cell Biol. 200(4):373-83.

Rome S, Forterre A, Mizgier ML, Bouzakri K (2019) Skeletal Muscle-Released Extracellular Vesicles: State of the Art. Front Physiol. 10:929.

Rückerl D, Jenkins SJ, Laqtom NN, Gallagher IJ, Sutherland TE, Duncan S, Buck AH, Allen JE (2012) Induction of IL-4Rα-dependent microRNAs identifies PI3K/Akt signaling as essential for IL-4-driven murine macrophage proliferation in vivo. Blood. 120(11):2307-16.

Savina A, Furlán M, Vidal M, Colombo MI (2003) Exosome release is regulated by a calcium-dependent mechanism in K562 cells. J Biol Chem. 278(22):20083-90.

Scheffer LL, Sreetama SC, Sharma N, Medikayala S, Brown KJ, Defour A, Jaiswal JK

(2014) Mechanism of Ca^{2+}-triggered ESCRT assembly and regulation of cell membrane

repair. Nat Commun. 5:5646.

Shao J, Li S, Liu Y, Zheng M (2020) Extracellular vesicles participate in

macrophage-involved immune responses under liver diseases. Life Sci. 240:117094.

Srere PA (1963) Citryl-CoA. An substrate for the citrate-cleavage enzyme. Biochim

Biophys Acta. 73:523-525.

Tanaka T, Narazaki M, Kishimoto T (2014) IL-6 in inflammation, immunity, and

disease. Cold Spring Harb Perspect Biol. 6(10):a016295.

Taylor J, Azimi I, Monteith G, Bebawy M (2020) Ca^{2+} mediates extracellular vesicle

biogenesis through alternate pathways in malignancy. J Extracell Vesicles.

9(1):1734326.

Ter-Ovanesyan D, Norman M, Lazarovits R, Trieu W, Lee JH, Church GM, Walt DR

(2021) Framework for rapid comparison of extracellular vesicle isolation methods. Elife.

10:e70725.

Trovato E, Di Felice V, Barone R (2019) Extracellular Vesicles: Delivery Vehicles of

Myokines. Front Physiol. 10:522.

Ueno M, Maeshige N, Hirayama Y, Yamaguchi A, Ma X, Uemura M, Kondo H, Fujino

H (2021) Pulsed ultrasound prevents lipopolysaccharide-induced muscle atrophy through inhibiting p38 MAPK phosphorylation in C2C12 myotubes. Biochem Biophys Res Commun. 570:184-190.

Van Deun J, Mestdagh P, Agostinis P, Akay Ö, Anand S, Anckaert J, Martinez ZA, Baetens T, Beghein E, Bertier L, Berx G, Boere J, Boukouris S, Bremer M, Buschmann D, Byrd JB, Casert C, Cheng L, Cmoch A, Daveloose D, De Smedt E, Demirsoy S, Depoorter V, Dhondt B, Driedonks TA, Dudek A, Elsharawy A, Floris I, Foers AD, Gärtner K, Garg AD, Geeurickx E, Gettemans J, Ghazavi F, Giebel B, Kormelink TG, Hancock G, Helsmoortel H, Hill AF, Hyenne V, Kalra H, Kim D, Kowal J, Kraemer S, Leidinger P, Leonelli C, Liang Y, Lippens L, Liu S, Lo Cicero A, Martin S, Mathivanan S, Mathiyalagan P, Matusek T, Milani G, Monguió-Tortajada M, Mus LM, Muth DC, Németh A, Nolte-'t Hoen EN, O'Driscoll L, Palmulli R, Pfaffl MW, Primdal-Bengtson B, Romano E, Rousseau Q, Sahoo S, Sampaio N, Samuel M, Scicluna B, Soen B, Steels A, Swinnen JV, Takatalo M, Thaminy S, Théry C, Tulkens J, Van Audenhove I, van der Grein S, Van Goethem A, van Herwijnen MJ, Van Niel G, Van Roy N, Van Vliet AR, Vandamme N, Vanhauwaert S, Vergauwen G, Verweij F, Wallaert A, Wauben M, Witwer KW, Zonneveld MI, De Wever O, Vandesompele J, Hendrix A (2017) EV-TRACK: transparent reporting and centralizing knowledge in extracellular vesicle research. Nat Methods. 14(3):228-232.

van Niel G, D'Angelo G, Raposo G (2018) Shedding light on the cell biology of extracellular vesicles. Nat Rev Mol Cell Biol. 19(4):213-228.

Whitham M, Parker BL, Friedrichsen M, Hingst JR, Hjorth M, Hughes WE, Egan CL,

Cron L, Watt KI, Kuchel RP, Jayasooriah N, Estevez E, Petzold T, Suter CM,

Gregorevic P, Kiens B, Richter EA, James DE, Wojtaszewski JFP, Febbraio MA (2018)

Extracellular Vesicles Provide a Means for Tissue Crosstalk during Exercise. Cell

Metab. 27(1):237-251.e4.

Wu H, Zhang T, Pan F, Steer CJ, Li Z, Chen X, Song G (2017) MicroRNA-206 prevents

hepatosteatosis and hyperglycemia by facilitating insulin signaling and impairing

lipogenesis. J Hepatol. 66(4):816-824.

Xin D, Li T, Chu X, Ke H, Yu Z, Cao L, Bai X, Liu D, Wang Z (2020) Mesenchymal

stromal cell-derived extracellular vesicles modulate microglia/macrophage polarization

and protect the brain against hypoxia-ischemic injury in neonatal mice by targeting

delivery of miR-21a-5p. Acta Biomater. 113:597-613.

Yamaguchi A, Maeshige N, Yan J, Ma X, Uemura M, Matsuda M, Nishimura Y,

Hasunuma T, Kondo H, Fujino H, Yuan ZM (2023) Skeletal myotube-derived

extracellular vesicles enhance itaconate production and attenuate inflammatory

responses of macrophages. Front Immunol. 14:1099799.

Yokota S, Miyamae T, Imagawa T, Iwata N, Katakura S, Mori M, Woo P, Nishimoto N,

Yoshizaki K, Kishimoto T (2005) Therapeutic efficacy of humanized recombinant

anti-interleukin-6 receptor antibody in children with systemic-onset juvenile idiopathic

arthritis. Arthritis Rheum. 52(3):818-25.

Zeghimi A, Escoffre JM, Bouakaz A (2015) Role of endocytosis in

sonoporation-mediated membrane permeabilization and uptake of small molecules: a

electron microscopy study. Phys Biol. 12(6):066007.

Zeng M, Zhu L, Li L, Kang C (2017) miR-378 suppresses the proliferation, migration

and invasion of colon cancer cells by inhibiting SDAD1. Cell Mol Biol Lett. 22:12.

Zhang H, Yu H, Yang H, Zhan Y, Liu X (2021) miR-378-3p alleviates contusion spinal

cord injury by negatively regulating ATG12. Int J Exp Pathol. 102(4-5):200-208.

Zhou Y, Shi J, Cui J, Deng CX (2008) Effects of extracellular calcium on cell

membrane resealing in sonoporation. J Control Release. 126(1):34-43.

Zomer A, Vendrig T, Hopmans ES, van Eijndhoven M, Middeldorp JM, Pegtel DM

(2010) Exosomes: Fit to deliver small RNA. Commun Integr Biol. 3(5):447-50.

Table 1 Upregulated miRNAs in myotube-derived extracellular vesicles by ultrasound

irradiation.

miRNA	logFC	FDR
miR-193a-3p	4.012954262	0.000742983
miR-138-5p	3.800680285	0.00141537
miR-223-3p	3.462174268	0.008790418
miR-362-3p	2.195592968	0.009437513
miR-34a-5p	2.105459261	0.009437513
miR-675-3p	1.876128033	0.009966738
miR-106b-5p	1.862814288	0.009966738
miR-30b-5p	1.566332803	0.011740976
miR-188-5p	1.559409694	0.015175451
miR-133b-3p	1.48161012	0.016332293
miR-1a-3p	1.450599776	0.016622751
miR-30a-5p	1.367918225	0.019014654
miR-27b-3p	1.330089099	0.01987041

668 **Table 2** Downregulated miRNAs in myotube-derived extracellular vesicles by

669 ultrasound irradiation.

miRNA	logFC	FDR
miR-128-2-5p	-3.254069169	0.000118723
miR-184-3p	-2.779741068	0.000118723
miR-615-3p	-2.393817511	0.000263853
miR-344d-3p	-2.324403579	0.00041415
miR-344d-4p	-2.308193263	0.000742983
miR-1964-3p	-2.267802775	0.00141537
miR-320-3p	-2.212812155	0.00141537
miR-128-3p	-2.031468208	0.003795561
miR-351-3p	-1.963143876	0.004305251
miR-1198-5p	-1.733517014	0.009437513
miR-501-3p	-1.72302403	0.009643704
miR-222-3p	-1.547695372	0.009966738
let-7a-5p	-1.538094973	0.009966738
miR-423-5p	-1.404879603	0.019014654

Figure 1 The cytotoxicity of ultrasound (US) irradiation on myotubes was investigated. Viability of myotubes was assessed by (A) MTT assay and (B) Zombie Red^{TM} staining at 24 h after US irradiation. (red: Zombie Red^{TM}, blue: DAPI). (C) Total protein content was measured by the Bradford method at 24 h after US irradiation. (D) Energy metabolism in C2C12 myotubes was measured by citrate synthase assay at 24 h after US irradiation. The US intensities of 1.0 W/cm^2, 2.0 W/cm^2, and 3.0 W/cm^2 were tested. Data are expressed as mean $\pm$ SEM. n = 4.

Figure 2 Characterization of extracellular vesicles (EVs) from ultrasound (US)-treated/untreated myotubes. EVs were isolated using ExoQuick reagent 12 h after US irradiation. (A) EV concentration in each group was quantified by a qNano system. (B) Size distribution of EVs in each group was investigated by a qNano system. The mode value was indicated as a red line. The US intensities of 1.0 W/cm^2, 2.0 W/cm^2, and 3.0 W/cm^2 were tested. Data are expressed as mean $\pm$ SEM. *p $<$ 0.01, vs. Control. n = 6.

Figure 3 Ca^{2+} mediates the promotive effect of ultrasound (US) on extracellular vesicle (EV) release from myotubes. (A) Intracellular Ca^{2+} levels were measured after US irradiation. The US intensities of 1.0 W/cm^2, 2.0 W/cm^2, and 3.0 W/cm^2 were tested. (B) Cell culture with Ca^{2+}-free medium decreased the intracellular Ca^{2+} level and canceled the facilitating effect of US on Ca^{2+} uptake by myotubes. (C) Cell culture with Ca^{2+}-free medium inhibited the facilitating effect of US on EV release from myotubes. (D) Size distribution of EVs in each group. EV concentration and size distribution were

quantified by a qNano system. (E) Cytotoxicity of cell culture with Ca^{2+}-free medium

was investigated by MTT assay. Control: untreated; US: 3.0 W/cm^2 US treatment;

Ca(-): Ca^{2+}-free culture; USCa(-): 3.0 W/cm^2 US treatment and Ca^{2+}-free culture. Data

are expressed as mean ± SEM. *p < 0.01 vs. Control, †p < 0.05 vs. 1W, ‡p< 0.01 vs.

US. n = 4.

Figure 4 Anti-inflammatory effect of extracellular vesicles (EVs) from ultrasound

(US)-treated myotubes on bone marrow-derived macrophages (BMDMs). (A) The

mRNA expression levels of *Il-1b* and *Il-6* were measured by qPCR. LPS: LPS-treated

BMDMs; CM: BMDMs treated with C2C12 conditioned medium and LPS; US-CM:

BMDMs treated with US-irradiated C2C12 conditioned medium and LPS; EV-depleted

CM: BMDMs treated with EV-depleted C2C12 conditioned medium and LPS;

USEV-depleted: BMDMs treated with EV-depleted C2C12 (US-irradiated) conditioned

medium and LPS. (B) When the concentration of EVs are equated, the enhancement of

anti-inflammatory effect of EVs by US was not observed. LPS: LPS-treated BMDMs;

CM: BMDMs treated with myotube EVs and LPS; diluted US-CM: BMDMs treated

with US-EVs at the same concentration as the EV group and LPS. (C) Cytotoxicity of

C2C12 conditioned medium on BMDMs was investigated by MTT assay. (D) EV

concentration in each condition was measured by a qNano system. Data are expressed

as mean ±SEM. *p < 0.05 vs. LPS, †p < 0.01 vs. CM, ‡p< 0.01 vs. Control, §p< 0.01

vs. US. n = 4.

Figure 5 miRNA-sequencing analysis in extracellular vesicles (EVs) from ultrasound

(US)-treated/untreated C2C12 myotubes. (A) miRNA characterization in EVs from

US-treated/untreated myotubes. (B) Percentage of miRNAs that were common to the control and US groups and those that were not. (C) Volcano plot of differentially expressed RNAs in the control group vs. US group. Blue dots represent miRNAs with statistically significant difference and red dots show miRNAs with no statistically significant difference between the control group vs. US group. (D) Top 10 abundant miRNAs and their proportion to total miRNA content in each group. n = 3.

Figure 6 In vitro ultrasound (US) irradiation system. The culture dish was placed on the probe of a US transducer (SONICTIZER SZ-100, Minato Medical Science Co., Ltd., Japan). US waves are sent out from the probe placed under the culture dish, and surplus energy is absorbed by the silicon.

C2C12 myotubes

Silicon

Culture dish

Probe head

Culture medium

US probe

US energy

Cell monolayer

Coupling gel