Visual mental imagery in typical imagers and in aphantasia: A millimeter-scale 7-T fMRI study

Jianghao Liu[1,2*], Minye Zhan[3], Dounia Hajhajate[1,4], Alfredo Spagna[5], Stanislas Dehaene[3,6], Laurent Cohen[1,7], Paolo Bartolomeo[1]

[1]Sorbonne Université, Inserm, CNRS, Paris Brain Institute, ICM, Hôpital de la Pitié-Salpêtrière, 75013 Paris, France

[2]Dassault Systèmes, Vélizy-Villacoublay, France

[3]Cognitive Neuroimaging Unit, Université Paris-Saclay,CEA, INSERM, CNRS ELR9003, NeuroSpin center, 91191 Gif/Yvette, France

[4]IRCCS SYNLAB SDN, Via E. Gianturco 113, 80143 Naples, Italy

[5]Department of Psychology, Columbia University in the City of New York, NY, 10027, USA

[6]Collège de France, Université Paris-Sciences-Lettres (PSL), 11 Place Marcelin Berthelot, 75005 Paris, France

[7]AP-HP, Hôpital de la Pitié Salpêtrière, Fédération de Neurologie, Paris, France

*Correspondence: jianghao.liu@icm-institute.org

Abstract

Most of us effortlessly describe visual objects, whether seen or remembered. Yet, around 4% of people report congenital aphantasia: they struggle to visualize objects despite being able to describe their visual appearance. What neural mechanisms create this disparity between subjective experience and objective performance? Aphantasia can provide novel insights into conscious processing and awareness. We used ultra-high field 7T fMRI to establish the neural circuits involved in visual mental imagery and perception, and to elucidate the neural mechanisms associated with the processing of internally generated visual information in the absence of imagery experience in congenital aphantasia. Ten typical imagers and 10 aphantasic individuals performed imagery and perceptual tasks in five domains: object shape, object color, written words, faces, and spatial relationships. In typical imagers, imagery tasks activated left-hemisphere frontoparietal areas, the relevant domain-preferring areas in the ventral temporal cortex partly overlapping with the perceptual domain-preferring areas, and a domain-general area in the left fusiform gyrus (the Fusiform Imagery Node). The results were valid for each individual participant. In aphantasic individuals, imagery activated similar visual areas, but the Fusiform Imagery Node was functionally disconnected from frontoparietal areas. Our results unveil the domain-general and domain-specific circuits of visual mental imagery, their functional disorganization in aphantasia, and support the general hypothesis that conscious visual experience - whether perceived or imagined - depends on the integrated activity of high-level visual cortex and frontoparietal networks.

Introduction

Visual mental imagery is the ability to experience visual information in the absence of actual external stimuli. However, approximately 4% of individuals report experiencing weak or absent visual mental imagery (Dance et al., 2022), a condition known as congenital aphantasia (Zeman et al., 2015). The nature of aphantasia remains elusive. Some perspectives suggest that aphantasia is a deficit in introspection or awareness of internal visual imagery (Liu & Bartolomeo, 2023), while others propose it as a deficiency in voluntarily-generated mental imagery (Zeman et al., 2015), or in general mental imagery capacities (Keogh & Pearson, 2018). Surprisingly, individuals with congenital aphantasia can typically provide accurate answers from memory to questions on the visual appearance of objects, despite the absence of subjective imagery experience. For example, these individuals can correctly indicate which fruit is darker red between strawberries or cherries (Liu & Bartolomeo, 2023), and perform many imagery-related tasks such as visual working memory and mental rotation (Pounder et al., 2022). Thus, aphantasia provides a unique window into the mechanisms underlying conscious processing and awareness. Why don't aphantasic individuals experience visual mental images?

Models of visual mental imagery are structured around two primary components: the frontoparietal (FP) cortex and the visual cortex. However, models diverge on the specific portions of the visual cortex crucial for visual mental imagery. While some emphasize the importance of early visual areas (EVA)(Pearson, 2019), others highlight the role of the high-level visual areas in the ventral temporal cortex (VTC)(Spagna et al., 2021). These competing models make different predictions on the possible neural underpinnings of aphantasia.

Many neuroimaging studies predominantly focused on the primary visual cortex (Kosslyn et al., 2006). For example, individual vividness was related to the activation level of the V1 (Cui et al., 2007) and imagined stimuli could be decoded or reconstructed from the representational pattern of the EVA (Naselaris et al., 2015; Senden et al., 2019; Thirion et

al., 2006), possibly driven by the signal from the deep cortical layers of V1 (Bergmann et al.,

2024). On the other hand, evidence from acquired brain lesions in neurological patients

demonstrating preserved imagery abilities after lesions to the EVA or their connections

(Bartolomeo et al., 1998, 2020) strongly suggests that the EVA are not necessary for visual

mental imagery. If, however, visual mental imagery experience does rely on EVA activity,

then this activity should be dysfunctional in aphantasia.

A recent meta-analysis of 27 imagery fMRI studies (Spagna et al., 2021) highlighted

imagery-related activity in FP regions and in a specific region of the left hemisphere fusiform

gyrus, which was active independent of the imagery domain. Spagna et al. labeled this

region the Fusiform Imagery Node (FIN). In line with the localization of the FIN, lesion

studies revealed domain-general imagery deficits following extensive damage to the left

temporal lobe (Bartolomeo et al., 2002; Liu et al., 2022; Moro et al., 2008; Thorudottir et al.,

2020). Nevertheless, no study has yet examined the functional properties of the putative FIN

during visual mental imagery, or its possible dysfunction in aphantasia.

A crucial yet understudied aspect is the role of domain-preferring VTC cortical

patches in visual mental imagery. The VTC contains cortical patches with relatively selective

activity for specific perceptual domains, such as faces, words, and colors (Cohen et al.,

2000; Kanwisher et al., 1997; Lafer-Sousa et al., 2016). Early fMRI studies showed that

imagery of faces and places reactivated the fusiform face area (FFA) and the

parahippocampal place area (PPA), respectively (Ishai et al., 2000; O'Craven & Kanwisher,

2000). Also, color information can be decoded from a color-biased region during color

imagery (Bannert & Bartels, 2018). Lesion studies demonstrated that brain damage to the

temporal cortex (but not to the occipital cortex) can result in deficits in imagery that are

specific to particular visual domains, such as object shape, object color, written words, faces,

and spatial relationships (Bartolomeo, 2002; Goldenberg, 1993). These findings underscore

the necessity for a more systematic investigation of domain-preferring cortex in visual mental

imagery. Do these regions contribute to imagery vividness in typical imagery or to its

absence in aphantasia?

Finally, a further open question concerns the role of FP networks. Clinical

(Bartolomeo, 2007), neuroimaging (Chica et al., 2012) and neurophysiological (Liu et al.,

2023; Spagna et al., 2022) evidence highlights the crucial role of FP network activity in

conscious perception. Can FP dysfunction result in the lack of experiential correlates of

access to offline visual information in aphantasia? This could be characterized by an

alteration of functional brain networks akin to those observed in some neurodevelopmental

disorders (Sokolowski & Levine, 2023). For example, Milton et al (2021) reported reduced

resting-state functional connectivity between prefrontal areas and the visual cortex in

aphantasic participants, compared to individuals reporting high imagery vividness. However,

it remains uncertain whether aphantasia displays impaired connectivity during attempted

mental imagery.

Neuroimaging studies have encountered challenges in distinguishing between

domain-general and domain-specific mechanisms in visual mental imagery for the following

reasons: i) the difficulty in demonstrating a causal link between activations (e.g. in EVA) and

imagery processes (Bartolomeo et al., 2020), ii) the dearth of studies that examine several

domains with naturalistic stimuli, which are essential for eliciting activity in high-level visual

cortex including both the FIN and domain-preferring areas, and iii) the limited spatial

resolution of conventional 3T fMRI, along with techniques like group averaging, which

hindered the ability to distinguish closely packed domain-preferring visual regions (Saxe et

al., 2006). For instance, the face, word, and body-preferring areas in the VTC are situated in

close proximity (Grill-Spector & Weiner, 2014). Moreover, given the individual variability in

the topographical organization of the VTC (Conway, 2018; Grill-Spector & Weiner, 2014),

assessing the degree of overlap of activations between imagery and perception requires the

use of methods with high spatial resolution in individual participants. Compared to functional

neuroimaging, lesion studies possess causal power (Bartolomeo et al., 2020), but are

affected by limitations including limited spatial resolution, disruption of white-matter

connections beyond the gray-matter lesions, and lesion-induced plasticity or reorganization.

Here, we aim to i) identify the neural circuits involved in domain-general and domain-

specific voluntary imagery in individual participants; ii) compare them with those engaged in

visual perception; iii) examine the neural processes associated with accurate information

processing in the absence of experiential correlates in aphantasia. We circumvent the

above-mentioned limitations by i) using ultra-high field 7-Tesla fMRI during tightly matched

imagery and perceptual tasks in five visual domains suggested by lesion studies

(Goldenberg, 1993): object shape, object color, written words, faces, and spatial

relationships, ii) studying typical imagers and individuals with aphantasia. Crucially, our tasks

involve visualizing real-world stimuli retrieved from long-term memory, without any visual

cues. We predicted that, under the EVA hypothesis, typical imagers would exhibit normal

activities including EVA activation, representational content and connectivity, whereas those

with aphantasia would demonstrate dysfunctional EVA activity or connectivity. Alternatively,

aphantasia could be associated with abnormal activity or connectivity of high-level visual

cortex, either in the domain-general or domain-preferring VTC, as suggested by lesion

localization in neurological patients with impaired visual mental imagery.

Results

Ten typical imagers (mean age = 29.28, SD = 8.47, 6 female) and 10 individuals with

congenital aphantasia (mean age = 28.69, SD = 8.27, 6 female) performed our enhanced

Batterie Imagerie / Perception - eBIP (Liu & Bartolomeo, 2023) while undergoing 7T fMRI

scanning at 1.2 mm isotropic resolution with large brain coverage. All participants were

native French speakers. The eBIP comprises 162 trials, each one testing participants'

performance on: (1) Mental imagery in response to auditory questions in five distinct

domains (Fig. 1A-E): shapes, colors, written words, faces, and spatial relationships of cities

on an imagined map of France; (2) a matched control task using abstract words to engage

non-visual semantics while minimizing the involvement of visual imagery (Fig. 1F); (3) a

perceptual task using the same items as the imagery tasks (Fig. 1G) but with the

simultaneous presentation of auditory questions and the relevant pictorial items. Within the

imagery and perceptual tasks, trials from the five domains were fully randomly presented. In

the imagery task, participants rated the vividness of their visual imagery at the end of each

trial using a 4-point Likert scale where a score of 4 indicated highly vivid imagery. In the

abstract word task and the perception task, the rating of visual vividness was substituted

with a confidence rating using a 4-point Likert scale, where a score of 4 represented

absolute certainty. Accuracy and response time (RT) were recorded.

We first present the behavioral results, then the fMRI findings in the domain-

preferring VTC areas, in the domain-general FIN, and finally in the EVA. For each region, we

compare perception vs. imagery, and typical imagers vs. aphantasic participants, using

univariate and multivariate methods, plus the study of functional connectivity.

Behavioral results

For typical imagers, the average trial-by-trial vividness score was 3.52 on a scale of 1 to 4,

while for aphantasic individuals the average score was 1.11 (Bayes Factor, BF = 5.387×10^{11},

see Fig. S1B for all behavioral results). Nevertheless, aphantasic individuals exhibited

comparable levels of accuracy to typical imagers in both imagery (Bayesian repeated

measures ANOVA with the factors of Group x Domain, the main Group effect BF = 0.25

without interaction) and perception (BF = 0.26 without interaction). Their RTs were 0.21s

slower on the imagery tasks and 0.43s slower on the perception tasks (BF = 30 and 85,

respectively), and they had lower confidence in their responses on perceptual tasks (BF =

3.59), consistent with similar findings from a more extensive cohort of 117 participants (Liu &

Bartolomeo, 2023). Importantly, however, there was no evidence of group difference on

either accuracy (BF = 0.46), RTs (BF = 0.41) or confidence scores (BF = 1.63) in the

abstract words task, suggesting that the group difference was specific to visual items.

Fig 1. Imagery and Perceptual Tasks.
Examples of trials of the eBIP. The eBIP comprises imagery tasks of five domains: shapes, colors, letters, faces, and spatial relationships (A-E), a control task with abstract words (F), and a perception task (G) in five audio-visual domains using the same items as in the imagery tasks. In the imagery tasks, participants heard a word indicating a particular imagery domain (e.g., "shape"), followed by 2 words (e.g. "beaver", "fox"). Participants were instructed to generate and maintain mental images as vivid as possible for each of these 2 words. Eight seconds after the second item, participants heard an attribute word (e.g. "long"). They then pressed one of two buttons indicating which of the items was most closely associated with the attribute (e.g. which of the animals they associate with the attribute "long"). In the shape imagery task, participants had to decide which of the 2 words designated a longer or rounder item. In the color imagery task, participants had to decide which fruit or vegetable had darker or lighter color. In the letter imagery task, participants had to imagine the shape of French words in lowercase and had to decide which word had ascenders (e.g., t, l, d) or descenders (e.g. j, p, y). In the famous faces imagery task, participants had to decide which of 2 named celebrities had a more round or oval face. In the map-of-France imagery task, participants had to decide which of 2 cities was located to the left or the right of Paris. In the non-imagery abstract word task, participants had to decide which of two abstract words (e.g., "routine", "convention") was semantically closer to a third word (e.g. "society"). In the perception task, the same stimuli used for the imagery tasks were presented in an audio-visual format.

Domain-preferring activations overlap during imagery and perception

In both imagery and perception, we localized domain-preferring regions by contrasting

activity elicited by each domain minus the other four domains in order to answer three

questions: Does local domain-preference prevail in imagery as it does in perception? Are

domain-preferring regions the same in imagery and in perception? Do typical and aphantasic

participants differ in those respects? There was substantial individual variability in the

location of activations, and each domain could activate multiple cortical patches, including in

the VTC, consistent with previous 3T and 7T studies with single-subject analyses (Zhan et

al., 2023; Zhen et al., 2015). For example, face-preferring fusiform patches showed

substantial individual variability during perception (Fig. S3B). Thus, we report individual

domain-preferring maps with a summary of regions for each group (thresholded at p<0.001

uncorrected for all individual maps, no data smoothing, and cluster size > 12 voxels).

In typical imagers, during **imagery tasks** (Fig. 2A displays the results for a

representative typical imager and see Fig. S3 for other individual activation maps; see Table

1 for a summary), i) shape imagery activated the fusiform gyrus (FG, 5 left and 2 bilateral out

of 10 participants) and lateral occipital complex (LOC, 3 right, 2 bilateral); ii) color imagery

activated medial FG (6 left, 1 right, 2 bilateral), parahippocampal gyrus (PHG, 4 left, 1 right,

3 bilateral), posterior occipitotemporal sulcus (OTS, 4 left, 6 bilateral), orbitofrontal cortex

(OFC, 2 left, 7 bilateral), inferior frontal sulcus (IFS, 2 left, 7 bilateral); iii) word imagery

activated posterior OTS (2 left, 1 right, 7 bilateral), lateral occipitotemporal cortex (LOTC, 4

left, 4 bilateral), dorsolateral prefrontal cortex (dlPFC) and Intraparietal sulcus (IPS); iv) face

imagery activated OTS (FFA, 2 left, 7 bilateral), middle superior temporal gyrus (mSTG, 2

left, 6 bilateral), ventral posterior cingulate cortex (vPCC, 10 bilateral), ventral medial

prefrontal cortex (vmPFC, 10 bilateral), OFC (adjacent and more lateral to the color imagery

OFC areas, 2 right, 7 bilateral); v) map imagery activated parahippocampal gyrus

(parahippocampal place area, PPA, 1 left, 1 right, 6 bilateral), LOC (7 left, 3 bilateral),

posterior parietal area (1 left, 9 bilateral), precuneus (10 bilateral), vPCC (10 bilateral).

Table 1. Number of participants exhibiting domain-specific activations during
imagery.

		Typical imagers				Aphantasic individuals			
		Total	Bilateral	Left	Right	Total	Bilateral	Left	Right
shape imagery	FG	7	2	5	0	6	2	3	1
	LOC	5	2	0	3	8	3	5	0
color imagery	FG	9	1	7	1	7	2	5	0
	PHG	8	3	4	1	4	2	2	0
	pOTS	10	6	4	0	7	2	5	0
	OFC	9	7	2	0	8	5	1	2
	IFS	9	7	2	0	8	6	2	0
word imagery	OTS	10	7	2	1	9	4	5	0
	LOTC	8	4	4	0	9	5	4	0
face imagery	OTS	9	7	2	0	10	9	1	0
	mSTG	8	6	2	0	9	3	6	0
	vPCC	10	10	0	0	10	10	0	0
	vmPFC	10	10	0	0	10	10	0	0
	OFC	9	7	0	2	8	5	2	1
map imagery	LOC	10	3	7	0	8	7	1	0
	p. parietal	10	9	1	0	10	10	0	0
	PPA	8	6	1	1	10	9	1	0
	Precuneus	10	10	0	0	10	10	0	0
	vPCC	10	10	0	0	10	10	0	0

See a list of glossary in supplementary materials. IFG: fusiform gyrus; IFS: inferior frontal sulcus; LOC: lateral occipital complex; LOTC: lateral occipitotemporal cortex; mSTG: middle superior temporal gyrus; PHG: parahippocampal gyrus; pOTS: posterior occipitotemporal sulcus; OFC: orbitofrontal cortex; vmPFC: ventral medial prefrontal cortex; vPCC: ventral posterior cingulate cortex; PPA: parahippocampal place area.

As expected, **perception tasks** evoked activity in both EVA and higher level visual areas (see Fig. 2B for a representative typical imager): i) shape perception activated the left LOC (Malach et al., 1995) and the bilateral medial FG (Kourtzi & Kanwisher, 2001); ii) color perception activated three patches in the bilateral medial FG, as well as the PHG (Lafer-Sousa et al., 2016) and the ventral OFC; iii) word perception activated the bilateral posterior FG (Cohen et al., 2000), and bilateral FP regions; iv) face perception activated the right OTS and bilateral amygdala (Kanwisher et al., 1997); v) the map of France activated the PPA, the bilateral vPCC, precuneus, and angular gyri, see (Epstein et al., 1999). Group-averaged results showing inter-individual consistency (if any) are displayed in Fig. S4 and Table S1 (for imagery) & S2 (for perception).

Attempted imagery in <u>aphantasia</u> evoked clear activations in domain-preferring

regions (see Fig. 2A for a representative aphantasic participant, and Fig. S3 for the

remaining participants; also see Table S5 for group-averaged results). Specifically, shape

imagery activated the FG (3 left, 1 right, 2 bilateral out of 10 participants) and the LOC (5

left, 3 bilateral); color imagery activated FG (5 left, 2 bilateral), the pOTS (5 left, 2 bilateral),

the OFC (1 left, 2 right, 5 bilateral), the IFS (2 left, 6 bilateral); word imagery activated the

OTS (5 left, 4 bilateral) and the LOTC (4 left, 5 bilateral); face imagery activated the OTS (1

left, 9 bilateral), mSTG (6 left, 3 bilateral), vPCC (10 bilateral), vmPFC (10 bilateral), OFC (2

left, 1 right, 5 bilateral); map imagery activated PPA (1 left, 9 bilateral), LOC (1 left, 7

bilateral), posterior parietal cortex (10 bilateral), precuneus (10 bilateral), vPCC (10 bilateral).

On **perception tasks**, aphantasic individuals activated similar domain-specific regions as

typical imagers (Table S6).

<u>Comparison between aphantasic and typical participants</u>

First, we compared **the extent of activation** between the two groups. We computed

the number of activated voxels in domain-preferring areas, taking into account the individual

variability in the location of activated regions. For each individual, we extracted the number

of voxels in the volumetric intersection between domain-specific activations and EVA, the

ventral visual pathway, and the dorsal visual pathway (see Methods for the specific areas),

respectively. Individual domain-preferring patches consisted of unsmoothed voxels. We

conducted a 3-way Bayesian ANOVA with the factor of Group x Region (including EVA and

the ventral and dorsal cortical visual pathways) x Domain. There was moderate evidence

supporting the absence of difference between aphantasic individuals and typical imagers,

with no interaction with domains (Main Group effect, BF = 0.21 without interaction effects;

see Fig.S5).

Second, we compared the **overlap between activations induced by imagery and**

by perception between the two groups. When studying unsmoothed individual patches, for

all five domains, there was some overlap between activations induced by imagery and by

perception, in high-level VTC visual areas, in FP, and in subcortical regions (e.g. amygdala

for faces). In contrast such overlap did not exist in EVA, despite domain-dependent EVA

activation during perception (Fig. 2B). For example, color perception evoked activation in

three color patches located bilaterally along the medial FG, the most anterior belonging to

the VTC and the other two to EVA, as previously described by Lafer-Sousa et al. (2016).

During color imagery tasks, only the anterior or central color patches were activated,

whereas the posterior color patch did not show any significant activation. In addition, the

location of areas where imagery and perception overlap was domain-dependent; e.g. colors

and faces engaged ventral patches whereas maps involved more dorsal patches (Fig. 2C).

In summary, typical imagers and individuals with aphantasia displayed similar

imagery-related activity in the relevant domain-preferring areas, with some overlapping with

perception-related activity in VTC patches.

Fig 2. Patterns of BOLD response in domain-preferring regions during imagery, and their overlap with perception in typical imagers and aphantasic individuals.

   (A) Domain-specific activation during single imagery domains in a representative typical imager (Typ 05) and in a representative aphantasic individual (Aph10), from the contrast of one domain > the remaining 4 domains. All individual-level maps were thresholded at p<0.001, without any data smoothing, and cluster size > 12 voxels. The different domains and the FIN are color-coded. Note that aphantasic individuals showed clear domain-specific imagery activation in high-level visual areas.

   (B) Domain-specific activations during perception (orange), during imagery (blue), and their overlap (purple), in a representative typical imager (Typ 05) and in a representative aphantasic individual (Aph 10), unsmoothed data. Only the ventral

view is displayed. Pc, Cc and Ac indicate the posterior, central, and anterior color-
biased regions, respectively. Face patches includes occipital face patch (OFA),
fusiform face patches (FFA1 & FFA2) and anterior face patch (AFP). For all five
domains, imagery-related activations overlapped with some of the perceptual-related
activations in high-level VTC visual areas, but not with EVA despite stimulus-
dependent EVA activation during perception.
(C) In typical imagers, box-and-whisker plots of the number of voxels showing domain-
specific activation during both mental imagery and perception, in EVA (V1, V2, and
V3), and in the ventral and dorsal cortical visual pathways. Boxplot shows values of
median, upper quartile, lower quartile, maximum and minimum, respectively. Such
unsmoothed voxels were present only in high-level visual areas, dependent on
domain, but not in EVA.
307
308
Univariate and multivariate activation of the FIN does not differ between typical
and aphantasic individuals

We then studied the activation of the FIN, a left VTC region which according to a meta-

analysis (Spagna et al., 2021) is reproducibly activated during imagery. We compared the

two groups using univariate measures, and multivariate methods probing representational

format, in order to answer three questions: Do the current data support the hypothesis that a

specific left VTC region is activated during imagery irrespective of content? Does this region

also contain domain-specific information? Do aphantasic individuals differ from typical

imagers in those respects?

FIN activation amplitude. In typical imagers, we identified domain-general regions by

comparing all averaged imagery domains minus the abstract word task. This contrast

showed a left-predominant set of regions (see Fig. 3A and Table S1) including the bilateral

inferior frontal gyrus (IFG) and dorsal premotor cortex (PMd), the left intraparietal sulcus,

and a region in the left posterior OTS (-41, -55, -10). This latter region was located at

coordinates nearly identical to those of the FIN, as identified in our previous meta-analysis of

fMRI studies of mental imagery (-40, -55, -11) (Spagna et al., 2021). **In aphantasia**, the

same contrast showed activations in the left IFG-IPS network, right dorsal FP areas, and,

importantly, also in the FIN (-45,-54,-10) (Fig. 3A and Table S5). How is the FIN activated

during perception compared to imagery? If so, does activation differ between groups? We

performed a ROI-based three-way Bayesian repeated measure ANOVA with the factors of

Group x Modality [Imagery/Perception] x Domain. There was strong evidence for an absence

of group differences of FIN activation amplitude (main Group effect, BF = 0.08; see Fig. 3A

the activity profile of the FIN across tasks). FIN activation was stronger for perception

compared to imagery (main Modality effects, BF = 194) and was stronger for words than

other domains (main Domain effect, BF = 5.65e5). No interaction effect was found (all BFs <

0.14).

Does FIN activity actually depend on the anatomically close visual word form area

(VWFA) activity? The FIN consisted of a single patch in all individual participants, unlike the

domain-preferring regions, which were organized in multiple patches (Fig. S3). Its peak

location was mesial, rostral and ventral to the VWFA (Bayesian t-tests on coordinates, two-

sided: X BF = 2.57, Y BF = 3.20, Z BF = 6.59, see Fig. S4 for individual maps), with partial

overlap of VWFA activation in the lateral OTS. We specifically tested for differences in

activation by identifying FIN-unique voxels, after excluding the voxels which overlapped with

the VWFA. There was extreme evidence for different activation profiles between the FIN

unique area and VWFA (BF = 2,629, see Fig. S6 for detailed statistics and activity profiles).

These results confirm that the FIN is a functionally unique region, different from VWFA.

The representational content of the FIN. We tested whether the FIN contains domain-

related information by computing representational dissimilarity matrices (RDMs) between the

multivoxel spatial activation patterns (Kriegeskorte, 2008). We computed the pairwise

correlations between the spatial patterns of BOLD response elicited by the 90 imagery items

and by the 30 perceptual items.

For imagery tasks, the RDMs of both groups featured small blocks around the

diagonal in the FIN (Fig. 3B, darker blue areas; Bayesian correlation values with a

theoretical diagonal matrix rs > 0.23; BFs > 5.89e44; n = 4005), suggesting the presence of

domain-related information in the FIN. In addition, the RDMs showed high similarity between

typical imagers and aphantasic individuals (r = 0.09; BF = 2.32e5). For perception tasks,

there were similar diagonal domain blocks patterns in the RDMs of FIN (rs > 0.27; BFs >

1.19e6; n = 435), which showed highly similar representation between groups (r = 0.24, BF =

1.44e5). However, the correlation between imagery and perceptual RDMs of the FIN (Fig.

S7A), where participants imagined or perceived the same items, was higher for typical

imagers than for aphantasic individuals (Bayesian t-test, BF = 8.15, Fig. S7B). Moreover, the

strength of this correlation was strongly correlated with individual vividness VVIQ scores only

in typical imagers (r = 0.82, BF = 14.04, Fig. S7C), i.e. the greater the similarity in voxel

patterns between imagery and perception of the same stimuli, the more vivid the mental

imagery experienced by participants.

FIN is functionally connected with FP areas in typical imagers, but not in
aphantasic individuals

We examined the functional connectivity of the FIN using psychophysiological interaction

(PPI) to map its task-specific functional network during the imagery tasks and test whether

aphantasia shows alteration of these connectivity within the imagery network.

For **domain-general connectivity**, we first studied typical imagers. We looked for

regions whose correlation with the FIN would be higher during Imagery than during the

Abstract words task, and found bilateral OFC regions (individual FINs as seed regions, Fig.

S8A). We then compared the connectivity of the FIN between the "all Imagery" and the "all

Perception" conditions. During imagery, we observed higher connectivity with the left OFC,

mSTS and the right anterior temporal lobe and during perception, bilateral posterior

occipitotemporal visual areas. The pattern of connectivity was very different in aphantasic

individuals. No brain region displayed increased functional connectivity with the FIN in either

the contrast of all Imagery vs. Abstract words or the contrast of all Imagery vs. all

Perception. Directly comparing patterns of FIN connectivity between the two groups

confirmed that typical imagers displayed stronger connectivity during imagery than for either

Abstract words (Fig. 3C, Table S10) or Perception (Fig. 3D) with the left anterior PFC, OFC,

and MTG/STG compared to aphantasic individuals.

For **domain-specific connectivity**, we compared each imagery or perceptual

domain minus Abstract words using PPI analysis. In typical imagers, we observed increased

FIN connectivity with dorsolateral FP areas, and with the relevant domain-preferring VTC

regions and subcortical areas in both domain-specific imagery and perceptual tasks

(Imagery: Fig. 3E, detailed report of ROIs in Table S3 and Fig. S8 for each domain;

Perception: Fig. 3F and Table S4). In contrast, in aphantasic individuals, when comparing

single imagery or perceptual domains with Abstract words, we did not observe any increase

in connectivity between the FIN and dorsal FP regions. Concerning specific imagery

domains, in aphantasic individuals the FIN displayed higher local connections with FG in

color imagery and right vPCC in map imagery (Fig. 3E and Table S7). No increased

connectivity was found for the imagery of shapes, faces, and maps. For perceptual domains,

the FIN showed only local higher connectivity with areas in the occipitotemporal region for

the perception of shapes, colors, words, and faces (Fig. 3F and Table S7). Word perception

induced also a higher FIN connection with right anterior PFC. No significant connectivity was

observed in the perception of maps. Importantly, this absence of measurable functional

connectivity pattern is consistent across all aphantasic individuals. A direct comparison

between the two groups revealed that left PMd and anterior PFC were regions consistently

more connected to the FIN in typical imagers than in aphantasic individuals during both

imagery and perception.

In task-residual connectivity, we regressed out the task-related activity (see Methods)

to obtain task residual data in imagery runs, and derive a proxy for resting-state functional

connectivity. We observed FIN connectivity with the bilateral supplementary motor area,

precuneus, V1, left middle frontal gyrus, FG and right IPS in typical imagers (Fig. S9C) and

with bilateral V1 and right MFG in aphantasic individuals. No significant group difference was

observed in task-residual connectivity.

 In summary, the FIN showed comparable activation and representational content

408 between typical imagers and aphantasic individuals. In typical imagers, the FIN is

409 functionally connected with FP areas and with the relevant domain-preferring regions during

410 both imagery and perception. Aphantasic individuals showed no measurable functional

411 connectivity between the FIN and other regions in either imagery or perception.

Fig 3. Activation profile, representational similarity and functional connectivity of the FIN between typical imagers and aphantasics

(A) Group-level contrast of all Imagery minus Abstract words identifies the FIN. Orange-colored patches indicate higher activation in imagery tasks than in the Abstract words task. All group-level maps underwent cluster-size thresholding using Monte-Carlo simulation with an alpha level of 0.05 and 5,000 simulations, and an initial threshold of $p<0.005$. The functional data was smoothed with a full width at half maximum of 6 mm. Group-level histogram of signal change in the FIN during imagery and perception for each domain. Error bars represent +/- 1 normalized SE across participants. Typ, typical imagers; Aph, aphantasic individuals.

(B) Group-averaged RDM and between-group correlation of imagery RDM for 90 different items and of perceptual RDM for 30 different items. The emergence of small diagonal domain blocks, indicated by the darker blue areas, shows higher similarity between items belonging to the same domain. The correlation coefficient is a measure of the resemblance between the mean RDMs of Typ and Aph groups.

(C) Group difference in the functional connectivity of the the FIN: all Imagery versus Abstract words. All group-level maps were thresholded at $p<0.005$ for cluster size correction. Orange shows regions with higher connectivity in typical imagers than in aphantasics. The FIN in typical imagers showed stronger connectivity during imagery with the left anterior PFC, OFC, and MTG/STG compared to aphantasic individuals.

(D) Group difference of imagery domain-general FIN connectivity: all Imagery versus all Perception. Orange shows regions with higher connectivity in typical imagers than in aphantasics.

(E) PPI for single Imagery domains minus Abstract words using the individual FIN as seed. In typical imagers, the FIN was functionally connected with FP areas and domain-specific VTC areas. In aphantasia, no FP region was significantly more connected to the FIN, and no region was significantly more connected to the FIN in the shape, word, and face imagery domains than during the Abstract words task. Comparisons between groups showed that the left PMd and anterior PFC regions were more connected to the FIN in typical than in aphantasic individuals across imagery domains. See 3D volume ROIs visualizations and detailed report for each domain in Fig. S8B.

(F) PPI for single Perceptual domains minus Abstract words using the individual FIN as seed. In aphantasia, no FP region was significantly more connected with the FIN and no difference in connectivity was measurable during map perception. See also 3D volume ROIs visualizations and detailed report for each domain in Fig. S9A.

No difference of V1 activity between typical imagers and aphantasic individuals during imagery

If EVA play a crucial role in mental imagery, then aphantasia might be associated with

abnormal EVA activity patterns. To investigate this possibility, we conducted the same

analyses for V1 as we did for the FIN.

For **activation amplitude**, we compared the activity in foveal and peripheral V1 ROIs

between the two groups using Bayesian repeated measures ANOVAs (group x task). As

expected, the foveal V1 was activated during the perceptual tasks, with no group differences

(BF = 0.53, Fig. 4A). In contrast, we observed deactivation in foveal V1 for the imagery tasks

and for the abstract word task in both groups. This deactivation had a lower amplitude in

aphantasic individuals than in typical imagers during the imagery tasks (BF = 7.41), and

during the abstract word task (BF = 3.47). No difference between tasks and groups was

found in the peripheral V1 (BF = 0.27).

For **representational content**, during imagery, there was no evidence of within-

domain similarity across trials in V1 (rs<0.04, BFs<1.09), and no significant correlation of

RDMs between groups (r = 0.05, BF = 3.24). In contrast, during perceptual tasks, RDMs for

V1 showed very strong evidence of within-domain similarity blocks in both groups (rs > 0.39,

BFs > 1.02e14), and very strong correlation between groups (r = 0.61, BF = 1.08e42). There

was moderate evidence to reject the correlation between imagery and perception

representation in V1 (BF = 0.24, Fig S7A), and no support for group differences (BF = 0.41).

Finally, we studied the functional connectivity of foveal and peripheral V1 seeds, and

found no significant group differences with either task-specific or task-residual analyses (all

ps>0.005, Fig. 4C).

A V1 activity profile

Foveal V1

Peripheral V1

B Imagery similarity   Perceptual similarity

Pearson r = 0.05 (BF = 3)   Pearson r = 0.61 (BF = 1.1e42)

Fig 4. Activation profile, representational similarity and functional connectivity of V1 between typical imagers and aphantasics

(A) Group-level histogram of signal change in the foveal V1 and peripheral V1 ROIs during imagery and perception for each domain. Error bars represent +/- 1 normalized SE across participants. Light and dark colors refer to imagery and perception, respectively. Note the deactivation in foveal V1 during imagery and the abstract words task. Typ, typical imagers; Aph, aphantasic individuals.

(B) Group-averaged RDM and between-group correlation of imagery RDM for 90 different items and of perceptual RDM for 30 different items. The darker blue blocks along the diagonal show within-domain similarity across trials. The correlation coefficient is a measure of the resemblance between the mean RDMs of the Typ and Aph groups.

Higher activation of a right IFG-SMG network in aphantasia

Most previous analyses were focused on the VTC and EVA. We then assessed whether differences between groups existed also in non-visual brain sectors, particularly in the frontal and parietal regions with an involvement in mental imagery. We used whole-brain ANOVAs

to compare brain activation. For the all Imagery tasks, aphantasic individuals showed higher

activation than typical imagers in the right-hemisphere IFG, supramarginal gyrus (SMG),

IPS, pMTG and a in a left occipital area (Fig 5A and Table S8). These same right-

hemisphere FP regions were *deactivated* in typical imagers for the same contrast. Notably,

during perception also, the same right-hemispheric IFG-SMG network showed higher

activation in aphantasic individuals than in typical imagers (Fig. 5B). Activation during the

control task with abstract words showed no activation difference between groups.

Concerning domain-specific imagery activation, no areas other than the right hemisphere

IFG-SMG network were differentially activated between groups.

504
505
Fig 5. Higher activation in right-hemisphere regions in aphantasia.
(A) Group difference in all Imagery tasks. Blue shows regions with higher activation in
aphantasic individuals than in typical imagers. Typ: typical imagers. Aph: aphantasic
individuals.
(B) Group difference in all Perception tasks. Regions with higher activation in aphantasic
individuals than in typical imagers are shown in blue.
512

513   Last, we sought to identify brain areas whose activity is modulated by subjective vividness

514   across the imagery trials. While there was no significant group-level effect of vividness, the

515   FIN appeared to be the most consistently modulated area by vividness when considering

516   individual participants (4 out of 6 typical imagers who showed significant modulated regions,

517   Fig. S10). No participant showed modulation of activity in EVA.

Discussion

We used ultra-high-field fMRI to systematically examine domain-general and domain-specific mechanisms of visual mental imagery in typical imagers and in individuals with congenital aphantasia, who claim not to experience any visual mental imagery during wakefulness. Our study involved comprehensive testing of visual mental imagery capabilities across five different domains, namely object shapes, object colors, faces, letters, and spatial relationships. In both typical imagers and aphantasic individuals, imagery tasks activated the relevant domain-preferring VTC patches for each of the five explored imagery domains. Importantly, imagery overlapped with perception only in the anterior VTC domain-preferring patches. In addition, we observed a domain-general cortical patch within the posterior lateral OTS in the left hemisphere, in a location consistent with the Fusiform Imagery Node (FIN) (Spagna et al., 2021). In aphantasic individuals, imagery and perception exhibited similar activation and representational content in high-level visual areas.

Beyond those commonalities, typical and aphantasic participants differed in three important respects, which should be key to understanding aphantasia. First, in the FIN, the correlation between imagery and perceptual representations was higher for typical imagers than for aphantasic individuals. Second, the FIN of aphantasics was functionally disconnected from frontoparietal areas. Third, aphantasic individuals showed enhanced deactivation of foveal V1 activity during imagery. We will now discuss in turn the role of the VTC, the FIN, and the EVA in conscious imagery and in aphantasia.

The role of domain-preferring ventral temporal areas. Dissociations in performance across various imagery domains among neurological patients (Bartolomeo, 2002; Goldenberg, 1993) suggest the existence of domain-specific circuits for visual mental imagery of object shape, object color, written words, faces, and spatial relationships, which may partly overlap with the corresponding domain-preferring circuits in visual recognition (Cohen et al.,

2000; Epstein et al., 1999; Kanwisher et al., 1997; Lafer-Sousa et al., 2016; Malach et al.,

1995). Our findings clarify the topographical organization of VTC regions thanks to high-

resolution precision imaging in individual participants (see Fig. S3). As mentioned above,

previous work had shown FFA and PPA activity during imagery of faces and places,

respectively (Ishai et al., 2000; O'Craven & Kanwisher, 2000). We replicated those findings,

and extended them by demonstrating imagery-related activity in the LOC for shape imagery,

in the VWFA for letter imagery, and in color-biased regions for color imagery. Importantly,

individual analyses allowed us to put to light the overlap of the domain-preferring patches

activated during imagery and during perception. This overlap was restricted to the VTC high-

level visual and associative areas. For instance, color imagery activated the anterior color-

biased patches but not the more posterior ones.

Beyond the occipitotemporal cortex, we also found domain-preferring imagery activity

in dorsal FP networks, in subcortical regions (such as in the amygdala for face imagery,

possibly encoding face-associated emotions), and adjacent OFC patches for faces and colors,

respectively. Face-preferring OFC patches had previously been described by Ref. (Barat et

al., 2018; Tsao et al., 2008) in monkeys. The previously unknown color-preferring OFC

patches may be related to the behavioral saliency of colors, particularly in detecting the

emotional aspects of faces and food choices, such as assessing the ripeness of fruits (Siuda-

Krzywicka et al., 2019).

A domain-general imagery node in the left fusiform gyrus. The FIN was

consistently active during visual mental imagery and during perception, independent of the

imagery domain. Left temporal activity was previously described at locations close to the FIN

during imagery (D'Esposito et al., 1997; Yomogida et al., 2004).

Anatomically, the FIN is contiguous to domain-preferring VTC regions such as VWFA

or the FFA, such that one may ask whether it is topographically distinct from those regions.

The present results localize the FIN around the left-hemispheric posterior OTS in 19

participants of 20. Individual analyses showed that the FIN, which was always restricted to a

single patch, tended to be more mesial, rostral and ventral than the VWFA. Its location was

sandwiched between the VWFA laterally and FFA mesially, with a possible partial overlap with

these regions.

Our findings substantiate the hypothesis that the FIN has a very specific involvement

in both imagery and perception, based on the following functional attributes. First, the FIN

showed domain-generality by its increased BOLD activation when performing imagery tasks

in all five domains: object shape, object color, written words, faces, and spatial relationships.

Second, the FIN, together with the left-hemisphere IFG/IPS (see Fig. 6A), coded semantic

content in its multivoxel patterns of activity. Third, the FIN showed increased functional

connectivity with FP networks and with the relevant domain-preferring regions during domain-

specific tasks, consistent with a role for the FIN as a semantic control hub (Jackson, 2021) for

task-relevant features in both imagery and perception. Moreover, the strong left lateralization

of the FIN, as well as that of domain-general FP areas, is in line with abundant evidence on

hemispheric asymmetries of voluntary generation of visual mental imagery (Farah, 1984; Liu

et al., 2022). Such hemispheric asymmetry is difficult to reconcile with the dominant model of

visual mental imagery, which puts an emphasis on the implication of non-lateralized EVA. It is

instead consistent with the predominant left-lateralization of the semantic system (Binder et

al., 2009; Fernandino et al., 2022), which provides a main input to voluntary visual mental

imagery. Thus, these results support the notion of a semantic contribution of the FIN to the

construction of mental images.

Still, the stronger activation of the FIN during imagery tasks than during abstract words

processing might have an alternative explanation: the FIN could merely be a semantic region

rostral to VWFA, specialized for concrete as opposed to abstract words. This possibility,

however, would be in line with the very definition of concrete words, which are words linked to

sensorimotor-based experience. As a consequence, word concreteness is highly correlated

with word imageability (r=.971, Khanna & Cortese, 2021). Moreover, the FIN appeared to be

critically involved in mental imagery in studies using a variety of contrasts, some of which

implemented non-verbal cues to evoke imagery (Spagna et al., 2021). For example, left

fusiform activations similar to ours have been found in studies using non-verbal stimuli such

as drawings (Mazard et al., 2005) or mathematical formulas (Pyke et al., 2017). In a further

study (Amalric & Dehaene, 2016), mathematicians rated their subjective imageability of

mathematical statements. These ratings positively correlated with activity in a left

inferotemporal region (Talairach coordinates: -54; -52; -1) close to the FIN (Talairach

coordinates: -41, -55, -10).

The separation between domain-general and domain-specific functions in the VTC is

likely to minimize the cost of long-distance wiring (Sporns & Betzel, 2016) thanks to locally

dense connections between domain-preferring regions and the FIN, and sparser connections

with more remote areas. The FIN, equipped with long-range connections with the perisylvian

language network and the anterior temporal lobe (Hajhajate et al., 2022), may thus act as a

central hub for global back-and-forth communication between visual areas and language-

related regions. Thus, the FIN may act as a bridge between semantic and visual information,

enabling the generation of mental images with a visual content. The observation of functional

connectivity between FIN and dlPFC / OFC dovetails nicely with the anatomical connectivity

of the FIN to these regions, and supports its role as a domain-general node at the confluence

of top-down influences from the FP networks and horizontal connections with the VTC domain-

preferring regions.

This hypothesis also offers an explanation for the deficits in visual mental imagery

observed in neurological patients, and yields testable predictions. Patients with lesions

affecting the domain-general FIN, or disconnecting it from semantic networks, are likely to

experience general imagery impairments (Bartolomeo, 2021; Moro et al., 2008). Lesions or

disconnections that specifically target domain-preferring regions may result in more domain-

specific patterns of mental imagery deficits (Bartolomeo et al., 2002). More posterior lesions

disconnecting VTC from visual input are instead likely to produce perceptual deficits with

preserved visual mental imagery (Bartolomeo, 2002, 2021).

The relationship between visual perception and visual mental imagery. Our

findings elucidate three important aspects of typical visual mental imagery, indicating that

imagery does share some neural substrates with visual perception (Dijkstra et al., 2019;

Mechelli et al., 2004), but with prominent differences. First, domain-preferring VTC regions

exhibited some overlap between imagery and perception in more rostral patches, whereas

more caudal patches only responded to perception. Second, we identified shared cortical

patterns of representation for semantic domain content between imagery and perception in

the high-level visual cortex. This finding aligns with previously reported similarities in

domain-preferring visual areas between imagery and perception (Cichy et al., 2012; Reddy

et al., 2010; Stokes et al., 2009). Notably, this representational overlap correlates with the

level of vividness in typical imagers, specifying an objective measurement of subjective

vividness across imagery domains (Dijkstra et al., 2017). Third, the FIN displayed different

functional connectivity patterns between imagery and perception. In imagery, it displayed

stronger connections with semantic networks, whereas in perception, it showed greater

connectivity with occipitotemporal areas. Taken together, this evidence emphasizes the

importance of high-level visual cortex for imagery and the common role of the FIN in

processing semantic and visual content for both imagery and perception.

The role of early visual areas in visual mental imagery. In both typical imagers

and aphantasic participants, we observed peripheral V1 activation during imagery and

perceptual tasks, as well as during the control task with abstract words. This activity might

result from orienting of spatial attention in response to auditorily presented stimuli, which is

often found in peripheral retinotopic locations of V1 (Brang et al., 2015; Cate et al., 2009),

even in the absence of external stimuli (Kastner et al., 1999). Importantly, however, *foveal*

V1 was active in perception but showed negative activity in imagery and in the abstract

words task. This can result from attention being endogenously directed toward internal

thoughts, which may inhibit foveal V1 to prevent potential interferences from external input.

These findings challenge standard models stressing the role of EVA in visual mental imagery

(Kosslyn et al., 2001; Pearson, 2019). However, the pattern we observed is quite consistent

with extensive neuroimaging evidence in neurotypical individuals, which shows that visual

mental imagery triggers activity in VTC and FP networks - but not in the EVA (Mechelli et al.,

2004; Spagna et al., 2021). Moreover, detailed studies of neurological patients provided

causal evidence through observations of disrupted imagery following left temporal damage

rather than following lesions restricted to the occipital cortex (Bartolomeo, 2002; Bartolomeo

et al., 2020; Liu et al., 2022). In different experimental contexts, mental imagery of colors

(Bergmann et al., 2024), or the expectation to see gratings (Aitken et al., 2020), have been

shown to modulate activity in the deep layers of V1. However, the comparable V1 activity

and connectivity of aphantasia in our study suggests that these V1 patterns have no causal

contribution to conscious imagery experience, consistent with early suggestions (Crick &

Koch, 1995).

Functional disconnection in aphantasia. Surprisingly, aphantasic individuals

exhibited the activation of similar brain networks during mental imagery as observed in

typical imagers. This was confirmed by univariate and RSA analyses of BOLD responses, in

both domain-general and domain-preferring VTC areas, and in the FP networks. Importantly,

aphantasic individuals could generate imagery-related representational patterns similar to

those of typical imagers, indicating the presence of relevant visual information in the high-

level visual cortex during attempted mental imagery. Consistent with this observation,

aphantasic individuals were able to perform as accurately as typical imagers on tests of

mental imagery (Liu & Bartolomeo, 2023). However, while typical imagers showed a

functional connection between the FIN and FP networks activity, aphantasic individuals did

not display such correlations. Specifically, in aphantasia the FIN was functionally

disconnected from the FP networks during both visual mental imagery and perception, which

might be in line with their slower behavioral responses. In partial agreement with our results,

Milton et al. (2021) found reduced resting-state functional connectivity between PFC and the

visual–occipital network in aphantasia. Reduced long-range connectivity is a hallmark of

various neurodevelopmental disorders, perhaps including aphantasia (Sokolowski & Levine,

2023). Within the framework of the Global Neuronal Workspace hypothesis of conscious

perception (Dehaene et al., 2006; Mashour et al., 2020), such a functional disconnection

between FIN and FP networks could be interpreted as depriving visual mental imagery of its

conscious experiential content. In other words, VTC activity by itself might be sufficient to

access visual information, but not to experience it as a conscious perception-like state. In

line with our findings, the Global Neuronal Workspace hypothesis underscores the

importance of long-range loops between different cortical areas in sustaining and

broadcasting neural information across the brain, with the dorsolateral PFC cortex playing an

essential role (Dehaene et al., 2006; Mashour et al., 2020). Hence, aphantasia,

characterized as a relatively "pure" deficit in conscious imagery experience, with a

paradoxically preserved ability to perform imagery tasks, offers a compelling testing ground

for theories of the brain mechanisms of conscious experience.

Aphantasic individuals exhibited impaired ability for object imagery but not for spatial

imagery (Blazhenkova & Pechenkova, 2019), as shown by OSIQ. However, our fMRI result

showed a consistent functional disconnection for spatial imagery as for the remaining

imagery domains in aphantasic individuals. This discrepancy may be due to differences in

the tasks. In the OSIQ, the Object part directly evaluates the subjective vividness of mental

images, for example, "My images are very vivid and photographic." On the other hand, the

Spatial part primarily pertains to spatial knowledge without explicitly requiring imagery

experience, for instance, "In high school, I had less difficulty with geometry than with art.".

These spatial questions differ substantially from our Map of France task, which requires

subjects to visualize a map and assess the spatial location of imagined cities. Aphantasic

individuals performed the task accurately, but when rating the trial-by-trial vividness they

reported almost no mental imagery at all.

Increased activity in aphantasia. In both imagery and perception, the present

group of aphantasic individuals exhibited greater activity in right-hemisphere IFG and SMG,

which are important components of the right-lateralized network for reorienting attention

(Bartolomeo & Seidel Malkinson, 2019; Corbetta et al., 2008) and its interaction with

conscious perception (Liu et al., 2023). Such abnormal activity may play a role in disrupting

the subjective experience of generating or maintaining mental imagery, for example by

interrupting ongoing activity in more dorsal FP networks (Corbetta et al., 2008). A possible

mechanism could be defective filtering of distracting events (Shulman et al., 2007), leading

to interference with internally generated models (Bartolomeo & Seidel Malkinson, 2022).

Several considerations support the notion that the imagery tasks of the eBIP were

capable of evoking visual mental images. First, using the same battery, Liu & Bartolomeo

(2023) found an inverse correlation between trial-by-trial subjective vividness and response

times on the eBIP. Higher levels of vividness were associated with faster response times.

Second, and more importantly, the eBIP imagery tasks induced activations of perceptual

domain-preferring VTC patches, confirming and extending early findings by O'Craven &

Kanwisher (2000) on imagery of faces and places. We obtained more systematic evidence

here across the five semantic domains we investigated. Third, tasks similar or identical to the

eBIP have often been used in clinical settings to assess domain-selective impairments.

Neurological patients with imagery deficits in specific domains showed impaired

performance on these tasks, which cannot be attributed to impaired semantic knowledge.

For example, patient VSB (2002) had a deficit in visual perception and visual imagery of

letters (as assessed by questions on the visual shape of letters, similar to those used in the

eBIP) as a consequence of a left temporal stroke. However, he could still answer the same

questions when he was allowed to mimic their writing, demonstrating preserved semantic

knowledge of letters.

Limitations of the present study include 1) The exclusion of bilateral anterior temporal

lobes due to limited brain coverage at 7T (Fig. S2A). 2) The impossibility to analyze our trial-

by-trial vividness scores, because of insufficient variability in the vividness ratings of typical

imagers (consistently high) and of aphantasic individuals (consistently low; Fig. S1B). (3)

The possibility that aphantasia might be a heterogeneous condition, in the absence of

diagnostic criteria that could identify potential subtypes with differing neural substrates.

Despite these limitations, our findings shed light on the left-predominant circuits of

individual-level visual mental imagery, encompassing the FIN, FP networks, and,

importantly, domain-preferring VTC regions. This evidence suggests the presence of distinct

domain-general and domain-preferring cortical components of visual mental

imagery(Spagna et al., 2024). Our results also demonstrate that visual mental imagery and

visual perception share similar neural mechanisms in the high-level visual cortex. Finally, we

identified a neural signature of aphantasia, which was associated with a functional

disconnection within the imagery network between the FIN and FP networks, despite

essentially normal behavioral performance, BOLD activity levels, and representational

content. Thus, the present results support the general hypothesis that conscious visual

experience - whether perceived or imagined - depends on the integrated activity of FP

networks and high-level visual cortex (Dehaene et al., 2006; Liu et al., 2023; Mashour et al.,

2020; van Vugt et al., 2018).

Acknowledgements

The research was supported by INSERM, CEA and specific funding from Dassault Systèmes. We are grateful to NeuroSpin support staff, and particularly to Bernadette Martins, the nurses and MR technicians at NeuroSpin who provided crucial help at various stages of data acquisition and analysis. The work of P.B. is supported by the Agence Nationale de la Recherche through ANR-16-CE37-0005 and ANR-10-IAIHU-06, and by the Fondation pour la Recherche sur les AVC through FR-AVC-017. J.L. expresses his gratitude for the 2023 European Workshop on Cognitive Neuropsychology Prize that was awarded to him in recognition of the work presented in this study.

Contributions

Conceptualization: J.L., P.B., L.C., S.D.; Data curation: J.L., D.H., M.Z.; Formal analysis: J.L., M.Z.; Investigation: J.L., D.H., M.Z.; Methodology: J.L., M.Z.; Funding acquisition: P.B., J.L.; Project administration: P.B.; Resources: S.D.; Software: J.L., M.Z.; Supervision: P.B.; Visualization: J.L.; Writing - original draft: J.L.; Writing - review & editing: All authors

Competing interest

The authors declare that they have no competing interests.

Materials & Correspondence

Requests for materials and correspondence should be addressed to J.L.

Method

1.   Participants

Ten typical imagers (mean age ± SD, 29.28 ± 8.47, 6 female) were recruited from the CNRS RISC volunteer database (https://www.risc.cnrs.fr/). All typical imagers had average or high vivid mental imagery with Vividness of Visual Imagery Questionnaire (VVIQ) score greater than 55 (mean ± SD, 71.70 ± 7.07, out of a total score of 80). Ten individuals with congenital aphantasia (mean age ± SD, 28.69 ± 8.27, 6 female) were recruited from French-language groups on aphantasia in various online social media. All aphantasic participants reported a *complete* life-long inability to generate visual mental imagery, and this was confirmed by an individual interview with author D.H. (a clinical psychologist) during the recruitment. All aphantasic individuals reported scored 16 out of 80 ("no image at all" for all questions) to the VVIQ. Interestingly, many aphantasics (8 out of 10) confirmed that they tend to perceive pictures with a *divide-and-label* strategy, akin to semantic conversion from visual features into content lists. While asked to visualize, they briefly recall the list of semantic labels to respond to the relevant question. Notably, they reported difficulties in remembering complex or unfamiliar pictures, which might reflect lower efficiency of the semantic encoding strategy in this case. All participants were right-handed with normal or corrected-to-normal vision and had no history of neurological/psychiatric disorders. Aphantasic participants did not differ from typical imagers in age (t = 0.35, p = 0.73, d = 0.16) or in education level (t =1.33, p = 0.20, d = 0.62). Participants provided written consent before the study and received monetary compensation after the study. The study was approved by CEA and, according to French bioethical law, by a randomly selected regional ethical committee for biomedical research (CPP 100055 to NeuroSpin center) and the study was carried out in accordance with the declaration of Helsinki.

2.   Introspective reports

Before the fMRI session, participants completed the French versions of the VVIQ (Santarpia et al., 2008) and Object-Spatial Imagery Questionnaire (OSIQ) (Blajenkova et al., 2006) questionnaires to assess the subjective vividness of their visual mental imagery. The OSIQ consists of two scales assessing preferences for representing and processing of object imagery about pictorial and high-resolution images of individual objects, and of spatial imagery about semantic images and spatial relations amongst objects (see Supplementary materials for French translations). The results confirmed the drastic reduction of imagery ability in aphantasic individuals compared to typical imagers (VVIQ, BF = infinity; OSIQ object, BF = $2.06{\times}10^9$), but no reduction in spatial ability (OSIQ spatial: BF = 0.46).

3.   Stimuli and fMRI experimental design

In the scanner, participants performed a longer version of the enhanced Battérie Imagerie-Perception - eBIP (Liu & Bartolomeo, 2023). The current version of the battery assesses (1) imagery of object shapes (Fig. 1A), object colors (Fig. 1B), faces (Fig. 1C), letters (Fig. 1D) and spatial relationships on an imaginary map of France (Fig. 1E); (2) a non-imagery control task using abstract words (Fig. 1F), and (3) an audio-visual perception task using the same items as in the imagery tasks (Fig. 1G). In the imagery tasks, participants heard a word indicating a particular imagery domain (e.g., "shape"), followed by 2 words, designating the items the participant is required to imagine (e.g. "beaver", "fox"). They were instructed to generate and maintain mental images as vivid as possible for each item. Eight seconds after the second items, they heard an attribute word (e.g. "long"). They then pressed one of two buttons indicating which of the items is most closely associated with the attribute (e.g. which of the animals they associate with the attribute "long", see Fig. 1A). Finally, they reported the overall vividness of their mental imagery in that trial on a 4-level Likert scale by pressing one of 4 buttons of an MR-compatible button box (Current Designs, Philadelphia, USA), where button 1 indicated "no image at all" and button 4 indicated a "vivid and realistic image". In the shape imagery task, participants had to decide which item was longer or rounder. In the color imagery task, participants had to decide which fruit or vegetable had darker or lighter color. In the letter imagery task, participants had to imagine the shape of French words in lowercase and had to decide which word had ascenders (e.g., t, l, d) or descenders (e.g. j, p, y). In the famous-faces imagery task, participants had to decide which celebrity had a more round or oval face. In the map-of-France imagery task, participants had to decide which city was located to the left or the right to Paris. In the non-imagery abstract-word task, participants had to decide which of two abstract words (e.g., "routine", convention") was semantically closer to a third word (e.g. "society"). While visual imagery cannot be completely excluded in this setting (or in any other settings), we selected abstract words for this task in order to minimize its engagement. In the perception task, the same stimuli used for the imagery tasks were presented in an audio-visual format. In the abstract-word and the perception tasks participants rated their confidence on a 4-level Likert scale, instead of rating vividness. The auditory stimulus were voice recordings of the corresponding French names per item. All voice recordings were generated in an online TTS engine (https://texttospeechrobot.com/; fr-FR_ReneeVoice) and digitized at a 44.1 kHz sampling rate. The images shown in the perception task were color photographs of the corresponding items on a gray background. The words in letters perception task were rendered in

"Monotype Corsiva" font to display words because in pilot testing some participants reported that this font was more natural and closer to their everyday visual experience. The resulting occipito-temporal activations in fMRI (-49, -59, -7; left fusiform; 1,989 mm3) were in line with the extensive neuroimaging literature on reading.

4. Data acquisition

The brain images are acquired using an ultra-high field 7-Tesla Magnetom scanner (Siemens, Erlangen, Germany) with a 1Tx/32Rx head coil (Nova medical, Wilmington, USA) at the NeuroSpin center of the French Alternative Energies and Atomic Energy Commission (CEA). No dielectric pads were used due to the limited space within the head coil with the headphone (OptoActive II, Optoacoustics, Israel). To minimize light reflections inside the head coil, a piece of black paper was inserted to cover the inner surface of the transmitter coil element. Stimuli were presented via a BOLDscreen 32 LCD screen (Cambridge Research Systems, UK, 69.84 x 39.29 cm, resolution=1920 x 1080 pixels, refresh rate=120 Hz, viewing distance ~200 cm), at the head-end of the scanner bore. Participants viewed the screen through a mirror attached to the head coil.

The fMRI scan session lasted 150 minutes in total, split into 5 runs. The first 3 runs consisted of 90 imagery-only trials (30 trials per run, each including 6 trials per domain, 12 min 14 s per run, 367 volumes with 2 volumes of resting at the beginning of each run). The absence of visual perceptual trials prevented potential spillovers of perceptual responses to imagery trials. The last 2 runs consisted of 36 trials each (6 abstract word trials, 15 imagery trials and 15 audio-visual perception trials per run, each including 3 trials per domain, 14 min 38 s per run, 439 volumes), in that order. Notably, items in the imagery trials and perceptual trials of the last two runs are identical. The inter-trial intervals were jittered between 3s and 7s. Importantly, the order of trials within each task type was fully randomized across domains. For each imagery domain, the order of items was counterbalanced, i.e., each item is presented once as the first item and once as the second item. Participants were instructed to keep their eyes open during scans and to pay attention to the fixation on a gray screen displayed at the beginning of each trial and during the intertrial period. During the imagery and abstract word tasks, they saw an empty gray screen without fixation cross. During the perception tasks, they saw visual stimuli while simultaneously hearing the corresponding auditory stimuli. The perception tasks also enabled the localization of domain-preferring regions.

Functional data were acquired with a 2D gradient-echo EPI sequence (TR = 2000 ms, TE =
28 ms, voxel size = 1.2 mm isotropic, multiband acceleration factor=2; encoding direction:
anterior to posterior, iPAT=3, flip angle = 75, partial Fourier=6/8, bandwidth=1488 Hz/Px,
echo spacing=0.78 ms, number of slices=70, no gap, reference scan mode: GRE, MB
LeakBlock kernel: off, fat suppression enabled). The slab is tilted upwards to cover most
parts of the brain except the anterior temporal lobe, and to avoid the eyeballs in the slab. To
correct for EPI distortion, a 5-volume functional run with the opposite phase encoding
direction (posterior to anterior) was acquired immediately before each task run. Participants
were instructed not to move between these pairs of two runs. Manual interactive shimming of
the B0 field was performed for all participants. The system voltage was set to 250 V for all
sessions, and the fat suppression was decreased per run to ensure the specific absorption
rate for all functional runs did not surpass 62%. To minimize artifacts and increase signal-to-
noise ratio around the ventral temporal cortex, the functional data acquisition slab was tilted
in a way that excluded the eyes and the ear canal signal dropout region, so that the ventral
temporal cortex especially the anterior occipital-temporal sulcus above the ear canal was
covered. However, part of the anterior temporal lobe was not able to be included (Fig.S2A).

High-resolution MP2RAGE anatomical images were obtained between the third and the
fourth functional runs (resolution=0.65 mm isotropic, TR=5000 ms, TE=2.51 ms,
TI1/TI2=900/2750 ms, flip angles=5/3, iPAT=2, bandwidth=250 Hz/Px, echo spacing=7 ms).

5.   Behavioral data analysis

Our behavioral data analysis was similar to that of a study using the same tasks with a more
extensive cohort of 117 participants (Liu & Bartolomeo, 2023). We conducted two Bayesian
repeated measures ANOVAs within each modality (imagery, perception), with the factors of
Group (Aphantasia, Typical imagers) and Domain (Shape, Color, Word, Face, Space). The
dependent variables for the imagery tasks were accuracy (arcsine-transformed proportions
of correct responses); response times (RTs) and trial-by-trial vividness scores (translated to
a 0-1 scale and arcsine-transformed), and for the perceptual tasks were accuracy, RTs and
trial-by-trial confidence scores (translated to a 0-1 scale and arcsine-transformed). For each
participant, we excluded trials with response times (RT) faster than 150 ms or exceeding
three SDs from the participant's mean. Statistical tests were performed using JASP 0.16.2
(https://jasp-stats.org/), and used the JASP default priors. A commonly accepted convention
is that Bayes factors (BF10 or BFs) between 3 and 10 indicate moderate evidence in favor of
the model in the numerator (H1); BFs between 10 and 30 indicate strong evidence; BFs
larger than 30 indicate very strong evidence. The inverse of these cut-offs values provides

moderate (0.33-0.1), strong (0.1-0.03), or very strong evidence (<0.03) for the model in the denominator (H0), i.e. the null hypothesis.

6.   fMRI data Preprocessing

We processed the fMRI data with BrainVoyager (Version 22.0.2.4572, Brain Innovation, Maastricht, The Netherlands, https://www.brainvoyager.com/), MATLAB (version R2018b), and NeuroElf 1.1 toolbox (http://neuroelf.net/) implemented in MATLAB. The functional data underwent EPI distortion correction using the posterior-to-anterior directional functional volumes (COPE plugin in BrainVoyager), where the in-plane voxel displacement map between the actual task run and the first volume of the preceding distortion correction run (in reversed phase encoding direction) was computed, and applied to the task run. The distortion-corrected data was then corrected for slice scan time (sinc interpolation, slice scanning order read from the slice time table in the DICOM headers), 3D rigid motion correction (trilinear for estimation, sinc for applying the correction, aligned to the first volume within each run), high-pass temporal filtering (GLM with Fourier basis set, number of cycles=3). No spatial smoothing was applied to the data at this stage.

The MP2RAGE anatomical data consisted of four image types: inversion 1, inversion 2, quantitative T1, uniform. To have a similar appearance to the conventional MPRAGE anatomical data, the uniform image was divided by the T1 image (an optional step), and the background noise was masked out by the inversion 2 image. The resulting anatomical image was resampled to 0.6 mm isotropic (framing cube dimension: 384×384×384), and transformed into Talairach space (TAL). All the coordinates reported in our manuscript are Talairach coordinates. For data visualization, the white matter-gray matter boundary was segmented in TAL space, and reconstructed as surface meshes.

For fMRI across-run co-registration, the fourth functional run was co-registered to the anatomical data, then all the other functional runs were manually co-registered to the fourth functional run. The across-run co-registration quality was inspected visually with animations looping through the first volumes across runs in TAL space. In traditional fMRI-to-anatomy coregistration, the contrast and resolution of the fMRI (T2*) is very different from the anatomical (T1) images (see Fig. S2A), and often includes imperfections such as EPI distortion (potentially not fully corrected by top-up distortion correction). In that scheme, the fMRI-to-anatomy coregistration algorithm is not perfect (especially with partial brain coverages, or the acquisition slab moved too much e.g. across different sessions), and

would induce machine errors everytime the coregistration is performed. In comparison, our

fMRI-to-fMRI coregistration approach minimizes the difference of image contrasts between

runs (always T2* and always the same resolution), and allows both machine and human

coregistration and quality assurance with high precision. For the choice of the 4th fMRI run,

participants initially performed the first three runs, then the T1 anatomical scan, and then the

4th and 5th runs. The first TR functional image of the 4th run was closest to the anatomical

scan, where participant's inter-run head movements (e.g. rest and stretching between runs,

plus the within-run head movements accumulated across the whole scanning session) would

be the smallest. After the quality checks, all functional images were then transformed into

TAL space.

After alignment across runs, the functional data of the main experiment were then spatially

smoothed with 6 mm FWHM for group-level univariate analysis due to the individual

anatomical/functional variability. No spatial smoothing was applied in the case of individual

analysis. All results in this study were computed in the volume space.

7.   Univariate analysis

Two different general linear models (GLMs) were defined with the following main predictors:

1) five domains in imagery-only trials (5 predictors in total) in the first 3 runs; 2) abstract word

and five domains in imagery and in perception trails (11 predictors in total) in the last 2 runs.

The period of predictors started from the first domain words to the attribute word before

rating periods. For all the GLM models above, the time courses were %-transformed, the

main predictors were convolved with a two-gamma hemodynamic response function, and the

6 parameters of participant's head motion were z-scored and entered as confounding

factors. AR(2) correction was used for correcting serial correlations. The two GLM models

were applied to both no-smoothed fMRI datasets for individual analysis and 6 mm FWHM

smoothed fMRI datasets for group-level analysis.

The contrast of all Imagery *versus* Abstract words, all Imagery *versus* all Perception*,* all

Perception *versus* Abstract words and single perceptual domain *versus* other four domains

was conducted with trials in run 4-5. The contrast of all Imagery *versus* baseline and single

imagery domain *versus* other four domains was conducted with trials in run 1-3.

The group random-effect GLM analysis was performed for each predictor set with smoothed

data. Cluster size thresholds for all group-level contrast maps in this study were estimated

using Monte-Carlo simulation (alpha level=0.05, numbers of iterations=5000, initial p<0.005),

with the BrainVoyager plugin Cluster-Level Statistical Threshold Estimator

(https://support.brainvoyager.com/brainvoyager/functional-analysis-statistics/46-tresholding-

multiple-comparisons-problem/226-plugin-help-cluster-thresholding), masked with the

common functional data coverage across 10 participants in each group. We used a p-value

of 0.005 instead of 0.001 as a necessary compromise between individual variability i.e. false

negative results, and the risk of false positive results. Our 7T fMRI data at 1.2 mm isotropic

resolution is highly robust at the individual-subject level because of the high image contrast

between white/gray matters and temporal signal-to-noise ratio (tSNR), but it has also

exacerbated inter-individual anatomical and functional differences. Besides, the Monte-Carlo

clustering method does not overly inflate the activated cluster sizes under different initial p

values in our data. For example, we made 3 Monte Carlo simulations under initial p=0.01,

0.005 and 0.001 for the contrast of all Imagery versus Abstract words with 5,000 iterations,

and observed consistent surviving clusters across different thresholds.

8.   Delineating individual functional foveal V1 and pheripheral V1

Due to the substantial functional variability across participants in EVA (Benson et al., 2022),,

we mapped the location of the foveal V1 and the pheripheral V1 in each individual based

on brain anatomy and functional activation during perception. For peripheral V1, the

individual brain was first co-registered to a functional visual atlas visfAtlas (Rosenke et al.,

2021) with linear transformation to each individual's brain anatomy through meticulous

manual alignment and thorough visual inspection, particularly in the vicinity of the calcarine

sulcus. The foveal V1 was manually delimited in the individual brain in the lateral occipital

lobe around the retro calcarine sulcus. The activity profiles of both V1 subregions confirmed

its activation during perception.

9.   Delineating EVA, ventral and dorsal visual pathway ROIs

To account for the variation in the location of the activation in different domains, we visually

delineated the EVA, ventral and dorsal visual pathway ROIs. The EVA consists of cuneus,

posterior part of lingual gyrus/pericalcarine. The ventral visual pathway ROI consists of

anterior part of lingual gyrus, fusiform gyrus and inferior temporal gyrus. The dorsal visual

pathway ROI consists of superior occipital cortex, inferior parietal lobule and posterior part of

superior parietal lobule. We then counted the voxels in the volumetric intersection of domain-
specific activation in the EVA, ventral, and dorsal ROIs, respectively, in each domain.

1005

10.   Representational similarity analysis (RSA)

To assess the group difference in representational information during imagery, we performed
RSA analysis within all imagery domain-general regions activated in the contrast of all
Imagery *versus* Abstract words, and additionally within V1. We obtained voxel-wise % signal
change from TR 4-5 (normalized by the TRs -2 to 0 as the baseline), corresponding to the
maximum activation of the first item and with minimal influence from the second item in each
trial. For imagery similarity analysis, we obtained brain activity from 90 imagery trials (18 per
domain) in run 1-3. For perceptual similarity analysis, we extracted brain activity from 30
perceptual trials (6 per domain) in run 4-5. In imagery-perceptual similarity comparison, we
used brain activity from imagery trials in run 4-5, which had the same items as in perceptual
trials. We calculated the correlation between the multivoxel patterns of each pair of stimuli
using the Pearson correlation coefficient to generate a representational dissimilarity matrix
(RDM). Before comparing between group, we first assessed within-group variability by
calculating the correlation between each participant's RDM and the mean RDM derived from
the remaining individuals within the same group. For either group, there was no evidence of
any differences in within-group consistency of these regions (all BFs < 2.51).

11.   Psychophysiological interactions (PPIs) analysis

To investigate task-specific functional connectivity, we conducted PPI analysis with trials in
run 4-5. We built each PPI design matrix by 1) generating a task contrast regressor of two
conditions, balanced by a regressor of the sum of two conditions, both convolved with the
hemodynamic response function (HRF); 2) extracting a demeaned time course from the
seed ROI; 3) generating an interaction regressor as an element-by-element product of the
HRF-convolved task contrast and seed ROI regressors; 4) adding the 6 parameters of
participant's z-scored head motion as confounding factors. Specifically, to investigate
domain-general connectivity, we grouped all five imagery domains as a single main condition
and all five perception domains as another main condition, and included a separate main
condition for Abstract words. In each PPI contrast, a task contrast regressor was
established, such as the contrast between all Imagery *versus* Abstract words, which was
defined by subtracting the main condition for Abstract words from the main condition for

Imagery. On domain-specific connectivity, we built the task contrast regressor by subtracting

the Abstract words condition from single domain condition, while keeping the other

imagery/perceptual domain conditions unchanged. In BrainVoyager, we conducted GLMs

using the PPI design matrix for each contrast separately on fMRI datasets that were

smoothed with 6 mm FWHM. The resulting group-level maps were thresholded at p<0.005

for cluster size correction.

12. Task-residual functional connectivity analysis

1041

In order to identify putative direct upstream or downstream areas within the same functional

structure of the FIN in the task, we further performed functional connectivity using task

residual data that exploits the variance remaining after removing the mean task-related

signal from a time series. We smoothed the task data of the run 1-3 at 6 mm FWHM,

regressed out the task-related activity by deconvolution analysis (12 stick predictors per

stimulus, covering the evolvement of the BOLD shape per trial) and head motion

parameters. Using individual FIN as the seed region, we averaged the extracted residual

time course across voxels, and correlated it with the residual time courses of all voxels in

each run, resulting in one correlation R map per run. The connectivity pattern across the

runs was stable for all participants. The R maps were Fisher's Z-transformed, averaged

across runs per participant for group-level comparisons. The resulting group-level maps

were thresholded at p<0.005 for cluster size correction using Monte-Carlo simulation (alpha

level=0.05, numbers of iterations=5000, initial p<0.005).

13. Individual trial-by-trial parametric modulation of vividness

1055

1056 We fitted individual GLM with trial-by-trial vividness ratings as an main predictor, applied to

1057 data from the run 1 to 3 with 6 mm FWHM smoothed fMRI datasets. There were 90 imagery

1058 trials in total to estimate the individual parametric modulation of vividness.

1059

References

Aitken, F., Menelaou, G., Warrington, O., Koolschijn, R. S., Corbin, N., Callaghan, M. F., & Kok, P. (2020). Prior expectations evoke stimulus-specific activity in the deep layers of the primary visual cortex. *PLOS Biology*, *18*(12), e3001023. https://doi.org/10/grzhbd

Amalric, M., & Dehaene, S. (2016). Origins of the brain networks for advanced mathematics in expert mathematicians. *Proc Natl Acad Sci U S A*, *113*(18), 4909–4917. https://doi.org/10.1073/pnas.1603205113

Bannert, M. M., & Bartels, A. (2018). Human V4 Activity Patterns Predict Behavioral Performance in Imagery of Object Color. *Journal of Neuroscience*, *38*(15), 3657–3668. https://doi.org/10.1523/JNEUROSCI.2307-17.2018

Barat, E., Wirth, S., & Duhamel, J.-R. (2018). Face cells in orbitofrontal cortex represent social categories. *Proceedings of the National Academy of Sciences*, *115*(47), E11158–E11167. https://doi.org/10.1073/pnas.1806165115

Bartolomeo, P. (2002). The Relationship Between Visual Perception and Visual Mental Imagery: A Reappraisal of the Neuropsychological Evidence. *Cortex*, *38*(3), 357–378. https://doi.org/10.1016/s0010-9452(08)70665-8

Bartolomeo, P. (2007). Visual neglect. *Curr Opin Neurol*, *20*(4), 381–386. https://doi.org/10.1097/WCO.0b013e32816aa3a3

Bartolomeo, P. (2021). Visual agnosia and imagery after Lissauer. *Brain: A Journal of Neurology*, *144*(9), 2557–2559. https://doi.org/10/grrsd6

Bartolomeo, P., Bachoud-Levi, A. C., Chokron, S., & Degos, J.-D. (2002). Visually- and motor-based knowledge of letters: Evidence from a pure alexic patient. *Neuropsychologia*, *40*(8), 1363–1371. https://doi.org/10/fm5k8g

Bartolomeo, P., Bachoud-Lévi, A.-C., De Gelder, B., Denes, G., Barba, G. D., Brugières, P., & Degos, J.-D. (1998). Multiple-domain dissociation between impaired visual perception and preserved mental imagery in a patient with bilateral extrastriate lesions. *Neuropsychologia*, *36*(3), 239–249. https://doi.org/10.1016/S0028-3932(97)00103-6

Bartolomeo, P., Hajhajate, D., Liu, J., & Spagna, A. (2020). Assessing the causal role of early visual areas in visual mental imagery. *Nature Reviews Neuroscience*, *21*(9), 517–517. https://doi.org/10.1038/s41583-020-0348-5

Bartolomeo, P., & Seidel Malkinson, T. (2019). Hemispheric lateralization of attention processes in the human brain. *Current Opinion in Psychology*, *29*, 90–96. https://doi.org/10/ggbr5v

Bartolomeo, P., & Seidel Malkinson, T. (2022). Building models, testing models: Asymmetric roles of SLF III networks?: Comment on "Left and right temporal-parietal junctions (TPJs) as 'match/mismatch' hedonic machines: A unifying account of TPJ function" by Doricchi et al. *Physics of Life Reviews*, *44*, 70–72. https://doi.org/10/grrsd8

Benson, N. C., Yoon, J. M. D., Forenzo, D., Engel, S. A., Kay, K. N., & Winawer, J. (2022). Variability of the Surface Area of the V1, V2, and V3 Maps in a Large Sample of Human Observers. *Journal of Neuroscience*, *42*(46), 8629–8646. https://doi.org/10.1523/JNEUROSCI.0690-21.2022

Bergmann, J., Petro, L. S., Abbatecola, C., Li, M. S., Morgan, A. T., & Muckli, L. (2024). Cortical depth profiles in primary visual cortex for illusory and imaginary experiences. *Nature Communications*, *15*(1), 1002. https://doi.org/10.1038/s41467-024-45065-w

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cerebral Cortex*, *19*(12), 2767–2796. https://doi.org/10/c2vp52

Blajenkova, O., Kozhevnikov, M., & Motes, M. A. (2006). Object-spatial imagery: A new self-report imagery questionnaire. *Applied Cognitive Psychology*, *20*(2), 239–263. https://doi.org/10.1002/acp.1182

Blazhenkova, O., & Pechenkova, E. (2019). The two eyes of the blind mind: Object vs. spatial aphantasia? *Russian Journal of Cognitive Science*, *6*(4), Article 4.

Brang, D., Towle, V. L., Suzuki, S., Hillyard, S. A., Di Tusa, S., Dai, Z., Tao, J., Wu, S., & Grabowecky, M. (2015). Peripheral sounds rapidly activate visual cortex: Evidence from electrocorticography. *Journal of Neurophysiology*, *114*(5), 3023–3028. https://doi.org/10.1152/jn.00728.2015

Cate, A. D., Herron, T. J., Yund, E. W., Stecker, G. C., Rinne, T., Kang, X., Petkov, C. I., Disbrow, E. A., & Woods, D. L. (2009). Auditory Attention Activates Peripheral Visual Cortex. *PLOS ONE*, *4*(2), e4645. https://doi.org/10.1371/journal.pone.0004645

Chica, A. B., Paz-Alonso, P. M., Valero-Cabré, A., & Bartolomeo, P. (2012). Neural Bases of the Interactions between Spatial Attention and Conscious Perception. *Cerebral Cortex*, *23*(6), 1269–1279. https://doi.org/10.1093/cercor/bhs087

Cichy, R. M., Heinzle, J., & Haynes, J. D. (2012). Imagery and perception share cortical representations of content and location. *Cereb Cortex*, *22*(2), 372–380. https://doi.org/10.1093/cercor/bhr106

Cohen, L., Dehaene, S., Naccache, L., Lehéricy, S., Dehaene-Lambertz, G., Hénaff, M.-A., & Michel, F. (2000). The visual word form area: Spatial and temporal characterization of an initial stage of reading in normal subjects and posterior split-brain patients. *Brain*, *123*(2), 291–307. https://doi.org/10.1093/brain/123.2.291

Conway, B. R. (2018). The Organization and Operation of Inferior Temporal Cortex. *Annual Review of Vision Science*, *4*(1), 381–402. https://doi.org/10.1146/annurev-vision-091517-034202

Corbetta, M., Patel, G., & Shulman, G. L. (2008). The Reorienting System of the Human Brain: From Environment to Theory of Mind. *Neuron*, *58*(3), 306–324. https://doi.org/10.1016/j.neuron.2008.04.017

Crick, F., & Koch, C. (1995). Are we aware of neural activity in primary visual cortex? *Nature*, *375*(6527), Article 6527. https://doi.org/10.1038/375121a0

Cui, X., Jeter, C. B., Yang, D., Montague, P. R., & Eagleman, D. M. (2007). Vividness of mental imagery: Individual variability can be measured objectively. *Vision Research*, *47*(4), 474–478. https://doi.org/10.1016/j.visres.2006.11.013

Dance, C. J., Ipser, A., & Simner, J. (2022). The prevalence of aphantasia (imagery weakness) in the general population. *Conscious Cogn, 97*, 103243. https://doi.org/10.1016/j.concog.2021.103243

Dehaene, S., Changeux, J.-P., Naccache, L., Sackur, J., & Sergent, C. (2006). Conscious, preconscious, and subliminal processing: A testable taxonomy. *Trends in Cognitive Sciences*, *10*(5), 204–211. https://doi.org/10.1016/j.tics.2006.03.007

D'Esposito, M., Detre, J. A., Aguirre, G. K., Stallcup, M., Alsop, D. C., Tippet, L. J., & Farah, M. J. (1997). A functional MRI study of mental image generation. *Neuropsychologia*, *35*(5), 725–730. https://doi.org/10.1016/S0028-3932(96)00121-2

Dijkstra, N., Bosch, S. E., & Gerven, M. A. J. van. (2017). Vividness of Visual Imagery Depends on the Neural Overlap with Perception in Visual Areas. *Journal of Neuroscience*, *37*(5), 1367–1373. https://doi.org/10.1523/JNEUROSCI.3022-16.2016

Dijkstra, N., Bosch, S. E., & van Gerven, M. A. J. (2019). Shared Neural Mechanisms of Visual Perception and Imagery. *Trends in Cognitive Sciences*, *23*(5), 423–434. https://doi.org/10.1016/j.tics.2019.02.004

Epstein, R., Harris, A., Stanley, D., & Kanswisher, N. (1999). The parahippocampal place area: Recognition, navigation, or encoding? *Neuron*, *23*(1), 115–125. https://doi.org/10/b7jhz8

Farah, M. J. (1984). The neurological basis of mental imagery: A componential analysis. *Cognition*, *18*(1), 245–272. https://doi.org/10.1016/0010-0277(84)90026-X

Fernandino, L., Tong, J.-Q., Conant, L. L., Humphries, C. J., & Binder, J. R. (2022). Decoding the information structure underlying the neural representation of concepts. *Proceedings of the National Academy of Sciences*, *119*(6), e2108091119. https://doi.org/10/gpgvhw

Goldenberg, G. (1993). The neural basis of mental imagery. *Baillieres Clin Neurol*, *2*(2), 265–286.

Grill-Spector, K., & Weiner, K. S. (2014). The functional architecture of the ventral temporal cortex and its role in categorization. *Nature Reviews Neuroscience*, *15*(8), Article 8. https://doi.org/10.1038/nrn3747

Hajhajate, D., Kaufmann, B. C., Liu, J., Siuda-Krzywicka, K., & Bartolomeo, P. (2022). The connectional anatomy of visual mental imagery: Evidence from a patient with left occipito-temporal damage. *Brain Structure and Function*. https://doi.org/10.1007/s00429-022-02505-x

Ishai, A., Ungerleider, L. G., & Haxby, J. V. (2000). Distributed neural systems for the generation of visual images. *Neuron*, *28*(3), 979–990. https://doi.org/10.1016/s0896-6273(00)00168-9

Jackson, R. L. (2021). The neural correlates of semantic control revisited. *NeuroImage*, *224*, 117444. https://doi.org/10.1016/j.neuroimage.2020.117444

Kanwisher, N., McDermott, J., & Chun, M. M. (1997). The Fusiform Face Area: A Module in Human Extrastriate Cortex Specialized for Face Perception. *Journal of Neuroscience*, *17*(11), 4302–4311. https://doi.org/10.1523/JNEUROSCI.17-11-04302.1997

Kastner, S., Pinsk, M. A., Weerd, P. D., Desimone, R., & Ungerleider, L. G. (1999). Increased Activity in Human Visual Cortex during Directed Attention in the Absence of Visual Stimulation. *Neuron*, *22*(4), 751–761. https://doi.org/10.1016/S0896-6273(00)80734-5

Keogh, R., & Pearson, J. (2018). The blind mind: No sensory visual imagery in aphantasia. *Cortex*, *105*, 53–60. https://doi.org/10.1016/j.cortex.2017.10.012

Khanna, M. M., & Cortese, M. J. (2021). How well imageability, concreteness, perceptual strength, and action strength predict recognition memory, lexical decision, and reading aloud performance. *Memory*, *29*(5), 622–636. https://doi.org/10.1080/09658211.2021.1924789

Kosslyn, S. M., Ganis, G., & Thompson, W. L. (2001). Neural foundations of imagery. *Nat Rev Neurosci*, *2*(9), 635–642. https://doi.org/10/cfmcjh

Kosslyn, S. M., Thompson, W. L., & Ganis, G. (2006). *The case for mental imagery*. Oxford University Press.

Kourtzi, Z., & Kanwisher, N. (2001). Representation of Perceived Object Shape by the Human Lateral Occipital Complex. *Science*, *293*(5534), 1506–1509. https://doi.org/10.1126/science.1061133

Kriegeskorte, N. (2008). Representational similarity analysis – connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*. https://doi.org/10/fh5dft

Lafer-Sousa, R., Conway, B. R., & Kanwisher, N. G. (2016). Color-Biased Regions of the Ventral Visual Pathway Lie between Face- and Place-Selective Regions in Humans, as in Macaques. *Journal of Neuroscience*, *36*(5), 1682–1697. https://doi.org/10.1523/JNEUROSCI.3164-15.2016

Liu, J., & Bartolomeo, P. (2023). Probing the unimaginable: The impact of aphantasia on distinct domains of visual mental imagery and visual perception. *Cortex*, *166*, 338–347. https://doi.org/10.1016/j.cortex.2023.06.003

Liu, J., Bayle, D. J., Spagna, A., Sitt, J. D., Bourgeois, A., Lehongre, K., Fernandez-Vidal, S., Adam, C., Lambrecq, V., Navarro, V., Seidel Malkinson, T., & Bartolomeo, P. (2023). Fronto-parietal networks shape human conscious report through attention gain and reorienting. *Communications Biology*, *6*(730), Article 1. https://doi.org/10.1038/s42003-023-05108-2

Liu, J., Spagna, A., & Bartolomeo, P. (2022). Hemispheric asymmetries in visual mental imagery. *Brain Structure and Function*, *227*(2), 697–708. https://doi.org/10.1007/s00429-021-02277-w

Malach, R., Reppas, J. B., Benson, R. R., Kwong, K. K., Jiang, H., Kennedy, W. A., Ledden, P. J., Brady, T. J., Rosen, B. R., & Tootell, R. B. (1995). Object-related activity

revealed by functional magnetic resonance imaging in human occipital cortex. *Proceedings of the National Academy of Sciences*, *92*(18), 8135–8139. https://doi.org/10.1073/pnas.92.18.8135

Mashour, G. A., Roelfsema, P., Changeux, J.-P., & Dehaene, S. (2020). Conscious Processing and the Global Neuronal Workspace Hypothesis. *Neuron*, *105*(5), 776–798. https://doi.org/10.1016/j.neuron.2020.01.026

Mazard, A., Laou, L., Joliot, M., & Mellet, E. (2005). Neural impact of the semantic content of visual mental images and visual percepts. *Cognitive Brain Research*, *24*(3), 423–435. https://doi.org/10.1016/j.cogbrainres.2005.02.018

Mechelli, A., Price, C. J., Friston, K. J., & Ishai, A. (2004). Where bottom-up meets top-down: Neuronal interactions during perception and imagery. *Cereb Cortex*, *14*(11), 1256–1265. https://doi.org/10.1093/cercor/bhh087

Milton, F., Fulford, J., Dance, C., Gaddum, J., Heuerman-Williamson, B., Jones, K., Knight, K. F., MacKisack, M., Winlove, C., & Zeman, A. (2021). Behavioral and Neural Signatures of Visual Imagery Vividness Extremes: Aphantasia versus Hyperphantasia. *Cerebral Cortex Communications*, *2*(2), tgab035. https://doi.org/10.1093/texcom/tgab035

Moro, V., Berlucchi, G., Lerch, J., Tomaiuolo, F., & Aglioti, S. M. (2008). Selective deficit of mental visual imagery with intact primary visual cortex and visual perception. *Cortex*, *44*(2), 109–118. https://doi.org/10.1016/j.cortex.2006.06.004

Naselaris, T., Olman, C. A., Stansbury, D. E., Ugurbil, K., & Gallant, J. L. (2015). A voxel-wise encoding model for early visual areas decodes mental images of remembered scenes. *Neuroimage*, *105*, 215–228. https://doi.org/10.1016/j.neuroimage.2014.10.018

O'Craven, K. M., & Kanwisher, N. (2000). Mental imagery of faces and places activates corresponding stiimulus-specific brain regions. *J Cogn Neurosci*, *12*(6), 1013–1023. https://doi.org/10.1162/08989290051137549

Pearson, J. (2019). The human imagination: The cognitive neuroscience of visual mental imagery. *Nature Reviews Neuroscience*, *20*(10), Article 10. https://doi.org/10.1038/s41583-019-0202-9

Pounder, Z., Jacob, J., Evans, S., Loveday, C., Eardley, A. F., & Silvanto, J. (2022). Only minimal differences between individuals with congenital aphantasia and those with typical imagery on neuropsychological tasks that involve imagery. *Cortex*, *148*, 180–192. https://doi.org/10.1016/j.cortex.2021.12.010

Pyke, A. A., Fincham, J. M., & Anderson, J. R. (2017). When math operations have visuospatial meanings versus purely symbolic definitions: Which solving stages and brain regions are affected? *NeuroImage*, *153*, 319–335.

Reddy, L., Tsuchiya, N., & Serre, T. (2010). Reading the mind's eye: Decoding category information during mental imagery. *Neuroimage*, *50*(2), 818–825. https://doi.org/10.1016/j.neuroimage.2009.11.084

Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K., & Goebel, R. (2021). A Probabilistic Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cerebral Cortex*, *31*(1), 603–619. https://doi.org/10.1093/cercor/bhaa246

Santarpia, A., Blanchet, A., Poinsot, R., Lambert, J. F., Mininni, G., & Thizon-Vidal, S. (2008). Évaluer la vivacité des images mentales dans différentes populations françaises. *Pratiques Psychologiques*, *14*(3), 421–441. https://doi.org/10.1016/j.prps.2007.11.001

Saxe, R., Brett, M., & Kanwisher, N. (2006). Divide and conquer: A defense of functional localizers. *NeuroImage*, *30*(4), 1088–1096. https://doi.org/10.1016/j.neuroimage.2005.12.062

Senden, M., Emmerling, T. C., van Hoof, R., Frost, M. A., & Goebel, R. (2019). Reconstructing imagined letters from early visual cortex reveals tight topographic correspondence between visual mental imagery and perception. *Brain Struct Funct*, *224*(3), 1167–1183. https://doi.org/10.1007/s00429-019-01828-6

Shulman, G. L., Astafiev, S. V., McAvoy, M. P., d'Avossa, G., & Corbetta, M. (2007). Right TPJ deactivation during visual search: Functional significance and support for a filter hypothesis. *Cerebral Cortex*, *17*(11), 2625–2633. https://doi.org/10/fhgdwq

Siuda-Krzywicka, K., Boros, M., Bartolomeo, P., & Witzel, C. (2019). The biological bases of colour categorisation: From goldfish to the human brain. *Cortex*, *118*, 82–106. https://doi.org/10.1016/j.cortex.2019.04.010

Sokolowski, H. M., & Levine, B. (2023). Common neural substrates of diverse neurodevelopmental disorders. *Brain*, *146*(2), 438–447. https://doi.org/10.1093/brain/awac387

Spagna, A., Bayle, D. J., Romeo, Z., Seidel-Malkinson, T., Liu, J., Yahia-Cherif, L., Chica, A. B., & Bartolomeo, P. (2022). The cost of attentional reorienting on conscious visual perception: An MEG study. *Cerebral Cortex*, bhac192. https://doi.org/10.1093/cercor/bhac192

Spagna, A., Hajhajate, D., Liu, J., & Bartolomeo, P. (2021). Visual mental imagery engages the left fusiform gyrus, but not the early visual cortex: A meta-analysis of neuroimaging evidence. *Neuroscience & Biobehavioral Reviews*, *122*, 201–217. https://doi.org/10.1016/j.neubiorev.2020.12.029

Spagna, A., Heidenry, Z., Miselevich, M., Lambert, C., Eisenstadt, B. E., Tremblay, L., Liu, Z., Liu, J., & Bartolomeo, P. (2024). Visual mental imagery: Evidence for a heterarchical neural architecture. *Physics of Life Reviews*, *48*, 113–131. https://doi.org/10.1016/j.plrev.2023.12.012

Sporns, O., & Betzel, R. F. (2016). Modular Brain Networks. *Annual Review of Psychology*, *67*(1), 613–640. https://doi.org/10.1146/annurev-psych-122414-033634

Stokes, M., Thompson, R., Cusack, R., & Duncan, J. (2009). Top-Down Activation of Shape-Specific Population Codes in Visual Cortex during Mental Imagery. *Journal of Neuroscience*, *29*(5), 1565–1572. https://doi.org/10.1523/JNEUROSCI.4657-08.2009

Thirion, B., Duchesnay, E., Hubbard, E., Dubois, J., Poline, J.-B., Lebihan, D., & Dehaene, S. (2006). Inverse retinotopy: Inferring the visual content of images from brain activation patterns. *NeuroImage*, *33*(4), 1104–1116. https://doi.org/10.1016/j.neuroimage.2006.06.062

Thorudottir, S., Sigurdardottir, H. M., Rice, G. E., Kerry, S. J., Robotham, R. J., Leff, A. P., & Starrfelt, R. (2020). The Architect Who Lost the Ability to Imagine: The Cerebral Basis of Visual Imagery. *Brain Sci*, *10*(2). https://doi.org/10.3390/brainsci10020059

Tsao, D. Y., Schweers, N., Moeller, S., & Freiwald, W. A. (2008). Patches of face-selective cortex in the macaque frontal lobe. *Nature Neuroscience*, *11*(8), Article 8. https://doi.org/10.1038/nn.2158

van Vugt, B., Dagnino, B., Vartak, D., Safaai, H., Panzeri, S., Dehaene, S., & Roelfsema, P. R. (2018). The threshold for conscious report: Signal loss and response bias in visual and frontal cortex. *Science*, *360*(6388), 537–542. https://doi.org/10/gc8jfv

Yomogida, Y., Sugiura, M., Watanabe, J., Akitsuki, Y., Sassa, Y., Sato, T., Matsue, Y., & Kawashima, R. (2004). Mental Visual Synthesis is Originated in the Fronto-temporal Network of the Left Hemisphere. *Cerebral Cortex*, *14*(12), 1376–1383. https://doi.org/10.1093/cercor/bhh098

Zeman, A., Dewar, M., & Della Sala, S. (2015). Lives without imagery – Congenital aphantasia. *Cortex*, *73*, 378–380. https://doi.org/10.1016/j.cortex.2015.05.019

Zhan, M., Pallier, C., Agrawal, A., Dehaene, S., & Cohen, L. (2023). Does the visual word form area split in bilingual readers? A millimeter-scale 7-T fMRI study. *Science Advances*, *9*(14), eadf6140. https://doi.org/10.1126/sciadv.adf6140

Zhen, Z., Yang, Z., Huang, L., Kong, X., Wang, X., Dang, X., Huang, Y., Song, Y., & Liu, J. (2015). Quantifying interindividual variability and asymmetry of face-selective regions: A probabilistic functional atlas. *NeuroImage*, *113*, 13–25. https://doi.org/10.1016/j.neuroimage.2015.03.010