1   **Correlating phylogenetic and functional diversity of the *nod*-free but nodulating**
2   ***Bradyrhizobium* phylogroup**

4   Lu Ling[1,2^], Alicia Camuel[3,4^], Sishuo Wang[1], Xiaojun Wang[2], Tianhua Liao[1], Jinjin Tao[1],
5   Xingqin Lin[2], Nico Nouwen[3,4], Eric Giraud[3,4*], Haiwei Luo[1,5*]

6
[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory
of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China
[2]Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, China
[3]IRD, Laboratoire des Symbioses Tropicales et Méditerranéennes (LSTM), UMR IRD/ Institut
Agro/INRAE/ Université Montpellier France
[4]PHIM Plant Health Institute, Université Montpellier, IRD, CIRAD, INRAE, Institut Agro,
Montpellier, France
[5]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong,
Shatin, Hong Kong SAR, China

16
17 [^]**These are co-first authors**: Lu Ling, Alicia Camuel

18
19 [*]**These are co-corresponding authors:** Eric Giraud (eric.giraud@ird.fr), Haiwei Luo
20 (haiweiluo@cuhk.edu.hk)

21
22 **Running title:** Expanded diversity of the Photosynthetic *Bradyrhizobium*

23
24 **Key words:** *Bradyrhizobium*, *Aeschynomene*, nodulation, population genomics
25

Abstract

Bradyrhizobium is a main rhizobial lineage of which most members nodulate legume plants using Nod factors (NFs) synthetized by the *nod* genes. However, members of the Photosynthetic supergroup (phylogroup) within *Bradyrhizobium* (PB) are *nod*-free but still capable of establishing nitrogen-fixing nodules with some tropical legumes of the *Aeschynomene* genus. These unusual findings are based on the genomic sequences of only 13 PB strains, and almost all were isolated from *Aeschynomene* nodules. Here, we investigate the diversity of *Bradyrhizobium* in grassland, forest, and rice field by *rpoB* amplicon sequencing and report that PB is mainly associated with rice root and rhizosphere. Moreover, we sequenced 209 new PB members isolated mostly from the rice field. The extended PB supergroup comprises three major clades: a basal clade with significant expansion of its diversity, followed by an intermediate clade composed by two strains, and a new clade exclusively represented by our new strains. Although the PB strains universally lack the canonical *nod* genes, all 28 assayed strains covering the broad diversity of these clades induced nodules on *Aeschynomene indica*. Interestingly, the three clades displayed significant differences in the efficiency of symbiosis, aligning well with their phylogenetic branching order. Our strain collection expands the ecological, phylogenetic and functional diversity of *nod*-free but nodulating *Bradyrhizobium*. With this expanded diversity, we conclude that the NF-independent nodulation of *Aeschynomene* is a common trait of this supergroup, in contrast to the photosynthetic trait originally thought as its unifying feature.

Introduction

The genus *Bradyrhizobium* is one of the largest and most diverse rhizobial genera and the primary symbiont of a wide range of legumes (Parker, 2015; Sprent et al., 2017). It encompasses seven phylogenetic supergroups (phylogroups) (Avontuur et al., 2019; Ormeno-Orrillo and Martinez-Romero, 2019), among which the Photosynthetic *Bradyrhizobium* (PB) supergroup is one of the most special as photosynthesis is a rare trait in rhizobia (Avontuur et al., 2019). Members belonging to this supergroup engage in a mutualistic relationship with some tropical semi-aquatic species of the *Aeschynomene* genus (Giraud and Fleischman, 2004), and they form nodules not only on roots but also on stems (Chaintreuil et al., 2013; Mornico et al., 2011). Further, none of the completely sequenced PB members carry *nodABC* genes encoding the enzymes synthesizing the core structure of the nodulation factors (NF), the only exception being *Bradyrhizobium* sp. ORS285, which can use both a *nod*-dependent and -independent pathway reliant to the *Aeschynomene* host plant (Bonaldi et al., 2011; Giraud et al., 2007). This indicates that the PB phylogroup uses a novel mechanism of interaction with legume hosts that differs from the traditional universal NF-based pathway (Lerouge et al., 1990; Oldroyd, 2013). Moreover, all known PB strains carry photosynthetic genes except for four deep-branching members isolated from French Guiana as represented by *Bradyrhizobium* sp. STM3843 (Miché et al., 2010; Mornico et al., 2011).

Despite these unique features, the research on PB has been very limited. Currently, there are only 13 PB strains that have their genomes sequenced, largely limiting genome-based analysis on this important *Bradyrhizobium* phylogroup. Although most PB strains were isolated from *Aeschynomene* spp. nodules, the PB phylogroup was also detected in paddy soil, rice roots as well as in lake water (Chaintreuil et al., 2000; Okubo et al., 2013; Piromyou et al., 2015; Van

Berkum and Eardly, 2002), suggesting that its members have much wider ecological niches than

previously thought. Here, we report that rice fields are an important reservoir of PB among the

terrestrial ecosystems through *rpoB* amplicon sequencing analysis. We isolated and genome-

sequenced 209 isolates predominantly from three niches of rice (within root, rhizosphere, bulk

soil) and discovered a novel deeply branching clade. We further show that phylogenomic

diversity of PB members match well with their symbiosis efficiency with *Aeschynomene indica*.

Materials and methods

All the methodological details were described in the Supplementary Text. Three plant

species (*Oryza sativa* indica, *Houttuynia cordata,* and *Camphora officinarum*) and soils, each

with three replicates, were collected from paddy field, grassland and forest, with each replicate

5-10 m apart from the others, respectively, in Hunan province, China (27.948 °N, 113.221 °E) in

July 2021. Three typical niches (bulk soil, rhizosphere and root) of each plant species (three

replicates) were each used for bacterial isolation using a modified arabinose-gluconate (MAG)

medium. Basic soil characteristics were also measured. The taxonomic affiliation of isolates was

determined by 16S rRNA gene analysis. Additionally, *Oryza sativa* japonica plants were

collected in Hong Kong, China (22.418 °N, 114.080 °E) in May 2022 for *Bradyrhizobium*

isolation following the same procedure.

DNA was extracted from fresh soil and plant root samples (0.25 g) using the DNeasy

PowerSoil Pro Kit (QIAGEN) according to the manufacturer's protocol and then sent to the

company (Magigene, Guangdong) for *rpoB* amplicon sequencing. As the quality and quantity of

DNA extracted from the second replicate of the rice root sample were poor and the amount of the

root sample was insufficient to support multiple DNA extractions, amplicon sequencing was not

performed on this sample. The quality control of the raw reads was performed with Trimmomatic

v0.39 (Bolger et al., 2014). Subsequently, the paired-end reads of the *rpoB* amplicon sequences

were processed with a denoising algorithm (DADA2) (Callahan et al., 2016) implemented in

QIIME2 (Bolyen et al., 2019) to perform sequence denoising, dereplication, and chimera

filtering to generate amplicon sequence variants (ASVs). The generated ASVs were filtered out

the non-*rpoB* sequences and assigned to each *Bradyrhizobium* supergroup as well as each of the

three clades of Photosynthetic supergroup through a phylogenetic placement method (Czech et

al., 2022), which was used to determine their relative abundance.

To distinguish each *Bradyrhizobium* supergroup, two *rpoB* gene trees, using the full length

(Fig. S1A) and amplified region (Fig. S1B) respectively, were constructed with IQ-Tree v2.2.0

(Minh et al., 2020) based on the sequence of *rpoB* genes retrieved from *Bradyrhizobium*

genomes (209 new genomes and 566 public genomes downloaded from the NCBI Genbank

database). In addition, to identify the diversity of Photosynthetic *Bradyrhizobium* in each sample,

a *rpoB* gene tree was also built by combining ASVs and amplified regions from Photosynthetic

Bradyrhizobium genomes (209 new genomes and 13 public genomes). Although phylogenetic

resolution of the *rpoB* gene faded when the short amplified region was used (Fig. S1B), members

from each supergroup remain clustered though broken into several subclades, suggesting the use

of the tree based on short amplified region has limited effect on amplicon sequence variants

(ASVs) assignment using the commonly used phylogenetic placement method (Janssen et al.,

2018).

A phylogenomic tree of *Bradyrhizobium* was built using IQ-Tree v2.2.0 (Minh et al., 2020)

with our 209 isolates and 566 public genomes (outgroup included) based on 123 shared single-

copy genes identified in a previous phylogenomic study of *Bradyrhizobium* (Tao et al., 2021). As

our isolates mainly fall into Photosynthetic *Bradyrhizobium* supergroup, we also performed

phylogenomic and comparative genomic analyses for this supergroup to understand their

phylogenetic and population structure. PopCOGenT (Arevalo et al., 2019) was used to delineate

genetically isolated populations for the 222 Photosynthetic *Bradyrhizobium* genomes (13 public

genomes and 209 new genomes). The population boundaries delineated by this method is based

on recent gene flow barriers, matching well with the idea that bacterial speciation often proceeds

rapidly (Arevalo et al., 2019). Next, we selected 28 strains spanning over the main clades and the

populations defined by PopCOGenT for symbiotic assays including nodulation and nitrogen

fixation capabilities on a tropical legume species *Aeschynomene indica*.

Results and discussion

<u>Photosynthetic supergroup of *Bradyrhizobium* is enriched in rice cropland compared to forest</u>

<u>and grassland</u>

Both the *rpoB* gene tree (Fig. S1A) and the phylogenomic tree (Fig. S2) grouped

Bradyrhizobium members into seven supergroups including the Photosynthetic *Bradyrhizobium*

(PB). Their largely congruent tree topological structures support the idea that *rpoB* is an

appropriate marker gene (Ogier et al., 2019) to distinguish *Bradyrhizobium* supergroups. We

therefore designed specific *rpoB* primers to investigate the relative abundance and diversity of

each *Bradyrhizobium* supergroup in different terrestrial ecosystems including three plant species

(One plant species specific to each ecosystem was chosen, *Oryza sativa* subsp. *Indica* for

cropland, *Houttuynia cordata* for grassland*,* and *Camphora officinarum* for forest). We show that

most *Bradyrhizobium* supergroups, including PB, are widely distributed in forest, grassland, and

cropland (Fig. 1A, Dataset S1). Among these, PB is notably enriched in the root and rhizosphere

niches of rice (Fig. 1B, Dataset S2), implying that members of this supergroup have the potential

to be used as plant growth promoting bacteria (PGPB) to promote rice growth and yield.

<u>Our culture collection contributes a novel deeply branching clade of the Photosynthetic</u>

<u>supergroup of *Bradyrhizobium*</u>

With the 209 newly isolated members (Dataset S3), the genome-based phylogeny of the PB

supergroup is split into three deeply branching clades (see Fig. S2 for their phylogenetic position

in the species tree of the entire *Bradyrhizobium* genus). Clade 1 represents the evolutionarily

basal lineage. It initially comprised only 12 strains (Dataset S4), but it now has its diversity

expanded by having 68 new strains representing several new lineages delineated as distinct

genetically isolated populations (Fig. 2, Dataset S3, S4; also discussed in the next section). Clade

2 has only two strains (the publicly available *Bradyrhizobium* sp. STM3843 and our newly

sequenced HKCCYLS1011 isolated from *Oryza sativa* subsp. *japonica*) and represents a

phylogenetic intermediate among the three clades. Clade 3 consists of 140 newly isolated strains.

The evolutionary branching order of the three clades is verified with two outgroup-independent

methods (Fig. 2, Fig. S3) and the outgroup-dependent method (Fig. S2). Collectively, our new

strains appreciably increase the existing phylogenetic diversity of the PB supergroup.

Comparative genomic analysis shows that except for *Bradyrhizobium* sp. ORS285, which is

known to use both a *nod*-dependent and -independent strategy for nodulation (Giraud et al.,

2007), all PB supergroup members lack the *nod* genes. Our *rpoB* amplicon analysis showed that

PB members in all analyzed ecosystems are dominated by the PB Clade 1, and distribution of the

newly isolated Clade 3 is mainly limited in rice-associated niches (Fig. 1B, Dataset S2). It is

worth mentioning that the ASVs assigned to Clade 3 are closely related to the cultured members

of Clade 3, but none of them have identical sequences to the cultures (Fig. S4). This suggests that

future work on cultivation is needed to further expand the fine-scale diversity within Clade 3.

These three major clades may have diverged functionally as each has a unique set of

ecologically relevant genes. For example, genes encoding C4-dicarboxylate transporter (*dctBD*)

are universally and exclusively found in Clade 1 (Fig. S5, Dataset S5), potentially enabling

Clade 1 strains to acquire C4-dicarboxylate compounds (e.g., malate and succinate) from the host

plants to fuel the energy-intensive nitrogen fixation (Liu et al., 2018). Other Clade 1-associated

genes, though not necessarily exclusively found in Clade 1, include those involved in carbon

metabolism and energy conservation processes, such as those encoding histidine utilization

enzymes (*hutCFHIU*), glutaconate CoA-transferase (*gctAB*), raffinose/stachyose/melibiose

transport system (*msmEFG*), O_2-independent ubiquinone biosynthesis genes (*ubiDTUV*) and

nitrate reductase (*narGHIJ* and *napABCDE*) (Fig. S5, Dataset S5). Specifically, the *hut* pathway

is a catabolic pathway that allows using histidine as a source of carbon, nitrogen and energy for

growth and to facilitate nitrogen fixation (Bender, 2012). The two non-homologous dissimilatory

nitrate reductases (*narGHIJ* and *napABCDE*) and the O_2-independent ubiquinone biosynthesis

genes (*ubiDTUV*) are both necessary for denitrification process (Gregory et al., 2003; Vo et al.,

2020).The *narGHIJ* carry out nitrate respiration primarily under anaerobic conditions (Gregory

et al., 2003), whereas the *napABCDE* reduce nitrate under both anaerobic and aerobic condition

(Gregory et al., 2003).

The genes prevalent to Clade 3 include the *mxa* gene cluster, *mgsABC*, and *mdh12* (Fig. S5,

Dataset S5, S6). These genes potentially facilitate the utilization of methanol, which is a waste

product during plant-cell wall degradation (Nemecek-Marshall et al., 1995), although a

physiological assay did not support this hypothesis (Fig. S6). This suggests that expression of

these genes might be regulated by unknown mechanisms. Another unique gene cluster in Clade 3

codes for a protein complex (*exoALOUWY*) (Fig. S5, Dataset S5, S6) probably involved in the

synthesis of the major components of exopolysaccharide (EPS), which may induce the immune

response of plants (De Sousa et al., 2021; Quelas et al., 2010). Plant LysM kinase receptors

perceive EPS, and depending on the compositions, EPS could either impair or promote

symbiosis between rhizobia and plants (De Sousa et al., 2021). Other Clade 3-associated genes

include malonate decarboxylase (*mdcABCDEG*), UDP-glucose/iron transport system (*STAR12*)

and tungstate transport system (*tupABC*) (Fig. S5, Dataset S5). The presence of these genes

implies potential unique carbon metabolism (Chohnan and Takamura, 2004) and transporters

(Hawkins et al., 2017) in this clade.

Clade 2 consists of only two strains, making it difficult to draw conclusive findings on their gene

composition. However, it is still interesting to note that as an intermediate clade, the two strains

in Clade 2 uniquely have a gene cluster for the type VI secretion system (T6SS) (Fig. S5, Dataset

S5), which is known to influence bacterial competitiveness and symbiosis with eukaryotes (De

Sousa et al., 2023; Tighilt et al., 2022). The effects on symbiosis with plants may be closely

related to the effectors secreted by T6SS that interact with the host plant and surrounding

microbiota (De Sousa et al., 2023). Other specific genes in these two strains include pyruvate

ferredoxin oxidoreductase (*porABC*) (Fig. S5, Dataset S5), which exclusively supports

fermentation under anaerobic conditions and releases energy at the same time (Wang et al., 2022).

<u>Fine-scale population structure of the Photosynthetic supergroup of *Bradyrhizobium*</u>

Apart from the broad diversity of the PB supergroup, we further asked whether fine-scale

phylogenetic differentiation correlates with the ecological niches where the PB supergroup

members were found. As geography may have an important impact on the genetic diversity of

rhizobial populations (Greenlon et al., 2019), we performed intensive isolation of PB members

from the same rice field to control for the potential impact of geographic separation on PB

population structure. Of the 209 new PB supergroup strains, 205 were collected from three rice

plants located in the same rice field. Bacterial isolation was performed from within the root,

rhizosphere, and bulk soil for each plant, collectively giving rise to 145, 23 and 41 PB strains,

respectively (Dataset S3). To facilitate the correlative analysis between fine-scale phylogenetic

groups and the niches, we assigned isolates' genomes into populations defined by PopCOGenT

(reported as "main cluster" or MC) (Arevalo et al., 2019). As bacterial members within a

population have significantly higher recombination rate than those across populations,

"population" defined here aligns with the "species" in higher eukaryotes. Additionally,

PopCOGenT can detect subpopulations (reported as 'subclusters'), which are under ongoing

differentiation within a population.

Using this approach, we show that i) all strains in Clade 3 share the membership of a single

population (MC1), ii) the two strains in Clade 2 each form a distinct population (MC id not

given), though they may be deemed as members of a single operational species as their genome-

wide average nucleotide identity (ANI) is 96.5% (Fig. S7, Dataset S7), exceeding the species

threshold of 95%, and iii) strains from Clade 1 fall into numerous populations, among which,

MC2, MC3, MC4, MC5 and MC6 are each exclusively comprised of the new strains, whereas

MC7, MC8 and the remaining six unassigned isolates each forming a distinct population are all

publicly available (Fig. 2). In Clade 1, the within- and between-population similarity is above

and below 95% ANI (Fig. S7, Dataset S7), respectively, consistent with the operational species

threshold of 95% ANI (Konstantinidis and Tiedje, 2005), whereas the 16S rRNA gene shows

little divergence with the between-population similarity generally above 99% (Fig. S7, Dataset

S7). Based on the available samples, intra-population subdivision occurred within MC1, MC2

and MC3. We found that each population and subpopulation have members sampled from

different plants and niches (Fig. 2), suggesting that the PB populations sampled from the rice

filed are not genetically subdivided according to the physical separation between the plant

individuals or following the niche separation between root, rhizosphere, and bulk soil.

We also found interesting associations between important metabolic pathways and

population identity. Remarkably, although the PB supergroup was named by the presence of the

photosynthetic genes (Avontuur et al., 2019), these genes are exclusively and universally found

in MC2, MC5, MC6, MC7, MC8 and the unassigned individuals in Clade 1 but completely

missing from MC3 and MC4 of Clade 1, Clade 2 and Clade 3 (MC1) (Fig. 2, Dataset S6). Our

result indicates that photosynthesis is not a characteristic trait that defines the PB supergroup.

Also interesting is the prevalence of a unique Type III secretion system (T3SS) in MC2, MC3

and MC4 but completely missing in other populations (Fig. 2, Fig. S8B, S8C, Dataset S6). It is

one of the six T3SS subtypes identified from known *Bradyrhizobium* phylotypes (Teulet et al.,

2022; Teulet et al., 2020). It was previously identified only in the PB strain *Bradyrhizobium*

oligotrophicum S58 (thus named 'S58-T3SS' subtype) and its function remains unknown (Teulet

et al., 2020). Since the general function of T3SS is to translocate effector proteins into host cells

that modulate the host immune response (Teulet et al., 2022), it cannot be excluded that this

T3SS type specifically identified in PB members plays an important role during their interaction

with their host plants (*Aeschynomene* spp. and rice).

The ability to nodulate *Aeschynomene indica* is a conserved trait shared by Photosynthetic

supergroup of *Bradyrhizobium* but differs between the major clades

All PB supergroup members have a *nif* gene cluster necessary for nitrogen fixation.

Duplication of the *nifH* gene which encodes one of the structural proteins of the nitrogenase

enzyme complex was observed in all PB strains (Fig. S8A), but the functional consequence is not

known. Across the PB supergroup members, an important difference was observed regarding

how the *nif* genes are structured. In most populations, all *nif* genes are co-located, along with

other genes (e.g., *sufBCDX*, *glbO* and *fixABCX*) potentially involved in nitrogen fixation. This is

not the case for a few members such as some basal lineages (e.g., *Bradyrhizobium* sp LMG 8443)

and some members assigned to several populations (e.g., SZCCHNR1015 in MC5 and

Bradyrhizobium sp. 83002 in MC7), where the *nif* genes are present in two contigs. This is likely

due to structural rearrangement, but a DNA assembly artefact cannot be ruled out.

A common feature observed of the PB *nif* cluster is the universal presence of the *nifV* gene

encoding for a homocitrate synthase that is involved in the biosynthesis of the nitrogenase

cofactor (FeMo-Co), which is absent in most other rhizobial lineages (Hakoyama et al., 2009;

Nouwen et al., 2017). An acetylene reduction assay on 28 representative PB strains, which cover

all three major clades, all newly identified populations (MC1 to MC6), most of the identified

subpopulations within each population, and the model PB strain ORS278 as control, confirmed

their ability to fix nitrogen under free-living conditions (Fig. S9A), a trait absent in most *nod*-

carrying rhizobia (Nouwen et al., 2017).

The plant inoculation experiments showed that all 28 representative strains were also able to

induce nodules on *A. indica* (Fig. 3B, 3C). However, a significant difference in symbiosis

efficiency was observed between the major clades. Clade 1 strains had the same symbiotic

properties as the model strain ORS278, inducing many nitrogen-fixing nodules that stimulated

the growth of the plants (Fig. 3A, 3B, Fig. S9B, S10A). In contrast, Clade 3 strains formed much

fewer nodules compared to Clade 1 strains and the measured nitrogenase enzyme activity was

lower under the symbiotic condition (Fig. 3B, Fig. S9B). Microscopic analysis showed that the

nodules elicited by Clade 3 strains displayed multiple aberrant phenotypes: i) some nodules

contained necrotic areas, ii) in others the central tissue was completely digested, and finally, iii)

the nodules that displayed less drastic symptoms contained mainly dead bacteria (Fig. 3C and

3F). These observations indicate that besides inducing fewer nodules, Clade 3 strains are unable

to maintain an effective chronic infection, which explains why they have no beneficial effect on

plant growth (Fig. 3A, 3C, 3F). Clade 2 represented by only two strains STM3843 and

HKCCYLS1011 exhibits a transitional pattern of phenotype, consistent with its intermediate

phylogenetic position between Clade 1 and Clade 3. The number (Fig. 3B) and phenotype (Fig.

3C, 3E) of the nodules elicited by strain STM3843 are comparable to those of Clade 1 strains

(Fig. 3B, 3C, 3D) though stimulation of plant growth is less (Fig. 3A), whereas the nodules

stimulated by strain HKCCYLS1011 (Fig. 3B, 3C, 3E) are like those of Clade 3 strains (Fig. 3B,

3C, 3F) and no effect on plant growth was observed (Fig. 3A).

The symbiosis efficiency differences observed between the major clades may result from an

inability of the Clade 3 strains to cope with the plant's immune response, and/or an over-

induction of the plant's defense mechanisms due to the absence or not properly recognized

symbiotic signal(s). However, it is remarkable that all strains from the PB supergroup, which

come from different origins and geographical locations, are capable of nodulating *A. indica*. The

ability to develop an NF-independent nodulation with *Aeschynomene* spp. thus appears to be a

common trait of PB supergroup members. Thus, unlike the *nod* genes of other rhizobial lineages

which are accessory genes acquired by horizontal transfer, the genes governing NF-independent

symbiosis may be essential genes belonging to the core genome of the PB supergroup.

Concluding remarks

The results of *rpoB* amplicon sequencing of samples from forest, grassland, and rice field

indicate that members of the Photosynthetic *Bradyrhizobium* (PB) are enriched in rice field. By

large-scale isolation and genome sequencing, we report a novel major clade, thereby greatly

expanding the phylogenetic diversity of the cultured members of the PB supergroup. An

important finding is that although all assayed phylogenetically diverse PB strains can nodulate

Aeschynomene spp., the branching order of the three major clades correlates nicely with their

symbiosis efficiency, with the clade taking the intermediate phylogenetic position showing the

symbiosis efficiency in between. It is therefore likely that the ability to establish symbiosis with

Aeschynomene plants, which often grow in the same wetlands as rice, is a key factor that shapes

the broad diversity of the PB supergroup. At the population level, no significant difference in

symbiosis efficiency was observed, suggesting that other traits such as photosynthesis and T3SS

that are uniquely associated with some but not all populations, might be among the important

drivers of population-level genetic differentiation and ecological adaptation. Collectively, our

study provides insights into the ecology and evolution of the Photosynthetic supergroup within

the globally dominant soil bacteria *Bradyrhizobium*, and additionally serves as a prime example

that links deep phylogenetic diversity of an ecologically relevant bacterial group to their major

phenotypic and functional diversity.

Acknowledgements

327   We thank Xiaoyuan Feng for helping with genome assembly. This work was supported by the

328   Hong Kong Research Grants Council Area of Excellence Scheme (AoE/M-403/16) and a grant

329   from the French National Research Agency ("ET-Nod"; ANR-20-CE20-0012).

Data availability

332   The genomic sequences and raw reads of the 209 newly sequenced *Bradyrhizobium* isolates have

333   been uploaded to NCBI (Project ID: PRJNA983111).

Competing interests

336   The authors declare no competing interests in relation to this work.

Reference

Arevalo, P., VanInsberghe, D., Elsherbini, J., Gore, J., and Polz, M.F. (2019). A Reverse Ecology

Approach Based on a Biological Definition of Microbial Populations. Cell *178*, 820-834 e814.

Avontuur, J.R., Palmer, M., Beukes, C.W., Chan, W.Y., Coetzee, M.P.A., Blom, J., Stepkowski,

T., Kyrpides, N.C., Woyke, T., Shapiro, N.*, et al.* (2019). Genome-informed *Bradyrhizobium*

taxonomy: where to from here? Syst Appl Microbiol *42*, 427-439.

Bender, R.A. (2012). Regulation of the histidine utilization (*hut*) system in bacteria. Microbiol

Mol Biol Rev *76*, 565-584.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics *30*, 2114-2120.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A.,

Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F.*, et al.* (2019). Reproducible, interactive,

scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology *37*, 852-

857.

Bonaldi, K., Gargani, D., Prin, Y., Fardoux, J., Gully, D., Nouwen, N., Goormachtig, S., and

Giraud, E. (2011). Nodulation of *Aeschynomene afraspera* and *A. indica* by photosynthetic

Bradyrhizobium Sp. strain ORS285: the *nod*-dependent versus the *nod*-independent symbiotic

interaction. Mol Plant Microbe Interact *24*, 1359-1371.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nature

Methods *13*, 581-583.

Chaintreuil, C., Arrighi, J.F., Giraud, E., Miche, L., Moulin, L., Dreyfus, B., Munive-Hernandez,

J.A., Villegas-Hernandez Mdel, C., and Bena, G. (2013). Evolution of symbiosis in the legume

genus *Aeschynomene*. New Phytol *200*, 1247-1259.

Chaintreuil, C., Giraud, E., Prin, Y., Lorquin, J., Ba, A., Gillis, M., de Lajudie, P., and Dreyfus, B.

(2000). Photosynthetic *bradyrhizobia* are natural endophytes of the African wild rice Oryza

breviligulata. Appl Environ Microbiol *66*, 5437-5447.

Chohnan, S., and Takamura, Y. (2004). Malonate decarboxylase in bacteria and its application for

determination of intracellular acyl-CoA thioesters. Microbes and Environments *19*, 179-189.

Czech, L., Stamatakis, A., Dunthorn, M., and Barbera, P. (2022). Metagenomic Analysis Using

Phylogenetic Placement-A Review of the First Decade. Front Bioinform *2*, 871393.

De Sousa, B.F.S., Castellane, T.C.L., Tighilt, L., Lemos, E.G.d.M., and Rey, L. (2021). Rhizobial

Exopolysaccharides and Type VI Secretion Systems: A Promising Way to Improve Nitrogen

Acquisition by Legumes. Frontiers in Agronomy *3*.

De Sousa, B.F.S., Domingo-Serrano, L., Salinero-Lanzarote, A., Palacios, J.M., and Rey, L.

(2023). The T6SS-Dependent Effector Re78 of *Rhizobium* etli Mim1 Benefits Bacterial

Competition. Biology (Basel) *12*.

Giraud, E., and Fleischman, D. (2004). Nitrogen-fixing symbiosis between photosynthetic

bacteria and legumes. Photosynthesis Research *82*, 115-130.

Giraud, E., Moulin, L., Vallenet, D., Barbe, V., Cytryn, E., Avarre, J.C., Jaubert, M., Simon, D.,

Cartieaux, F., Prin, Y.*, et al.* (2007). Legumes symbioses: absence of *Nod* genes in photosynthetic

Bradyrhizobia. Science *316*, 1307-1312.

Greenlon, A., Chang, P.L., Damtew, Z.M., Muleta, A., Carrasquilla-Garcia, N., Kim, D., Nguyen,

H.P., Suryawanshi, V., Krieg, C.P., Yadav, S.K.*, et al.* (2019). Global-level population genomics

reveals differential effects of geography and phylogeny on horizontal gene transfer in soil

bacteria. Proc Natl Acad Sci U S A *116*, 15200-15209.

Gregory, L.G., Bond, P.L., Richardson, D.J., and Spiro, S. (2003). Characterization of a nitrate-

respiring bacterial community using the nitrate reductase gene (*narG*) as a functional marker.

Microbiology (Reading) *149*, 229-237.

Hakoyama, T., Niimi, K., Watanabe, H., Tabata, R., Matsubara, J., Sato, S., Nakamura, Y., Tabata,

S., Jichun, L., Matsumoto, T.*, et al.* (2009). Host plant genome overcomes the lack of a bacterial

gene for symbiotic nitrogen fixation. Nature *462*, 514-517.

Hawkins, J.P., Geddes, B.A., and Oresnik, I.J. (2017). Succinoglycan Production Contributes to

Acidic pH Tolerance in *Sinorhizobium* meliloti Rm1021. Mol Plant Microbe Interact *30*, 1009-

1019.

Janssen, S., McDonald, D., Gonzalez, A., Navas-Molina, J.A., Jiang, L., Xu, Z.Z., Winker, K.,

Kado, D.M., Orwoll, E., Manary, M.*, et al.* (2018). Phylogenetic Placement of Exact Amplicon

Sequences Improves Associations with Clinical Information. mSystems *3*,

10.1128/msystems.00021-00018.

Konstantinidis, K.T., and Tiedje, J.M. (2005). Genomic insights that advance the species

definition for prokaryotes. Proc Natl Acad Sci U S A *102*, 2567-2572.

Lerouge, P., Roche, P., Faucher, C., Maillet, F., Truchet, G., Prome, J.C., and Denarie, J. (1990).

Symbiotic host-specificity of *Rhizobium* meliloti is determined by a sulphated and acylated

glucosamine oligosaccharide signal. Nature *344*, 781-784.

Liu, A., Contador, C.A., Fan, K., and Lam, H.M. (2018). Interaction and Regulation of Carbon,

Nitrogen, and Phosphorus Metabolisms in Root Nodules of Legumes. Front Plant Sci *9*, 1860.

Miché, L., Moulin, L., Chaintreuil, C., Contreras-Jimenez, J.L., Munive-Hernández, J.A., Del

Carmen Villegas-Hernandez, M., Crozier, F., and Béna, G. (2010). Diversity analyses of

Aeschynomene symbionts in Tropical Africa and Central America reveal that *nod*-independent

stem nodulation is not restricted to photosynthetic *bradyrhizobia*. Environmental microbiology

12, 2152-2164.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,

and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Molecular Biology and Evolution *37*, 1530-1534.

Mornico, D., Miche, L., Bena, G., Nouwen, N., Vermeglio, A., Vallenet, D., Smith, A.A., Giraud,

E., Medigue, C., and Moulin, L. (2011). Comparative genomics of *aeschynomene* symbionts:

insights into the ecological lifestyle of *nod*-independent photosynthetic *bradyrhizobia*. Genes

(Basel) *3*, 35-61.

Nemecek-Marshall, M., MacDonald, R.C., Franzen, J.J., Wojciechowski, C.L., and Fall, R.

(1995). Methanol Emission from Leaves (Enzymatic Detection of Gas-Phase Methanol and

Relation of Methanol Fluxes to Stomatal Conductance and Leaf Development). Plant physiology

108, 1359-1368.

Nouwen, N., Arrighi, J.-F., Cartieaux, F., Chaintreuil, C., Gully, D., Klopp, C., and Giraud, E.

(2017). The role of rhizobial (*NifV*) and plant (FEN1) homocitrate synthases in

Aeschynomene/photosynthetic *Bradyrhizobium* symbiosis. Scientific Reports *7*, 448.

Ogier, J.-C., Pagès, S., Galan, M., Barret, M., and Gaudriault, S. (2019). *rpoB*, a promising

marker for analyzing the diversity of bacterial communities by amplicon sequencing. BMC

Microbiology *19*, 171.

Okubo, T., Fukushima, S., Itakura, M., Oshima, K., Longtonglang, A., Teaumroong, N., Mitsui,

H., Hattori, M., Hattori, R., Hattori, T.*, et al.* (2013). Genome analysis suggests that the soil

oligotrophic bacterium Agromonas oligotrophica (*Bradyrhizobium oligotrophicum*) is a nitrogen-

fixing symbiont of *Aeschynomene indica*. Appl Environ Microbiol *79*, 2542-2551.

Oldroyd, G.E.D. (2013). Speak, friend, and enter: signalling systems that promote beneficial

symbiotic associations in plants. Nature Reviews Microbiology *11*, 252-263.

Ormeno-Orrillo, E., and Martinez-Romero, E. (2019). A Genomotaxonomy View of the

Bradyrhizobium Genus. Front Microbiol *10*, 1334.

Parker, M.A. (2015). The spread of *Bradyrhizobium* lineages across host legume clades: from

Abarema to Zygia. Microb Ecol *69*, 630-640.

Piromyou, P., Greetatorn, T., Teamtisong, K., Okubo, T., Shinoda, R., Nuntakij, A., Tittabutr, P.,

Boonkerd, N., Minamisawa, K., and Teaumroong, N. (2015). Preferential association of

endophytic *bradyrhizobia* with different rice cultivars and its implications for rice endophyte

evolution. Appl Environ Microbiol *81*, 3049-3061.

Quelas, J.I., Mongiardini, E.J., Casabuono, A., Lopez-Garcia, S.L., Althabegoiti, M.J., Covelli,

J.M., Perez-Gimenez, J., Couto, A., and Lodeiro, A.R. (2010). Lack of galactose or galacturonic

acid in *Bradyrhizobium* japonicum USDA 110 exopolysaccharide leads to different symbiotic

responses in soybean. Mol Plant Microbe Interact *23*, 1592-1604.

Sprent, J.I., Ardley, J., and James, E.K. (2017). Biogeography of nodulated legumes and their

nitrogen-fixing symbionts. New Phytol *215*, 40-56.

Tao, J., Wang, S., Liao, T., and Luo, H. (2021). Evolutionary origin and ecological implication of

a unique *nif* island in free-living *Bradyrhizobium* lineages. ISME J *15*, 3195-3206.

Teulet, A., Camuel, A., Perret, X., and Giraud, E. (2022). The Versatile Roles of Type III

Secretion Systems in *Rhizobium*-Legume Symbioses. Annu Rev Microbiol *76*, 45-65.

Teulet, A., Gully, D., Rouy, Z., Camuel, A., Koebnik, R., Giraud, E., and Lassalle, F. (2020).

Phylogenetic distribution and evolutionary dynamics of nod and T3SS genes in the genus

Bradyrhizobium. Microbial genomics *6*.

Tighilt, L., Boulila, F., De Sousa, B.F.S., Giraud, E., Ruiz-Argueso, T., Palacios, J.M., Imperial,

J., and Rey, L. (2022). The *Bradyrhizobium* Sp. LmicA16 Type VI Secretion System Is Required

for Efficient Nodulation of Lupinus Spp. Microb Ecol *84*, 844-855.

Van Berkum, P., and Eardly, B.D. (2002). The aquatic budding bacterium Blastobacter

denitrificans is a nitrogen-fixing symbiont of *Aeschynomene indica*. Appl Environ Microbiol *68*,

1132-1136.

Vo, C.D., Michaud, J., Elsen, S., Faivre, B., Bouveret, E., Barras, F., Fontecave, M., Pierrel, F.,

Lombard, M., and Pelosi, L. (2020). The O(2)-independent pathway of ubiquinone biosynthesis

is essential for denitrification in *Pseudomonas* aeruginosa. J Biol Chem *295*, 9021-9032.

Wang, Y., Chen, X., Spengler, K., Terberger, K., Boehm, M., Appel, J., Barske, T., Timm, S.,

Battchikova, N., Hagemann, M.*, et al.* (2022). Pyruvate:ferredoxin oxidoreductase and low

abundant ferredoxins support aerobic photomixotrophic growth in cyanobacteria. Elife *11*,

e71339.

Figure legends

Figure 1. The relative abundance of each *Bradyrhizobium* supergroup (A) and each major clade

within the Photosynthetic supergroup of *Bradyrhizobium* (B). The relative abundance of each

Bradyrhizobium supergroup / PB clade was determined by dividing the number of reads assigned

to each supergroup / clade by the total number of filtered reads assigned to *Bradyrhizobium* / PB.

Abbreviations: *B. elkanii*, *B. jicamae* and *B. japonicum* represent *Bradyrhizobium elkanii*,

Bradyrhizobium jicamae and *Bradyrhizobium japonicum* supergroups, respectively.

Figure 2. The phylogenomic tree and population delineation of the "photosynthetic"

Bradyrhizobium (PB) supergroup. The phylogenomic tree was rooted using the minimum

variance (MV) method. The 209 PB strains sequenced in the present study were indicated by red

dots in the innermost layer surrounding the phylogeny. The ultrafast bootstrap values higher than

or equal to 95% calculated by IQ-Tree were labeled on the nodes with black circles. The

phylogenomic tree of the Photosynthetic *Bradyrhizobium* based on the minimal ancestor

deviation (MAD) rooting method was displayed in Fig. S3.

Figure 3. Symbiotic properties of several representative strains of the main populations

identified in the PB supergroup. (A) Comparison of growth of the *A. indica* plants (leaf

phenotype) non-inoculated (NI) or inoculated with different representative strains of PB. The six

MC1 representative strains are presented in the following order: SZCCHNR1021,

SZCCHNR1085, SZCCHNS30121, SZCCHNGR1005, SZCCHNRI2010 and SZCCHNF30212.

For the other MCs, only one representative strain is shown: SZCCHNRI2049 (MC2),

SZCCHNS30592 (MC3), SZCCHNRI3016 (MC4), SZCCHNS3002 (MC5) and SZCCHNS2021

(MC6), Photos were taken 17 days after inoculation. (B) Nodule number on *A. indica* plants

induced by different representative strains of PB at 17 days after inoculation. Box plots show the

results of one of the two experiments performed independently (4 plants each). (C) Symbiotic

phenotypes of some representative strains tested on *A. indica*. Column 1 and 2: photo of roots

and nodules. Scale bars: column 1, 0.5 cm; column 2, 0.2 cm. Column 3: Micro-sections of

nodules observed using light microscopy. Scale bars: 250 µm. (D) Confocal microscopy images

of micro-sections of nodules elicited by Clade 1 strains. The nodules formed by SZCCHNR1043

from MC3 and ORS278 are shown as examples. The central nodule tissue is intracellularly

infected with live spherical bacteroids stained green by SYTO9. (E) Confocal microscopy

images of micro-section of nodules elicited by Clade 2 strains (STM3843 and HKCCYLS1011).

The nodules elicited by STM3843 look normal but a mix of live and dead bacteria (stained red

by propidium iodide) can be noticed while the nodules formed by HKCCYLS1011 displayed a

central infected tissue that is digested. (F) Confocal microscopy images of micro-section of

nodules elicited by Clade 3 strains displaying different phenotypes: nodule containing mainly

dead bacteria (SZCCHNS30121), nodule with a completely digested central tissue

(SZCCHNRI2010); and nodule with a necrotic area and a digested zone (SZCCHNGR1005).

Scale bars for (D), (E) and (F): column 1, 200 µm; column 2, 10 µm.

Tree scale: 0.1

Legend from outer to inner

Outer

Functional gene cluster
- Methanol into formaldehyde
- Photosynthetic genes
- T3SS gene clusters
- *nodABC*

Plant
- *Oryza sativa* indica 1
- *Oryza sativa* indica 2
- *Oryza sativa* indica 3
- *Oryza sativa* japonica
- *Houttuynia cordata*
- *Camphora officinarum*

Niche
- Root
- Rhizosphere
- Soil

Sub_cluster
- Sub_cluster 1
- Sub_cluster 2
- Sub_cluster 3
- Sub_cluster 4
- Sub_cluster 5
- Sub_cluster 6

- strains used for symbiosis assay
- sequenced in this study

Background > Main cluster
- MC1
- MC2
- MC3
- MC4
- MC5
- MC6
- MC7
- MC8

Inner

Clade 1

Clade 2

Clade 3

Figure S1. *RpoB* gene tree constructed by full length (A) and amplified region (B). Strains from Xanthobacteraceae were used as an outgroup. The black circles on the nodes indicate ultrafast bootstrap values higher than or equal to 95% calculated by IQ-Tree. The 209 PB strains sequenced in the present study are indicated by red dots in the outermost layer surrounding the tree.

Figure S2. The maximum-likelihood phylogenomic tree of *Bradyrhizobium*. Strains from Xantho-bacteraceae were used as an outgroup. The tree was constructed using the 123 orthologous genes. identified in a previous study (Tao et al., 2021). The black circles on the nodes indicate ultrafast bootstrap values higher than or equal to 95% calculated by IQ-Tree. The 209 PB strains sequenced in the present study are indicated by red dots in the outermost layer surrounding the tree.

Figure S3. The phylogenomic tree of the photosynthetic *Bradyrhizobium* is based on the minimal ancestor deviation (MAD) rooting method. Solid circles in the phylogeny indicate nodes with IQ-Tree's ultrafast bootstrap values ≥ 95%.

Figure S4. The ASVs (amplicon sequence variants) and *rpoB* genes (amplified region from photosynthetic *Bradyrhizobium* genomes) tree. This gene tree was rooted by the minimum variance (MV) method. The 222 *rpoB* genes from PB genomes in this study were divided into each clade and main cluster (MC) according to Fig. 2. The ultrafast bootstrap values higher than or equal to 95% calculated by IQ-Tree were labeled on the nodes with black circles.

Figure S5. The phyletic pattern of specific genes between three PB clades. The solid circles and blank in the right panel represent the presence and absence of the genes, respectively. The genome tree in the left panel was displayed in rectangular mode. Detailed information of specific genes was shown in Dataset S5. Histidine metabolism *hutCFHIU*, signaling and cellular processes *mdoGH*, raffinose/stachyose/melibiose transport system *msmEFG*, multidrug efflux system *triABC*, styrene and butanoate metabolism *gctAB*, glycerolipid metabolism *dhaKLM*, O₂-independent ubiquinone biosynthesis *ubiXTUVD*, nitrogen metabolism (denitrification) *narGHIJ napABCDE*, photosynthetic genes 1) porphyrin metabolism *bchCFMOXYZJ* and *chlBHILNPG* 2) carotenoid biosynthesis *crtCDIF* 3) light-harvesting complex *pufAB* 4) photosynthetic reaction center *pufML* and *puhA*, C4-dicarboxylate transport *dctBD*, adhesin transport system *lapBCE*, Type III secretion system, pyruvate metabolism *porABC*, multicomponent Na⁺:H⁺ antiporter *mnhBD*, glucose/mannose transport system *gtsBC*, formate dehydrogenase *fdoHI*, mannopine transport system *attA1A2BC*, Type VI secretion system, tungstate transport system *tupABC*, UDP-glucose/iron transport system *fetAB*, malonate decarboxylase *mdcABCDEG*, C1 metabolism *mgsABC mxaFI* and *mxaACGKL*, succinoglycan biosynthesis *exoALOUWY*, two-component system *adeRS*.

Figure S6. Growth of representative strains from the populations delineated by PopCOGenT for the PB of *Bradyrhizobium* on methanol as a sole carbon source, with the presence of lanthanide (Ln) species (Ce^{2+}, 30 uM). The reference strain *Bradyrhizobium diazoefficiens* USDA110 was used as a positive control. Error bars indicate the standard deviation of the mean from three replicates.

Figure S7. The heatmap of the pairwise identity of 16S rRNA genes and the whole-genome average nucleotide identity (ANI) of all PB members.

Figure S8. Comparison of the genomic context of the *nif* gene cluster (*nif* island) (A) and T3SS gene cluster (B) in the representative strains of PB of *Bradyrhizobium*. Gene functions are distinguished by different colors. The visualization of gene arrangement is performed with DNA-features-viewer v3.0.3 (Zulkower and Rosser, 2020). (C) The phylogenetic tree of the *rhcN* protein from *Bradyrhizobium*. The *rhcN* families were defined according to Teulet et al. (2020). The gene tree was rooted using the minimum variance (MV) method. The different colored branches correspond to the distinct genetic organization of the T3SS clusters to which the *rhcN* gene belongs. The *rhcN* in the strain *Bradyrhizobium* sp. 36 is not shown as it belongs to a different type of T3SS, likely a result of HGT from distantly related bacteria according to Teulet et al. (2020). Black circles in the phylogeny indicate nodes with IQ-Tree's ultrafast bootstrap values ≥ 95%.

Figure S9. Ability of several representative strains of the main populations identified in PB supergroup to fix nitrogen during their free-living and symbiotic states. (A) Free-living nitrogen fixation after 8 days of culture in vacutainer tube. (B) Nitrogen fixation of *A. indica* plants inoculated with different PB representative strains at 17 days post-inoculation. In (A) and (B), ORS278 is used as a positive control and USDA110 (a member of *B. japonicum* supergroup) is used as a negative control in (A). NI: non inoculated. MC: Main Cluster

A

B

Figure S10. Complementary data of Fig. 2 showing symbiotic properties of several representative strains of the main populations identified in PB supergroup. (A) Comparison of the growth of the *A. india* plant (leaf phenotype) non-inoculated (NI) or inoculated with different representative strains of PB. All representative strains tested from each MC are present in this order : MC1 - SZCCHNR1021; SZCCHNR1085; SZCCHNS30121; SZCCHNGR1005; SZCCHNRI2010; SZCCHNF30212; MC2 - SZCCHNG3015; SZCCHNRI2049; SZCCHNS30581; SZCCHNS1012; SZCCHNR1019; MC3 - SZCCHNR1043; SZCCHNR1053; SZCCHNS30592; SZCCHNRI3043; MC4 - SZCCHNRI3016; SZCCHNS3053; SZCCHNS3016; MC5 - SZCCHNR3003; SZCCHNRI1009; SZCCHNR1015; SZCCHNS3002 and MC6 - SZCCHNS2021; SZCCHNS2022; SZCCHNR1002. ORS278 is used as control. (B) Confocal microscopy images of micro-section of nodules elicited by the other Clade 1 strains tested after staining with SYTO9 (green, live bacteria), propidium iodide (red, infected plant nuclei and dead bacteria or bacteria with compromised membranes) and calcofluor (blue, plant cell wall). Scale bars: column 1, 200 µm; column 2, 10 µm.

Figure S11. Gel electrophoresis image of DNA from PB strains amplified with the specific *rpoB* primer set BR2106F/BR2516R.

Supplementary Information

Correlating phylogenetic and functional diversity of the *nod*-free but nodulating

***Bradyrhizobium* phylogroup**

Lu Ling[^], Alicia Camuel[^], Sishuo Wang, Xiaojun Wang, Tianhua Liao, Jinjin Tao, Xingqin LIN,

Nico Nouwen, Eric Giraud[*], Haiwei Luo[*]

[^]**These are co-first authors**: Lu Ling, Alicia Camuel

[*]**These are co-corresponding authors:** Eric Giraud (eric.giraud@ird.fr), Haiwei Luo

(haiweiluo@cuhk.edu.hk)

This PDF file includes:

14   **Supplementary Text.** Methods

15   **Supplementary references.**

16   **Figures S1 to S11.**

Supplementary Text: Methods

Sample niche classification

Excess soil was vigorously shaken away from the plant roots to remove loose soil, leaving

approximately 1 mm of soil attached to the roots, which constitutes the rhizosphere compartment

(Edwards et al., 2015). To separate the 1 mm of rhizosphere soil directly from the roots, roots

were cut into 5 cm lengths and two grams were placed in a sterile flask containing 50 mL of

sterile phosphate buffered saline (PBS) solution and vortexed several times (15 seconds). The

PBS solution used to wash the roots was transferred to a 50 mL Falcon tube and then centrifuged

at 10,000 x g for one minute. The supernatant was discarded, and the remaining soil samples

were saved as rhizosphere samples (Edwards et al., 2015). The roots were then resuspended and

vortexed three times (15 seconds each) before being transferred to new Falcon tubes as root

samples. Soil away from the root was stored as bulk soil.

Bacterial isolation

The root, rhizosphere, and bulk soil samples were placed in the incubation chamber for three

to four days at 28°C and 70% relative humidity under the day/night (16/8 h) cycle. All root

samples were washed with sterile Milli-Q water to remove any adhering soil, then subjected to

the surface sterilization by immersion in 75% ethanol for three minutes, followed by 5% sodium

hypochlorite for five minutes, and rinsed five times with sterile distilled water (Mbai et al., 2013).

The washed roots were then aseptically dried on the clean bench before being ground in a

sterilized mortar and pestle with PBS buffer (pH 6.8). A series of fivefold and tenfold dilutions

were prepared and then inoculated (100 µL) on modified arabinose-gluconate (MAG) media

without any nitrogen source (1.0 g DL-arabinose, 1.0 g sodium gluconate, 1.0 g yeast extract, 2

mL KH$_2$PO$_4$ solution (110 g/L), 4 mL Na$_2$SO$_4$ solution (62.5 g/L), 1 mL MgSO$_4$•7H$_2$O solution

(180 g/L), 1 mL CaCl$_2$ solution (13 g/L), 1 mL FeCl$_3$•6H$_2$O solution (6.7 g/L) and 15 g agar were

added. The medium was made up to 1 L with Milli-Q water, the pH was adjusted to 6.6 with

KOH, and then autoclaved at 121 ℃ for 15 - 30 minutes.) (Tao et al., 2021). To prepare soil

inoculations, 5.0 g of fresh soil was mixed with 45 mL of sterile deionized water in a 50 mL

Falcon tube. One mL of soil suspension was serially diluted 10-fold after mixing with a vortex

mixer, and 100 μL of diluted samples were spread on the MAG media (Tao et al., 2021). The

isolation plates were then placed in a 28 ℃ incubator to allow bacteria to grow. After one week,

colonies with specific morphology (small colonies with white or pink color) were picked and

purified on fresh MAG media with the addition of 2 mL NH$_4$Cl solution (160 g/L). Purified

strains were preserved at -80 ℃ in glycerol suspensions (30% v/v).

To identify the taxonomy of each strain, colony polymerase chain reaction (PCR) was

performed. Briefly, bacterial colonies were mixed with the 10% (w/v) Chelex solution [Bio-Rad,

USA] and incubated at 90 ℃ for 20 minutes to release DNA from bacterial colonies as templates.

The universal bacteria 16S rRNA primers 27F (5′-AGRGTTYGATYMTGGCTCAG-3′) and

1492R (5′-GGYTACCTTGTTACGACTT-3′), Premix Taq [Takara Bio, USA] and RNA/DNAse-

free water were used in the PCR. The PCR condition included an initial denaturation at 95 ℃ for

five minutes, followed by 35 cycles of amplification (95 ℃ for 45 s, 55 ℃ for 45 s and 72 ℃ for

90 s) and a final extension at 72 ℃ for 10 minutes (Tao et al., 2021). PCR products were run on

0.8% agarose and viewed by the automatic analysis system of electrophoresis gel imaging.

Positive PCR products were sequenced in BGI Genomics. The taxonomic information of each

isolate was analyzed by comparing the 16S rRNA gene sequences with *Bradyrhizobium* in

EzBioCloud (Yoon et al., 2017).

Primer design for detecting *Bradyrhizobium* diversity

It is well known that the closely related genera are difficult to be distinguished by the

traditional 16S rRNA gene sequences due to their slow evolutionary rate (Vos et al., 2012). In

contrast, the *rpoB* gene has a higher resolution and can be used as a marker gene to distinguish

closely related species (Vos et al., 2012). Therefore, we designed specific *Bradyrhizobium rpoB*

primers for amplicon sequencing to investigate the relative abundance and diversity of

Bradyrhizobium supergroups in different samples. The primer sequences were BR2106F

(CCGRTSACGCCBGACAAG) and BR2516R (TGTCGCCCTTCYTGACGAYR), producing a

~410 bp sequence. The primers were validated to amplify various *Bradyrhizobium* members as

well as strains within the PB supergroup with specificity (Fig. S11). PCR conditions included

denaturation at 95 °C for three minutes, followed by 35 cycles of 95 °C for 30 seconds, 61 °C for

45 seconds and 72 °C for 45 seconds, with a final extension at 72 °C for 10 minutes.

Amplicon sequence analysis

The quality and quantity of the DNA extracted from the soil and root samples were assessed

using a NanoDropTM 2000 [Thermo Fisher, USA]. The amplicon sequence variants (ASVs)

obtained from the *rpoB* amplicon sequences after denoising were subsequently filtered out the

non-*rpoB* amplicon by conducting BLASTN searches against the NCBI nr database. The aligned

ASVs with an e-value lower than 1e-10 but without alignment with 'DNA-directed RNA

polymerase subunit beta' in the nr database were removed.The reference *rpoB* sequence was

extracted from the genomic sequence and then aligned. The alignment was trimmed to retain the

amplified region and used as the reference alignment. The ASVs were then aligned with the

reference alignment using PaPaRa v2.5 (Berger and Stamatakis, 2012). The aligned ASVs were

subsequently placed on the phylogenetic tree (Fig. S1B) using EPA-ng v0.3.8 (Barbera et al.,

2019) and assigned to seven *Bradyrhizobium* supergroups, including Soil 1, Soil 2,

Bradyrhizobium elkanii, *Bradyrhizobium jicamae*, Kakadu, Photosynthetic and *Bradyrhizobium*

japonicum supergroups. The relative abundance of each *Bradyrhizobium* supergroup was

determined by dividing the number of reads assigned to each *Bradyrhizobium* supergroup by the

total number of filtered reads assigned to *Bradyrhizobium* (Fig. 1a, Dataset S1). Similarly, the

relative abundance of each clade of Photosynthetic *Bradyrhizobium* (PB) was calculated by

dividing the number of reads assigned to each PB clade by the total number of filtered reads

assigned to PB (Fig. 1b, Fig. S3, Dataset S2).

<u>Basic soil characteristics</u>

Soil samples were separated to measure the basic soil characteristics after all the samples

were brought back to the laboratory. Soil water content was determined by drying fresh soil

(10.00 g) at 105°C for 6 hours. Soil pH was measured at the ratio of 1:5 (w/w) of soil-to-

deionized water using a pH electrode. The potassium dichromate oxidation process combined

with the heating method was used to measure the soil organic carbon (SOC). Total nitrogen (TN)

was determined by the kjeldah method (Bao, 2000). Total phosphorus (TP) was identified based

on the sodium hydroxide melting-molybdenum antimony colorimetric method (Bao, 2000).

Available phosphorus (AP) was ascertained using the molybdenum antimony anti-colorimetric

method (Bao, 2000). Soil nitrite nitrogen (NO_2^-), nitrate nitrogen (NO_3^-) and ammonium nitrogen

(NH_4^+) were extracted with KCl solution (2 mol L^{-1}) and detected by continuous flow analysis

(Tian et al., 2014). Dissolved organic carbon and nitrogen (DOC and DON) were analyzed using

a TOC/TN analyzer (Shimadzu, Analytical Sciences, Kyoto, Japan). Microbial biomass carbon

and nitrogen (MBC and MBN) were determined by chloroform fumigation extraction

(Durenkamp et al., 2010; Wu et al., 1990). Fe and Mo contents were measured by using

inductively coupled plasma mass spectrometry (ICP-MS). These sample-associated metadata

were provided in Dataset S3.

Genome sequencing, assembly and annotation

Bacteria Genomic DNA Extraction Kit [OMEGA Bacterial DNA Kit D3350] was used to

extract genomic DNA from each of the 209 isolates. NanoDrop^{TM} 2000 [Thermo Fisher, USA]

was used to assess the quality of the extracted DNA samples with the following criteria,

A260/A280 > 1.8, A260/A230 > 2.0 and A260/A270 > 1.0. Whole genome sequencing was

performed in Wuhan Huada Gene Biotechnology Company by using the MGISEQ-2000

PE150+150+10+10 platform (paired-end reads of 150 bp). The BBMerge method in the BBmap

package v38.79 was used to identify the untrimmed adapters associated with the raw reads

(Bushnell et al., 2017). Subsequently, Trimmomatic v0.39 was used to trim adapters and low-

quality reads (Bolger et al., 2014). Reads with less than 40 bp in length were discarded, and the

quality of the remaining reads was assessed by FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Contigs were assembled using

SPAdes v3.10.1 based on the remaining high-quality paired-end reads with default parameters

(Bankevich et al., 2012). Those contigs with lengths greater than 1,000 bp and a k-mer coverage

greater than five were kept for further analysis. The quality of the genome assemblies (Dataset

S8) was assessed by CheckM v1.0.7 (Parks et al., 2015). In general, 68 newly sequenced strains

were clustered with 12 publicly available strains as Clade 1, strains STM3843 and the newly

sequenced HKCCYLS1011 were clustered as Clade 2, and the remaining 140 newly sequenced

strains were clustered as Clade 3. Genomes from Clade 3 exhibited a significantly lower GC

content ($63.67 \pm 0.07\%$ vs. $65.39 \pm 0.22\%$, $p < 0.001$, phylogenetic ANOVA) and a similar

genome size (7.45 ± 0.14 Mb vs. 7.56 ± 0.33 Mb, $p = 0.746$, phylogenetic ANOVA) compared to

Clade 1 (Datasets S2 and S7).

We noted an assembly error related to the *nif* island that was supposed to harbor two *nifH*

genes. This was due to the presence of two *nifH* gene copies with nearly identical sequences,

which can lead to mis-assembly in short-read sequencing technologies such as Illumina or

MGISEQ-2000 (Tørresen et al., 2019). To address this issue, we mapped all raw reads to the

assembled genome and observed that the sequencing depth of the *nifH* gene was twice that of

other nearby genes. We further conducted Nanopore sequencing on three phylogenetically

distantly related PB strains (SZCCHNR3119, SZCCHNS2021 and SZCCHNS1050), and had

their genomes assembled using Flye v2.6 (Kolmogorov et al., 2019). Analysis of these genomes

confirmed the presence of two *nifH* genes separated by approximately 43 genes in these strains.

To correct this error in the remaining PB strains, we re-assembled their genomes using the

reference-guided method (Cabuk and Unlu, 2022) in Spades v3.10.1 with the parameter of "--

untrusted-contigs" (Bankevich et al., 2012). For each assembled genome, genes were predicted

using Prokka v1.12 (Seemann, 2014) and annotated using the Cluster of Orthologous Genes

(COG) database (Galperin et al., 2020) and the Kyoto Encyclopedia of Genes and Genomes

(KEGG) database (Kanehisa et al., 2023; Kanehisa et al., 2004) .

<u>Phylogenomic tree construction for all available *Bradyrhizobium* lineages</u>

A phylogenomic tree for all available *Bradyrhizobium* lineages (209 new genomes and 566

public genomes downloaded from the NCBI Genbank database) was constructed by IQ-TREE

v2.2.0 (Minh et al., 2020). The parameter "-s alignment -spp partition -m MFP -mset

LG,WAG,JTT -mrate E,G,I,G+I -bb 1000" was applied so that each gene was allowed to have its

own best-fit substitution model, automatically selected by the ModelFinder implemented in IQ-

TREE v2.2.0 (Minh et al., 2020). The branch support was assessed by 1000 ultrafast bootstrap

approximations (Hoang et al., 2017).

Phylogenomic and comparative genomic analyses for Photosynthetic *Bradyrhizobium*

OrthoFinder v2.3.4 (Emms and Kelly, 2019) was used to identify orthologous gene families

for all Photosynthetic *Bradyrhizobium* strains. Each of the identified 3,300 single-copy ortholog

families was aligned at the amino acid level using MAFFT v7.487 (Katoh and Standley, 2013),

and trimmed using trimAl v1.4.rev15 with the parameters "-automated1 -resoverlap 0.55 -

seqoverlap 60" (Capella-Gutiérrez et al., 2009). A maximum likelihood phylogenomic tree of the

PB supergroup was built based on the concatenated alignment of 3,300 single-copy orthologs

using IQ-TREE v2.2.0 with 1,000 ultrafast bootstrap replicates (Hoang et al., 2017). The best-fit

evolutionary model for each ortholog was determined by ModelFinder implemented in IQ-TREE

v2.2.0 (Minh et al., 2020).

As shown in Fig. S2, the Kakadu supergroup was considered to be a sister lineage to the

Photosynthetic supergroup. However, the long branches connecting these two supergroups

suggested that using the Kakadu supergroup as an outgroup to root the species tree of the

Photosynthetic supergroup may not be appropriate. Therefore, the Photosynthetic supergroup

phylogenomic tree was rooted using outgroup-independent rooting methods, specifically the

minimum variance (MV) (Mai et al., 2017) and the minimal ancestor deviation (MAD) (Tria et

al., 2017) methods. The MV method identifies the root with the minimum variance of root-to-tip

distances, while the MAD method considers all branches as plausible root positions, calculates

the relative deviation from the clock-likeness for each candidate, and determines the root with

the minimal mean relative deviation from the molecular clock interpretation of all branches (Tria

et al., 2017). The trees based on these two outgroup-independent methods indicated the same

root position (Fig. 2 and Fig. S3). To compare continuous traits of strains while incorporating the

evolution of those traits, a phylogenetic ANOVA (Garland et al., 1993) was conducted using the

'phylANOVA' function with 1,000 simulations implemented in the R package 'phytools' (Revell,

2023).

The similarity of the PB strains was measured at both 16S rRNA gene and whole-genome

levels. For the former, pairwise 16S rRNA gene identity was calculated using BLAST (Boratyn

et al., 2013), and strains were clustered using the complete linkage method based on sequence

identity. For the latter, the whole-genome average nucleotide identity was estimated by FastANI

v1.2 (Jain et al., 2018).

<u>Symbiotic analysis on *Aeschynomene indica*</u>

The 28 strains used for symbiotic analysis were grown in AG medium (Sadowsky et al.,

1987) at 28°C on Petri dishes for one week. Bacteria were harvested from the plate and

resuspended in 10 mL of sterile water and the OD_{600} was adjusted to 1.0. *A. indica* plants were

cultured as previously described (Okazaki et al., 2016). Each strain was inoculated into four

plants with 1 mL of bacterial suspension and the symbiotic properties (number of nodules per

plant and nitrogenase enzyme activity) were analyzed at 17 days after inoculation (Bonaldi et al.,

2010). Cytological analysis of nodules elicited by one strain representative of the populations

(MC2 to MC6), two strains for the Clade 2 and three strains for the MC1 population was

performed as described (Songwattana et al., 2021). Each strain was assessed in duplicate.

<u>In vitro nitrogenase enzyme activity</u>

Bacteria were grown in 9 mL vacuette® tubes (Greiner Bio-One GmbH) containing 2 mL of

0.8% agar BNM-B medium with 10 mM succinate and 10 mM arabinose as carbon source at

28 °C (Nouwen et al., 2017). At the beginning of the experiment, 10% acetylene was added to

the vacuette® tubes and after eight days of incubation, the amount of ethylene produced by the

bacterial culture was measured by gas chromatography (Giraud et al., 2000).

<u>Methanol metabolism</u>

We selected 17 strains covering the major populations predicted by PopCOGenT. The

reference strain *Bradyrhizobium* diazoefficiens USDA110 was used as a positive control. Each

strain was inoculated into 14 mL test tubes containing 5 mL minimal salts medium,

supplemented with 30 μM $CeCl_2$ (lanthanide chlorides) and 0.5% methanol as the sole carbon

source (Wang et al., 2019) and incubated at 28 °C with reciprocal shaking at 280 rpm. Growth

was monitored spectrophotometrically by measuring the optical density at 600 nm (OD_{600}).

Supplementary references

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,

Nikolenko, S.I., Pham, S., Prjibelski, A.D.*, et al.* (2012). SPAdes: a new genome assembly

algorithm and its applications to single-cell sequencing. J Comput Biol *19*, 455-477.

Bao, S. (2000). Soil agrochemical analysis (China agriculture press Beijing).

Barbera, P., Kozlov, A.M., Czech, L., Morel, B., Darriba, D., Flouri, T., and Stamatakis, A.

(2019). EPA-ng: Massively Parallel Evolutionary Placement of Genetic Sequences. Syst Biol *68*,

365-369.

Berger, S.A., and Stamatakis, A. (2012). PaPaRa 2.0: a vectorized algorithm for probabilistic

phylogeny-aware alignment extension. Heidelberg Institute for Theoretical Studies *12*.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics *30*, 2114-2120.

Bonaldi, K., Gourion, B., Fardoux, J., Hannibal, L., Cartieaux, F., Boursot, M., Vallenet, D.,

Chaintreuil, C., Prin, Y., Nouwen, N.*, et al.* (2010). Large-scale transposon mutagenesis of

photosynthetic *Bradyrhizobium* sp. strain ORS278 reveals new genetic loci putatively important

for *nod*-independent symbiosis with *Aeschynomene indica*. Mol Plant Microbe Interact *23*, 760-

770.

Boratyn, G.M., Camacho, C., Cooper, P.S., Coulouris, G., Fong, A., Ma, N., Madden, T.L.,

Matten, W.T., McGinnis, S.D., Merezhuk, Y.*, et al.* (2013). BLAST: a more efficient report with

usability improvements. Nucleic Acids Res *41*, W29-33.

Bushnell, B., Rood, J., and Singer, E. (2017). BBMerge–accurate paired shotgun read merging

via overlap. PLOS ONE *12*, e0185056.

Cabuk, U., and Unlu, E.S. (2022). A combined de novo assembly approach increases the quality

of prokaryotic draft genomes. Folia Microbiol (Praha) *67*, 801-810.

Capella-Gutiérrez, S., Silla-Martínez, J.M., and Gabaldón, T. (2009). trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics *25*, 1972-

1973.

Durenkamp, M., Luo, Y., and Brookes, P.C. (2010). Impact of black carbon addition to soil on

the determination of soil microbial biomass by fumigation extraction. Soil Biol Biochem *42*,

2026-2029.

Edwards, J., Johnson, C., Santos-Medellin, C., Lurie, E., Podishetty, N.K., Bhatnagar, S., Eisen,

J.A., and Sundaresan, V. (2015). Structure, variation, and assembly of the root-associated

microbiomes of rice. Proc Natl Acad Sci U S A *112*, E911-920.

Emms, D.M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for

comparative genomics. Genome Biol *20*, 238.

Galperin, M.Y., Wolf, Y.I., Makarova, K.S., Vera Alvarez, R., Landsman, D., and Koonin, E.V.

(2020). COG database update: focus on microbial diversity, model organisms, and widespread

pathogens. Nucleic Acids Research *49*, D274-D281.

Garland, T., Dickerman, A.W., Janis, C.M., and Jones, J.A. (1993). Phylogenetic Analysis of

Covariance by Computer-Simulation. Systematic Biology *42*, 265-292.

Giraud, E., Hannibal, L., Fardoux, J., Vermeglio, A., and Dreyfus, B. (2000). Effect of

Bradyrhizobium photosynthesis on stem nodulation of *Aeschynomene sensitiva*. Proc Natl Acad

Sci U S A *97*, 14795-14800.

Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., and Vinh, L.S. (2017). UFBoot2:

Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution *35*, 518-522.

Jain, C., Rodriguez, R.L., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S. (2018). High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat

Commun *9*, 5114.

Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M., and Ishiguro-Watanabe, M. (2023).

KEGG for taxonomy-based analysis of pathways and genomes. Nucleic Acids Res *51*, D587-

D592.

Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y., and Hattori, M. (2004). The KEGG resource

for deciphering the genome. Nucleic Acids Res *32*, D277-280.

Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol *30*, 772-780.

Kolmogorov, M., Yuan, J., Lin, Y., and Pevzner, P.A. (2019). Assembly of long, error-prone reads

using repeat graphs. Nat Biotechnol *37*, 540-546.

Mai, U., Sayyari, E., and Mirarab, S. (2017). Minimum variance rooting of phylogenetic trees

and implications for species tree reconstruction. PLOS ONE *12*, e0182238.

Mbai, F., Magiri, E., Matiru, V., Nganga, J., and Nyambati, V. (2013). Isolation and

characterization of bacterial root endophytes with potential to enhance plant growth from

Kenyan Basmati rice. American International Journal of Contemporary Research.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,

and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Molecular Biology and Evolution *37*, 1530-1534.

Nouwen, N., Arrighi, J.-F., Cartieaux, F., Chaintreuil, C., Gully, D., Klopp, C., and Giraud, E.

(2017). The role of rhizobial (*NifV*) and plant (FEN1) homocitrate synthases in

Aeschynomene/photosynthetic *Bradyrhizobium* symbiosis. Scientific Reports *7*, 448.

Okazaki, S., Tittabutr, P., Teulet, A., Thouin, J., Fardoux, J., Chaintreuil, C., Gully, D., Arrighi,

J.F., Furuta, N., Miwa, H.*, et al.* (2016). *Rhizobium*-legume symbiosis in the absence of *Nod*

factors: two possible scenarios with or without the T3SS. ISME J *10*, 64-74.

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., and Tyson, G.W. (2015). CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res *25*, 1043-1055.

Revell, L.J. (2023). phytools 2.0: An updated R ecosystem for phylogenetic comparative

methods (and other things). *In press*, 2023.2003.2008.531791.

Sadowsky, M.J., Tully, R.E., Cregan, P.B., and Keyser, H.H. (1987). Genetic Diversity in

Bradyrhizobium japonicum Serogroup 123 and Its Relation to Genotype-Specific Nodulation of

Soybean. Appl Environ Microbiol *53*, 2624-2630.

Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. Bioinformatics *30*, 2068-

2069.

Songwattana, P., Chaintreuil, C., Wongdee, J., Teulet, A., Mbaye, M., Piromyou, P., Gully, D.,

Fardoux, J., Zoumman, A.M.A., Camuel, A.*, et al.* (2021). Identification of type III effectors

modulating the symbiotic properties of *Bradyrhizobium* vignae strain ORS3257 with various

Vigna species. Sci Rep *11*, 4874.

Tao, J., Wang, S., Liao, T., and Luo, H. (2021). Evolutionary origin and ecological implication of

a unique *nif* island in free-living *Bradyrhizobium* lineages. ISME J *15*, 3195-3206.

Tian, X.F., Hu, H.W., Ding, Q., Song, M.H., Xu, X.L., Zheng, Y., and Guo, L.D. (2014).

Influence of nitrogen fertilization on soil ammonia oxidizer and denitrifier abundance, microbial

biomass, and enzyme activities in an alpine meadow. Biology and Fertility of Soils *50*, 703-713.

Tørresen, O.K., Star, B., Mier, P., Andrade-Navarro, M.A., Bateman, A., Jarnot, P., Gruca, A.,

Grynberg, M., Kajava, A.V., Promponas, V.J.*, et al.* (2019). Tandem repeats lead to sequence

assembly errors and impose multi-level challenges for genome and protein databases. Nucleic

Acids Research *47*, 10994-11006.

Tria, F.D.K., Landan, G., and Dagan, T. (2017). Phylogenetic rooting using minimal ancestor

deviation. Nat Ecol Evol *1*, 193.

Vos, M., Quince, C., Pijl, A.S., de Hollander, M., and Kowalchuk, G.A. (2012). A comparison of

rpoB and 16S rRNA as markers in pyrosequencing studies of bacterial diversity. PLOS ONE *7*,

e30600.

Wang, L., Suganuma, S., Hibino, A., Mitsui, R., Tani, A., Matsumoto, T., Ebihara, A., Fitriyanto,

N.A., Pertiwiningrum, A., Shimada, M.*, et al.* (2019). Lanthanide-dependent methanol

dehydrogenase from the legume symbiotic nitrogen-fixing bacterium *Bradyrhizobium*

diazoefficiens strain USDA110. Enzyme Microb Technol *130*, 109371.

Wu, J., Joergensen, R.G., Pommerening, B., Chaussod, R., and Brookes, P.C. (1990).

Measurement of Soil Microbial Biomass C by Fumigation Extraction - an Automated Procedure.

Soil Biol Biochem *22*, 1167-1169.

Yoon, S.H., Ha, S.M., Kwon, S., Lim, J., Kim, Y., Seo, H., and Chun, J. (2017). Introducing

EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome

assemblies. Int J Syst Evol Microbiol *67*, 1613-1617.

331