Phylogenetic and population structure of the *nod*-free but nodulating *Bradyrhizobium* phylogroup

Lu Ling[1,2,^], Alicia Camuel[3,4,^], Sishuo Wang[1], Xiaojun Wang[2], Xinqin Lin[2], Nico Nouwen[3,4], Eric Giraud[3,4,*], Haiwei Luo[1,5,*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

[2]Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, China

[3]IRD, Laboratoire des Symbioses Tropicales et Méditerranéennes (LSTM), UMR IRD/ SupAgro/INRA/ Université de Montpellier /CIRAD, TA-A82/J– Campus de Baillarguet 34398 Montpellier cedex 5, France

[4]PHIM Plant Health Institute, Université Montpellier, IRD, CIRAD, INRAE, Institut Agro, Montpellier, France

[5]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China

[^]**These are co-first authors**: Lu Ling, Alicia Camuel

[*]**These are co-corresponding authors:** Eric Giraud (eric.giraud@ird.fr), Haiwei Luo (haiweiluo@cuhk.edu.hk)

Key words: *Bradyrhizobium*, *Aeschynomene*, nodulation, population genomics

Abstract

Bradyrhizobium is a main rhizobial lineage of which the majority of its members nodulate legume plants using Nod factors (NFs) encoded by the *nod* genes. However, the "photosynthetic" phylogroup within *Bradyrhizobium* (PB) does not contain *nod* genes but are still capable of establishing a nitrogen-fixing symbiosis with some tropical legumes of the *Aeschynomene* genus. Furthermore, members of this phylogroup colonize rice roots and promote plant growth. Despite this unique combination of ecological features, genomes of only 13 PB members are sequenced, and almost all are isolated from *Aeschynomene* spp. nodules. Here, we sequenced 208 new PB members predominantly associated with rice. The extended PB phylogroup comprises three main clades: a basal clade with significant expansion of its diversity, followed by the diversification of a singleton clade from a new clade exclusively represented by our strains. Although the PB strains universally lack the canonical *nod* genes, all 27 assayed strains representing the broad diversity of these clades induced nodules on *Aeschynomene indica*. Interestingly, significant differences in the efficiency of symbiosis between the main clades were observed and the singleton clade showed an intermediate symbiotic phenotype aligning well with its intermediate phylogenetic position. Our strain collection expands the phylogenetic and ecological diversity of the *nod*-free but nodulating *Bradyrhizobium* and shows that the NF-independent nodulation of *Aeschynomene* is a common trait of this phylogroup, in contrast to the photosynthetic trait originally thought as its unifying feature.

Introduction

Bradyrhizobium is one of the largest and most diverse rhizobial genera and the primary

symbiont of a wide range of legumes [1, 2]. The genus *Bradyrhizobium* encompasses seven

phylogenetic supergroups (phylogroups) [3, 4], among which the "photosynthetic"

Bradyrhizobium (PB) supergroup is one of the most special. First, these bacteria are

photosynthetic which is a very rare property in rhizobia [3]. Second, members belonging to this

supergroup can form nitrogen-fixing nodules not only on roots but also on stems of some tropical

semi-aquatic species of the *Aeschynomene* genus [5]. Finally, the most original feature of the PB

strains is the fact that these nodulating bacteria lack the canonical *nodABC* genes encoding the

enzymes that synthesize the core structure of the nodulation factors (NF).

The available evidence indicates that the PB phylogroup takes a novel symbiotic strategy

that differs from the traditional universal NF-based pathway [6, 7]. Although several attempts

have been made to identify the symbiotic determinants of PB strains by mutagenesis, the

mechanism has not yet been discovered [8]. This mystery has become more intriguing since the

recent discovery of another NF-independent symbiotic process present in a wide variety of non-

photosynthetic *Bradyrhizobium* strains [9]. This last process relies on a Type III secretion system

(T3SS) which secretes, into the plant cells, effectors (T3Es) able of triggering nodulation [10-

12]. The absence of T3SS in most PB strains indicates that they use a distinct symbiotic

mechanism that was shown specific to this phylogroup [12].

Despite these unique features, the research on photosynthetic *Bradyrhizobium* has been

very limited. Currently, there are only 13 PB strains that have their genomes sequenced, largely

limiting comparative genomics analysis on this important *Bradyrhizobium* phylogroup. Although

most PB strains were isolated from *Aeschynomene* spp. nodules, the PB phylogroup was also

detected in paddy soil, rice roots as well as in lake water [13-16], suggesting that its members

have much wider ecological niches than previously thought. Here, we cultured and sequenced

the genomes of 208 isolates (Fig. 1) predominantly associated with three niches of rice (within

root, rhizosphere, bulk soil). We then conducted genomic analyses and physiological assays to

reveal the phylogenetic and population structure of the photosynthetic *Bradyrhizobium*

supergroup.

Results and discussion

<u>The new genomes expand the phylogenetic diversity of the photosynthetic supergroup of</u>

<u>*Bradyrhizobium*</u>

With the 208 newly isolated members (see Supplementary Text for photosynthetic

Bradyrhizobium isolation), the phylogeny of the PB supergroup is split into three deep-branching

clades (see Fig. S1 for their phylogenetic position in the species tree of the entire

Bradyrhizobium genus). Clade 1 represents the evolutionarily basal lineage. It initially comprised

only 12 strains, but it now has its diversity expanded by having 68 new strains representing

several new lineages delineated as distinct genetically isolated populations (Fig. 1, Datasets S1

and S2; also discussed in the next section). Clade 2 is a singleton clade with only one strain

(*Bradyrhizobium* sp. STM3843) available and represents a phylogenetic intermediate among the

three clades. Clade 3 is new and consists exclusively of 140 new strains. The evolutionary

branching order of the three clades is verified with two outgroup-independent methods (Fig. 1,

Fig. S7A) and the outgroup-dependent method (Fig. S1). Collectively, our new strains

appreciably increase the existing phylogenetic diversity of the PB supergroup. Comparative

genomic analysis shows that except for *Bradyrhizobium* sp. ORS285, which is known to have

nod genes and use both a *nod*-dependent and -independent strategy for nodulation [17, 18], all

PB supergroup members lack the *nod* genes.

<u>Fine-scale population structure of the photosynthetic supergroup of *Bradyrhizobium*</u>

Apart from the broad diversity of the PB supergroup, we further asked whether fine-scale

phylogenetic differentiation correlates with the ecological niches where the PB supergroup

members were found. Of the 208 new PB supergroup strains, 205 were collected from three rice

plants located in the same rice field with each plant 5-10 m apart from the others. The remaining

three strains were isolated from grassland and forest. Bacterial isolation was performed from

within the root, rhizosphere, and bulk soil for each plant, collectively giving rise to 145, 23 and

40 PB strains, respectively (Dataset S1). To facilitate the correlative analysis between fine-scale

phylogenetic groups and the niches, we applied the population genomics software PopCOGenT

[19], which defines the boundaries of populations (reported as "main cluster" or MC) between

which gene flow is limited, distinguishes recent gene flow from historical events, and detects

subpopulations (reported as 'subclusters') under ongoing differentiation within a population.

Using this approach, we showed that all strains in Clade 3 share the membership of a

single population (MC1), whereas strains from Clade 1 fall into many populations. In Clade 1,

MC2, MC3, MC4, MC5 and MC6 are each exclusively comprised of the new strains, whereas

MC7 and the remaining unassigned isolates are from published resources. In Clade 1, the within-

and between-population similarity was above and below 95% ANI (Fig. S2), respectively,

consistent with the operational species threshold of 95% ANI [20], whereas the 16S rRNA gene

shows little divergence with the between-population similarity generally above 99% (Fig. S2,

Dataset S3). Based on the available samples, intra-population subdivision occurred within MC1,

MC2 and MC4. We found that each population and subpopulation have members sampled from

different plants and niches (Fig. 1), suggesting that the PB populations sampled from the rice

field are not separated or subdivided according to the physical separation between plant

individuals and the three typical niches (root, rhizosphere, and bulk soil).

Rather, we found an interesting association between important metabolic pathways and

population identity. Remarkably, although the photosynthetic supergroup was named by the

presence of the photosynthesis genes [3], these genes are exclusively and universally found in

MC2, MC3, MC6 and MC7 but completely missing from MC1, MC4 and MC5 (Figure 1,

dataset S4) as well as STM3843 as previously reported [21, 22]. Our result indicates that

photosynthesis is far from being a common characteristic trait that defines the PB phylogroup.

Also interesting is the prevalence of a unique Type III secretion system (T3SS) in MC2,

MC4 and MC5 but completely missing in other populations (Fig. 1, Fig. S3B and S3C, Dataset

S4). It is one of the six T3SS subtypes identified from known *Bradyrhizobium* phylotypes [23]. It

was previously identified only in the PB strain *Bradyrhizobium oligotrophicum* S58 (thus named

'S58-T3SS' subtype) and its function remains unknown [23]. Since the main function of T3SS in

rhizobia is to translocate effector proteins into host cells that modulate the host immune response

or trigger nodulation [24], it cannot be excluded that this T3SS type specifically identified in PB

members plays a role during their interaction with their host plants (*Aeschynomene* spp. and

rice).

We also found that *mxa* gene sets participating in methanol metabolism were specific to

MC1 (Dataset S4), which suggests that these MC1 strains may form intimate associations with

plants considering that methanol is a waste product formed during plant-cell wall degradation

[25]. Our methanol metabolism experiment showed that MC1 strains could not use methanol as a

sole C source (Fig. S4), hinting that the expression of *mxa* genes in these MC1 strains might be

under regulation of unknown mechanisms.

The ability to nodulate *Aeschynomene indica* is a conserved trait shared by all PBs but differs

between the three major PB clades

Since the new PB strains were isolated from non-legume environments, we thought that it

would be very interesting to investigate (i) whether they are capable of performing nitrogen

fixation under free-living and symbiotic states, (ii) whether they are capable of nodulating

Aeschynomene indica, the typical host plant of the previously isolated PB strains, and (iii)

whether there is a difference in nodulation ability across the clades and the populations. The

detailed assay procedures can be seen in Supplementary Text.

All PB supergroup members have a *nif* gene cluster necessary for nitrogenase fixation.

Interestingly, duplication of the *nifH* gene which encodes one of the structural proteins of the

nitrogenase enzyme complex was observed in all PB strains (Fig. S3A), although the functional

consequence is not known. Across the PB supergroup members, an important difference was

observed regarding how the *nif* genes are structured. In most populations, all *nif* genes are co-

located, along with other genes potentially involved in nitrogen fixation. This is not the case for a

few members such as some basal lineages (e.g., *Bradyrhizobium* sp LMG 8443) and some

members assigned to several populations (e.g., SZCCHNR1015 in MC3 and *Bradyrhizobium* sp.

83002 in MC7), where the *nif* genes were found in two contigs. This is likely due to structural

rearrangement, but a DNA assembly artefact cannot be ruled out.

Another original feature observed of the PB *nif* cluster is the universal presence of the

nifV gene encoding for a homocitrate synthase that is involved in the biosynthesis of the

nitrogenase cofactor (FeMo-Co), and which is absent in most other rhizobial lineages [26, 27].

An acetylene reduction assay on 27 representative PB strains covering all three major clades, all

newly identified populations (MC1 to MC6) within Clade 1 and Clade 3, and most of the

identified subpopulations within each population, confirmed their ability to fix nitrogen under

free-living conditions (Fig. S5A), a trait absent in most rhizobia [27].

The plant inoculation experiments showed that all 27 representative strains were also able

to induce nodules on *A. indica* (Fig. 2B, 2C). However, a significant difference in symbiosis

efficiency was observed between the major clades. Clade 1 strains had the same symbiotic

properties as the model strain ORS278, inducing a large number of nitrogen-fixing nodules that

stimulated the growth of the plants (Fig. 2A, 2B, Fig. S5B and Fig. S6A). In contrast, Clade 3

strains formed a much smaller number of nodules compared to Clade 1 strains and the measured

nitrogenase enzyme activity was lower under the symbiotic condition (Fig. 2B, Fig. S5B).

Microscopic analysis showed that the nodules elicited by Clade 3 strains displayed multiple

altered phenotypes: i) some nodules contained necrotic areas, ii) in others the central tissue was

completely digested, and finally, iii) the nodules that displayed less drastic symptoms contained

mainly dead bacteria (Fig. 2C and 2F). These observations indicate that besides inducing fewer

nodules, Clade 3 strains are unable to maintain an effective chronic infection, which explains

why they have no beneficial effect on plant growth (Fig. 2A, 2C and 2F). Interestingly, the

singleton Clade 2 represented by strain STM3843 which is phylogenetically positioned between

Clade 1 and Clade 3, showed an intermediate phenotype. The number and phenotype of the

nodules is comparable to those of Clade 1 strains but stimulation of plant growth is less (Fig. 2A,

2B and 2E).

The symbiotic differences observed between the major clades may result from an

inability of the Clade 3 strains to cope with the plant's immune response, and/or an over-

induction of the plant's defense mechanisms due to the absence or not properly recognized

symbiotic signals. However, it is remarkable that all strains from the PB phylogroup, which

come from different origins and geographical locations, are capable of nodulating *A. indica*.

Concluding remarks

The present study greatly expands the phylogenetic and ecological diversity of the

"photosynthetic" *Bradyrhizobium* supergroup. The photosynthetic trait, which was thought to be

a unifying character of these bacteria, in fact displays a patchy distribution within the

supergroup. In contrast, the ability to develop a NF-independent symbiosis with *Aeschynomene*

spp., despite varying in nodulation efficiency between major clades, appears to be a universal

trait of the phylogroup. We therefore propose to rename this phylogroup as "*nod*-free but

nodulating *Bradyrhizobium*".

The conservation of the nodulation character in such an ecologically diverse phylogroup

and the fact that these strains were mainly isolated from the rice environment lacking selective

pressure from legumes suggest that the symbiotic determinants governing the NF-independent,

T3SS-independent symbiosis may be also essential to their life. This could explain the failures so

far to identify these symbiotic determinants using loss-of-function approaches. Although it is

beyond our goal to reveal these symbiotic determinants in the present study, the resulting new

knowledge leads us to consider tackling this difficult problem using other methods (e.g., gain-of-

function approaches). Furthermore, these data lay the ground for a new evolutionary concept of

an alternative symbiotic process between rhizobia and legumes in which key symbiotic genes

can be inherited vertically rather than transferred horizontally as described for *nod* genes.

In addition to their ability to use a unique symbiotic process with legumes, PB strains

may also have the original function of colonizing rice roots and improving plant growth [17, 18,

28]. This new wealth of genomic sequences of strains isolated from rice could therefore be of

great value in the future to improve our understanding of this plant growth stimulating effect.

Acknowledgements

We thank Xiaoyuan Feng for helping with genome assembly and Jinjin Tao for discussion.

This work was supported by the Hong Kong Research Grants Council Area of Excellence

Scheme (AoE/M-403/16) and a grant from the French National Research Agency ("ET-Nod";

ANR-20-CE20-0012).

Data availability

The genomic sequences and raw reads of the 208 newly sequenced *Bradyrhizobium* isolates

are in the process of uploading to NCBI (Project ID: PRJNA983111) and will be released in the

next version of the manuscript.

Competing interests

The authors declare no competing interests in relation to this work.

References

1.   Parker MA. The spread of Bradyrhizobium lineages across host legume clades: from

Abarema to Zygia. *Microb Ecol* 2015; **69:** 630-640.

2.   Sprent JI, Ardley J, James EK. Biogeography of nodulated legumes and their nitrogen-fixing

symbionts. *New Phytol* 2017; **215:** 40-56.

3.   Avontuur JR, Palmer M, Beukes CW, Chan WY, Coetzee MPA, Blom J, et al. Genome-

informed Bradyrhizobium taxonomy: where to from here? *Syst Appl Microbiol* 2019; **42:** 427-

439.

4.   Ormeno-Orrillo E, Martinez-Romero E. A Genomotaxonomy View of the Bradyrhizobium

Genus. *Front Microbiol* 2019; **10:** 1334.

5.   Giraud E, Fleischman D. Nitrogen-fixing symbiosis between photosynthetic bacteria and

legumes. *Photosynthesis Research* 2004; **82:** 115-130.

6.   Lerouge P, Roche P, Faucher C, Maillet F, Truchet G, Prome JC, et al. Symbiotic host-

specificity of Rhizobium meliloti is determined by a sulphated and acylated glucosamine

oligosaccharide signal. *Nature* 1990; **344:** 781-784.

7.   Oldroyd GED. Speak, friend, and enter: signalling systems that promote beneficial

symbiotic associations in plants. *Nature Reviews Microbiology* 2013; **11:** 252-263.

8.   Bonaldi K, Gourion B, Fardoux J, Hannibal L, Cartieaux F, Boursot M, et al. Large-scale

transposon mutagenesis of photosynthetic Bradyrhizobium sp. strain ORS278 reveals new

genetic loci putatively important for nod-independent symbiosis with Aeschynomene indica. *Mol*

Plant Microbe Interact 2010; **23:** 760-770.

9.   Okazaki S, Tittabutr P, Teulet A, Thouin J, Fardoux J, Chaintreuil C, et al. Rhizobium-

legume symbiosis in the absence of Nod factors: two possible scenarios with or without the

T3SS. *ISME J* 2016; **10:** 64-74.

10. Teulet A, Busset N, Fardoux J, Gully D, Chaintreuil C, Cartieaux F, et al. The rhizobial type

III effector ErnA confers the ability to form nodules in legumes. *Proc Natl Acad Sci U S A* 2019;

116: 21758-21768.

11. Ratu STN, Teulet A, Miwa H, Masuda S, Nguyen HP, Yasuda M, et al. Rhizobia use a

pathogenic-like effector to hijack leguminous nodulation signalling. *Sci Rep* 2021; **11:** 2034.

12. Camuel A, Teulet A, Carcagno M, Haq F, Pacquit V, Gully D, et al. Widespread

Bradyrhizobium distribution of diverse Type III effectors that trigger legume nodulation in the

absence of Nod factor. *ISME J* 2023; **In press**.

13. Chaintreuil C, Giraud E, Prin Y, Lorquin J, Ba A, Gillis M, et al. Photosynthetic

bradyrhizobia are natural endophytes of the African wild rice Oryza breviligulata. *Appl Environ*

Microbiol 2000; **66:** 5437-5447.

14. Piromyou P, Greetatorn T, Teamtisong K, Okubo T, Shinoda R, Nuntakij A, et al. Preferential

association of endophytic bradyrhizobia with different rice cultivars and its implications for rice

endophyte evolution. *Appl Environ Microbiol* 2015; **81:** 3049-3061.

15. Van Berkum P, Eardly BD. The aquatic budding bacterium Blastobacter denitrificans is a

nitrogen-fixing symbiont of Aeschynomene indica. *Appl Environ Microbiol* 2002; **68:** 1132-

1136.

16. Okubo T, Fukushima S, Itakura M, Oshima K, Longtonglang A, Teaumroong N, et al.

Genome analysis suggests that the soil oligotrophic bacterium Agromonas oligotrophica

(Bradyrhizobium oligotrophicum) is a nitrogen-fixing symbiont of Aeschynomene indica. *Appl*

Environ Microbiol 2013; **79:** 2542-2551.

17. Bonaldi K, Gargani D, Prin Y, Fardoux J, Gully D, Nouwen N, et al. Nodulation of

Aeschynomene afraspera and A. indica by photosynthetic Bradyrhizobium Sp. strain ORS285:

the nod-dependent versus the nod-independent symbiotic interaction. *Mol Plant Microbe Interact*

2011; **24:** 1359-1371.

18. Giraud E, Moulin L, Vallenet D, Barbe V, Cytryn E, Avarre JC, et al. Legumes symbioses:

absence of Nod genes in photosynthetic Bradyrhizobia. *Science* 2007; **316:** 1307-1312.

19. Arevalo P, VanInsberghe D, Elsherbini J, Gore J, Polz MF. A Reverse Ecology Approach Based on a Biological Definition of Microbial Populations. *Cell* 2019; **178:** 820-834 e814.

20. Konstantinidis KT, Tiedje JM. Genomic insights that advance the species definition for prokaryotes. *Proc Natl Acad Sci U S A* 2005; **102:** 2567-2572.

21. Mornico D, Miche L, Bena G, Nouwen N, Vermeglio A, Vallenet D, et al. Comparative genomics of aeschynomene symbionts: insights into the ecological lifestyle of nod-independent photosynthetic bradyrhizobia. *Genes (Basel)* 2011; **3:** 35-61.

22. Miché L, Moulin L, Chaintreuil C, Contreras-Jimenez JL, Munive-Hernández JA, Del Carmen Villegas-Hernandez M, et al. Diversity analyses of Aeschynomene symbionts in Tropical Africa and Central America reveal that nod-independent stem nodulation is not restricted to photosynthetic bradyrhizobia. *Environ Microbiol* 2010; **12:** 2152-2164.

23. Teulet A, Gully D, Rouy Z, Camuel A, Koebnik R, Giraud E, et al. Phylogenetic distribution and evolutionary dynamics of nod and T3SS genes in the genus Bradyrhizobium. *Microb Genom* 2020; **6**.

24. Teulet A, Camuel A, Perret X, Giraud E. The Versatile Roles of Type III Secretion Systems in Rhizobium-Legume Symbioses. *Annu Rev Microbiol* 2022; **76:** 45-65.

25. Nemecek-Marshall M, MacDonald RC, Franzen JJ, Wojciechowski CL, Fall R. Methanol Emission from Leaves (Enzymatic Detection of Gas-Phase Methanol and Relation of Methanol Fluxes to Stomatal Conductance and Leaf Development). *Plant Physiol* 1995; **108:** 1359-1368.

26. Hakoyama T, Niimi K, Watanabe H, Tabata R, Matsubara J, Sato S, et al. Host plant genome overcomes the lack of a bacterial gene for symbiotic nitrogen fixation. *Nature* 2009; **462:** 514-517.

27. Nouwen N, Arrighi J-F, Cartieaux F, Chaintreuil C, Gully D, Klopp C, et al. The role of

rhizobial (NifV) and plant (FEN1) homocitrate synthases in Aeschynomene/photosynthetic

Bradyrhizobium symbiosis. *Scientific Reports* 2017; **7:** 448.

28. Cai W, Ou F, Staehelin C, Dai W. Bradyrhizobium sp. strain ORS278 promotes rice growth

and its quorum sensing system is required for optimal root colonization. *Environ Microbiol Rep*

2020; **12:** 656-666.

 Figure legends

Figure 1. The phylogenomic tree and population delineation of the "photosynthetic"

Bradyrhizobium (PB) supergroup. The phylogenomic tree was rooted using the minimum

variance (MV) method. The 208 PB strains sequenced in the present study are indicated by black

dots in the innermost layer surrounding the phylogeny. The ultrafast bootstrap values higher than

or equal to 95% calculated by IQ-Tree were labeled on the nodes with black circles.

Figure 2. Symbiotic properties of several representative strains of the main populations

identified in the PB supergroup. (A) Comparison of growth of the *A. indica* plants (leaf

phenotype) non-inoculated (NI) or inoculated with different representative strains of PB. The six

MC1 representative strains are presented in the following order: SZCCHNR1021,

SZCCHNR1085, SZCCHNS30121, SZCCHNGR1005, SZCCHNRI2010 and SZCCHNF30212.

For the other MCs, only one representative strain is shown: SZCCHNRI2049 (MC2),

SZCCHNS3002 (MC3), SZCCHNS30592 (MC4), SZCCHNRI3016 (MC5) and SZCCHNS2021

(MC6), Photos were taken 17 days after inoculation. (B) Nodule number on *A. indica* plants

induced by different representative strains of PB at 17 days after inoculation. Box plots show the

results of one of the two experiments performed independently (4 plants each). (C) Symbiotic

phenotypes of some representative strains tested on *A. indica*. Column 1 and 2: photo of roots

and nodules. Scale bars: column 1, 0.5 cm; column 2, 0.2 cm. Column 3: Micro-sections of

nodules observed using light microscopy. Scale bars: 250 µm. (D) Confocal microscopy images

of micro-sections of nodules elicited by Clade 1 strains. The nodules formed by

SZCCHNRI3016 from MC4 and ORS278 are shown as examples. The central nodule tissue is

intracellularly infected with live spherical bacteroids stained green by SYTO9. (E) Confocal

microscopy images of micro-section of nodules elicited by Clade 2 strain STM3843. The nodules

look normal, but a mix of live and dead bacteria (stained red by propidium iodide) can be

noticed. (F) Confocal microscopy images of micro-section of nodules elicited by Clade 3 strains

displaying different phenotypes: nodule containing mainly dead bacteria (SZCCHNS30121),

nodule with a completely digested central tissue (SZCCHNRI2010); and nodule with a necrotic

area and a digested zone (SZCCHNGR1005). Scale bars for (D), (E) and (F): column 1, 200 µm;

column 2, 10 µm.

Tree scale: 0.1

Functional gene clusters
- Methanol into formaldehyde
- Photosynthetic genes
- T3SS gene clusters
- *nodABC*

Plants
- *Oryza sativa* indica 1
- *Oryza sativa* indica 2
- *Oryza sativa* indica 3
- *Houttuynia cordata*
- *Camphora officinarum*

Niches
- Root
- Rhizosphere
- Soil

Sub_clusters
- Sub_cluster 1
- Sub_cluster 2
- Sub_cluster 3
- Sub_cluster 4
- Sub_cluster 5
- Sub_cluster 6

- strains used for symbiosis assay
- sequenced in this study

Background > Main clusters
- MC1
- MC2
- MC3
- MC4
- MC5
- MC6
- MC7

Outer

Inner

MC7 MC6 MC5 MC4 MC3 MC2

Clade1 Clade2 Clade3

Main cluster 1 (MC1)