Phage predation and antibiotic exposure are inversely associated with disease severity and shape pathogen genetic diversity in cholera patients

N. Madi[1,2], E. T. Cato[3], Md. Abu Sayeed[3], K. Islam[4], Md. I.UI. Khabir[4], Md. T. R. Bhuiyan[4], Y. Begum[4], M.M. Rashid[4], A. Creasy-Marrazzo[3], L. Brinkley[3], M. Kamat[4], A. Cuénod[1], L. S. Bailey[5], K. B. Basso[5], F. Qadri[4], A. I. Khan[4*], B. J. Shapiro[1,2,6,7*], E. J. Nelson[3*]

[1] Department of Microbiology & Immunology, McGill University, Montréal, Canada
[2] Département de Sciences Biologiques, Université de Montréal, Montréal, Canada
[3] Departments of Pediatrics and Environmental and Global Health, University of Florida, Gainesville, FL, USA
[4] Infectious Diseases Division (IDD) & Nutrition and Clinical Services Division (NCSD), International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh
[5] Department of Chemistry, University of Florida, Gainesville, FL, USA
[6] McGill Genome Centre, McGill University, Montréal, Canada
[7] McGill Centre for Microbiome Research, McGill University, Montréal, Canada

[*] Corresponding authors

Email: eric.nelson@ufl.edu; jesse.shapiro@mcgill.ca; ashrafk@icddrb.org

Keywords: Cholera, diarrhoea, diarrhea, *Vibrio cholerae*, bacteriophage, phage, phage resistance, antibiotics, antimicrobial resistance, AMR, predator, prey, metagenomics

ABSTRACT

Nearly a century has passed since d'Hérelle and colleagues discovered phages that infect *Vibrio cholerae* and influence the clinical and epidemiologic trajectories of cholera outbreaks. Despite an increasingly detailed picture of the molecular mechanisms of resistance and counter-resistance that shape phage and bacterial interactions, we lack an understanding of how these interactions play out during natural infection, how they are affected by antibiotic exposure, and how they relate to clinical outcomes. To fill these gaps, we conducted a nation-wide study of diarrheal disease patients in the cholera-endemic setting of Bangladesh. A total of 2574 stool samples were collected from enrolled patients at hospital admission and screened for *V. cholerae* and virulent phages (ICP1, 2, or 3). All 282 culture positive samples, and an additional 107 culture negative but PCR positive samples, were analyzed by shotgun metagenomic sequencing. From these metagenomes, we estimated the relative abundances of *V. cholerae,* phages, and members of the gut microbiome, while accounting for antibiotic exposure determined by quantitative mass spectrometry. Consistent with d'Hérelle's thesis, we observed higher phage to *V. cholerae* ratios in patients with mild dehydration, demonstrating in the modern era that phages are indicative of disease severity. Antibiotics were associated with less *V. cholerae* and with mild disease; ciprofloxacin in particular was associated with several known antibiotic resistance genes. Phage resistance genes in the *V. cholerae* integrative conjugative element (ICE) were associated with lower phage to *V. cholerae* ratios. In the absence of detectable ICEs, phages shaped genetic diversity, selecting for nonsynonymous point mutations in the *V. cholerae* genome. Together, our results point to both antibiotics and phages as inverse correlates of disease severity, while also selecting for resistance genes or mutations within cholera patients.

INTRODUCTION

The secretory diarrheal disease cholera is caused by the Gram negative bacterium *Vibrio cholerae* (*Vc*) and can progress to life-threatening hypovolemic shock in less than 12 hours (Andrews et al. 2017). Cholera remains a major public health problem due to inadequate sanitation and safe drinking water. Global estimates of cholera burden are 1.3-4.0 million cases and 21,000 to 143,000 deaths annually (Ali et al. 2015). As of January 2023, there were over 30 countries with active outbreaks necessitating the WHO to escalate the response to its highest level (grade three) ("Cholera – Global Situation" report, 2023). Rehydration is the primary life-saving maneuver for cholera patients. With effective rehydration, mortality rates fall from 20% and above to less than 1%; antibiotics reduce stool volume and duration of diarrhea but are considered supportive and reserved for patients with more severe disease to reduce selection for antibiotic resistance (Islam et al. 1995; Dromigny et al. 2002; Weill et al. 2017; Nelson et al. 2011, 2009). Nevertheless, antibiotic-resistant *V. cholerae* have emerged and spread globally (Das et al. 2020; Weill et al. 2017; Lassalle et al. 2022). Mechanisms of resistance are diverse and reside in the core genome and on mobile genetic elements, including the SXT integrative conjugative element (ICE) (Rivard, Colwell, and Burrus 2020). SXT is a ~100 kb ICE and can harbor resistance to sulfamethoxazole and trimethoprim, ciprofloxacin (*qnr$_{vc}$*), trimethoprim (*dfra31*) and streptomycin (*aph(6)*) (Waldor, Tschäpe, and Mekalanos 1996; Dalsgaard et al. 1999; Beaber, Hochhut, and Waldor 2002; Creasy-Marrazzo et al. 2022). Recent work has also shown that the ICE can encode diverse phage resistance mechanisms (LeGault et al. 2021).

V. cholerae is targeted by numerous lytic bacteriophages (phages) (Nelson et al. 2008; Boyd et al. 2021; K. D. Seed et al. 2011). With rising levels of antibiotic resistance, phages are a promising alternative or complementary therapy. The first clinical trials of phage therapy cocktails occurred during the Cholera Phage Inquiry from 1927 to 1936 (Summers 1993; D'Herelle and Malone 1927). In a *proto* randomized controlled trial, the Inquiry found the odds of mortality were reduced by 58% among those with phage therapy, with an absolute reduction in

mortality of 10% (95% CI 0.27-0.64) (Pasricha, de Monte, and O'Flynn 1936) (reanalyses by E.

J. Nelson). Despite these early findings, there is a lack of evidence in the modern era that links

phage predation with disease severity during *natural* infection in humans. A collection of

tangentially related studies support the hypothesis that phage mitigate severity:

Environmentally, phages in the aquatic environment are negatively correlated with cholera

cases in Dhaka, Bangladesh over time, suggesting a role for phages in influencing epidemic

dynamics (Faruque, Naser, et al. 2005). Clinically, a higher percentage of cholera patients shed

phage towards the end of an outbreak (Faruque, Islam, et al. 2005). Computationally, models

predict that phage can dampen outbreaks (Jensen et al. 2006). Experimentally, animal studies

found phage exposure was negatively associated with burden of colonization and disease

severity (Zahid et al. 2008; Jaiswal et al. 2013; Yen, Cairns, and Camilli 2017; Nelson et al.

2008). In this setting, a key unanswered mechanistic question is how phage, antibiotics, and

associated resistance factors interact to impact disease severity.

To address this question, we conducted a national prospective longitudinal study in the

cholera endemic setting of Bangladesh. Stool samples were collected at hospital admission

from diarrheal disease patients and screened for *V. cholerae* or *V. cholerae* phage. From these

positive samples, we sequenced shotgun metagenomes to quantify the relative abundances of

V. cholerae, phages, and gut microbiome taxa, while accounting for antibiotic exposure

determined by mass spectrometry. We demonstrate that severe dehydration is inversely

associated with phage ICP1. We also find evidence for non-linear predator-prey interactions

within infected hosts, but these interactions are suppressed at high levels of azithromycin.

Known phage-resistance elements (ICEs) are associated with lower phage:*V. cholerae* ratios. In

the absence of detectable ICEs, phages select for nonsynonymous point mutations in the *V.*

cholerae genome, with potential evolutionary consequences. Together, our results support a

hierarchy of selective pressures and resistance mechanisms evolving within individual cholera

patients.

RESULTS & DISCUSSION

Study overview. A total of 2574 stool samples were collected from enrolled participants from

March to December 2018; collection continued to April 2019 at one site (icddr,b). A total of 282

samples (10.9%) were culture-positive for *V. cholerae*. Among culture-negative samples, 10% of

samples were randomly selected and screened by PCR for *V. cholerae* or phage (ICP1,2,3)

which generated an additional 107 positive samples (**Table S1**). Stool metagenomes were

successfully sequenced from 88.4% (344/389) of these samples; 35%(122) were from the

icddr,b. Detection rates for *V. cholerae,* ICP1, ICP2, and ICP3 were 55% (190), 18% (61), 1%

(4) and 8% (28), respectively. Select antibiotics (**Table S2**) were quantified in stool using liquid

chromatography-mass spectrometry.

Metagenomic correlates of disease severity and succession. In the context of enrollment at

hospital admission, we expected to sample patients at different stages in their disease

progression, beginning with high levels of *Vc* followed by *E. coli* and, in most cases, a return to

a normal microbiome in which anaerobic bacteria dominate (David et al. 2015). We

hypothesized that this ecological succession would be accompanied by a progression from

severe to mild dehydration. Consistent with this hypothesis, we identified *Vc* as an indicator

species of severe dehydration (Methods). *Vc was* more likely to be present and higher in

relative abundance in patients with severe dehydration, while two *Bifidobacterium* species and

E. coli were indicators of mild dehydration (**Figure 1A, Table S3**). In a time series of cholera

patients, the phage ICP1 peaked on the first day of infection (David et al. 2015) yet intriguingly

ICP1 was an indicator of mild dehydration in our analysis (**Figure 1A**). This contrasts with ICP3,

which despite being less frequently detected in our study (28 samples with >0.1% ICP3 reads,

compared to 61 samples with >0.1% ICP1), was an indicator of severe dehydration. This

suggests that different phages can have contrasting disease associations. For subsequent

analyses, we focus on the more prevalent ICP1 phage.

Although the distribution of ICP1 relative abundance is variable and less clearly

associated with dehydration status than *Vc* (**Figure S1**), higher ratios of ICP1 to *Vc* were

associated with mild dehydration (**Figure 1B**). This ratio is particularly informative, as it provides

insight into how strongly ICP1 might suppress its bacterial host. These analyses suggest that

ICP1 plays a role in modulating disease severity, or that it could be associated with later stages

of diseases than previously thought. Our study design prevents us from confidently

disentangling these possibilities; however, we recorded self-reported duration of diarrhea,

providing an approximate control for disease progression. We found higher relative abundances

of ICP1 associated with mild dehydration at early stages of disease (duration of diarrhea <72h)

but not at late stages (**Figure S1B and D**). We therefore cannot exclude a model in which ICP1

suppresses *Vc* at early disease stages to reduce disease severity; nor can we exclude ICP1 as

a non-causal biomarker of dehydration severity. In either case, ICP1 could provide a clinically

useful indicator of disease severity.

Figure 1. Dehydration severity is inversely associated with higher ICP1:*Vc* ratios in stool metagenomes. (A) Relative abundances of the most prevalent species in patients with severe, moderate, or mild dehydration; these conventions equate to the World Health Organization (WHO) conventions of severe, some and no dehydration, respectively. Significant indicator species for severe or mild dehydration are shown in red or blue bold text, respectively. See Table S3 for full indicator species results. (B) The ICP1:*V. cholerae* ratio is higher in patients with mild dehydration. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test, adjusted for multiple tests using the Benjamini-Hochberg (BH) method. Only significant *P*-values (<0.05) are shown. Only 323 samples with *V. cholerae*>0% of metagenomic reads were included, with 165 from severe, 128 from moderate, and 30 from mild cases. A pseudocount of one was added to the ratio before log transformation. In (A) and (B) the solid horizontal line is the median and the boxed area is the interquartile range. (C) Redundancy analysis (RDA) showing relationships among the seven most dominant bacterial species and explanatory variables: phages, patient metadata, and antibiotic concentration. Angles between variables (arrows) reflect their correlations; arrow length is proportional to effect size. Samples (points) are colored by dehydration severity. All displayed variables are significant (*P*<0.05, permutation test) except ICP2, ICP3, and doxycycline (Table S4). For the RDA: R^2=0.25 and adjusted R^2=0.184, permutation test *P* = 0.001). To improve readability, collection date and location are not shown (see Figure S3 for these details).

Antibiotic exposure is associated with mild disease and resistance genes. To visualize the

complex relationships between disease severity, bacteria, and phage in the context of antibiotic

exposure, we used redundancy analysis (RDA; **Figure 1C; Table S4**). For simplicity, the seven

most dominant bacterial species identified by principal components analysis were included

(**Figure S2**). As explanatory variables, we used the most important clinical data, chosen by

forward selection beginning with phages and antibiotic concentrations (**Figure S3**). In

accordance with the indicator species analysis (**Figure 1A, Table S3**), *Vc* was positively

correlated with severe dehydration (**Figure 1C**). ICP1 was moderately associated with *Vc*,

consistent with a phage's reliance on its host for replication, but less associated with severe

dehydration. Both azithromycin (AZI) and ciprofloxacin (CIP) were negatively correlated with *V.*

cholerae and severe dehydration, suggesting their role in suppressing cholera infection and

disease.

To test the hypothesis that antibiotic exposure selects for antibiotic resistance genes

(ARGs), we assessed the relative abundances of 634 known ARGs conferring resistance to 34

antibiotic classes using deepARG (Arango-Argoty et al. 2018) applied to the metagenomic data.

To identify associations between these ARGs and gut microbes, we ran a multiple factor

analysis (MFA) with the 37 most dominant species from a PCA (20 highest coordinates on both

axes, Methods) and all the ARGs. ARGs were primarily associated with three different

taxonomic clusters dominated by (i) *V. cholerae*, (ii) *Escherichia* spp. and *Shigella* spp., and (iii)

Prevotella spp. and *Bifidobacterium* spp. (**Figure S4**). The *V. cholerae* cluster was associated

with 12 ARGs: *ompT, ompU, varG, dfrA1, dfrA16, aph3, aph6, mexI, floR, tet34, tet35*, and a

chloramphenicol exporter. Several of these targets were previously linked to antibiotic

resistance in *V. cholerae* (Creasy-Marrazzo et al. 2022; Monir et al. 2023). To determine if these

ARGs correlated with antibiotic exposure, we compared the relative abundance of ARGs

(normalized to *V. cholerae* reads) at different levels of antibiotic exposure: above or below the

minimum inhibitory concentration (MIC) set by experiments under aerobic vs anaerobic

conditions (**Table S5**) (Methods). CIP exposure was associated with higher relative abundance

of *dfrA16* (Wilcoxon test, BH corrected *P*=0.0178 and 0.0026 for anaerobic and aerobic MIC,

respectively), *aph6* (BH corrected *P*=0.084 for anaerobic and 0.0552 for aerobic MIC) (**Figure**

S5). No significant correlations were observed with azithromycin or doxycycline. Together, these

results provide evidence that exposure to certain antibiotics selects for resistance genes in the

human gut.

Predator-prey dynamics between phage and pathogen. Offset oscillations between *Vc* and

its phages in the aquatic environment have been observed at the broader population level in

Bangladesh, supporting the hypothesis that these predator-prey dynamics influence the

progression of epidemics (Faruque, Islam, et al. 2005; Jensen et al. 2006). This hypothesis has

not been revisited in nearly two decades, and it remains unclear if the dynamics of aquatic

phages are mirrored by phages within infected patients. To address these questions, we

focused on a one-year time series from our most sampled site, icddr,b, in Dhaka, and tracked

the mean relative abundances of *Vc* and ICP1 across patients over time. Within the context of a

limited sample size, the abundance of *V. cholerae* was high from March through July 2018;

ICP1 abundance was low in March 2018 at the beginning of the epidemic, increased in July,

and spiked in August as *V. cholerae* declined (**Figure 2A, B**). ICP1 and *V. cholerae* were

inversely correlated from September 2018 onward, with a spike in *V. cholerae* corresponding to

a decline of ICP1 in September, and a peak in ICP1 associated with a decline of *V. cholerae* in

November. *V. cholerae* and ICP1 were inversely correlated within patients over time (Spearman

r=-0.25, P=0.017; **Figure 2C**), mirroring the trend previously observed between recorded

cholera cases and environmental phage concentrations (Faruque, Naser, et al. 2005). Sampling

sites outside Dhaka had sparser time series, but also generally supported alternating peaks of

Vc and ICP1 (**Figure S6**). Our results suggest that phages may contribute to suppressing *Vc*

not just in the environment, but also within patients. The degree of coupling between phage

concentrations in the environment and within patients could therefore be relevant to epidemic

dynamics and deserves further study.

Figure 2. Offset temporal dynamics of *V. cholerae* and phage ICP1. Relative abundances of (A) *V.
cholerae* and (B) ICP1 over time (binned by month) in metagenomes sampled from the icddr,b (94
samples with *Vc*>0.5% or ICP1>0.1% of metagenomic reads). Red line shows the mean. The boxplots
contain 16 samples from 2018-03, 25 samples from 2018-04,17 samples from 2018-05, 5 samples from
2018-06, 4 samples from 2018-07, 3 samples from 2018-08, 4 samples from 2018-09, 9 samples from
2018-10, 2 samples from 2018-11, 8 samples from 2019-03, and 1 sample from 2019-04. (C) Spearman
correlation between *V. cholerae* and ICP1 in the same samples as in A and B. The solid horizontal line is
the median and the boxed area is the interquartile range. Panels (D) and (E) show GAM results, fit to data
from all 344 samples. (D) *V. cholerae* declines in relative abundance with higher relative abundance of
ICP1. (E) The relationship between ICP1 and *V. cholerae* is affected by azithromycin (AZI) concentration.
The illustrated AZI concentrations show regular intervals between the minimum (0 ug/ml) and maximum
(78 ug/ml) observed values.

Antibiotic exposure suppresses predator-prey dynamics. Our data provide a unique

opportunity to determine if and how antibiotics interact with phage to suppress *V. cholerae*

within patients. To do so, we modeled the relationships between ICP1, *V. cholerae* and

antibiotic exposure within each patient. We fit a generalized additive model (GAM) for *V.*

cholerae relative abundance as a function of ICP1, antibiotic exposure, and their interaction,

including dehydration status as a random effect. We fit GAMs with all antibiotics and their

interaction with ICP1, as well as separate models with each antibiotic, alone or in combination,

and compared them based on their AIC (**Table S6**). The most parsimonious model (with the

lowest AIC) included the interaction between AZI and ICP1 as a predictor of *V. cholerae*

abundance. *V. cholerae* was affected by AZI both directly (GAM, P=0.00236, Chi-square test)

and in interaction with ICP1 (GAM, P=0.026, Chi-square test); the effect of ICP1 alone was not

significant. Consistent with previous reports that AZI suppresses *V. cholerae (Alexandrova et al.*

2019) and with our RDA results (**Figure 1C**), we found a negative relationship between *V.*

cholerae and AZI (**Figure 2D**). This is also consistent with our observation that no annotated

ARGs were associated with AZI (**Figure S7**), suggesting that *V. cholerae* in our samples is

generally AZI-sensitive. The relationship between *V. cholerae* and ICP1 was quadratic: at low

ICP1 abundance, the relationship was positive but became negative at higher ICP1 abundance

(**Figure 2E**). This alternation between positive and negative correlations is consistent with

predator-prey dynamics within infected patients (Carr et al. 2019). However, at higher

concentrations of AZI, the quadratic relationship was flattened, effectively suppressing the

phage-bacteria interaction.

Integrative and conjugative elements (ICEs) are associated with phage suppression.

SXTs are ~100kb integrative and conjugative elements (ICEs) that have been associated with

resistance to both antibiotics and phages and were first discovered in *V. cholerae* (Waldor,

Tschäpe, and Mekalanos 1996). SXTs have conserved 'core' genes along with variable

'hotspots' encoding different genes; for example, hotspot 5 is a ~17kb region associated with

phage resistance. At the time of our sampling, ICEVchInd5 and ICEVchind6 were the two most

prevalent ICEs in Bangladesh (LeGault et al. 2021). These ICEs differ in their anti-phage

systems: ICEVchInd5 (henceforth "ind5") encodes a type 1 bacteriophage exclusion (BREX)

system while ICEVchInd6 ("ind6") encodes several other restriction-modification systems

(LeGault et al. 2021).

We enumerated ICEs in metagenomes by mapping reads against reference sequences

for ind5 (NCBI accession GQ463142.1) and ind6 (accession MK165649.1). An ICE was defined

as present when 90% of its length was covered by at least one metagenomic read (**Figure**

S8A). We found that 59% (144/244) of the samples contained ICEVchind5, 10.6% (26/244)

contained ICEVchind6, and 22.1% (54/244) had no detectable ICE. The lack of ICE detections

was not due to the lack of *V. cholerae* in a metagenome because ICE-negative samples did not

contain fewer *V. cholerae* reads (**Figure S8B**).

Resistance mechanisms on ICEs have been shown to suppress phage *in vitro*, but their

relevance within human infection remains unclear. We found that metagenomes without a

detectable ICE (denoted as ICE-) were associated with higher phage to *V. cholerae* ratios

(**Figure 3**). The effect was strongest for ICP1, which had the largest sample size (**Figure 3A**).

This observation is consistent with ICE-encoded mechanisms suppressing phage within

patients. Higher ICP1:*Vc* ratios, which tend to occur in ICE- patients, were also associated with

mild dehydration (**Figure 3B**). The phage:*Vc* ratios varied according to the precise phage-ICE

combination, suggesting that different ICE hotspots might confer resistance to different phages

(**Figure 3, Figure S9**). Together, these results demonstrate a role for ICEs in suppressing

phages during human infections, complementing and generally confirming the predictions of

laboratory experiments (LeGault et al. 2021). That said, the suppression is not complete, and

further experiments are needed to dissect the underlying causal relationships.

Figure 3. Integrative conjugative elements (ICEs) are associated with lower ICP1:*V. cholerae* ratios in patient metagenomes. (A) Distribution of ICP1:*Vc* ratios across patients with different ICE profiles. (B) The same data as (A) binned into boxplots according to dehydration status. (C) Distribution of phage:*Vc* ratios, including the sum of all phages (ICP1, ICP2, ICP3). (D) The same data as (C) binned into boxplots according to dehydration status. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test adjusted with the Benjamini-Hochberg (BH) method. Only *P*-values < 0.1 are shown. Only samples with sufficient *Vc* or ICP1 were included (224 samples with *Vc*>0.5% or phages >0.1% of metagenomic reads), of which 54 samples were ICE-, 26 were ind6+ and 144 were ind5+. For clarity, the Y-axes were log10 transformed after adding one to the ratios. The solid horizontal line is the median and the boxed area is the interquartile range.

Hypermutation generates V. cholerae genetic diversity. In addition to variation in gene content in ICEs and other mobile elements, resistance to phages and antibiotics may be conferred by point mutations (single nucleotide variants; SNVs) that existed before or emerged *de novo* during infection. Although we cannot exclude mixed infections by different *Vc* strains as a source of diversity, there was no evidence for more than one strain co-infecting a patient in our study population (**Figure S10**). We previously found a generally low level of *V. cholerae* genetic diversity within individual cholera patients (Levade et al. 2017), with the exception of hypermutation events characterized by DNA repair defects and hundreds of SNVs in the *Vc* genome, primarily transversion mutations (Levade et al. 2021). Here, with a much larger sample size, we can better quantify the frequency of hypermutator phenotypes, and test if within-host *V. cholerae* diversity is associated with phages or antibiotics – both of which could potentially select for resistance mutations. After quality filtering the metagenomic reads, we used InStrain (Olm et al. 2021) to quantify within-host *V. cholerae* diversity in 133 samples that passed the post-instrain filter (Methods). To identify likely hypermutators, we independently counted samples with defects (nonsynonymous mutations) in any of 65 previously defined DNA repair genes (Jolivet-Gougeon et al. 2011) or a high rate of SNVs (25 or more) (Levade et al. 2021). We found that 35% of samples (47/133) had both a high SNV count and nonsynonymous mutations in DNA repair genes – making them likely hypermutator samples. Higher SNV counts were significantly associated with DNA repair defects (Fisher's exact test, *P*=2.2e-16), consistent with these defects resulting in higher mutation rates within patients. The number of SNVs was not significantly confounded by *V. cholerae* genome coverage (**Figure S11A**). Consistent with our previous study (Levade et al. 2021), putative hypermutators had a distinct mutational profile, enriched in transversions (**Figure S11B**). This suggests that hypermutation is common and detectable in over one third of cholera patient stool metagenomes. We also observed that samples with high numbers of SNVs but no DNA repair defects had somewhat higher levels of phage ICP1 (**Figure S11D**). This suggested that phage could select for

resistance mutations even in non-mutators. Nevertheless, for subsequent analysis, we

considered all SNVs together, regardless of whether they were generated by hypermutation or

not.

Phages, not antibiotics, are associated with V. cholerae within-host diversity. We asked if

V. cholerae within-host diversity was associated with phage or antibiotics – both likely strong

selective pressures on bacterial populations. To do so, we fit GLMMs with phages and

antibiotics as predictors of the number of high-frequency nonsynonymous (NS) SNVs in the *V.*

cholerae genome. We focused on higher-frequency SNVs (>10% within a sample) as more

likely to be beneficial, and NS SNVs as those more likely to have fitness effects. We fit several

models with different combinations of predictors: from a model with all antibiotics and their

interaction with ICP1 to separate models with each antibiotic and its interaction with ICP1, we

added *V. cholerae* abundance as a fixed effect to the model to control for any coverage effect

(**Table S7**). The most parsimonious model included *V. cholerae* abundance and the interaction

between *V. cholerae* and ICP1 as predictors of the number of high-frequency NS SNVs. Adding

antibiotics or their interaction with ICP1 did not improve the model (**Table S7**), suggesting a

limited role for antibiotics in selecting for point mutations within patients.

In the model, *V. cholerae* relative abundance and the interaction between *V. cholerae*

and ICP1 both had significant effects (GLMM, Wald test, *P*=0.00246 and *P*=0.00494

respectively). The negative relationship between *V. cholerae* and the number of high-frequency

NS SNVs (**Figure 4A**) is not easily explained by coverage artifacts, since the total number of

SNVs is not associated with *V. cholerae* relative abundance (**Figure S11A**). The number of

high-frequency NS SNVs rises with increasing ICP1 – but only when *V. cholerae* abundance is

high (**Figure 4A**). As a control, we ran the same GLMM on NS SNVs without a minimum

frequency cutoff and found no significant effects, suggesting that the interaction between ICP1

and *V. cholerae* on SNV count is specific to high-frequency mutations, rather than low-

frequency mutations that are more likely neutral or deleterious. These data support a scenario in

which ICP1 selects for NS SNVs when the *V. cholerae* population is large enough to respond

efficiently to selection – for example, at the beginning of an infection or in the absence of

antibiotics.

If phage select for beneficial mutations, we would expect these mutations to increase in

frequency at higher phage abundance. We lack time-series data from individual patients, but the

relative abundance of phage provides a proxy for the combined effect of time and strength of

phage selection. To test this expectation, we fit a GAM with the average within-sample

frequency of SNVs as a function of ICP1, antibiotics, and their interactions. We included the

fixed effect of ICE presence/absence as another factor that could provide phage or antibiotic

resistance, as well as mutation type to differentiate among non-synonymous (NS), synonymous

(S), and intergenic (I) mutations. We fit GAMs with all antibiotics and their interaction with ICP1,

as well as simpler models with each antibiotic separately, and compared them based on AIC

(**Supplementary Table S8**). The most parsimonious model included the interaction between

ICP1, ICE and mutation type, but not antibiotics. ICP1 was a strong predictor of higher

frequency NS SNVs in the absence of a detectable ICE (**Figure 4B**). Samples in this analysis

were unambiguous in terms of their ICE presence/absence patterns (**Figure S12A, B**). To

confirm this model prediction, we compared the distribution of the average frequency of NS

SNVs between ICP1-positive and ICP1-negative samples. We found that NS SNV frequency

was significantly higher in ICP1-positive samples when the ICE was not detected (Wilcoxon test,

P=0.0094) (**Figure 4C**). This effect was strongest for NS mutations, making them the likely

targets of ICP1 selection. The slight (not significant, P>0.05) trends for synonymous and

intergenic SNVs could be due to genetic linkage with nonsynonymous mutations. Together,

these results suggest that, in the absence of detectable ICE-encoded phage resistance, ICP1

selects for non-synonymous point mutations that may confer alternative modes of resistance.

Figure 4. ICP1 selects for non-synonymous point mutations in the *V. cholerae* genome in the absence of ICE. (A) Results of a GLMM modeling high frequency nonsynonymous SNV counts as a function of *V. cholerae* (*Vc*) and ICP1 standardized relative abundances. In the bottom panel, shades of gray indicate *Vc* relative abundance at the mean or +/- 1 standard deviation (SD) across samples. (B) GAM output with the mean mutation frequency as a function of the interaction between ICP1, ICE and mutation type (non-synonymous; NS, synonymous; S, or intergenic; I). Significant effects are shown with a star (*P*<0.05). The model was fit using 130 samples that passed the post-Instrain filter (Methods). (C) Boxplots of mutation frequency in the presence or absence of ICP1 and/or ICEs. The single significant comparison is indicated with a star (Wilcoxon test, *P*=0.0094). Boxplots include 130 samples, of which 32 are ICP1+ (ICP1>=0.1% of reads) and 98 are ICP- (ICP1<0.1% of reads). The solid horizontal line is the median and the boxed area is the interquartile range.

To explore which point mutations might be specifically associated with phage resistance,

we focused on the *V. cholerae* genes that accumulated repeated mutations under ICP1

pressure. We previously observed that the secreted hemolysin toxin gene *hlyA* was mutated

more often than expected by chance in cholera patients, but we were unable to identify an

underlying selective pressure (Levade et al. 2021). In the larger sample reported here, *hlyA* was

among the genes most frequently mutated in patients with relatively high *Vc* to ICP1 ratios

(**Table S9**). This gene, along with others mutated at either high or low levels of phage (**Table**

S9; Table S10) – including several with membrane or virulence-related functions – provide

candidate phage resistance mechanisms that can be explored in future experiments.

CONCLUSION

The tripartite interactions between pathogens, phage, and antibiotics have been studied in the

laboratory, *in silico* with mathematical models, and to a lesser extent in the field, but how these

factors interact during human infection remains an open question. Our objective was to

characterize these interactions in the context of cholera. We analyzed more than 300 stool

metagenomes from symptomatic patients enrolled at hospital admission across Bangladesh

during an entire seasonal outbreak period. We discovered that higher levels of phage (ICP1)

relative to *V. cholerae* were associated with mild dehydration, providing a biomarker of disease

severity. The presence of phage alone did not strongly correlate with severity. However, the

ratio of phage to *V. cholerae* did strongly correlate with severity, suggesting that "effective

predation" is a key metric to be used when considering phage as a biomarker.

As predicted by laboratory experiments showing that phage resistance elements on

ICEs can protect against ICP1 (LeGault et al. 2021), we found that patient samples without

detectable ICEs were associated with higher phage to *V. cholerae* ratios. While we have

previously reported loss of ICE genes within cholera patients (Levade et al. 2017), lack of ICE

detection in metagenomes could also be explained by colonization of a strain lacking the ICE or

encoding an ICE variant with low similarity to those previously sequenced. In the absence of

detectable ICE, phage (ICP1) were associated with increased rates of nonsynonymous point

mutations in the *V. cholerae* genome. Many of these mutations likely arose by hypermutation,

which generates deleterious mutations – but may also rapidly confer phage resistance, as

observed in experiments with *Pseudomonas fluorescens (Pal et al. 2007)*. *V. cholerae* can

evolve resistance to phage ICP2 within patients, yet the resistance mutations in a surface

protein reduce its potential for onward transmission (K. S. Seed et al. 2014). Whether the

mutations associated with ICP1 in our study mediate similar fitness tradeoffs for *V. cholerae*

remains unknown. Antibiotic exposure was not associated with increased rates of point

mutations, but was associated with less *V. cholerae* and less severe disease. Azithromycin

appeared to be particularly effective at suppressing *V. cholerae* and was not associated with

any known resistance genes in metagenomes. By contrast, ciprofloxacin exposure was

associated with several known resistance genes and was less effective at suppressing *V.*

cholerae. These predictions can be tested in genome-wide association studies, among other

approaches.

Our study has several limitations which provide opportunities for future research. First,

samples were collected at a single time point (hospital admission) which allows us to establish

statistical correlations, but in the absence of time-series or interventional experiments, we

cannot infer causality. Notably, our results are consistent with ICP1 suppressing *V. cholerae*

and reducing disease severity, but time-series studies of individual patients, or randomized

controlled trials, will be needed to show causality. Second, shotgun metagenomic sequencing

provides rich data on bacteria and phages in the gut microbiome, but is limited to the most

abundant taxa, such as ICP1 and *V. cholerae*. Phages ICP2 and ICP3 were less prevalent in

our study population, and less abundant within patients, making it challenging to infer their

interactions with *V. cholerae* and associations with disease severity. Short read sequencing also

made it challenging to link ICE variants with *V. cholerae* genomes, so we relied on read

mapping approaches. From a clinical perspective, quality of assessments of dehydration may

have varied by provider and location which could represent potential confounders and reduce

effect sizes. Finally, our study lacked information about host genetics or immunity, which also

contribute to disease outcomes (Harris et al. 2008; Nelson et al. 2009). Future studies

combining rich patient metadata, time series, clinical interventions, long-read metagenomics,

and isolate genome sequencing will complement and expand upon our study.

Despite these limitations, our study implicates both phages and antibiotics as

determinants of cholera disease severity, and paves the way for future enquiries into their

interacting roles in disease progression and recovery. We propose a hierarchy of selective

pressures acting on *V. cholerae* in the gut. In the absence of resistance genes, antibiotics are

effective at suppressing cholera and reducing disease severity. In the absence of effective

antibiotics, lytic phages suppress *V. cholerae* when the bacteria do not encode phage

resistance in the ICE. Finally, in the absence of ICE-encoded resistance, phages may select for

point mutations conferring phage resistance and potentially longer-term fitness consequences.

MATERIALS AND METHODS

Ethics Statement. The samples analyzed were collected within the confines of two previously

published IRB approved clinical studies conducted in Bangladesh: (i) The mHealth Diarrhea

Management (mHDM) cluster randomized controlled trial (IEDCR IRB/2017/10; icddr,b

ERC/RRC PR-17036; University of Florida IRB 201601762) (Khan et al. 2020a). (ii) The

National Cholera Surveillance (NCS) study (icddr,b ERC/RRC PR-15127) (Khan et al. 2020b);

See supplementary materials for further details.

Study Design. The study design was a prospective longitudinal study of patients presenting

with diarrhoeal disease at five Bangladesh Ministry of Health and Family Welfare district

hospitals (both mHDM and NCS sites) and two centralized NCS hospitals (BITID; icddr,b) from

March 2018 to December 2018. See supplementary materials.

Sample collection. Stool samples were collected at hospital admission. Aliquots for transport

and subsequent culture were stabbed into Cary-Blair transport media; aliquots for molecular

analysis were preserved in RNAlater. See supplementary materials.

Microbiological and molecular analysis. Culture was performed via standard methods

(Balows 2003); total nucleic acid (tNA) was extracted from the RNAlater preserved samples

using standard methods. Cholera samples for subsequent metagenomic analysis were identified

by screening all samples by culture for *V. cholerae*. Among culture negative samples, a random

10% of the remaining samples were screened for *V. cholerae* specific phage (ICP1, 2, 3) by

PCR using total nucleic acid (tNA) extracts. From samples positive by culture or PCR,

sequencing libraries were prepared using the NEB Ultra II shotgun kit and sequenced on

illumina NovaSeq 6000 S4, pooling 96 samples per lane, yielding a mean of >30 million paired-

end 150bp reads per sample.

Antibiotic detection by liquid chromatography mass spectrometry (LC-MS/MS). Those

cholera samples identified for metagenomic analysis were also analyzed by qualitative and

quantitative LC-MS/MS for antibiotics (Alexandrova et al. 2019; Creasy-Marrazzo et al. 2022).

While the target list for the qualitative analyses was broad, the list for the quantitative analyses

was narrow: Ciprofloxacin, Doxycycline/Tetracycline, and Azithromycin. Standard curves were

made for each quantitative target by preparing a dilution series of the three native and isotopic

forms of the quantitative targets (Ciprofloxacin, Doxycycline, Azithromycin); for quantitative LC-

MS/MS, clinical samples were spiked with the isotopes as internal references. See

supplementary materials.

Metagenomic data analysis. We taxonomically classified short reads using Kraken2 (Wood,

Lu, and Langmead 2019) and Bracken v.2.5 (Lu et al. 2017). Reads were assembled using

MEGAHIT v.1.2.9 (Li et al. 2015) and binned with DAS tool (Sieber et al. 2018). To characterize

diversity within *V. cholerae*, we used StrainGE (van Dijk et al. 2022) and InStrain v.1.5.7 (Olm et

al. 2021). To identify antibiotic resistance genes in metagenomes, we used deepARG v 1.0.2

(Arango-Argoty et al. 2018). See supplementary materials for details.

Statistical analyses. Statistics and visualizations were done in R studio version 1.2.5042. See

supplementary materials for details.

Data availability. All sequencing data are deposited in the NCBI SRA under BioProject

PRJNA976726. See supplementary materials for further information.

Code availability. Computer code for this paper will be available at

https://github.com/Naima16/Cholera-phage-antibiotics.

Acknowledgements

We thank the patients for participating in this study and the clinical and laboratory teams that

collected the samples. We are grateful to S. Flora and colleagues at the Institute of

Epidemiology, Disease Control and Research (IEDCR), Ministry of Health and Family Welfare,

Government of Bangladesh who collaborated on the original clinical studies in which the

samples analyzed herein were collected. We are also grateful to R. Autrey and K. Berquist for

their administrative expertise at the University of Florida. AIK and FQ were the principal

investigators in Bangladesh and PIs of the ERC/RRC approvals at the icddr,b. This collective

research infrastructure and support was invaluable to the success of this study. We thank

members of the Nelson, Khan, Qadri and Shapiro labs for discussions that improved the

manuscript.

Financial Support

This work was supported by the National Institutes of Health grants to EJN [R21TW010182] and

KBB [S10 OD021758-01A1] and internal support from the Emerging Pathogens Institute at the

University of Florida and the Departments of Pediatrics/ Children's Miracle Network (Florida).

BJS and NM were supported by a Canadian Institutes for Health Research Project Grant. AC

was supported by a Postdoctoral Mobility Fellowship of the Swiss National Science Foundation

[P500PB_214356].

Disclaimer

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Potential conflicts of interest. All authors: No reported conflicts.

REFERENCES

Alexandrova, Ludmila, Farhana Haque, Patricia Rodriguez, Ashton C. Marrazzo, Jessica A. Grembi, Vasavi Ramachandran, Andrew J. Hryckowian, et al. 2019. "Identification of Widespread Antibiotic Exposure in Cholera Patients Correlates with Clinically Relevant Microbiota Changes." *The Journal of Infectious Diseases*, June. https://doi.org/10.1093/infdis/jiz299.

Ali, Mohammad, Allyson R. Nelson, Anna Lena Lopez, and David A. Sack. 2015. "Updated Global Burden of Cholera in Endemic Countries." *PLoS Neglected Tropical Diseases* 9 (6): e0003832.

Andrews, Jason R., Daniel T. Leung, Shahnawaz Ahmed, Mohammed Abdul Malek, Dilruba Ahmed, Yasmin Ara Begum, Firdausi Qadri, Tahmeed Ahmed, Abu Syed Golam Faruque, and Eric J. Nelson. 2017. "Determinants of Severe Dehydration from Diarrheal Disease at Hospital Presentation: Evidence from 22 Years of Admissions in Bangladesh." *PLoS Neglected Tropical Diseases* 11 (4): e0005512.

Arango-Argoty, Gustavo, Emily Garner, Amy Pruden, Lenwood S. Heath, Peter Vikesland, and Liqing Zhang. 2018. "DeepARG: A Deep Learning Approach for Predicting Antibiotic Resistance Genes from Metagenomic Data." *Microbiome* 6 (1): 23.

Balows, Albert. 2003. "Manual of Clinical Microbiology 8th Edition: P. R. Murray, E. J. Baron, J. H. Jorgenson, M. A. Pfaller, and R. H. Yolken, Eds., ASM Press, 2003, 2113 Pages, 2 Vol, 2003 + Subject & Author Indices, ISBN: 1-555810255-4, US$ 189.95." *Diagnostic Microbiology and Infectious Disease* 47 (4): 625.

Beaber, John W., Bianca Hochhut, and Matthew K. Waldor. 2002. "Genomic and Functional Analyses of SXT, an Integrating Antibiotic Resistance Gene Transfer Element Derived from Vibrio Cholerae." *Journal of Bacteriology* 184 (15): 4259–69.

Boyd, Caroline M., Angus Angermeyer, Stephanie G. Hays, Zachary K. Barth, Kishen M. Patel, and Kimberley D. Seed. 2021. "Bacteriophage ICP1: A Persistent Predator of Vibrio Cholerae." *Annual Review of Virology* 8 (1): 285–304.

Carr, Alex, Christian Diener, Nitin S. Baliga, and Sean M. Gibbons. 2019. "Use and Abuse of Correlation Analyses in Microbial Ecology." *The ISME Journal* 13 (11): 2647–55.

"Cholera – Global Situation." WHO Report. 2023. https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON437.

Creasy-Marrazzo, Ashton, Morteza M. Saber, Manasi Kamat, Laura S. Bailey, Lindsey Brinkley, Emilee Cato, Yasmin Begum, et al. 2022. "Genome-Wide Association Studies Reveal Distinct Genetic Correlates and Increased Heritability of Antimicrobial Resistance in Vibrio Cholerae under Anaerobic Conditions." *Microbial Genomics* 8 (12). https://doi.org/10.1099/mgen.0.000905.

Dalsgaard, A., A. Forslund, N. V. Tam, D. X. Vinh, and P. D. Cam. 1999. "Cholera in Vietnam: Changes in Genotypes and Emergence of Class I Integrons Containing Aminoglycoside Resistance Gene Cassettes in Vibrio Cholerae O1 Strains Isolated from 1979 to 1996." *Journal of Clinical Microbiology* 37 (3): 734–41.

Das, Bhabatosh, Jyoti Verma, Pawan Kumar, Amit Ghosh, and Thandavarayan Ramamurthy. 2020. "Antibiotic Resistance in Vibrio Cholerae: Understanding the Ecology of Resistance Genes and Mechanisms." *Vaccine* 38 Suppl 1 (February): A83–92.

David, Lawrence A., Ana Weil, Edward T. Ryan, Stephen B. Calderwood, Jason B. Harris, Fahima Chowdhury, Yasmin Begum, Firdausi Qadri, Regina C. LaRocque, and Peter J. Turnbaugh. 2015. "Gut Microbial Succession Follows Acute Secretory Diarrhea in Humans." *mBio* 6 (3): e00381–15.

D'Herelle, F., and R. H. Malone. 1927. "A Preliminary Report of Work Carried out by the Cholera Bacteriophage Enquiry." *Indian Medical Gazette* 62 (11): 614–16.

Dijk, Lucas R. van, Bruce J. Walker, Timothy J. Straub, Colin J. Worby, Alexandra Grote, Henry L. Schreiber 4th, Christine Anyansi, et al. 2022. "StrainGE: A Toolkit to Track and Characterize Low-Abundance Strains in Complex Microbial Communities." *Genome Biology* 23 (1): 74.

Dromigny, Jacques-Albert, Olivat Rakoto-Alson, Davidra Rajaonatahina, René Migliani, Justin Ranjalahy, and Philippe Mauclére. 2002. "Emergence and Rapid Spread of Tetracycline-Resistant Vibrio Cholerae Strains, Madagascar." *Emerging Infectious Diseases* 8 (3): 336–38.

Faruque, Shah M., M. Johirul Islam, Qazi Shafi Ahmad, A. S. G. Faruque, David A. Sack, G. Balakrish Nair, and John J. Mekalanos. 2005. "Self-Limiting Nature of Seasonal Cholera Epidemics: Role of Host-Mediated Amplification of Phage." *Proceedings of the National Academy of Sciences of the United States of America* 102 (17): 6119–24.

Faruque, Shah M., Iftekhar Bin Naser, M. Johirul Islam, A. S. G. Faruque, A. N. Ghosh, G. Balakrish Nair, David A. Sack, and John J. Mekalanos. 2005. "Seasonal Epidemics of Cholera Inversely Correlate with the Prevalence of Environmental Cholera Phages." *Proceedings of the National Academy of Sciences of the United States of America* 102 (5): 1702–7.

Harris, Jason B., Regina C. LaRocque, Fahima Chowdhury, Ashraful I. Khan, Tanya Logvinenko, Abu S. G. Faruque, Edward T. Ryan, Firdausi Qadri, and Stephen B. Calderwood. 2008. "Susceptibility to Vibrio Cholerae Infection in a Cohort of Household Contacts of Patients with Cholera in Bangladesh." Edited by Albert I. Ko. *PLoS Neglected Tropical Diseases* 2 (4): e221.

Islam, M. S., A. K. Siddique, A. Salam, K. Akram, R. N. Majumdar, K. Zaman, N. Fronczak, and S. Laston. 1995. "Microbiological Investigation of Diarrhoea Epidemics among Rwandan Refugees in Zaire." *Transactions of the Royal Society of Tropical Medicine and Hygiene* 89 (5): 506.

Jaiswal, Abhishek, Hemanta Koley, Amit Ghosh, Anup Palit, and Banwarilal Sarkar. 2013. "Efficacy of Cocktail Phage Therapy in Treating Vibrio Cholerae Infection in Rabbit Model." *Microbes and Infection / Institut Pasteur* 15 (2): 152–56.

Jensen, Mark A., Shah M. Faruque, John J. Mekalanos, and Bruce R. Levin. 2006. "Modeling the Role of Bacteriophage in the Control of Cholera Outbreaks." *Proceedings of the National Academy of Sciences of the United States of America* 103 (12): 4652–57.

Jolivet-Gougeon, Anne, Bela Kovacs, Sandrine Le Gall-David, Hervé Le Bars, Latifa Bousarghin, Martine Bonnaure-Mallet, Bernard Lobel, François Guillé, Claude-James Soussy, and Peter Tenke. 2011. "Bacterial Hypermutation: Clinical Implications." *Journal of Medical Microbiology* 60 (Pt 5): 563–73.

Khan, Ashraful I., Jasmine A. Mack, M. Salimuzzaman, Mazharul I. Zion, Hasnat Sujon, Robyn L. Ball, Stace Maples, et al. 2020a. "Electronic Decision Support and Diarrhoeal Disease Guideline Adherence (mHDM): A Cluster Randomised Controlled Trial." *The Lancet. Digital Health* 2 (5): e250–58.

Khan, Ashraful I., Md Mahbubur Rashid, Md Taufiqul Islam, Mokibul Hassan Afrad, M. Salimuzzaman, Sonia Tara Hegde, Md Mazharul I. Zion, et al. 2020b. "Epidemiology of Cholera in Bangladesh: Findings From Nationwide Hospital-Based Surveillance, 2014-2018." *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America* 71 (7): 1635–42.

Lassalle, Florent, Salah Al-Shalali, Mukhtar Al-Hakimi, Elisabeth Njamkepo, Ismail Mahat Bashir, Matthew J. Dorman, Jean Rauzier, et al. 2022. "Genomic Epidemiology of the Cholera Outbreak in Yemen Reveals the Spread of a Multi-Drug Resistance Plasmid between Diverse Lineages of Vibrio Cholerae." *bioRxiv*. https://doi.org/10.1101/2022.08.24.504966.

LeGault, Kristen N., Stephanie G. Hays, Angus Angermeyer, Amelia C. McKitterick, Fatema-Tuz

Johura, Marzia Sultana, Tahmeed Ahmed, Munirul Alam, and Kimberley D. Seed. 2021. "Temporal Shifts in Antibiotic Resistance Elements Govern Phage-Pathogen Conflicts." *Science* 373 (6554). https://doi.org/10.1126/science.abg2166.

Levade, Inès, Ashraful I. Khan, Fahima Chowdhury, Stephen B. Calderwood, Edward T. Ryan, Jason B. Harris, Regina C. LaRocque, et al. 2021. "A Combination of Metagenomic and Cultivation Approaches Reveals Hypermutator Phenotypes within Vibrio Cholerae-Infected Patients." *mSystems* 6 (4): e0088921.

Levade, Inès, Yves Terrat, Jean-Baptiste Leducq, Ana A. Weil, Leslie M. Mayo-Smith, Fahima Chowdhury, Ashraful I. Khan, et al. 2017. "Vibrio Cholerae Genomic Diversity within and between Patients." *Microbial Genomics* 3 (12). https://doi.org/10.1099/mgen.0.000142.

Li, Dinghua, Chi-Man Liu, Ruibang Luo, Kunihiko Sadakane, and Tak-Wah Lam. 2015. "MEGAHIT: An Ultra-Fast Single-Node Solution for Large and Complex Metagenomics Assembly via Succinct de Bruijn Graph." *Bioinformatics* 31 (10): 1674–76.

Lu, Jennifer, Florian P. Breitwieser, Peter Thielen, and Steven L. Salzberg. 2017. "Bracken: Estimating Species Abundance in Metagenomics Data." *PeerJ Computer Science* 3 (January): e104.

Monir, Md Mamun, Mohammad Tarequl Islam, Razib Mazumder, Dinesh Mondal, Kazi Sumaita Nahar, Marzia Sultana, Masatomo Morita, et al. 2023. "Genomic Attributes of Vibrio Cholerae O1 Responsible for 2022 Massive Cholera Outbreak in Bangladesh." *Nature Communications* 14 (1): 1154.

Nelson, Eric J., Ashrafuzzaman Chowdhury, James Flynn, Stefan Schild, Lori Bourassa, Yue Shao, Regina C. LaRocque, Stephen B. Calderwood, Firdausi Qadri, and Andrew Camilli. 2008. "Transmission of Vibrio Cholerae Is Antagonized by Lytic Phage and Entry into the Aquatic Environment." Edited by Frederick M. Ausubel. *PLoS Pathogens* 4 (10): e1000187.

Nelson, Eric J., Jason B. Harris, J. Glenn Morris Jr, Stephen B. Calderwood, and Andrew Camilli. 2009. "Cholera Transmission: The Host, Pathogen and Bacteriophage Dynamic." *Nature Reviews. Microbiology* 7 (10): 693–702.

Nelson, Eric J., Danielle S. Nelson, Mohammed A. Salam, and David A. Sack. 2011. "Antibiotics for Both Moderate and Severe Cholera." *The New England Journal of Medicine* 364 (1): 5–7.

Olm, Matthew R., Alexander Crits-Christoph, Keith Bouma-Gregson, Brian A. Firek, Michael J. Morowitz, and Jillian F. Banfield. 2021. "inStrain Profiles Population Microdiversity from Metagenomic Data and Sensitively Detects Shared Microbial Strains." *Nature Biotechnology* 39 (6): 727–36.

Pal, Csaba, María D. Maciá, Antonio Oliver, Ira Schachar, and Angus Buckling. 2007. "Coevolution with Viruses Drives the Evolution of Bacterial Mutation Rates." *Nature* 450 (7172): 1079–81.

Pasricha, C. L., A. J. H. de Monte, and E. G. O'Flynn. 1936. "Bacteriophage in the Treatment of Cholera." *Indian Medical Gazette* 71 (2): 61–68.

Rivard, Nicolas, Rita R. Colwell, and Vincent Burrus. 2020. "Antibiotic Resistance in Vibrio Cholerae: Mechanistic Insights from IncC Plasmid-Mediated Dissemination of a Novel Family of Genomic Islands Inserted at trmE." *mSphere* 5 (4). https://doi.org/10.1128/mSphere.00748-20.

Seed, K. D., K. L. Bodi, A. M. Kropinski, H. W. Ackermann, S. B. Calderwood, F. Qadri, and A. Camilli. 2011. "Evidence of a Dominant Lineage of Vibrio Cholerae-Specific Lytic Bacteriophages Shed by Cholera Patients over a 10-Year Period in Dhaka, Bangladesh." *mBio* 2 (1): e00334–10 – e00334–10.

Seed, K. S., M. Yen, B. J. Shapiro, I. J. Hilaire, R. C. Charles, J. E. Teng, L. C. Ivers, J. Boncy, J. B. Harris, and A. Camilli. 2014. "Evolutionary Consequences of Intra-Patient Phage Predation on Microbial Populations." *eLife* 3: e03497.

Sieber, Christian M. K., Alexander J. Probst, Allison Sharrar, Brian C. Thomas, Matthias Hess,

Susannah G. Tringe, and Jillian F. Banfield. 2018. "Recovery of Genomes from Metagenomes via a Dereplication, Aggregation and Scoring Strategy." *Nature Microbiology* 3 (7): 836–43.

Summers, W. C. 1993. "Cholera and Plague in India: The Bacteriophage Inquiry of 1927-1936." *Journal of the History of Medicine and Allied Sciences* 48 (3): 275–301.

Waldor, Matthew K., Helmut Tschäpe, and John J. Mekalanos. 1996. "A New Type of Conjugative Transposon Encodes Resistance to Sulfamethoxazole, Trimethoprim, and Streptomycin in Vibrio Cholerae O139." *Journal of Bacteriology* 178 (14): 4157–65.

Weill, François-Xavier, Daryl Domman, Elisabeth Njamkepo, Cheryl Tarr, Jean Rauzier, Nizar Fawal, Karen H. Keddy, et al. 2017. "Genomic History of the Seventh Pandemic of Cholera in Africa." *Science* 358 (6364): 785–89.

Wood, Derrick E., Jennifer Lu, and Ben Langmead. 2019. "Improved Metagenomic Analysis with Kraken 2." *Genome Biology* 20 (1): 257.

Yen, Minmin, Lynne S. Cairns, and Andrew Camilli. 2017. "A Cocktail of Three Virulent Bacteriophages Prevents Vibrio Cholerae Infection in Animal Models." *Nature Communications* 8 (February): 14187.

Zahid, M. S. H., S. M. N. Udden, A. S. G. Faruque, S. B. Calderwood, J. J. Mekalanos, and S. M. Faruque. 2008. "Effect of Phage on the Infectivity of Vibrio Cholerae and Emergence of Genetic Variants." *Infection and Immunity* 76 (11): 5266–73.