Title: Phage predation is a biomarker for disease severity and shapes pathogen genetic diversity in cholera patients

Authors: N. Madi[1,2], E. T. Cato[3], Md. Abu Sayeed[3], A. Creasy-Marrazzo[3], A. Cuénod[1,2], K. Islam[4], Md. I.Ul. Khabir[4], Md. T. R. Bhuiyan[4], Y. Begum[4], E. Freeman[3], A. Vustepalli[3], L. Brinkley[3], Kamat[4], L. S. Bailey[5], K. B. Basso[5], F. Qadri[4], A. I. Khan[4*], B. J. Shapiro[1,2,6*], E. J. Nelson[3*]

Affiliations:

[1] Department of Microbiology & Immunology, McGill University, Montréal, Canada
[2] McGill Genome Centre, McGill University, Montréal, Canada
[3] Departments of Pediatrics and Environmental and Global Health, University of Florida, Gainesville, FL, USA
[4] Infectious Diseases Division (IDD) & Nutrition and Clinical Services Division (NCSD), International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh
[5] Department of Chemistry, University of Florida, Gainesville, FL, USA
[6] McGill Centre for Microbiome Research, McGill University, Montréal, Canada

[*] Corresponding authors

Email: ashrafk@icddrb.org; jesse.shapiro@mcgill.ca; eric.nelson@ufl.edu

The PDF file includes:

Index
Materials and methods
Results
Figures: S1 to S19
Tables: S1 to S12
References

Keywords: Cholera, diarrhoea, diarrhea, *Vibrio cholerae*, bacteriophage, phage, phage resistance, antibiotics, antimicrobial resistance, AMR, predator, prey, metagenomics, co-evolution, evolution

Enumerations: Abstract: 194 words. Manuscript 4594 words (less figures). Inserts 5. Refs. 58.

Abstract

A century has passed since the discovery that virulent bacteriophages are associated with improved survival of cholera patients. Despite an increasingly detailed picture of the molecular mechanisms that govern phage-bacterial interactions, we lack an understanding of how these interactions impact disease severity. Here we report a year-long, nation-wide study of diarrheal disease patients in Bangladesh. Among the subset of cholera patients, we quantified *Vibrio cholerae* (prey) and its virulent phages (predators) using metagenomics and quantitative PCR, while accounting for antibiotic exposure using quantitative mass spectrometry. Virulent phage (ICP1) and antibiotics suppressed *V. cholerae* to varying degrees and were inversely associated with severe dehydration; aspects of these effects were dependent on resistance mechanisms. In the absence of anti-phage defenses, predation was 'effective' with a high predator to prey ratio that correlated with increased genetic diversity among the prey. In the presence of anti-phage defenses, predation was 'ineffective' with a low predator to prey ratio that correlated with increased genetic diversity among the predators. Our results link phage predation to disease severity, support phage-bacteria coevolution within patients, and suggest the ratio of phage to pathogen can serve as a biomarker for clinical, diagnostic, and epidemiologic applications.

One Sentence Summary

A nation-wide prospective study of cholera patients in Bangladesh identifies effective phage predation and antibiotics as markers of disease severity and selective pressures for cognate resistance mechanisms.

MAIN TEXT

Introduction

The secretory diarrheal disease cholera is caused by the Gram negative bacterium *Vibrio cholerae* (*Vc*) and can progress to life-threatening hypovolemic shock in less than 12 hours (*1*). Cholera remains a major public health problem due to inadequate sanitation and restricted access to safe drinking water. Global estimates of cholera burden are 1.3-4.0 million cases and 21,000 to 143,000 deaths annually (*2*). In 2023, there were over 30 countries with active outbreaks necessitating the WHO to escalate the response to its highest level (*3*). Rehydration is the primary life-saving intervention for cholera patients. With effective rehydration, mortality rates fall from over 20% to less than 1%. Antibiotics reduce stool volume and duration of diarrhea but are considered supportive and reserved for patients with more severe disease to reduce selection for antibiotic resistance (*4-8*). Nevertheless, antibiotic-resistant *Vc* have emerged globally (*6, 9, 10*). Mechanisms of resistance are diverse and reside in the core genome, plasmids of the incompatibility type C (*11, 12*) and on mobile genetic elements, including the ~100 kb SXT integrative conjugative element (ICE) that can harbor resistance to sulfamethoxazole and trimethoprim, ciprofloxacin (*qnr$_{vc}$*), trimethoprim (*dfra31*) and streptomycin (*aph(6)*) (*13-16*). Recent work has also shown that the ICE can encode diverse phage resistance mechanisms, with distinct hotspots of gene content variation encoding different resistance genes (*17*).

With rising levels of antibiotic resistance, virulent bacteriophages are a promising alternative or complementary therapy to antibiotics. Phages and bacteria coevolve, with each partner selecting for adaptations in the other that generates genetic diversity for both predator and prey (*18, 19*). Given the extensive arsenal of resistance and counter-resistance mechanisms among *V. cholerae* and its phages, coevolution between predator and prey occurs (*17, 19-24*). Yet it remains unclear how these interactions impact disease severity during natural infection, with and without antibiotic exposure. Virulent phages specifically targeting *Vc* include

ICP1 (*Myoviridae*), ICP2 (*Podoviridae*), and ICP3, which (*Podoviridae*) (*21, 25, 26*); these

phages are most common and considered to be found only in symptomatic or asymptomatic

patients during acute infection or the convalescent period. The first clinical trials of phage

therapy occurred during the Cholera Phage Inquiry from 1927 to 1936 (*27, 28*). In a *proto*

randomized controlled trial, the Inquiry found the odds of mortality were reduced by 58% among

those with phage therapy, with an absolute reduction in mortality of 10% (95% CI 0.27-0.64)

(*29*) (reanalysis by E. J. Nelson). Despite these early findings, there is a lack of evidence in the

modern era that links phage predation with disease severity during natural infection in humans.

However, indirect studies do support a link: Environmentally, virulent phages in the aquatic

environment were negatively correlated with cholera cases in Bangladesh over time, suggesting

a role for phages in influencing epidemic dynamics (*30*). Clinically, a higher percentage of

cholera patients shed virulent phage towards the end of an outbreak (*31*). Computationally,

models predict that phage predation can dampen cholera outbreaks (*32*). Experimentally,

animal studies found phage predation was inversely associated with colonization and severity

(*25, 33-35*). The key unanswered question is if and how virulent phages, antibiotics, and

bacterial evolution interact to impact infection in a meaningful manner for clinicians, public

health officials, and most importantly patients.

To address this question, we conducted a national prospective longitudinal study in the

cholera endemic setting of Bangladesh. Stool samples were collected at hospital admission

from diarrheal disease patients and screened for *Vc*, antibiotics, and prevalent virulent cholera

phages ICP1, 2 and 3. We hypothesized that: (i) virulent phages and antibiotics would suppress

Vc and be inversely associated with disease severity, (ii) this suppression would be lifted for *Vc*

encoding anti-phage genes on ICEs, (iii) phages would select for other potential resistance

mutations in the absence of ICE-encoded resistance, and (iv) phages would be under selection

to escape suppression by ICEs. We provide broad support for these hypotheses, paving the

way for mechanistic experimental studies, the development of the phage:pathogen ratio as a

biomarker of disease severity, and further dissection of the longer-term consequences of phage

predation on pathogen evolution.

Results and Discussion

Study overview. A total of 2574 stool samples were collected from enrolled participants

admitted for diarrheal disease at seven hospitals across Bangladesh from March to December

2018; collection continued to April 2019 at one site (icddr,b). Three groups of cholera samples

were analyzed: (i) *Vc* culture-positive (282/2574; 10.9%)*,* (ii) *Vc* culture-negative but phage

(ICP1,2,3) PCR-positive (127/2292; 5.5%; 80 included; 47 excluded for DNA < 1ng/µl), and (iii) a

random 10% of *Vc* culture-negative and phage PCR-negative samples that were *Vc* PCR-

positive (14.8%; 37/250; 27 included; 10 excluded for DNA < 1ng/µl)*.* Stool metagenomes were

successfully sequenced from 88.4% (344/389) of samples from these three groups, 35% of

which were from the icddr,b site. Detection rates by PCR were low for ICP2 (0.5%) and ICP3

(1.6%) in culture-negative samples and were not targeted for sequencing. Based on

metagenomic read mapping to a taxonomic database, detection rates for *Vc,* ICP1, ICP2, and

ICP3 were 55%, 18%, 1% and 8%, respectively. These ranked detection rates were supported

by an analysis of *Vc* phages identified in metagenomic assemblies. As expected, the prophage

encoding the cholera toxin (CTXphi) was identified in most assemblies, with ICP1 being the next

most prevalent. While some additional putative phages were detected, none were prevalent

enough to merit further analysis (**Fig S1**). For both *Vc* and ICP1, relative abundances in

metagenomes correlated with absolute quantification by qPCR (**Fig S2**). Five antibiotics (**Table**

S2) were prioritized for detection in stool using liquid chromatography-mass spectrometry

(LCMS); of these, azithromycin, ciprofloxacin and doxycycline were quantified.

Metagenomic correlates of disease severity and succession. At hospital admission, we

expected patients to present at different stages of disease, with an ecological succession of *Vc*

followed by the facultative anaerobe *Escherichia coli,* then by a healthy flora of mostly anaerobic

bacteria (*36*). We hypothesized that these stages of succession would be associated with

different disease severity. As expected, we identified *Vc* as an indicator species of severe

dehydration. *Vc* was relatively more abundant in patients with severe dehydration, while two

Bifidobacterium species, *E. coli*, and ICP1 were indicators of mild dehydration (**Fig. 1A, Table**

S3). ICP3 was an indicator of severe dehydration despite being less frequently detected in our

study (28 samples with >0.1% ICP3 reads, compared to 61 samples with >0.1% ICP1). This

suggests that different phages can have contrasting disease associations. Given the robust

sample size for ICP1, we focus on ICP1 for subsequent analyses.

The distribution of ICP1 relative abundance was variable and less clearly associated

with dehydration status than *Vc* (**Fig. S3**). Deeper investigation revealed that it was not simply

the presence of phage that mattered. Rather, the ratio of ICP1 to *Vc* was the critical

determinant. Higher ratios were inversely associated with dehydration severity (**Fig. 1B**); the

same results were obtained using qPCR instead of metagenomics to quantify the ratio (**Fig S4**).

This simple ratio is a marker of 'effective' phage predation that could be used in clinical,

diagnostic and epidemiological applications.

These observations suggest that ICP1 plays a role in modulating disease severity, or

could be associated with later stages of disease, when patients are recovering. Consistent with

the former possibility, ICP1 was previously associated with early, rather than late stages of

disease, peaking on the first day of infection in a time series of cholera patients (*36*). Given that

we collected only one sample per patient (at hospital admission), we were limited in

disentangling these possibilities. Nevertheless, we recorded self-reported duration of diarrhea,

providing an approximate proxy for disease progression. We found that higher relative

abundances of ICP1 were associated with mild dehydration at early stages of disease (duration

of diarrhea <72h) but not at late stages (**Fig. S3B and D**). We therefore put forth a model in

which ICP1 suppresses *Vc* at early disease stages in a way that reduces disease severity by

suppressing *Vc*, or by other indirect mechanisms. In either case, ICP1 serves as a meaningful

biomarker of disease severity.

Fig. 1. Dehydration severity is inversely associated with higher ICP1:*Vc* ratios in stool
metagenomes. (A) Relative abundances of the most prevalent species in patients with severe, moderate,
or mild dehydration; these conventions equate to the World Health Organization (WHO) conventions of
severe, some and no dehydration, respectively. Significant indicator species for severe or mild
dehydration are shown in red or blue bold text, respectively ($P<0.05$ in a permutation test with 9999
iterations as implemented in the indicator species function in R). See Table S3 for full indicator species
results. (B) The ICP1:*V. cholerae* ratio is higher in patients with mild dehydration. *P*-values are from a
Kruskal-Wallis test with Dunn's post-hoc test, adjusted for multiple tests using the Benjamini-Hochberg
(BH) method. Only significant *P*-values (<0.05) are shown. Only 323 samples with *V. cholerae*>0% of
metagenomic reads were included, with 165 from severe, 128 from moderate, and 30 from mild cases. A
pseudocount of one was added to the ratio before log transformation. For supporting analyses using
qPCR data, see Fig. S4. In (A) and (B) the solid horizontal line is the median and the boxed area is the
interquartile range. (C) Redundancy analysis (RDA) showing relationships among the seven most
dominant bacterial species and explanatory variables: phages, patient metadata, and antibiotic
concentration. Angles between variables (arrows) reflect their correlations; arrow length is proportional to
effect size. Samples (points) are colored by dehydration severity. All displayed variables are significant
($P<0.05$, permutation test) except ICP2, ICP3, and doxycycline (Table S4). For the RDA: $R^2=0.25$ and
adjusted $R^2=0.184$, permutation test $P = 0.001$. To improve readability, collection date and location are
not shown (see Fig. S6 for these details).
184
Antibiotic exposure is inversely associated with disease severity. To visualize the complex

relationships between disease severity, bacteria, and phage in the context of antibiotic

exposure, we used redundancy analysis (RDA; **Fig. 1C; Table S4**). For simplicity, the seven

most dominant bacterial species identified by principal components analysis were included (**Fig.**

S5). As explanatory variables, we visualized those clinical data with strong effects, chosen by

forward selection beginning with phages and antibiotic concentrations (**Fig. S6**). In accordance

with the indicator species analysis (**Fig. 1A, Table S3**), higher *Vc* relative abundance was

positively correlated with severe dehydration (**Fig. 1C**). ICP1 was moderately associated with

Vc, consistent with a phage's reliance on its host for replication, but less associated with severe

dehydration. The antibiotics azithromycin (AZI) and, to a lesser extent ciprofloxacin (CIP), were

negatively correlated with *Vc* and severe dehydration, suggesting their role in suppressing

cholera infection and disease. Supporting previous reports that AZI suppresses *Vc* (*37*), AZI

was most strongly anticorrelated with *Vc* in our cohort (**Fig. 1C**). We did not identify annotated

antibiotic resistance genes associated with AZI exposure (**Fig. S7**) at established thresholds

(**Table S5**), suggesting that AZI resistance is relatively rare in our cohort. This does not exclude

the possibility that resistance to AZI is controlled by non-canonical resistance genes absent

from standard databases. In contrast, CIP exposure was significantly associated with the

resistance genes *dfrA* and *aph6* (**Fig. S8),** which are both associated with *Vc* in our

metagenomes (**Fig S9**) and have previously been linked with CIP resistance in *Vc (16, 38)*.

Taken together, these results suggest CIP exposure selects for resistance genes within

patients, possibly explaining why CIP may be less effective at suppressing *Vc* than AZI.

Fig. 2. Interactions between *V. cholerae*, phage ICP1, and azithromycin. GAM results, fit with *Vc* as a function of antibiotic concentrations and ICP1 relative abundance in 344 metagenomes. (A) *V. cholerae* declines in relative abundance with higher relative abundance of azithromycin (AZI) in µg/ml. (B) The relationship between ICP1 and *V. cholerae* is affected by AZI concentration (µg/ml); the illustrated AZI concentrations show regular intervals between the minimum (0 µg/ml) and maximum (78 µg/ml) observed values. Both effects of AZI (A) and the ICP1-AZI interaction (B) are significant (Chi-square test, *P*<0.05). For details on GAM outputs see Table S6. For equivalent analyses using qPCR data, see Fig. S10.

Antibiotic exposure suppresses predator-prey dynamics. We next asked if and how

antibiotics interact with phage to suppress *Vc* within patients. To do so, we modeled the

relationships between ICP1, *Vc* and antibiotic exposure within each patient. We fit generalized

additive models (GAMs) of *Vc* (relative abundance from metagenomics or absolute abundance

from qPCR) as a function of ICP1, antibiotic exposure, and their interaction, including

dehydration status as a random effect. We fit GAMs with all antibiotics and their interaction with

ICP1, as well as separate models with each antibiotic, alone or in combination, and compared

them based on their Akaike Information Criterion (AIC; **Tables S6, S7**). The most parsimonious

models (with the lowest AIC) using either metagenomics or qPCR data showed a significant

negative relationship between *Vc* and AZI (**Fig. 2A, S10**), consistent with the RDA results (**Fig.**

1C) and with known patterns of *Vc* suppression by AZI during infection (*37*). The relationship

between *Vc* and ICP1 was quadratic in both metagenomics- and qPCR-based models: at low

ICP1 abundance, the relationship was positive but became negative at higher ICP1 abundance

(**Fig. 2B, S10**). This alternation between positive and negative correlations is consistent with

predator-prey dynamics within infected patients (*39*). However, at higher concentrations of AZI,

the quadratic relationship flattened, effectively suppressing the phage-bacteria interaction. In the

future, these interactions could be interrogated using patient time-series data and laboratory

experiments challenging *Vc* with antibiotics and phages.

Integrative and conjugative elements (ICEs) are associated with phage suppression.

SXTs are approximately 100kb integrative and conjugative elements (ICEs) that are associated

with resistance to both antibiotics and phages and were first discovered in *Vc* (*13*). SXTs have

conserved 'core' genes along with variable 'hotspots' encoding different genes; for example,

hotspot 5 is a ~17kb region associated with phage resistance. At the time of our sampling,

ICEVchInd5 (abbreviated here as *ind5*) and ICEVchInd6 (*ind6*) were the two most prevalent ICE

types in Bangladesh (*17*). These ICEs differ in their anti-phage systems: *ind5* encodes a type 1

bacteriophage exclusion (BREX) system while *ind6* encodes several other different restriction-

modification systems (*17*).

We enumerated ICEs in metagenomes by mapping reads against reference sequences

for *ind5* (NCBI accession GQ463142.1) and *ind6* (accession MK165649.1). An ICE was defined

as present when 90% of its length was covered by at least one metagenomic read (**Fig. S11A**).

We found that 59% (144/244) of samples with *Vc*>0.5% or phages >0.1% of metagenomic

reads contained *ind5*, 10.6% (26/244) contained *ind6*, and 22.1% (54/244) had no detectable

ICE. The lack of ICE detections was not due to the lack of *Vc* in a metagenome because ICE-

negative samples did not contain fewer *Vc* reads (**Fig. S11B**).

Resistance mechanisms on ICEs have been shown to suppress phage *in vitro (17)*, but

their relevance within human infection remains unclear. We found that metagenomes without a

detectable ICE (denoted as ICE-) were associated with higher phage to *Vc* ratios (**Fig. 3**). The

effect was strongest for ICP1, which had the largest sample size (**Fig. 3A**). This observation is

consistent with ICE-encoded mechanisms suppressing phage within patients. Higher ICP1:*Vc*

ratios, which occurred more often in ICE- patients, were also associated with mild dehydration

(**Fig. 3B**). The phage:*Vc* ratios varied according to the precise phage-ICE combination,

suggesting that different ICE variants might confer resistance to different phages (**Fig. 3, Fig.**

S12). The same associations were evident using qPCR-based quantification of phage:bacteria

ratios (**Fig S13**). Together, these results implicate ICEs in suppressing phages during human

infections, complementing and generally confirming the predictions of laboratory experiments

(*17*). That said, the suppression is not complete, and further experiments are needed to dissect

the underlying causal relationships.

Fig. 3. Integrative conjugative elements (ICEs) are associated with lower ICP1:*V. cholerae* ratios in patient metagenomes. (A) Distribution of ICP1:*Vc* ratios across patients with different ICE profiles. (B) The same data as (A) binned into boxplots according to dehydration status. (C) Distribution of phage:*Vc* ratios, including the sum of all phages (ICP1, ICP2, ICP3). (D) The same data as (C) binned into boxplots according to dehydration status. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test adjusted with the Benjamini-Hochberg (BH) method. Only *P*-values < 0.1 are shown. Only samples with appreciable *Vc* or ICP1 were included (224 samples with *Vc*>0.5% or phages >0.1% of metagenomic reads), of which 54 samples were ICE-, 26 were ind6+ and 144 were ind5+. The Y-axes were log10 transformed after adding one to the ratios. The solid horizontal line is the median and the boxed area is the interquartile range. For supporting analyses using qPCR data, see Fig. S13.

Hypermutation generates V. cholerae genetic diversity. In addition to variation in gene

content in ICEs and other mobile elements, we hypothesized that resistance to phages and

antibiotics may have been conferred by point mutations (single nucleotide variants; SNVs) that

existed before or emerged *de novo* during infection. Although we cannot fully exclude mixed

infections by different *Vc* strains as a source of within-patient diversity, we found no evidence for

more than one strain co-infecting a patient in our study population (**Fig. S14**). We previously

found a low level of *Vc* genetic diversity within individual cholera patients (*40*) – on the order of

0-3 detectable SNVs – with the exception of hypermutation events characterized by DNA repair

defects and hundreds of SNVs in the *Vc* genome, primarily transversion mutations (*41*).

Hypermutation generates deleterious mutations, but may also rapidly confer phage resistance,

as shown experimentally with *Pseudomonas fluorescens (18)*. Here, we were able to better

quantify the frequency of hypermutators because of sufficient sample size, and test if within-host

Vc diversity is associated with phage or antibiotic exposure – both of which could potentially

select for resistance mutations to each factor.

287   To identify hypermutators in metagenomes, we identified *Vc* populations with defects

(nonsynonymous mutations) in any of 65 previously defined DNA repair genes (*42*) or with a

relatively high number of SNVs (25 or more) (*41*). We used InStrain (*43*) to quantify *Vc* within-

host diversity in 133 samples passing stringent quality filters (Methods) and found that 35% of

samples (47/133) had both a high SNV count and nonsynonymous mutations in DNA repair

genes – making them likely to contain hypermutators. Higher SNV counts were significantly

associated with DNA repair defects (Fisher's exact test, *P*<2.2e-16), consistent with their

resulting in higher mutation rates within patients. The number of SNVs was not confounded by

Vc genome coverage (**Fig. S15A**). Consistent with our previous study (*41*), putative

hypermutators had a distinct mutational profile, enriched in transversions (**Fig. S15B,C**). This

suggests that hypermutators are common, occurring in over one third of cholera patients. We

also observed that samples with high numbers of SNVs but no DNA repair defects had higher

levels of phage ICP1 (**Fig. S15D**), suggesting that phage may select for resistance mutations in

hypermutators and non-mutator populations alike. For subsequent analysis, we considered all

SNVs together, regardless of whether they were generated by hypermutation.

Phages, not antibiotics, are associated with Vc within-host diversity. We asked if *Vc*

within-host diversity was dependent on phage or antibiotic exposure – both strong selective

pressures. To test this hypothesis, we fit GLMMs with phages and antibiotics as predictors of

the number of high-frequency nonsynonymous (NS) SNVs in the *Vc* population within a patient.

We focused on higher-frequency SNVs (>10% within a sample) as likely beneficial mutations

(unlikely to rise to such high frequency by chance if neutral or deleterious), and on NS SNVs as

these are more likely to have fitness effects. We fit several models with different combinations of

predictors: from a model with all antibiotics and their interaction with ICP1 to separate models

with each antibiotic and its interaction with ICP1. We added *Vc* abundance as a fixed effect to

the model to control for any coverage bias in SNV calling (**Table S8**). The most parsimonious

model included *Vc* abundance and the interaction between *Vc* and ICP1 as predictors of the

number of high-frequency NS SNVs. Adding antibiotics, or their interaction with ICP1, did not

improve the model (**Table S8**), suggesting a limited role for antibiotics in selecting for *Vc* point

mutations within patients.

In the model, *Vc* relative abundance and the interaction between *Vc* and ICP1 both had

significant effects (GLMM, Wald test, P=0.00246 and P=0.00494 respectively). The negative

relationship between *Vc* and the number of high-frequency NS SNVs (**Fig. 4A**) was not easily

explained by sequencing coverage, since the total number of SNVs is not associated with *Vc*

relative abundance (**Fig. S15A**). The number of high-frequency NS SNVs rose with increasing

ICP1 – but only when *Vc* abundance was relatively high (**Fig. 4A**). As a control, we ran the

same GLMM on NS SNVs without a minimum frequency cutoff and found no significant effects,

suggesting that the interaction between ICP1 and *Vc* on SNV count is specific to high-frequency

mutations, rather than low-frequency mutations that are more likely neutral or deleterious.

These data support a scenario in which ICP1 selects for NS SNVs when the *Vc* population is

large enough to respond efficiently to selection – for example, at the beginning of an infection or

in the absence of antibiotics.

If phages select for beneficial mutations, we expect these mutations to increase in

frequency at higher phage predation. We lack time-series data from individual patients, but the

relative abundance of phage provides a proxy for the combined effects of the strength and

duration of phage selection. To test this hypothesis, we fit a GAM with the average within-

sample frequency of SNVs as a function of ICP1, antibiotics, and their interactions. We included

the fixed effect of ICE presence/absence as another factor that could provide phage or antibiotic

resistance, as well as mutation type to differentiate among non-synonymous (NS), synonymous

(S), and intergenic (I) mutations. We fit GAMs with all antibiotics and their interaction with ICP1,

as well as models with each antibiotic separately (**Table S9**). The most parsimonious model

included the interaction between ICP1, ICE and mutation type, but not antibiotics. ICP1 was a

strong predictor of higher frequency NS SNVs in the absence of a detectable ICE (**Fig. 4B**).

Samples in this analysis were unambiguous in terms of their ICE presence/absence patterns

(**Fig. S16**).

To confirm this model prediction, we compared the distribution of the average frequency

of NS SNVs between ICP1-positive and ICP1-negative samples. Consistent with the model, NS

SNV frequency was significantly higher in ICP1-positive samples when the ICE was not

detected (Wilcoxon test, P=0.0094) making them the likely targets of selection by ICP1

predation (**Fig. 4C**). Qualitatively similar results were found when ICP1 was quantified by qPCR

instead of metagenomics (**Fig S17**). Together, these results suggest that, in the absence of

detectable ICE-encoded phage resistance, ICP1 selects for nonsynonymous point mutations

(**Fig 5A**). We identified several *Vc* genes, including some with membrane or virulence-related

functions, that were mutated at higher levels ICP1:*Vc* ratios (**Table S10**); these provide

candidate phage resistance mechanisms that can be explored in future experiments. In

contrast, the secreted hemolysin gene, *hlyA,* that we previously observed (*41)* to be mutated

more often by chance in cholera patients was among the genes most frequently mutated in

patients with relatively low ICP1:*Vc* ratios (**Table S11**). This suggests that *hlyA* sequence

evolution may be affected directly or indirectly by phage predation.

As *Vc* evolves as a function of ICP1, we expect ICP1 evolution to be impacted by *Vc*

(**Fig 5A**). Despite the generally low genetic diversity of ICP1 (*21)*, were were able to quantify

SNVs in the ICP1 genome in 45 samples. This sample size was too low to fit sophisticated

models, but simple correlations allowed us to draw tentative conclusions. First, we ruled out

sequencing coverage as source of bias in SNV calling (Spearman correlation between ICP1

relative abundance and number of SNVs, $p > 0.1$ for all SNV categories). Next, we observed a

negative correlation between the ICP1:*Vc* ratio and the number of NS SNVs in the ICP1

genome – a correlation that is only significant when *Vc* encodes an *ind5* ICE (**Fig 5B, S18**). This

is consistent with our observation that *ind5* is associated with lower ICP1:*Vc* ratios in our cohort

(**Fig. 3**), potentially suppressing ICP1 and applying selection to escape suppression via NS

mutations. Some of these NS mutations may be beneficial to the phage, rising to high frequency

along with ICP1 – which is indeed what we observe: the mean frequency of NS SNVs in ICP1

increases with the ICP1:*Vc* ratio, only in the presence of *ind5* (**Fig 5C, S19**). Several ICP1

genes, mostly hypothetical proteins, repeatedly acquired NS mutations in the presence of *ind5*,

providing candidate escape mutations to test in future work (**Table S12**).

Fig. 4. ICP1 selects for non-synonymous point mutations in the *V. cholerae* genome in the absence of ICE. (A) Results of a GLMM modeling high frequency nonsynonymous SNV counts as a function of *V. cholerae* (*Vc*) and ICP1 standardized relative abundances. In the bottom panel, shades of gray indicate *Vc* relative abundance at the mean or +/- 1 standard deviation (SD) across samples. (B) GAM output with the mean mutation frequency as a function of the interaction between ICP1, ICE and mutation type (non-synonymous; NS, synonymous; S, or intergenic; I). Significant effects are shown with a star (Chi-square test, *P*<0.05). The model was fit using 130 samples that passed the post-InStrain filter (Methods). (C) Boxplots of mutation frequency in the presence or absence of ICP1 and/or ICEs. The single significant comparison is indicated with a star (Wilcoxon test, *P*=0.0094). Boxplots include 130 samples, of which 32 are ICP1+ (ICP1>=0.1% of reads) and 98 are ICP- (ICP1<0.1% of reads). The solid horizontal line is the median and the boxed area is the interquartile range. For supporting analysis using qPCR data, see Fig. S17.

The tripartite interactions between pathogens, phage, and antibiotics have been studied in the laboratory, *in silico* with mathematical models, and to a lesser extent in the field, but how these factors interact during human infection remains an open question. Our objective was to characterize these interactions in the context of cholera. We analyzed more than 300 stool

metagenomes from cholera patients enrolled at hospital admission across Bangladesh during

an entire outbreak season. We found that high predator (ICP1) to prey (*Vc*) ratios were

associated with less severe dehydration. We also demonstrate how *Vc* and ICP1 interact within

patients, with ICP1 selecting for potential phage resistance point mutations in the absence of

ICE-based anti-phage defenses, and ICE-encoding *Vc* selecting for point mutations in the

phage genome (**Figure 5**). This apparent coevolution between predator and prey likely has

longer-term consequences for cholera disease severity and transmission. Antibiotics,

particularly azithromycin, also played a role in suppressing *Vc* and could mask phage-bacteria

interactions. Ciprofloxacin was associated with known antibiotic resistance genes, but we found

no evidence that antibiotics select, as ICP1 does, for high-frequency nonsynonymous point

mutations. Thus, although resistance mechanisms to certain phages and antibiotics colocalize

to the ICE (*17*), they impose distinct selective pressures that could be exploited to improve the

efficacy of antibiotics by combining them with phage therapy.

Fig. 5. Phage-bacteria coevolution and disease severity in cholera patients. (A) The illustration
summarizes the main conclusions supported by our analyses. Patients with mild dehydration (left) have
higher ratios of phage ICP1 to pathogen *Vibrio cholerae*, while the opposite is true in patients with severe
dehydration (right). Although the causal relationship between the ICP1:*Vc* ratio and dehydration remains
unclear, it provides a potential biomarker for disease severity. At high ICP1:*Vc* ratios, and in the absence
of anti-phage genes encoded on the ICE, a higher frequency of nonsynonymous point mutations
(asterisks on white circular genomes) are selected for in the *Vc* genome, with potential evolutionary
consequences: some mutations may provide phage resistance while others may simply provide greater
genetic diversity upon which future natural selection can act. When ICE (primarily the *ind5* variant) is
encoded by *Vc*, its anti-phage properties impose strong selection on ICP1, selecting for nonsynonymous
point mutations in the ICP1 which may eventually rise to high frequency. For simplicity, the effects of
antibiotics are not shown, but antibiotics such as azithromycin can suppress *Vc*, thereby masking
interactions with ICP1. **(B)** The number of nonsynonymous (NS) SNVs detected in the ICP1 genome is
negatively correlated with the ICP1:*Vc* ratio in the presence of *ind5.* The Spearman correlation coefficient
and *p*-value are shown. **(C)** The mean frequency of NS SNVs in the ICP1 genome is positively correlated
with the ICP1:*Vc* ratio in the presence of *ind5.* See Figures S18 and S19 for equivalent plots in ICE- and
ind6 samples, and for synonymous and intergenic SNVs.

Our study has several limitations. First, samples were collected at a single time point

(upon hospital admission) which allowed us to establish statistical correlations, but we cannot

infer causality in the absence of time-series or interventional clinical studies. Second, our

sample size allowed us to study the interaction between *Vc* and ICP1, but the sample size for

ICP2 and ICP3 was insufficient for similar statistical analyses. Third, we prioritized common

antibiotics for mass spectrometry, and we cannot exclude a role for other unmeasured

antibiotics. Fourth, due to logistical limitations, we extracted DNA from a bacterial pellet plus a

small amount of supernatant from each sample. This allowed us to capture both intracellular

and cell-bound phages, along with free phage particles, but sequencing each fraction separately

could yield further insights into distinct phage populations. From a clinical perspective, the

measurement of dehydration status was categorical and could be improved in future studies by

incorporating digital tools to quantify the degree of dehydration (*44-46*). Finally, our study lacked

information about host genetics or immunity, which also influence disease severity (*8, 47*).

Future studies combining rich patient metadata, a time-series design, long-read metagenomics,

and isolate genome sequencing will complement and expand upon these findings.

CONCLUSION

We propose that an index of effective phage predation, quantified as the phage:bacteria ratio, might be used as a tool for physicians to assess disease severity, and potentially prognosticate a disease course. While we show here that this ratio is associated with cholera disease severity, its predictive value should be studied in larger cohorts sampled over the course of infection. Whether this biomarker can be generalized to phages other than ICP1 and diseases other than cholera, and whether the association with disease severity changes as predator and prey coevolve over time, remains unclear. The potential of phage therapy and prophylaxis has long been recognized, and a combination of ICP1, 2, and 3 prevents cholera in animal models (*35*). However, just as hypermutators can drive the evolution of resistance to combinations of antibiotics (*48*), they may also help pathogen populations survive combinations of phages, while increasing their potential to evolve resistance to future antimicrobial treatments. Phages therapy cocktails will therefore need to be updated regularly to remain effective against currently circulating coevolved bacteria, and creative new strategies are needed to minimize the unwanted evolution of resistance.

Materials and Methods

Ethics Statement. The samples analyzed were collected within two previously published IRB approved clinical studies in Bangladesh: (i) The mHealth Diarrhea Management (mHDM) cluster randomized controlled trial (IEDCR IRB/2017/10; icddr,b ERC/RRC PR-17036; University of Florida IRB 201601762) (*46*). (ii) The National Cholera Surveillance (NCS) study (icddr,b ERC/RRC PR-15127) (*49*); See supplementary materials for further details.

Study Design. The study design was a prospective longitudinal study of patients presenting with diarrheal disease at five Bangladesh Ministry of Health and Family Welfare district hospitals (both mHDM and NCS sites) and two centralized NCS hospitals (BITID; icddr,b) from March

2018 to December 2018. Sites were distributed geographically nation-wide (*50*). See

supplementary materials.

Sample collection. Stool samples were collected at hospital admission. Aliquots for transport

and subsequent culture were stabbed into Cary-Blair transport media; aliquots for molecular

analysis were preserved in RNAlater (Invitrogen). See supplementary materials.

Microbiological and molecular analysis. Culture was performed via standard methods (*51*);

total nucleic acid (tNA) was extracted from the RNAlater preserved samples using standard

methods. Criteria for subset subsequent shotgun metagenomic sequencing were: (i) culture

positivity, (ii) Phage (ICP1,2,3) detection by PCR among culture-negative samples, or (iii) *Vc*

detection by PCR among a random 10% of culture-negative and phage (ICP1,2,3) negative

samples. Sequencing libraries were prepared using the NEB Ultra II shotgun kit and sequenced

on illumina NovaSeq 6000 S4, pooling 96 samples per lane, yielding a mean of >30 million

paired-end 150bp reads per sample. Among samples identified for metagenomic analysis,

qPCR was performed to determine absolute abundances of *Vc*, ICP1, ICP2, and ICP3 using

standard methods.

Antibiotic detection by liquid chromatography mass spectrometry (LC-MS/MS). Those

cholera samples identified for metagenomic analysis were analyzed by qualitative and

quantitative LC-MS/MS for antibiotics. Based on prior research (*16, 37*), the target list prioritized

5 common antibiotics: ciprofloxacin, doxycycline/tetracycline, and azithromycin were analyzed

quantitatively, and metronidazole and nalidixic acid were analyzed qualitatively. Standard

curves were made for each quantitative target by preparing a dilution series of the three native

and isotopic forms of the quantitative targets; clinical samples were spiked with the isotopes of

the quantitative targets as internal references. See supplementary materials.

Metagenomic data analysis. We taxonomically classified short reads using Kraken2 (*52*) and

Bracken v.2.5 (*53*). Reads were assembled using MEGAHIT v.1.2.9 (*50*) and binned with DAS

tool (*54*). We inferred probably phage contigs using geNomad v1.7.0 (*55*) and predicted their

likely bacterial hosts with iPhoP v1.3.1 (*56*). To characterize intra-patient *Vc* diversity, we used

StrainGE (*57*) and InStrain v.1.5.7 (*43*). To identify antibiotic resistance genes in metagenomes,

we used deepARG v 1.0.2 (*58*). See supplementary materials for details.

Statistical analyses. Statistics and visualizations were done in R studio version 1.2.5042. See

supplementary materials for details.

Data availability. All sequencing data are deposited in the NCBI SRA under BioProject

PRJNA976726. See supplementary materials for further information.

Code availability. Computer code for this paper will be available at

https://github.com/Naima16/Cholera-phage-antibiotics.

Acknowledgements: We thank the patients for participating in this study as well as the clinical

and laboratory teams who collected the samples. We are grateful to S. Flora and colleagues at

the Institute of Epidemiology, Disease Control and Research (IEDCR), Ministry of Health and

Family Welfare, Government of Bangladesh who collaborated on the original clinical studies in

which the samples analyzed herein were collected. We are also grateful to R. Autrey, B.

Johnson and K. Berquist for their administrative expertise at the University of Florida. AIK and

FQ were the principal investigators in Bangladesh and PIs of the ERC/RRC approvals at the

icddr,b. This collective research infrastructure and support was invaluable to the success of this

study. We thank members of the Nelson, Khan, Qadri and Shapiro labs, and Frederique Le

Roux, for discussions that improved the manuscript.

Funding: This work was supported by the National Institutes of Health grants to EJN

[R21TW010182] and KBB [S10 OD021758-01A1] and internal support from the Emerging

Pathogens Institute at the University of Florida and the Departments of Pediatrics/ Children's

Miracle Network (Florida). BJS and NM were supported by a Canadian Institutes for Health

Research Project Grant. AC was supported by a Postdoctoral Mobility Fellowship of the Swiss

National Science Foundation [P500PB_214356]. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions:

Conceptualization: FW, AIK, BJS, EJN
Methodology: NM, MAS, ETC, MTRB, YB, ACM, MK, AC, EF, AV, LSB, KBS, BJS, EJN
Investigation: NM, MAS, ETC, KI, MIUK, YB, ACM, MK, AC, EF, AV, LSB, BJS, EJN
Visualization: NM, BJS, EJN
Funding acquisition: KBS, FW, AIK, BJS, EJN
Project administration: MTRB, YB, KBS, AIK, BJS, EJN
Supervision: KBS, FQ, AIK, BJS, EJN
Writing – original draft: NM, ETC, MAS, ACM, MK, LSB, KSB, AIK, BJS, EJN
Writing – review & editing: NM, AIK, AC, BJS, EJN

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary

materials.

Supplementary materials:

Materials and methods

Figures: S1 to S19

Tables: S1 to S12

References

Data Files S1 to S5

References

1. J. R. Andrews *et al.*, Determinants of severe dehydration from diarrheal disease at hospital presentation: Evidence from 22 years of admissions in Bangladesh. *PLoS Negl Trop Dis* **11**, e0005512 (2017).

2. M. Ali, A. R. Nelson, A. L. Lopez, D. A. Sack, Updated global burden of cholera in endemic countries. *PLoS Negl Trop Dis* **9**, e0003832 (2015).

3. Cholera – Global situation (2023). WHO Report. (available at https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON437).

4. M. S. Islam *et al.*, Microbiological investigation of diarrhoea epidemics among Rwandan refugees in Zaire. *Trans R Soc Trop Med Hyg* **89**, 506 (1995).

5. J. A. Dromigny, O. Rakoto-Alson, D. Rajaonatahina, R. Migliani, J. Ranjalahy, P. Mauclere, Emergence and rapid spread of tetracycline-resistant *Vibrio cholerae* strains, Madagascar. *Emerg Infect Dis* **8**, 336-338 (2002).

6. F. X. Weill *et al.*, Genomic history of the seventh pandemic of cholera in Africa. *Science* **358**, 785-789 (2017).

7. E. J. Nelson, D. S. Nelson, M. A. Salam, D. A. Sack, Antibiotics for both moderate and severe cholera. *N Engl J Med* **364**, 5-7 (2011).

8. E. J. Nelson, J. B. Harris, J. G. Morris, Jr., S. B. Calderwood, A. Camilli, Cholera transmission: the host, pathogen and bacteriophage dynamic. *Nat Rev Microbiol* **7**, 693-702 (2009).

9. B. Das, J. Verma, P. Kumar, A. Ghosh, T. Ramamurthy, Antibiotic resistance in *Vibrio cholerae*: Understanding the ecology of resistance genes and mechanisms. *Vaccine* **38 Suppl 1**, A83-A92 (2020).

10. F. Lassalle *et al.*, Genomic epidemiology reveals multidrug resistant plasmid spread between *Vibrio cholerae* lineages in Yemen. *Nat Microbiol* **8**, 1787-1798 (2023).

561   11.   N. Rivard, R. R. Colwell, V. Burrus, Antibiotic Resistance in *Vibrio cholerae*: Mechanistic

562   Insights from IncC Plasmid-Mediated Dissemination of a Novel Family of Genomic

563   Islands Inserted at trmE. *mSphere* **5**, (2020).

564   12.   S. J. Ambrose, C. J. Harmer, R. M. Hall, Compatibility and entry exclusion of IncA and

565   IncC plasmids revisited: IncA and IncC plasmids are compatible. *Plasmid* **96-97**, 7-12

566   (2018).

567   13.   M. K. Waldor, H. Tschape, J. J. Mekalanos, A new type of conjugative transposon

568   encodes resistance to sulfamethoxazole, trimethoprim, and streptomycin in *Vibrio*

569   *cholerae* O139. *J Bacteriol* **178**, 4157-4165 (1996).

570   14.   A. Dalsgaard, A. Forslund, N. V. Tam, D. X. Vinh, P. D. Cam, Cholera in Vietnam:

571   changes in genotypes and emergence of class I integrons containing aminoglycoside

572   resistance gene cassettes in *Vibrio cholerae* O1 strains isolated from 1979 to 1996. *J*

573   *Clin Microbiol* **37**, 734-741 (1999).

574   15.   J. W. Beaber, B. Hochhut, M. K. Waldor, Genomic and functional analyses of SXT, an

575   integrating antibiotic resistance gene transfer element derived from *Vibrio cholerae*. *J*

576   *Bacteriol* **184**, 4259-4269 (2002).

577   16.   A. Creasy-Marrazzo *et al.*, Genome-wide association studies reveal distinct genetic

578   correlates and increased heritability of antimicrobial resistance in *Vibrio cholerae* under

579   anaerobic conditions. *Microbial Genomics* **8**, (2022).

580   17.   K. LeGault *et al.*, Temporal shifts in antibiotic resistance elements govern phage-

581   pathogen conflicts. *Science* **373**, 2020.2012.2016.423150 (2021).

582   18.   C. Pal, M. D. Macia, A. Oliver, I. Schachar, A. Buckling, Coevolution with viruses drives

583   the evolution of bacterial mutation rates. *Nature* **450**, 1079-1081 (2007).

584   19.   M. Yen, A. Camilli, Mechanisms of the evolutionary arms race between *Vibrio cholerae*

585   and Vibriophage clinical isolates. *Int Microbiol* **20**, 116-120 (2017).

20.   K. D. Seed, S. M. Faruque, J. J. Mekalanos, S. B. Calderwood, F. Qadri, A. Camilli,

Phase variable O antigen biosynthetic genes control expression of the major protective

antigen and bacteriophage receptor in *Vibrio cholerae* O1. *PLoS Pathog* **8**, e1002917

(2012).

21.   C. M. Boyd, A. Angermeyer, S. G. Hays, Z. K. Barth, K. M. Patel, K. D. Seed,

Bacteriophage ICP1: A Persistent Predator of *Vibrio cholerae*. *Annu Rev Virol* **8**, 285-

304 (2021).

22.   K. D. Seed, D. W. Lazinski, S. B. Calderwood, A. Camilli, A bacteriophage encodes its

own CRISPR/Cas adaptive response to evade host innate immunity. *Nature* **494**, 489-

491 (2013).

23.   K. D. Seed *et al.*, Evolutionary consequences of intra-patient phage predation on

microbial populations. *Elife* **3**, e03497 (2014).

24.   R. C. Molina-Quiroz, A. Camilli, C. A. Silva-Valenzuela, Role of Bacteriophages in the

Evolution of Pathogenic Vibrios and Lessons for Phage Therapy. *Adv Exp Med Biol*

1404, 149-173 (2023).

25.   E. J. Nelson *et al.*, Transmission of *Vibrio cholerae* is antagonized by lytic phage and

entry into the aquatic environment. *PLoS Pathog* **4**, e1000187 (2008).

26.   K. D. Seed *et al.*, Evidence of a dominant lineage of *Vibrio cholerae*-specific lytic

bacteriophages shed by cholera patients over a 10-year period in Dhaka, Bangladesh.

MBio **2**, e00334-00310 (2011).

27.   W. C. Summers, Cholera and plague in India: the bacteriophage inquiry of 1927-1936. *J*

Hist Med Allied Sci **48**, 275-301 (1993).

28.   F. D'Herelle, R. Malone, A preliminary report of work carried out by the cholera

bacteriophage enquiry. *Indian Medical Gazette*, 614-617 (1927).

29.   C. L. Pasricha, A. J. H. de Monte, E. G. O'Flynn, Bacteriophage in the treatment of

cholera. *Ind. Med. Gaz.* **71**, 61-68 (1936).

30. S. M. Faruque *et al.*, Seasonal epidemics of cholera inversely correlate with the

prevalence of environmental cholera phages. *Proc Natl Acad Sci U S A* **102**, 1702-1707

(2005).

31. S. M. Faruque *et al.*, Self-limiting nature of seasonal cholera epidemics: Role of host-

mediated amplification of phage. *Proc Natl Acad Sci U S A* **102**, 6119-6124 (2005).

32. M. A. Jensen, S. M. Faruque, J. J. Mekalanos, B. R. Levin, Modeling the role of

bacteriophage in the control of cholera outbreaks. *Proc Natl Acad Sci U S A* **103**, 4652-

4657 (2006).

33. M. S. Zahid, S. M. Udden, A. S. Faruque, S. B. Calderwood, J. J. Mekalanos, S. M.

Faruque, Effect of phage on the infectivity of *Vibrio cholerae* and emergence of genetic

variants. *Infect Immun* **76**, 5266-5273 (2008).

34. A. Jaiswal, H. Koley, A. Ghosh, A. Palit, B. Sarkar, Efficacy of cocktail phage therapy in

treating *Vibrio cholerae* infection in rabbit model. *Microbes Infect* **15**, 152-156 (2013).

35. M. Yen, L. S. Cairns, A. Camilli, A cocktail of three virulent bacteriophages prevents

Vibrio cholerae infection in animal models. *Nat Commun* **8**, 14187 (2017).

36. L. A. David *et al.*, Gut microbial succession follows acute secretory diarrhea in humans.

MBio **6**, e00381-00315 (2015).

37. L. Alexandrova *et al.*, Identification of Widespread Antibiotic Exposure in Patients With

Cholera Correlates With Clinically Relevant Microbiota Changes. *J Infect Dis* **220**, 1655-

1666 (2019).

38. M. M. Monir *et al.*, Genomic attributes of *Vibrio cholerae* O1 responsible for 2022

massive cholera outbreak in Bangladesh. *Nat Commun* **14**, 1154 (2023).

39. A. Carr, C. Diener, N. S. Baliga, S. M. Gibbons, Use and abuse of correlation analyses

in microbial ecology. *ISME J* **13**, 2647-2655 (2019).

40. I. Levade *et al.*, *Vibrio cholerae* genomic diversity within and between patients. *Microb*

Genom **3**, (2017).

41.   I. Levade *et al.*, A Combination of Metagenomic and Cultivation Approaches Reveals

639   Hypermutator Phenotypes within *Vibrio cholerae*-Infected Patients. *mSystems* **6**,

640   e0088921 (2021).

641   42.   A. Jolivet-Gougeon *et al.*, Bacterial hypermutation: clinical implications. *J Med Microbiol*

642   **60**, 563-573 (2011).

643   43.   M. R. Olm, A. Crits-Christoph, K. Bouma-Gregson, B. A. Firek, M. J. Morowitz, J. F.

644   Banfield, inStrain profiles population microdiversity from metagenomic data and

645   sensitively detects shared microbial strains. *Nat Biotechnol* **39**, 727-736 (2021).

646   44.   A. C. Levine *et al.*, A comparison of the NIRUDAK models and WHO algorithm for

647   dehydration assessment in older children and adults with acute diarrhoea: a prospective,

648   observational study. *Lancet Glob Health* **11**, e1725-e1733 (2023).

649   45.   A. C. Levine *et al.*, External validation of the DHAKA score and comparison with the

650   current IMCI algorithm for the assessment of dehydration in children with diarrhoea: a

651   prospective cohort study. *Lancet Glob Health* **4**, e744-751 (2016).

652   46.   A. I. Khan *et al.*, Electronic decision support and diarrhoeal disease guideline adherence

653   (mHDM): a cluster randomised controlled trial. *Lancet Digit Health* **2**, e250-e258 (2020).

654   47.   J. B. Harris *et al.*, Susceptibility to *Vibrio cholerae* Infection in a Cohort of Household

655   Contacts of Patients with Cholera in Bangladesh. *PLoS Negl Trop Dis* **2**, e221 (2008).

656   48.   B. Seed, Purification of genomic sequences from bacteriophage libraries by

657   recombination and selection in vivo. *Nucleic Acids Res* **11**, 2427-2445 (1983).

658   49.   A. I. Khan, F. Qadri, Epidemiology of cholera in Bangladesh: Findings from Nationwide

659   Hospital-based Surveillance, 2014-2018. *CID*, (2019).

660   50.   D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: an ultra-fast single-node

661   solution for large and complex metagenomics assembly via succinct de Bruijn graph.

662   *Bioinformatics* **31**, 1674-1676 (2015).

663   51.   *Manual of Clinical Microbiology. 8th ed.* P. Murray, E. Baron, J. Jorgensen, M. Pfaller, R.

664   Yolken, Eds., (American Society for Microbiology Press, Washington, D.C., 2003).

665   52.   D. E. Wood, J. Lu, B. Langmead, Improved metagenomic analysis with Kraken 2.

666   *Genome Biol* **20**, 257 (2019).

667   53.   J. Lu, F. P. Breitwieser, P. Thielen, S. L. Salzberg, Bracken: estimating species

668   abundance in metagenomics data. *PeerJ Comput. Sci* **3**, (2017).

669   54.   C. M. K. Sieber *et al.*, Recovery of genomes from metagenomes via a dereplication,

670   aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).

671   55.   A. P. Camargo *et al.*, Identification of mobile genetic elements with geNomad. *Nat*

672   *Biotechnol*, (2023).

673   56.   S. Roux *et al.*, iPHoP: An integrated machine learning framework to maximize host

674   prediction for metagenome-derived viruses of archaea and bacteria. *PLoS Biol* **21**,

675   e3002083 (2023).

676   57.   L. R. van Dijk *et al.*, StrainGE: a toolkit to track and characterize low-abundance strains

677   in complex microbial communities. *Genome Biol* **23**, 74 (2022).

678   58.   G. Arango-Argoty, E. Garner, A. Pruden, L. S. Heath, P. Vikesland, L. Zhang, DeepARG:

679   a deep learning approach for predicting antibiotic resistance genes from metagenomic

680   data. *Microbiome* **6**, 23 (2018).

Supplementary Materials for

Phage predation is a biomarker for disease severity and shapes pathogen genetic diversity in cholera patients

Authors: N. Madi[1,2], E. T. Cato[3], Md. Abu Sayeed[3], A. Creasy-Marrazzo[3], A. Cuénod[1,2], K. Islam[4], Md. I.UI. Khabir[4], Md. T. R. Bhuiyan[4], Y. Begum[4], E. Freeman[3], A. Vustepalli[3], L. Brinkley[3], Kamat[4], L. S. Bailey[5], K. B. Basso[5], F. Qadri[4], A. I. Khan[4*], B. J. Shapiro[1,2,6*], E. J. Nelson[3*]

Affiliations:

[1] Department of Microbiology & Immunology, McGill University, Montréal, Canada
[2] McGill Genome Centre, McGill University, Montréal, Canada
[3] Departments of Pediatrics and Environmental and Global Health, University of Florida, Gainesville, FL, USA
[4] Infectious Diseases Division (IDD) & Nutrition and Clinical Services Division (NCSD), International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh
[5] Department of Chemistry, University of Florida, Gainesville, FL, USA
[6] McGill Centre for Microbiome Research, McGill University, Montréal, Canada

[*] Corresponding authors

Email: ashrafk@icddrb.org; jesse.shapiro@mcgill.ca; eric.nelson@ufl.edu

The PDF file includes:

Index
Materials and methods
Results
Figures: S1 to S19
Tables: S1 to S12
References

Other Supplementary Materials for this manuscript include the following:
Data S1 to S5

Materials and Methods

Ethics Statement. The samples analyzed were collected within the context of two previously published IRB approved clinical studies conducted in Bangladesh: the mHealth Diarrhea Management (mHDM) cluster randomized controlled trial (IEDCR IRB/2017/10; icddr,b ERC/RRC PR-17036; University of Florida IRB 201601762; clinicaltrials.gov NCT03154229) (*1*) and a hospital-based National Cholera Surveillance (NCS) study (icddr,b ERC/RRC PR-15127) (*2*); the consent processes are described within these published studies.

Study Design. The study design was a national prospective longitudinal study of patients presenting with diarrhoeal disease at five Bangladesh Ministry of Health and Family Welfare district hospitals (both mHDM and NCS sites) and two centralized hospitals (NCS sites alone) (BITID, icddr,b) from March 2018 to December 2018. Recruitment was based on the census of diarrheal disease patients that sought care at the hospitals. For the mHDM study, inclusion criteria were patients two-months of age or older, and presentation with acute diarrhea defined as three or more episodes of loose stools in the 24 hours prior to admission and a duration of disease less than seven days. For the NCS study, patients with diarrheal disease of all ages were included. For those under 2 months of age, diarrhea was defined as a change in stool habit from 'usual' (increased frequency with less formed stool). For those 2 months and older, diarrhea was defined as 3 or more loose or liquid stools within 24 hours or 3 loose/liquid stools or fewer causing dehydration in the last 24 hours. There was no exclusion based on enrollment in a second study (e.g., mHDM vs NCS). Patient metadata is in Data File S1.

Sample Collection. For the mHDM study, the intent was to collect four stool samples per study site per day. For the NCS study, the intent was to collect daily stool samples from two participants less than 5 years old and two participants 5 years of age and older; if the target number for an age group was not met, samples were collected from the other age group to achieve a total of 4 samples per day. For the samples collected at the five district hospitals and BITID, an aliquot was placed in transport media (Cary-Blair) and transferred to the icddr,b laboratory for culture, and a 0.5 ml aliquot was added to 1.3 ml of RNAlater to stabilize the sample for subsequent analysis; cold-chain below 4°C was not consistently available at these sites. For samples collected at the icddr,b, samples were stabilized in RNAlater; culture was performed directly. Samples were stored at the centralized icddr,b laboratory at -80°C.

Antibiotic detection by liquid chromatography mass spectrometry (LC-MS/MS).

LC-MS/MS methodology for both qualitative and quantitative approaches. The approach was based on prior studies (*3, 4*). Stool supernatants from the primary collection were obtained by centrifugation without filtration to minimize loss. Proteins were precipitated (1:7 ratio (v/v) of

water:methanol). Supernatants were diluted with methanol and water (1:1 v/v) in 1% formic acid for liquid chromatography, and 5 µl of supernatant was injected for analysis. LC-MS/MS was performed on a 2.1 x 150-mm Hypersil Gold aQ column (particle size, 3 µm) using a high-performance liquid chromatography system (Thermo UltiMate 3000 series) with an LTQ XL ion trap mass spectrometer (Thermo Fisher Scientific). Mobile phases were 1% formic acid in water (A) and 1% formic acid in methanol (B) and held at a constant 5%B for 2 minutes before ramping to 95%B at 15 minutes where it was held for an additional minute before returning to starting conditions for a total run time of 25 min. Blanks were run in between every two samples, as well as before and after quality control samples and standards.

Eluent was ionized using electrospray ionization (ESI) in positive mode at a spray voltage of 5 kV, a nitrogen sheath gas flow rate of 8 L min^{-1}, and capillary temperature of 300°C. Two scan events were programmed to perform an initial scan from *m/z* 100 to 1000, which was followed by targeted collision induced dissociation based on a retention time and mass list. Retention time windows ranged from 1.7 minutes to 5.1 min, depending on the elution range of the standards at high and low concentrations. Masses were targeted for the most abundant adduct or ion associated with each antibiotic (typically the [M+H]$^{+}$ ion) with a *m/z* 1 window.

LC-MS/MS methodology specific to qualitative analysis. The target list for the qualitative analysis was narrowed from prior work that assayed for antibiotics and non-antibiotics (*3, 4*): acetaminophen, Metronidazole, Zofran, Furazolidone, Nalidixic Acid, Sulfamethoxazole, Trimethoprim, Omeprazole, Ciprofloxacin, Cephalexin, Penicillin V (M+H), Amoxicillin, Penicillin V, Doxycycline or Tetracycline, Ceftriaxone, Erythromycin, and Azithromycin. Based on this prior work, we narrowed our analysis to the common clinical relevant agents of ciprofloxacin, azithromycin, doxycycline/tetracycline (same mass), nalidixic acid and metronidazole. Qualitative data analysis was performed using extracted ion chromatograms and MSMS matching with a control antibiotic MSMS library using Xcalibur 2.2 SP 1.48 (Thermo Fisher Scientific).

LC-MS/MS methodology specific to quantitative analysis. The target list for the quantitative analysis was furthered narrowed to: Ciprofloxacin, Azithromycin and Tetracycline/Doxycycline. A standard curve was made for each quantitative target by preparing a dilution series of a mix of the three native forms of the quantitative targets (Ciprofloxacin, Doxycycline, Azithromycin). The dilutions were: 0.5, 0.25, 0.125, 0.063, 0.05, 0.02, 0.01 µg/ml. All samples, including standards, were spiked with a 0.25 µM mix of isotope labeled control antibiotics (Ciprofloxacin d-8 (Millipore-Sigma), doxycycline d-5 (Isosciences), Azithromycin d-5 (LGC standards).

A calibration curve was generated by plotting the ratio of the area under the curve for analyte peaks in the ion chromatograms to the AUC of the isotope peaks against known

concentrations of the seven standards. The linear line of best fit produced by this plot was generated and used to extrapolate the quantity of drug in mHDM samples. The AUC of chromatogram peaks produced by the target compound was compared to that of the peak produced by the isotope spike in to generate a ratio. This ratio was then converted to a concentration (in units of µg/ml) using the calibration curve equation. Data analysis was performed manually by viewing the extracted ion chromatograms and MSMS from each sample and matching with an in-house antibiotic library using 2.2 SP 1.48 (Thermo Fisher Scientific). The detection of a drug was confirmed when a peak in the chromatogram at the proper retention time window was identified and the most abundant adduct or ion associated with that drug (typically the $[M+H]^+$ ion) with a m/z 1 window. Measured antibiotic concentrations are in Data File S2.

Molecular analyses. Standard methods were used to extract total nucleic acid (tNA) from the samples stabilized in RNAlater (Invitrogen). In brief, the samples were thawed at room temperature and centrifuged for 5 minutes at max speed. All but 50 µl of the supernatant was removed; the supernatant was refrozen for subsequent mass spectrometry. The remaining pellet, with the 50 µl supernatant, was combined with 380 mg of glass beads and 1 ml InhibitEx buffer and processed using the QIAamp Fast DNA Stool Mini Kit. Elutions were performed with 50 µl of Qiagen ATE buffer followed by a second elution with 50 µl ATE, quantification (Nanodrop) and storage at -80°C. For qPCR, we used established *tcpA* primers for *V. cholerae* (*4, 5*) and 16S ribosomal gene (rDNA) primers as a positive control for sample processing and extraction (*6*). Primer pairs were developed for the virulent bacteriophages ICP1 (gp58.2), ICP2 (gp24), and ICP3 (gp19); primer sequences are conserved between Southeast Asia and African isolates. Reactions used Luna qPCR master mix (New England Biolabs) with 25 µl volumes, and the method was a two-step reaction with denaturation at 98°C and annealing/extension at 60°C for all primers except ICP2 GP24 F/R (annealing/extension 63°C). Assays were performed on an Azure Cielo Real-time qPCR System and two Bio-Rad CFX96 Touch Real-Time PCR Detection Systems. Samples were analyzed in 96-well formats with two replicates except for 16SrDNA (one replicate); a Ct threshold less than 28 was used to define positivity (*4, 5, 7*). For quality control, samples that were negative for 16S rDNA were excluded from subsequent analyses. A third replicate was performed when: both replicates were positive but the difference in Ct values was more than 4, or one replicate was positive and one was negative and the difference in Ct values was greater than four. After the third replicate was performed, outliers were removed and the Ct values were averaged and the final determination of positivity was scored (Ct<28). Primers were validated using standard curves of (i) synthetic templates and (ii) biologic agents for *V. cholerae*, ICP1, ICP2, ICP3. Conventional methods were used for PCR.

<u>Metagenomic analyses</u>

Short read classification using kraken2/braken. We classified all short-read data with a Kraken2 database (Wood, Lu, and Langmead 2019) containing bacterial, archaeal, and viral domains, along with the human genome and a collection of known vectors (UniVec_Core) from NCBI, in July 2020. A Bracken database (Lu et al. 2017) was also built with a read length of 150 bp and the default k-mer length of 35. Kraken2 and Bracken version 2.5 were run with default parameters.

Phage identification from metagenomic assemblies. We characterised phages in 224 metagenomes with either >0.5% *Vc* or >0.1% ICP phage relative abundances. We used geNomad (v1.7.0, min-score 0.7, max-fdr 0.1) (*8*) to identify (pro)-phages from 83,928 contigs identified as 'viral'. iPhoP (v1.3.1, min-score 90%) (*8*) was used to predict the bacterial hosts of the identified phage contigs, resulting in 627 contigs classified as '*Vibrio* phages'. We used two different approaches to further characterise the predicted '*Vibrio* phage' sequences: (i) BLASTn to compare these sequences against known *Vibrio* phages in the INPHARED (*9*) database and reference sequences of the TLC (*10*) and PLE satellite phages (*11*) (% identity >= 80%, query coverage >= 90%), and (ii) vContact2 (*12*) to build similarity networks including the putative *Vibrio* phages and genomes from the INPHARED NCBI RefSeq databases. We used clusters in the vContact2 network to harmonise the naming of the INPHARED database genomes (eg. all INPHARED genomes which clustered together with the ICP2 reference genomes, were called 'ICP2').

Quality filtering. Before calling SNVs within the *V. cholerae* population, metagenomes were decontaminated of human and PhiX reads by mapping the reads against the GRCh 38 assembly of the human genome and the PhiX genomes, with bowtie2 version 2.3.5 (*13*). Unmapped reads were used for subsequent analyses. Then, we used iu_filter_quality_minoche from the Illumina Utils package (v2.10) with default parameters (*14*) to trim adapters, remove duplicates and for quality filtering on the short reads.

SNV profiling. Short reads were assembled with MEGAHIT (*15*) version 1.2.9 with the default parameters (k-mers length: k=21 et maxk=99 and min contig length of 1000. The contigs were grouped into bins with concoct (v1.0.0) (*16*) and MetaBAT 2 (v2.12.1) (*17*) with default parameters. Bins were then aggregated together using DAS Tool version 1.1.0 (*18*). Then we used drep (*19*) version 2.0.0 with S_ani 0.98, to de-replicate the bins and identify unique metagenome-assembled genomes (MAGs) with a secondary clustering threshold of 98% and minimum completeness of 75% and maximum contamination of 25%. To prevent mismapping of reads from other species to the *V. cholerae* genome, we competitively mapped reads from all

samples with sufficient *V. cholerae* or phages (*Vc*>0.5% of reads or phages>0.1% of metagenomic reads) against the concatenation of the 677 unique MAGs and ICP1 genome from NCBI (reference NC_015157.1, accessed 2021/12/01), using bowtie2 v 2.3.5. To characterize within-patient *V. cholerae* and ICP1 genetic diversity, we profiled the resulting bam files using inStrain v1.5.7 (*20*) with default parameters (minimum coverage of 5 to call a variant). We also ran gene-level profiling with instrain using genes annotated with prodigal. We used prodigal v2.6.3 (*21*) with the default parameters to predict genes in the *Vc* MAG and ICP1 and annotated them with eggNOG-Mapper v2 with default parameters and eggNOG database v2 downloaded on April 2021 (*22*).

To infer potential mixed infections, we used strainGST from the strainGE toolkit (*23*). To do so, we compiled a reference database of publicly available *V. cholerae* assemblies, including (i) all complete genomes from NCBI RefSeq (n=106, accessed 2023/05/12) and (ii) all assemblies published on NCBI Genbank between 2015/01/01 and 2023/05/12 and originating from samples collected between 2015/01/01 and 2019/12/31 (n=758).

To reduce false positive SNVs, we applied a stringent post-InStrain filter: all positions with coverage <20 were removed. We removed the first and last 100 positions of every scaffold, as well as positions with coverage below 0.3*median and above 3*median (median coverage across all the positions in the sample). We also removed sites that did not pass the coverage filter in more than 2 samples. Finally, we removed 420 sites that were variable in a *V. cholerae* isolate genome sequenced alongside the metagenomes. These sites were considered prone to sequencing error since they varied in an isolate genome that theoretically should contain no variation. After applying all these quality filters, we were left with 130 samples with SNV calls.

Hypermutator definition. Hypermutators were identified as samples with one or more nonsynonymous mutations in DNA repair genes (DNA repair defects) in *V. cholerae*, defined previously (*24*).

Antibiotic resistance gene identification. Antibiotic resistance genes (ARGs) were predicted from short reads using deepARG (*25*) v 1.0.2 with default parameters. Deeparg is a deep learning model that can predict ARGs from short-read metagenomic data; it uses DeepARG-DB, a merged database from 3 databases: Antibiotic Resistance Genes Database [ARDB], Comprehensive Antibiotic Resistance Database [CARD], and UniProt. This database was downloaded in December 2021. We used the relative abundance of ARGs normalized by the 16S rRNA gene content in the sample. We assessed the relative abundances of 634 known ARGs conferring resistance to 34 antibiotic classes. To identify associations between these ARGs and

gut microbes, we ran a multiple factor analysis (MFA) with the 37 most dominant species from a PCA (20 highest coordinates on both axes) and all the ARGs.

SXT ICE identification. We used Bowtie2/2.3.5 with the 'very-sensitive' option to map short reads against the two most prevalent SXT ICEs in Bangladesh at the time of our sampling (*26*): ICEVchind5 (NCBI accession GQ463142.1) and ICEVchind6 (accession MK165649.1) reference sequences downloaded from NCBI. The ICE was considered as present in a metagenome when 90% of its length was covered by at least one read. Using this criterion, 144 samples (59%) were ind5+, 26 (10.6%) ind6+, and 54 (22.1%) ICE-.

Statistical analyses. All statistics and visualizations were done in R studio version 1.2.5042.To reduce dimensions of the species composition table, we ran a principal component analysis (PCA) on the Hellinger transformed abundance data (5719 species*344 samples) (decostand function and rda function in the vegan R library) and selected the 20 most dominant species on each axis (37 in total).

To understand relationships between taxa in the microbiome, phages, antibiotics and patient metadata, we used a Redundancy analysis (RDA) on bacterial species abundances (rda function from vegan R package). Only the 7 most dominant species identified with the PCA were included: *v*, *Escherichia coli*, *Bacteroides vulgatus*, *Bifidobacterium longum*, *Shigella flexneri*, *Bifidobacterium breve*, and *Streptococcus mitis*. As explanatory variables, we used three antibiotics: azithromycin (AZI), ciprofloxacin (CIP), Doxycycline (DOX) as well as the three phages (ICP1, ICP2, ICP3), along with the most contributing patient metadata, selected with forward selection method (ordiR2step function from the vegan R library). These metadata are: the area where the sample was collected, date of sampling, age in years, vomiting state (yes or no) and dehydration status (severe, moderate or mild). We began the forward selection with a model with phages and antibiotic concentrations (µg/ml). The RDA was run on log-chord transformed abundances, and a permutation test with 999 iterations was used to assess the statistical significance of both the model and of each explanatory variable (anova.cca function from vegan package in R). The explained variation R^2 and adjusted R^2 were estimated with the RsquareAdj function in the vegan R package.

To identify species associated with each degree of dehydration, we used the indicator species analysis (*27*) as implemented in the multipatt R function from the indicspecies R library with (no group combination, duleg parameter set to TRUE and 9999 permutations) on the log-chord transformed species table (334 samples, 37 dominant species from the PCA) using the decostand and rda functions from the R vegan library). Reported P-values correspond to a permutation test with 999 iterations.

To study associations between ARGs and species, we ran a Multiple factor analysis (MFA) (*28*) using the mfa function from the FactoMineR R package, with the 37 most dominant species and all 634 resistance genes identified by deepARG. The visualization was done with the fviz_mfa_var function from the factoextra R library.

To model the relationships between phage, antibiotics and *V. cholerae* within each patient, we fit a generalized additive model (GAM) with *V. cholerae* relative abundance as a function of ICP1, antibiotics, and the interaction between them. We added the degree of dehydration as a random effect because it improved the model (smallest Akaike information criteria (AIC) compared to the other models). We fit several GAMs with different combinations of predictors: from a model with all antibiotics and their interaction with ICP1 to separate models with each antibiotic and its interaction with ICP1, and compared them based on their AIC (AICtab function from the bblme R package). The most parsimonious model was retained (the one with the smallest AIC). The GAMs were fitted using a beta error distribution with log-link function because *Vc* relative abundance is a continuous value between 0 and 1. We evaluated the selected model fit by inspecting residual distributions and fitted-observed value plots using the gam.check function from the mgcv R package. All P-values reported for the GAMs correspond to the Chi-square test from gam.summary function from mgcv R package. We also reported the adjusted R^2 (from the same function) as an evaluation of the goodness of fit (**Table S6**). To support the results from the models fitted on metagenomic relative abundances, we repeated the same methods with the qPCR-based absolute abundances of *Vc* and ICP1. Since the response (*Vc* absolute abundance from qPCR) is a continuous variable, we used additive models (Gaussian GAMs) after log-transforming the response (Figure S10, Table S7).

We tested whether phages or antibiotics select for potentially adaptive mutations in *Vc* by fitting generalized linear mixed models (GLMMs) (glmmtmb function from glmmTMB R package) with phages and antibiotics as predictors of the number of high-frequency nonsynonymous SNVs in the *V. cholerae* genome. We added *Vc* abundance as a fixed effect to the model to control for any coverage effects. We focused on higher-frequency SNVs (>10% within a sample) as more likely to be beneficial, and nonsynonymous SNVs as those more likely to have fitness effects. We fit several models with different combinations of predictors: from a model with all antibiotics and their interaction with ICP1 to separate models with each antibiotic and its interaction with ICP1, and compared them based on their AIC (AICtab function from bblme R package). The most parsimonious model was retained (the one with the smallest AIC). Because the response is count data, we fit three different count GLMMs: Poisson, negative binomial1 and negative binomial2, implemented in the glmmTMB R package, and then selected the model with the lowest AIC as

described (*29*). The selected model was fitted with the nbinom2 error distribution. All continuous predictors were standardized to zero mean and unit variance before analyses to improve the model's convergence. We evaluated the selected model fits by inspecting the residuals using the DHARMa library in R (simulateResiduals and plot functions). All the *P*-values reported for the GLMMs correspond to the Wald *P*-values reported by the glmm.summary function from the R package glmmTMB. As an evaluation of the goodness of fit, we compared the GLMMs and the corresponding null models (the same model but with no fixed effects other than the intercept) with likelihood-ratio tests (anova function from the stats R package), and reported the corresponding P-values (**Table S8**). We also reported the marginal R^2 as a measure of the variance explained by fixed effects, estimated with the r2 function from the performance R package. We used GLMMs because they are currently more flexible than GAMs in the range of count models that they can fit (https://bbolker.github.io/mixedmodels-misc/glmmFAQ.html). Furthermore, they can deal with overdispersion with two versions of negative binomial distributions: negative binomial 1 and negative binomial 2, respectively with linear and quadratic parametrization (*30*).

To further test the hypothesis that phages or antibiotics select for adaptive mutations in *V. cholerae*, we examined the relationship of the frequency of nonsynonymous SNVs in *V. cholerae* with phages and antibiotics. We fit a generalized additive model (GAM) (gam function in mgcv R library) with the average frequency of nonsynonymous SNVs as a function of ICP1, antibiotics, and their interactions, and *Vc* and ICP1 relative abundances from metagenomic data. We also included the fixed effect of ICE presence/absence as another factor that could provide phage or antibiotic resistance as well as mutation type to differentiate among non-synonymous (NS), synonymous (S), and intergenic (I) mutations. We fit GAMs with all antibiotics and their interaction with ICP1, as well as several simpler models with each antibiotic separately, and compared them based on their AIC. The GAMs were fitted using a beta error distribution with a log-link function because the average minor allele frequency of nonsynonymous SNVs is a continuous value between 0 and 0.5. We evaluated the selected model fit by inspecting residual distributions and fitted-observed value plots using the gam.check function from the mgcv R package. All the P-values reported for the GAMs correspond to the Chi-square test from the gam.summary function of the mgcv R package. We also reported the adjusted R2 (from the same function) as an evaluation of the goodness of fit (**Table S9**).

FIGURE S1

Fig S1: Inferred *Vibrio* phages in 224 metagenome assemblies. (a) At least one predicted *Vibrio* phage contig matched a CTXphi, ICP1, ICP3, TLC, ICP2 and K139 genome using BLASTn in 189, 63, 28, 13, 4 and 1 samples, respectively. **(b)** Phages ICP3, ICP2 and K139 are unambiguously assigned using vContact2. Contigs matching the CTXphi reference genome using BLASTn clustered either with the CTXphi reference viral cluster, or formed singletons (observed in only one sample). Similarly, most contigs matching an ICP1 reference genome using BLASTn were assigned by vContact2 to Viral Cluster 4, which did not include any reference phage genomes, but numerous connections to the reference ICP1 genome. **(c)** For 321 inferred *Vibrio* phage contigs from 210 samples, there was no BLASTn hit to a known *Vibrio* phage. Of these, 245 formed singletons, 18 were labeled as 'outliers' and one was placed between two viral sequence clusters. The remaining 57 contigs with no BLASTn hit clustered into 16 viral clusters, 14 of which exclusively comprised '*Vibrio* phages' identified in this analysis (mean cluster size = 2.8 contigs, range 2-5). Eighteen contigs with no BLASTn hit clustered with the TLC satellite phage (which flanks the CTXphi prophage in the *V. cholerae* genome), and one each with CTXphi and in a cluster predominantly including *Erwinia* phages.

FIGURE S2

Fig S2. Correlations between qPCR- and metagenomics-based quantification of *Vc* and ICP1. Raw qPCR cycle threshold (Ct) values plotted against relative abundances in metagenomes for (A) *Vc* and (B) ICP1. Inferred number of (C) *Vc* colony forming units (CFU) per ml of stool and (D) ICP1 plaque forming units (PFU) per ml, both at a Ct threshold of 28. Red lines show Pearson correlations, with coefficients and *p*-values noted in the figure panels.

FIGURE S3

Fig S3. ICP1:*V. cholerae* ratios among patients with different dehydration status binned by self-reported duration of diarrhea. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test, adjusted for multiple tests using the Benjamini-Hochberg (BH) method. Only significant (*P*<0.05) and marginally significant *P*-values (<0.1) are shown. 323 samples with *V. cholerae*>0% of metagenomic reads were included, with 165 from severe, 128 from moderate, and 30 from mild cases. The solid horizontal line is the median and the boxed area is the interquartile range.

FIGURE S4

Fig S4. The ICP1:*V. cholerae* ratio quantified with qPCR is higher in patients with mild dehydration. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test, adjusted for multiple tests using the Benjamini-Hochberg (BH) method. Only significant *P*-values (<0.05) are shown. Only 248 samples with *V. cholerae*>0% from qPCR data were included to exclude zeroes from the denominator of the ratio, with 138 from severe, 94 from moderate, and 16 from mild cases. A pseudocount of one was added to the ratio before log transformation. This figure is a version of Figure 1B using qPCR data.

FIGURE S5

Fig. S5. Principal component analysis (PCA) of the taxonomic composition of cholera patient stool samples. This analysis was run to reduce species composition dimensions from 5719 annotated with kraken2/braken to the 37 most dominant (with the 20 most dominant on the two first axes). Circles indicate samples; arrows indicate species. Samples are colored by their *V. cholerae* and ICP1 relative abundances (percentage of metagenomic reads). Yellow: patients with *V. cholerae* < 0.5% and ICP1 < 0.1%, red: *V. cholerae* < 0.5% and ICP1 >= 0.1%, green: *V. cholerae* >= 0.5% and ICP1 < 0.1%, purple: *V. cholerae* >= 0.5% and ICP1 >= 0.1%.

Fig S6. The two first axes from the RDA on the 7 most dominant species identified with the PCA (Figure S4), phages (ICP1, ICP2, ICP3), patient metadata, and antibiotic concentrations. Three quantified antibiotics are represented: azithromycin (AZI), ciprofloxacin (CIP), Doxycycline (DOX). Patient metadata are: the area where the sample was collected, date of sampling, age in years, vomiting state (yes or no) and dehydration status (severe, moderate or mild). All variables are shown here; the collection date and area code were omitted in Fig. 1C for clarity.

FIGURE S7

Fig. S7. Distribution of the relative abundance of *V. cholerae*-associated ARGs in patients with different exposures to azithromycin (AZI). AZI>=MIC (detected; D) and AZI<MIC (not detected; ND) under anaerobic conditions (A) or aerobic conditions (B). AZI MIC=1 µg/ml under aerobic conditions and 8 µg/ml under anaerobic conditions. The Y-axis is the relative abundance of ARGs in metagenomes normalized by 16S rRNA gene reads and by *V. cholerae* reads. Only V. cholerae-positive samples are plotted (V. cholerae>0 reads). P-values are from a Wilcoxon test. No comparisons (D vs. ND) are significant (*P*<0.05) after BH correction for 24 tests.

FIGURE S8

Fig. S8. Distribution of the relative abundance of *V. cholerae*-associated ARGs in patients with different exposures to ciprofloxacin (CI). CIP>=MIC (detected; D) and CIP<MIC (not detected; ND) under anaerobic conditions (A) or aerobic conditions (B). CIP MIC=0.016 µg/ml under aerobic conditions and 0.063 µg/ml under anaerobic conditions. The Y-axis is the relative abundance of ARGs in metagenomes normalized by 16S rRNA gene reads and by *V. cholerae* reads. Only *V. cholerae*-positive samples are plotted (*V. cholerae*>0 reads). *P*-values are from a Wilcoxon test. In red: BH corrected *P*<0.05 after correction for 24 tests.

Fig S9. MFA showing correlations between bacterial species and antimicrobial resistance genes (ARGs) in patient gut microbiomes. The 37 most dominant species from the PCA (Fig S5) and all the resistance genes annotated with deepARG were included in the ordination. 12 ARGs clustered with *V. cholerae*: OMPU, OMPT, VARG, TET34, TET35, DFRA1, DFRA16, FLOR, Chloramphenicol exporter, Aph6, Aph3, MEXI. Only the greatest correlations on each axis are shown for clarity.

FIGURE S10

Fig S10. Interactions between *V. cholerae*, phage ICP1, and azithromycin tested with qPCR data. (A) *V. cholerae* declines in absolute abundance with higher relative abundance of azithromycin (AZI) in µg/ml. (B) The relationship between ICP1 and *V. cholerae* abundances estimated with qPCR is affected by AZI concentration (µg/ml); the illustrated AZI concentrations show regular intervals between the minimum (0 µg/ml) and maximum (78 µg/ml) observed values. GAM results fit to data from 329 samples with qPCR data. The response variable (*Vc* absolute abundance from qPCR) was log-transformed after adding a pseudocount of 1 and a Gaussian distribution was used. The predictors are ICP1 abundance from qPCR and antibiotics concentration (µg/ml). As with metagenomics data, we fit several GAMs with different combinations of predictors: from a model with all antibiotics and their interaction with ICP1 to separate models with each antibiotic and its interaction with ICP1 and compared them based on their Akaike information criteria (AIC). The most parsimonious model (shown) was retained (the one with the smallest AIC). Model details are provided in Table S7.

FIGURE S11

Fig S11. A) Breadth of metagenomic read coverage of the ICEs ind5 and ind6. Samples were identified as ind5-positive or ind6-positive based on read mapping against the reference ICE sequences. The ICE was considered as present when 90% of the reference ICE length was covered by at least one read. Only 2 samples had 100% ind5 and >90% ind6, which were considered ind5-positive. B) Boxplot showing that ICE-negative samples are not associated with lower relative abundance of *V. cholerae*, suggesting that the lack of detecting ICEs is not due to low *Vc* abundance. The *P*-value is from a Kruskal-Wallis test.

FIGURE S12

Fig S12. (A) Distribution of ICP2:*Vc* ratios across patients with different ICE profiles. (B) The same data as (A) binned into boxplots according to dehydration status. (C) Distribution of ICP3:*Vc* ratios across patients with different ICE profiles, (D) The same data as (C) binned into boxplots according to dehydration status. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test adjusted with the Benjamini-Hochberg (BH) method. Only *P*-values < 0.1 are shown. Only samples with sufficient *Vc* or ICP were included. For clarity, the Y-axes were log10 transformed after adding one to the ratios. The solid horizontal line is the median and the boxed area is the interquartile range.

FIGURE S13

Fig S13. Integrative conjugative elements (ICEs) are associated with lower ICP1:*V. cholerae* ratios using qPCR data. (A) Distribution of ICP1:*Vc* ratios across patients with different ICE profiles. (B) The same data as (A) binned into boxplots according to dehydration status. (C) Distribution of ICP2:*Vc* ratios. (D) The same data as (C) binned into boxplots according to dehydration status. (E) Distribution of ICP3:*Vc* ratios, (F) The same data as (E) binned into boxplots according to dehydration status. (G) Distribution of phage:*Vc* ratios, including the sum of all phages (ICP1, ICP2, ICP3). (H) The same data as (G) binned into boxplots according to dehydration status. Only samples with appreciable *Vc* or ICP1 were included (215 samples with *Vc*>0 or phages >0 from qPCR data), of which 50 samples were ICE-, 25 were ind6+ and 140 were ind5+. The Y-axes were log10 transformed after adding one to the ratios to improve readability. The solid horizontal line is the median and the boxed area is the interquartile range. These results support the findings based on metagenomic data shown in Figure 3 and S12 using qPCR data.

FIGURE S14

Fig S14. Mixed infections by more than on *V. cholerae* strain is unlikely in our patients. (A) The distribution of the number of distinct strains detected across samples. In 260/344 samples, strainGST identified only one reference strain. In 83/344 samples, no reference strain could be identified with confidence, likely due to low coverage of *V. cholerae* (these samples all had <1% <u>Vc</u> reads). (B) The relative abundance of *Vc* (% reads) in samples with zero, one, or two strains identified. In one sample, strainGST identified two reference strains, but with a low confidence score and at low *Vc* abundance.

FIGURE S15

Fig S15. A) Low *V. cholerae* relative abundance is associated with DNA repair mutations independently of the number of SNVs. Mutators are defined as having a high (H) number of SNVs (25 or more) in the *V. cholerae* genome, along with one or more nonsynonymous mutations in a DNA repair gene (+) resulting in a predicted defect in DNA repair. Non-mutators have neither a high number of SNVs nor a DNA repair defect. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test adjusted with the Benjamini-Hochberg (BH) method. Only *P*-values < 0.1 are shown. (B and C) Transversion mutations, particularly G->T and C->A, are more common in mutators compared to non-mutators. D) Relative abundance of phage ICP1 in samples with different *V. cholerae* mutation profiles. Kruskall-wallis test (*P*= 0.07197). n=133, with 47 belonging to mutators, 70 to non-mutators, 14 to "high SNVs/no DNA repair defect" and 2 to "low SNVs/with DNA repair defect" groups.

FIGURE S16

Fig S16. ICE- are readily distinguishable from ind5+ and ind6+ samples. Breadth of ICE coverage across patients with different ICE profiles. (A) Breadth of ind5 coverage. (B) Breadth of ind6 coverage. Note that the few samples with ambiguous breadth of ICE coverage (in the 40-80% range) were not included in the InStrain analysis, and are not included here. *P*-values are from a Kruskal-Wallis test with Dunn's post-hoc test adjusted with the Benjamini-Hochberg (BH) method. n=131 with 24 belonging to ICE-, 18 to ind6+, and 89 to ind5+ groups.

Fig S17. Boxplots of mutation frequency in the presence or absence of ICP1 and/or ICEs. Boxplots include 128 samples, of which 41 are ICP1+ (ICP1 detected by qPCR at a Ct cutoff of 28) and 87 are ICP- (Ct below 28). The Y-axes were log10 transformed after adding a pseudocount of 1 to improve readability. The solid horizontal line is the median and the boxed area is the interquartile range. This figure is a version of Fig 4C with qPCR data.

FIGURE S18

Fig S18. Number of SNVs in ICP1 as a function of the ICP1:*Vc* ratio and the ICE type encoded by *Vc*. Single nucleotide variant (SNV) count per sample in the ICP1 genome is plotted as a function of the ICP1:*Vc* ratio for different ICE profiles. All 45 samples for which mutations in ICP1 were called with inStrain were included in this analysis, of which 18 ICE-, 20 ind5 and 7 ind6. Non-synonymous (NS), synonymous (S), and intergenic (I) SNVs are plotted in separate rows. The only significant correlation (*p* < 0.05; top right) is shown with the trendline in gray and Spearman correlation coefficient and *p*-value.

FIGURE S19

Fig S19. SNVs in ICP1 as a function of the ICP1:Vc ratio and the ICE type encoded by *Vc*. The mean single nucleotide variant (SNV) frequency per sample in the ICP1 genome is plotted as a function of the ICP1:*Vc* ratio for different ICE profiles. This analysis includes 29 samples for which high frequency mutations in ICP1 were called by inStrain, of which 11 ICE-, 13 *ind5* and 5 *ind6*. Non-synonymous (NS), synonymous (S), and intergenic (I) SNVs are plotted in separate rows. The only significant correlation ($p <$ 0.05; top right) is shown with the trendline in gray and Spearman correlation coefficient and *p*-value. Only SNV frequencies >0.1 are included.

SUPPLEMENTARY TABLES

Table S1. Primers used for PCR and qPCR.

Target	Primer Name	Sequence 5'-3'	Reference
ompW (Vibrio cholerae)	ompW_F	CACCAAGAAGGTGACTTTAT	(*31*)
	ompW_R	GAACTTATAACCACCCGCG	
tcpA (Vibrio cholerae)	*tcpA set2*_F	ACACGATAAGAAAACCGGTCA	(*4*)
	*tcpA set2*_R	GCCTTGGTCATATTCTGCGA	
ICP1	gp58.2_F	CAAAGGCAGCAGGTAGGACA	This study
	gp58.2_R	CCCTTCAAGCCGTAGTTGGT	
ICP2	gp24_F	AGAAGTCGCAAACGGGGTAC	This study
	gp24_R	AACGTGGTTCTCGTGAGTGG	
ICP3	gp19_F	AGACCAACGCCGACTGTTAG	This Study
	gp19_R	CGATACCACGGAAAGCCTGT	
16S rDNA1	Maeda_1048_1067_F	GTGSTGCAYGGYTGTCGTCA	(*6*)
	Maeda_1175_1194_R A	ACGTCRTCCMCACCTTCCTC	

Table S2. LC MS/MS targets and parameters.

Antibiotic	RT (min)	Extraction Ions
Azithromycin	11.4 – 13.1	750,591
Ciprofloxacin	9.5 – 11.2	332
Doxycycline	8.9 – 14.0	445,428
Nalidixic Acid	12.3 – 14.4	233
Azithromycin-d5	11.4 – 13.1	754,595
Ciprofloxacin-d8	9.5 – 11.2	339
Doxycycline-d5	8.9 – 14.0	450,433
Metronidazole	3.5-5.5	172

Table S3. Indicator species analysis. For each group, we report the indicator value (stat) ranging between 0-1, with 1 being the perfect indicator species meaning that the species occurs exclusively in one group. P-values are from a permutation test with 9999 iterations as implemented in the indicator species function in R (multipatt from indicspecies library). The analysis was run on all the data (344 samples) and the 37 most dominant species from the PCA in Figure S5, and 24 species were selected as indicators of the patient's dehydration status.

Group	stat	*p*
Group 1: mild dehydration		
1. *Bifidobacterium longum*	0.372	**1e-05**
2. *Bifidobacterium breve*	0.304	**6e-05**
3. *Escherichia coli*	0.266	**0.00057**
4. *Enterococcus aecium*	0.214	0.00437
5. *Bifidobacterium kashiwanohense*	0.2	0.00818
6. *Salmonella enterica*	0.183	0.01262
7. ICP1	0.167	**0.02641**
Group 2: moderate dehydration		
1. *Streptococcus pasteurianus*	0.23	0.00272
2. *Roseburia intestinalis*	0.21	0.00596
3. *Streptococcus gallolyticus*	0.16	0.03152
Group 3: severe dehydration		
1. Vibrio cholerae	0.333	**6e-05**
2. *Fusobacterium mortiferum*	0.31	0.00011
3. *Faecalibacterium prausnitzii*	0.291	0.00020
4. *Prevotella ruminicola*	0.284	0.00028
5. *Prevotella dentalis*	0.281	0.00030
6. *Brachyspira pilosicoli*	0.249	0.00084
7. *Collinsella aerofaciens*	0.246	0.00149
8. *Bacteroides thetaiotaomicron*	0.234	0.00221
9. *Bacteroides vulgatus*	0.226	**0.00309**
10. *Prevotella intermedia*	0.209	0.00560
11. *Bacteroides fragilis*	0.201	0.00823
12. *Roseburia[Eubacterium] rectale*	0.194	0.01063
13. ICP3	0.189	**0.01050**
14. *Roseburia lachnospiraceae bacterium* GAM79	0.168	0.02409

Table S4. Variables used in the Redundancy analysis (RDA) showing relationships among the seven most dominant bacterial species and explanatory variables: phages, patient metadata, and antibiotic concentration. First column: explanatory variables (CIP:, Ciprofloxacin, DOX:Doxycycline, and AZI: Azithromycin), second column: p-values from a permutation test with 999 iterations.

Variable	p^a
CIP	0.073
DOX	0.701
AZI	0.001***
ICP1	0.012*
ICP2	0.336
ICP3	0.104
Area_Code	0.001***
Dehydration status	0.001***
Collection date	0.003**
Vomiting	0.002**
Age_in_Years	0.006**

[a] Permutation test (anova function from the vegan R package).

Table S5. Antibiotic resistance thresholds. Minimal inhibitory concentrations (MICs) were established under aerobic or anaerobic conditions in (Creasy-Marrazzo et al. 2022). Concentrations are in units of µg/ml.

	Aerobic		Anaerobic
Ciprofloxacin	0.016		0.063
Azithromycin	1		8
Doxycycline	0.13		0.13

Table S6. Generalized additive models included in the model selection using metagenomic data. Generalized additive models included in the model selection. GAMs were fit with *Vc* relative abundance as a function of ICP1, antibiotics and their interactions. The selection was based on ∆AIC. We report ∆AIC, R syntax for the formula, the predictors (fixed effects) and the corresponding P-values (Chi-square test), as well as the *P*-value corresponding to dehydration random effect (RE) and the adjusted r-squared of the model. nt: not tested.

model	∆AIC	Formula	ICP1	AZI	ICP1*AZI	Dehyd (RE)	R2
Gam1 (selected)	0	Vc ~ s(ICP1) + s(AZI) + te(ICP1, AZI) + s(Dehydration_Status, bs = "re")	0.614	0.002	0.026	3.66e-07	0.031
Gam2	1.3	Vc ~ s(ICP1) + s(AZI) + s(Dehydration_Status, bs = "re")	0.077	0.004	Not tested	2.76e-07	0.035
Gam3	3.9	Vc ~ s(ICP1) + s(AZI) + s(CIP) + s(DOX) + te(ICP1, AZI) + te(ICP1, CIP) + te(ICP1, DOX) +s(Dehydration_Status, bs = "re")	0.606	0.002	0.025	4.08e-07	0.027
Gam4	5.2	Vc ~ s(ICP1) + s(AZI) + s(CIP) + s(DOX) + s(Dehydration_Status, bs = "re")	0.078	0.003	nt	3.06e-07	0.032
Gam5	11.7	Vc ~ s(ICP1) + s(DOX) + te(ICP1, DOX) + s(Dehydration_Status, bs = "re")	0.059	nt	nt	3.65e-08	0.035
Gam6	11.9	Vc ~ s(ICP1) + s(CIP) + te(ICP1, CIP) + s(Dehydration_Status, bs = "re")	0.664	nt	4.35e-08	4.35e-08	0.022
Gam7	12.4	Vc ~ s(ICP1) + s(AZI) + te(ICP1, AZI, by = Dehydration_Status)					
Gam1(B)[a]	29.3	Vc ~ s(ICP1) + s(AZI) + te(ICP1, AZI)	0.914	0.003	0.015	nt	-0.017

[a] Without dehydration RE

Table S7. Generalized additive models included in the model selection using qPCR data. Generalized additive models included in the model selection. GAMs were fit with *Vc* as a function of ICP1 (absolute abundances from qPCR with CT=28), antibiotics and their interactions, the same methods as for the models fit with metagenomics data were used. The selection was based on ∆AIC. We report ∆AIC, R syntax for the formula, the predictors (fixed effects) and the corresponding P-values (Chi-square test), as well as the *P*-value corresponding to dehydration random effect (RE) and the adjusted r-squared of the model. nt: not tested.

model	∆AIC	formula	ICP1	AZI	ICP1*AZI	Dehyd(RE)	R2
Gam3	0	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + s(CIP) + s(DOX) + te(qPCR_ICP1_PFUml_CT28, AZI) +te(qPCR_ICP1_PFUml_CT28, CIP) + te(qPCR_ICP1_PFUml_CT28, DOX) + s(Dehydration_Status,bs = "re")	0.622	0.001	0.017	< 2e-16	0.172
Gam4	0.4	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + s(CIP) + s(DOX) + s(Dehydration_Status, bs = "re")	0.023	0.001	nt	< 2e-16	0.172
Gam1	0.9	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + te(qPCR_ICP1_PFUml_CT28, AZI) + s(Dehydration_Status, bs = "re")	0.636	0.001	0.015	< 2e-16	0.163
Gam2	1.3	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + s(Dehydration_Status, bs = "re")	0.018	0.001	nt	< 2e-16	0.163
Gam7	5.3	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + te(qPCR_ICP1_PFUml_CT28, AZI, by = Dehydration_Status)	0.265	0.004	<0.003 *	nt	0.169
Gam5	11.8	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(DOX) + te(qPCR_ICP1_PFUml_CT28, DOX) + s(Dehydration_Status, bs = "re")	0.019	nt	nt	< 2e-16	0.136
Gam6	11.9	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(CIP) + te(qPCR_ICP1_PFUml_CT28, CIP) + s(Dehydration_Status, bs = "re")	0.786	nt	nt	< 2e-16	0.134
Gam8 (Gam1 without RE)	30.6	log_qpcr_vc ~ s(qPCR_ICP1_PFUml_CT28) + s(AZI) + te(qPCR_ICP1_PFUml_CT28, AZI)	0.393	0.0001	0.031	nt	0.079

*For all dehydration status levels

Table S8. Generalized linear mixed models. GLMMs with *Vc* SNV count as a function of phage relative abundance from metagenomics and antibiotics and their interaction. We report ΔAIC, the R syntax for the formula, the predictors and the corresponding P-values (Wald test). As an evaluation of the goodness of fit, we report the adjusted R-squared (R2) and *P*-value from the comparison of the models and null models (exactly the same model but with no fixed terms) (likelihood-ratio test, anova function from stats package in R). Antibiotics terms are not shown (not significant, *P*>0.05, GLMM, wald test). nt : not tested.

Model	dAIC	Formula	Vc	ICP1	ICP1*Vc	R2	P/null model
Mod1 (Selected)	0.0	SNV_nb ~ Vc + Vc:ICP1	0.002	nt	0.004	0.377	7e-04
Mod2	0.6	SNV_nb ~ ICP1 + Vc + Vc:ICP1	0.007	0.266	0.012	0.408	1.19e-03
Mod3	1.6	SNV_nb ~ ICP1 + Vc + AZI + Vc:ICP1	0.008	0.267	0.015	0.409	0.002
Mod4	3.2	SNV_nb ~ ICP1 + Vc + CIP + AZI + Vc:ICP1	0.007	0.273	0.015	0.412	0.004
Mod5	5.0	SNV_nb ~ Vc	0.002	nt	nt	0.228	0.006
Mod6	5.2	SNV_nb ~ ICP1 + Vc + CIP + DOX + AZI + Vc:ICP1	0.00745	0.27359	0.01554	0.412	0.008
Mod6	5.9	SNV_nb ~ ICP1 + Vc + CIP + DOX + AZI + ICP1:CIP + ICP1:AZI + Vc:ICP1	0.0428	0.1634	0.0385	0.458	0.008
Mod7	7.0	SNV_nb ~ ICP1 + Vc + CIP + AZI + DOX + Vc:ICP1:CIP + Vc:ICP1:AZI + Vc:ICP1:DOX	0.015	0.732	nt	0.442	0.012
Mod8	7.1	SNV_nb ~ ICP1 + Vc + CIP + DOX + AZI + ICP1:AZI + Vc:ICP1	0.047	0.275	0.059	0.422	0.015
Mod9	12.1	SNV_nb ~ ICP1	nt	0.506	nt	0.018	0.527

Table S9. Generalized additive models. GAMs used to select the most parsimonious model. The response is the average frequency of NS mutations in *V. cholerae* and the predictors are ICP1 (relative abundance from metagenomics), antibiotics, ICE presence/absence and mutation type (NS: non-synonymous, S:synonymous and I:intergenic) as well as their interactions. We defined a variable (ICE.by.mut) as the combination between ICE and mutation type with 9 levels (mutation type*ICE) to represent the interaction between mutation type and ICE factors. Model selection was based on ΔAIC. We report ΔAIC, the R syntax for the formula, the predictors and the corresponding P-values (Chi-square test) and the adjusted r-squared of the model. nt: not tested.

	ΔAIC	formula	P (ICP1*ICE*mut)			P (antbx *mut)	R2
			Ind5*NS	Ind6*NS	ICE-*NS		
Mod1	0.0	mean_freq ~ s(ICP1, by = ICE.by.mut)	0.1336	0.642	0.0469	nt	0.021
Mod2	5.8	mean_freq ~ s(ICP1, by = ICE.by.mut) + s(AZI, by = mutation_type) + s(CIP, by = mutation_type) + DOX * mutation_type	0.13242	0.58891	0.04677	>0.05	0.060
Mod3	9.0	mean_freq ~ s(ICP1, by = ICE.by.mut) + s(AZI, by = mutation_type) + s(CIP, by = mutation_type) + DOX * mutation_type + te(ICP1, AZI, by = mutation_type) + te(ICP1, CIP, by = mutation_type) + DOX:ICP1:mutation_type	<2e-16	<2e-16	<2e-16	<2e-16 (AZI)	0.067
Mod4	10.9	mean_freq ~ s(ICP1, by = ICE.by.mut) + s(AZI, by = mutation_type) + s(CIP, by = mutation_type)	0.1223	0.6346	0.0496	>0.05	0.002

Table S10. Most frequently mutated *Vc* genes in samples with high ICP1:*Vc* ratios.

Mutation count denotes the total number of SNVs and patient count denotes the number of patients in which a mutation was observed. Genes were identified with prodigal v2.6.3 and annotated with eggNOG-Mapper v2 as well as manually with blastx against NCBI database within the Basic Local Alignment Search Tool (BLAST) online tool. Genes that contain mutations in samples with ICP1>*Vc* exclusively are shown in bold. Only the top 15 genes are shown, See Data File S3 for complete list.

COG (eggnog)	PFAM (eggnog) annotation	NCBI annotation	Mutation count	Patient count
Q	HemolysinCabind_Peptidase_M10_C	retention module-containing protein WP_001191814.1	5	2
M	**Peptidase_S13**	**D-alanyl-D-alanine carboxypeptidase/D-alanyl-D-alanine-endopeptidase (genbank : AAF93798.1)**	**4**	**1**
V	ACR_tran	multidrug resistance protein, putative AAF93795.1	4	1
O	PPC_Peptidase_M9_Peptidase_M9_N	TPA: collagenase HAS4622795.1	4	1
P	BPD_transp_1	ABC transporter permease subunit WP_000252168.1	4	1
K	**MerR_1**	**MerR family transcriptional regulator WP_000226962.1**	**3**	**1**
C	CCG_FADoxidase_C_FAD_binding_4_Fer4_7_Fer4_8	FAD-binding and (Fe-S)-binding domain-containing protein WP_000188699.1	3	1
C	**Gp_dh_C_Gp_dh_N**	**glyceraldehyde 3-phosphate dehydrogenase GenBank: AAF96741.1**	**3**	**1**
S	DUF3302	DUF3302 domain-containing protein WP_000478180.1	2	1
IQ	PPbinding	Chain A, 3-oxoacyl-[acyl-carrier-protein] synthase 2 PDB: 4JRH_A	2	1
C	**Fer4_12_Radical_SAM**	**methyl-accepting chemotaxis protein WP_000383592.1**	**2**	**2**
Q	FtsX_MacB_PCD	ABC transporter permease WP_000645916.1	2	1
C	**CCG_Fer4_8**	**anaerobic glycerol-3-phosphate dehydrogenase subunit GlpC WP_001014995.1**	**2**	**1**
GM	CoA_binding_3_Polysacc_synt_2	nucleoside-diphosphate sugar epimerase/dehydratase WP_000494952.1	2	1
F	PpxGppA	guanosine-5'-triphosphate,3'-diphosphate diphosphatase WP_000076046.1	2	1

Table S11. Most frequently mutated *Vc* genes in samples with low ICP1:*V.c* ratios. Mutation count denotes the total number of SNVs and patient count denotes the number of patients in which a mutation was observed. Genes were identified with prodigal v2.6.3 and annotated with eggNOG-Mapper v2 as well as manually with blastx against NCBI database within the Basic Local Alignment Search Tool (BLAST) online tool. Genes that contain mutations in samples with ICP1<*Vc* exclusively are shown in bold. Only the top 15 genes are shown, see Data File S4 for complete list.

COG category (eggnog)	PFAM (eggnog) annotation	NCBI annotation	Mutation count	Patient count
O	PPC_Peptidase_M9_Peptidase_M9_N	TPA: collagenase HAS4622795.1	56	15
MQ	**ACD_ADPrib_exo_Tox_Anthrax_toxA_ Hydrolase_4_MLD_Peptidase_C80_Rtx A**	**TPA: MARTX multifunctional-autoprocessing repeats-in-toxin holotoxin RtxA HAS4620517.1**	**14**	**6**
PT	**GGDEF_Hemerythrin**	**GGDEF domain-containing protein WP_001190450.1**	**12**	**4**
T	**EAL_GGDEF**	**EAL domain-containing protein WP_000160066.1**	**11**	**2**
S	**Glyco_hydro_129**	**hypothetical protein VC_A0254 AAF96165.1**	**11**	**2**
S	**Betaprism_lec_Hemolysin_N_Leukocid in**	**Cytolysin vcc (ncbi) WP_001125271.1 (hlyA gene) [Levade et al 2021]**	**10**	**3**
C	FixO	cytochrome-c oxidase, cbb3-type subunit II WP_000097777.1	9	1
L	**Fapy_DNA_glyco_H2TH_zfFPG_IleRS**	**bifunctional DNA-formamidopyrimidine glycosylase/DNA-(apurinic or apyrimidinic site) lyase WP_001114647.1**	**9**	**3**
I	AcylCoA_dh_1_AcylCoA_dh_M_AcylCoA _dh_N_DUF1974	acyl-CoA dehydrogenase FadE WP_000404358.1	8	4
L	Phage_int_SAM_4	hypothetical protein WP_000222725.1	8	4
M	ACD_PG_binding_1_Pesticin_Phage_GP D	Type VI secretion system tip protein VgrG WP_000212125.1	7	1
K	LacI_Peripla_BP_3	DNA-binding transcriptional regulator CytR WP_000224452.1	7	1
T	EAL	EAL domain-containing protein WP_000547542.1	7	2
E	SDH_alpha_SDH_beta	sodium:solute symporter WP_001882525.1	7	4
S	Inovirus_Gp2	inovirus Gp2 family protein WP_000149629.1	7	4

Table S12. Most frequently mutated ICP1 genes in samples with ICE *ind5* detected. Mutation count denotes the total number SNVs (NS SNVs with frequency>0.1) and patient count denotes the number of patients in which a mutation was observed. Only samples with the ICE *ind5* are included (13 samples out of the 27 samples for which high frequency NS mutations were called in ICP1 with inStrain). Genes were annotated manually with blastx against NCBI database within the Basic Local Alignment Search Tool (BLAST) online tool. Most of the genes had no hit with eggNOG-Mapper. Only the top 15 genes are shown; see Data File S5 for the complete list (41 genes).

NCBI annotation	Mutation count	Patient count
hypothetical protein ViPhICP1_gp184 YP_004251125.1	33	3
hypothetical protein ViPhICP1_gp147 YP_004251088.1	5	2
anaerobic ribonucleotide reductase small subunit YP_004251055.1	4	3
hypothetical protein ViPhICP1_gp150 YP_004251091.1	4	2
acyl carrier protein YP_004251038.1	4	3
hypothetical protein ViPhICP1_gp226 YP_004251167.1	3	3
nicotinamide phosphoribosyl transferase YP_004251171.1	3	3
hypothetical protein ViPhICP1_gp139 YP_004251080.1	2	1
hypothetical protein ViPhICP1_gp166 YP_004251107.1	2	2
hypothetical protein ViPhICP1_gp018 YP_004250959.1	2	2
hypothetical protein ViPhICP1_gp175 YP_004251116.1	2	1
hypothetical protein ViPhICP1_gp199 YP_004251140.1	2	2
ribonucleoside diphosphate reductase, beta chain YP_004251147.1	2	2
hypothetical protein ViPhICP1_gp068 YP_004251009.1	2	2
hypothetical protein ViPhICP1_gp070 YP_004251011.1	2	1

REFERENCES

1. A. I. Khan *et al.*, Electronic decision support and diarrhoeal disease guideline adherence (mHDM): a cluster randomised controlled trial. *Lancet Digit Health* **2**, e250-e258 (2020).

2. A. I. Khan, F. Qadri, Epidemiology of cholera in Bangladesh: Findings from Nationwide Hospital-based Surveillance, 2014-2018. *CID*, (2019).

3. A. Creasy-Marrazzo *et al.*, Genome-wide association studies reveal distinct genetic correlates and increased heritability of antimicrobial resistance in *Vibrio cholerae* under anaerobic conditions. *Microbial Genomics* **8**, (2022).

4. L. Alexandrova *et al.*, Identification of Widespread Antibiotic Exposure in Patients With Cholera Correlates With Clinically Relevant Microbiota Changes. *J Infect Dis* **220**, 1655-1666 (2019).

5. J. A. Grembi, K. Mayer-Blackwell, S. P. Luby, A. M. Spormann, High-Throughput Multiparallel Enteropathogen Detection via Nano-Liter qPCR. *Front Cell Infect Microbiol* **10**, 351 (2020).

6. H. Maeda *et al.*, Quantitative real-time PCR using TaqMan and SYBR Green for Actinobacillus actinomycetemcomitans, Porphyromonas gingivalis, Prevotella intermedia, tetQ gene and total bacteria. *FEMS Immunol Med Microbiol* **39**, 81-86 (2003).

7. K. E. Flaherty *et al.*, High-throughput low-cost nl-qPCR for enteropathogen detection: A proof-of-concept among hospitalized patients in Bangladesh. *PLoS One* **16**, e0257708 (2021).

8. S. Roux *et al.*, iPHoP: An integrated machine learning framework to maximize host prediction for metagenome-derived viruses of archaea and bacteria. *PLoS Biol* **21**, e3002083 (2023).

9.   R. Cook *et al.*, INfrastructure for a PHAge REference Database: Identification of Large-Scale Biases in the Current Collection of Cultured Phage Genomes. *Phage (New Rochelle)* **2**, 214-223 (2021).

10.   F. Hassan, M. Kamruzzaman, J. J. Mekalanos, S. M. Faruque, Satellite phage TLCphi enables toxigenic conversion by CTX phage through dif site alteration. *Nature* **467**, 982-985 (2010).

11.   K. D. Seed, D. W. Lazinski, S. B. Calderwood, A. Camilli, A bacteriophage encodes its own CRISPR/Cas adaptive response to evade host innate immunity. *Nature* **494**, 489-491 (2013).

12.   H. Bin Jang *et al.*, Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat Biotechnol* **37**, 632-639 (2019).

13.   B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

14.   A. E. Minoche, J. C. Dohm, H. Himmelbauer, Evaluation of genomic high-throughput sequencing data generated on Illumina HiSeq and genome analyzer systems. *Genome Biol* **12**, R112 (2011).

15.   D. Li, C. M. Liu, R. Luo, K. Sadakane, T. W. Lam, MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**, 1674-1676 (2015).

16.   J. Alneberg *et al.*, Binning metagenomic contigs by coverage and composition. *Nat Methods* **11**, 1144-1146 (2014).

17.   D. D. Kang *et al.*, MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

18.   C. M. K. Sieber *et al.*, Recovery of genomes from metagenomes via a dereplication, aggregation and scoring strategy. *Nat Microbiol* **3**, 836-843 (2018).

19.   M. R. Olm, C. T. Brown, B. Brooks, J. F. Banfield, dRep: a tool for fast and accurate genomic comparisons that enables improved genome recovery from metagenomes through de-replication. *ISME J* **11**, 2864-2868 (2017).

20.   M. R. Olm, A. Crits-Christoph, K. Bouma-Gregson, B. A. Firek, M. J. Morowitz, J. F. Banfield, inStrain profiles population microdiversity from metagenomic data and sensitively detects shared microbial strains. *Nat Biotechnol* **39**, 727-736 (2021).

21.   D. Hyatt, G. L. Chen, P. F. Locascio, M. L. Land, F. W. Larimer, L. J. Hauser, Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

22.   C. P. Cantalapiedra, A. Hernandez-Plaza, I. Letunic, P. Bork, J. Huerta-Cepas, eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Mol Biol Evol* **38**, 5825-5829 (2021).

23.   L. R. van Dijk *et al.*, StrainGE: a toolkit to track and characterize low-abundance strains in complex microbial communities. *Genome Biol* **23**, 74 (2022).

24.   A. Jolivet-Gougeon *et al.*, Bacterial hypermutation: clinical implications. *J Med Microbiol* **60**, 563-573 (2011).

25.   G. Arango-Argoty, E. Garner, A. Pruden, L. S. Heath, P. Vikesland, L. Zhang, DeepARG: a deep learning approach for predicting antibiotic resistance genes from metagenomic data. *Microbiome* **6**, 23 (2018).

26.   K. LeGault *et al.*, Temporal shifts in antibiotic resistance elements govern phage-pathogen conflicts. *Science* **373**, 2020.2012.2016.423150 (2021).

27.   M. Dufrene, Pireere Legendre, Species Assemblages and Indicator Species: The Need for a Flexible Asymmetrical Approach. *Ecological Monographs* **67**, 345-366 (1997).

28.   P. J., Analyse Factorielle Multiple Appliquée Aux Variables Qualitatives et Aux Données Mixtes. *Revue de Statistique Appliquee* **4**, 5-37 (2002).

29.   M. E. Brooks *et al.*, Modeling zero-inflated count data with glmmTMB. *bioRxiv*, 132753

        (2017).

30.   J. W. Hardin, J. M. Hilbe, *Generalized Linear Models and Extensions*. (Stata Press), vol.

        Vol. 4th ed.

31.   B. Nandi, R. K. Nandy, S. Mukhopadhyay, G. B. Nair, T. Shimada, A. C. Ghose, Rapid

        method for species-specific identification of *Vibrio cholerae* using primers targeted to the

        gene of outer membrane protein OmpW. *J Clin Microbiol* **38**, 4145-4151 (2000).

DATA FILES

File S1. Patient metadata.

File S2. Antibiotic concentrations in stool samples.

File S3. Full list of *Vc* genes containing mutations in samples where ICP1 > *Vc* (% of reads).

File S4. Full list of *Vc* genes containing mutations in samples where ICP1 < *Vc* (% of reads).

File S5. Full list of ICP1 genes containing mutations in samples containing ICE *ind5*.