Hexokinase regulates Mondo-mediated longevity via the PPP and organellar dynamics

Raymond Laboy[1,2], Marjana Ndoci[1], Syed Shamsh Tabrez[1], Maximilian Vonolfen[1,2], Eugen

Ballhysa[1,2], Tim Droth[1], Klara Schilling[1,2], Anna Löhrke[1], Ilian Atanassov[1], Adam Antebi[1,2*]

1 Department of Molecular Genetics of Ageing, Max Planck Institute for Biology of Ageing,

Cologne, Germany

2 Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases

(CECAD), University of Cologne, Germany

*For correspondence: aantebi@age.mpg.de

Abstract

The transcriptional complex Mondo/Max-like, MML-1/MXL-2, acts as a convergent

transcriptional regulatory output of multiple longevity pathways in *Caenorhabditis elegans*.

These transcription factors coordinate nutrient sensing with carbohydrate and lipid metabolism

across the evolutionary spectrum. While most studies have focused on the downstream outputs,

little is known about the upstream inputs that regulate these transcription factors in a live

organism. Here, we found that knockdown of various glucose metabolic enzymes decreases

MML-1 localization in the nucleus and identified two hexokinase isozymes, *hxk-1* and *hxk-2,*

as the most vigorous regulators of MML-1 function. Upon hexokinase knockdown, MML-1

redistributes to mitochondria and lipid droplets (LD), and concomitantly, transcriptional targets

are downregulated and germline longevity is abolished. Further, we found that *hxk-1* regulates

MML-1 through mitochondrial β-oxidation, while *hxk-2* regulates MML-1 by modulating the

pentose phosphate pathway (PPP) and its coordinated association with lipid droplets. Similarly,

inhibition of the PPP rescues mammalian MondoA nuclear translocation and transcriptional

function upon starvation. These studies reveal how metabolic signals and organellar

communication regulate a key convergent metabolic transcription factor to promote longevity.

Key words: MondoA/ChREBP, MML-1, lipid droplet, metabolism, aging, transcription

Introduction

Aging is a complex multifactorial process associated with organismal homeostasis demise and an increased risk of age-related diseases. One key aspect of aging is the dysregulation of gene expression and metabolism (Fischer et al., 2022). Many transcription factors that regulate metabolic pathways have been shown to control lifespan in the evolutionary spectrum (Clancy et al., 2001; Green et al., 2021; Lapierre et al., 2013; Selman et al., 2008; Wang et al., 2008). Changes in the activity of specific transcription factors have been shown to impact gene expression patterns and alter metabolic processes, leading to a decline in cellular function and an increased risk of age-related diseases (Benhamed et al., 2012; Carroll et al., 2015; Denechaud et al., 2008; Du & Zheng, 2021; Tong et al., 2009). Thus, understanding the interplay between metabolism and transcriptional regulation is crucial for gaining insight into the mechanisms underlying aging and developing novel strategies for promoting healthy aging and reducing the risk of age-related diseases.

MondoA and the Carbohydrate Response Element Binding Protein (ChREBP) are paralogous transcription factors that work in heterodimeric complexes with Max-like protein X (Mlx) and have been identified as key regulators of glucose metabolism and energy homeostasis. These transcription factors recognize and bind hexanucleotide tandem sequences (CANNTG) in the promoter regions called Enhancer box (E-box) sequences (Massari & Murre, 2000) of various genes involved in glucose uptake, utilization, and storage (Billin et al., 2000; Cha-Molstad et al., 2009; Sans et al., 2006; Stoltzman et al., 2008; Yamashita et al., 2001). In addition to their role in glucose metabolism, MondoA/ChREBP have also been shown to regulate lipid metabolism and cellular stress responses (Iizuka et al., 2006; Ishii et al., 2004; Ma et al., 2006). Studies have suggested that changes in MondoA/ChREBP activity with age may contribute to age-related metabolic dysfunction, leading to a decline in glucose tolerance and increased risk of type 2 diabetes and obesity (Ghasemi et al., 2015; Iizuka et al., 2006; Richards et al., 2017; Yamamoto-Imoto et al., 2022).

In *Caenorhabditis elegans,* the transcriptional complex formed by MML-1 (MondoA/ChREBP homolog) and MXL-2 (Mlx homolog) is part of an integrated helix-loop-helix (HLH) network required for lifespan extension in diverse longevity pathways (Johnson et al., 2014; Nakamura et al., 2016). However, how these transcription factors integrate multiple upstream signals to regulate lifespan is still unknown. This work describes how metabolic inputs regulate MML-1 subcellular localization and function. We found that HXK-1 and HXK-2 regulate MML-1 cellular distribution and are required for germline longevity by regulating mitochondrial β-oxidation and the pentose phosphate pathway (PPP), respectively.

68 Moreover, inhibition of MML-1 nuclear translocation results in the relocalization of this
69 transcription factor to mitochondria associated with lipid droplets (LD). This study provides
70 key insights into how glucose and lipid metabolism directly impact transcriptional regulation
71 and aging.

72
Results

Hexokinases are required for MML-1 subcellular localization and longevity

MondoA reportedly responds to glucose 6-phosphate (G6P) and other hexose
phosphate sugars in cultured cells by translocating to the nucleus and regulating the expression
of its downstream targets (Stoltzman et al., 2008, 2011). Nevertheless, little is known about
MondoA regulation in an organism-wide setting, and most subsequent work has focused on
the molecular mechanisms downstream of this transcription factor. Hence, we decided to
investigate how MML-1 is regulated by upstream glucose metabolism in *C. elegans*. To
address this, we performed a targeted RNA interference (RNAi)-based screen on genes
encoding enzymes involved in glucose metabolism. Overall, we found a preponderance of
decreased intestinal MML-1 nuclear localization when knocking down such genes (**Figure 1A-**
B, Supplementary Table S1). Knockdown of most of the enzymes involved in glycolysis, as
well as the pyruvate dehydrogenase complex, which links glycolysis to the tricarboxylic acid
(TCA) cycle via acetyl-CoA, decreased MML-1 nuclear accumulation (**Supplementary Table**
S1). Interestingly, whereas most enzymes in the TCA cycle also decreased nuclear MML-1,
inhibition of the oxoglutarate dehydrogenase complex (OGDC) subunits *ogdh-1* and *dld-1* and
the succinyl-CoA synthase (SCS) *suca-1* and *sucg-1* subunits increased MML-1 nuclear
localization. We also observed that knockdown of enzymes involved in the oxidative and
nonoxidative branches of the PPP, phosphogluconate dehydrogenase (PGDH) *T25B9.9* and
transketolase (TKT) *tkt-1* genes, respectively, triggered a strong increase in MML-1 nuclear
accumulation (**Figure 1A-B**). These data suggest that overall changes in glucose metabolism
regulate MML-1 and that this transcription factor may sense the divergence of G6P to different
branches of this network.

Our screen revealed that hexokinase knockdown had the most potent effect in
decreasing MML-1 nuclear localization. This enzyme is involved in the first step of glucose
metabolism, phosphorylating glucose to G6P, and is a critical mediator of cellular metabolism.
Among the three *C. elegans* hexokinase genes, *hxk-1* and *hxk-2* more strongly affected MML-1
nuclear localization in two independent MML-1::GFP reporter strains (**Figure 1B,**
Supplementary Figure 1A), while *hxk-3* had just a small effect on MML-1 nuclear

localization, probably due to its low expression in adult worms (Hutter & Suh, 2016). To test

whether the phosphotransferase activity of hexokinase is responsible for regulating MML-1

nuclear localization, we used two drugs to inhibit hexokinases pharmacologically.

3-Bromopyruvate (3-BrP) is an alkylating agent and HKII inhibitor, and 2-deoxy-glucose (2-

DG) is an analog of glucose phosphorylated to 2-DG6P, which cannot be further metabolized,

thus inhibiting hexokinase function. Supplementation of both drugs significantly decreased

MML-1 nuclear localization compared to control (**Supplementary Figure 1B-C**), indicating

that hexokinase activity drives MML-1 nuclear translocation. We also tested the specificity of

RNAi knockdown by using strains where the three endogenous hexokinase loci were tagged

with mKate2. We used these strains to evaluate the protein levels by western blot and found

that each RNAi was specific to each isozyme (**Supplementary Figure 1D-E**). As glycolysis

is essential during development, we measured the size (area and length) of worms fed with

hexokinase RNAi egg-on. *hxk-1i* and *hxk-2i* displayed no significant change in worm growth,

while *hxk-3i* treated animals were somewhat smaller (**Supplementary Figure 1F-G**). To

determine if our observations were due to a change in food intake, we counted pharyngeal

pumping rate but found no significant difference compared to the control (**Supplementary**

Figure 1H), nor did we see an effect on worm motility compared to control (**Supplementary**

Figure 1I). Moreover, hexokinase knockdown did not affect the steady-state levels of *mml-1*

mRNA (**Supplementary Figure 1J**). Taken together, these data indicate no RNAi cross-

reactivity among the hexokinase isozymes and that the effects on MML-1 nuclear localization

are not due to a decrease in food consumption or developmental phenotype.

MML-1 works in a transcriptional complex with MXL-2, and previous studies have

demonstrated that MML-1 and MXL-2 are required for lifespan extension in multiple longevity

pathways, including germline-less *glp-1(e2141)* mutant (Nakamura et al., 2016). Hence, we

tested whether hexokinases influenced MML-1 subcellular localization in this longevity model.

As expected, MML-1 nuclear localization was increased in *glp-1(e2141)* as reported previously

(Nakamura et al., 2016), and upon *hxk-1* and *hxk-2* knockdown, MML-1 nuclear localization

was abolished in *glp-1(e2141)* mutants (**Figure 1C-D**). Changes in MML-1 nuclear

localization would be expected to modulate its transcriptional activity. To test this idea, we

next analyzed the expression of downstream targets reportedly linked to gonadal longevity

(Nakamura et al., 2016). We observed decreased expression of *mdl-1*, *lgg-2*, *swt-1*, and *fat-5*

under *hxk-1i* and *hxk-2i,* but not under *hxk-3i* (**Figure 1E, Supplementary Figure 1K**),

consistent with decreased MML-1 transcriptional activity. Since hexokinase knockdown

decreases MML-1 nuclear localization in the *glp-1(e2141)* longevity model, we reasoned that

the lifespan might also be abolished. Accordingly, *hxk-1i* and *hxk-2i* significantly decreased
glp-1(e2141) but did not affect wildtype lifespan (**Figure 1F**). These data indicate that
hexokinases are upstream regulators of MML-1 nuclear localization and transcriptional
function in germline-mediated longevity.

Next, we tested whether the effect of hexokinase knockdown on MML-1 nuclear
localization and longevity was shared by other longevity pathways. On the one hand, we
observed that in *isp-1(qm150)* mutants, a model for mitochondrial longevity, *hxk-1* and *hxk-2*
knockdown decreased MML-1 nuclear localization and abolished lifespan extension (**Figure
1G-H**). On the other hand, while *mml-1* was required for the insulin signaling *daf-2(e1370)*
mutant longevity, *hxk-1* and *hxk-2* knockdown did not alter MML-1 nuclear localization nor
suppress the lifespan in this background (**Supplementary Figure 1L-M**). Further, we found
that MML-1 nuclear localization was increased in *raga-1(ok701)* mutants, a model for
mTORC1 longevity, and *mml-1* was required for lifespan extension (**Supplementary Figure
1N-O**). However, *hxk-1* and *hxk-2* knockdown did not suppress *raga-1(ok701)* longevity
(**Supplementary Figure 1P**). Collectively, these data suggest that while MML-1 acts as a
convergent factor in these four longevity pathways, it only requires HXK-1 and HXK-2 for
germline and mitochondrial longevity, but not for insulin and mTOR longevity.

C. elegans hexokinases are differentially expressed

Mammals harbor four hexokinases that differ in tissue expression, subcellular
localization, and affinity to glucose and G6P. However, knowledge of the worm isozymes is
limited. We used the hexokinase tagged with mKate2 strains to investigate the tissue-specific
expression and subcellular localization of the different isozymes by confocal microscopy. We
found that HXK-1 was expressed in the cytosol of neurons, pharynx, gonadal sheath, and
coelomocytes (**Figure 2A**). HXK-2 had a reticular expression pattern in neurons, muscle,
intestine, and hypodermis (**Figure 2B**). HXK-3 signal was diffused and highly expressed in
the pharynx, muscle, hypodermis, and intestine (**Supplementary Figure 2A**).

Human HKI and HKII have been shown to bind reversibly to the mitochondria (Sui &
Wilson, 1997). Binding depends on an intrinsic hydrophobic N-terminal sequence critical for
inserting into the outer mitochondrial membrane (Pastorino et al., 2002). Due to the reticular
expression pattern, HXK-2 was a plausible candidate to be mitochondrially localized. To
corroborate this idea, we crossed the HXK-2::mKate2 strain with a mitochondrial reporter that
expresses GFP with a mitochondrial targeting sequence under the body wall muscle promoter
myo-3. Indeed, we observed that full-length HXK-2 strongly co-localized with mitochondria

(**Figure 2C**). We also generated an HXK-2 mutant that lacks the first ten amino acids after the

initiator methionine (d10 HXK-2) and found that mitochondrial localization was abolished

entirely (**Figure 2C**). In summary, we found that hexokinases are differentially expressed in

C. elegans and identified HXK-2 as the worm mitochondrial hexokinase.

Hexokinase knockdown increases MML-1 mitochondrial localization

176   Since we observed that *mml-1* mRNA levels were unaffected by *hxk* knockdown and

saw a decrease in MML-1 nuclear localization upon hexokinase knockdown, we wondered

whether MML-1 might re-localize to another subcellular compartment. First, we crossed the

MML-1::GFP reporter strain with two compartmental markers and examined colocalization by

confocal microscopy. We looked at the lysosome/endosome reporter strain LMP-1::mKate2

but saw no evident colocalization in intestinal cells upon *hxk-2* knockdown or control

conditions (**Supplementary Figure 2B-C**). We had previously shown that MML-1 localizes

with mitochondria in the hypodermis (Nakamura et al., 2016); however, mitochondrial

subcellular localization in other tissues had not been examined. Hence, we crossed the

MML-1::GFP strain with the mitochondrial reporter TOMM-20::mCherry under the intestinal

promoter *ges-1* and confirmed that MML-1 localized with mitochondria in the intestine

(**Figure 2D-E**). Under normal conditions, around 60-80% of the total MML-1 was nuclear, and

15% co-localized with mitochondria. Interestingly, MML-1 mitochondrial localization was

increased to ~62% of total MML-1 upon *hxk-2i* (**Figure 2F**). Taken together, our data indicate

that upon *hxk-2* knockdown, MML-1 is excluded from the nucleus and re-localizes to a subset

population of mitochondria.

Inhibition of mitochondrial β-oxidation rescues MML-1 nuclear localization under hxk-1i

194   Many genes that regulate MML-1 localization are involved in mitochondrial

metabolism, such as OGDC and SCS, and inhibiting these complexes could activate stress

response pathways. Hence, we assessed whether MML-1 nuclear localization was associated

with the activation of mitochondrial stress response and oxidative stress. As expected, we

observed increased expression of *hsp-6*::*GFP*, a reporter for mitochondrial stress, upon

knockdown of mitochondrial components *ogdh-1*, *sucl-1*, and *sucl-2* as well as in control

worms treated with antimycin A (AA), a complex III inhibitor (**Supplementary Figure 2D**).

However, inducing mitochondrial stress with AA did not affect MML-1 nuclear localization

under control conditions and *hxk-2* knockdown (**Supplementary Figure 2E**). Next, we sought

to test whether an increase in oxidative stress affects MML-1 nuclear localization. We exposed worms to the ROS-generating compound paraquat and observed induction of the oxidative reporter *gst-4p*::GFP (**Supplementary Figure 2F**); however, we did not see any effect on MML-1 nuclear localization at the concentrations tested (**Supplementary Figure 2G**). Taken together, these data indicate that MML-1 regulation by glucose metabolism genes is independent of mitochondrial and oxidative stress.

Hexokinase downregulation might be expected to cause a reduction in the reliance on glycolysis and increase fatty acid metabolism as fuel for cellular maintenance (**Figure 2G**). Mitochondrial β-oxidation is an important pathway that generates acetyl-CoA for the TCA cycle through the catabolism of fatty acids to produce reductants and energy. Hence, we sought to investigate whether decreasing fatty acid oxidation could affect MML-1 nuclear localization upon hexokinase knockdown. First, we measured the neutral lipid content by feeding the worms with the labeled fatty acid C1-BODIPY-C12 and found increased total lipid storage under hexokinase knockdown in the wildtype and *glp-1(e2141)* backgrounds (**Supplementary Figure 2H**). Next, we measured the oxygen consumption rate (OCR) upon hexokinase knockdown under basal conditions and with etomoxir supplementation, a mitochondrial β-oxidation inhibitor that acts on the carnitine palmitoyltransferase (CPT). Inhibition of the CPT decreases acyl-CoA import to mitochondria, and changes in OCR are a correlative measure of mitochondrial β-oxidation of fatty acids (Ramachandran et al., 2019). We found no difference in basal OCR under *hxk-1*, *hxk-2*, and *mml-1* knockdown in the wildtype background (**Figure 2H**). *glp-1(e2141)* animals showed decreased basal OCR compared to wildtype, whereas *hxk-1* and *mml-1* knockdown significantly increased the basal OCR compared to *glp-1(e2141)* (**Figure 2H**). Interestingly, we saw a greater percentage of OCR reduction upon adding etomoxir under *hxk-1* knockdown in wildtype and *hxk-1*, *hxk-2*, and *mml-1* knockdown in *glp-1(e2141)* (**Figure 2I**), indicating higher mitochondrial β-oxidation levels. We confirmed this result using the transcriptional reporter *acs-2p*::GFP, a mitochondrial acyl-CoA synthase that is induced and required for fatty acid β-oxidation in *C. elegans* (Van Gilst et al., 2005), and found increased expression upon *hxk-1* and *hxk-2* knockdown (**Supplementary Figure 2I**).

We next wondered whether this increase in fatty acid oxidation was responsible for decreased MML-1 nuclear localization under hexokinase knockdown. First, we performed a double knockdown of *hxk-1* or *hxk-2* combined with *acs-2*. MML-1 nuclear localization was rescued upon *acs-2i* under *hxk-1i* (**Figure 2J**). Interestingly, *acs-2i* did not rescue MML-1

nuclear localization under *hxk-2i*. We also pharmacologically inhibited fatty acid oxidation by
supplementing etomoxir and again saw the rescue of MML-1 nuclear localization under *hxk-*
1i, but no effect upon *hxk-2i* (**Figure 2K**). These data suggest that HXK-1 normally promotes
MML-1 nuclear localization by decreasing mitochondrial β-oxidation and reveal that the two
hexokinases might regulate MML-1 nuclear localization through somewhat independent
pathways.

hxk-2 regulates MML-1 function through increased PPP

G6P can be diverted as a substrate for metabolic pathways other than glycolysis. Thus,
we wanted to test whether MML-1 could sense other branches of glucose metabolism. From
our RNAi screen, we observed increased MML-1 nuclear localization upon knockdown of two
enzymes from the PPP, *T25B9.9* and *tkt-1* (**Figure 1A-B**). The PPP is a cytosolic metabolic
pathway involved in the interconversion of 3 to 7-carbon sugars that generate precursors for
the biosynthesis of lipids, amino acids, and nucleotides (**Figure 3A**). It also maintains redox
homeostasis through NADPH production. To further address these findings, we tested the
effect of pharmacological inhibition of the PPP on MML-1 nuclear localization using
6-aminonicotinamide (6-AN), which is a competitive inhibitor of the G6P dehydrogenase
(G6DH) and phosphogluconate dehydrogenase (6PGD). These two enzymes generate NADPH
in the PPP. In agreement with our previous results, we observed increased MML-1 nuclear
localization after supplementation with 1 mM of 6-AN (**Figure 3B**). Moreover, we found that
hxk-2i, but not *hxk-1i*, significantly increased the NADPH levels (**Figure 3C**). Additionally,
we found that *hxk-2i* had increased TKT-1 enzymatic activity compared to the control (**Figure
3D**). These data suggest that inhibition of the mitochondrial hexokinase enhances the PPP.
Next, we sought to test the epistatic interaction of hexokinase and PPP knockdown on
MML-1 nuclear localization. For these experiments, we performed double RNAi of *hxk-1* and
hxk-2 in combination with the 6PGD *T25B9.9* and transketolase *tkt-1*. Interestingly, knocking
down these enzymes rescued MML-1 nuclear localization under *hxk-2i* (**Figure 3E-F**) but did
not rescue the localization under *hxk-1i* (**Supplementary Figure 3A**). To evaluate MML-1
function, we measured its transcriptional readouts and found that knocking down the PPP under
hxk-2i rescued the expression of *lgg-2*, *mdl-1*, *swt-1*, and *fat-5* (**Figure 3G, Supplementary
Figure 3B-D**); however, it had no effect under *hxk-1i* (**Supplementary Figure 3E-H**).
Moreover, we observed lower levels of G6P upon double knockdown of hexokinases with
T25B9.9 and *tkt-1* compared to control (**Figure 3H, Supplementary Figure 3I**). Taken

together, these data suggest that the PPP activity is increased upon knockdown of the

mitochondrial hexokinase HXK-2, and inhibiting the PPP is sufficient to rescue MML-1

function independent of G6P levels.

Finally, we asked whether reducing the PPP could impact *glp-1(e2141)* lifespan upon

hxk-2 knockdown. Inhibition of the PPP has been reported to extend the lifespan in *C. elegans*

by activating the mitochondrial UPR^{mt} (Bennett et al., 2017). As reported previously, we found

a significant lifespan extension upon *tkt-1* knockdown in wildtype (**Figure 3I**). Moreover,

knockdown of the PPP in the *glp-1(e2141)* background had no additive effect on longevity,

suggesting that *glp-1* and *tkt-1i* may share common mechanisms in extending lifespan.

Consistent with our data, knockdown of *T25B9.9* and *tkt-1* rescued *glp-1(e2141)* longevity

upon *hxk-2i* (**Figure 3I, Supplementary Figure 3J**). Collectively, these data indicate that an

increase in the PPP under *hxk-2* knockdown suppresses MML-1 function, and reducing PPP

restores longer life, placing PPP epistatically downstream of *hxk-2*.

MML-1 interactome uncovers potential regulators of its function and localization

Given the mitochondrial localization of MML-1 and possibly other organelles, we

decided to identify potential MML-1 binding partners to better understand its subcellular

distribution. We used CRISPR/Cas9 to tag the endogenous MML-1 locus with a 3xFLAG tag.

MML-1::3xFLAG was immunoprecipitated from whole-worm lysates, and potential

interactors were identified by mass spectrometry. We detected ca.1600 to 2000 different

proteins from three biological replicates and observed that the biological replicates clustered

together (**Supplementary Figure 4A-B**). We detected several proteins in the

immunoprecipitation (IP), including MXL-2, an established MML-1 binding partner (Pickett

et al., 2007) (**Figure 4A**). Analysis of the top 50 co-enriched candidates in MML-1 IPs revealed

enrichment for nuclear and mitochondrial proteins (**Supplementary Figure 4C**). We also

found multiple enriched proteins from different organelles, including lipid droplets (LD).

Mitochondria and LD dynamically interact in highly active metabolic tissues, and it has

been shown that mitochondria associated with LD have specific metabolic behavior compared

to cytosolic mitochondria (*e.g.*, different dynamics, motility, and capacity to burn

carbohydrates and lipids) (Gordaliza-Alaguero et al., 2019; Rambold et al., 2015). LD are

organelles involved in multiple roles serving as nutrient reservoirs and participating in

signaling pathways, and their biogenesis or degradation is tightly coupled to carbohydrate

metabolism (**Figure 4B**). Interestingly, PLIN-1, one of the most abundant LD proteins, was

co-enriched in MML-1 IP (**Figure 4A**). Hence, we wanted to test whether MML-1 localized to

these organelles. First, we used LipiBlue, a dye to specifically visualize LD *in vivo* (Tatenaka
et al., 2019), and found that all LipiBlue-stained structures were positive with the LD resident
proteins DHS-3::GFP and PLIN-1::mCherry (**Supplementary Figure 4D-F**), indicating that
this dye could be used as a reliable method for visualization of LD in *C. elegans*. Next, we
examined LipiBlue-stained animals co-expressing MML-1::GFP and TOMM-20::mCherry
and saw that MML-1 localized with LipiBlue-positive structures (**Figure 4C**). Interestingly,
MML-1 appeared localized to mitochondria associated with LD (**Figure 4C-D**). Moreover, this
colocalization was significantly increased upon *hxk-2* knockdown (**Figure 4E**).

Inhibition of transaldolase and enzymes from the PPP have been shown to activate a
fasting-like response by increasing lipolysis and enhancing the breakdown of LD (Bennett et
al., 2017), including higher expression of the adipose triglyceride lipase ATGL-1
(**Supplementary Figure 4G**). Therefore, we wondered whether the LD size correlates with
MML-1 nuclear localization. We knocked down *hxk-2* and *tkt-1* to decrease and increase
MML-1 nuclear localization, respectively. We observed a significant reduction in diameter and
the total amount of LDs upon knockdown of *tkt-1*, while *hxk-2i* did not have an effect (**Figure
4F, Supplementary Figure 4H).** Double knockdown of *hxk-2* and *tkt-1*, however, resulted in
smaller and fewer LDs than the control. We also wanted to test whether the HXK-1/fatty acid
β-oxidation axis regulation of MML-1 also converged on LD size regulation. Interestingly, we
found that knockdown of *acs-2i* and *hxk-1i acs-2i* double knockdown resulted in a mild but
significant increase in LD size (**Supplementary Figure 4I**), supporting the notion that the two
hexokinases regulate MML-1 via distinct mechanisms. Collectively, these data suggest that
MML-1 can localize with a specific subpopulation of mitochondria associated with LD and
that inhibition of the PPP can rescue MML-1 nuclear localization upon *hxk-2i* correlated with
decreasing LD size.

To test whether LDs directly regulated MML-1 localization, we knocked down *dgat-2*
and *atgl-1*, two enzymes involved in LD metabolism. DGAT-2 is a diacylglycerol (DAG)
transferase that promotes LD droplet expansion, while ATGL-1 is a rate-limiting enzyme in
LD breakdown (**Figure 4B**). First, we measured the neutral lipid composition and found that
atgl-1i increased lipid levels while *dgat-2i* worms did not show significant changes under *ad*
libitum conditions (**Supplementary Figure 4J**). Next, we tested MML-1 nuclear localization
and observed that *dgat-2i* increased MML-1 nuclear localization while *atgl-1i* resulted in
decreased levels of nuclear MML-1 (**Figure 4G**). Finally, we did an epistatic analysis of LD
metabolism and the PPP and found that MML-1 nuclear localization in *atgl-1i* was rescued

upon *tkt-1* knockdown (**Figure 4H**), thus suggesting that LD metabolism is upstream or parallel to the PPP in MML-1 regulation.

The PPP regulates MondoA function in mammalian cells

We next asked whether the PPP regulation of MML-1 we observed in *C. elegans* was conserved in mammals. Specifically, we wanted to determine whether inhibition of the PPP is sufficient to induce MondoA translocation to the nucleus under contrasting conditions where MondoA is mostly cytosolic (starvation media) or split between the nucleus and cytoplasm (high-glucose growing media). To test this, we cultured the HEK293T cells in high-glucose media with 6-AN and subsequently transferred the cells to either high-glucose media or starvation media (**Supplementary Figure 5A**). First, we observed decreased NADPH/NADP+ ratio in cells treated with 6-AN and under starvation conditions (**Supplementary Figure 5B**). Next, we found that in both conditions, MondoA nuclear localization was increased upon inhibition of the PPP (**Figure 5A-B**), while there was no evident regulation of MondoA at the transcript level (**Figure 5C**). Consistent with MondoA increased nuclear localization, we observed increased expression of two of the best-described downstream targets of MondoA, *TXNIP* and *ARRDC4* (**Figure 5D, Supplementary Figure 5C**). These findings indicate that the PPP regulates mammalian MondoA localization and transcriptional function.

Discussion

Nutrient sensing pathways and metabolism play a central role in modulating aging by establishing resilience and survival metabolic states. Although much work has focused on identifying mediators of these processes, we still know little about how organisms coordinate feedback loops of metabolic pathways and transcriptional responses to establish such states. Here, we present novel carbohydrate and lipid metabolism mechanisms regulating the transcription factor MML-1/Mondo. We observed that MML-1 senses multiple steps of glucose metabolism and found that hexokinases regulate MML-1 function through different mechanisms in *C. elegans*. On the one hand, we found that under *hxk-2* knockdown, MML-1 re-localizes to mitochondria and LD, abolishing the longevity of germline-less worms and mutants with reduced mitochondrial function. Moreover, we could rescue MML-1 function by decreasing the PPP independent of the G6P levels. On the other hand, we found that *hxk-1* knockdown decreased MML-1 function by upregulating mitochondrial β-oxidation. Thus, we found two metabolic branches that converge on MML-1 to trigger a transcriptional response

and regulate longevity. We note that while MML-1 is required in four different longevity
models, namely *glp-1,isp-1, daf-2,* and *raga-1, hxk-1* and *hxk-2* were required only for *glp-1*
and *isp-1* mediated longevity. Presumably, the signal transduction wiring is different in *raga-1*
and *daf-2*, or their inputs into MML-1 work downstream or in parallel to the hexokinases.

MondoA and ChREBP transcription factors have well-established central roles in
glucose metabolism (Ma et al., 2006; Sans et al., 2006; Stoltzman et al., 2008; Yamashita et
al., 2001), but whether and how glucose metabolic circuits affect Mondo regulation in a whole
animal model is little explored. By performing a systematic RNAi-based targeted screen of
enzymes involved in glucose anabolism and catabolism, we found that MML-1 senses multiple
steps of glucose metabolism in the worm. Among the genes tested, we found MML-1 nuclear
localization to be broadly sensitive to the downregulation of glycolytic enzymes, and the
knockdown of hexokinase had the most profound effect. Hexokinase carries out the first step
in glucose metabolism, the phosphorylation of glucose to form G6P, which can be shunted to
multiple metabolic pathways, including glycolysis, the PPP, glycogenesis, and trehalose
production. Our findings are consistent with previous work in cultured cells showing that
MondoA localization is regulated by HKII, which is thought to stimulate nuclear localization
via the production of G6P (Sans et al., 2006). Additionally, MondoA senses other
phosphorylated hexoses, including allose and 3-*O*-methylglucose (Stoltzman et al., 2011).

Consistent with our genetic findings above, pharmacological inhibition of glycolysis
by supplementation with 3-BrP and 2-DG suppressed MML-1 nuclear localization. In contrast
to MML-1, MondoA has been shown to accumulate in the nucleus upon 2-DG in rat L6 cells
(Stoltzman et al., 2008). A possible explanation for this difference is that our treatment
represents a chronic exposure to 2-DG from development until adulthood, while the
experiments in cell culture may reflect an acute response. Overall, 2-DG has been used as a
glycolytic inhibitor because it can be phosphorylated by hexokinase to 2-DG 6-phosphate;
however, it cannot be further metabolized in glycolysis. Further evidence shows that 2-DG not
only functions as a catabolic inhibitor but can also be directed to the PPP or other pathways
(Chi et al., 1987; Ralser et al., 2008). Moreover, low concentrations of 2-DG have been shown
to extend lifespan by restricting glucose metabolism, while higher concentrations of 2-DG and
glucose shortened the lifespan in worms (Schulz et al., 2007, Lee et al., 2009), suggesting a
threshold for the benefits of glucose metabolism inhibition in lifespan.

Additionally, we found an important link between glycolysis and pyruvate metabolism
regulating MML-1 localization. Under aerobic conditions, pyruvate is imported into the

mitochondria and converted to acetyl-CoA to generate ATP and other reducing molecules.

Downregulation of the mitochondrial pyruvate carrier *mpc-1* and the pyruvate dehydrogenase

complex (**Supplementary Table S1**) also decreased MML-1 nuclear localization and function.

Further, we found MPC-1 enriched in the MML-1 interactome. Interestingly, loss of *MPC1* in

yeast has been shown to accumulate pyruvate, lower TCA cycle intermediates, and reduce

chronological aging (Bricker et al., 2012; Orlandi et al., 2014). *MPC1* similarly regulates the

disposition of glycolytic and TCA intermediates in flies and mammals (Bricker et al., 2012).

Pyruvate can also enter the TCA cycle through pyruvate carboxylase PYC-1, which converts

pyruvate to oxaloacetate. However, knockdown of *pyc-1* did not affect MML-1 localization

(**Supplementary Table S1**), suggesting that MML-1 may sense pyruvate metabolism mainly

through oxidative decarboxylation of acetyl-CoA.

We also found that MML-1 responds strongly to alterations in the TCA cycle. The TCA

cycle allows organisms to oxidize carbohydrates, fatty acids, and amino acids to provide energy

and intermediates for the biosynthesis of macromolecules. Interestingly, the knockdown of two

subunits of the OGDC, *ogdh-1* and *dld-1*, increased MML-1 nuclear localization. The OGDC

is a rate-limiting enzyme within the mitochondrial TCA cycle that decarboxylates

2-oxoglutarate to succinyl-CoA. It has been shown that supplementation of 2-oxoglutarate and

knockdown of *ogdh-1* extends lifespan through inhibition of ATP synthase, decreasing oxygen

consumption, and increasing autophagy in an mTOR-dependent manner (Chin et al., 2014).

Whether MML-1 is required for this lifespan extension has yet to be explored. Conceivably,

the mitochondrial localization of MML-1 could help sense 2-oxoglutarate levels, and the

accumulation of this metabolite could signal translocation to the nucleus. Moreover, glutamine

can be converted to glutamate and subsequently to 2-oxoglutarate during glutaminolysis and

represents an important mechanism to replenish the TCA cycle under different metabolic

challenges. In this case, OGDC plays a pivotal role in channeling anaplerotic reactions into the

TCA cycle from glutamine and glutamate and other amino acids that can ultimately be

converted to 2-oxoglutarate (*i.e.*, histidine, proline, and arginine) (Owen et al., 2002).

Interestingly, increased glutamine-dependent anaplerosis has been shown to regulate MondoA

activity in BxPC-3 cells (Kaadige et al., 2009), though the molecular mechanism remains

unclear. Whether glutamine, glutamate supplementation, or knockdown of glutamate

dehydrogenase can regulate MML-1 nuclear localization remains to be determined.

On the other hand, the TCA cycle can operate in reverse to restore TCA intermediate

levels (Dalziel & Londesborough, 1968). Increased levels of 2-oxoglutarate due to inhibition

of OGDC, or glutamine anaplerosis, can undergo reductive carboxylation by isocitrate

dehydrogenase to produce isocitrate. After that, aconitase can convert isocitrate to citrate,
which can be exported from the mitochondria for *de novo* lipogenesis. MML-1 could help
orchestrate the metabolic rewiring of the TCA cycle with the fatty acid metabolism
transcriptional response. Notably, ChREBP has been shown to regulate many lipogenic
enzymes, including acetyl-CoA carboxylase (ACC), the fatty acid synthase (FAS), and
ATP-citrate lyase (ACLY) (Ishii et al., 2004; Postic et al., 2007). Accordingly, the MML-1
transcriptome shows clear regulation of several fatty acid metabolic enzymes (Nakamura et al.,
2016).

We observed that PPP enzymes strongly influenced MML-1 nuclear localization and
function. The PPP is a cytosolic pathway involved in the interconversion of sugars to generate
precursors for the biosynthesis of lipids, amino acids, and nucleotides and maintain redox
homeostasis. We found that knockdown of PPP components, PGDH *T25B9.9* and transketolase
tkt-1, enhanced MML-1 nuclear localization, increased the transcription of MML-1 target
genes, and restored longevity to *glp-1(e2141)* animals under *hxk-2i*. The PPP is the primary
source of NADPH, which plays a vital role in many cellular processes, including fatty acid,
nucleotide, neurotransmitter, and cholesterol metabolism, and is an essential reducing agent
(Ju et al., 2020). We found that pharmacological inhibition with 6-AN of the two enzymes
producing NADPH in the PPP increased MML-1 nuclear localization. We also found increased
levels of NADPH and lower levels of G6P under *hxk-2* knockdown. Importantly, these
observations suggest that residual G6P may be diverted to the PPP under low mitochondrial
hexokinase and further imply that metabolites other than G6P could regulate MML-1 nuclear
localization. What this other metabolite could be remains elusive, but a simplifying notion is
that MML-1 nuclear localization is promoted by glycolysis or PPP intermediates. Blocking
entry into the PPP or TCA might be expected to accumulate such metabolites and promote
Mondo nuclear localization. Conversely, depletion of these metabolites would disrupt Mondo
nuclear localization and activity.

Another possibility is that Mondo organellar distribution is regulated at another scale.
Notably, a major output of the PPP and NADPH metabolism is the production of lipids.
Interestingly, we found that under *tkt-1* knockdown, fat levels decreased, and LDs were smaller
in diameter, suggesting that a reduction in LDs could signal MML-1 to translocate to the
nucleus. Consistently, transaldolase *tald-1* knockdown has been shown to increase lifespan and
decrease intestinal fat levels in the worm through increased activity of adipose triglyceride
lipase ATGL-1 (Bennett et al., 2017). This enzyme is involved in the metabolism of LD under
starvation to mobilize stored fats for energy production (Lee et al., 2014).

Interestingly, although both HXK-1 and HXK-2 positively regulate MML-1 function,
they do so through different mechanisms **(Figure 5E)**. We observed that decreasing the PPP
rescued MML-1 nuclear localization and *glp-1* longevity under *hxk-2* knockdown, independent
of the G6P levels, but not under *hxk-1* knockdown. Conversely, inhibition of mitochondrial
fatty acid β-oxidation is sufficient to rescue MML-1 nuclear localization under *hxk-1*
knockdown, but not under *hxk-2* knockdown. Interestingly, *acs-2* knockdown abolished *glp-1*
longevity (data not shown)*,* consistent with previous work showing that NHR-49, a
transcription factor that drives *acs-2* expression, is required for *glp-1* longevity (Ratnappan et
al., 2014). Thus, inhibiting fatty acid β-oxidation promotes MML-1 nuclear localization under
hxk-1i but abolishes lifespan extension, potentially due to epistatic effects on other
transcription factors or processes.
How both hexokinases activate different metabolic pathways to regulate MML-1
remains unknown and could work at multiple levels. First, there is evidence that differences in
the subcellular location of hexokinases may result in the compartmentalization of glucose
metabolism, with the channeling of G6P to different pathways (John et al., 2011; Wilson,
2003). Mitochondrial hexokinase has preferential access to ATP generated in the mitochondria
and provides efficient glycolysis coupling with further pyruvate oxidation by the TCA cycle
and oxidative phosphorylation (OXPHOS). Furthermore, mammalian HKI and HKII have a
dynamic localization, shuttling between the mitochondrial outer membrane and the cytoplasm
(John et al., 2011). Both mammalian hexokinases have been reported to reversibly bind to
mitochondria through their N-terminal sequence (Roberts et al., 2014) and via their interaction
with VDAC1 (Lindén et al., 1982). This dual localization allows the cell to adapt to different
metabolic requirements and maintain energetic balance. Under high levels of G6P, HKII
shuttles from the mitochondria to the cytoplasm, causing cells to use glycogen as an energy
source (John et al., 2011). The subcellular localization also plays other significant roles in
different signaling pathways. For example, mitochondria-bound hexokinase has anti-apoptotic
effects (Gottlob et al., 2001), and increased N-acetylglucosamine upon pathogen exposure
releases hexokinase from the mitochondria and activates innate immune responses (Wolf et al.,
2016). Here we found that *C. elegans* HXK-2 is associated with mitochondria and that this
localization is abolished by deleting the first ten amino acids. However, we saw no effect upon
vdac-1 knockdown nor dynamic localization of HXK-2 between mitochondria and cytosol
(data not shown). Taken together, these observations suggest that the nematode has dedicated
mitochondrial (HXK-2) and cytosolic (HXK-1 and HXK-3) hexokinases.

We also found differences in the tissue expression of the different hexokinase isozymes
of the nematode: HXK-1 was mainly found in the pharynx, neurons, gonadal sheath, and
coelomocytes, while HXK-2 was present in the hypodermis, body wall muscle, and intestine.
Consistently, RNAseq data from different tissues in *C. elegans* shows that *hxk-2* is the main
isozyme in the body wall muscle and intestine (Hutter & Suh, 2016). At the same time, *hxk-1*
is the main one expressed in pharyngeal muscle and neurons. This differential distribution
could explain how *hxk-1* and *hxk-2* differentially regulate MML-1 through β-oxidation and
PPP pathways. Notably, we scored MML-1 nuclear localization in the intestine in our
experiments, where *hxk-2* but not *hxk-1* is expressed. In the future, it will be important to see
whether *hxk-1* cell autonomously impacts *mml-1* nuclear localization within the same tissue,
such as the pharynx.

In any case, our observation that HXK-1 affects intestinal MML-1 nuclear localization
suggests cell non-autonomous signaling. What might be the nature of such cell non-
autonomous signaling? One possibility is the sterol hormone signaling pathway, which acts in
somatic reproductive tissue to regulate *glp-1(e2141)* longevity. Decreasing the activity of the
nuclear hormone receptor DAF-12, or its ligands dafachronic acids, abolishes the longevity of
germline-less worms (Gerisch et al., 2001; Yamawaki et al., 2010). Interestingly, carbohydrate
metabolism directly impacts the production of dafachronic acids as NADPH is required for the
last step of dafachronic acid biosynthesis by DAF-9 (Motola et al., 2006; Penkov et al., 2015).
Furthermore, DAF-9 is expressed in the somatic gonad (spermatheca) and XXX neurons in
adult animals (Gerisch & Antebi, 2004). Conceivably, hexokinase downregulation could affect
the production of NADPH and dafachronic acids regulating DAF-12 activity, thereby affecting
MML-1 function. It will be interesting to see whether dafachronic acid supplementation can
rescue MML-1 nuclear localization upon *hxk-1i* or *hxk-2i*.

The nervous system also plays a crucial role in integrating metabolic status and
systemic responses in a cell non-autonomous manner (Mutlu et al., 2020; Zhang et al., 2019).
Inhibition of the mTOR signaling pathway has been shown to extend lifespan across the
evolutionary spectrum (Kennedy & Lamming, 2016), and recent work has shown that restoring
*raga-1/*RagA or *rsks-1/*S6K in the nervous system is enough to entirely suppress the lifespan
extension conferred by mutation of these mTOR signaling factors (Zhang et al., 2019). In cells,
HKII has been shown to shuttle from the mitochondria and directly interact with mTOR under
glucose deprivation and inhibit its function resulting in an induction of autophagy (Roberts et
al., 2014). This interaction is dependent on the mTOR signaling (TOS) motif present in HKII

required for binding to Raptor. Interestingly, this TOS motif (KDIDI) is conserved in the
nematode HXK-1; however, HXK-2 and HXK-3 lack this sequence. Our finding that *raga-1*
mutants are epistatic to hexokinase knockdown for life span extension is consistent with a role
of mTOR signaling downstream of hexokinase. It would be interesting to study whether the
interaction between hexokinase and mTOR is conserved in *C. elegans* and whether decreasing
the main neuronal hexokinase (HXK-1) would result in mTOR-dependent inhibition of MML-
1.

Proteins tend to co-localize near their substrates to maximize specificity and regulate
signaling pathways. In particular, MondoA was first reported to be associated with the
mitochondria outer membrane in primary human SkMC cells through protein-protein
interaction (Sans et al., 2006). We have previously shown that this localization is conserved in
C. elegans (Nakamura et al., 2016). Mitochondria-bound MondoA has been proposed to have
preferential access to newly synthesized G6P from HKII to coordinate metabolism and
transcriptional response (Wilde et al., 2019). However, the identity of the mitochondrial factor
tethering MondoA at the mitochondria remains unknown.

Interestingly, our proteomics analysis of MML-1 immunoprecipitation showed
enrichment with proteins from different cellular compartments, including mitochondria, ER,
LD, and nucleolus, suggesting an important role of Mondo in integrating diverse organellar
signals. Localization to mitochondria was particularly conspicuous. Multiple nuclear
transcription factors are localized in the mitochondria, although dissecting their nuclear and
mitochondrial function is challenging (Leigh-Brown et al., 2010). On the one hand, nuclear
transcription factors have been shown to directly bind the mitochondrial genome to regulate
transcription, such as the cAMP response element-binding (CREB) protein (Marinov et al.,
2014). Interestingly, *mml-1(ok849)* loss-of-function mutants showed a decrease in all 12
mitochondrial-encoded genes compared to wildtype (Nakamura et al., 2016). However, this
could be indirect by regulating the mitochondrial transcription factor TFAM or mitochondrial
biogenesis (Li et al., 2005). On the other hand, transcription factors like p53 have been shown
to translocate to the mitochondrial outer membrane under apoptotic signals to interact with
Bcl-2 and promote membrane permeabilization and apoptosis (Marchenko et al., 2007). Future
studies will help elucidate whether MML-1/MondoA mitochondrial localization regulates
other cellular processes independent of its nuclear transcription function.

LD represents the main lipid storage in cells to balance metabolism and energy
demands. These organelles are essential for many signaling pathways and are required for
survival during fasting by mobilizing lipids for energy production. In *C. elegans,* LDs are

mainly localized in the intestine and hypodermis and are composed almost exclusively of TAG

(Vrablik et al., 2015). We found PLIN-1, one of the most abundant LD proteins, enriched in

the MML-1 interactome. Interestingly, we found that MML-1 colocalizes with LD *in vivo* in

the intestine. MondoA was previously seen in large-scale proteomics associated with LD

(Krahmer et al., 2018). It was shown that MondoA and Mlx localized with LD through

amphipathic helices in the C-terminus in SUM159 cells (Mejhert et al., 2020). The researchers

proposed a model in which MondoA/Mlx colocalization with LD inhibits its transcriptional

activity, limiting glucose-dependent transcription. Work by Brunet and colleagues suggests

that MUFA intercalation into LD and peroxisomes somehow promotes longevity, though

downstream transcription factors are little explored (Papsdorf et al., 2023). Mitochondria and

LD physically interact with one another to regulate metabolism (Boutant et al., 2017). As

MondoA has also been shown to localize with mitochondria, it would be interesting to

understand the role of this organelle in regulating MondoA's association with LD. Although

the proteins involved in these contact sites are just beginning to be uncovered, both

PLIN1/PLIN-1 and ACSL1/ACS-13 have been proposed to tether mitochondria and LD

(Gordaliza-Alaguero et al., 2019).

Mitochondria bound to LD facilitate the coordination of TAG metabolism and fatty

acid oxidation. For example, lipases (*e.g.*, ATGL) in the LDs break down TAG to release free

fatty acids that can be activated in the mitochondria's outer membrane by ACSL1 to fatty

acyl-CoA. This acyl-CoA can be imported into the mitochondria for β-oxidation. We propose

a model **(Figure 5E)** in which MML-1 interacts with specific mitochondria associated with LD

to orchestrate a transcriptional response depending on fuel utilization. Under normal

conditions, glucose stimulates MML-1 translocation to the nucleus to activate the transcription

of glycolytic enzymes. In contrast, upon blockage of glycolysis (*e.g.,* hexokinase knockdown),

the cell utilizes fatty acids to produce ATP, re-localizing and inhibiting MML-1 at the

mitochondria associated with LD.

Materials and Methods

Worm growth and RNAi treatment

C. elegans worm strains were grown and maintained on nematode growth medium

(NGM) plates seeded with *E. coli* OP50 strain as the primary food source. Worms were grown

at 20 °C unless stated otherwise. *glp-1(e2141)* strains were maintained at 15 °C. All worm

strains are listed in **Supplementary Table S4**. Worms were synchronized by 4 to 6 h egg lays

at 20 °C. For experiments including *glp-1(e2141)*, egg lays were performed at 15 °C, and the

eggs were upshifted to 25 °C for 52 to 56 h to induce the germline-less phenotype. After *glp-1*

induction, worms were downshifted to 20 °C until the day of analysis or collection.

The NGM media for RNAi knockdown was supplemented with 1 mM isopropyl

β-1-D-thiogalactopyranoside (IPTG) and 100 µg mL^{-1} ampicillin. The plates were seeded with

the RNAi-expressing *E. coli* HT115 (DE3) strain transformed with the L4440 vector from the

Ahringer and Vidal libraries (Kamath & Ahringer, 2003; Rual et al., 2004), or cloned for this

work. Bacteria containing luciferase RNAi cloned into the L4440 vector were used as controls.

Worms were grown on RNAi plates egg-on in all the experiments.

Cell culture experiments

HEK293T (ATCC) cells were maintained in DMEM medium 4.5 g/L D-glucose

(Gibco) supplemented with 10% FBS (Gibco). Cells were tested for Mycoplasma

contamination by sequencing (Eurofins Genomics). For blocking the PPP, cells were

supplemented with 16 and 100 µM 6-AN for 24 to 48 h. After, cells were washed and added

fresh DMEM medium or transferred to starvation media (NaCl 140 mM, CaCl$_2$ 1 mM, MgCl$_2$

1 mM, HEPES pH 7.4 20 mM, BSA 1%) for 6 h. Cells were collected for immunofluorescence,

RNA extraction, and metabolite quantification.

For addressing MondoA localization, cells were fixed with 4% paraformaldehyde (PFA) for

15 min at room temperature. Following fixation, the samples were washed 3 times with 500

µL DPBS, followed by 500 µL of methanol 99% for 5 min at -20 °C, and washed twice with

500 µL of DPBS and three times with TBST 1X. Cells were blocked with 10% goat serum

diluted in DPBS for 1 h at room temperature. Cells were incubated with 200 µL of the Anti-

MondoA antibody (Bethyl Laboratories, A303-195A) 1:500 diluted in goat serum overnight at

4 °C. Next, cells were washed 3 times with DPBS containing Triton x-100 (PBST) and

similarly incubated with 200 µL of Alexa-488 secondary antibody (Invitrogen, A21206) 1:500

diluted in the goat serum at room temperature and covered in aluminum foil for 2 h. After

washing 3 times with 500 µL of PBST, the samples were covered with a mounting medium

containing DAPI. The cells were cured for 24 h at room temperature in the dark and then stored

at 4 °C.

Demographic analysis

Lifespan analyses were conducted by synchronizing worms by egg lay. All animals

were kept at 20 °C egg on, or transferred to 20 °C on day one of adulthood for the experiments

including *glp-1(e2141)*. More than 120 worms were used per strain/condition with 20 to 30

worms per 6 cm plate. Worms were transferred every other day to fresh plates until they

stopped laying eggs and were scored every time they were transferred. Animals with internal

hatching and vulva protrusions were censored from the analysis. All lifespans were performed

by blinding the genotypes/RNAi. Demographic parameters like mean, median, and maximum

lifespan were calculated and plotted using GraphPad Prism 9 software.

Oxygen consumption rate measurements

Oxygen consumption rates (OCR) measurements in *C. elegans* were based on

(Koopman et al., 2016). Worms were synchronized by egg lay. 10 to 15 adult worms were

transferred to a Seahorse XF96 Cell Culture Microplate (Agilent) with 200 μL of M9 buffer.

OCR was measured using the Seahorse XFe96 Analyzer (Agilent). To measure mitochondrial

fatty acid β-oxidation, basal oxygen consumption was measured followed by injection of

etomoxir (final well concentration 500 μM) and finally injection of sodium azide (final well

concentration 40 mM), all measured five times. OCR was normalized to the number of worms

with at least six technical replicates.

Body measurements, pharyngeal pumping rates, and motility assay

Worms were imaged or recorded using the Leica stereo microscope MDG41 equipped

with Leica DFC3000G monochrome camera with a 6.3X magnification. For measuring the

body length and area, >20 worms were transferred to a drop of sodium azide 50 mM, imaged,

and analyzed in Fiji imaging software. To measure the pharyngeal pumping rates, worms were

transferred to a new plate, acclimated on the plates without a lid for 10 min, and then recorded

for 30 s. Worm motility was assessed by placing >20 worms in a drop of M9 buffer and

recording for 30 s using the Leica M80 Binocular Microscope equipped with the Leica MC170

HD camera. Videos were analyzed to count the thrashing of each worm.

Molecular cloning

Plasmids were constructed by classical cloning or Gibson Assembly. Restriction

enzymes from NEB were used according to the manufacturer's instructions. Custom primers

were designed with SnapGene v.5.3.2 and NEBuilder Assembly Tool v.2.5.4 (**Supplementary**

Table S5). For classical cloning, digested plasmids and amplicons were ligated using the T4

DNA Ligase according to the manufacturer's instructions (New England Biolabs). For Gibson

Assembly, digested plasmid and amplicon were assembled using NEBuilder HiFi DNA

Assembly Master Mix (New England Biolabs) according to the manufacturer's instructions.

Plasmids were transformed into *E. coli* DH5α (Life Technologies) strain using the heat shock

protocol and plated on LB media supplemented with ampicillin.

RNA extraction and RT-qPCR

Worms were synchronized by egg lay for 4 h and collected once they reached the young

adult stage in 15 mL conical tubes. The worm pellet was subjected to 4 cycles of freeze and

thaw with liquid nitrogen and a water bath at ~37 °C. To lyse the worms, 200 μL of 1.0 mm

Zirconia/Silica beads (FisherScientific) were added and transferred to the TissueLyser LT

(Qiagen) for 30 min, full speed at 4 °C. Samples were incubated for 5 min at RT. After, 200

μL chloroform was added and centrifuged at 12,000 g for 15 min at 4 °C. The top aqueous

phase was transferred to a new 1.5 mL tube and mixed with 200 μL of ethanol. The RNA

purification was performed using the RNeasy Mini Kit (Qiagen) according to the

manufacturer's instructions. The concentration and purity of the RNA samples were quantified

using the NanoPhotometer NP80 spectrophotometer (Implen).

cDNA was prepared after extracting total RNA from young adult worms or cell extracts

using the iScript cDNA Synthesis Kit (BioRad) according to the manufacturer's instructions.

Primers, cDNA, and Power SYBR Green Master Mix (Applied Biosystems) were transferred

to a 384-well plate using the JANUS automated workstation (PerkinElmer) with four technical

replicates per genotype/gene. The reaction was quantified using the ViiA 7 Real-Time PCR

system machine (Applied Biosystems). *cdc-42* was used as an endogenous control for the

quantification. The complete primer sequences are in **Supplementary Table S5**.

Oxidative stress assay

Worms were grown on NGM plates until day 1 adults and then transferred to NGM

supplemented with paraquat at 15 and 35 mM for 6 h. As control, we used the CL2166 strain

that contains the transcriptional reporter *gst-4p*::GFP. MML-1::GFP nuclear localization was

measured in the intestine.

Protein extraction and Western blot analysis

Worms were harvested from a mixed population or synchronized by egg lay or

bleaching with M9 buffer and washed three times. The worm pellet was snap-frozen in liquid

nitrogen and stored at -80 °C until extraction. The pellet was resuspended in liquid lysis buffer

supplemented with cOmplete ULTRA EDTA-free protease inhibitors (Roche) and PhosSTOP

phosphatase inhibitors (Roche) and sonicated with Bioruptor Plus (Diagenode S.A.) coupled

to Minichiller 300 (Huber) for 15 cycles of 30 sec sonication, 30 sec rest at 4 °C. The total

protein extract was cleared by spinning down at 20,817 g for 10 min at 4 °C. Supernatant was

transferred to a new tube, and protein was quantified using Pierce BCA Protein Assay Kit

(ThermoFisher Scientific) according to the manufacturer's instructions. Equal amounts of

protein were mixed with 4X loading sample buffer containing DTT 50 mM and boiled at 95

°C for 10 min. Samples were loaded onto an SDS-PAGE PERCENTAGE gel (BioRad), and

proteins were then transferred onto a nitrocellulose membrane (BioRad) using the Trans-Blot

Turbo Transfer System (BioRad). Membranes were blocked for 1 h at RT with 5% milk in

TBST. Primary and secondary antibodies were diluted in 5% bovine serum albumin (BSA) in

TBST and incubated for 2-18 h at 4 °C. Antibodies: RFP Tag Polyclonal Antibody 1:1000

(ThermoFisher Scientific, R10367), Monoclonal ANTI-FLAG M2, Clone M2 1:1000 (Sigma

Aldrich, F1804), Anti-beta Actin Antibody 1:1000 (Abcam, Ab8224), Anti-rabbit IgG (H+L)

1:5000 (Invitrogen, G21234), Anti-mouse IgG 1:5000 (Invitrogen, G21040), Rabbit Anti-

Mouse IgG (Ligh Chain Specific) (D3V2A) mAB (HRP Conjugate) 1:5000 (CST, 58802). The

membranes were washed three times with TBST for 10 min at RT in-between incubation with

the antibodies and after the secondary. The signals were detected using Western Lightning

Plus-ECL, Enhanced Chemiluminescence Substrate (PerkinElmer), and imaged in the

ChemiDoc MP Imaging System with Image Lab software (BioRad).

Immunoprecipitation

Worms were collected from a mixed population of twenty 10 cm plates, transferred into

15 mL conical tubes with M9 buffer, and washed three times. Worms were snap-frozen in

liquid nitrogen and stored at -80 °C until extraction. The worm pellet was resuspended in lysis

buffer supplemented with cOmplete ULTRA EDTA-free protease inhibitors (Roche) and

PhosSTOP phosphatase inhibitors (Roche). Worms were sonicated with Bioruptor Plus

(Diagenode S.A.) coupled to Minichiller 300 (Huber) for 30 cycles of 30 sec sonication, 30 sec

rest at 4 °C. The total extract was cleared by spinning down at 20,817 g for 15 min at 4 °C.

Supernatant was transferred to a new tube, and protein was quantified using Pierce BCA

Protein Assay Kit (ThermoFisher Scientific) according to the manufacturer's instructions.

The Anti-FLAG M2 magnetic beads (Sigma) were resuspended and equilibrated with

TBST buffer. A total of 7 mg of total protein extract in 800 μL was mixed with the magnetic

beads and incubated overnight at 4 °C. The next day, the supernatant was removed and stored

for WB analysis. The beads were washed 3 times with 1 mL of cold washing buffer #1

(Tris/HCl, pH 7.4 20 mM, NaCl 300 mM, EDTA 1 mM, NP-50 0.5%), and 3 times with 1 mL

of cold washing buffer #2 (Tris/HCl, pH 7.4 20 mM, NaCl 300 mM, EDTA 1 mM). In the last

wash, 500 μL of washing buffer #2 was added, and 100 μL were stored for WB analysis.

To elute samples for mass spectrometry, 100 μL of elution buffer (Trypsin 5 ng/μL,

Tris/HCl, pH 7.5 50 mM, Tris(2-carboxyethyl)phosphine (TCEP) 1 mM, Chloroacetamide

(CAA) 5 mM) were added and incubated for 30 min at RT. The supernatant was transferred to

a new 0.5 mL tube and incubated overnight at 37 °C. The next day, formic acid was added to

a final concentration of 1% to stop the digestion. Peptides were stored at -20 °C until

purification.

The peptides were cleaned and purified with StageTip. A 30 μg C18-SD tip was

hydrated with 200 μL of methanol, followed by 200 μL of 40% acetonitrile (ACN)/0.1%

formic acid (FA). 200 μL of 0.1% FA was added to equilibrate the C18-SD. The digested

peptides were dissolved in 0.1% FA and added to the C18-SD and washed once with 0.1% FA.

To elute the peptides, 80 μL of 40% ACN/0.1% FA was added, and elutes were collected in

0.5 μL tubes and dried in a centrifugal vacuum Concentrator Plus (Eppendorf) for 45 min at

45 °C.

LC-MS/MS analysis

Peptides were separated on a 25 cm, 75 μm internal diameter PicoFrit analytical column

(New Objective) packed with 1.9 μm ReproSil-Pur 120 C18-AQ media (Dr. Maisch) using an

EASY-nLC 1200 (ThermoFisher Scientific). The column was maintained at 50 °C. Buffer A

and B were 0.1% FA in water and 0.1% FA in 80% CAN, respectively. Peptides were separated

on a segmented gradient from 6 to 31% buffer B for 57 min and from 31 to 44% buffer B for

5 min at 250 nl/min. Eluting peptides were analyzed on an Orbitrap Fusion Tribrid mass

spectrometer (ThermoFisher Scientific). Peptide precursor m/z measurements were carried out

at 60,000 resolution in the 350 to 1,500 m/z range. The most intense precursors with charge

states from 2 to 7 were selected for HCD fragmentation using 27% normalized collision energy.

The cycle time was set to 1 sec. The m/z values of the peptide fragments were measured at a

resolution of 50,000 using an AGC target of 2e5 and 86 ms maximum injection time. Upon

fragmentation, precursors were put on a dynamic exclusion list for 45 sec.

The raw data were analyzed with MaxQuant v.1.6.1.0 (Cox & Mann, 2008). Peptide

fragmentation spectra were searched against the canonical and isoform sequences of the *C.*

elegans reference proteome (proteome ID UP000001940, downloaded December 2018 from

UniProt). Methionine oxidation and protein N-terminal acetylation were set as variable

modifications; cysteine carbamidomethylation was set as fixed modification. The digestion

parameters were set to "specific" and "Trypsin/P". The minimum number of peptides and razor

peptides for protein identification was 1; the minimum number of unique peptides was 0.

Protein identification was performed at a peptide spectrum match and protein false discovery

rate of 0.01. The "second peptide" option was on. Successful identification was transferred

between the different raw files using the "Match between runs" option. Label-free

quantification (LFQ) (Cox et al., 2014) was performed using an LFQ minimum ratio count of

2. LFQ intensities were filtered for at least two valid values in at least one group and imputed

from a normal distribution with a width of 0.3 and a downshift of 1.8. Differential expression

analysis was performed using *limma* package v.3.34.9 (Ritchie et al., 2015) in R v.3.4.3 (R

Core Team, 2017).

G6P and NADP$^+$/NADPH measurements and TKT activity assay

G6P and NADP$^+$/NADPH were measured from total worm extracts or cell extracts

using the Glucose-6-Phosphate Assay Kit (Sigma MAK014) and the NADP/NADPH-Glo

Assay (Promega G9081), respectively, following the manufacturer's instructions. The

Transketolase Activity Assay Kit (Fluorometric) (Sigma MAK420) was used to measure TKT

activity in worm extracts following the manufacturer's instructions. Samples were normalized

to total protein concentration.

glp-1(e2141) strains were used to avoid signals from the germline. Briefly, worms were

synchronized as mentioned previously, collected in a 15 mL conical tube, and washed three

times with M9 buffer. Worms were snap-frozen in liquid nitrogen and sonicated using the

Bioruptor Plus (Diagenode S.A.) coupled to Minichiller 300 (Huber) for 15 cycles of 30 sec

sonication, 30 sec rest at 4 °C. Worms' extracts were cleared by centrifugation at 20,817 g for

15 min at 4 °C. Samples were always kept on ice. As mentioned, part of the samples was used

to quantify protein, and the rest was used to quantify the metabolites.

For G6P colorimetric detection, samples were run in technical duplicates in a 96-well
plate. The standard curve was made using the G6P Standard solution provided by diluting with
ddH$_2$O to generate 0 (blank), 2, 4, 6, 8, and 10 nmol/well standards. Samples were combined
with the reaction mix in each well and mixed for 30 min at RT, protecting the plate from the
light. The absorbance was measured at 450 nm using the POLARstar OMEGA Plate Reader
with Luminescence (BGM Labtech). For the analysis, all samples were corrected for the
background. The standard curve was generated with the appropriate values obtained by the
standards mentioned previously, and the amount of G6P in each sample was calculated with
the standard curve. Values were normalized to protein levels.

For the NADP$^+$/NADPH bioluminescent assay, samples were run in technical
duplicates in a 96-well plate. 50 µL of worm lysis in M9 buffer was incubated with 50 µL base
solution with 1% DTAB. Samples were split into two: one to measure NADP$^+$ and the other to
measure NADPH. Samples were incubated with or without 25 µL 0.4 N of HCl to measure
NADP$^+$ or NADPH, respectively, at 60 °C for 15 min, and then the plate was equilibrated for
10 min at RT. Next, 25 µL of Trizma base was added to each well of acid-treated samples to
neutralize the acid, and 50 µL of HCl/Trizma solution was added to each well of base-treated
samples. An equal amount of the NADP/NADPH-Glo Detection Reagent was added to each
well and mixed at RT for 30 min by protecting the plate from the light. The luminescence was
recorded using the POLARstar OMEGA Plate Reader with Luminescence (BGM Labtech).
Samples were corrected for background luminescence. The relative luminescent units were
standardized to the protein concentration.
820

Brightfield and fluorescent imaging
Brightfield and fluorescent images with higher resolution were acquired using the Zeiss
Axio Imager Z1 Fluorescent Microscope equipped with the Digital Microscope Camera Axio
Cam Mono 506 and the IcC5 true color camera. For fluorescent microscopy, the microscope
used a Colibri 7 LED Light Source and the filter sets GFP (Set 38 HE), TR (Set 45), and TL.
The software used was Zeiss Zen v2.3.69.1017. Brightfield and fluorescent images were also
acquired with the Leica stereo microscope MDG41 equipped with Leica DFC3000G
monochrome camera. The microscope was equipped with the Leica EL6000 external light
source. The software used was the Leica Application Suite X v3.7.5.24914.
All dyes were added to the bacterial lawn, incubated egg on, and protected from the
light until the day of imaging. Final solutions were made by diluting in M9 buffer. LipiBlue

(Dojindo) was resuspended in 100 μL of DMSO to make a 100 μM stock. The final

concentration used was 1 μM. C1-BODIPY-C12 500/510 (ThermoFisher Scientific) was

dissolved in DMSO to make a 5 mM stock. The final concentration used was 1 μM. Worms

were mounted on a slide with a 5% agarose pad using sodium azide 50 mM. Levamisole 2 mM

was used as an anesthetic to image mitochondria.

Nuclear localization of MML-1::GFP was quantified in intestinal cells from day one

adult worms. Worms were anesthetized with sodium azide 40 mM. At least 20 worms were

imaged per genotype/condition. Quantification of the transcription factor nuclear localization

was performed by manually selecting each nucleus and calculating the pixel intensity per area.

The same area was used to calculate the cytosolic and background signal. Values were

corrected by background subtraction. Nuclear localization was calculated by dividing the

nuclear signal from the cytosolic signal.

Confocal Imaging

Confocal images were acquired with Leica TCS SP8 confocal microscope equipped

with a white light laser and a 63X 1.4 NA oil objective. The microscope was also equipped

with HyD detectors for fluorescent images and a PMT detector for the bright field. The LAS

X Life Science software was used to acquire images and 3D reconstruction. When multiple

fluorescent proteins or dyes were imaged, sequential scanning was used to reduce the bleed-

through of the signal from one channel to another. All images used for colocalization analysis

were deconvoluted with the installed LIGHTNING package to optimize the images and reduce

background noise.

Statistical analysis and software

The results are presented as mean + SD or SEM, as the figure legends indicate. The

number of biological replicates is shown in the figure legends as "N", while the number of

worms is presented as "n". Before comparing groups, the data were tested for Gaussian

distribution using the Kolmogorov-Smirnov normality test with Dallal-Wilkinson-Lilliefor *p*-

value. For normally distributed data, unpaired *t*-test with Welch's correction or one-way

ANOVA with Brown-Forsythe and Welch's corrections were calculated in GraphPad Prism 9

(GraphPad software). For lifespan analysis, *p*-values were calculated using Log-rank (Mantel-

Cox) test. Statistical data from all the lifespans are included in **Supplementary Table S2**.

Statistical significance for qPCR experiments was calculated with two-tailed *t*-test. For

colocalization analysis, Costes' randomization was performed to determine statistical

significance. Significance levels are ns = not significant $p>0.05$, * $p<0.05$, ** $p<0.01$, ***
$p<0.001$, **** $p<0.0001$.

Image analysis was performed using open-source Fiji package software. Additional
JACoP (Bolte & Cordelières, 2006) plugin was used for colocalization analysis. Densitometry
analyses were performed using GelAnalyzer v.19.1. Graphs were generated in GraphPad Prism
9 and using custom R scripts.

Acknowledgments
We would like to thank the CGC (University of Minnesota) and SunyBiotech for some
of the strains used in this study. We also thank Katrin Wollenweber and Nadine Hochhard for
their technical assistance and the Proteomics, Bioinformatics, and FACS & Imaging Core
Facilities at the MPI AGE for their support. Special thanks to Özlem Karalay, Andrea Annibal,
Roberto Ripa, Sarah Kreuz, and Victoria Martínez Miguel for the discussions and comments.
This work was funded by the Max Planck Society and the Cologne Graduate School for Ageing
Research (CGA).

Authors contributions
R.L. and A.A. designed the experiments and wrote the manuscript. R.L. performed
most of the experiments. M.N. established and performed the MondoA localization upon PPP
inhibition in cell culture, S.S.T. performed *hxk-1* lipid droplet and *raga-1* aging experiments;
M.V. acquired and processed the samples for proteomics and repeated biological replicates,
E.B. measured the NADPH/NADP$^+$ levels in mammalian cells, K.S. carried out *raga-1* aging
experiments, I.A. performed proteomic analysis, T.D. and A.L. provided technical assistance.

Competing interests
The authors declare no competing interests.

Figures

Figure 1. Hexokinases are required for germline longevity by regulating MML-1 localization and function. A. Screening for MML-1 regulators upon knockdown of glucose metabolism enzymes. For clarity, we only mapped the candidate genes involved in glycolysis,

the TCA cycle, and PPP. See **Supplementary Table S1** for a complete list of genes tested.

MML-1::GFP (dhIs989) nuclear localization was measured in intestinal cells of day one adult

worms grown in the RNAi egg-on. Each circle represents one gene encoding an isozyme or

subunit of the complex. Circles are color-coded to indicate the increase (red) or decrease (blue)

of MML-1::GFP nuclear localization upon knockdown of the enzymes compared to internal

luci controls. DHAP, dihydroxyacetone phosphate; PEP, phosphoenolpyruvate; TCA cycle,

tricarboxylic acid cycle; PPP, pentose phosphate pathway. **B.** Quantification of MML-1 nuclear

localization upon knockdown of glucose metabolism genes. Only genes that statistically

affected MML-1 nuclear localization compared to the internal *luci* control are depicted

($p<0.05$; N≥2). **C-D.** Representative images (**C**) and quantification (**D**) of MML-1 nuclear

localization in wildtype and the germline-less *glp-1(e2141)* longevity background upon

knockdown of *luci* (control), *hxk-1i*, *hxk-2i*, and *mml-1i*. MML-1::GFP was quantified in nuclei

from intestinal cells (white arrows) in day one adult worms (N=3). Scale bars, 10 μm. **E.**

Relative mRNA levels of MML-1 downstream target *mdl-1* in young adult worms measured

by qPCR (N=3). **F.** Lifespans of wildtype and *glp-1(e2141)* upon *hxk-1* and *hxk-2* knockdown.

Lifespans for *hxk-1* and *hxk-2* knockdown were performed in the same experiment and plotted

separately for clarity; therefore, control lifespans are shared between plots. **G.** Quantification

of MML-1 nuclear localization in wildtype and *isp-1(qm150)* mutants upon *luci*, *hxk-1i*, and

hxk-2i in day one adult worms (N=3). **H.** Lifespan of wildtype and *isp-1(qm150)* upon *hxk-1*

and *hxk-2* knockdown (N=2). Statistical significance was calculated with *t*-test in **A** and **B,**

one-way ANOVA in **D** and **G**, two-tailed *t*-test in **E**, and Log-rank (Mantel-Cox) test in **F** and

H. **D** and **G** bars represent mean ± SD, and **E** bars represent mean ± SEM.

Supplementary Figure 1. Hexokinases regulate MML-1 nuclear localization and
transcriptional function. A. MML-1 nuclear localization in the fosmid line wgIs198 was
quantified in intestinal cells of day one adult worms using upon *hxk-1* and *hxk-2* knockdown
(N=3). **B**. Quantification of MML-1 nuclear localization in intestinal cells of day one adult
wildtype and *glp-1(e2141)* worms supplemented with the glycolytic inhibitor 3-bromopyruvate
(3-BrP) 5, 10, and 20 mM egg-on (N=2). **C**. Quantification of MML-1 nuclear localization in
intestinal cells of day one adult *glp-1(e2141)* worms supplemented with the glycolytic inhibitor
2-deoxy-glucose (2-DG) 50 mM egg-on (N=2). **D-E.** Representative Western blot (**D**) and
densitometry analysis (**E**) of hexokinases basal levels under *hxk-1i*, *hxk-2i*, *hxk-3i*, and *mml-1i*.
Protein levels were normalized to actin as a loading control (N=3). **F-H.** Measurements of body
size area (**F**), length (**G**), and pharyngeal pumping rate (**H**) of day one adult worms grown
under hexokinase knockdown (N=3). **I**. Motility assay of day one adult worms grown under
hexokinase knockdown (N=2). **J.** Relative *mml-1* mRNA levels under hexokinase knockdown
measured by qPCR (N=3). **K.** Relative mRNA expression of MML-1 downstream targets
lgg-2, *swt-1*, and *fat-5* measured by qPCR (N=3). **L.** Quantification of MML-1 nuclear
localization in wildtype and *daf-2(e1370)* mutants upon *luci*, *hxk-1i*, and *hxk-2i* in day one
adult worms (N=3). **M.** Lifespan of wildtype and *daf-2(e1370)* upon *hxk-1* and *hxk-2*
knockdown (N=2). **N.** Quantification of MML-1 nuclear localization in wildtype and *raga-*
1(ok701) mutants in day one adult worms (N=3). **O.** Lifespan of wildtype, *raga-1(ok701)*, *mml-*
1(ok849) and *raga-1 mml-1* double mutant (N=3). **P.** Lifespan of wildtype and *raga-1(ok701)*
upon *hxk-1* and *hxk-2* knockdown (N=3). Statistical significance was calculated with *t*-test in
C and **N**, one-way ANOVA in **A-B** and **E-L**, and Log-rank (Mantel-Cox) test in **M** and **O-P**.
A-C, **F-J**, **L**, and **N** bars represent mean ± SD, and **E** and **K** bars represent mean ± SEM.
946

Figure 2. Enhanced mitochondrial β-oxidation suppresses MML-1 nuclear localization upon *hxk-1* knockdown. A-B. Tissue expression of HXK-1::mKate2 (**A**) and HXK-2::mKate2 (**B**) isozymes. The primary tissues are identified with arrows. Scale bars, 20 μm. **C.** Colocalization of mitochondrial reporter *myo-3p*::mitoGFP with full-length HXK-2::mKate2 and d10 HXK-2::mKate2 lacking the first ten residues of the N-terminus. Scale bars, 10 μm. **D.** Representative confocal images of day one adult worms expressing MML-1::GFP and the mitochondrial reporter *ges-1p*::TOMM-20::mCherry under *luci* and *hxk-2i*. Scale bars, 10 μm. **E.** Intensity plot of the region of interest (ROI) from *hxk-2i* (**D**). **F.** Colocalization of MML-1::GFP and TOMM-20::mCherry in *luci* and *hxk-2i*. The colocalization analysis was

calculated with the thresholded Manders' colocalization coefficients tM1 (GFP channel) and
tM2 (mCherry channel) (N=3). **G.** When glycolysis is reduced, cells rely on fatty acids for
energy production. Free fatty acids are activated to acyl-CoA and transported to the
mitochondria through the CPT in the mitochondrial outer membrane. The acyl-CoA is then
metabolized to generate acetyl-CoA that can enter the TCA cycle to produce energy. MPC,
mitochondrial pyruvate carrier; CPT, carnitine palmitoyltransferase. **H-I.** Oxygen consumption
rate (OCR) under basal conditions (**H**) and after the addition of the mitochondrial β-oxidation
inhibitor etomoxir (**I**). Mitochondrial β-oxidation is calculated as the percentage of decreased
OCR after injecting etomoxir (N=3). **J.** MML-1 nuclear localization of day one adult worms
grown with double knockdown of *hxk-1* or *hxk-2* combined *acs-2i* (N=3). **K.** MML-1 nuclear
localization of day one adult worms grown upon *hxk-1* and *hxk-2* knockdown with
supplementation of etomoxir 100 μM egg-on (N=3). Statistical significance was calculated
with *t*-test in **F**, and one-way ANOVA in **H-K**. **F** and **J-K** bars represent mean ± SD, **H-I** bars
represent mean ± SEM.
972

Supplementary Figure 2. MML-1 regulation by glucose metabolism is independent of mitochondrial and oxidative stress response. A. Localization of HXK-3::mKate2 isozyme. The primary tissues are identified with arrows. Scale bars, 20 μm. **B.** Representative confocal images of day one adult worms expressing MML-1::GFP and the lysosome/endosome reporter LMP-1::mKate2 under *luci* and *hxk-2i*. Scale bars, 10 μm. **C.** Intensity plots of the ROIs from **B** (n>10 MML-1 positive puncta per worm). **D.** Quantification of mitochondrial stress response *hsp-6p*::GFP transcriptional reporter upon knockdown of candidate genes regulating MML-1 nuclear localization. Worms were imaged on day one of adulthood. As a positive control for induction of mitochondrial stress, worms were supplemented with antimycin A (AA) 10 μg/mL for 24 h before imaging (N=3). **E.** Quantification of MML-1::GFP nuclear localization of worms grown under *luci* and *hxk-2i* supplemented with AA at the L4 stage, and nuclear localization was measured after 24 h (N=3). **F.** Quantification of day one adult worms expressing *gst-4p*::GFP transcriptional reporter exposed to paraquat for 6 h (N=3). **G.** Quantification of MML-1::GFP nuclear localization of day one adult worms exposed to

 paraquat for 6 h (N=3). **H.** Neutral lipid content was measured on day one of adulthood by feeding worms with C1-BODIPY-C12 egg-on (N=3). **I.** Quantification of day one adult worms expressing the transcriptional reporter *acs-2p*::GFP under hexokinase knockdown (N=3). Statistical significance was calculated with one-way ANOVA in **D-G**. **D-G** bars represent mean ± SD.

Figure 3. HXK-2 regulates MML-1 localization and function through inhibition of the PPP. A. The PPP is a cytosolic pathway that metabolizes G6P to produce NADPH and ribose-5P, and interconverts 3 to 7 carbon sugars. 6-Aminonicotinamide (6-AN) is a PPP inhibitor that acts on the NADPH-producing enzymes GSPD-1 and T25B9.9. **B.** Quantification of MML-1 nuclear localization of day one adult worms supplemented with 500 μM and 1 mM of 6-AN egg-on (N=3). **C.** Quantification of NADP$^+$ and NADPH levels in day one adult *glp-1(e2141)* worms grown under hexokinase knockdown. *T25B9.9i* was used as a control. Metabolite levels were normalized to total protein concentration (N=3). **D.** Transketolase enzymatic activity assay on day one adult *glp-1(e2141)* worms grown under hexokinase knockdown. *tkt-1i* was used as a control. Activity levels were normalized to total protein concentration (N=3). **E-F.** Representative images (**E**) and quantification (**F**) of MML-1 nuclear

localization of day one adult worms upon double knockdown of *hxk-2* with the PPP enzymes *T25B9.9* and *tkt-1* (N=3). Scale bars, 20 μm. **G.** Relative mRNA levels of MML-1 downstream target *mdl-1* in young adult worms upon double knockdown of *hxk-2* in combination with enzymes of the PPP measured by qPCR (N=3). **H.** Quantification of G6P levels relative to total protein measured on day one adult *glp-1(e2141)* upon double knockdown of *hxk-2* in combination with enzymes of the PPP (N=4). **I.** Lifespan analysis of *glp-1(e2141)* with *hxk-2i* and *tkt-1i* single and double knockdowns. Lifespan was performed in the same experiment as **Supplementary Figure 3J**; therefore, control lifespans are shared between plots (N=3). Statistical significance was calculated with a one-way ANOVA test in **B-D** and **F-H**, and Log-rank (Mantel-Cox) test in **I**. **B** and **F** bars represent mean ± SD, **C-D** and **G-H** bars represent mean ± SEM.

Supplementary Figure 3. HXK-1 and HXK-2 regulate MML-1 through parallel or independent pathways. A. MML-1 nuclear localization of day one adult worms upon double knockdown of *hxk-1* with enzymes of the PPP (N=3). **B-D.** Relative mRNA levels of MML-1 downstream targets *lgg-2* (**B**), *swt-1* (**C**), and *fat-5* (**D**) in young adult worms with double knockdown of *hxk-2* with enzymes of the PPP measured by qPCR (N=3). **E-H.** Relative mRNA levels of MML-1 downstream targets *mdl-1* (**E**), *lgg-2* (**F**), *swt-1* (**G**), and *fat-5* (**H**) in young adult worms with double knockdown of *hxk-1* with enzymes of the PPP measured by qPCR (N=3). **I.** Quantification of G6P levels relative to total protein measured on day one adult *glp-1(e2141)* worms grown under *hxk-1i* in combination with enzymes from the PPP (N=4). **J.** Lifespan analysis of *glp-1(e2141)* with *hxk-2i* and *T25B9.9i* single and double knockdowns. Lifespan was performed in the same experiment as **Figure 3I**; therefore, control lifespans are shared between plots (N=3). Statistical significance was calculated with a one-way ANOVA test in **A-I**, and Log-rank (Mantel-Cox) test in **J**. **A** bars represent mean ± SD, **B-I** bars represent mean ± SEM.

1036

Figure 4. *hxk-2* knockdown increases MML-1 colocalization with LD. A. Anti-FLAG immunoprecipitation (IP) from whole-worm lysates of animals expressing MML-1 endogenously tagged with 3xFLAG in wildtype background. IP was analyzed by mass spectrometry (LC-MS/MS). The 25 top candidates co-enriched with MML-1 compared to non-transgenic wildtype control are shown as log FC, including MML-1 (blue) and its established binding partner MXL-2 (red). **B.** Crosstalk between glycolysis, PPP, and lipid droplet (LD) metabolism. On the one hand, LD expansion is favored by increased fatty acid synthesis and subsequent formation of triacylglycerol (TAG) by the diacylglycerol acetyltransferase DGAT-2. On the other hand, LD consumption releases free fatty acids from TAG by the rate-limiting lipase ATGL-1. **C.** Confocal microscopy of day one adult worms expressing MML-1::GFP and the mitochondrial reporter TOMM-20::mCherry stained with the LD dye LipiBlue under *luci* and *hxk-2i* (N=3). Scale bars, 10 μm. **D.** Intensity plot of the region of interest (ROI) from *hxk-2i* (**C**). **E.** Colocalization of MML-1::GFP and LD in *luci* and *hxk-2i*. The colocalization analysis was calculated with the thresholded Manders' colocalization coefficients tM1 (GFP channel) and tM2 (LipiBlue channel) (N=3). **F.** Quantification of LD diameter in day one adult worms expressing the LD reporter PLIN-1::mCherry upon knockdown of *hxk-2* and *tkt-1* (N=3). **G.** Quantification of MML-1 nuclear localization under knockdown of the LD metabolic enzymes *dgat-2* and *atg-1* (N=3). **H.** Quantification of MML-1 nuclear localization upon double knockdown of *tkt-1* and *atgl-1* (N=3). Statistical significance was calculated with a one-way ANOVA test in **E-H**. **E-H** bars represent mean ± SD.

Supplementary Figure 4. MML-1 interactome is enriched with proteins from different

organelles. A. Quantified proteins in each biological replicate of wildtype and *mml-1::3xFLAG*

anti-FLAG immunoprecipitation (IP). **B**. Principal component analysis (PCA) with each

biological replicate of the genotypes from proteomics. **C.** Top candidates co-enriched in

MML-1::FLAG IP grouped by predicted subcellular localization (uniprot.org). **D.**

Representative images of day one adult worms expressing the LD proteins DHS-3::GFP and

PLIN-1::mCherry reporters grown with the LD dye LipiBlue egg-on. Scale bars, 10 μm. **E-F.**

Intensity plots of the marked ROI from **D** showing DHS-3::GFP (**E**) and PLIN-1::mCherry (**F**)

colocalization with LipiBlue. **G.** Expression of the ATGL-1::GFP translational reporter

measured on day one adult worms grown under *T25B9.9* and *tkt-1* RNAi egg-on (N=3). **H.**

Quantification of LD reporter PLIN-1::mCherry intensity in day one adult worms upon

knockdown of *hxk-2* and *tkt-1* (N=3). **I.** Quantification of LD diameter in day one adult worms

expressing the LD reporter PLIN-1::mCherry upon knockdown of *hxk-1* and *acs-2* (N=3). **J.**

Quantification of neutral lipid content in day one adult worms supplemented with

C1-BODIPY-C12 egg-on under *dgat-2* and *atgl-1* knockdown (N=3). Statistical significance

was calculated with a one-way ANOVA test in **G-J**. **G-J** bars represent mean ± SD.

1076

1077

Figure 5. PPP regulates MondoA localization and transcriptional function in HEK293T

cells. A. Immunofluorescence of MondoA in HEK293T cells under high glucose media (HG)

and starvation (S) conditions supplemented with the PPP inhibitor 6-aminonicotinamide (6-

AN) 100 μM. Scale bars, 20 μm. **B.** Quantification of MondoA nuclear localization in

HEK293T cells treated with 6-AN 16 and 100 μM under HG and S conditions (N=3). **C.**

Relative mRNA levels of MondoA in HEK293T cells treated with 6-AN 16 and 100 μM under

HG and S conditions measured by qPCR (N=3). **D.** Relative mRNA levels of MondoA

downstream target *TXNIP* in HEK293T cells treated with 6-AN 16 and 100 μM under HG and

S conditions measured by qPCR (N=3). Statistical significance was calculated with one-way

ANOVA in **B-D**. **B** bars represent mean ± SD, **C-D** bars represent mean ± SEM. **E.** Working

model of MML-1 regulation by hexokinases. HXK-1 and HXK-2 positively regulate MML-1

through different downstream mechanisms. On the one hand, mitochondrial β-oxidation

inhibition rescues MML-1 nuclear localization upon *hxk-1i*. On the other hand, the PPP

regulates MML-1 subcellular localization between LD-mitochondria and the nucleus under

hxk-2i, which is required for the lifespan extension mediated by germ-line loss. PPP, pentose

phosphate pathway; LD, lipid droplet.

1094

Supplementary Figure 5. Inhibition of PPP NADPH-producing enzymes by 6-AN induces

MondoA downstream targets. A. Experimental design for inhibition of the PPP in HEK293T

cells. Cells were grown in high glucose media (HG) supplemented with 6-AN for 24 h, and

then the media was replaced with either HG or starvation (S) media for 6 h. Cells were collected

for immunostaining, RNA extraction, and metabolite quantification. **B.** Quantification of

NADP$^+$ and NADPH levels in HEK293T cells treated with 6-AN 16 and 100 μM under HG

and S conditions (N=3). **C.** Relative mRNA levels of MondoA downstream target *ARRDC4* in

HEK293T cells treated with 6-AN 16 and 100 μM under HG and S conditions measured by

qPCR (N=3). Statistical significance was calculated with one-way ANOVA in **B-C**. **B-C** bars

represent mean ± SEM.

1105

References

Benhamed, F., Denechaud, P. D., Lemoine, M., Robichon, C., Moldes, M., Bertrand-Michel,

J., Ratziu, V., Serfaty, L., Housset, C., Capeau, J., Girard, J., Guillou, H., & Postic, C.

(2012). The lipogenic transcription factor ChREBP dissociates hepatic steatosis from

insulin resistance in mice and humans. *Journal of Clinical Investigation*, *122*(6), 2176–

2194. https://doi.org/10.1172/JCI41636

Bennett, C. F., Kwon, J. J., Chen, C., Russell, J., Acosta, K., Burnaevskiy, N., Crane, M. M.,
Bitto, A., Vander Wende, H., Simko, M., Pineda, V., Rossner, R., Wasko, B. M., Choi,
H., Chen, S., Park, S., Jafari, G., Sands, B., Perez Olsen, C., … Kaeberlein, M. (2017).
Transaldolase inhibition impairs mitochondrial respiration and induces a starvation-like
longevity response in Caenorhabditis elegans. In *PLoS Genetics* (Vol. 13, Issue 3).
https://doi.org/10.1371/journal.pgen.1006695

Billin, A. N., Eilers, A. L., Coulter, K. L., Logan, J. S., & Ayer, D. E. (2000). MondoA, a
Novel Basic Helix-Loop-Helix–Leucine Zipper Transcriptional Activator That
Constitutes a Positive Branch of a Max-Like Network. *Molecular and Cellular Biology*,
20(23), 8845–8854. https://doi.org/10.1128/mcb.20.23.8845-8854.2000

Bolte, S., & Cordelières, F. P. (2006). A guided tour into subcellular colocalization analysis
in light microscopy. *Journal of Microscopy*, *224*(3), 213–232.
https://doi.org/10.1111/j.1365-2818.2006.01706.x

Boutant, M., Kulkarni, S. S., Joffraud, M., Ratajczak, J., Valera-Alberni, M., Combe, R.,
Zorzano, A., & Cantó, C. (2017). Mfn2 is critical for brown adipose tissue thermogenic
function. *The EMBO Journal*, *36*(11), 1543–1558.
https://doi.org/10.15252/embj.201694914

Bricker, D. K., Taylor, E. B., Schell, J. C., Orsak, T., Boutron, A., Chen, Y.-C., Cox, J. E.,
Cardon, C. M., Van Vranken, J. G., Dephoure, N., Redin, C., Boudina, S., Gygi, S. P.,
Brivet, M., Thummel, C. S., & Rutter, J. (2012). A Mitochondrial Pyruvate Carrier
Required for Pyruvate Uptake in Yeast, Drosophila, and Humans. *Science*, *337*(July),
96–100. https://doi.org/10.1126/science.1218099

Carroll, P. A., Diolaiti, D., McFerrin, L., Gu, H., Djukovic, D., Du, J., Cheng, P. F.,
Anderson, S., Ulrich, M., Hurley, J. B., Raftery, D., Ayer, D. E., & Eisenman, R. N.
(2015). Deregulated Myc Requires MondoA/Mlx for Metabolic Reprogramming and
Tumorigenesis. *Cancer Cell*, *27*(2), 271–285. https://doi.org/10.1016/j.ccell.2014.11.024

Cha-Molstad, H., Saxena, G., Chen, J., & Shalev, A. (2009). Glucose-stimulated expression
of Txnip is mediated by carbohydrate response element-binding protein, p300, and
histone H4 acetylation in pancreatic beta cells. *Journal of Biological Chemistry*,
284(25), 16898–16905. https://doi.org/10.1074/jbc.M109.010504

Chi, M. M. Y., Pusateri, M. E., Carter, J. G., Norris, B. J., McDougal, D. B., & Lowry, O. H.
(1987). Enzymatic assays for 2-deoxyglucose and 2-deoxyglucose 6-phosphate.
Analytical Biochemistry, *161*(2), 508–513. https://doi.org/10.1016/0003-2697(87)90481-
7

Chin, R. M., Fu, X., Pai, M. Y., Vergnes, L., Hwang, H., Deng, G., Diep, S., Lomenick, B.,

Meli, V. S., Monsalve, G. C., Hu, E., Whelan, S. A., Wang, J. X., Jung, G., Solis, G. M.,

Fazlollahi, F., Kaweeteerawat, C., Quach, A., Nili, M., … Huang, J. (2014). The

metabolite α-ketoglutarate extends lifespan by inhibiting ATP synthase and TOR.

Nature, *510*(7505), 397–401. https://doi.org/10.1038/nature13264

Clancy, D. J., Gems, D., Harshman, L. G., Oldham, S., Stocker, H., Hafen, E., Leevers, S. J.,

& Partridge, L. (2001). Extension of life-span by loss of CHICO, a Drosophila insulin

receptor substrate protein. *Science*, *292*(5514), 104–106.

https://doi.org/10.1126/science.1057991

Cox, J., Hein, M. Y., Luber, C. A., Paron, I., Nagaraj, N., & Mann, M. (2014). Accurate

proteome-wide label-free quantification by delayed normalization and maximal peptide

ratio extraction, termed MaxLFQ. *Molecular and Cellular Proteomics*, *13*(9), 2513–

2526. https://doi.org/10.1074/mcp.M113.031591

Cox, J., & Mann, M. (2008). MaxQuant enables high peptide identification rates,

individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.

Nature Biotechnology, *26*(12), 1367–1372. https://doi.org/10.1038/nbt.1511

Dalziel, K., & Londesborough, J. C. (1968). The mechanisms of reductive carboxylation

reactions: Carbon dioxide or bicarbonate as substrate of nicotinamide-adenine

dinucleotide phosphate-linked isocitrate dehydrogenase and malic enzyme. *The*

Biochemical Journal, *110*(2), 223–230. https://doi.org/10.1042/bj1100223

Denechaud, P. D., Dentin, R., Girard, J., & Postic, C. (2008). Role of ChREBP in hepatic

steatosis and insulin resistance. *FEBS Letters*, *582*(1), 68–73.

https://doi.org/10.1016/j.febslet.2007.07.084

Du, S., & Zheng, H. (2021). Role of FoxO transcription factors in aging and age-related

metabolic and neurodegenerative diseases. *Cell and Bioscience*, *11*(1), 1–17.

https://doi.org/10.1186/s13578-021-00700-7

Fischer, F., Grigolon, G., Benner, C., & Ristow, M. (2022). Evolutionarily Conserved

Transcription Factors As Regulators of Longevity and Targets for Geroprotection.

Physiological Reviews, *102*(3), 1449–1494. https://doi.org/10.1152/physrev.00017.2021

Gerisch, B., & Antebi, A. (2004). Hormonal signals produced by DAF-9/cytochrome P450

regulate C. elegans dauer diapause in response to environmental cues. *Development*,

131(8), 1765–1776. https://doi.org/10.1242/dev.01068

Gerisch, B., Weitzel, C., Kober-Eisermann, C., Rottiers, V., & Antebi, A. (2001). A

Hormonal Signaling Pathway Influencing C. elegans Metabolism, Reproductive

Development, and Life Span. *Developmental Cell*, *1*(6), 841–851.
https://doi.org/10.1016/S1534-5807(01)00085-5

Ghasemi, A., Aghajani, H., Fallah, S., Assadi, M., & Seifi, M. (2015). C771G (His241Gln)
polymorphism of MLXIPL Gene, TG levels and coronary artery disease: A case control
study. *Anatolian Journal of Cardiology*, *15*(1), 8–12.
https://doi.org/10.5152/akd.2014.5135

Gordaliza-Alaguero, I., Cantó, C., & Zorzano, A. (2019). Metabolic implications of
organelle–mitochondria communication. *EMBO Reports*, *20*(9), 1–27.
https://doi.org/10.15252/embr.201947928

Gottlob, K., Majewski, N., Kennedy, S., Kandel, E., Robey, R. B., & Hay, N. (2001).
Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step
of glycolysis and mitochondrial hexokinase. *Genes and Development*, *15*(11), 1406–
1418. https://doi.org/10.1101/gad.889901

Green, C. L., Lamming, D. W., & Fontana, L. (2021). Molecular mechanisms of dietary
restriction promoting health and longevity. *Nature Reviews Molecular Cell Biology*,
0123456789. https://doi.org/10.1038/s41580-021-00411-4

Hutter, H., & Suh, J. (2016). GExplore 1.4: An expanded web interface for queries on
Caenorhabditis elegans protein and gene function . *Worm*, *5*(4), e1234659.
https://doi.org/10.1080/21624054.2016.1234659

Iizuka, K., Miller, B., & Uyeda, K. (2006). Deficiency of carbohydrate-activated
transcription factor ChREBP prevents obesity and improves plasma glucose control in
leptin-deficient (ob/ob) mice. *American Journal of Physiology - Endocrinology and
Metabolism*, *291*(2), 358–364. https://doi.org/10.1152/ajpendo.00027.2006

Ishii, S., Iizuka, K., Miller, B. C., & Uyeda, K. (2004). Carbohydrate response element
binding protein directly promotes lipogenic enzyme gene transcription. *Proceedings of
the National Academy of Sciences of the United States of America*, *101*(44), 15597–
15602. https://doi.org/10.1073/pnas.0405238101

John, S., Weiss, J. N., & Ribalet, B. (2011). Subcellular localization of hexokinases I and II
directs the metabolic fate of glucose. *PLoS ONE*, *6*(3).
https://doi.org/10.1371/journal.pone.0017674

Johnson, D. W., Llop, J. R., Farrell, S. F., Yuan, J., Stolzenburg, L. R., & Samuelson, A. V.
(2014). The Caenorhabditis elegans Myc-Mondo/Mad Complexes Integrate Diverse
Longevity Signals. *PLoS Genetics*, *10*(4). https://doi.org/10.1371/journal.pgen.1004278

Ju, H. Q., Lin, J. F., Tian, T., Xie, D., & Xu, R. H. (2020). NADPH homeostasis in cancer:

functions, mechanisms and therapeutic implications. *Signal Transduction and Targeted Therapy*, *5*(1), 1–12. https://doi.org/10.1038/s41392-020-00326-0

Kaadige, M. R., Looper, R. E., Kamalanaadhan, S., & Ayer, D. E. (2009). Glutamine-dependent anapleurosis dictates glucose uptake and cell growth by regulating MondoA transcriptional activity. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(35), 14878–14883. https://doi.org/10.1073/pnas.0901221106

Kamath, R. S., & Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis elegans. *Methods*, *30*(4), 313–321. https://doi.org/10.1016/S1046-2023(03)00050-1

Kennedy, B. K., & Lamming, D. W. (2016). The Mechanistic Target of Rapamycin: The Grand ConducTOR of Metabolism and Aging. *Cell Metabolism*, *23*(6), 990–1003. https://doi.org/10.1016/j.cmet.2016.05.009

Koopman, M., Michels, H., Dancy, B. M., Kamble, R., Mouchiroud, L., Auwerx, J., Nollen, E. A. A., & Houtkooper, R. H. (2016). A screening-based platform for the assessment of cellular respiration in Caenorhabditis elegans. *Nature Protocols*, *11*(10), 1798–1816. https://doi.org/10.1038/nprot.2016.106

Krahmer, N., Najafi, B., Schueder, F., Quagliarini, F., Steger, M., Seitz, S., Kasper, R., Salinas, F., Cox, J., Uhlenhaut, N. H., Walther, T. C., Jungmann, R., Zeigerer, A., Borner, G. H. H., & Mann, M. (2018). Organellar Proteomics and Phospho-Proteomics Reveal Subcellular Reorganization in Diet-Induced Hepatic Steatosis. *Developmental Cell*, *47*(2), 205-221.e7. https://doi.org/10.1016/j.devcel.2018.09.017

Lapierre, L. R., De Magalhaes Filho, C. D., McQuary, P. R., Chu, C. C., Visvikis, O., Chang, J. T., Gelino, S., Ong, B., Davis, A. E., Irazoqui, J. E., Dillin, A., & Hansen, M. (2013). The TFEB orthologue HLH-30 regulates autophagy and modulates longevity in Caenorhabditis elegans. *Nature Communications*, *4*. https://doi.org/10.1038/ncomms3267

Lee, S.-J., Murphy, C.T., & Kenyon, C. (2009). Glucose Shortens the Lifespan of *Caenorhabditis elegans* by Down-Regulating Aquaporin Gene Expression. *Cell Metabolism, 10*(5), 379-391. https://doi.org/10.1016/j.cmet.2009.10.003

Lee, J. H., Kong, J., Jang, J. Y., Han, J. S., Ji, Y., Lee, J., & Kim, J. B. (2014). Lipid Droplet Protein LID-1 Mediates ATGL-1-Dependent Lipolysis during Fasting in Caenorhabditis elegans. *Molecular and Cellular Biology*, *34*(22), 4165–4176. https://doi.org/10.1128/mcb.00722-14

Leigh-Brown, S., Enriquez, J. A., & Odom, D. T. (2010). Nuclear transcription factors in mammalian mitochondria. *Genome Biology*, *11*(7), 1–9. https://doi.org/10.1186/gb-

2010-11-7-215

Li, F., Wang, Y., Zeller, K. I., Potter, J. J., Wonsey, D. R., O'Donnell, K. A., Kim, J.,

Yustein, J. T., Lee, L. A., & Dang, C. V. (2005). Myc Stimulates Nuclearly Encoded

Mitochondrial Genes and Mitochondrial Biogenesis. *Molecular and Cellular Biology*,

25(14), 6225–6234. https://doi.org/10.1128/mcb.25.14.6225-6234.2005

Lindén, M., Gellerfors, P., & Dean Nelson, B. (1982). Pore protein and the hexokinase-

binding protein from the outer membrane of rat liver mitochondria are identical. *FEBS*

Letters, *141*(2), 189–192. https://doi.org/10.1016/0014-5793(82)80044-6

Ma, L., Robinson, L. N., & Towle, H. C. (2006). ChREBP·Mlx is the principal mediator of

glucose-induced gene expression in the liver. *Journal of Biological Chemistry*, *281*(39),

28721–28730. https://doi.org/10.1074/jbc.M601576200

Marchenko, N. D., Wolff, S., Erster, S., Becker, K., & Moll, U. M. (2007).

Monoubiquitylation promotes mitochondrial p53 translocation. *EMBO Journal*, *26*(4),

923–934. https://doi.org/10.1038/sj.emboj.7601560

Marinov, G. K., Wang, Y. E., Chan, D., & Wold, B. J. (2014). Evidence for site-specific

occupancy of the mitochondrial genome by nuclear transcription factors. *PLoS ONE*,

9(1), 1–24. https://doi.org/10.1371/journal.pone.0084713

Massari, M. E., & Murre, C. (2000). *Helix-Loop-Helix Proteins: Regulators of Transcription*

in Eucaryotic. *20*(2), 429–440. https://doi.org/10.1128/mcb.20.2.429-440.2000

Mejhert, N., Kuruvilla, L., Gabriel, K. R., Elliott, S. D., Guie, M. A., Wang, H., Lai, Z. W.,

Lane, E. A., Christiano, R., Danial, N. N., Farese, R. V., & Walther, T. C. (2020).

Partitioning of MLX-Family Transcription Factors to Lipid Droplets Regulates

Metabolic Gene Expression. *Molecular Cell*, *77*(6), 1251-1264.e9.

https://doi.org/10.1016/j.molcel.2020.01.014

Motola, D. L., Cummins, C. L., Rottiers, V., Sharma, K. K., Li, T., Li, Y., Suino-Powell, K.,

Xu, H. E., Auchus, R. J., Antebi, A., & Mangelsdorf, D. J. (2006). Identification of

Ligands for DAF-12 that Govern Dauer Formation and Reproduction in C. elegans. *Cell*,

124(6), 1209–1223. https://doi.org/10.1016/j.cell.2006.01.037

Mutlu, A. S., Gao, S. M., Zhang, H., & Wang, M. C. (2020). Olfactory specificity regulates

lipid metabolism through neuroendocrine signaling in Caenorhabditis elegans. *Nature*

Communications, *11*(1), 1–15. https://doi.org/10.1038/s41467-020-15296-8

Nakamura, S., Karalay, Ö., Jäger, P. S., Horikawa, M., Klein, C., Nakamura, K., Latza, C.,

Templer, S. E., Dieterich, C., & Antebi, A. (2016). Mondo complexes regulate TFEB

via TOR inhibition to promote longevity in response to gonadal signals. *Nature*

Communications, *7*. https://doi.org/10.1038/ncomms10944

Orlandi, I., Coppola, D. P., & Vai, M. (2014). Rewiring yeast acetate metabolism through

mpc1 loss of function leads to mitochondrial damage and decreases chronological

lifespan. *Microbial Cell*, *1*(12), 393–405. https://doi.org/10.15698/mic2014.12.178

Owen, O. E., Kalhan, S. C., & Hanson, R. W. (2002). The key role of anaplerosis and

cataplerosis for citric acid cycle function. *Journal of Biological Chemistry*, *277*(34),

30409–30412. https://doi.org/10.1074/jbc.R200006200

Papsdorf, K., Miklas, J. W., Hosseini, A., Cabruja, M., Morrow, C. S., Savini, M., Yu, Y.,

Silva-garcía, C. G., Haseley, N. R., Murphy, L. M., Yao, P., Launoit, E. De, Dixon, S.

J., Snyder, M. P., Wang, M. C., Mair, W. B., & Brunet, A. (2023). Lipid droplets and

peroxisomes are co-regulated to drive lifespan extension in response to mono-

unsaturated fatty acids. *Nature Cell Biology*. https://doi.org/10.1038/s41556-023-01136-

6

Pastorino, J. G., Shulga, N., & Hoek, J. B. (2002). Mitochondrial binding of hexokinase II

inhibits Bax-induced cytochrome c release and apoptosis. *Journal of Biological*

Chemistry, *277*(9), 7610–7618. https://doi.org/10.1074/jbc.M109950200

Penkov, S., Kaptan, D., Erkut, C., Sarov, M., Mende, F., & Kurzchalia, T. V. (2015).

Integration of carbohydrate metabolism and redox state controls dauer larva formation in

Caenorhabditis elegans. *Nature Communications*, *6*.

https://doi.org/10.1038/ncomms9060

Pickett, C. L., Breen, K. T., & Ayer, D. E. (2007). A C. elegans Myc-like network cooperates

with semaphorin and Wnt signaling pathways to control cell migration. *Developmental*

Biology, *310*(2), 226–239. https://doi.org/10.1016/j.ydbio.2007.07.034

Postic, C., Dentin, R., Denechaud, P. D., & Girard, J. (2007). ChREBP, a transcriptional

regulator of glucose and lipid metabolism. *Annual Review of Nutrition*, *27*, 179–192.

https://doi.org/10.1146/annurev.nutr.27.061406.093618

R Core Team. (2017). *R: A language and environment for statistical computing* (3.4.3). R

Foundation for Statistical Computing. https://www.r-project.org/

Ralser, M., Wamelink, M. M., Struys, E. A., Joppich, C., Krobitsch, S., Jakobs, C., &

Lehrach, H. (2008). A catabolic block does not sufficiently explain how 2-deoxy-D-

glucose inhibits cell growth. *Proceedings of the National Academy of Sciences of the*

United States of America, *105*(46), 17807–17811.

https://doi.org/10.1073/pnas.0803090105

Ramachandran, P. V., Savini, M., Folick, A. K., Hu, K., Masand, R., Graham, B. H., &

Wang, M. C. (2019). Lysosomal Signaling Promotes Longevity by Adjusting Mitochondrial Activity. *Developmental Cell*, *48*(5), 685-696.e5. https://doi.org/10.1016/j.devcel.2018.12.022

Rambold, A. S., Cohen, S., & Lippincott-Schwartz, J. (2015). Fatty acid trafficking in starved cells: Regulation by lipid droplet lipolysis, autophagy, and mitochondrial fusion dynamics. *Developmental Cell*, *32*(6), 678–692. https://doi.org/10.1016/j.devcel.2015.01.029

Ratnappan, R., Amrit, F.R.G., Chen, S.-W., Gill, H., Ward, J., Yamamoto, K.R., Olse, C.P., & Ghazi, A. (2014). Germline signals deploy NHR-49 to modulate fatty-acid β-oxidation and desaturation in somatic tissues of *C. elegans. PLoS Genetics*, *10*(12), e1004829. https://doi.org/10.1371/journal.pgen.1004829.

Richards, P., Ourabah, S., Montagne, J., Burnol, A. F., Postic, C., & Guilmeau, S. (2017). MondoA/ChREBP: The usual suspects of transcriptional glucose sensing; Implication in pathophysiology. *Metabolism: Clinical and Experimental*, *70*, 133–151. https://doi.org/10.1016/j.metabol.2017.01.033

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Research*, *43*(7), e47. https://doi.org/10.1093/nar/gkv007

Roberts, D. J., Tan-Sah, V. P., Ding, E. Y., Smith, J. M., & Miyamoto, S. (2014). Hexokinase-II Positively Regulates Glucose Starvation-Induced Autophagy through TORC1 Inhibition. *Molecular Cell*, *53*(4), 521–533. https://doi.org/10.1016/j.molcel.2013.12.019

Rual, J. F., Ceron, J., Koreth, J., Hao, T., Nicot, A. S., Hirozane-Kishikawa, T., Vandenhaute, J., Orkin, S. H., Hill, D. E., van den Heuvel, S., & Vidal, M. (2004). Toward improving Caenorhabditis elegans phenome mapping with an ORFeome-based RNAi library. *Genome Research*, *14*(10 B), 2162–2168. https://doi.org/10.1101/gr.2505604

Sans, C. L., Satterwhite, D. J., Stoltzman, C. A., Breen, K. T., & Ayer, D. E. (2006). MondoA-Mlx Heterodimers Are Candidate Sensors of Cellular Energy Status: Mitochondrial Localization and Direct Regulation of Glycolysis. *Molecular and Cellular Biology*, *26*(13), 4863–4871. https://doi.org/10.1128/mcb.00657-05

Schulz, T.J., Zarse, K., Voigt, A., Urban, N., Birringer, M., & Ristow, M. (2007). Glucose Restriction Extends *Caenorhabditis elegans* Life Span by Inducing Mitochondrial Respiration and Increasing Oxidative Stress. *Cell Metabolism*, *6*(4), 280-293.

https://doi.org/10.1016/j.cmet.2007.08.011

Selman, C., Lingard, S., Choudhury, A. I., Batterham, R. L., Claret, M., Clements, M.,

Ramadani, F., Okkenhaug, K., Schuster, E., Blanc, E., Piper, M. D., Al-Qassab, H.,

Speakman, J. R., Carmignac, D., Robinson, I. C. A., Thornton, J. M., Gems, D.,

Partridge, L., & Withers, D. J. (2008). Evidence for lifespan extension and delayed age–

related biomarkers in insulin receptor substrate 1 null mice. *The FASEB Journal*, *22*(3),

807–818. https://doi.org/10.1096/fj.07-9261com

Stoltzman, C. A., Kaadige, M. R., Peterson, C. W., & Ayer, D. E. (2011). MondoA senses

non-glucose sugars: Regulation of thioredoxin-interacting protein (TXNIP) and the

hexose transport curb. *Journal of Biological Chemistry*, *286*(44), 38027–38034.

https://doi.org/10.1074/jbc.M111.275503

Stoltzman, C. A., Peterson, C. W., Breen, K. T., Muoio, D. M., Billin, A. N., & Ayer, D. E.

(2008). Glucose sensing by MondoA:Mlx complexes: A role for hexokinases and direct

regulation of thioredoxin-interacting protein expression. *Proceedings of the National*

Academy of Sciences of the United States of America, *105*(19), 6912–6917.

https://doi.org/10.1073/pnas.0712199105

Sui, D., & Wilson, J. E. (1997). Structural determinants for the intracellular localization of

the isozymes of mammalian hexokinase: Intracellular localization of fusion constructs

incorporating structural elements from the hexokinase isozymes and the green

fluorescent protein. *Archives of Biochemistry and Biophysics*, *345*(1), 111–125.

https://doi.org/10.1006/abbi.1997.0241

Tatenaka, Y., Kato, H., Ishiyama, M., Sasamoto, K., Shiga, M., Nishitoh, H., & Ueno, Y.

(2019). Monitoring Lipid Droplet Dynamics in Living Cells by Using Fluorescent

Probes. *Biochemistry*, *58*(6), 499–503. https://doi.org/10.1021/acs.biochem.8b01071

Tong, X., Zhao, F., Mancuso, A., Gruber, J. J., & Thompson, C. B. (2009). The glucose-

responsive transcription factor ChREBP contributes to glucose-dependent anabolic

synthesis and cell proliferation. *Proceedings of the National Academy of Sciences of the*

United States of America, *106*(51), 21660–21665.

https://doi.org/10.1073/pnas.0911316106

Van Gilst, M. R., Hadjivassiliou, H., & Yamamoto, K. R. (2005). A Caenorhabditis elegans

nutrient response system partially dependent on nuclear receptor NHR-49. *Proceedings*

of the National Academy of Sciences of the United States of America, *102*(38), 13496–

13501. https://doi.org/10.1073/pnas.0506234102

Vrablik, T. L., Petyuk, V. A., Larson, E. M., Smith, R. D., & Watts, J. L. (2015). Lipidomic

and proteomic analysis of Caenorhabditis elegans lipid droplets and identification of ACS-4 as a lipid droplet-associated protein. *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids*, *1851*(10), 1337–1345. https://doi.org/10.1016/j.bbalip.2015.06.004

Wang, M. C., O'Rourke, E. J., & Ruvkun, G. (2008). Fat metabolism links germline stem cells and longevity in C. elegans. *Science*, *322*(5903), 957–960. https://doi.org/10.1126/science.1162011

Wilde, B. R., Ye, Z., Lim, T. Y., & Ayer, D. E. (2019). Cellular acidosis triggers human mondoa transcriptional activity by driving mitochondrial atp production. *ELife*, *8*, 1–25. https://doi.org/10.7554/eLife.40199

Wilson, J. E. (2003). Isozymes of mammalian hexokinase: Structure, subcellular localization and metabolic function. *Journal of Experimental Biology*, *206*(12), 2049–2057. https://doi.org/10.1242/jeb.00241

Wolf, A. J., Reyes, C. N., Liang, W., Becker, C., Shimada, K., Wheeler, M. L., Cho, H. C., Popescu, N. I., Coggeshall, K. M., Arditi, M., & Underhill, D. M. (2016). Hexokinase Is an Innate Immune Receptor for the Detection of Bacterial Peptidoglycan. *Cell*, *166*(3), 624–636. https://doi.org/10.1016/j.cell.2016.05.076

Yamamoto-Imoto, H., Minami, S., Shioda, T., Yamashita, Y., Sakai, S., Maeda, S., Yamamoto, T., Oki, S., Takashima, M., Yamamuro, T., Yanagawa, K., Edahiro, R., Iwatani, M., So, M., Tokumura, A., Abe, T., Imamura, R., Nonomura, N., Okada, Y., … Yoshimori, T. (2022). Age-associated decline of MondoA drives cellular senescence through impaired autophagy and mitochondrial homeostasis. *Cell Reports*, *38*(9), 110444. https://doi.org/10.1016/j.celrep.2022.110444

Yamashita, H., Takenoshita, M., Sakurai, M., Bruick, R. K., Henzel, W. J., Shillinglaw, W., Arnot, D., & Uyeda, K. (2001). A glucose-responsive transcription factor that regulates carbohydrate metabolism in the liver. *Proceedings of the National Academy of Sciences of the United States of America*, *98*(16), 9116–9121. https://doi.org/10.1073/pnas.161284298

Yamawaki, T. M., Berman, J. R., Suchanek-Kavipurapu, M., McCormick, M., Gaglia, M. M., Lee, S. J., & Kenyon, C. (2010). The somatic reproductive tissues of C. elegans promote longevity through steroid hormone signaling. *PLoS Biology*, *8*(8), 45–46. https://doi.org/10.1371/journal.pbio.1000468

Zhang, Y., Lanjuin, A., Chowdhury, S. R., Mistry, M., Silva-García, C. G., Weir, H. J., Chia-Lin, L., Escoubas, C. C., Tabakovic, E., & Mair, W. B. (2019). Neuronal TORC1

1417   modulates longevity via ampk and cell nonautonomous regulation of mitochondrial

1418   dynamics in C. Elegans. *ELife*, *8*, 1–24. https://doi.org/10.7554/eLife.49158

1419