Mechanisms of cilia regeneration in *Xenopus* multiciliated epithelium *in vivo*.

Venkatramanan G. Rao[1], Vignesharavind Subramanianbalachandar[1], Magdalena M. Magaj[1,3,4]
Stefanie Redemann[1,3,4], Saurabh S. Kulkarni[1,2,3] *

1- Department of Cell Biology, University of Virginia, Charlottesville, VA 22903
2- Department of Biology, University of Virginia, Charlottesville, VA 22903
3- Center for Membrane & Cell Physiology, University of Virginia, Charlottesville, VA 22903
4- Department of Molecular Physiology and Biological Physics, University of Virginia, Charlottesville, VA 22903

*- Author for correspondence (sk4xq@virginia.edu)

Keywords

Cilia, transition zone, regeneration, *Xenopus*, B9d1, multiciliated cell, ciliary tip, IFT

SUMMARY

Cilia regeneration is a physiological event, and while studied extensively in unicellular organisms, it remains poorly understood in vertebrates. In this study, using *Xenopus* multiciliated cells (MCCs) as a model, we demonstrate that, unlike unicellular organisms, deciliation removes the transition zone (TZ) and the ciliary axoneme. While MCCs immediately begin the regeneration of the ciliary axoneme, surprisingly, the assembly of TZ is delayed. However, ciliary tip proteins, Sentan and Clamp, localize to regenerating cilia without delay. Using cycloheximide (CHX) to block new protein synthesis, we show that the TZ protein B9d1 is not a component of the cilia precursor pool and requires new transcription/translation, providing insights into the delayed repair of TZ. Moreover, MCCs in CHX treatment assemble fewer (~ 10 vs. ~150 in controls) but near wild-type length (ranging between 60 to 90%) cilia by gradually concentrating ciliogenesis proteins like IFTs at a select few basal bodies. Using mathematical modeling, we show that cilia length compared to cilia number influences the force generated by MCCs more. In summary, our results question the requirement of TZ in motile cilia assembly and provide insights into how cells determine organelle size and number.

INTRODUCTION

Cilia perform critical functions as the signaling hub (primary cilia) and force generator to create extracellular fluid flow (motile cilia) during development. Dysfunction of cilia can manifest into a constellation of defects like polycystic kidney disease[1,2], congenital heart disease[1], primary ciliary dyskinesia[2], retinal degeneration, etc., known as ciliopathies[3]. Nevertheless, how cells assemble a cilium remains a central outstanding question in the field.

Cilium is an extracellular hairlike organelle composed of a microtubule-based cytoskeleton called the ciliary axoneme covered with a membrane[4]. Cilium is anchored to the cell via a basal body (modified centriole) and enclosed by a specialized ciliary membrane[4,5]. The ciliary axoneme consists of the ciliary shaft and ciliary tip. The ciliary shaft is the main body of the axoneme and is made of 9+0 or 9+2 microtubule doublets in sensory or motile cilia, respectively[4]. The structure of the ciliary tip can vary across species and ciliary types; however, it is usually made of microtubule singlets in motile cilia[6,7]. The transition zone (TZ) is a structural junction between the basal body and the axoneme proposed to form a molecular sieve-like barrier[8,9]. It acts like a "ciliary gate" regulating intraciliary protein traffic[8,10,11]. To understand how cells construct such a complex structure, it is necessary to analyze the spatial and temporal order of the assembly of each component in the context of the entire cilium.

The presumption in the field is that cilia assemble bottom-up from the basal body, followed by the TZ and then the ciliary axoneme[12–14]. Ciliogenesis is initiated once the cells exit the cycling phase. The centriole now undergoes maturation to form the basal body to template ciliary growth by recruiting ciliary vesicles to the distal end. The modified centriole at this stage may build a TZ and dock to the cell membrane or dock to the cell membrane, followed by building a TZ[15–17]. Post TZ formation, the axoneme grows by recruitment of proteins at the base of cilia by Intra-flagellar transport proteins (IFT) that ferry ciliary cargo to the tip of the cilia[18,19].

However, in motile cilia and MCCs, more evidence is needed because cilium assembly happens in the context of other intracellular events like basal body migration and cytoskeleton remodeling[20–22], making it challenging to tease out the spatial and temporal sequence of events. Alternatively, one can study ciliogenesis by removing mature cilia and observing the cells reassemble the entire structure without the confounding signals from basal body maturation or cytoskeleton remodeling[20,21,23]. While it is important to remember that regeneration of cilia may not be identical to *de novo* assembly, cilia regeneration studies in *Chlamydomonas reinhardtii*

103 have provided significant insights into ciliogenesis, e.g., cargo transport, the presence of
104 precursor pool, regulation of ciliary gene expression.[18,23,24]
105
The deciliation-regeneration approach offers clear advantages to understanding ciliogenesis.
However, in vertebrates, this has been hindered by a few challenges. First, ciliated cells, including
cells with primary cilia or multiple cilia (MCCs), are often present in tissues that are not accessible
for inducing controlled deciliation and offer technical challenges to studying regeneration by
imaging. The use of cell culture could circumvent this challenge. However, a detailed study in
which primary cilia or MCCs have been deciliated and regenerated is absent in the literature.
These obstacles can be overcome by studying MCCs of the *Xenopus* embryonic epidermis, which
are external and amenable to controlled deciliation, and studying cilia regeneration by imaging
[25,26]. Previous studies have demonstrated that *Xenopus* multiciliated epithelium can be deciliated
using the Ca^{2+} shock in an *in vivo* setting, and these cilia regenerate within hours [25,27,28]. Using
this *in vivo* vertebrate system of deciliation and regeneration of cilia, our study is the first to explore
the fundamental questions of spatial and temporal dynamics of motile cilia assembly.
118
119 **RESULTS AND DISCUSSION**
120
Cilia regeneration on *Xenopus* mucociliary epithelium
The MCCs of the *Xenopus* embryonic epidermis can be consistently deciliated within 10-15
seconds of exposure to a deciliation buffer, as confirmed by the loss of acetylated (Ac.) α-tubulin
signal (Fig. 1A-B). The recovery of Ac. α-tubulin signal was observed within an hour, indicating
the initiation of cilia regeneration, with cilia reaching wild-type length by 6 hours (Fig. 1C).
Compared to biciliate *Chlamydomonas*, where both cilia can attain full length in 90 minutes, the
regeneration of cilia in MCCs of *Xenopus* takes longer, raising questions about the underlying
mechanisms. One possibility is that the delay in cilia regeneration in MCCs may be due to the
requirement to reassemble hundreds of cilia instead of just two in *Chlamydomonas*. Alternatively,
the entire MCC may be replaced by a stem cell population rather than regenerating cilia.
131
The mucociliary epithelium of the *Xenopus* embryonic epidermis is similar in structure and
function to the mammalian airway. Damage to cilia or MCCs in the mammalian airway is
suggested to be repaired by replacing the damaged MCCs using basal stem cells[29,30].
Regeneration of cilia *per se* in mammalian airway MCCs has not yet been reported in the
literature. Therefore, we investigated whether *Xenopus* MCCs regenerate cilia or depend on stem
cell-based replacement of damaged MCCs.
138
We performed live imaging of cilia regeneration to distinguish between the two hypotheses. We
used *Xenopus* stem cell explants (animal caps), which auto-differentiate into an embryonic
multiciliated epidermis. We harvested animal caps from *Xenopus* embryos injected with
membrane RFP (to label cell and ciliary membrane) and grew them on fibronectin-coated slides.
We deciliated the animal caps using the same protocol as the embryos and immediately began
imaging MCCs using confocal microscopy. All MCCs showed a complete absence of cilia,
indicating successful deciliation**.** Approximately 15 minutes post-deciliation, we could observe
membrane RFP signal corresponding to regenerating cilia in MCCs. MCC specification,
differentiation, radial intercalation, basal body amplification, migration, docking, and cilia
formation from basal stem cells in *Xenopus* epithelium can take several hours (Personal
observation). Thus, our results indicate that *Xenopus* multiciliated epithelium regenerates cilia in
the same MCCs and does not undergo stem cell-based renewal of damaged MCCs. (Fig. S1A;
Supplement Video 1).
152
Deciliation affects the F-actin network but does not affect basal body number and polarity.

Next, we investigated whether the deciliation affects only the ciliary axoneme or cilia-associated
structures, such as the apical F-actin network, basal body number, and polarity. Our results show
that deciliation affected the apical F-actin network significantly (calculated as cortical and medial
actin) (Fig. S1B). We used Chibby or Centrin to label basal bodies and Clamp to label rootlets to
assess the effect of deciliation on basal body number and polarity [26]. Our results show that
deciliation does not affect basal body number and polarity. (Fig. S1C, D).
160
161 <u>**The Transition zone (TZ) is removed by deciliation.**</u>
Next, we determined the location where the deciliation treatment severed cilia. Unicellular models
such as *Chlamydomonas* and *Tetrahymena* lose cilia distal to the TZ and below the central pair
(CP) microtubules[10,31–33]. Therefore, we hypothesized that *Xenopus* MCCs adopt a similar
mechanism and would lose the ciliary axonemes distal to the TZ. To test the hypothesis, we
labeled the TZ with an antibody to B9d1, a bona fide TZ protein[11,34–37]. To our surprise, B9d1 was
lost from the deciliated MCCs (Fig. 2A, 0 hr. timepoint), suggesting that TZ was removed along
with the ciliary axonemes. We did Transmission Electron Microscopy (TEM) to visualize the TZ
and validate this unexpected result. TZ can be seen as an electron-dense H-shaped structure
above the basal bodies [10,38–42]. In the deciliated embryos, we could see the basal bodies; however,
the H-shaped structure corresponding to the TZ was lost entirely (Fig. 2B, 0 mins. timepoint).
Taken together, these data demonstrate that, unlike unicellular organisms, deciliation in *Xenopus*
MCCs removes the TZ with the ciliary axoneme.
174
While the TZ is lost with deciliation, the actual excision may happen at two distinct locations, distal
to the basal body (between the basal body and TZ) and distal to the TZ (between TZ and CP),
separately releasing the TZ and the axoneme into the medium[31,32,42]. Alternatively, deciliation may
happen only distal to the basal bodies, in which case the TZ stays with the ciliary
axoneme[10,31,40,42]. A previous proteomic study where authors deciliated *Xenopus* embryonic
multiciliated epithelium found TZ proteins in their ciliary axoneme preparation, suggesting that the
TZ is removed with the ciliary axonemes in *Xenopus* MCCs, supporting the later hypothesis [28].
We examined whether the B9d1 signal was associated with cilia post-deciliation to test the
hypothesis. To that end, we performed an *in-situ* deciliation of embryos using deciliation buffer on
poly-L-lysine coated coverslips. Once removed, cilia stick to the coverslip, stained for B9d1 and
Ac. α-tubulin (Fig. S2A, B). Surprisingly, we did not see any B9d1 signal with cilia (Fig. S2B,
chemical deciliation). Notably, B9d1 is a soluble protein closely associated with the membrane-
associated MKS protein complex in the TZ [9,34,43]. We examined ciliary axonemes with EM to
examine if the detergent in the deciliation buffer is stripping the ciliary membrane and leading to
the loss of the B9d1 signal. As expected, we could not observe any ciliary membrane (Fig. S2D)
supporting our hypothesis that loss of B9d1 is likely the result of loss of ciliary membrane.
Therefore, to understand if the TZ remains associated with the axoneme after deciliation, we
adopted an alternative approach of mechanical deciliation[44] (Fig. S2A). The advantage of this
approach is that we did not use any detergent. We observed that cilia removed from the embryo
consistently showed a B9d1 signal (Fig. S2C). Thus, we demonstrate that the site of deciliation in
Xenopus cilia is distal to the basal body, and the TZ remains attached to the ciliary axoneme.
196
197 <u>**TZ is dispensable for the initiation of cilia assembly during regeneration.**</u>
Removal of the TZ during deciliation gave us a unique opportunity not afforded by any other
vertebrate or invertebrate model to study the temporal relationship between TZ and axoneme
assembly. The TZ is thought to form a molecular sieve-like barrier and serve as a "ciliary gate,"
regulating protein traffic into and out of the cilium[8,9]. Mutations in or knock-down/out of TZ proteins
impair cilia assembly and function[11,34,36,40,45,46]. Further, recent studies have shown that
Intraflagellar transport (IFT) trains assemble at the TZ, suggesting that TZ is essential for ciliary
assembly and is predicted to precede ciliary axoneme assembly[10,22]. However, no direct evidence
to support this hypothesis has been published.

206
Using our unique cilia regeneration model, we directly tested this hypothesis. We examined the
restoration of the B9d1 signal as a marker to track the reassembly of TZ during the first three
hours of cilia regeneration. As expected, both the B9d1 and Ac. α-tubulin signals were completely
lost after deciliation. One hour into regeneration, we were surprised to see the recovery of the Ac.
α-tubulin but no detectable B9d1 signal. The B9d1 signal began to appear after 2 hours of
regeneration and further increased at 3 hours. (Fig. 2A). We quantified the length of the recovered
B9d1 signal in the MCCs. In control embryos (pre-deciliation), the length of the B9d1 signal was
1.33 μM (± 0.33, SEM); this signal increased from no signal at 1 hour to 0.55 μM (± 0.16, SEM)
at 2 hours and 0.73 μM (±0.021, SEM) at 3 hours after deciliation (Fig. 2C). This result was
unexpected because, contrary to the previous speculation, cells appeared to initiate ciliary
axoneme assembly in the absence of B9D1.
218
One limitation of our results is that we have used only one protein, B9d1, to track the loss and
reassembly of TZ. We attempted to localize multiple TZ proteins in *Xenopus,* but our attempts
were unsuccessful as antibodies available to *Xenopus* are particularly limited. To overcome this
shortcoming, we performed electron microscopy to directly analyze the early axoneme and TZ
assembly steps during regeneration. Specifically, we performed the TEM on samples taken at 0,
20, and 90 minutes after deciliation (Fig. 2B). At 20 minutes after deciliation, a small ciliary
axoneme was present; however, the H-shaped structure indicating the presence of mature TZ
was not yet visible. The H-shaped structure of the TZ began to appear 90 minutes post-deciliation.
The TEM data supported the results from our B9d1 immunofluorescence experiments. However,
one of the limitations of the TEM approach is that the sections are very thin (~75nm), and we
could miss the H-shaped structure when looking at a single section. To address this concern, we
performed electron tomography of regenerating cilia at different time points (Pre., 0, 20 mins, 1
hr., 3 hrs., and 6 hrs.) to examine the timing of TZ assembly (Supplementary Videos 2-5).
Consistent with the TEM data, electron tomography results demonstrated that MCCs could initiate
ciliary axoneme assembly without mature TZ. The TZ was absent from the 20-minute tomogram
and assembled between 1 and 3 hours after deciliation; by 6 hours, the TZ looked
indistinguishable from controls. Our results suggest that the cells can incorporate TZ (or
components of TZ) once the axoneme is assembled, suggesting a dynamic nature of TZ proteins.
An earlier study in *Chlamydomonas* also reports the dynamic nature of a TZ protein, CEP290,
incorporated into the pre-assembled axoneme[31]. During the mating process, there was an
exchange of CEP290 from the wild-type cilia to the cilia of the CEP290 deficient mutant during
the mating process [31]. While the previous study showed one TZ protein, CEP290, to be
dynamically assembled in the cilia, and our study shows the incorporation of another TZ protein,
B9D1, we speculate that this could apply to many TZ proteins where they could be dynamically
assembled in cilia.
244
The structure of the ciliary axoneme is different at the proximal end, where the TZ is located, and
the distal end, where the ciliary tip is located. Like the TZ, the ciliary tip comprises of a distinct
protein complex of unknown function[6,7,39,47,48]. Since the TZ showed a delayed regeneration, we
asked if the ciliary tip proteins assembled after the ciliary shaft was built. To test this speculation,
we used mScarlet-Sentan[49] and RFP-Clamp[50] proteins known to localize to the tip of motile cilia.
As reported earlier[49], we observed that mScarlet-Sentan and RFP-Clamp localized to ciliary tips
in control *Xenopus* MCCs (Fig. 3A, S3A). After deciliation, ciliary Sentan and Clamp signals were
lost as expected. However, when regeneration began, we noticed that small stubs of ciliary
axonemes at 1 hr. post-deciliation were decorated with the Sentan and Clamp signals (Fig. 3A, B
and S3A, B). We measured both the Clamp and Sentan signals at different time points during
regeneration. We observed that, like pre-deciliation controls, Sentan and Clamp signals became
stronger at the ciliary tips as regeneration progressed. (Fig. 3B, S3B). Our data suggests that

ciliary tip proteins like Sentan and Clamp are trafficked immediately into regenerating cilia and
gradually begin to concentrate at the tips.
259
TZ protein B9d1 is newly synthesized during cilia regeneration.
We next addressed why there was a delay in assembling TZ. The clue came from an earlier study
where photobleaching of B9D1 and some other TZ proteins (in a pre-existing TZ) did not recover
to pre-bleaching levels[51]. This study indicated that unlike some proteins needed for axoneme
maintenance, e.g., IFTs[18], the TZ proteins such as B9d1 may not be needed for cilia maintenance
and thus are not present in the cytosolic pool. Consistently, we did not observe any B9d1 signal
near the base of the cilia or in the cytoplasm. Therefore, we hypothesized that post-deciliation
MCCs newly synthesize TZ proteins, such as B9D1, resulting in delayed assembly of TZ.
268
To test this hypothesis, we treated deciliated embryos with either vehicle (DMSO) alone or
Cycloheximide (CHX), a protein synthesis inhibitor. We collected embryos every hour for the first
3 hours and quantified the B9d1 signal in the regenerating cilia (Fig. 4). By 3 hours, control MCCs
showed a significant increase in the B9d1 signal. However, treatment with CHX completely
abolished B9d1 recovery. These results suggest that B9d1 is absent in the ciliary precursor pool
and requires new transcription/translation during regeneration.
275
Synthesis of a new pool of ciliary proteins is required to support cilia regeneration.
Our results from the previous experiments raised the question: does deciliation trigger new protein
synthesis in MCCs? From the live imaging experiments, we noted that cilia regeneration begins
within approximately 12-15 minutes after deciliation. *Chlamydomonas* also begin cilia reassembly
within minutes after deciliation. Studies have shown that *Chlamydomonas* uses the precursor pool
of proteins for rapid cilia reassembly; however, this precursor pool is insufficient to assemble full-
length cilia [52,53]. Loss of cilia triggers the synthesis of new mRNA and proteins to support
subsequent cilia assembly [53]. In contrast to *Chlamydomonas*, sea urchins contain excess proteins
that can support full regeneration of cilia up to four times [54]. Therefore, we speculated that
Xenopus MCCs might have a precursor pool of unassembled proteins that allow the cell to begin
cilia reassembly quickly. However, it remains unknown whether *Xenopus* MCCs require new
transcription and translation of some/all proteins to complete cilia regeneration or whether the
unassembled ciliary protein pool is enough to build all cilia.
289
To test this hypothesis, we blocked protein synthesis immediately post-deciliation using
Cycloheximide (CHX); thus, the precursor pool is the only source for building new cilia. (Fig. 5A).
Chlamydomonas has only two cilia, but in *X. tropicalis* MCCs, there are ~150 cilia. Therefore,
CHX treatment could result in either the cell having a sufficient protein pool to regenerate all cilia
comparable to vehicle control or a limited pool that results in MCCs choosing between the number
and length of cilia.
296
We observed that embryos treated with CHX showed shorter cilia than controls during
regeneration (Fig. 5B, C, D, S4A, B). Specifically, there was no significant difference in the ciliary
length in the first hour between control and CHX treatment (vehicle vs. CHX, 2.75±1.04μm vs.
2.24±1.12μm, mean±SD, Fig. 5D, and S4A and B). However, cilia did not grow to the same length
in CHX treatment as vehicle treatment at 3 and 6 hrs. (3 hrs. vehicle vs. CHX, 8.53±1.74 μm, vs.
5.40±1.73 μm, and 6 hrs. vehicle vs. CHX, 14.87±2.38μm vs. 8.8± 2.14μm; mean±SD; Fig. 5D
and Fig. S4A and B). Thus, ciliary assembly is compromised during regeneration with CHX
treatment.
305
Next, we measured the number of cilia per MCC during cilia regeneration. While we could count
the cilia number at 1 hr. in both treatments (1 hr. vehicle 41.61±12.71 vs CHX; 33.78±17.27 cilia/

MCC, Fig. 5C, S4C), it was challenging to count the cilia in vehicle-treated samples after 3hrs.
and 6hrs. due to the long cilia that clumped/entangled during the staining process. However, the
basal body number did not change with the vehicle and CHX treatment (Fig S5A and B). Assuming
all the basal bodies can assemble cilia, we can estimate 125-150 cilia/MCC in vehicle treatment
(Fig S5B). In contrast to the controls, we noted that the cilia number at 1 hr. time point in CHX-
treated MCCs was higher than the 3 hrs. (1 hr. CHX vs. 3 hrs. CHX; 33.78±17.27 vs. 14±6 cilia/
MCC, Fig. 5B, E, S4C). This number was further reduced by the end of 6 hrs. (8 ± 4 cilia/MCC;
mean±SD, Fig. 5C, E, Fig S4C), suggesting that the CHX treatment reduced the cilia number
without affecting the basal body number.
317
We speculated that this dramatic reduction in cilia number and length in CHX treatment could be
due to the blocking of new protein synthesis and the ubiquitin-dependent proteasomal
degradation of the existing ciliary protein pool in the cell. To test the contribution of protein
degradation on the regeneration of cilia length and number, we blocked protein degradation using
the drug MG132 (Fig 5A)[55]. Treating the cells with MG132 with CHX did not significantly increase
the number of cilia at any time point (Fig. 5C, E, and Fig. S4C). However, CHX+MG132 led to a
significant increase in cilia length compared to CHX treatment, by an average of 3μm by 6 hours.
(3 hrs. CHX vs CHX+ MG132; 5.4 ± 1.73μm vs. 6.85±1.53 μm, 6 hrs. CHX vs CHX+ MG132; 8.8
±2.14μm vs. 11.93±2.53 μm; mean ± SD; Fig. 5C, D, S4A, B).
327
Overall, these experiments reveal an interesting observation: with a limited protein pool, MCCs
prefer longer cilia over more cilia. We speculated that MCCs optimize the length and number of
cilia to generate optimal beating force. Therefore, the reason for MCCs to prefer length over
number was unclear. A longer cilium can exert more beating force over a shorter cilium up to a
certain length. However, increasing length beyond a threshold can increase the hydrodynamic
drag [56,57]. However, in MCCs, this calculation is complex because both cilia length and number
play important roles in generating the optimal force necessary for mechanical coupling with
surrounding MCCs [58]. To understand this relationship, we used mathematical modeling to
generate a force matrix for different numbers and lengths of cilia per cell (Supplement Table 1).
Our modeling data shows that the force exerted by the beating cilia increases more for a unit
increase in length than the number. For example, one 12-μm-long cilium will exert higher force
than 12 cilia of 1-μm length (Supplementary Table 1). When applied to the number and lengths
in our data, we show that the theoretical force generated by vehicle-treated MCCs increased
during regeneration. Further, as expected, it was higher than the CHX and CHX+MG132 treated
MCCs (Fig. 6A, B, Supplementary Table 1). Within the CHX-treated samples (CHX and
CHX+MG132), there was an increase in force generation from 1hr. to 3hrs., and 3hrs. to 6 hrs.
despite a gradual decrease in the number of cilia. (Fig. S4C, Fig. 6A, B). Interestingly, the
reduction in cilia number is correlated with an increase in cilia length, supporting the theoretical
calculations that force generation has a quadratic relationship with cilia length compared to a
linear relationship with cilia number. However, one could argue that MCCs have more cilia
building blocks as a precursor pool. As a result, over 6 hrs. of regeneration, there was an increase
in total cilia quantity, which led to increased force generation (cilia quantity = avg. number of cilia
X avg. cilia length). Therefore, we measured total cilia quantity over time in CHX and
CHX+MG132 treatments. Our data shows that total cilia quantity in CHX treatment remained
unchanged (75.66μm at 1 hr., 77.16μm at 3 hrs., and 72.78μm at 6 hrs.), suggesting that the
ciliary proteins were likely redistributed from shortened cilia to support fewer but longer cilia. This
phenomenon was also observed in *Chlamydomonas* when one of the two flagella is removed in
the presence of CHX, and the intact flagellum is shortened to support the growth of the amputated
flagellum [18,59]. On the other hand, there was a slight increase in total cilia quantity over time in
CHX+MG132 treatment (89.56μm at 1 hr., 108.41μm at 3 hrs., and 122.64μm at 6 hrs.). However,
this increase was minimal (21.04% from 1 hr. to 3hrs. and 13.12% from 3 hrs. to 6 hrs.) compared

to the calculated force generation (214.7% from 1 hr. to 3 hrs. and 163.06% from 3 hrs. to 6 hrs.),
which was more proportionate to the change in cilia length (241.7% from 1 hr. to 3 hrs. and
174.41% from 3 hrs. to 6 hrs.). Therefore, we propose that cilia length is a major determinant of
force generation. While our model is based on the previously published literature [56,57,58], it is still
a simplified model and does not consider other factors, such as the coordination of beating cilia,
the mucus viscosity, and the spacing/distribution of ciliary arrays on the cell, which could affect
the effective force transmission.
366
Our model would also benefit from a direct test to measure bead movement as a readout of force
generation by these MCCs. However, there are technical challenges involved in accurately
measuring the bead movement over the surface of single MCCs with different cilia lengths and
numbers. Even if we overcome this challenge, we do not understand when the motility machinery
is fully incorporated into a regenerating cilium. Moreover, in the CHX treatment, whether the
machinery proteins are present as precursors or require new transcription/translation similar to
the TZ protein B9D1. Nonetheless, mathematical modeling coupled with the experimental data
on cilia length and number sheds light on an important cell biology question of how cells balance
the number and size of organelles to optimize functional output.
376
MCCs redistribute proteins among basal bodies to regenerate fewer but longer cilia.
Given the total cilia quantity in CHX treatment remains unchanged from 1 hr. to 6 hrs., and MCCs
cannot synthesize new proteins, we hypothesized that existing proteins essential for ciliogenesis
accumulate at a few basal bodies to support fewer longer cilia. A similar phenomenon was
observed previously in *Chlamydomonas*; when *Chlamydomonas* loses one of its two flagella, it
resorbs the longer flagellum to redistribute the material to the regenerating flagella to support its
assembly until both flagella reach the same length [18,52]. However, unlike *Chlamydomonas* cells,
Xenopus MCCs have hundreds of cilia, and basal bodies are not physically connected, raising
the possibility of distinct mechanisms.
386
IFT proteins are necessary to assemble the axoneme by transporting the axonemal proteins to
the tip of the growing structure [19,27,60]. Ciliogenesis and cilia regeneration occur by accumulating
several ciliary proteins, including IFTs at the base of cilia in *Chlamydomonas* and *Xenopus* [10,18,27].
Therefore, we examined the pool of three IFT proteins (IFT20, IFT80, and IFT43) at the base of
regenerating cilia as a proxy for essential ciliogenesis proteins required for cilia regeneration in
the vehicle and CHX-treated embryos (Fig. 7, Fig. S6, S7). In pre-deciliation-and 0 hr. deciliated
MCCs, all basal bodies were decorated with IFT proteins (IFT20, IFT80, and IFT43) signal,
demonstrating that IFTs were similarly distributed among all basal bodies. In vehicle-treated
embryos, all basal bodies remained equally decorated with IFT signal throughout regeneration as
expected (Fig. 7A, S6A, S7A). In contrast, in the CHX-treated cells, we observed that IFT proteins
gradually enriched at a subset of the basal bodies as regeneration progressed. Interestingly, the
basal bodies that were saturated with the IFT signal were also ciliated at all time points (Fig. 7A",
B, S6A", B, S7A", B), suggesting that in the absence of new protein synthesis, IFT proteins enrich
a few basal bodies and support the assembly of long cilia. It must be noted that the non-ciliated
basal bodies at 6 hrs. possess some IFT-GFP signal (IFT20, IFT80, and IFT43); we speculate
this could be due to multiple reasons. One, some minimal amount of IFT is always associated
with basal bodies. Second, these IFT molecules are in the process of getting redistributed. Note
that 6 hours is sufficient to regenerate full-length cilia in vehicle-treated MCCs, but we do not
know if this dynamic applies to CHX treatment. Third, this results from ectopic overexpression,
where an excess of proteins could be associated with non-ciliated basal bodies.
407
While *Chlamydomonas* also redistributes the proteins between cilia to coordinate their
regeneration, significant differences between the model systems raise fascinating questions
about how cells sense information and make decisions that may require quantitative data [61,62].

For example, *Chlamydomonas* has only two basal bodies that are proximally placed and
physically connected. Thus, the basal bodies share the protein pool required for assembly. On
the other hand, *Xenopus* MCCs nucleate hundreds of basal bodies. While they are connected via
the cytoskeletal network [20,26,63], they are not physically connected like *Chlamydomonas*.
Moreover, each cilium has its pool of proteins at the basal body to support the assembly. Given
hundreds of basal bodies in *Xenopus* MCCs, how does inter-basal body transport operate? One
possibility is that proteins are transported by diffusion to capture mechanism from the cytosol to
cilia [27,64]. However, another possibility is the direct transport between basal bodies using a
cytoskeleton (microtubule) network[65,66]. Hibbard et al. tested this possibility by depolymerizing
microtubules with nocodazole and cold shock and showed that IFT recruitment at basal bodies
could be independent of cortical microtubules [27]. However, this experiment was not carried out
when the cell was experiencing a shortage of proteins, as is the case for CHX treatment. Our
results also show that cells ensure that at least a few basal bodies can assemble near wild-type
length cilia, raising questions about how these basal bodies are selected. One possibility is that
ciliary axonemes must reach a minimum length to be selected to grow longer. This is not an issue
in wild-type cells where proteins are abundant; however, in a protein-deficient state, only cilia that
reached the minimum length continue assembling, whereas others resorb because of competition
for the available pool of proteins.

429
In conclusion, our study reveals new and significant findings and raises exciting questions about
the mechanisms of cilia regeneration in vertebrates. We show that the location for severing the
ciliary axoneme in MCCs was distal to the basal body, which differs from unicellular organisms
like *Chlamydomonas*. Most interestingly, MCCs initiate the assembly of the ciliary tip and ciliary
shaft before the assembly of the TZ, after which both grow concurrently (Fig. 8). One possibility
is that there is a lack of a cytosolic pool for one or more proteins that are rate-limiting for TZ
assembly. These findings raise intriguing questions about the spatial-temporal sequence of cilia
assembly and the importance of TZ in ciliary assembly. For example, is the TZ dispensable for
motile ciliogenesis? When are other components needed for ciliary stability (e.g., microtubule
inner proteins) transported into the cilium? How do MCCs decide the spatial location of the central
pair microtubules without TZ? Moreover, the study also highlights the importance of new
transcription/translation in completing cilia regeneration. Without protein synthesis during cilia
regeneration, MCCs produce few longer-length cilia over multiple short-length cilia. Our
mathematical model suggests that this optimization helps MCCs exert more force per cell. This
study also provides important insights into the cellular decision-making process that regulates the
size and number of organelles to maximize their function.

446
447 **MATERIALS AND METHODS**

448
Animal husbandry and *in-vitro* fertilization
Frogs (*Xenopus tropicalis*) were bred and housed in a vivarium using protocols approved by the
University of Virginia Institutional Animal Care and Use Committee (IACUC). Embryos needed for
experiments were generated using *in-vitro* fertilization as described before [26,67]. Briefly, the testes
from male frogs were crushed in 1x MBS (pH-7.4) with 0.2% BSA and added to eggs obtained
from the female frogs. After 3 minutes of incubation, freshly made 0.1x MBS (pH-7.8) was added,
and the eggs were incubated for 10 more minutes till contraction of the animal pole of the eggs
was visible. The jelly coat was removed using 3% cysteine in 1/9 x MR solution (pH 7.8-8.0) for 6
minutes. The embryos were either microinjected with DNA/RNA and raised at 25°C or 28°C till
they reached appropriate stages for fixation and staining or mucociliary organoid (animal cap)
dissection. Embryos were staged as described previously [68].

460
Animal cap / mucociliary organoid dissection

Embryos at stage 10 were used for animal cap dissection. The animal pole from the stage 10
embryos was dissected using a fine hair tool in a pool of Danilchik's for Amy (DFA) medium. The
dissected tissue was trimmed to remove the equatorial cells from the sides. This tissue was then
laid on a coverslip or glass slide coated with fibronectin (Sigma Cat No. F1141) immersed in DFA
solution. The tissue was allowed to grow and spread on the fibronectin-coated slides/coverslips
overnight till they reached stage 28.
468
DNA, RNA, and microinjections
Plasmids used in this study are listed in the key resource table. For generating Chibby-BFP, the
full-length Chibby gene was cloned to the N-terminus of mtagBFP2 in the pCS2+ vector using
Gibson assembly. RNA used in this study was generated by linearizing the plasmids and *in-vitro*
transcribed using the mMessage and mMachine SP6 transcription kit (Thermo Fisher Scientific).
Microinjections were done using glass needles using the Pico-liter microinjection system (Warner
instruments). All the microinjections were done at the 1-cell stage except for Figure 2, where the
membrane RFP was injected in 1 of the four blastomeres at the four-cell stage.
477
Chemical deciliation of embryos and mucociliary organoids and drug treatments
Embryos, after they reach stage 28, were used for deciliation experiments. A basket was designed
for deciliation by cutting a 50 ml centrifuge tube. One of the open ends of the tube was heat-
sealed with a muslin cloth. A 6-well plate was filled with deciliation solution (75 mM CaCl$_2$, 0.02%
NP40 in 1/3 MR solution). The rest of the five wells were filled with 1/9x MR. Anesthetized
embryos were transferred to this container and dipped in a deciliation solution (75 mM CaCl$_2$,
0.02% NP40 alternative in 1/3 MR solution) in a 6-well plate for 10-15 seconds. Post-deciliation,
embryos with the container were dipped/washed with 1/9x MR in the rest of the wells of the 6-well
plate. The embryos were then transferred to a petri dish with 1/9x MR with gentamicin and
incubated at 28°C. Embryos were collected during various stages of cilia regeneration, as
mentioned in the text and figure legends. For animal caps, the glass slide/coverslip was removed
from the DFA medium and dipped in the deciliation solution for 15 seconds. Post-deciliation, the
glass side/coverslip was dipped in 1/9x MR for 30 seconds and transferred to DFA until the cilia
were regenerated to the appropriate stage. Control or deciliated embryos were transferred to 1/9
X MR solution containing 10 mg/ml of cycloheximide (CHX) or DMSO for cycloheximide
experiments. MG132 was used to prevent ubiquitin-mediated protein degradation; we used 10
μM of MG132 dissolved in DMSO for treatment post-regeneration.
495
Mechanical deciliation
Poly-L-lysine-coated cover glass was used for the mechanical deciliation of embryos. 25-30
embryos at NF stage 28-30 were transferred to the poly-L-lysine-coated cover glass, and any
residual 1/9 x MR was removed. A mechanical pressure was applied by a glass slide, placing it
on top of the embryos for a minute. After a minute, the embryos were discarded, and the cilia
stuck to the coverglass were fixed in chilled methanol for 10 minutes. And rehydrated using 1x
PBS, followed by blocking and staining with anti-Ac. α tubulin antibody and anti-B9D1 antibody.
503
Immunofluorescence staining and imaging.
The *X. tropicalis* embryos used for the study were fixed post-stage 28 with appropriate fixative
agents. For most experiments, 4% paraformaldehyde (PFA) was used as a fixative, and 100%
chilled methanol was used to fix the embryos for anti-B9d1 staining. After fixation, the embryos
were washed three times with PBST (1x PBS with 0.2% Triton X-100) for 10 mins each and then
incubated in a blocking solution (3% BSA in PBST) for 1 hour. Appropriate antibody was added
to the embryos, incubated for 1 hour at room temperature, and rewashed three times for 10 mins
each with PBST. A conjugated secondary antibody was used to stain embryos for 1 hour. The
embryos were washed thrice with PBST, stained with phalloidin in PBST for 45 minutes, and
washed once post-staining. The embryos were mounted and imaged. Confocal imaging was

performed using the Leica DMi8 SP8 microscope with a 40x oil immersion objective (1.3 NA). Images were captured at 1x or 4x zoom and adjusted (brightness and contrast), analyzed, cropped in Fiji, and assembled in Adobe Illustrator software.

Sample preparation for electron tomography

Xenopus stage 26-28 embryos Pre and post deciliation were fixed in 2.5% glutaraldehyde and 2% paraformaldehyde in 0.1M sodium cacodylate buffer pH-7.4 for 1 hour, rinsed in the buffer, then post-fixed in 1% osmium tetroxide. Subsequently, embryos were rinsed in buffer and stained in 2% aqueous uranyl acetate for another hour. Embryos were rinsed and dehydrated in an ethanol series (15min 30% ethanol, 15min 50% ethanol, 15min 70% ethanol, 15min 90% ethanol, 3x 30min 100% ethanol). Dehydrated embryos were infiltrated with EPON Araldite (Electron Microscopy Sciences) and polymerized overnight at 60 C. Serial semi-thick sections (200 nm) were cut using an Ultracut UCT Microtome (Leica Microsystems, Vienna, Austria). Sections were collected on piliform-coated copper slot grids and poststained with 2% uranyl acetate in 70% methanol, followed by Reynold's lead citrate57.

Data acquisition by electron tomography.

Colloidal gold particles (15 nm; Sigma-Aldrich) were attached to both sides of semi-thick sections collected on copper slot grids to serve as fiducial markers for subsequent image alignment. For dual-axis electron tomography, a series of tilted views were recorded using a TECNAI F20 transmission electron microscope (FEI Company, Eindhoven, The Netherlands) operated at 200 kV. Images (4K x 4K) were captured every 1h over a ±60 range and a pixel size of 2.3nm using a TVIPS CCD camera. We used the IMOD software package (http://bio3d.colourado.edu/imod) for image processing, which contains all the programs needed to calculate electron tomograms. The tilted views were aligned using the positions of the colloidal gold particles as fiducial markers. Tomograms were computed for each tilt axis using the R-weighted back-projection algorithm. Reconstructed tomograms were flattened.

Image analysis, quantification, and statistics

All the experiments were repeated three times. All the measurements and analyses were performed on at least three embryos. Sample size, indicated by "n" values, is included in the figure legends. For length measurement (cilia length, ciliary tip length, and TZ length), at least three cells per embryo were chosen randomly; however, it was ensured that only those cilia/signals that were distinct and not overlapping (or tangled) with neighboring ones were measured. For statistical analysis, Prism ver. 10 was used. The type of analysis, p-values, and significance are included in the figure legends. The F-actin intensity measurements were done using Fiji. The cortical actin refers to the actin network along the MCC boundary, while the medial actin refers to the F-actin network at the apical surface of the MCC. The intensity measurements were performed using a line or polygon tool in Fiji, as described in an earlier study[20]. For orientation calculations, images were taken after aligning the embryos horizontally from anterior on the left to posterior on the right and stained for Clamp-GFP (rootlet) and Centrin-RFP (basal body). The Clamp-GFP orientations were calculated in MATLAB R2021a using image processing and circular statistics toolboxes. The cilia orientations were drawn by hand from rootlet to basal body using the 'drawline' command. The angles were calculated with respect to the global X-axis in the cartesian coordinate system (first point starting at 0,0) using the command 'atan2d', which gives the angles from 0^o to $+180^o$ in a counterclockwise direction (quadrants I and II) and 0^o to -180^o in a clockwise direction (Quadrants IV and III). 26 and 21 multiciliated cells were analyzed for control and deciliated embryos, respectively, with an average of 112 cilia orientations (~60-200 cilia per cell) in each cell for each condition. For the IFT-GFP (IFT20, IFT80, and IFT43) fluorescence intensity calculations, we adopted an unbiased approach to measure the IFT intensities. Firstly, the three channels were separated using Fiji. The cilia and centrin channels were merged, and a circular ROI between 0.646-0.680 μm^2 was drawn around the ciliated and non-ciliated centrin-RFP puncta

566 (4-5 basal bodies from each category). The mean gray value of the IFT-GFP channel was
567 calculated, and the values were normalized from 0-1. The normalization was done using the
568 following formula:
569
570 Normalized value = (Mean gray value – Min value in the data set) / (Max value in the dataset- Min
571 value in the data set)
572
573 **Theoretical estimate of force applied by a single Multiciliated Cell (MCC):**
574 Boselli et al. derived a simple force model for a cilium from resistive force theory [58]using a
575 simplified assumption of cilium as a rigid rod rotating around the base, making an arc of varying
576 lengths from 0 to length of the cilium (l) during each cycle. By ignoring force contribution during
577 the recovery stroke, the effective force contribution from a cilium during a cycle is simplified as
578

$$f = \frac{\zeta \times l \times V_c}{12}$$

580
581 where l is the length of the cilium,
582   V_c is the velocity approximated as ~4×ν×l where ν is the beat frequency (assumed as 25
583 Hz and l is the length of the cilium) and

$$\zeta = \frac{4\pi\mu}{\ln{(\sqrt{e\varepsilon})}}$$

is the transverse drag coefficient,
where μ is the viscosity of the liquid (assumed as 0.001 Pa. s for water)
e is the natural log (e=2.718281828459)
ε is the cilium aspect ratio (length/thickness), the thickness of the cilium is assumed as
0.2 μm.
Force contribution from the entire MCC is approximated as F = f × N × C_o
where f is the force contribution from a single cilium where N is the total number of cilia in the cell,
and C_o is the coupling coefficient, which is defined as 0.55 based on
593
Boselli et al.'s force calculations[58] for far-field estimates show that ~50-60 % of cilia don't
contribute to a cell's final average lateral force component due to entanglement and phase shifts[58].
Hence, a value of C_o = 0.55 is used for the calculations, reflecting close to the actual force
contribution from each cell. Using the formula described force contribution from a single MCC was
simulated for varying lengths (1-15 μm assuming all cilia in the cell have the same length) and
number of cilia (1-150) and visualized as surface plots.
600
601 **Key resource table**

Antibodies			
Name	**Marker**	**Make and Cat. No.**	**Remark**
Mouse Anti-Acetylated α-tubulin (6-11B1)	Ciliary axoneme	Sigma	PFA fix, 1:1000
Rabbit mAb Acetyl-α-Tubulin (Lys40) (D20G3)	Ciliary axoneme	Cell Signaling technology (#5335)	Methanol Fix, 1: 500
Rabbit mAb Acetyl-α-Tubulin (Lys40) (D20G3) (Alexa Fluor® 647 Conjugate)	Ciliary axoneme	Cell Signaling Technology (#81502)	PFA fix, 1: 25
Mouse pAb Anti-B9d1	Transition Zone	Sigma (SAB1409114)	Methanol Fix, 1:50

Chicken anti Mouse conjugated to Alexa fluor 488	NA	Invitrogen	1:500
Anti-Rabbit conjugated to Alexa fluor 647	NA	Jackson	1:500
Phalloidin conjugated to Alexa fluor 488	Actin	Invitrogen	1:50
Phalloidin conjugated to Alexa fluor 647	Actin	Invitrogen	1:50

DNA/RNA			
Name	**Marker**	**Conc. injected**	**Source**
Chibby-tagBFP2	Basal body	100 pg (RNA)	This study
mScarlet-Sentan	Cilia tip	200 pg(RNA)	This study
GFP-Clamp	Rootlets	200 pg (RNA)	[69]
Centrin-RFP	Basal body	100 pg (RNA)	[69]
MemRFP	Cell and Ciliary membrane	100 pg(RNA)	[50]
IFT43-GFP	Cilia base/Cilia	150pg (RNA)	[27]
IFT20-GFP	Cilia base/Cilia	150pg (RNA)	[27]
IFT80-GFP	Cilia base/Cilia	150pg (RNA)	[27]

Drugs/Chemicals			
Name	**Solvent**	**Make and Cat No.**	**Remark**
Cycloheximide	DMSO	Sigma	10μg/ml
MG132	DMSO	Sigma	10 μM

602
603
604

ACKNOWLEDGMENTS
The authors would like to express their gratitude to Dr. Bob Bloodgood and the members of the
Kulkarni Lab for their invaluable feedback and thorough review of this manuscript. Appreciation
is also extended to Dr. Mustafa Khokha for the initial discussions and intellectual guidance for the
project. Acknowledgment is also due to the Molecular Electron Microscopy Core and the
Advanced Microscopy Facility at the University of Virginia. Lastly, we are grateful for the NIH
Pathway to Independence, K99/R00 (K99HL133606 and R00HL133606) grant awarded to
Saurabh Kulkarni.

FIGURE LEGENDS

Figure 1: Deciliation and regeneration of cilia from the mucociliary epithelium of *Xenopus*
***tropicalis*.**
A) Cartoon depiction of the deciliation and regeneration protocol of *X. tropicalis* embryos.
B) Pre-deciliated embryos (Pre.) and embryos during regeneration (0 hr.,1 hr., 3 hrs., 6 hrs.)
stained for cilia (Ac. α-tubulin in magenta) and F-actin (phalloidin in green). For each time point,
the top panel shows the entire embryo, the lower left panel is a zoomed image of the multiciliated

epithelium, and the lower right panel is an SEM image of the embryonic epithelium. Black arrows in the SEM image point to MCCs.

C) Cilia length pre- and during regeneration. The values in parenthesis indicate the number of cilia measured from three trials using 10 embryos. **** - p<0.0001, *** - p<0.001, ns - not significant, ANOVA analysis was followed by Tukey's multiple comparison test.

Figure 2: Loss and regeneration of transition zone

A) MCCs labeled with B9d1 antibody (TZ, green) and Ac. α-tubulin (cilia, magenta) in embryos pre- and during cilia regeneration (0 hr.,1 hr., 3 hrs.). MCCs are depicted by a white dashed outline.

B) Representative TEM images of cilia pre-, and during the regeneration (0 mins, 20 mins, and 90 mins). The TZ can be seen as an 'H' shaped structure between the axoneme and basal body (indicated by a black arrow). Post-deciliation samples show a complete loss of TZ. TZ is absent after 20 mins of cilia regeneration. Premature electron-dense TZ structure can be seen after 90 minutes of cilia regeneration.

C) The length of the B9d1 signal was measured pre- and during cilia regeneration (at 2 hrs. and 3 hrs.) The B9d1 signal was absent at earlier time points (0 hr. and 1 hr.). The values in parenthesis indicate the number of cilia measured from three trials using 10 embryos. **** - p<0.0001; ns - not significant, One-way ANOVA analysis followed by Tukey's multiple comparison test.

Figure 3: Ciliary tip protein Sentan is localized to the ciliary axoneme during the early stages of regeneration.

A) MCCs are labeled with mScarlet-Sentan decorating ciliary tips (magenta) and membrane-GFP (cilia, green) at various stages of cilia regeneration. Sentan can be seen at the tips of cilia in pre-deciliated samples. After 1 hr. post-deciliation, Sentan spans the entire length of ciliary axonemes. At 2 hrs., Sentan starts accumulating at the ciliary tips. At 3 and 6 hr. Sentan signal appears similar to pre-deciliated samples.

B) The Sentan signal length (left panel) and cilia length (right panel) were measured and compared among different time points. The values in parenthesis indicate the number of cilia measured from three trials using 9 embryos. **** - p<0.0001; *** - p<0.0005; * - p<0.05 ns - not significant, Kruskal-Wallis test, followed by Dunn's test.

Figure 4: TZ protein B9d1 requires new protein synthesis during regeneration.

A) Embryos treated with DMSO alone (Vehicle) or cycloheximide (CHX) in DMSO were stained for B9d1 (green, TZ) and Ac. α-Tubulin (magenta, cilia) that were collected pre-deciliation and during cilia regeneration (0 hr., 1 hr., 2 hrs., 3 hrs.). The cell outline is marked with a white dashed line, and the dashed square box in the B9d1 channel is zoomed and depicted in the inset. While cilia can regenerate in both treatments (controls and CHX), the B9D1 signal is recovered only in controls and remains completely absent in the CHX-treated samples at 2 hrs. And 3 hrs. timepoints. Ax: Axoneme, TZ: transition zone, BB: Basal body

B) The percent of MCCs positive for the B9d1 signal (green) and the absent B9d1 signal (magenta). Numbers in parentheses indicate the number of MCCs counted from three trials using 9 embryos. *** - p<0.001, chi-square test using VassarStats to test the significance.

Figure 5: Complete cilia regeneration requires new protein synthesis.

A) Deciliated embryos were treated with DMSO (Vehicle), cycloheximide (CHX), and CHX with MG132 in DMSO and collected during different time points of cilia regeneration (0 hr., 1hr., 3 hrs., 6 hrs.).

B) Pre-deciliated and 0 hr. deciliated MCCs stained for with Ac. α-tubulin antibody (cilia, magenta) and Phalloidin (F-actin, green).

C) MCCs treated with vehicle, CHX, and CHX+MG132 at 1hr., 3hrs., and 6 hrs.
D) Graph showing the cilia lengths at different times pre-deciliation and during cilia regeneration
in Vehicle, CHX, and CHX+MG132. The number in parenthesis indicates the number of cilia
counted from three trials using 9-10 embryos. ****- $p < 0.0001$, ns = not significant, Kruskal Wallis
test, followed by Dunn's multiple comparison test.
E) Graph showing the effect of CHX and CHX with MG132 treatments on cilia number per MCC
during regeneration. The number in parenthesis indicates the number of MCCs counted from
three trials using 9-10 embryos. ****- $p < 0.0001$, **- $p < 0.01$, ns = not significant, Kruskal Wallis
test, followed by Dunn's multiple comparison test.
680

Figure 6: Mathematical model describing the contribution of cilia length and number to
force generation in MCCs.
A) The visualized 3-D plot shows a linear relationship between the number of cilia and the force
generated by MCC, while a quadratic relationship exists between the length of cilia and the force
generated by MCC. The individual force values for different cilia lengths and numbers are given
in Supplementary Table 1.
B) The cartoon depiction of the data shows that, a decrease in cilia number and an increase in
cilia length were correlated with increased force generation (calculated from the model) over time
in both CHX and CHX+MG132 treatments.
690

Figure 7: Distribution of ciliary precursor pool (IFT43-GFP) in MCCs.
A) Embryos injected with IFT43-GFP (green) and Centrin-RFP (orange hot, basal bodies) were
deciliated at stage 28 (0 hr.) and were split into two experiments (DMSO and CHX). A', A'') The
embryos in both sets were allowed to regenerate cilia for 6 hours. After fixation, the embryos were
stained for cilia(magenta). Note that the control MCCs at all time points (1 hr., 3 hrs., and 6 hrs.)
have multiple cilia regenerating, and the IFT43-GFP intensity is uniform at every basal body. In
contrast, the number of regenerating cilia decreases with time in CHX-treated samples, and the
IFT43-GFP is enriched at a few ciliated basal bodies.
B) The intensity of IFT43-GFP associated with ciliated (green) vs. non-ciliated (gray) basal bodies
in the same MCC (in CHX-treated samples) at different time points during cilia regeneration. A
total of 8-10 basal bodies per MCC (4-5 ciliated and 4-5 non-ciliated) and 5 MCCs were chosen,
and the mean gray value was estimated and normalized to the maximum and the minimum values
in the set. The value in parenthesis indicates the number of basal bodies analyzed (with and
without IFT43-GFP) from 9 embryos from three independent trials. Note the significant difference
in the IFT43-GFP signal intensity at ciliated vs. non-ciliated basal bodies at all time points. **** -
$p < 0.0001$, Kruskal Wallis test, followed by Dunn's multiple comparison test.
707

Figure 8: Model for cilia regeneration in *Xenopus tropicalis* embryos
Cilia regeneration in *Xenopus tropicalis* embryonic MCCs is multi-staged and is depicted in the
cartoon. Post-deciliation, the TZ is lost with cilia. During stage 1 of regeneration (0-2 hours), the
cilia grow without the TZ. Also, the cilia tip proteins Sentan and Clamp are immediately trafficked
into the ciliary axoneme. By stage 2 (2-4 hours), a partial transition zone appears, and ciliary tip
proteins begin to accumulate at the distal end of the axoneme. The cilia look like pre-deciliated
samples by the end of regeneration.
715
716

717   **SUPPLEMENTARY FIGURE LEGENDS**
718

719   **Figure S1: Deciliation affects apical F-actin, but the basal body number is unaffected.**
720   A) Montage of regenerating cilia (mem-RFP) in the animal caps. The mem-RFP signal specks
721   can be seen emerging from the cell's surface by ~12 mins, eventually growing into beating cilia
722   (see Supplement video 1).

B) Pre- and post-deciliation MCCs that are stained for cilia(magenta) and F-actin(green) are depicted. Cortical and medial F-actin intensity significantly differs in Pre and 0-hr. deciliated samples. The values in parenthesis indicate the number of MCCs measured from three trials from 9-10 embryos. **** - p<0.0001, Mann-Whitney test.

C) The number of basal bodies labeled with Chibby-GFP (cyan) is unaffected after deciliation. The values in parenthesis indicate the number of MCCs measured from three trials from 9-10 embryos. ns = not significant, Mann-Whitney test.

D) Deciliation does not affect basal body polarity. Basal body polarity was determined by measuring the orientation of rootlets labeled with Clamp-GFP (green) with their respective basal body labeled with Centrin-RFP (magenta) and represented in the rose plot. The directionality was measured in 27 pre-deciliated cells and 22 MCCs post-deciliation from three trials from 9-10 embryos in each category.

Figure S2: Transition zone is removed with cilia during deciliation.
A) Schematic of the chemical and mechanical deciliation methods.

B) Cilia with chemical deciliation lose the B9d1 signal, possibly due to the loss of membrane with detergent in the deciliation buffer, whereas C) the B9d1 signal is maintained with mechanical deciliation.

D) Electron micrographs of cilia from the chemical deciliation method lack the ciliary membrane and show splaying of axonemal microtubules.

Figure S3: Clamp is localized to the ciliary axoneme during the early stages of regeneration.
B) MCCs are labeled with RFP-Clamp (ciliary tip and base, magenta) and memGFP (cilia, green) at various stages of cilia regeneration. After 1 hr. post-deciliation, Clamp signal can be seen in the ciliary axonemes. At 2 hrs., Clamp signal starts accumulating at the ciliary tips. At 3 and 6 hr., Clamp signal appears more like pre-deciliated samples.

B) Clamp signal length (left panel) and cilia length (right panel) were measured and compared among different time points. The values in parenthesis indicate the number of cilia measured from three trials using 9-10 embryos. **** - p<0.0001; * - p< 0.05; ns - not significant, Kruskal-Wallis test, followed by Dunn's test.

Figure S4: Quantification of cilia length and number on CHX and CHX+MG132 treatment.
A) Graph depicting the length of regenerating cilia in the vehicle, CHX, and CHX+MG132 treatments. The cilia lengths at every time point, treatment, and the 'n' values are shown in Fig 5D.

B) The Table shows the average cilia length in 3 treatments and 4 time points. Values are represented as mean±SD. The "n" values are shown in Fig. 5D.

C) Table showing variation in cilia number in 3 treatments and 3 time points. Values are represented as mean±SD. The n values and statistical differences are shown in Fig. 5E. ND— Not determined.

Figure S5: Blocking new protein synthesis does not affect the basal body during regeneration.
A) Stage 28 embryos labeled Chibby-BFP (green, basal bodies) were transferred to DMSO containing 1/9 MR (vehicle) or cycloheximide (CHX) containing 1/9 MR post-deciliation. The embryos were allowed to regenerate and collected at 3 and 6 hrs. Embryos were fixed and stained for cilia (magenta) using Ac. α-Tubulin.

B) The Chibby puncta were counted in all samples (deciliation and regeneration), and the quantification of the basal body number is shown in the graph. The values in parenthesis indicate the number of cells counted from 8 embryos from three trials. ns - not significant, Kruskal-Wallis test, followed by Dunn's test.

Figure S6: Distribution of ciliary precursor pool (IFT80-GFP) in MCCs.
A) Embryos injected with IFT80-GFP (green) and Centrin-RFP (orange hot, basal bodies) were deciliated at stage 28 (0 hr.) and were split into two experiments (DMSO and CHX). A', A'') The embryos in both sets regenerated cilia for 6 hours. After fixation, the embryos were stained for cilia(magenta). Note that the control MCCs at all time points (1 hr., 3 hrs., and 6 hrs.) have multiple cilia regenerating, and the IFT80-GFP intensity is uniform at every basal body. In contrast, the number of regenerating cilia decreases with time in CHX-treated samples, and the IFT80-GFP is enriched at a few ciliated basal bodies.
B) The intensity of IFT80-GFP associated with ciliated (green) vs. non-ciliated(gray) basal bodies in the same MCC (in CHX-treated samples) at different time points during cilia regeneration. A total of 8-10 basal bodies per MCC (4-5 ciliated and 4-5 non-ciliated) and 5 MCCs were chosen, and the mean gray value was estimated and normalized to the maximum and the minimum values in the set. The value in parenthesis indicates the number of basal bodies analyzed (with and without IFT80-GFP) from 9 embryos from three independent trials. Note the significant difference in the IFT80-GFP signal intensity at ciliated vs. non-ciliated basal bodies at all time points. **** - $p<0.0001$, Kruskal Wallis test, followed by Dunn's multiple comparison test.

Figure S7: Distribution of ciliary precursor pool (IFT20-GFP) in MCCs.
A) Embryos injected with IFT20-GFP (green) and Centrin-RFP (orange hot, basal bodies) were deciliated at stage 28 (0 hr.) and were split into two experiments (DMSO and CHX). A', A'') The embryos in both sets regenerated cilia for 6 hours. After fixation, the embryos were stained for cilia(magenta). Note that the control MCCs at all time points (1 hr., 3 hrs., and 6 hrs.) have multiple cilia regenerating, and the IFT20-GFP intensity is uniform at every basal body. In contrast, the number of regenerating cilia decreases with time in CHX-treated samples, and the IFT20-GFP is enriched at a few ciliated basal bodies.
B) The intensity of IFT20-GFP associated with ciliated (green) vs. non-ciliated(gray) basal bodies in the same MCC (in CHX-treated samples) at different time points during cilia regeneration. A total of 8-10 basal bodies per MCC (4-5 ciliated and 4-5 non-ciliated) and 5 MCCs were chosen, and the mean gray value was estimated and normalized to the maximum and the minimum values in the set. The value in parenthesis indicates the number of basal bodies analyzed (with and without IFT20-GFP) from 9 embryos from three independent trials. Note the significant difference in the IFT20-GFP signal intensity at ciliated vs. non-ciliated basal bodies at all time points. **** - $p<0.0001$, Kruskal Wallis test, followed by Dunn's multiple comparison test.

SUPPLEMENTARY VIDEO LEGENDS

Supplement Video 1: Mucociliary epithelium regenerates cilia in the same MCC.
Live imaging of animal caps dissected from embryos injected with membrane-RFP RNA. Imaging was started approximately 2 minutes post-deciliation. The time stamp on the video is when the imaging was started and not the time of deciliation.

Supplement video 2 - 6: Cilia regeneration tomogram data.
Tomograms from control and regenerating MCCs. Cilium from the control sample shows the presence of TZ, indicated by an 'H' shaped electron-dense structure (video 2). Note that this structure is missing in samples after 20 minutes of cilia regeneration (video 3) and begins to appear after 1 hour (video 4). By the end of 3 hours (video 5) and 6 hours (video 6) of cilia regeneration, the tomograms reveal complete regeneration of the transition zone.

SUPPLEMENTARY DATA TABLE

826 **Table 1**: Force data computed for different cilia numbers and lengths with maximum length of
827 15um and maximum number of 150 cilia.
828

829 **REFERENCES**
830

1.   Klena, N.T., Gibbs, B.C., and Lo, C.W. (2017). Cilia and Ciliopathies in Congenital Heart
Disease. Cold Spring Harb Perspect Biol *9*. 10.1101/cshperspect.a028266.
2.   Noone, P.G., Leigh, M.W., Sannuti, A., Minnix, S.L., Carson, J.L., Hazucha, M.,
Zariwala, M.A., and Knowles, M.R. (2004). Primary Ciliary Dyskinesia. Am J Respir Crit
Care Med *169*, 459–467. 10.1164/rccm.200303-365OC.
3.   Afzelius, B.A. (2004). Cilia-related diseases. J Pathol *204*, 470–477.
https://doi.org/10.1002/path.1652.
4.   Ishikawa, T., Ueno, H., Omori, T., and Kikuchi, K. (2021). Cilia and centrosomes:
Ultrastructural and mechanical perspectives. Semin Cell Dev Biol *110*, 61–69.
https://doi.org/10.1016/j.semcdb.2020.03.007.
5.   Nicastro, D., Schwartz, C., Pierson, J., Gaudette, R., Porter, M.E., and McIntosh, J.R.
(2006). The Molecular Architecture of Axonemes Revealed by Cryoelectron Tomography.
Science (1979) *313*, 944–948. 10.1126/science.1128618.
6.   Reynolds, M.J., Phetruen, T., Fisher, R.L., Chen, K., Pentecost, B.T., Gomez, G., Ounjai,
P., and Sui, H. (2018). The Developmental Process of the Growing Motile Ciliary Tip
Region. Sci Rep *8*, 7977. 10.1038/s41598-018-26111-2.
7.   Legal, T., Tong, M., Black, C., Valente Paterno, M., Gaertig, J., and Bui, K.H. (2023).
Molecular architecture of the ciliary tip revealed by cryo-electron tomography. Preprint,
10.1101/2023.01.03.522627 10.1101/2023.01.03.522627.
8.   Kee, H.L., Dishinger, J.F., Lynne Blasius, T., Liu, C.J., Margolis, B., and Verhey, K.J.
(2012). A size-exclusion permeability barrier and nucleoporins characterize a ciliary pore
complex that regulates transport into cilia. Nat Cell Biol *14*, 431–437. 10.1038/ncb2450.
9.   Okazaki, M., Kobayashi, T., Chiba, S., Takei, R., Liang, L., Nakayama, K., and Katoh, Y.
(2020). Formation of the B9-domain protein complex MKS1–B9D2–B9D1 is essential as
a diffusion barrier for ciliary membrane proteins. Mol Biol Cell *31*, 2259–2268.
10.1091/mbc.E20-03-0208.
10.   van den Hoek, H., Klena, N., Jordan, M.A., Alvarez Viar, G., Righetto, R.D., Schaffer,
M., Erdmann, P.S., Wan, W., Geimer, S., Plitzko, J.M., et al. (2022). In situ architecture
of the ciliary base reveals the stepwise assembly of intraflagellar transport trains. Science
(1979) *377*, 543–548. 10.1126/science.abm6704.
11.   Li, C., Jensen, V.L., Park, K., Kennedy, J., Garcia-Gonzalo, F.R., Romani, M., de Mori,
R., Bruel, A.L., Gaillard, D., Doray, B., et al. (2016). MKS5 and CEP290 Dependent
Assembly Pathway of the Ciliary Transition Zone. PLoS Biol *14*, 1–29.
10.1371/journal.pbio.1002416.
12.   Wu, Z., Chen, H., Zhang, Y., Wang, Y., Wang, Q., Augière, C., Hou, Y., Fu, Y., Peng, Y.,
Durand, B., et al. (2024). Cep131-Cep162 and Cby-Fam92 complexes cooperatively
maintain Cep290 at the basal body and contribute to ciliogenesis initiation. PLoS Biol *22*,
e3002330.
13.   Wu, Z., Pang, N., Zhang, Y., Chen, H., Peng, Y., Fu, J., and Wei, Q. (2021). CEP290 is
essential for the initiation of ciliary transition zone assembly. PLoS Biol *18*, e3001034.
14.   Vieillard, J., Paschaki, M., Duteyrat, J.-L., Augière, C., Cortier, E., Lapart, J.-A., Thomas,
J., and Durand, B. (2016). Transition zone assembly and its contribution to axoneme

formation in Drosophila male germ cells. Journal of Cell Biology *214*, 875–889.
10.1083/jcb.201603086.
15.   Garcia-Gonzalo, F.R., and Reiter, J.F. (2012). Scoring a backstage pass: Mechanisms of
ciliogenesis and ciliary access. Journal of Cell Biology *197*, 697–709.
10.1083/jcb.201111146.
16.   Sorokin, S. (1962). CENTRIOLES AND THE FORMATION OF RUDIMENTARY
CILIA BY FIBROBLASTS AND SMOOTH MUSCLE CELLS . Journal of Cell Biology
15, 363–377. 10.1083/jcb.15.2.363.
17.   Reiter, J.F., Blacque, O.E., and Leroux, M.R. (2012). The base of the cilium: roles for
transition fibres and the transition zone in ciliary formation, maintenance and
compartmentalization. EMBO Rep *13*, 608–618. https://doi.org/10.1038/embor.2012.73.
18.   Craft, J.M., Harris, J.A., Hyman, S., Kner, P., and Lechtreck, K.F. (2015). Tubulin
transport by IFT is upregulated during ciliary growth by a cilium-autonomous mechanism.
Journal of Cell Biology *208*, 223–237. 10.1083/jcb.201409036.
19.   Wingfield, J.L., Mengoni, I., Bomberger, H., Jiang, Y.-Y., Walsh, J.D., Brown, J.M.,
Picariello, T., Cochran, D.A., Zhu, B., Pan, J., et al. (2017). IFT trains in different stages
of assembly queue at the ciliary base for consecutive release into the cilium. Elife *6*,
e26609. 10.7554/eLife.26609.
20.   Kulkarni, S.S., Griffin, J.N., Date, P.P., Liem, K.F., and Khokha, M.K. (2018). WDR5
Stabilizes Actin Architecture to Promote Multiciliated Cell Formation. Dev Cell *46*, 595-
610.e3. https://doi.org/10.1016/j.devcel.2018.08.009.
21.   Gakovic, M., Shu, X., Kasioulis, I., Carpanini, S., Moraga, I., and Wright, A.F. (2011).
The role of RPGR in cilia formation and actin stability. Hum Mol Genet *20*, 4840–4850.
10.1093/hmg/ddr423.
22.   Molla-Herman, A., Ghossoub, R., Blisnick, T., Meunier, A., Serres, C., Silbermann, F.,
Emmerson, C., Romeo, K., Bourdoncle, P., Schmitt, A., et al. (2010). The ciliary pocket:
an endocytic membrane domain at the base of primary and motile cilia. J Cell Sci *123*,
1785–1795. 10.1242/jcs.059519.
23.   Rannestad, J. (1974). THE REGENERATION OF CILIA IN PARTIALLY
DECILIATED TETRAHYMENA. J Cell Biol *63*, 1009–1017.
24.   Silflow, C.D., and Lefebvre, P.A. (2001). Assembly and Motility of Eukaryotic Cilia and
Flagella. Lessons from Chlamydomonas reinhardtii . Plant Physiol *127*, 1500–1507.
10.1104/pp.010807.
25.   Werner, M.E., and Mitchell, B.J. (2013). Chapter Ten - Using Xenopus Skin to Study
Cilia Development and Function. In Cilia, Part B, W. F. B. T.-M. in E. Marshall, ed.
(Academic Press), pp. 191–217. https://doi.org/10.1016/B978-0-12-397944-5.00010-9.
26.   Kulkarni, S., Marquez, J., Date, P., Ventrella, R., Mitchell, B.J., and Khokha, M.K.
(2021). Mechanical stretch scales centriole number to apical area via Piezo1 in
multiciliated cells. Elife *10*, e66076. 10.7554/eLife.66076.
27.   Hibbard, J.V.K., Vazquez, N., Satija, R., and Wallingford, J.B. (2021). Protein turnover
dynamics suggest a diffusion-to-capture mechanism for peri-basal body recruitment and
retention of intraflagellar transport proteins. Mol Biol Cell *32*, 1171–1180.
10.1091/mbc.E20-11-0717.
28.   Sim, H.J., Yun, S., Kim, H.E., Kwon, K.Y., Kim, G.H., Yun, S., Kim, B.G., Myung, K.,
Park, T.J., and Kwon, T. (2020). Simple Method to Characterize the Ciliary Proteome of
Multiciliated Cells. J Proteome Res *19*, 391–400. 10.1021/acs.jproteome.9b00589.

29. Tadokoro, T., Wang, Y., Barak, L.S., Bai, Y., Randell, S.H., and Hogan, B.L.M. (2014).
IL-6/STAT3 promotes regeneration of airway ciliated cells from basal stem cells.
Proceedings of the National Academy of Sciences *111*, E3641–E3649.
10.1073/pnas.1409781111.
30. Parekh, K.R., Nawroth, J., Pai, A., Busch, S.M., Senger, C.N., and Ryan, A.L. (2020).
Stem cells and lung regeneration. American Journal of Physiology-Cell Physiology *319*,
C675–C693. 10.1152/ajpcell.00036.2020.
31. Craige, B., Tsao, C.C., Diener, D.R., Hou, Y., Lechtreck, K.F., Rosenbaum, J.L., and
Witman, G.B. (2010). CEP290 tethers flagellar transition zone microtubules to the
membrane and regulates flagellar protein content. Journal of Cell Biology *190*, 927–940.
10.1083/jcb.201006105.
32. Parker, J.D.K., Hilton, L.K., Diener, D.R., Rasi, M.Q., Mahjoub, M.R., Rosenbaum, J.L.,
and Quarmby, L.M. (2010). Centrioles are freed from cilia by severing prior to mitosis.
Cytoskeleton *67*, 425–430. 10.1002/cm.20454.
33. Dean, S., Moreira-Leite, F., Varga, V., and Gull, K. (2016). Cilium transition zone
proteome reveals compartmentalization and differential dynamics of ciliopathy
complexes. Proceedings of the National Academy of Sciences *113*, E5135–E5143.
10.1073/pnas.1604258113.
34. Chih, B., Liu, P., Chinn, Y., Chalouni, C., Komuves, L.G., Hass, P.E., Sandoval, W., and
Peterson, A.S. (2012). A ciliopathy complex at the transition zone protects the cilia as a
privileged membrane domain. Nat Cell Biol *14*, 61–72. 10.1038/ncb2410.
35. Tony Yang, T., Su, J., Wang, W.J., Craige, B., Witman, G.B., Bryan Tsou, M.F., and
Liao, J.C. (2015). Superresolution pattern recognition reveals the architectural map of the
ciliary transition zone. Sci Rep *5*, 1–13. 10.1038/srep14096.
36. Garcia-Gonzalo, F.R., Corbit, K.C., Sirerol-Piquer, M.S., Ramaswami, G., Otto, E.A.,
Noriega, T.R., Seol, A.D., Robinson, J.F., Bennett, C.L., Josifova, D.J., et al. (2011). A
transition zone complex regulates mammalian ciliogenesis and ciliary membrane
composition. Nat Genet *43*, 776–784. 10.1038/ng.891.
37. Dowdle, W.E., Robinson, J.F., Kneist, A., Sirerol-Piquer, M.S., Frints, S.G.M., Corbit,
K.C., Zaghloul, N.A., van Lijnschoten, G., Mulders, L., Verver, D.E., et al. (2011).
Disruption of a Ciliary B9 Protein Complex Causes Meckel Syndrome. The American
Journal of Human Genetics *89*, 589. https://doi.org/10.1016/j.ajhg.2011.09.013.
38. Ringo, D.L. (1967). STRUCTURE APPARATUS MOTION OF THE AND IN
CHLAMYDOMONAS Light Microscopy. J Cell Biol, 543–571.
39. Portman, R.W., LeCluyse, E.L., and Dentler, W.L. (1987). Development of microtubule
capping structures in ciliated epithelial cells. J Cell Sci *87*, 85–94. 10.1242/jcs.87.1.85.
40. Awata, J., Takada, S., Standley, C., Lechtreck, K.F., Bellvé, K.D., Pazour, G.J., Fogarty,
K.E., and Witman, G.B. (2014). NPHP4 controls ciliary trafficking of membrane proteins
and large soluble proteins at the transition zone. J Cell Sci *127*, 4714–4727.
10.1242/jcs.155275.
41. Greenan, G.A., Vale, R.D., and Agard, D.A. (2020). Electron cryotomography of intact
motile cilia defines the basal body to axoneme transition. Journal of Cell Biology *219*.
10.1083/jcb.201907060.
42. Diener, D.R., Lupetti, P., and Rosenbaum, J.L. (2015). Proteomic Analysis of Isolated
Ciliary Transition Zones Reveals the Presence of ESCRT Proteins. Current Biology *25*,
379–384. https://doi.org/10.1016/j.cub.2014.11.066.

43.   Garcia-gonzalo, F.R., Corbit, K.C., Sirerol-piquer, M.S., Otto, E.A., Noriega, T.R., Seol,
966   A.D., Robinson, J.F., Bennett, C.L., Josifova, D.J., García-verdugo, J.M., et al. (2012).
967   HHS Public Access. *43*, 776–784. 10.1038/ng.891.A.

968   44.   Kiesel, P., Alvarez Viar, G., Tsoy, N., Maraspini, R., Gorilak, P., Varga, V., Honigmann,
969   A., and Pigino, G. (2020). The molecular structure of mammalian primary cilia revealed
970   by cryo-electron tomography. Nat Struct Mol Biol *27*, 1115–1124. 10.1038/s41594-020-
971   0507-4.

972   45.   Wang, L., Wen, X., Wang, Z., Lin, Z., Li, C., Zhou, H., Yu, H., Li, Y., Cheng, Y., Chen,
973   Y., et al. (2022). Ciliary transition zone proteins coordinate ciliary protein composition
974   and ectosome shedding. Nat Commun *13*, 3997. 10.1038/s41467-022-31751-0.

975   46.   Craige, B., Tsao, C.-C., Diener, D.R., Hou, Y., Lechtreck, K.-F., Rosenbaum, J.L., and
976   Witman, G.B. (2010). CEP290 tethers flagellar transition zone microtubules to the
977   membrane and regulates flagellar protein content. Journal of Cell Biology *190*, 927–940.
978   10.1083/jcb.201006105.

979   47.   Dentler, W.L. (1984). Attachment of the cap to the central microtubules of Tetrahymena
980   cilia. J Cell Sci *66*, 167–173. 10.1242/jcs.66.1.167.

981   48.   Sanders, M.A., and Salisbury, J.L. (1994). Centrin plays an essential role in microtubule
982   severing during flagellar excision in Chlamydomonas reinhardtii. Journal of Cell Biology
983   *124*, 795–805. 10.1083/jcb.124.5.795.

984   49.   Kubo, A., Yuba-Kubo, A., Tsukita, S., Tsukita, S., and Amagai, M. (2008). Sentan: a
985   novel specific component of the apical structure of vertebrate motile cilia. Mol Biol Cell
986   *19*, 5338–5346. 10.1091/mbc.e08-07-0691.

987   50.   Werner, M.E., Mitchell, J.W., Putzbach, W., Bacon, E., Kim, S.K., and Mitchell, B.J.
988   (2014). Radial intercalation is regulated by the par complex and the microtubule-
989   stabilizing protein CLAMP/Spef1. Journal of Cell Biology *206*, 367–376.
990   10.1083/jcb.201312045.

991   51.   Takao, D., Wang, L., Boss, A., and Verhey, K.J. (2017). Protein interaction analysis
992   provides a map of the spatial and temporal organization of the ciliary gating zone. Current
993   Biology *27*, 2296–2306.

994   52.   Rosenbaum, J.L., and Child, F.M. (1967). Flagellar regeneration in protozoan flagellates. J
995   Cell Biol *34*, 345–364. 10.1083/jcb.34.1.345.

996   53.   Marshall, W.F., and Rosenbaum, J.L. (2001). Intraflagellar transport balances continuous
997   turnover of outer doublet microtubules : implications for flagellar length control . Journal
998   of Cell Biology *155*, 405–414. 10.1083/jcb.200106141.

54.   Auclair, W., and Siegel, B.W. (1966). Cilia Regeneration in the Sea Urchin Embryo:
Evidence for a Pool of Ciliary Proteins. Science (1979) *154*.

55.   Tasca, A., Helmstädter, M., Brislinger, M.M., Haas, M., Mitchell, B., and Walentek, P.
(2021). Notch signaling induces either apoptosis or cell fate change in multiciliated cells
during mucociliary tissue remodeling. Dev Cell *56*, 525-539.e6.
10.1016/j.devcel.2020.12.005.

56.   Bottier, M., Thomas, K.A., Dutcher, S.K., and Bayly, P. V (2019). How does cilium
length affect beating? Biophys J *116*, 1292–1304.

57.   Khona, D.K., Rao, V.G., Motiwalla, M.J., Varma, P.C.S., Kashyap, A.R., Das, K.,
Shirolikar, S.M., Borde, L., Dharmadhikari, J.A., and Dharmadhikari, A.K. (2013).
Anomalies in the motion dynamics of long-flagella mutants of Chlamydomonas
reinhardtii. J Biol Phys *39*, 1–14.

58.   Boselli, F., Jullien, J., Lauga, E., and Goldstein, R.E. (2021). Fluid mechanics of mosaic
ciliated tissues. Phys Rev Lett *127*, 198102.

59.   Rosenbaum, J.L., Moulder, J.E., and Ringo, D.L. (1969). FLAGELLAR ELONGATION
AND SHORTENING IN CHLAMYDOMONAS : The Use of Cycloheximide and
Colchicine to Study the Synthesis and Assembly of Flagellar Proteins . Journal of Cell
Biology *41*, 600–619. 10.1083/jcb.41.2.600.

60.   Quidwai, T., Wang, J., Hall, E.A., Petriman, N.A., Leng, W., Kiesel, P., Wells, J.N.,
Murphy, L.C., Keighren, M.A., Marsh, J.A., et al. (2021). A WDR35-dependent coat
protein complex transports ciliary membrane cargo vesicles to cilia. Elife *10*, e69786.
10.7554/eLife.69786.

61.   Hendel, N.L., Thomson, M., and Marshall, W.F. (2018). Diffusion as a Ruler: Modeling
Kinesin Diffusion as a Length Sensor for Intraflagellar Transport. Biophys J *114*, 663–
674. https://doi.org/10.1016/j.bpj.2017.11.3784.

62.   Fai, T.G., Mohapatra, L., Kar, P., Kondev, J., and Amir, A. (2019). Length regulation of
multiple flagella that self-assemble from a shared pool of components. Elife *8*, e42599.
10.7554/eLife.42599.

63.   Sedzinski, J., Hannezo, E., Tu, F., Biro, M., and Wallingford, J.B. (2016). Emergence of
an Apical Epithelial Cell Surface In Vivo. Dev Cell *36*, 24–35.
https://doi.org/10.1016/j.devcel.2015.12.013.

64.   Hibbard, J.V.K., Vázquez, N., and Wallingford, J.B. (2022). Cilia proteins getting to work
– how do they commute from the cytoplasm to the base of cilia? J Cell Sci *135*,
jcs259444. 10.1242/jcs.259444.

65.   Conkar, D., Bayraktar, H., and Firat-Karalar, E.N. (2019). Centrosomal and ciliary
targeting of CCDC66 requires cooperative action of centriolar satellites, microtubules and
molecular motors. Sci Rep *9*, 14250. 10.1038/s41598-019-50530-4.

66.   Conkar, D., Culfa, E., Odabasi, E., Rauniyar, N., Yates III, J.R., and Firat-Karalar, E.N.
(2017). The centriolar satellite protein CCDC66 interacts with CEP290 and functions in
cilium formation and trafficking. J Cell Sci *130*, 1450–1462. 10.1242/jcs.196832.

67.   Khokha, M.K., Chung, C., Bustamante, E.L., Gaw, L.W.K., Trott, K.A., Yeh, J., Lim, N.,
Lin, J.C.Y., Taverner, N., Amaya, E., et al. (2002). Techniques and probes for the study of
Xenopus tropicalis development. Developmental Dynamics *225*, 499–510.
https://doi.org/10.1002/dvdy.10184.

68.   Zahn, N., James-Zorn, C., Ponferrada, V.G., Adams, D.S., Grzymkowski, J., Buchholz,
D.R., Nascone-Yoder, N.M., Horb, M., Moody, S.A., Vize, P.D., et al. (2022). Normal
Table of Xenopus development: a new graphical resource. Development *149*, dev200356.
10.1242/dev.200356.

69.   Kim, S.K., Zhang, S., Werner, M.E., Brotslaw, E.J., Mitchell, J.W., Altabbaa, M.M., and
Mitchell, B.J. (2018). CLA MP/Spef1 regulates planar cell polarity signaling and
asymmetric microtubule accumulation in the Xenopus ciliated epithelia. Journal of Cell
Biology *217*, 1633–1641. 10.1083/jcb.201706058.

1051

1052

1053

Figure 1

A

Stage 25-28 embryos

15 s.

Deciliation — Wash — 1/9 x MR solution — Wash — Incubate at 28°C — Regeneration

Fix in 4% PFA

Pre. | 0 hr. | 1 hr. | 3 hr. | 5 hr. | 6 hr

Start — Regeneration — End

B Deciliation and regeneration of cilia

Pre.
Ac. tub
F-actin
Ac. tub
500 μm
20 μm
10 μm²
EHT = 3.00 kV | Signal A = SE2 | Date :24 Sep 2020
WD = 14.3 mm | Mag = 2.50 K X | Time :14:42:54

0 hr.
500 μm
20 μm
10 μm²
EHT = 3.00 kV | Signal A = SE2 | Date :24 Sep 2020
WD = 14.3 mm | Mag = 2.50 K X | Time :14:45:54

1 hr.
500 μm
20 μm
10 μm²
EHT = 3.00 kV | Signal A = SE2 | Date :24 Sep 2020
WD = 14.4 mm | Mag = 2.50 K X | Time :14:51:46

3 hr.
500 μm
20 μm
13 μm²
EHT = 3.00 kV | Signal A = SE2 | Date :24 Sep 2020
WD = 14.4 mm | Mag = 2.50 K X | Time :14:56:32

6 hr.
500 μm
20 μm
10 μm²
EHT = 3.00 kV | Signal A = SE2 | Date :24 Sep 2020
WD = 14.4 mm | Mag = 2.50 K X | Time :15:19:16

C

Cilia length (μm)

**** | **** | *** | ns

Control (34) | 0 hr (34) | 1 hr (30) | 3 hr (35) | 5 hr (34) | 6 hr (33)

Figure 2

A   Loss and regeneration of transition zone

Pre.   B9d1   Ac. tub.   5 µm

0 hr.   5 µm

1 hr.   5 µm

3 hrs.   5 µm

B   Pre.   0 mins.   20 mins.   90 mins.

C   B9d1 length in regenerating cilia

Length of B9d1 signal (µm)

**** **** ns

Pre. (60)   2 hrs. (60)   3 hrs. (60)

Figure 3

A Recovery of Sentan signal during regeneration

Pre — mScar-SNTN / memGFP — 10 µm

1 hr. — 10 µm

2 hrs. — 10 µm

3 hrs. — 10 µm

6 hrs. — 10 µm

B

Sentan length (µm)

| ns | ns | ns |

**** *

Pre. (28), 1 hr. (29), 2 hrs. (30), 3 hrs. (30), 6 hrs. (30)

Cilia length (µm)

**** ns ***

Pre. (29), 1 hr. (29), 2 hrs. (30), 3 hrs. (29), 6 hrs. (29)

A   Figure 4

Pre — Ac. tub | B9d1 | Ax, TZ, BB

0 hr.

Controls | CHX

1 hr. | **1 hr.**

2 hr. | **2 hr.**

3 hr. | **3 hr.**

B   Quantification of MCCs with B9d1 signal

- Percent MCCs with no B9d1 signal
- Percent MCCs with B9d1 signal

3 hr. CHX (49)
3 hr. DMSO (40)
2 hr. CHX (34)
2 hr. DMSO (45)
1 hr. CHX (26)
1 hr. DMSO (28)
Control (30)

****
****
****

Figure 5

A.
Stage 26-28 embryos

Deciliate
- 1/9 MR with DMSO (Vehicle)
- 1/9 MR with 10μg/ml CHX
- 1/9 MR with 10μg/ml CHX +10μM MG132

Fix
0 hr. 1 hr. 3 hrs. 6 hrs.

B. Pre   0 hr.
Ac.tub
F-Actin
10 μm   10 μm

C.   Vehicle   CHX   CHX+ MG132
1 hr   10 μm   10 μm   10 μm
3 hr   10 μm   10 μm   10 μm
6 hr   10 μm   10 μm   10 μm

D. Cilia length

Pre.   CHX
Vehicle   CHX + MG132

ns   ns   ****   ****   ****   ****

Cilia length (μm)

(50) (78) (91) (78) (74) (76) (73) (66) (79) (72)
        1 hr.   3 hrs.   6 hrs.

E. Cilia number

ns   ns   ns   ns

Cilia per MCC

(36) (37) (37) (38) (39) (38) (39)
    1 hr.   3 hrs.   6 hrs.

Figure 6

A

Ciliary force prediction per MCC during regeneration

150
100
50
0

Number of cilia

Ciliary length

B

Regeneration

1 hr.   3 hrs.   6 hrs.

Vehicle

Deciliation

Pre.   0 hr

1 hr.
Length:3 μm
Number:42
Force: 6.79 pN

3 hrs.
Length:9 μm
Number:150
Force: 162.49 pN

6 hrs.
Length:15 μm
Number:150
Force: 403.5 pN

CHX

1 hr.
Length:2 μm
N umber:34
Force: 2.79 pN

3 hrs.
Length:5 μm
Number:14
Force:5.42 pN

6hrs.
Length:9 μm
Number:8
Force: 8.67 pN

CHX + MG132

1 hr.
Length:3 μm
Number:32
Force: 5.17 pN

3 hrs.
Length:7 μm
Number:16
Force: 11.1 pN

6 hrs.
Length:12 μm
Number:10
Force: 18.10 pN

Length:15 μm
Number:150
Force: 403.5 pN

Figure 7

A — Distribution of cytosolic pool of IFT43-GFP in regenerating MCCs

Pre. — Centrin-RFP / IFT43-GFP — Ac. tub / IFT43-GFP — 0 hr.

A' Vehicle

1 hr.

3hrs.

6hrs.

A'' CHX

B — Quantification of IFT43-GFP at the basal bodies in regenerating MCCs from CHX treated embryos

● Ciliated basal body   ● Non-ciliated basal body

Normalized IFT43 intensity

**** — 1 hr. CHX (75) (76)

**** — 3 hrs. CHX (75) (77)

**** — 6 hrs. CHX (70) (80)

Figure 8

Figure S1

A — Cilia regeneration montage

0 min — Mem-RFP
9 mins
12 mins
18 mins
27 mins
10 μm

B — Effect of deciliation on MCC apical F-actin network

Pre- deciliation
Ac. tub
F-actin

Deciliated
F-actin
10 μm
10 μm

Intensity (AU)
Cortical
Pre   Decil
(29)
****

Medial
Pre   Decil
(29)
****

C — Effect of deciliation on basal body number

Pre- deciliation
Ac. tub
Chibby-BFP

Deciliated
10 μm
10 μm

Basal body number
ns
(24)
Pre.   Decil.

D — Effect of deciliation on basal body polarity

Pre- deciliation
Centrin-RFP
Clamp-GFP
Clamp-GFP

Deciliated
Clamp-GFP
Clamp-GFP
10 μm
10 μm

Deciliated
Pre

Figure S2

A — Methods of deciliation

Transfer embryos to poly-L-Lysine coated coverslips.

Chemical deciliation

Add deciliation buffer.

1 min.

Remove deciliation buffer and embryos

Fix the coverslip in methanol and stain.

Mechanical deciliation

Gently press embryos with a glass slide.

1 min.

Remove the glass slide and embryos.

Fix the coverslip in methanol and stain

B — Chemical deciliation

Ac. tubulin
B9d1

Mem. lost

TZ

10 µm

C — Mechanical deciliation

Ac. tubulin
B9d1

Mem. intact

TZ

10 µm

D — Loss of ciliary membrane upon chemical deciliation

A Recovery of Clamp signal during cilia regeneration

Pre — RFP-Clamp — memGFP

1 hr.

2 hrs.

3 hrs.

6 hrs.

B

Clamp length (μm) — ns, ****, ns, ns

Pre. (35), 1 hr. (35), 2 hrs. (35), 3 hrs. (35), 6 hrs. (35)

Cilia length (μm) — ****, *, *, ****

Pre. (36), 1 hr. (42), 2 hrs. (38), 3 hrs. (49), 6 hrs. (39)

A

Quantification of cilia length upon CHX and MG132 treatment

B

	Cilia length in µm		
	Vehicle	**CHX**	**CHX+ MG132**
Pre	15.61 ± 2.29	ND	ND
1 hr.	2.746 ± 1.045	2.242 ± 1.118	2.836 ± 1.494
3 hrs.	8.529 ± 1.736	5.4 ± 1.733	6.846 ± 1.534
6 hrs.	14.87 ± 2.379	8.798 ± 2.140	11.93 ± 2.529

C

	Cilia number		
	Vehicle	CHX	CHX+ MG132
1 hr.	41.61 ± 12.71	33.78 ± 17.27	31.65 ± 16.31
3 hrs.	ND	14.29 ± 6.164	15.85 ± 5.729
6 hrs.	ND	8.289 ± 3.986	10.28 ± 6.061

A. Quantification of BB number with CHX treatment

B.

A Distribution of cytosolic pool of IFT80-GFP in regenerating MCCs

A' Vehicle

A'' CHX

B Quantification of IFT80-GFP at the basal bodies in regenerating MCCs from CHX treated embryos

● Ciliated basal body ● Non-ciliated basal body

Figure S7

A Distribution of cytosolic pool of IFT20-GFP in regenerating MCCs

Pre.
Centrin-RFP
IFT20-GFP
Ac. tub
IFT20-GFP
10 μm

0 hr.
10 μm

A' Vehicle

A'' CHX

1 hr.
10 μm
10 μm

3 hr.
10 μm
10 μm

6 hr.
10 μm
10 μm

B Quantification of IFT20-GFP at the basal bodies in regenerating MCCs from CHX treated embryos

● Ciliated basal body ● Non-ciliated basal body

Normalized IFT20 intensity
****
1 hr. CHX
(75) (75)

****
3 hrs. CHX
(75) (74)

****
6 hrs. CHX
(75) (75)