1 **Characterization of cotton virus A, a novel and distinct member of the genus**
2 ***Caulimovirus* with endogenous viral elements in *Gossypium* spp.**
3

4 Michael West-Ortiz [1], Douglas Stuehler [2], Emma Pollock [1,3], Jennifer R. Wilson [1,^],
5 Stephanie Preising [1], Olufemi Alabi [4], Marc Fuchs [5], Michelle Heck [1,6,*], Alejandro Olmedo-
6 Velarde [1,*]
7
[1] Section of Plant Pathology and Plant-Microbe Biology, School of Integrative Plant
Science, Cornell University, Ithaca, NY, 14853, USA
[2] Oak Ridge Institute for Science and Education, USDA Agricultural Research Service,
Ithaca, NY 14853, USA
[3] Tompkins Seneca Tioga Board of Cooperative Educational Programs, New Visions
Life Sciences Program, Ithaca High School, Ithaca, NY 14850 USA
[4] Department of Plant Pathology & Microbiology, Texas A&M AgriLife Research and
Extension Center, Weslaco, TX, 78596, USA
[5] Section of Plant Pathology and Plant-Microbe Biology, School of Integrative Plant
Science, Cornell University, Geneva, NY, 14456, USA
[6] USDA Agricultural Research Service, Emerging Pests and Pathogens Research Unit,
Ithaca, NY, 14853, USA
20
21 **Present Address:**
22 ^ USDA-Agricultural Research Service, Corn, Soybean & Wheat Quality Research Unit,
23 Wooster, OH, 44691, USA
24
25 * Corresponding authors: Alejandro Olmedo-Velarde (ao383@cornell.edu); Michelle
26 Heck (michelle.heck@usda.gov)

28 Keywords: cotton, caulimovirus, DNA virus, endogenous viral elements

Abstract

The complete genome sequence of cotton virus A, a new virus infecting cotton
(*Gossypium* spp.), was determined using high-throughput sequencing, PCR and rolling
circle amplification. The 7,482-nt genome is a circular dsDNA molecule that codes for
six open reading frames similar to other members of the *Caulimovirus* genus (family
Caulimoviridae) but presenting key genome organization differences. P3, P5 and P6 are
presumed movement, coat, and reverse transcriptase proteins, respectively. P1, P2 and
P4 showed no homology to virus proteins. Endogenous viral elements were found in *G.
hirsutum* and *G. tomentosum*, but not in *G. barbadense*, *G. arboreum,* or *G. herbaceum.*

Cotton (*Gossypium spp.*) is one of the world's most important natural fibers. The
cotton industry's global, yearly economic impact has been estimated to be more than
$600 billion dollars (Khan et al. 2020). Cotton leafroll dwarf virus (CLRDV) is an
emerging viral pathogen of cotton that is transmitted by the cotton aphid, *Aphis gossypii*,
and the causative agent of cotton blue disease (CBD) in South America (Corrêa et al.,
2005), and cotton leafroll dwarf disease (CLRDD) in the USA (Brown et al. 2019, Alabi
et al. 2020; Iriarte et al. 2020; Thiessen et al. 2020). Symptoms, including stunting, leaf
rolling, interveinal chlorosis and reduced boll sets, vary depending on biotic and abiotic
factors such as cultivar, environment, and stage of growth, and can be aggravated by
other underlying stresses. The diversity of symptoms and incidence of asymptomatic
plants infected by CLRDV in the USA highlights a need to clarify if there are additional
viral agents in the field co-infecting cotton with CLRDV that may alter disease progress.
Therefore, we characterized virus populations associated with CLRDV-infected plants in
Mississippi.
62
In 2019, one *G. barbadense* and nine *G. hirsutum* plants displaying leaf rolling
and distortion were uprooted and collected from Stoneville, MS and hand-carried to
Cornell University in Ithaca, NY under USDA-APHIS-PPQ permit # P526P-22-06971.
Once at Cornell, the plants were potted in 2-gallon pots containing 50% Cornell soil mix
and 50% coarse perlite, and maintained in insect-proof cages in the greenhouse with bi-
weekly fertilization, pest management and pruning as needed. In 2022, leaf midrib and
petiole samples were collected from each plant. Additionally in 2022, six *Gossypium*
spp. samples comprising leaf midribs and petioles were also collected from plants
growing at Stoneville, MS, and displaying the symptoms described above in addition to
interveinal chlorosis. Total RNA was extracted from all samples using the Spectrum™
Total RNA Kit (Sigma-Aldrich, USA), as per the manufacturer's instructions. Three
composite samples were prepared by pooling 2, 3 and 5 total RNA samples,
respectively, using the 2019 samples, whereas the 2022 samples were pooled into a
fourth composite sample. After depleting ribosomal RNA from the composite samples
using the RiboMinus Plant kit for RNA-seq (ThermoFisher Scientific), cDNA libraries
were constructed using the TruSeq RNA library prep kit (Illumina). The libraries were
sequenced on an Illumina® NovaSeq 6000 system to obtain paired-end reads (2 × 100
bp) at the Genomics High-Throughput Sequencing (HTS) Facility at the University of
California, Irvine.
82
The data obtained from ~ 66-78 million raw reads per composite sample were
processed following previously described workflow (Larrea-Sarmiento et al., 2022). The
identity of contigs larger than 500 bp was annotated by BLASTx searches of the NCBI
virus sequence database as of September 2022. The annotated virus contigs revealed
sequence similarity to CLRDV and caulimoviruses. The four caulimovirus contigs (2,043
to 7,217 nt), one from each of the four composite samples, shared ~98% nucleotide
identity with each other, had significant matches with dahlia mosaic virus (DMV) and
other caulimoviruses, but were sufficiently divergent suggesting they represented a
sequence of a potential new virus within the family *Caulimoviridae*.
92

Since some caulimovirus genomes have been found integrated into their host
genome and regarded as endogenous viral elements (EVEs) (Teycheney et al., 2020),
DNase-RT-PCR assays were performed with contig-specific primers (Table S1) on
DNase-treated RNA using all the 2019 and 2022 samples, as previously detailed
(Olmedo-Velarde et al., 2022) to detect viral RNA and exclude amplification of DNA
endogenous viral elements. This assay included RNA extracted from an uninfected *G.*
hirsutum seedling as a negative control. Caulimovirus-like transcripts were detected in
all the ten 2019, and four 2022 samples, except for the negative control (Figure 1A).
Sanger sequencing of purified PCR amplicons revealed they had 100% nucleotide
identity with the Illumina sequence contigs. Furthermore, rolling circle amplification
(RCA) was used to amplify the episomal circular DNA virus genome and distinguish it
from possible EVEs in *Gossypium*. Total nucleic acids were extracted using the OPS
Synergy 2.0 Plant DNA Extraction Kit (OPS Diagnostics, Lebanon, NJ, USA), and
subjected to RCA using Equiphi29 DNA polymerase (Thermoscientific, Whatman, PA),
as per the manufacturer's instructions. RCA reactions were digested with *KpnI*. A ~2.1
kbp DNA fragment was observed by agarose gel electrophoresis for all samples (data
not shown). The amplicons were gel-purified and directly sequenced at PlasmidSaurus
(Eugene, OR) using Oxford Nanopore sequencing. The consensus sequence (30x
coverage) showed 99% nucleotide identity to the caulimovirus Illumina contig sequence,
confirming the episomal presence of a putative new caulimovirus in the cotton samples.
113
To obtain the complete genome sequence of the novel virus and validate the
HTS results, one 2022 sample was selected for further analysis and five contig-specific
overlapping primer sets (Table S1) were designed to fill gaps in the circular genome of
the virus. PCR assays were performed using ten-fold diluted RCA products as the
template. The amplicons were cloned, and three clones per amplicon were sequenced
using Sanger and Nanopore sequencing. The complete genome of the novel virus
comprises a 7,482-nt dsDNA circular molecule (GenBank accession). We consider the
sequence to represent a new caulimovirus for which the common name cotton virus A
(CotV-A) and species name Caulimovirus gossypii are proposed.
123
The genome of CotV-A starts with the conserved tRNAMet sequence 5′-
TGGTATCAGAGCC-3′ and shares 66.7% nucleotide sequence identity with grapevine
pararetrovirus (OP886324) with 22% query coverage. Six putative ORFs (>300 bp)
were identified in the complete CotV-A genome sequence (Figure 1B) using ORF Finder
(https://www.ncbi.nlm.nih.gov/orffinder/). ORF1 is at positions 38-406 nt; ORF2, 346-
951 nt; ORF3, 948-2021 nt; ORF4, 2018-2347 nt; ORF5, 2440-3672 nt; and
ORF6,4014-5948 nt. The predicted amino acid sequences of the putative proteins (P1
to P6) encoded by ORF 1-6 were searched against the Pfam and GenBank databases
using the InterPro and BLASTP servers, respectively, for functional annotation of the
conserved domains/motifs. P1 (122 aa, 14.4 kDa), P2 (201 aa, 24.4 kDa) and P4 (109
aa, 12.8 kDa) did not display homology to any known caulimovirus or known virus
proteins. In contrast, P3 (357 aa, 41 kDa) represents a putative movement protein (MP)
(PF01107), P5 is a putative viral coat protein (CP) (410 aa, 49.5 kDa) and P6 (644 aa,
75.9 kDa) is a typical polyprotein, akin other caulimoviruses, containing aspartic
protease (PF02160, aa 22-229), reverse transcriptase (RT) (PF00078, aa 285-470) and
RNase H1 (PF17919, aa 534-629) motifs. CotV-A has three intergenic regions (IGRs)

(Figure 1B); the shorter IGR is 92 bp in length separating ORFs 4 and 5, the
intermediate IGR is 341 bp and is between ORFs 5 and 6, whereas the long IGR (1571
bp) lies between ORFs 1 and 6. In the intermediate and long IGR, tentative TATA-like
boxes were found in addition to a polyadenylation signal in the large IGR, which is a
feature typical of caulimoviruses (Petrzik et al., 1998). The nonanucleotide
'TATAAAAAA' (nt 3859-3867) was found in the intermediate IGR, and 'TATAAAT' (nts
6683-6689, 7264-7270, 7395-7401), 'TATAAAAT' (nt 6997-7004) and the
polyadenylation signal 'AATAA' (nt 7472-7476) were found in the long IGR.
148
Pairwise sequence comparisons showed that the overall nucleotide sequence
identity between CotV-A and other caulimoviruses ranged from 62.92% to 66.74%. The
amino acid sequence identities of the CotV-A putative P3, P5 and P6 with homologs of
other caulimoviruses ranged from 23.15% to 54.40% (Table S2). To gain insight into the
taxonomic position of CotV-A and its relatedness to members of the family
Caulimoviridae, phylogenetic analysis using the maximum-likelihood algorithm based on
the RT-RNaseH protein region sequence was performed using the best model of protein
evolution (rtREV+G+I) in MEGA 11 v11.0.13 (Tamura et al., 2021) after 1,000 bootstrap
replicates. The results showed the clustering of CotV-A into the *Caulimovirus* genus
clade in a subclade also grouping grapevine pararetrovirus and plant-associated
caulimovirus 1 (Figure 2). For members within the genus *Caulimovirus*, the species
demarcation criterion requires divergence of more than 20% in the RT-RNase
nucleotide region. Using pairwise comparisons, the sequence identities of this
nucleotide region of CotV-A and caulimovirus homologs were all below 66.6%. This
result provides evidence that CotV-A is a putative new species in the genus
Caulimovirus.
165
To evaluate CotV-A integration into the cotton genome, BLASTn searches were
performed using the CotV-A genome sequence as query against chromosome-level
genome sequences of the four commercially grown *Gossypium* species, *G. hirsutum*, *G.*
barbadense, *G. arboreum* and *G. herbaceum.* Also, chromosome-level genome
sequences of *G. tomentosum*, a very close wild *Gossypium* species was analyzed. All
the *Gossypium* spp. genome sequences used in this study were from the CottonGen
database (www.cottongen.org/) (Yu et al., 2021). Additionally, mapping of quality
trimmed virus reads was performed against the *Gossypium* spp. genome sequences
using the Geneious mapper (Kearse et al., 2012). A total of four structurally unique
EVEs were found in *G. hirsutum* and *G. tomentosum.* These contained the same
predicted conserved protein domains as CotV-A (Figure 3A). No EVE was found in the
other *Gossypium* species. In *G. hirsutum*, chromosomes A04 (11 out of the 13 genomes
analyzed) and D12 (1 out of the 13 genomes analyzed) contained EVEs as inverted
tandem repeat sequences (EVE 1 and 2, Figure 3B) which presented ~99% nucleotide
identity, and 84.5-100% protein identities to CotV-A (Table S3). Also, a single
integration event was found in chromosome D03 (13 out of the 13 genomes analyzed)
which was not only truncated (EVE 3, Figure 3B) but also presented ~80% nucleotide
identity, and 54-94% protein identities to CotV-A. Interestingly, EVE 1, which is either
inserted in the sense direction (chromosome A04), or in the anti-sense direction
(chromosome D12), contains two additional ORFs which were not present in the
episomal CotV-A genome (Figure 3B). BLASTx searches of the region containing both

ORFs showed that the concatenated predicted protein sequence shares low (<27%)
identity to the P6 of grapevine pararetrovirus and other caulimoviruses. P6 is a
multifunctional protein involved in viroplasm formation, and other virus cycle processes
as well as suppression of antiviral defenses (Pooggin and Ryabova 2018). In EVE 2, in
each tandem repeat on chromosomes A4 and D12, the CP and the polyprotein
sequences were interrupted, and spread across two predicted ORFs (Figure 3A and
3B). *In G. tomentosum*, chromosomes A03 (2 out of 2 genomes analyzed) contained a
single virus integration event (EVE 4) which lacks ORF 1 and 2, and similar to EVE 2,
the CP and the polyprotein sequences are each coded across two ORFs (Figure 3A and
3B). EVE 4 presented ~79% nucleotide identity, and 71-89% protein identities to CotV-A
(Table S3). The genomic locations in the chromosomes where EVE 1-3 are inserted
were analyzed in the *G. hirsutum* genomes described above using bedtools v2.27.1
(Quinlan et al., 2010) and chromoMap v4.1.1 (Anand & Rodriguez Lopez, 2022). EVE
insertion sites were found to be highly conserved across the available genomes (Figure
S2), even in chromosome D12, where the entire region of the chromosome appears to
present in the anti-sense direction as the insertion site on A4 (Figure S2A).
203
To infer the evolutionary relationships between CotV-A and the EVEs,
phylogenetic analyses were performed as detailed above using the MP and peptidase
sequences, which are the only proteins with caulimovirid homologs present among the
episomal virus and the EVEs. For both proteins, phylogenetic analyses showed that
EVE4 is more closely related to an ancestor virus of CotV-A, than the other EVEs and
CotV-A (Figure S1). Together with the absence of an EVE 4 homolog in chromosome
A03 of *G. hirsutum*, these results support the hypothesis that an ancestor of CotV-A
infected both *Gossypium* species after speciation took place, and at different time
frames, with the infection and subsequent viral element integration into *G. hirsutum*
likely being more recent. Intriguingly, the phylogenic analysis of the MP shows that
CotV-A and its EVEs are more closely related to soymoviruses than to caulimoviruses
(Figure S1A), suggesting that CotV-A could be a recombinant virus resulting from co-
infection of a soymovirus and a caulimovirus.
217
Due to the unique attributes that CotV-A presents, such as the lack of ORFs
presenting homology to proteins involved in aphid transmission (caulimovirus P2 and
P3) and viroplasm formation (caulimovirus P6), further molecular and biological studies
are required to understand if it is pathogenic in cotton. In addition, the EVEs in the
genomes of *G. hirsutum* and *G. tomentosum* should be further corroborated because
the analysis was performed using publicly available genomic data, which may represent
incomplete or inaccurate cotton genome assemblies. It will also be interesting to
determine whether CotV-A is aphid transmissible, and forms viroplasms akin to other
caulimoviruses. Moreover, further experiments are needed to determine if abiotic factors
such as heat and drought, and/or biotic factors such as insect feeding and/or CLRDV or
other virus co-infection, trigger the formation of the episomal form of CotV-A from the
EVEs.
230
231
232
233

Acknowledgements and Conflicts of Interest.

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under No. Grant DGE–2139899. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. The research was funded by NIFA CARE grant 8062-22410-007-014-I and USDA ARS project number 8062-22410-007-000-D. We are grateful to members of the Heck Lab for helpful discussion and Chad Thomas, Lisa Scanlon and Julie Bojanowski for their help in maintaining the cotton plants used in the study. We are grateful to Jodi Scheffler from the Crop Genetics Research Unit (USDA Agricultural Research Service) for her helpful and invaluable assistance in plant and sample collection.

References

Alabi, OJ, T Isakeit, R Vaughn, D Stelly, K Conner, BC Gaytán, C Villegas, C Hitzelberger, L De Santiago, C Monclova-Santana, and JK Brown. 2020. First Report of Cotton leafroll dwarf virus Infecting Upland Cotton (Gossypium hirsutum) in Texas. Plant Dis, 104: 998.

Anand, L, Rodriguez-Lopez, CM, 2022. ChromoMap: an R package for interactive visualization of multi-omics data and annotation of chromosomes. BMC Bioinformatics 23, 33.

Aramaki, T, et al. 2020. KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold. Bioinformatics 36(7): 2251–2252.

Avelar, S, DW Schrimsher, K Lawrence, and JK Brown. 2019. First Report of Cotton leafroll dwarf virus Associated with Cotton Blue Disease Symptoms in Alabama. Plant Dis, 103: 592.

Brown, S., Conner, K., Hagan, A., Jacobson, A., Koebernick, J., Lawrence, K., et al. (2019). Report of a research review and planning meeting on cotton leafroll dwarf virus. Cotton Incorporated. Available at: https://www.cottoninc.com/wp-content/uploads/2019/11/10-19-CLRDV-Research-Review-Meeting-Report-Nichols.pdf (Accessed June 13, 2023).

Corrêa, R. L., T. F. Silva, J. L. Simões-Araújo, P. A. Barroso, M. S. Vidal, and M. F. Vaslin. 2005. Molecular characterization of a virus from the family Luteoviridae associated with cotton blue disease. Arch Virol, 150: 1357-67.

Iriarte, FB, KK Dey, IM Small, KN Conner, GK O'Brien, L Johnson, C Savery, E Carter, D Sprague, RL Nichols, DL Wright, MJ Mulvaney, and M Paret. 2020. First Report of Cotton Leafroll Dwarf Virus in Florida. Plant Dis, 104: 2744.

Kearse, M, Moir, R, Wilson, A, Stones-Havas, S, Cheung, M, Sturrock, S, et al. (2012). Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28, 1647–1649.

Khan, MA, Wahid, A, Ahmad, M, Tahir, MT, Ahmed, M, Ahmad, S, & Hasanuzzaman, M 2020. World cotton production and consumption: An overview. *Cotton production and uses: Agronomy, crop protection, and postharvest technologies*, 1-7.

Olmedo-Velarde, A, Loristo, J, Kong, A, Waisen, P, Wang, K-H, Hu, J and Melzer, M.
2022. Examination of the Virome of Taro Plants Affected by a Lethal Disease, the
Alomae-Bobone Virus Complex, in Papua New Guinea. Viruses 14(7): 1410.
Larrea-Sarmiento, AE, Olmedo-Velarde, A, Wang, X, Borth, W, Domingo, R,
Matsumoto, TK, Suzuki, JY, Wall, MM, Melzer, MJ and Hu, J. 2022. Genetic
Diversity of Viral Populations Associated with Ananas Germplasm and
Improvement of Virus Diagnostic Protocols. Pathogens, 11(12): 1470.
Petrzik, K, Beneš, V, Mráz, I, Honetšlegrová-Fránová, J, Ansorge, W, Špak, J. 1998.
Strawberry Vein Banding Virus—Definitive Member of the Genus Caulimovirus.
Virus Genes 16, 303–305.
Pooggin, MM, Ryabova, LA. 2018. Ribosome Shunting, Polycistronic Translation, and
Evasion of Antiviral Defenses in Plant Pararetroviruses and Beyond. Front
Microbiol. 9: 644.
Quinlan, AR, Hall, IM, 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26(6): 841–842.
Tamura, K, Stecher, G, and Kumar, S. 2021. MEGA11: Molecular Evolutionary Genetics
Analysis version 11. Mol Biol Evol, 38:3022-3027
Teycheney, PY, Geering, ADW, Dasgupta, I, Hull, R, Kreuze, JF, Lockhart, B, Muller, E,
Olszewski, N, Pappu, H, Pooggin, MM, Richert-Pöggeler, KR, Schoelz, JE, Seal,
S, Stavolone, L, Umber, M, 2020. Report Consortium I. ICTV Virus Taxonomy
Profile: Caulimoviridae. J Gen Virol, 101(10):1025-1026.
Thiessen, LD, Schappe, T, Zaccaron, M, Conner, K, Koebernick, J, Jacobson, A
Huseth, A. 2020. First Report of Cotton Leafroll Dwarf Virus in Cotton Plants
Affected by Cotton Leafroll Dwarf Disease in North Carolina. Plant Dis, 104:
3275.
Yu, J, Jung, S, Cheng, C-H, Lee, T, Zheng, P, Buble, K, Crabb, J, Humann, J, Hough,
H, Jones, D, Campbell, JT, Udall, J, Main, D. 2021. CottonGen: The Community
Database for Cotton Genomics, Genetics, and Breeding Research. Plants 10,
2805.
307
308
309
310

Figure 1. Detection of virus transcripts using DNase-RT-PCR assays (A) and a cartoon of the genome organization of cotton virus A (CotV-A) (B). A) RNA extracts of 2019 (lanes 1-10) and 2022 (lanes 1-6) samples were DNase-digested and subjected to RT-PCR using movement protein (MP) gene-specific primers to amplify a 474 bp product (Table S1). S = *Gossypium hirsutum* seedling. NTC = non-template control. B) A cartoon of the genome organization and predicted open reading frames (ORF) coding for six putative proteins that have conserved virus protein domains: MP (PF01107), aspartic protease (PF02160), reverse transcriptase (PF00078) and RNase H1 (PF17919). The genome is drawn to scale.

324
Figure 2. Phylogenetic relationships of the RT-RNase H protein sequence of cotton
virus A (CotV-A) in relation to other members of the family *Caulimoviridae*. The
maximum-likelihood method with the best model of protein evolution (rtREV+G+I) was
used with 1,000 bootstrap replicates as percentage values for branch support. Predicted
amino acid sequences were used and the respective GenBank accession number is
shown with each virus name. The alignment was generated using MUSCLE and
implemented in MEGA v.11.0.13. The green circle indicates CotV-A.

Figure 3. Illustration representing endogenous virus elements (EVEs) in *Gossypium* spp. genomes that likely originated from infection by ancestor of cotton virus A (CotV-A). EVEs were detected *in silico* in *G. hirsutum* and *G. tomentosum* chromosome-level genomes available in the CottonGen database. A) EVEs 1-4 present unique, structurally different gene organizations compared to CotV-A. B) EVE 1 and 2 in *G. hirsutum* were found as inverted tandem repeats in chromosomes A04 (11 out of 13 genomes) and D12 (1 out of 13 genomes). EVE 3 was found in chromosome D03 (13 out of 13 genomes) as a single integration event and was considerably shorter than the episomal CotV-A genome. C) EVE 4 found in *G. tomentosum* that was found as a single integration event in chromosome A03.

A

Movement protein (PF01107) 🟩
Aspartic Protease (PF02160) 🟥
Reverse Transcriptase (PF00078) 🟧
RNase H1 (PF17919) 🟨

Episomal Form

Integrated Forms

EVE 1
EVE 2
EVE 3

EVE 4

Gossypium hirsutum

Gossypium tomentosum

B

Chr A04
EVE 1
EVE 2

Chr D12
EVE 2
EVE 1

Chr D03
EVE 3

C

Chr AO3
EVE 4