Homeostatic Synaptic Plasticity of Miniature Excitatory Postsynaptic Currents in Mouse Cortical

Cultures Requires Neuronal Rab3A

Andrew G. Koesters[1*], Mark M. Rich[2], and Kathrin L. Engisch[3]

[1]Department of Pharmacology, Physiology, and Neurobiology, University of Cincinnati College
of Medicine, Cincinnati, OH 45267, USA
[2]Department of Neuroscience, Cell Biology and Physiology, Boonshoft School of Medicine,
Wright State University, Dayton, OH 45345
[3]Department of Neuroscience, Cell Biology and Physiology, Boonshoft School of Medicine and
the College of Science and Mathematics, Wright State University, Dayton, OH 45435
12

*Andrew G. Koesters –Corresponding author
Department of Pharmacology, Physiology, and Neurobiology, University of Cincinnati College of
Medicine, Cincinnati, OH 45267, USA
koesteag@ucmail.uc.edu, ph 513-558-2819

Author contributions: A.G.K., K.L.E. and M.M.R. designed research; A.G.K. performed research;
A.G.K. and K.L.E. analyzed data; A.G.K, K.L.E., and M.M.R. wrote the paper.
20
21   ORCID iDs: ; A.G.K, https://orcid.org/0000-0003-3281-188X; M.M.R., https://orcid.org/0000-
22   0002-6956-5531; K.L.E., https://orcid.org/0000-0002-1058-5343
23

The authors have no competing interests.

Keywords: Homeostatic Synaptic Plasticity; Synaptic Scaling; mEPSCs; Rab3; AMPA receptors

Acknowledgements: The authors thank E. Whitlock and H. Ghouse for their contributions to
confocal image analyses. This work was partially supported by NINDS P01 NS057228, Tim Cope,
PI, KE, PI project 4.

Abstract

Following prolonged activity blockade, amplitudes of miniature excitatory postsynaptic currents

(mEPSCs) increase, a form of plasticity termed "homeostatic synaptic plasticity." We previously

showed that a presynaptic protein, the small GTPase Rab3A, is required for full expression of

the increase in miniature endplate current amplitudes following prolonged blockade of action

potential activity at the mouse neuromuscular junction in vivo (Wang et al., 2011), but it is

unknown whether this form of Rab3A-dependent homeostatic plasticity shares any

characteristics with central synapses. We show here that homeostatic synaptic plasticity of

mEPSCs is impaired in mouse cortical neuron cultures prepared from Rab3A$^{-/-}$ and mutant mice

expressing a single point mutation of Rab3A, Rab3A Earlybird mice. To determine if Rab3A is

involved in the well-established homeostatic increase in postsynaptic AMPA-type receptors

(AMPARs), we performed a series of experiments in which electrophysiological recordings of

mEPSCs and confocal imaging of synaptic AMPAR immunofluorescence were assessed within

the same cultures. We found that the increase in postsynaptic AMPAR levels was more variable

than that of mEPSC amplitudes, suggesting other factors may contribute to the homeostatic

increase in synaptic strength. Finally, we demonstrate that Rab3A is acting in neurons because

only selective loss of Rab3A in neurons, not glia, disrupted the homeostatic increase in mEPSC

amplitudes. This is the first demonstration that a protein thought to function presynaptically is

required for homeostatic synaptic plasticity of quantal size.

Introduction

One of the most studied phenomena triggered by prolonged activity blockade is the

increase in amplitudes of miniature excitatory postsynaptic currents (mEPSCs) in neurons. First

demonstrated in cultures of dissociated cortical neurons (Turrigiano et al., 1998) and spinal

cord neurons (O'Brien et al., 1998), the compensatory response to prolonged loss of activity has

been dubbed "homeostatic synaptic plasticity" (Turrigiano and Nelson, 2004; Pozo and Goda,

2010). These first two studies provided evidence that glutamate receptors were also increased

after prolonged inactivity, via an increased response to exogenously applied glutamate

(Turrigiano et al., 1998) and increased immunofluorescent labeling of GluA1 receptors at

synaptic sites (O'Brien et al., 1998). The accompanying increase in synaptic AMPA receptors has

been confirmed with immunofluorescence in multiple studies of hippocampal and cortical

cultures treated with TTX, including (Wierenga et al., 2005; Hou et al., 2008; Ibata et al., 2008;

Jakawich et al., 2010; Tatavarty et al., 2013; Xu and Pozzo-Miller, 2017; Dubes et al., 2022). It is

now well accepted that the homeostatic increase in mEPSC amplitudes in neurons is due to an

increase in postsynaptic AMPA receptors.

In our previous work, we found that a TTX cuff applied for 48 hrs around the sciatic

nerve in mice led to an increase in the amplitude of the miniature endplate currents (mEPCs)

recorded in tibialis anterior muscles. Surprisingly, and unlike at central synapses, we could find

no evidence that the acetylcholine receptors levels at the neuromuscular junction (NMJ) were

increased in the TTX-cuff treated mice (Wang et al., 2005). This result led us to search for

presynaptic molecules that might homeostatically regulate the presynaptic quantum. In

previous studies in chromaffin cells, we identified the small GTPase Rab3A, a presynaptic vesicle

protein, as a regulator of synaptic vesicle fusion pore opening (Wang et al., 2008), so we

examined whether deletion of Rab3A (Rab3A$^{-/-}$) might prevent homeostatic upregulation of

mEPC amplitude at the NMJ. The results were clear: the homeostatic increase in mEPC

amplitude induced by the TTX cuff was strongly reduced in the Rab3A$^{-/-}$ mouse, and was

completely abolished in the Earlybird mutant (Rab3A$^{Ebd/Ebd}$), which has a single point mutation

in Rab3A that causes a shift towards early awakening due to a shorter circadian period, that is

more dramatic than that of the Rab3A deletion mouse (Kapfhamer et al., 2002; Wang et al.,

2011). These results led us to conclude that the homeostatic increase in mEPC amplitude at the

NMJ is a presynaptic phenomenon.

It has remained somewhat of a mystery what the role of Rab3A is in synaptic

transmission. The Rab3A$^{-/-}$ mouse has minimal phenotypic abnormalities, with evoked synaptic

transmission and mEPSCs essentially normal in hippocampal slices (Geppert et al., 1994). At the

Rab3A$^{-/-}$ NMJ, reductions in evoked transmission were detected, but only under conditions of

reduced extracellular calcium (Coleman et al., 2007). There are some modest changes in short

term plasticity during repetitive stimulus trains, with increased depression observed in

response to moderate frequencies in hippocampal slices (Geppert et al., 1994) and at the

mammalian NMJ (Coleman and Bykhovskaia, 2009), but increased facilitation in response to

high frequency stimulation at the mammalian NMJ (Coleman and Bykhovskaia, 2010). The most

dramatic effect of loss of Rab3A is the abolishment of a presynaptic form of long-term

potentiation (LTP) at the mossy fiber-CA3 synapse that is induced by 25 Hz stimulation in the

presence of NMDA blockers (Weisskopf et al., 1994; Castillo et al., 1997). Our result that

prolonged inactivity-induced plasticity of mEPC amplitude at the mammalian NMJ is disrupted

with loss of function of Rab3A further cements a role for Rab3A in activity-dependent plasticity

of synaptic strength. However, a crucial question that remains is whether the TTX-cuff induced

effect on the amplitude of the spontaneous synaptic current at the NMJ shares underlying

mechanisms with central synapses. We set out to answer this question for the more well-

studied phenomenon of homeostatic plasticity of mEPSCs in dissociated mouse cortical neurons

prepared from wild type, Rab3A$^{-/-}$ mice and Rab3a$^{Ebd/Ebd}$ mice treated with TTX for 48 hrs. We

show here that homeostatic synaptic plasticity of mEPSC amplitude was strongly reduced in

cultures from Rab3A$^{-/-}$ mice, and completely abolished in cultures from Rab3A$^{Ebd/Ebd}$ mice,

supporting a shared mechanism with the neuromuscular junction.

Results

We previously reported that mixed cultures of cortical neurons and glia prepared from

postnatal day 0-2 mouse pups responded to a block of action potential-mediated activity by a

48 hr TTX treatment with an increase in mEPSC amplitude (Hanes et al., 2020). Here, we asked

the question whether cortical cultures prepared from mice lacking the small GTPase Rab3A, or,

expressing a point mutation of Rab3A, Rab3A Earlybird, have an altered homeostatic plasticity

response to a prolonged loss of network activity. The Rab3A Earlybird mutation was discovered

in a mutagenesis screen in mice for genes involved in the rest-activity cycle (Kapfhamer et al.,

2002). In that study, the authors concluded that the Earlybird mutation was likely dominant-

negative, as the shift in circadian period was substantially more dramatic than that observed in

the Rab3A$^{-/-}$ mice. As described in the Introduction, the effects of loss of Rab3A have been

modest, so we included experiments using the Rab3A Earlybird mutant mice to see if there

would be a more robust phenotype, and in doing so, strengthen our conclusions based on the

Rab3A deletion mice. To obtain Rab3A$^{-/-}$ and Rab3A$^{Ebd/Ebd}$ homozygotes, we established two

mouse colonies of heterozygous breeders with cultures prepared from pups derived from a

final breeding pair of homozygotes. Although we backcrossed Rab3A$^{+/-}$ with Rab3A$^{+/Ebd}$ for 11

generations, clear differences in mEPSC amplitudes in untreated cultures (see below) and in

calcium current amplitudes in adrenal chromaffin cells (unpublished obs.) remained. Therefore,

throughout this study we keep the two strains separate and there are two Rab3A$^{+/+}$ or 'wild

type' phenotypes.

We found that loss of Rab3A resulted in a greatly impaired homeostatic increase after

activity blockade. Example current traces of spontaneously occurring mEPSCs recorded from

pyramidal neurons in untreated (CON) 13-14 DIV cortical cultures and sister cultures treated

with 500 nM TTX for 48 hr prepared from wild type and Rab3A$^{-/-}$ mice in the Rab3A$^{+/-}$ colony are

shown in Figures 1A and B, respectively. Average mEPSC waveforms from the same recordings

are shown in Figures 1C and D. The mean mEPSC amplitudes for 30 control and 23 TTX-treated

neurons from Rab3A$^{+/+}$ cultures are displayed in the box and whisker plot in Figure 1E; after

activity blockade the average mEPSC amplitude increased from 13.9 ± 0.7 pA to 18.2 ± 0.9 pA.

In contrast, in cultures prepared from Rab3A$^{-/-}$ mice, the average mEPSC amplitude was not

significantly increased, for 25 untreated cells and 26 TTX-treated cells (Figure 1F, 13.6 ± 0.1 pA

vs. 14.3 ± 0.6 pA). We found that TTX treatment also resulted in an increase in mEPSC

frequency in cultures prepared from Rab3A$^{+/+}$ mice, as shown in Figure 1G (CON, 2.26 ± 0.37

sec^{-1}; TTX, 4.62 ± 0.74 sec^{-1}). This TTX-induced frequency increase was strongly reduced in

cultures from Rab3A$^{-/-}$ mice, but there were a few outliers with very high frequencies still

present after TTX treatment (Figure 1H, CON, 2.74 ± 0.49 Hz; TTX, 3.23 ± 0.93 Hz).

We next examined homeostatic plasticity in cultures prepared from wild type mice in

the Rab3A$^{+/Ebd}$ colony. Example current traces of mEPSCs, and average mEPSC waveforms, are

shown in Figures 2A and C, respectively, for cortical cultures prepared from wild type mice in

the Rab3A$^{+/Ebd}$ colony. Treatment with TTX for 48 hr led to a significant increase in the average

mEPSC amplitude of 23 TTX-treated cells when compared to 20 untreated cells (Figure 2E, CON,

11.0 ± 0.6 pA; TTX, 15.0 ± 1.3 pA). We note here that while the two wild type strains respond

very similarly to activity blockade, the mean mEPSC amplitude in untreated cultures was

significantly different in the two wild type strains, 13.9 ± 0.7 pA, wild type from Rab3A$^{+/-}$ colony;

11.0 ± 0.6 pA, wild type from Rab3A$^{+/Ebd}$ colony (p = 0.004, Kruskal-Wallis test).

We found a complete disruption of homeostatic plasticity in cortical cultures prepared

from Rab3A$^{Ebd/Ebd}$ mice, as can been seen in viewing example mEPSC traces and average mEPSC

waveforms (Figures 2B and D, respectively). The lack of TTX effect was confirmed in a

comparison of mEPSC amplitude means for 21 untreated and 22 TTX-treated cells (Figure 2F,

CON, 15.1 ± 1.0 pA vs. TTX, 14.6 ± 1.1 pA). There was a trend towards increased mEPSC

frequency in TTX-treated cultures from this wild type strain, but the difference was not as

robust as for the wild type cultures from Rab3A$^{+/-}$ colony, and did not reach statistical

significance (Figure 2G, CON, 1.15 ± 0.19 sec^{-1}; TTX, 2.54 ± 0.55 sec^{-1}). This trend was not

observed in neurons from cultures prepared from Rab3A$^{Ebd/Ebd}$ mice, likely due to an increase in

frequency in untreated cells—frequencies remained high in TTX-treated cells (Figure 2H, CON,

1.71 ± 0.41 sec^{-1}; TTX, 3.05 ± 0.80 sec^{-1}). Our results show that the homeostatic increase in

mEPSC amplitude after activity blockade is disrupted in both Rab3A$^{-/-}$ and the Rab3A$^{Ebd/Ebd}$

cortical neurons, strongly supporting a crucial role of functioning Rab3A in this process.

The disruption in homeostatic plasticity differs for Rab3A$^{-/-}$ and Rab3A$^{Ebd/Ebd}$. In the

Rab3A$^{-/-}$ data set, mEPSCs from untreated cultures were indistinguishable from mEPSCs from

Rab3A$^{+/+}$ untreated cultures, demonstrating loss of Rab3A has no impact on basal activity

mEPSC amplitudes. In the Rab3A$^{Ebd/Ebd}$ data set, mEPSC amplitudes from untreated cultures

were significantly larger than those of untreated cultures from wild type mice, as can be seen

comparing the CON values in Figures 2E and F (CON, Rab3A$^{+/+}$, 11.0 ± 0.7 pA, vs. CON,

Rab3A$^{Ebd/Ebd}$, 15.1 ± 1.0 pA, p = 0.0027, Kruskal-Wallis test). The increase in mEPSC amplitude in

cultures from Rab3A$^{Ebd/Ebd}$ mice is consistent with the increase in mEPC amplitude we observed

at the Rab3A$^{Ebd/Ebd}$ NMJ (Wang et al., 2011).

The small GTPase Rab3A is generally thought to function presynaptically to regulate

synaptic vesicle trafficking, possibly in an activity-dependent manner (Castillo et al., 1997;

Lonart et al., 1998; Leenders et al., 2001; Schluter et al., 2006; Coleman and Bykhovskaia, 2009;

Tian et al., 2012). In contrast, homeostatic plasticity of mEPSC amplitude has been attributed to

an increase in postsynaptic receptors on the surface of the dendrite (O'Brien et al., 1998;

Turrigiano et al., 1998). At the NMJ in vivo, we could find no evidence for an increase in AChRs

after TTX block of the sciatic nerve in vivo (Wang et al., 2005). Is Rab3A required for the

homeostatic increase in surface AMPA-type glutamate receptors that has been confirmed by

multiple studies (see Introduction)?

We explored whether Rab3A might be acting to increase postsynaptic AMPA receptors.

In some studies of homeostatic synaptic plasticity it has been found that application of a Ca^{2+}-

permeable AMPA receptor-specific inhibitor reversed the homeostatic increase in mEPSC

amplitude (Ju et al., 2004; Thiagarajan et al., 2005; Sutton et al., 2006). Figure 3A shows that

the *TTX-induced increase* in mean mEPSC amplitude was nearly identical in a set of 11 CON and

11 TTX-treated cells before and after treatment with 1-Naphthyl acetyl spermine trihydrochloride

(NASPM), a synthetic analogue of the Ca^{2+}-permeable AMPA receptor inhibitor philanthotoxin

(before NASPM, CON 12.9 ± 3.5 pA; TTX, 17.5 ± 3.1 pA; after NASPM, CON 11.9 ± 2.6 pA; TTX

16.1 ± 3.5 pA). This result indicates that the TTX-induced increase in mEPSC amplitude does not

depend on Ca^{2+}-permeable receptors, since the effect of TTX clearly remained when their

presence was removed by acute NASPM application. We do not think that there was any

technical issue with the NASPM application, because overall, mEPSC amplitudes were reduced

modestly in both untreated and TTX-treated cultures (Figure 3B, CON, before NASPM, 12.9 ±

3.5 pA; after NASPM, 11.9 ± 2.6 pA; TTX, before NASPM, 17.5 ± 3.1 pA; after NASPM, 16.1 ± 3.5

pA). Furthermore, NASPM consistently reduced mEPSC frequency (Figure 3C, CON, before

NASPM, 1.84 ± 0.55 sec^{-1}; after NASPM, 1.56 ± 0.53 sec^{-1}; TTX, before NASPM, 4.40 ± 3.51 sec^{-1};

after NASPM, 2.68 ± 2.25 sec^{-1}). Similar to the data in Figure 1, frequency is significantly

increased after TTX treatment (p = 0.04), but this difference is no longer significant after

NASPM application (p = 0.20). One explanation for the effect on frequency is that there are

synaptic sites that express only homomers of Ca^{2+}-permeable AMPA receptors (GluA2-lacking

AMPARs); mEPSCs from these sites would be expected to be completely blocked by NASPM

(see cartoon description in Figure 3D, Left). We cannot rule out an alternative explanation, that

presynaptic Ca^{2+}-permeable AMPA receptors normally enhance release probability, and NAPSM

prevents this action. The magnitude of the frequency reduction following acute NASPM appears

to be greater after TTX treatment, suggesting that loss of activity could promote the

establishment of synaptic sites that contain only Ca^{2+}-permeable receptors. However, these

new sites do not appear to contribute to the increase in mEPSC amplitude (Figure 3A). The lack

of effect of NASPM rules out the contribution of Ca^{2+}-permeable AMPA receptors GluA1, GluA3

and GluA4 (Hollmann et al., 1991; Burnashev et al., 1992; Jonas and Burnashev, 1995), as well

as Ca^{2+}-permeable forms of Kainate receptors GluA5 and GluA6 (Koike et al., 1997; Sun et al.,

2009) to homeostatic synaptic plasticity.

Having established that homomeric Ca^{2+}-permeable receptors are not contributing to

the homeostatic increase in mEPSC amplitude, we turned to immunofluorescence and confocal

imaging to assess whether GluA2 receptor expression, which will identify GluA2 homomers and

GluA2 heteromers, was increased in our wild type mouse cortical cultures following 48 hr

treatment with TTX. Since mEPSCs necessarily report synaptic levels of receptors, we used

VGLUT1-immunofluorescence to identify synapses on pyramidal primary apical dendrites

labeled with MAP-2 immunofluorescence. We focused on the primary dendrite of pyramidal

neurons as a way to reduce variability that might arise from being at widely ranging distances

from the cell body, or, from inadvertently sampling dendritic regions arising from inhibitory

neurons. In addition, it has been shown that the increase in receptors is most dramatic in this

region (Turrigiano et al., 1998). Figure 4 (Top) shows two pyramidal neurons, one from an

untreated culture on the left, and one from a TTX-treated culture on the right, prepared from

Rab3A$^{+/+}$ mice. To measure characteristics of synaptically located GluA2 receptor clusters, we

zoomed in on the primary dendrite (region in white rectangle). The zoomed regions for single

confocal sections of the selected area are shown below and contain pairs of VGLUT1- and

GluA2-immunofluorescent clusters indicated by white trapezoidal frames. A dendrite typically

required ~10 confocal sections to be fully captured, and the total number of synaptic pairs for

all the sections imaged on a dendrite averaged 20 (20.4 $\pm$ 6.5 (SD); range, 11-38) so this is an

atypically high number of pairs within a single section; these particular dendrites and sections

were selected for illustration purposes. In addition to the synaptic pairs, we observed many

GluA2-immunofluorescent clusters not associated with VGLUT1 immunofluorescence (a few are

indicated with white arrows), most probably extrasynaptic receptors. There were also GluA2-

positive clusters present outside of MAP2-positive dendrites, which may be located on

astrocytes (Fan et al., 1999). Finally, GluA2 immunofluorescent clusters close to VGLUT1

immunofluorescence but not located along any apparent MAP2-positive neurites suggests the

presence of axon-axonal contacts, although VGLUT1 has also been detected in astrocytes

(Ormel et al., 2012). Only sites that contained both VGLUT1 and GluA2 immunofluorescence

close to the primary MAP2-postive dendrite or a secondary branch along this primary dendrite

were selected for analyses. In the experiment from which these images were collected, we

analyzed the distance from the cell body for each synapse. The average distance from the cell

body, for dendrites from the untreated cell, was 38.5 ± 2.8 μm; for the TTX-treated cell, it was

42.4 ± 3.2 μm (p = 0.35, Kruskal-Wallis test). Since the greatest distance from the cell body is

set by the outer limit of the zoom window, and that window was placed adjacent to the cell

body, distances in other experiments would be in this same range.

Variability in the magnitude of the homeostatic response from culture to culture is

averaged out in physiological experiments by the pooling of data from many cells across many

cultures. To reduce the necessity for many cultures, we chose to pair experiments in the same

cultures by recording mEPSCs from one set of coverslips, and processing another set of

coverslips from the same culture for immunofluorescence. We also wanted to determine if the

homeostatic response of mEPSC amplitudes varied from culture to culture, and if so, whether

the GluA2 receptor levels varied in parallel. We completed this matched paradigm of physiology

and immunofluorescence for 3 cultures prepared from Rab3A$^{+/+}$ mice. We first present the

results as means for each culture, for control and TTX-treated dishes, for mEPSC data (Figure

5A) and imaging data (Figure 5B). Levels of GluA2 immunofluorescence at synaptic sites were

quantified by the size of the GluA2-immunofluorescent receptor cluster, because this showed

the greatest response to activity blockade; the average intensity value, and the integral of

intensity values across the area of the cluster, are included in a summary of the data in Table 1.

Three of three cortical cultures showed an increase in mean mEPSC amplitude after TTX

treatment. In contrast, one of the same three cultures showed a decrease in mean GluA2

receptor cluster size. Therefore, the decrease in receptor levels in Culture #3 was due to

something other than a lack of homeostatic effect on mEPSCs in that culture. Because there

was a clear trend towards increased GluA2 receptor cluster size, but the receptor response

appears more variable, we performed immunofluorescence measurements on two additional

cortical cultures, both of which showed modest increases after TTX treatment (Figure 5C). We

confirmed that there was not an increase in GluA1 receptor levels in two matched cultures that

showed robust homeostatic increases in mEPSC amplitudes (Supplementary Figure 1A, culture

mean mEPSC amplitudes; Supplementary Figure 1B, culture mean GluA1 receptor cluster sizes).

We next pooled data from the multiple cultures to examine effects of activity blockade

on GluA2 receptors at the level of cell means. Because the average number of synaptic sites

identified in the primary dendrites in three cultures was 20.4, we also took 20 mEPSC samples

from each cell, the 21^{st} to the 40^{th} mEPSC recorded, to minimize any differences in variability

between GluA2 receptors and mEPSCs that might be due to a difference in sampling. In the

data pooled from the 3 matched experiments, neurons from Rab3A$^{+/+}$ cultures showed a

significant increase in mean mEPSC amplitudes following activity blockade (Figure 5D; Table 1).

This result indicates that the homeostatic response averaged across the 3 cultures was very

similar to, but slightly smaller than, that of the data set presented in Figure 1. The mean for size

of GluA2 immunofluorescent receptor clusters showed a trend towards higher values after

activity blockade (18.1%) that was of similar magnitude to the mEPSC amplitude increase

(19.7%), but did not reach statistical significance for 3 cultures (Figure 5E) or 5 cultures (Figure

5F; Table 1). As expected, mean mEPSC amplitude was not increased following activity

blockade in the data pooled from a new set of 3 Rab3A$^{-/-}$ cultures (Table 1). For

immunofluorescence results from the same Rab3A$^{-/-}$ cultures, no trend towards higher values

was apparent for GluA2 receptor cluster characteristics (Table 1). However, given the variability

of the GluA2 response in cultures from Rab3A$^{+/+}$ mice, it would require a much higher number

of Rab3A$^{+/+}$ and Rab3A$^{-/-}$ cultures to make any firm conclusions about the role of Rab3A in the

homeostatic response of GluA2 receptors.

Having used VGLUT1 immunofluorescence to mark synaptic sites, we could also ask

whether there was a presynaptic effect of activity blockade on the size, intensity or integral of

the VGLUT1 signal at sites apposed to the previously examined GluA2 postsynaptic sites. In the

pooled data from cultures prepared from Rab3A$^{+/+}$ mice, we found no change in the size of the

VGLUT1-positive regions, and trends towards reduced intensity and integral (Table 2). These

data do not support the idea that activity blockade caused an increase in the amount of

VGLUT1 per vesicle, although we cannot rule out that a simultaneous reduction in the number

of vesicles obscured the effect, since we did not have an independent label for synaptic

vesicles. In cultures prepared from Rab3A$^{-/-}$ mice, the size, intensity and integral of the VGLUT1-

positive regions did not significantly change (Table 2).

We have demonstrated here for the first time that the synaptic vesicle protein Rab3A

influences the homeostatic regulation of quantal amplitude. What are possible ways that

Rab3A could exert its effect? It has previously been shown that exogenous addition of TNFα to

hippocampal cultures causes an increase in surface expression of GluA1 receptors (although not

GluA2 receptors), and that the homeostatic increase in mEPSC amplitudes is abolished in

cultures prepared from the TNFα deletion mouse (Stellwagen et al., 2005; Stellwagen and

Malenka, 2006). Furthermore, neurons from TNFα$^{+/+}$ mice plated on glial feeder layers derived

from TNFα$^{-/-}$ mice fail to show the increase in mEPSC amplitude after TTX treatment, indicating

that the TNFα inducing the receptor increases following activity blockade comes from the glial

cells (Stellwagen and Malenka, 2006). It has recently been shown that the source of TNFα is

astrocytes rather than microglia (Heir et al., 2024). Rab3A has been detected in astrocytes

(Maienschein et al., 1999; Hong et al., 2016), so to determine whether Rab3A is acting via

regulating TNFα release from astrocytes, we performed experiments similar to those of

Stellwagen and Malenka (2006) (schema illustrated in Figure 6, Left). Under our culturing

conditions, the glial feeder layers would be expected to be composed mainly of astrocytes (Heir

et al., 2024). We compared the effect of activity blockade on mEPSC amplitudes recorded from

cortical neurons from Rab3A$^{+/+}$ mice plated on Rab3A$^{+/+}$ glial feeder layers (Figure 6A); neurons

from Rab3A$^{+/+}$ mice plated on Rab3A$^{-/-}$ glia (Figure 6B), and neurons from Rab3A$^{-/-}$ mice plated

on Rab3A$^{+/+}$ glia (Figure 6C). If Rab3A is required for TNFα release from glia, then Rab3A$^{+/+}$

neurons plated on Rab3A$^{-/-}$ glia should not show a homeostatic increase in mEPSC amplitude

after treatment with TTX, and any cultures with Rab3A$^{+/+}$ glia should have a normal homeostatic

response. We found the opposite result: mEPSC amplitudes increased dramatically in cultures

where Rab3A was present in neurons (WT on WT, CON, 13.3 ± 0.5 pA, TTX, 16.7 ± 1.2 pA; WT

on KO, CON, 13.3 ± 1.0 pA, TTX, 18.8 ± 1.4 pA), but the increase was greatly diminished in

cultures where Rab3A was present only in glia (KO on WT, CON, 15.2 ± 1.1 pA, TTX, 16.9 ± 0.7

pA). In the glial feeder layer culture, the loss of Rab3A in the neurons may have caused an

increase in mEPSC amplitude at baseline, similar to the Earlybird mutant in dissociated cultures,

but this difference was not significant (CON, WT on WT, 13.3 ± 0.5 pA; CON, KO on WT, 15.2 ±

1.1 pA, p = 0.23, Kruskal-Wallis test), so it is also possible this can be attributed to culture to

culture variability. Regarding mEPSC frequency, in cultures with Rab3A$^{+/+}$ neurons, frequency

trended upward after activity blockade, whereas in the culture with Rab3A$^{-/-}$ neurons,

frequency was already increased (although not significantly, CON, WT on WT, 2.54 $\pm$ 0.57 sec^{-1};

CON, KO on WT, 4.46 $\pm$ 1.21 sec^{-1}, p = 0.23, Kruskal-Wallis test), and trended downward after

activity blockade.

In summary, in the absence of Rab3A from glia, the homeostatic plasticity of mEPSC

amplitude following activity blockade was normal, whereas in the absence of Rab3A in neurons,

homeostatic plasticity was greatly diminished. This result makes it highly unlikely that Rab3A is

required for the release of TNFα, or another factor from glia, that induces a homeostatic

upregulation of postsynaptic receptors and thereby increases mEPSC amplitude following TTX

treatment. Neuronal Rab3A appears to mediate the homeostatic increase in mEPSC amplitude

following activity blockade.

Discussion

We found that homeostatic synaptic plasticity of mEPSC amplitude in dissociated mixed

cultures of mouse cortical neurons and glia behaved remarkably similar to the mouse NMJ in

response to loss of Rab3A function: in cultures from Rab3A$^{-/-}$ mice, the increase in mEPSC

amplitudes following prolonged network silencing by TTX was strongly diminished, and in

cultures from Rab3A$^{Ebd/Ebd}$ mice, basal mEPSC amplitude was increased compared to that of

wild-type cultures and was not further modified by TTX treatment. These results suggest that

normal function of the synaptic vesicle protein Rab3A is required for the homeostatic increase

of mEPSC amplitude in cortical cultures. At the NMJ, the effect of Rab3A on homeostatic

upregulation of mEPC amplitude was not mediated by upregulation of postsynaptic receptors.

In neurons, the increase in mEPSC amplitude following block of activity was not due to

increases in Ca^{2+}-permeable GluA1 homomers because acute NASPM application was unable to

reverse the TTX-induced increase in amplitude. Imaging of immunofluorescence of GluA2

receptors revealed the response was more variable than that of mEPSC amplitude. The simplest

explanation for our findings at both the NMJ and in neurons is that Rab3A regulates a

presynaptic contributor to alteration of quantal amplitude during homeostatic synaptic

plasticity.

Is the lack of homeostatic plasticity in the Earlybird mutant due to occlusion?

We found that mEPSC amplitude was increased in untreated cultures prepared from

Rab3A$^{Ebd/Ebd}$ mice, compared to mEPSC amplitude in cultures from wild type mice. If there is a

limit to how large an mEPSC can become, it is possible that the mutant Rab3A does not affect

homeostatic plasticity, only occludes it. Similar increases in mEPSC amplitude at baseline,

combined with a failure to increase further after activity blockade, were noted in cultures

prepared from mECP2, AKAP, Homer1a, and Arc/Arg3.1 deletion mice (Shepherd et al., 2006;

Hu et al., 2010; Xu and Pozzo-Miller, 2017; Sanderson et al., 2018) as well as in cultures treated

with ubiquitin proteasome inhibitors (Jakawich et al., 2010) or microRNA 186-5p (Silva et al.,

2019), suggesting the possibility of a generalized phenomenon in which increased mEPSC

amplitude is induced when homeostatic regulation is disrupted. Since we did not observe such

an increase in mEPSC amplitude at baseline in cultures from Rab3A$^{-/-}$ mice, it remains a

possibility that the point mutant may bind to novel partners, causing activities that would not

be either facilitated or inhibited by Rab3A. Still, it is a strong coincidence that the novel activity

of the mutant Earlybird Rab3A would affect mEPSC amplitude, the same characteristic that is

modulated by activity blockade in a Rab3A dependent manner. Although we cannot rule out

occlusion, an alternate interpretation is that the presence of the mutant protein mimics the

condition of activity blockade.

Homeostatic plasticity of postsynaptic receptors was less consistent than that of mEPSC

amplitude

While we found that the effect of TTX treatment on mEPSC amplitude and GluA2

receptor levels were of similar magnitude, the effect on GluA2 receptor levels was not

statistically significant for the pooled data sets acquired in the same cultures, nor in a larger

data set containing 5 cultures. In light of the extensive literature documenting increases in

GluA1 or GluA2 receptors, or both, after activity blockade, the less than convincing increase in

GluA2 receptor levels in the current work came as a surprise. The reason for the lack of

significance was that the increase in receptors was more variable. We are aware of three

possible explanations for greater variability. 1) The number of synaptic sites sampled was

smaller for the imaging data set. This was because we limited our analysis to synaptic receptors

on the primary dendrite, where the increase in sensitivity to exogenously applied glutamate

was shown to be greatest (see Figure 3 in (Turrigiano et al., 1998)). To address this issue, we

limited our sample size of mEPSCs in the matched experiments to the same number per cell as

the mean sample size per cell for imaging, but there was still a discrepancy in the effect of TTX

on mEPSCs vs. receptors. 2) The small sample of only 3 cultures was not enough to reveal the

increase in receptors. We attempted to address this by adding 2 more cultures. Although both

additional cultures showed increases in GluA2 receptor cluster size, the magnitudes of the

increases were not large, and when cells from these cultures were included in the pooled data

set, the p value was not substantially improved. The only remaining possibility is: 3) The

complexity involved in processing cultures for immunofluorescence introduces variability that

obscures the true effect. One can get around technical issues related to immunofluorescence

processing by performing Western blots, but then any changes detected could be due to

upregulation of extrasynaptic receptors. What remains unclear is why our data were impacted

by this but data of other studies were not. Some other studies have used the number of

synaptic sites or puncta as the sample size (for example, (Hou et al., 2008; Ibata et al., 2008;

Wang et al., 2019)). When we analyzed our data after including as few as 6 randomly selected

cluster size measurements per cell, a Kruskal-Wallis test gave highly significant p values for 3

cultures (p = 0.001, n = 174 sites (6 *29 cells)) and 5 cultures (p = 0.005, n = 294 sites (6 *49

cells)). We found only a single study that had comparable sample sizes for receptors and

mEPSC amplitudes, and the p value was < 0.01 for mEPSC amplitudes (n = 9 and 10 cells for

CON and TTX, respectively), but five times higher (< 0.05) for receptor data (n = 13 and 14 cells

for CON and TTX) (Xu and Pozzo-Miller, 2017). This single example is consistent with our

suggestion that measurements of changes in receptors after activity blockade are more variable

than measurements of mEPSC amplitudes. In sum, our data suggest that TTX treatment of

mouse cortical cultures consistently leads to an increase in mEPSC amplitudes, and sometimes

leads to an increase in synaptic GluA2 receptor cluster size. Increased receptors likely

contribute to increases in mEPSC amplitudes, but our current data, coupled with our previous

findings at the mouse neuromuscular junction, support the idea that there are additional

sources contributing to the homeostatic increase in quantal size.

Neuronal, not glial, Rab3A is required for homeostatic regulation of quantal amplitude

Astrocytic release of TNFα was shown to mediate the increase in mEPSC amplitudes in

prolonged TTX-treated cultures of dissociated hippocampal neurons and cultures of

hippocampal slices (Stellwagen and Malenka, 2006; Heir et al., 2024), but we found that loss of

Rab3A in glia did not disrupt the increase in mEPSC amplitude after activity blockade. Further,

we previously showed that the homeostatic increase in NMJ mEPC amplitude was completely

normal in the absence of TNFα (Wang et al., 2011). This leaves neuronal Rab3A as the potential

controller of homeostatic regulation of quantal amplitude.

Is Rab3A acting in the presynaptic cell to cause homeostatic increase in mEPSC amplitude?

Our current results support the surprising possibility that the presynaptic vesicle protein

Rab3A either regulates postsynaptic AMPARs or regulates a presynaptic contributor to

homeostatic synaptic plasticity. To our knowledge, there is no evidence localizing Rab3A to

postsynaptic dendritic sites. Other presynaptic molecules, such as SNARE proteins,

synaptotagmins, and NSF, have been identified in the postsynaptic compartment (Lledo et al.,

1998; Nishimune et al., 1998; Osten et al., 1998; Song et al., 1998; Araki et al., 2010; Kennedy et

al., 2010; Suh et al., 2010; Jurado et al., 2013; Hussain and Davanger, 2015; Gu et al., 2016; Wu

et al., 2017). A recent report implicates Rab3A in the localization of plasma membrane proteins,

including the EGF receptor, in rafts in HEK cells and Jurkat-T cells, which indicates Rab3A can be

found in other specialized locations besides the presynaptic nerve terminal (Diaz-Rohrer et al.,

2023). Without additional experiments targeting Rab3A deletion to the pre- or post-synaptic

compartment, we cannot rule out that Rab3A is regulating homeostatic plasticity in mouse

cortical cultures by acting postsynaptically on receptor trafficking. However, given our previous

finding of a lack of any increase in AChR after TTX treatment the NMJ (Wang et al., 2011), the

simplest interpretation is that in these two situations, Rab3A is required for an increase in the

presynaptic quantum rather than an increase in postsynaptic receptors.

The presynaptic quantum, or the amount of transmitter released during a single fusion

event, is an important contributor to quantal size. The amount of transmitter released during

vesicle fusion can be affected by the kinetics of the fusion pore opening (Chang et al., 2017).

We previously used amperometric measurements in mouse adrenal chromaffin cells to show

loss of Rab3A increased the occurrence of very small amplitude fusion pore feet (Wang et al.,

2008). Paired with our finding that there is an increase in the occurrence of very slow-rising,

abnormally shaped mEPCs at the neuromuscular junction of Rab3A$^{-/-}$ mice, our data suggest

that small synaptic vesicles may have a fusion pore step under certain circumstances (see also

(Chiang et al., 2018)). If activity blockade causes an increase in mEPSC amplitude due to a more

rapid opening of a fusion pore, or a larger fusion pore conductance, it is possible that Rab3A is

required for this modulation.

Another way to increase the presynaptic quantum is to increase levels of the transmitter

transporter (Daniels et al., 2004; De Gois et al., 2005; Wilson et al., 2005; Hartman et al., 2006).

However, we found that immunofluorescence for the glutamate transporter, VGLUT1, was not

increased after activity blockade under our experimental conditions. A similar lack of increase in

VGLUT1 levels, relative to synapsin, was previously observed in hippocampal cultures treated

with NBQX, an AMPA receptor blocker (see supplemental data, (Wilson et al., 2005)). Taken

together with our findings, it appears that there is not strong evidence that the amount of

VGLUT1 at synapses is a contributing factor to the increase in mEPSC amplitude in neuronal

cultures after activity blockade with TTX.

Finally, the amount of transmitter released by a vesicle could be increased due to larger

vesicle size. Larger vesicle diameter has been observed following activity disruption in multiple

paradigms, including ocular deprivation, cortical ablation, and immobilization (Kawana et al.,

1971; Lloret and Saavedra, 1975; Cheresharov et al., 1978). Loss of Rab3 family members is

associated with increased vesicle size (Riedel et al., 2002; Schonn et al., 2010). Finally, an

increase in volume has been shown to increase miniature endplate junctional current

amplitude at the Drosophila NMJ (Karunanithi et al., 2002). Rab3A could be required for an

activity-dependent increase in vesicle size, but the increase in diameter needed for a 25%

increase in transmitter, and therefore mEPSC amplitude, would be difficult to detect.

The number of molecules shown to be required for homeostatic synaptic plasticity of

quantal size in neuronal cultures may have surpassed ~20 (see (Koesters et al., 2024) for a

recent summary), but fall into two main categories: molecules involved in glutamate receptor

expression or trafficking, and signaling molecules. To our knowledge, this is the first evidence

477 that a presynaptic protein is participating in this process. Homeostatic synaptic plasticity is

478 becoming increasingly implicated in essential normal functions such as sleep (Tononi and Cirelli,

479 2014; Diering et al., 2017; Torrado Pacheco et al., 2021), and pathological conditions such as

480 epilepsy, neuropsychiatric disorders, Huntington's, and alcohol use disorder (Trasande and

481 Ramirez, 2007; Fernandes and Carvalho, 2016; Wang et al., 2017; Lovinger and Abrahao, 2018;

482 Smith-Dijak et al., 2019; Lignani et al., 2020; Suzuki et al., 2021; Kavalali and Monteggia, 2023).

483 Therefore, our identification of a novel Rab3A-dependent regulatory pathway in homeostatic

484 synaptic plasticity may have important therapeutic implications. Our findings also further

485 cement the importance of Rab3A in activity-dependent modulation of synaptic strength.

487 <u>Materials and Methods</u>

488 *Animals.* To determine the role of Rab3A in homeostatic plasticity in mouse cortical cultures,

489 we employed two distinct genetic mouse strains with altered Rab3A function. Rab3A deletion

490 mice were generated as follows: Rab3A$^{+/-}$ heterozygous mice were bred and genotyped as

491 previously described (Kapfhamer et al., 2002; Wang et al., 2008), and maintained as a colony

492 using multiple heterozygous breeding pairs. Homozygous Rab3A$^{+/+}$ or Rab3A$^{-/-}$ mouse pups

493 were obtained by breeding Rab3A$^{+/+}$ pairs and Rab3A$^{-/-}$ pairs, respectively, so that the pups

494 were the first-generation progeny of a homozygous mating. This protocol minimizes the

495 tendency of multiple generations of Rab3A$^{+/+}$ and Rab3A$^{-/-}$ homozygous breedings to produce

496 progeny that are more and more genetically distinct. Rab3A$^{Ebd/Ebd}$ mice were identified in an

497 EU-mutagenesis screen of C57BL/6J mice, and after a cross to C3H/HeJ, were backcrossed for 3

498 generations to C57BL/6J (Kapfhamer et al., 2002). Rab3A$^{+/Ebd}$ heterozygous mice were bred at

Wright State University and genotyped in a two-step procedure: 1. a PCR reaction (forward

primer: TGA CTC CTT CAC TCC AGC CT; reverse primer: TGC ACT GCA TTA AAT GAC TCC T)

followed by 2. a restriction digest with enzyme Bsp1286I (New England Biolabs) that

distinguishes the Earlybird mutant by its different base-pair products. Rab3A$^{+/-}$ mice were

backcrossed with Rab3A$^{+/+}$ mice from the Earlybird heterozygous colony for 11 generations in

an attempt to establish a single wild type strain, but differences in mEPSC amplitude and

adrenal chromaffin cell calcium currents persisted, likely due to genes that are close to the

Rab3A site, resulting in two wild type strains: 1. Rab3A$^{+/+}$ from the Rab3A$^{+/-}$ colony, and 2.

Rab3A$^{+/+}$ from the Rab3A$^{+/Ebd}$ colony.

Primary Culture of Mouse Cortical Neurons. Primary dissociated cultures of mixed neuronal and

glia populations were prepared as previously described (Hanes et al., 2020). Briefly, postnatal

day 0-2 (P0-P2) Rab3A$^{+/+}$, Rab3A$^{-/-}$ or Rab3A$^{Ebd/Ebd}$ neonates were euthanized by rapid

decapitation, as approved by the Wright State University Institutional Animal Care and Use

Committee, and brains were quickly removed. Each culture was prepared from the cortices

harvested from two animals; neonates were not sexed. Cortices were collected in chilled

Neurobasal-A media (Gibco) with osmolarity adjusted to 270 mOsm and supplemented with 40

U/ml DNAse I (ThermoFisher Scientific). The tissues were digested with papain (Worthington

Biochemical) at 20 U/ml at 37°C for 20 minutes followed by trituration with a sterile, fire-

polished Pasteur pipette, then filtered through a 100 µm cell strainer, and centrifuged at 1100

rpm for 2 minutes. After discarding the supernatant, the pellet was resuspended in room

temperature Neurobasal-A media (270 mOsm), supplemented with 5% fetal bovine serum for

glia growth, and 2% B-27 supplement to promote neuronal growth (Gibco), L-glutamine, and

gentamicin (ThermoFisher Scientific). Neurons were counted and plated at 0.15 * 10^6

cells/coverslip onto 12 mm coverslips pre-coated with poly-L-lysine (BioCoat, Corning). This

plating density produces a "high density" culture characterized by a complex mesh of neuronal

processes criss-crossing the field of view, completely filling the space between cell bodies (see

Figure 4). We found in preliminary experiments that an increase in mEPSC amplitude in mouse

cortical cultures was inconsistent when cell density was low, likely due to lower levels of

baseline activity, although it is a limitation of this study that we did not directly measure activity

levels in cultures prepared from Rab3A$^{+/+}$, Rab3A$^{-/-}$ or Rab3A$^{Ebd/Ebd}$ mice, and therefore do not

know if differences in results are due to differences in activity levels. While unlikely for Rab3A$^{+/+}$

and Rab3A$^{-/-}$ studies, given the mEPSC amplitude in untreated cultures was relatively small, it

remains possible that in Rab3A$^{Ebd/Ebd}$ studies, the larger mEPSC amplitude in untreated cultures

was due to a loss of activity in these cultures. The culture media for the first day (0 DIV) was the

same as the above Neurobasal-A media supplemented with FBS, B-27, L-glutamine, and

gentamicin, and was switched after 24 hours (1 DIV) to media consisting of Neurobasal-A (270

mOsm), 2% B-27, and L-glutamine without FBS to avoid its toxic effects on neuronal viability

and health (Stellwagen and Malenka, 2006). Half of the media was changed twice weekly and

experiments were performed at 13-14 DIV. Two days prior to experiments, tetrodotoxin (TTX)

(500 nM; Tocris), a potent Na$^+$ channel blocker, was added to some cultures to chronically

silence all network activity and induce homeostatic synaptic plasticity mechanisms, while

untreated sister cultures served as controls. Cultures prepared from mutant mice were

compared with cultures from wild-type mice from their respective colonies. Note that the

cultures comprising the Rab3A$^{+/+}$ data here are a subset of the data previously published in

Hanes et al., 2020. This smaller data set was restricted to the time period over which cultures

were also prepared from Rab3A$^{-/-}$ mice.

Preparation of Glial Feeder Layers. Glial feeder layers were prepared from the cortices of P0-P2

Rab3A$^{+/+}$ or Rab3A$^{-/-}$ mouse pups as described previously (Stellwagen and Malenka, 2006).

Briefly, cortices were dissected and cells were dissociated as described above. Cell suspensions

of mixed neuronal and glial populations were plated onto glass coverslips pre-coated with poly-

L-lysine in Dulbecco's Modified Eagle Media (ThermoFisher Scientific) supplemented with 5%

FBS (to promote glial proliferation and to kill neurons), L-glutamine, and gentamicin, and

maintained in an incubator at 37°C, 5% CO_2; cultures were maintained in this manner for up to

1 month to generate purely glial cultures (all neurons typically died off by 7 DIV). Culture media

was replaced after 24 hours, and subsequent media changes were made twice weekly,

replacing half of the culture media with fresh media. Feeder layers were not used for neuronal

seeding until all native neurons were gone and glia approached 100% confluency (visually

inspected).

Plating of Neurons on Glial Feeder Layers. Cortical neurons were obtained as described above.

The cell pellet obtained was resuspended in Neurobasal-A (osmolarity adjusted to 270 mOsm)

containing B27 (2%, to promote neuronal growth), L-glutamine, and 5-fluorodeoxyuridine (FdU,

a mitotic inhibitor; Sigma). Addition of FdU was used to prevent glial proliferation and

contamination of the feeder layer with new glia, promoting only neuronal growth on the feeder

layers (FdU-containing media was used for the maintenance of these cultures and all

subsequent media changes). Glial culture media was removed from the feeder layer cultures,

and the neuronal cell suspension was plated onto the glial feeder cultures. The culture strategy

used to distinguish the relative roles of neuronal and glial Rab3A is outlined in Figure 8, Left. At

1 DIV, all of the culture media was removed and replaced with fresh Neurobasal-A media

containing FdU described above, and half of the media was replaced twice per week for all

subsequent media changes. Cultures were maintained in a 37°C, 5% CO_2 incubator for 13-14

DIV.

Whole-Cell Voltage Clamp to Record mEPSCs. At 13-14 DIV, mEPSCs from TTX-treated and

untreated sister cultures of Rab3A$^{+/+}$ or Rab3A$^{-/-}$ neurons from the Rab3A$^{+/-}$ colony, or Rab3A$^{+/+}$

or Rab3A$^{Ebd/Ebd}$ neurons from the Rab3A$^{+/Ebd}$ colony, were recorded via whole-cell voltage clamp

to assess the role of Rab3A in homeostatic synaptic plasticity. Recordings were taken from

pyramidal neurons, which were identified visually by a prominent apical dendrite; images were

taken of all cells recorded from. Cells were continuously perfused with a solution consisting of

(in mM): NaCl (115), KCl (5), CaCl$_2$ (2.5), MgCl$_2$ (1.3), dextrose (23), sucrose (26), HEPES (4.2),

pH = 7.2 (Stellwagen and Malenka, 2006). On the day of recording, the osmolarity of the media

from the cultures was measured (normally 285 – 295 mOsm) and the perfusate osmolarity was

adjusted to match the culture osmolarity to protect against osmotic shock to the neurons. To

isolate glutamatergic mEPSCs, TTX (500 nM) and picrotoxin (50 μM), were included in the

perfusion solution to block action potentials and GABAergic currents, respectively. The NMDA

receptor antagonist, APV, was not included in the perfusion solution because in pilot studies, all

mEPSCs were blocked by 10 μM CNQX and 50 μm picrotoxin, demonstrating no APV-sensitive

mEPSCs were present (data not shown). Previous studies have described NMDA mEPSCs in

neuronal cultures, but recordings were performed in 0 Mg^{2+} (Watt et al., 2000; Sutton et al.,

2006). In our recordings, the presence of extracellular Mg^{2+} (1.3 mM) <u>and</u> TTX would mean that

the majority of NMDA receptors are blocked by extracellular Mg^{2+} at resting Vm. Patch

electrodes (3.5 – 5 MΩ) were filled with an internal solution containing (in mM): K-gluconate

(130), NaCl (10), EGTA (1), $CaCl_2$ (0.132), $MgCl_2$ (2), HEPES (10), pH = 7.2. Osmolarity was

adjusted to 10 mOsm less than the perfusion solution osmolarity, which we have found

improves success rate of achieving low access resistance recordings. Neurons were clamped at

a voltage of -60 mV using an Axopatch 200B patch-clamp (Axon Instruments), recorded from for

2 – 5 minutes, and data were collected with Clampex 10.0/10.2 (Axon Instruments).

NASPM application. The antagonist of Ca^{2+}-permeable AMPA receptors, N-naphthyl

acetylspermine (NASPM, 20 μM; Tocris), was applied during recordings in a subset of

experiments. NASPM is a synthetic analog of Joro Spider Toxin (JSTX) (Koike et al., 1997), and

chemicals in this family block Ca^{2+} permeable glutamate receptors (Blaschke et al., 1993;

Herlitze et al., 1993; Iino et al., 1996). The presence of the edited GluA2 subunit disrupts both

the Ca^{2+} permeability and the sensitivity to JSTX and related substances (Blaschke et al., 1993;

Bochet et al., 1994; Jonas and Burnashev, 1995). Therefore, 20 μM NASPM is expected to

completely block AMPA receptors containing GluA1, GluA3 or GluA4 subunits (Hollmann et al.,

1991; Burnashev et al., 1992), or the Kainate receptors containing GluA5 or GluA6 (Koike et al.,

1997; Sun et al., 2009), but be ineffective against any heteromer or homomer containing GluA2

subunits. Because the effect of the spider toxins and their analogues are use-dependent

(Herlitze et al., 1993; Iino et al., 1996; Koike et al., 1997), NASPM was applied with a

depolarizing high K^+ solution (25 mM KCl, 95 mM NaCl) which we expected to trigger release of

glutamate and opening of glutamate-activated receptors, allowing entry of NASPM into the

pore. Baseline recordings were performed for 2 minutes in our standard perfusate, followed by

perfusion of NASPM + High K$^+$ solution for 45 seconds, and then perfusion of a NASPM only

solution for 5 minutes, after which recording was recommenced for 5 minutes (because we

found in pilot experiments that frequency was reduced following NASPM application).

Analysis of mEPSCs. Miniature excitatory postsynaptic currents (mEPSCs) were manually

selected using Mini Analysis software (Synaptosoft), a standard program used for mEPSC event

detection and analysis. Records were filtered at 2 kHz using a low-pass Butterworth filter prior

to selection. The program detection threshold was set at 3 pA but the smallest mEPSC selected

was 4.01 pA. A fully manual detection process means that the same criterion ("does this *look*

like an mEPSC?") is applied to all events, not just the false positives or the false negatives,

which prevents the bias from being primarily at one end or the other of the range of mEPSC

amplitudes. It is important to note that when performing the MiniAnalysis process, the

researcher did not know whether a record was from an untreated cell or a TTX-treated cell.

Immunofluorescence, microscopy, and data analysis. Primary cultures of mouse cortical

neurons were grown for 13-14 DIV. Antibodies to GluA2 (mouse ab against N-terminal, EMD

Millipore) were added directly to live cultures at 1:40 dilution, and incubated at 37 °C in a CO_2

incubator for 45 minutes. Cultures were rinsed 3 times with PBS/5% donkey serum before being

fixed with 4% paraformaldehyde. After 3 rinses in PBS/5% donkey serum, cultures were

incubated in CY3-labeled donkey-anti-mouse secondary antibodies for 1 hour at room

temperature, rinsed in PBS/5% donkey serum, permeabilized with 0.2% saponin, and incubated

in chick anti-MAP2 (1:2500, AbCAM) and rabbit anti-VGLUT1 (1:4000, Synaptic Systems) for 1

hour at room temperature in PBS/5% donkey serum. After rinsing with PBS/5% donkey serum,

coverslips were incubated with 488-anti chick and CY5-anti rabbit secondary antibodies for 1

hour at room temperature, rinsed, blotted to remove excess liquid, and inverted on a drop of

Vectashield (Vector Labs). Coverslips were sealed with nail polish and stored at 4 °C for < 1

week before imaging. All secondary antibodies were from Jackson Immunoresearch and were

used at 1:225 dilution.

Coverslips were viewed on a Fluoview FV1000 laser scanning confocal microscope with a

60x oil immersion, 1.35 NA objective. Once a pyramidal neuron was identified and a single

confocal section of the cell acquired (see Top, Figure 4), Fluoview 2.1 software was used to

zoom in on the primary dendrite (5X) and a series of confocal sections were taken every 0.5

µm. Images were analyzed offline with ImagePro 6 (Cybernetics). The composite image was

used to locate synaptic sites containing both VGLUT1 and GluA2 immunoreactivity in close

apposition to each other and to the primary dendrite or a secondary branch identified with

MAP2 immunoreactivity. An area of interest (AOI) was manually drawn around the GluA2

cluster in the confocal section in which it was the brightest. The AOIs for a dendrite were saved

in a single file; the AOI number and the confocal section it was associated with were noted for

later retrieval. For quantification, AOIs were loaded, an individual AOI was called up on the

appropriate section, and the count/measurement tool used to apply a threshold. Multiple

thresholds were used, 400-550, but we found that within a given experiment, the magnitude of

the TTX effect was not altered by different thresholds, therefore, all data presented here use a

threshold of 450. Pixels within the cluster that were above the threshold were automatically

outlined, and size, average intensity, and integral of the outlined region saved to Origin2020

(OriginLab) file for calculating mean values. For analyzing VGLUT1 sites, previous AOIs were

examined, and if not directly over the VGLUT1 site, moved to encompass it (this happened

when GluA2 and VGLUT1 were side by side rather than overlapping). The threshold was 200 for

all VGLUT1 data except Rab3A$^{+/+}$ Culture #1, where threshold was 400. There was no threshold

that worked for both that culture and the other cultures, with too much of the background

regions being included for Culture #1 with a threshold of 200, and too little of the synaptic site

area being included for the other Cultures with a threshold of 400.

Statistics. In our previous publication (Hanes et al., 2020), we described the characteristics of

homeostatic plasticity of mEPSC amplitudes in mouse cortical cultures using multiple mEPSC

amplitude quantiles from each cell to create data sets with thousands of samples. We

compared the cumulative distribution functions for mEPSC amplitudes in untreated vs. TTX-

treated cultures, as well as sorting the values from smallest to largest, and plotted TTX values

vs. control values in the rank order plot first described in (Turrigiano et al., 1998) to

demonstrate synaptic scaling. Finally, we plotted the ratio of TTX values/control values vs.

control values to show that the ratio, which represents the effect of TTX, increased with

increasing control values, which we termed "divergent scaling." Here, we wanted to avoid the

inflation of sample size caused by pooling multiple mEPSCs per cell in order to statistically

compare the effects of TTX in cultures of wild type mice, Rab3A$^{-/-}$ mice and Rab3A$^{Ebd/Ebd}$ mutant

mice. Therefore, we computed the mean mEPSC amplitude for each cell and compared means

from untreated and TTX-treated cultures with the non-parametric Kruskal-Wallis test, with N =

the number of cells. The overall means are presented ± SEM in the text, and p values are

displayed above each plot. The same process was carried out for the characteristics of synaptic

GluA2 receptor clusters and VGLUT1 sites in imaging experiments, with N = the number of

dendrites (one dendrite was sampled for each cell). Although we quantified three

characteristics of the GluA2 clusters and VGLUT1 sites, size, average intensity, and integral, we

have not corrected p values for multiple comparisons. Means are presented as box plots + data,

with the box at 25 and 75 percentiles, whiskers at 10 and 90 percentiles, a dot indicating the

mean, and a line indicating the median. In Figure 3, mEPSC amplitudes were measured within

the same cell before and after NASPM application, so a paired t test was used to compare the

pre- and post-NASPM mEPSC amplitudes. In Figure 5A-C the mean mEPSC amplitude for

untreated and TTX-treated coverslips in the same culture are compared with a paired t test.

Statistics were computed and plots were created using Origin2020 (OriginLab).

References

Araki Y, Lin DT, Huganir RL (2010) Plasma membrane insertion of the AMPA receptor GluA2 subunit is
regulated by NSF binding and Q/R editing of the ion pore. Proc Natl Acad Sci U S A 107:11080-
11085.
Blaschke M, Keller BU, Rivosecchi R, Hollmann M, Heinemann S, Konnerth A (1993) A single amino acid
determines the subunit-specific spider toxin block of alpha-amino-3-hydroxy-5-methylisoxazole-
4-propionate/kainate receptor channels. Proc Natl Acad Sci U S A 90:6528-6532.
Bochet P, Audinat E, Lambolez B, Crepel F, Rossier J, Iino M, Tsuzuki K, Ozawa S (1994) Subunit
composition at the single-cell level explains functional properties of a glutamate-gated channel.
Neuron 12:383-388.
Burnashev N, Monyer H, Seeburg PH, Sakmann B (1992) Divalent ion permeability of AMPA receptor
channels is dominated by the edited form of a single subunit. Neuron 8:189-198.
Castillo PE, Janz R, Sudhof TC, Tzounopoulos T, Malenka RC, Nicoll RA (1997) Rab3A is essential for
mossy fibre long-term potentiation in the hippocampus. Nature 388:590-593.
Chang CW, Chiang CW, Jackson MB (2017) Fusion pores and their control of neurotransmitter and
hormone release. J Gen Physiol 149:301-322.
Cheresharov L, Ovtscharoff W, Manolov S (1978) Ultrastructural changes in the anterior horn synapses
of rat spinal cord under different locomotor conditions. J Neural Transm 42:9-21.
Chiang CW, Chang CW, Jackson MB (2018) The Transmembrane Domain of Synaptobrevin Influences
Neurotransmitter Flux through Synaptic Fusion Pores. J Neurosci 38:7179-7191.
Coleman WL, Bykhovskaia M (2009) Rab3a-mediated vesicle recruitment regulates short-term plasticity
at the mouse diaphragm synapse. Mol Cell Neurosci 41:286-296.
Coleman WL, Bykhovskaia M (2010) Cooperative regulation of neurotransmitter release by Rab3a and
synapsin II. Mol Cell Neurosci 44:190-200.
Coleman WL, Bill CA, Bykhovskaia M (2007) Rab3a deletion reduces vesicle docking and transmitter
release at the mouse diaphragm synapse. Neuroscience 148:1-6.
Daniels RW, Collins CA, Gelfand MV, Dant J, Brooks ES, Krantz DE, DiAntonio A (2004) Increased
expression of the Drosophila vesicular glutamate transporter leads to excess glutamate release
and a compensatory decrease in quantal content. J Neurosci 24:10466-10474.
De Gois S, Schafer MK, Defamie N, Chen C, Ricci A, Weihe E, Varoqui H, Erickson JD (2005) Homeostatic
scaling of vesicular glutamate and GABA transporter expression in rat neocortical circuits. J
Neurosci 25:7121-7133.
Diaz-Rohrer B, Castello-Serrano I, Chan SH, Wang HY, Shurer CR, Levental KR, Levental I (2023) Rab3
mediates a pathway for endocytic sorting and plasma membrane recycling of ordered
microdomains. Proc Natl Acad Sci U S A 120:e2207461120.
Diering GH, Nirujogi RS, Roth RH, Worley PF, Pandey A, Huganir RL (2017) Homer1a drives homeostatic
scaling-down of excitatory synapses during sleep. Science 355:511-515.
Dubes S, Soula A, Benquet S, Tessier B, Poujol C, Favereaux A, Thoumine O, Letellier M (2022) miR-124-
dependent tagging of synapses by synaptopodin enables input-specific homeostatic plasticity.
EMBO J 41:e109012.

Fan D, Grooms SY, Araneda RC, Johnson AB, Dobrenis K, Kessler JA, Zukin RS (1999) AMPA receptor
protein expression and function in astrocytes cultured from hippocampus. J Neurosci Res
57:557-571.
Fernandes D, Carvalho AL (2016) Mechanisms of homeostatic plasticity in the excitatory synapse. J
Neurochem 139:973-996.
Geppert M, Bolshakov VY, Siegelbaum SA, Takei K, De Camilli P, Hammer RE, Sudhof TC (1994) The role
of Rab3A in neurotransmitter release. Nature 369:493-497.
Gu Y, Chiu SL, Liu B, Wu PH, Delannoy M, Lin DT, Wirtz D, Huganir RL (2016) Differential vesicular sorting
of AMPA and GABAA receptors. Proc Natl Acad Sci U S A 113:E922-931.
Hanes AL, Koesters AG, Fong MF, Altimimi HF, Stellwagen D, Wenner P, Engisch KL (2020) Divergent
Synaptic Scaling of Miniature EPSCs following Activity Blockade in Dissociated Neuronal Cultures.
J Neurosci 40:4090-4102.
Hartman KN, Pal SK, Burrone J, Murthy VN (2006) Activity-dependent regulation of inhibitory synaptic
transmission in hippocampal neurons. Nat Neurosci 9:642-649.
Heir R, Abbasi Z, Komal P, Altimimi HF, Franquin M, Moschou D, Chambon J, Stellwagen D (2024)
Astrocytes Are the Source of TNF Mediating Homeostatic Synaptic Plasticity. J Neurosci 44.
Herlitze S, Raditsch M, Ruppersberg JP, Jahn W, Monyer H, Schoepfer R, Witzemann V (1993) Argiotoxin
detects molecular differences in AMPA receptor channels. Neuron 10:1131-1140.
Hollmann M, Hartley M, Heinemann S (1991) Ca2+ permeability of KA-AMPA--gated glutamate receptor
channels depends on subunit composition. Science 252:851-853.
Hong Y, Zhao T, Li XJ, Li S (2016) Mutant Huntingtin Impairs BDNF Release from Astrocytes by Disrupting
Conversion of Rab3a-GTP into Rab3a-GDP. J Neurosci 36:8790-8801.
Hou Q, Zhang D, Jarzylo L, Huganir RL, Man HY (2008) Homeostatic regulation of AMPA receptor
expression at single hippocampal synapses. Proc Natl Acad Sci U S A 105:775-780.
Hu JH, Park JM, Park S, Xiao B, Dehoff MH, Kim S, Hayashi T, Schwarz MK, Huganir RL, Seeburg PH,
Linden DJ, Worley PF (2010) Homeostatic scaling requires group I mGluR activation mediated by
Homer1a. Neuron 68:1128-1142.
Hussain S, Davanger S (2015) Postsynaptic VAMP/Synaptobrevin Facilitates Differential Vesicle
Trafficking of GluA1 and GluA2 AMPA Receptor Subunits. PLoS One 10:e0140868.
Ibata K, Sun Q, Turrigiano GG (2008) Rapid synaptic scaling induced by changes in postsynaptic firing.
Neuron 57:819-826.
Iino M, Koike M, Isa T, Ozawa S (1996) Voltage-dependent blockage of Ca(2+)-permeable AMPA
receptors by joro spider toxin in cultured rat hippocampal neurones. J Physiol 496 (Pt 2):431-
437.
Jakawich SK, Neely RM, Djakovic SN, Patrick GN, Sutton MA (2010) An essential postsynaptic role for the
ubiquitin proteasome system in slow homeostatic synaptic plasticity in cultured hippocampal
neurons. Neuroscience 171:1016-1031.
Jonas P, Burnashev N (1995) Molecular mechanisms controlling calcium entry through AMPA-type
glutamate receptor channels. Neuron 15:987-990.
Ju W, Morishita W, Tsui J, Gaietta G, Deerinck TJ, Adams SR, Garner CC, Tsien RY, Ellisman MH, Malenka
RC (2004) Activity-dependent regulation of dendritic synthesis and trafficking of AMPA
receptors. Nat Neurosci 7:244-253.
Jurado S, Goswami D, Zhang Y, Molina AJ, Sudhof TC, Malenka RC (2013) LTP requires a unique
postsynaptic SNARE fusion machinery. Neuron 77:542-558.
Kapfhamer D, Valladares O, Sun Y, Nolan PM, Rux JJ, Arnold SE, Veasey SC, Bucan M (2002) Mutations in
Rab3a alter circadian period and homeostatic response to sleep loss in the mouse. Nat Genet
32:290-295.

Karunanithi S, Marin L, Wong K, Atwood HL (2002) Quantal size and variation determined by vesicle size
776   in normal and mutant Drosophila glutamatergic synapses. J Neurosci 22:10267-10276.

777   Kavalali ET, Monteggia LM (2023) Rapid homeostatic plasticity and neuropsychiatric therapeutics.
778   Neuropsychopharmacology 48:54-60.

779   Kawana E, Akert K, Bruppacher H (1971) Enlargement of synaptic vesicles as an early sign of terminal
780   degneration in the rat caudate nucleus. J Comp Neurol 142:297-307.

781   Kennedy MJ, Davison IG, Robinson CG, Ehlers MD (2010) Syntaxin-4 defines a domain for activity-
782   dependent exocytosis in dendritic spines. Cell 141:524-535.

783   Koesters AG, Rich MM, Engisch KL (2024) Diverging from the Norm: Reevaluating What Miniature
784   Excitatory Postsynaptic Currents Tell Us about Homeostatic Synaptic Plasticity. Neuroscientist
785   30:49-70.

786   Koike M, Iino M, Ozawa S (1997) Blocking effect of 1-naphthyl acetyl spermine on Ca(2+)-permeable
787   AMPA receptors in cultured rat hippocampal neurons. Neurosci Res 29:27-36.

788   Leenders AG, Lopes da Silva FH, Ghijsen WE, Verhage M (2001) Rab3a is involved in transport of synaptic
789   vesicles to the active zone in mouse brain nerve terminals. Mol Biol Cell 12:3095-3102.

790   Lignani G, Baldelli P, Marra V (2020) Homeostatic Plasticity in Epilepsy. Front Cell Neurosci 14:197.

791   Lledo PM, Zhang X, Sudhof TC, Malenka RC, Nicoll RA (1998) Postsynaptic membrane fusion and long-
792   term potentiation. Science 279:399-403.

793   Lloret IL, Saavedra JP (1975) Enlargement of synaptic vesicles in degenerating nerve endings: a
794   comparison between cat and monkey. J Neurocytol 4:1-6.

795   Lonart G, Janz R, Johnson KM, Sudhof TC (1998) Mechanism of action of rab3A in mossy fiber LTP.
796   Neuron 21:1141-1150.

797   Lovinger DM, Abrahao KP (2018) Synaptic plasticity mechanisms common to learning and alcohol use
798   disorder. Learn Mem 25:425-434.

799   Maienschein V, Marxen M, Volknandt W, Zimmermann H (1999) A plethora of presynaptic proteins
800   associated with ATP-storing organelles in cultured astrocytes. Glia 26:233-244.

801   Nishimune A, Isaac JT, Molnar E, Noel J, Nash SR, Tagaya M, Collingridge GL, Nakanishi S, Henley JM
802   (1998) NSF binding to GluR2 regulates synaptic transmission. Neuron 21:87-97.

803   O'Brien RJ, Kamboj S, Ehlers MD, Rosen KR, Fischbach GD, Huganir RL (1998) Activity-dependent
804   modulation of synaptic AMPA receptor accumulation. Neuron 21:1067-1078.

805   Ormel L, Stensrud MJ, Bergersen LH, Gundersen V (2012) VGLUT1 is localized in astrocytic processes in
806   several brain regions. Glia 60:229-238.

807   Osten P, Srivastava S, Inman GJ, Vilim FS, Khatri L, Lee LM, States BA, Einheber S, Milner TA, Hanson PI,
808   Ziff EB (1998) The AMPA receptor GluR2 C terminus can mediate a reversible, ATP-dependent
809   interaction with NSF and alpha- and beta-SNAPs. Neuron 21:99-110.

810   Pozo K, Goda Y (2010) Unraveling mechanisms of homeostatic synaptic plasticity. Neuron 66:337-351.

811   Riedel D, Antonin W, Fernandez-Chacon R, Alvarez de Toledo G, Jo T, Geppert M, Valentijn JA, Valentijn
812   K, Jamieson JD, Sudhof TC, Jahn R (2002) Rab3D is not required for exocrine exocytosis but for
813   maintenance of normally sized secretory granules. Mol Cell Biol 22:6487-6497.

814   Sanderson JL, Scott JD, Dell'Acqua ML (2018) Control of Homeostatic Synaptic Plasticity by AKAP-
815   Anchored Kinase and Phosphatase Regulation of Ca(2+)-Permeable AMPA Receptors. J Neurosci
816   38:2863-2876.

817   Schluter OM, Basu J, Sudhof TC, Rosenmund C (2006) Rab3 superprimes synaptic vesicles for release:
818   implications for short-term synaptic plasticity. J Neurosci 26:1239-1246.

819   Schonn JS, van Weering JR, Mohrmann R, Schluter OM, Sudhof TC, de Wit H, Verhage M, Sorensen JB
820   (2010) Rab3 proteins involved in vesicle biogenesis and priming in embryonic mouse chromaffin
821   cells. Traffic 11:1415-1428.

Shepherd JD, Rumbaugh G, Wu J, Chowdhury S, Plath N, Kuhl D, Huganir RL, Worley PF (2006)
Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. Neuron 52:475-484.
Silva MM, Rodrigues B, Fernandes J, Santos SD, Carreto L, Santos MAS, Pinheiro P, Carvalho AL (2019)
MicroRNA-186-5p controls GluA2 surface expression and synaptic scaling in hippocampal
neurons. Proc Natl Acad Sci U S A 116:5727-5736.
Smith-Dijak AI, Nassrallah WB, Zhang LYJ, Geva M, Hayden MR, Raymond LA (2019) Impairment and
Restoration of Homeostatic Plasticity in Cultured Cortical Neurons From a Mouse Model of
Huntington Disease. Front Cell Neurosci 13:209.
Song I, Kamboj S, Xia J, Dong H, Liao D, Huganir RL (1998) Interaction of the N-ethylmaleimide-sensitive
factor with AMPA receptors. Neuron 21:393-400.
Stellwagen D, Malenka RC (2006) Synaptic scaling mediated by glial TNF-alpha. Nature 440:1054-1059.
Stellwagen D, Beattie EC, Seo JY, Malenka RC (2005) Differential regulation of AMPA receptor and GABA
receptor trafficking by tumor necrosis factor-alpha. J Neurosci 25:3219-3228.
Suh YH, Terashima A, Petralia RS, Wenthold RJ, Isaac JT, Roche KW, Roche PA (2010) A neuronal role for
SNAP-23 in postsynaptic glutamate receptor trafficking. Nat Neurosci 13:338-343.
Sun HY, Bartley AF, Dobrunz LE (2009) Calcium-permeable presynaptic kainate receptors involved in
excitatory short-term facilitation onto somatostatin interneurons during natural stimulus
patterns. J Neurophysiol 101:1043-1055.
Sutton MA, Ito HT, Cressy P, Kempf C, Woo JC, Schuman EM (2006) Miniature neurotransmission
stabilizes synaptic function via tonic suppression of local dendritic protein synthesis. Cell
125:785-799.
Suzuki K, Kim JW, Nosyreva E, Kavalali ET, Monteggia LM (2021) Convergence of distinct signaling
pathways on synaptic scaling to trigger rapid antidepressant action. Cell Rep 37:109918.
Tatavarty V, Sun Q, Turrigiano GG (2013) How to scale down postsynaptic strength. J Neurosci 33:13179-
13189.
Thiagarajan TC, Lindskog M, Tsien RW (2005) Adaptation to synaptic inactivity in hippocampal neurons.
Neuron 47:725-737.
Tian M, Xu CS, Montpetit R, Kramer RH (2012) Rab3A mediates vesicle delivery at photoreceptor ribbon
synapses. J Neurosci 32:6931-6936.
Tononi G, Cirelli C (2014) Sleep and the price of plasticity: from synaptic and cellular homeostasis to
memory consolidation and integration. Neuron 81:12-34.
Torrado Pacheco A, Bottorff J, Gao Y, Turrigiano GG (2021) Sleep Promotes Downward Firing Rate
Homeostasis. Neuron 109:530-544 e536.
Trasande CA, Ramirez JM (2007) Activity deprivation leads to seizures in hippocampal slice cultures: is
epilepsy the consequence of homeostatic plasticity? J Clin Neurophysiol 24:154-164.
Turrigiano GG, Nelson SB (2004) Homeostatic plasticity in the developing nervous system. Nat Rev
Neurosci 5:97-107.
Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB (1998) Activity-dependent scaling of
quantal amplitude in neocortical neurons. Nature 391:892-896.
Wang G, Zhong J, Guttieres D, Man HY (2019) Non-scaling regulation of AMPA receptors in homeostatic
synaptic plasticity. Neuropharmacology 158:107700.
Wang JKT, Langfelder P, Horvath S, Palazzolo MJ (2017) Exosomes and Homeostatic Synaptic Plasticity
Are Linked to Each other and to Huntington's, Parkinson's, and Other Neurodegenerative
Diseases by Database-Enabled Analyses of Comprehensively Curated Datasets. Front Neurosci
11:149.
Wang X, Thiagarajan R, Wang Q, Tewolde T, Rich MM, Engisch KL (2008) Regulation of quantal shape by
Rab3A: evidence for a fusion pore-dependent mechanism. J Physiol 586:3949-3962.

Wang X, Wang Q, Yang S, Bucan M, Rich MM, Engisch KL (2011) Impaired activity-dependent plasticity of
quantal amplitude at the neuromuscular junction of Rab3A deletion and Rab3A earlybird mutant
mice. J Neurosci 31:3580-3588.
Wang X, Li Y, Engisch KL, Nakanishi ST, Dodson SE, Miller GW, Cope TC, Pinter MJ, Rich MM (2005)
Activity-dependent presynaptic regulation of quantal size at the mammalian neuromuscular
junction in vivo. J Neurosci 25:343-351.
Watt AJ, van Rossum MC, MacLeod KM, Nelson SB, Turrigiano GG (2000) Activity coregulates quantal
AMPA and NMDA currents at neocortical synapses. Neuron 26:659-670.
Weisskopf MG, Castillo PE, Zalutsky RA, Nicoll RA (1994) Mediation of hippocampal mossy fiber long-
term potentiation by cyclic AMP. Science 265:1878-1882.
Wierenga CJ, Ibata K, Turrigiano GG (2005) Postsynaptic expression of homeostatic plasticity at
neocortical synapses. J Neurosci 25:2895-2905.
Wilson NR, Kang J, Hueske EV, Leung T, Varoqui H, Murnick JG, Erickson JD, Liu G (2005) Presynaptic
regulation of quantal size by the vesicular glutamate transporter VGLUT1. J Neurosci 25:6221-
6234.
Wu D, Bacaj T, Morishita W, Goswami D, Arendt KL, Xu W, Chen L, Malenka RC, Sudhof TC (2017)
Postsynaptic synaptotagmins mediate AMPA receptor exocytosis during LTP. Nature 544:316-
321.
Xu X, Pozzo-Miller L (2017) EEA1 restores homeostatic synaptic plasticity in hippocampal neurons from
Rett syndrome mice. J Physiol 595:5699-5712.

Figure 1. Loss of Rab3A disrupted the TTX-induced increase in amplitudes of mEPSCs recorded in cultured mouse cortical neurons; the increase in frequency was more variable but also appears to be reduced.

(A) Ten second example traces recorded at -60 mV in pyramidal cortical neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in cultures prepared from Rab3A$^{+/+}$ mice from the Rab3A$^{+/-}$ colony. (B) Ten second example traces recorded at -60 mV in pyramidal cortical neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in cultures

prepared from Rab3A$^{-/-}$ mice. (C & D) Average traces for the recordings shown in A and B, respectively. (E) Box plots for average mEPSC amplitudes from untreated cells and TTX-treated cells in cultures prepared from Rab3A$^{+/+}$ mice (Rab3A$^{+/-}$ colony; CON, N = 30 cells, 13.9 ± 0.7 pA; TTX, N = 23 cells, 18.2 ± 0.9 pA; from 11 cultures). (F) Box plots for average mEPSC amplitudes from untreated cells and TTX-treated cells in cultures prepared from Rab3A$^{-/-}$ mice (CON, N = 25 cells, 13.6 ± 0.7 pA; TTX, N = 26 cells, 14.3 ± 0.6 pA); from 11 cultures. (G) Box plots for average mEPSC frequency for same Rab3A$^{+/+}$ cells as in (E) (CON, 2.26 ± 0.37 sec^{-1}; TTX, 4.62 ± 0.74 sec^{-1}). (H) Box plots for average mEPSC frequency for same Rab3A$^{-/-}$cells as in (F) (CON, 2.74 ± 0.49 sec^{-1}; TTX, 3.23 ± 0.93 sec^{-1}). Box plot parameters: box ends, 25th and 75th percentiles; whiskers, 10th and 90th percentiles; open circles represent means from individual cells, bin size 1 pA, 0.5 sec^{-1}; line, median; dot, mean. p values (shown on the graphs) are from Kruskal-Wallis test. For all p-values, * with underline indicates significance with $p < 0.05$.

Figure 2. Normally functioning Rab3A was required for TTX-induced homeostatic plasticity of mEPSC amplitudes in cultured mouse cortical neurons.

(A) Ten second example traces recorded at -60 mV in pyramidal cortical neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in cultures prepared from Rab3A$^{+/+}$ mice from the Rab3A$^{+/Ebd}$ colony. (B) Ten second example traces recorded at -60 mV in pyramidal cortical neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in cultures

prepared from Rab3A$^{Ebd/Ebd}$ mice. (C & D) Average traces for the recordings shown in A and B, respectively. (E) Box plots for average mEPSC amplitudes from untreated cells and TTX-treated cells from cultures prepared from Rab3A$^{+/+}$ mice (Rab3A$^{+/Ebd}$ colony; CON, N = 20 cells, 11.0 ± 0.6 pA; TTX, N = 23 cells, 15.1 ± 1.2 pA; from 6 cultures). (F) Box plots for average mEPSC amplitudes from untreated cells and TTX-treated cells from cultures prepared from Rab3A$^{Ebd/Ebd}$ mice (CON, N = 21 cells, 15.1 ± 1.0 pA; TTX, N = 22 cells, 14.6 ± 1.1 pA; from 7 cultures.) (G) Box plots for average mEPSC frequency for same Rab3A$^{+/+}$ cells as in (E) (CON, 1.15 ± 0.19 sec^{-1}; TTX, 2.54 ± 0.55 sec^{-1}). (H) Box plots for average mEPSC frequency for same Rab3A$^{Ebd/Ebd}$ cells as in (F) (CON, 1.71 ± 0.41 sec^{-1}; TTX, 3.05 ± 0.80 sec^{-1}). Box plot parameters: box ends, 25th and 75th percentiles; whiskers, 10th and 90th percentiles; open circles represent means from individual cells; bin size 1 pA, 0.5 sec^{-1}; line, median; dot, mean. p values (shown on the graphs) are from Kruskal-Wallis test. For all p-values, * with underline indicates significance with $p < 0.05$.

Figure 3

Figure 3. Homeostatic plasticity of mEPSC amplitudes in mouse cortical cultures treated with TTX for 48 hr was unchanged by acute inhibition of Ca^{2+} permeable AMPA receptors by NASPM (20 μm).

(A) Box plot comparison of the TTX effect on mEPSC amplitudes in the same pyramidal neurons before and after application of 20 μM NASPM. (N = 11 cells from 3 cultures; Pre-NASPM, CON: 12.9 $\pm$ 1.1 pA; TTX: 17.5 $\pm$ 0.9 pA; post-NASPM, CON: 11.9 $\pm$ 0.8 pA; TTX: 16.1 $\pm$ 1.0 pA). (B) Line series plot of mEPSC amplitudes before and after acute perfusion with 20 μM NASPM for untreated and TTX-treated pyramidal neurons; same cells as in (A); CON, pre-NASPM: 12.9 $\pm$ 1.1 pA; post-NASPM: 11.9 $\pm$ 0.8 pA; TTX, pre-NASPM: 17.5 $\pm$ 0.9 pA; post-NASPM: 16.1 $\pm$ 0.1 pA. (C) Line series plot of mEPSC frequency before and after acute perfusion with 20 μM NASPM for untreated and TTX-treated pyramidal neurons; same cells as in (A); CON, pre-NAPSM: 1.84 $\pm$ 0.55 sec^{-1}; post-NASPM; 1.56 $\pm$ 0.53 sec^{-1}; TTX, pre-NASPM: 4.40 $\pm$ 1.06 sec^{-1}; post-NASPM, 2.68 $\pm$ 0.68 sec^{-1}. (D) A proposed mechanism for why NASPM had a robust effect on frequency without greatly affecting amplitude. A dendrite with spines (Top) is expanded on three postsynaptic sites (Middle) to show possible types of AMPA receptor distributions: Left, at a site comprised only of Ca^{2+}-permeable AMPA receptors (NASPM-sensitive, GluA2-lacking receptors (black)), NASPM would completely inhibit the mEPSC, causing a decrease in frequency to be measured in the overall population; Right, at a site comprised only of Ca^{2+}-impermeable AMPA receptors (NASPM-insensitive, GluA2-containing receptors (green)), NASPM would have no effect on mEPSC amplitude; Middle, at a site comprised of a mix of Ca^{2+}-permeable and Ca^{2+}-impermeable AMPA receptors, NASPM would partially inhibit the mEPSC. Diagram produced in BioRender.com (2024). Box plot parameters: box ends, 25th and 75th percentiles; whiskers, 10th and 90th percentiles; open circles represent means from individual cells; bin size 1 pA; line, median; dot, mean. p values (shown on the graphs) are from Kruskal-Wallis test. Line series plot p values are from student's paired t test. For all p-values, * with underline indicates significance with $p < 0.05$.

Figure 4. Identification of synaptic GluA2 receptor immunofluorescence on primary dendrites of pyramidal neurons in high density mouse cortical co-cultures prepared from Rab3A[+/+] mice.

(Top) Non-zoomed single confocal sections collected with a 60X oil immersion objective of recognizably pyramidal-shaped neurons, presumed to be excitatory neurons, selected for synaptic GluA2 analysis. A neuron was selected from an untreated coverslip (CON, Left) and a TTX-treated coverslip (TTX, Right) from the same culture prepared from Rab3A[+/+] animals (= Rab3A[+/+] Culture #2 in Figure 5). The white rectangular boxes indicate 5X zoomed areas shown in the images below. Note that the depth of the single confocal section of the non-zoomed neuron image is not at the same depth as the confocal section in zoomed dendritic image, so

some features are not visible in both images. Scale bar, 20 μm. (Bottom) 5X zoom single confocal sections selected for demonstration purposes because they had an unusually high number of identified synaptic pairs along the primary dendrite contained within a single confocal section. Synaptic pairs, highlighted with white trapezoids, were identified based on close proximity of GluA2 (red) and VGLUT1 (white) immunofluorescence, apposed to the MAP2 immunofluorescent primary dendrite (green). Some apparent synaptic pairs are not highlighted with a white trapezoid because there was a different confocal section in which the two immunofluorescent sites were maximally bright. There were GluA2 positive clusters located on the primary dendrites that were not apposed to VGLUT1-positive terminals (4 of these non-synaptic GluA2 clusters are highlighted with white arrows in the MAP2-GluA2 and MAP2-GluA2-VGLUT1 panels). Scale bar, 10 μm.

Figure 5. Comparison of mEPSC amplitudes and GluA2 receptor cluster areas in matched and unmatched mouse cortical cultures prepared from Rab3A$^{+/+}$ mice and treated with TTX for 48 hr.

(A) Culture averages of mEPSC amplitudes for untreated (CON) and TTX-treated coverslips (TTX) in each of 3 Rab3A$^{+/+}$ mouse cortical co-cultures. Culture #1, CON, N = 6, 14.2 ± 2.2 pA; TTX, N = 6, 15.9 ± 1.9 pA; Culture #2, CON, N = 7, 13.8 ± 2.4 pA; TTX, N = 8, 17.7 ± 1.8 pA; Culture #3, CON, N = 10, 13.4 ± 0.8 pA; TTX, N = 9, 15.7 ± 1.0 pA. (B) Culture averages of GluA2 receptor cluster size in the same 3 cultures as shown in (A). Culture #1, CON, N = 10, 0.73 ± 0.09 μm^2; TTX, N = 9, 0.89 ± 0.09 μm^2; Culture #2, CON, N = 9, 0.91 ± 0.12 μm^2; TTX, N = 9, 1.42 ± 0.24 μm^2; Culture #3, CON, N = 10, 1.05 ± 0.11 μm^2; TTX, N = 9, 0.95 ± 0.15 μm^2. (C) Two additional culture averages are included that did not have corresponding mEPSC recordings. Culture #4, CON, N = 10, 0.58 ± 0.07 μm^2: TTX, N = 10, 0.71 ± 0.04 μm^2; Culture 5, CON, N =10, 0.89 ± 0.05 μm^2; TTX, N = 10, 0.95 ± 0.08 μm^2. (D) Box plots of cell mean mEPSC amplitudes pooled from same 3 cultures as shown in (A). CON, N = 23 cells, 13.7 ± 0.9 pA; TTX, N = 24 cells, 16.4 ± 0.9 pA. (E) Box plots of dendrite mean GluA2 receptor cluster size pooled from same 3 cultures as shown in (A) and (B). CON, N = 29 dendrites, 0.90 ± 0.06 μm^2; TTX, N = 28 dendrites, 1.08 ± 0.10 μm^2. (F) Box plots of dendrite mean GluA2 receptor cluster size pooled from the same 5

cultures shown in (C). CON, N = 49 dendrites, 0.83 ± 0.04 μm^2; TTX, N = 48 dendrites, 0.98 ± 0.07 μm^2. Line series, Student's paired T test; box plots, Kruskal-Wallis test. Box plot parameters: box ends, 25th and 75th percentiles; whiskers, 10 and 90 percentiles; open circles indicate means of individual cells/dendrites; bin size 1 pA, 0.5 μm^2; line, median; dot, mean. For all p-values, * with underline indicates significance with $p < 0.05$.

.

Figure 6. Rab3A in neurons, not astrocytes, was required for full TTX-induced homeostatic plasticity.

(A)-(C), mEPSC amplitude (Middle) and frequency (Right) data from dissociated cortical neurons plated on an astrocyte feeder layer, each prepared separately from the type of mice depicted in the schema (Left): (A) Neurons from Rab3A$^{+/+}$ mice plated on astrocytes from Rab3A$^{+/+}$ mice. Box plots for mEPSC amplitudes (CON, N = 17 cells, 13.3 ± 0.5 pA; TTX, N = 20 cells, 16.7 ± 1.2 pA) and mEPSC frequency in the same recordings (mean, CON, 2.54 ± 0.57 sec^{-1}; TTX, 3.48 ± 0.64 sec^{-1}); from 4 cultures. (B) Neurons from Rab3A$^{+/+}$ mice plated on astrocytes from Rab3A$^{-/-}$ mice. Box plots for average mEPSC amplitude (CON, N = 11 cells, 13.3 ± 1.0 pA; TTX, N = 11 cells,

18.8 ± 1.4 pA) and average mEPSC frequency in the same recordings (CON, 2.01 ± 0.41 sec^{-1}; TTX, 4.47 ± 1.53 sec^{-1}); from 2 cultures. (C) Neurons from Rab3A$^{-/-}$ neurons plated on astrocytes from Rab3A$^{+/+}$ mice. Box plots for average mEPSC amplitude (CON, N = 14 cells, 15.2 ± 1.1 pA; TTX, N = 11 cells, 16.9 ± 0.7 pA) and mEPSC frequency in the same recordings (CON, 4.47 ± 1.21 sec^{-1}; TTX, 3.02 ± 0.70 sec^{-1}); from 3 cultures. Box plot parameters: box ends, 25th and 75th percentiles; whiskers, 10th and 90th percentiles; open circles represent means from individual cells; line, median; dot, mean. p values (shown on the graphs) are from Kruskal-Wallis test. p-values denoted with * and underline indicates significance of $p < 0.05$.

Supplemental Figure 1. Comparison of mEPSC amplitudes and GluA1 receptor cluster areas in matched mouse cortical cultures prepared from Rab3A$^{+/+}$ mice and treated with TTX for 48 hr.

(A) Culture averages of mEPSC amplitudes for untreated (CON) and TTX-treated coverslips (TTX) in each of 2 Rab3A$^{+/+}$ mouse cortical co-cultures. Culture #2, CON, N = 7, 13.8 $\pm$ 2.4 pA; TTX, N = 8, 17.7 $\pm$ 1.8 pA (+28.3%); Culture #6, CON, N = 6, 13.0 $\pm$ 1.5 pA; TTX, N = 6, 18.5 $\pm$ 2.0 pA (+42.3%). (B) Culture averages of GluA1 receptor cluster size in the same 2 cultures as shown in (A). Culture #2, CON, N = 10, 0.37 $\pm$ 0.04 μm^2; TTX, N = 10, 0.40 $\pm$ 0.06 μm^2 (+8.1%) Culture #6, CON, N = 10, 0.35 $\pm$ 0.05 μm^2; TTX, N = 10, 0.33 $\pm$ 0.04 μm^2 (-5.7%) "Culture #2" is the same Culture #2 depicted in Figure 5, different coverslips were processed for GluA1 immunofluorescence labeling. GluA1 receptor cluster intensity did not increase in Culture #2 (CON, 364 vs. TTX, 357) or Culture #6 (CON, 357 vs. TTX, 363). Live cultures were exposed to GluA1 antibody against the extracellular domain (ABN241, purchased from EMD Millipore, now available from Millipore Sigma) before being fixed and processed with secondary antibodies.

Table 1. Comparison of mEPSC amplitude and GluA2 receptor cluster characteristics in mouse cortical cultures prepared from Rab3A[+/+] mice, and Rab3A[-/-] mice.

Quantity	Rab3A[+/+]				Rab3A[-/-]			
	CON Mean, SE (N, cultures)	TTX Mean, SE (N, cultures)	% change	p value	CON Mean, SE (N, cultures)	TTX Mean, SE (N, cultures)	% change	p value
mEPSC amplitude	13.7 ± 0.9 pA (23 cells, 3 cultures)	16.4 ± 0.9 pA (24 cells, 3 cultures)	+19.7	*p = 0.02	14.9 ± 0.8 pA (21 cells, 3 cultures)	13.5 ± 0.9 pA (21 cells, 3 cultures)	-9.4	p = 0.18
GluA2 cluster size	0.83 ± 0.04 μm^2 (49 cells, 5 cultures)	0.98 ± 0.07 μm^2 (48 cells, 5 cultures)	+18.1	p = 0.21	0.93 ± 0.05 μm^2 (30 cells, 3 cultures)	0.91 ± 0.05 μm^2 (29 cells, 3 cultures)	-2.2	p = 0.75
GluA2 cluster intensity	733 ± 14 a.u. (49 cells, 5 cultures)	738 ± 13 a.u. (48 cells, 5 cultures)	+0.7	p = 0.69	766 ± 12 a.u. (30 cells, 3 cultures)	776 ± 15 a.u. (29 cells, 3 cultures)	+1.3	p = 0.46
GluA2 cluster integral	311021 ± 19282 a.u. (49 cells, 5 cultures)	365366 ± 27080 a.u. (48 cells, 5 cultures)	+17.4	p = 0.22	369436 ± 23439 a.u. (30 cells, 3 cultures)	364237 ± 25833 a.u. (29 cells, 3 cultures)	-0.8	p = 0.86

Table 2. Comparison of mEPSC amplitude and VGLUT1 positive presynaptic site characteristics in mouse cortical cultures prepared from Rab3A$^{+/+}$ and Rab3A$^{-/-}$ mice. mEPSC data are identical to that in Table 1, reproduced here for comparison purposes.

Quantity	Rab3A$^{+/+}$				Rab3A$^{-/-}$			
	CON Mean, SE	TTX Mean, SE	% change	p value	CON Mean, SE	TTX Mean, SE	% change	p value
mEPSC amplitude	13.7 ± 0.9 pA (23 cells, 3 cultures)	16.4 ± 0.9 pA (24 cells, 3 cultures)	+19.7	*p = 0.02	14.9 ± 0.8 pA (21 cells, 3 cultures)	13.5 ± 0.9 pA (21 cells, 3 cultures)	-9.4	p = 0.18
VGLUT1 site size	1.17 ± 0.08 µm^2 (29 cells, 3 cultures)	1.07 ± 0.06 µm^2 (24 cells, 3 cultures)	-7.7	p = 0.97	0.89 ± 0.03 µm^2 (30 cells, 3 cultures)	0.95 ± 0.04 µm^2 (29 cells, 3 cultures)	+6.7	p = 0.55
VGLUT1 site intensity	699 ± 39 a.u. (29 cells, 3 cultures)	639 ± 31 a.u. (28 cells, 3 cultures)	-8.6	p = 0.28	554 ± 13 a.u. (30 cells, 3 cultures)	570 ± 19 a.u. (29 cells, 3 cultures)	+2.9	p = 0.18
VGLUT1 site integral	430787 ± 36818 a.u. (29 cells, 3 cultures)	351653 ± 21689 a.u. (28 cells, 3 cultures)	-18.4	p = 0.13	257159 ± 18265 a.u. (30 cells, 3 cultures)	289857 ± 18521 a.u. (29 cells, 3 cultures)	+12.7	p = 0.21