1

Homeostatic Synaptic Plasticity of Miniature Excitatory Postsynaptic Currents in Mouse Cortical

Cultures Requires Neuronal Rab3A.

Andrew G. Koesters[1*], Mark M. Rich[2], and Kathrin L. Engisch[3]

[1]Department of Pharmacology and Systems Physiology, University of Cincinnati College of
Medicine, Cincinnati, OH 45267, USA
[2]Department of Neuroscience, Cell Biology and Physiology, Boonshoft School of Medicine,
Wright State University, Dayton, OH 45345
[3]Department of Neuroscience, Cell Biology and Physiology, Boonshoft School of Medicine and
the College of Science and Mathematics, Wright State University, Dayton, OH 45435
14

*Andrew G. Koesters –Corresponding author
Department of Pharmacology and Systems Physiology, University of Cincinnati College of
Medicine, Cincinnati, OH 45267, USA
koesteag@ucmail.uc.edu, ph 513-558-2819

Author contributions: A.G.K., K.L.E. and M.M.R. designed research; A.G.K. performed research;
A.G.K. and K.L.E. analyzed data; A.G.K, K.L.E., and M.M.R. wrote the paper.

ORCID iDs: ; A.G.K, https://orcid.org/0000-0003-3281-188X; M.M.R., https://orcid.org/0000-
0002-6956-5531; K.L.E., https://orcid.org/0000-0002-1058-5343

The authors have no competing interests.

Keywords: Homeostatic Synaptic Plasticity; Synaptic Scaling; mEPSCs; Rab3A; AMPA receptors

Abstract

Following prolonged activity blockade, amplitudes of miniature excitatory postsynaptic currents

(mEPSCs) increase, a form of homeostatic plasticity termed "synaptic scaling." We previously

showed that a presynaptic protein, the small GTPase Rab3A, is required for full expression of

the increase in miniature endplate current amplitudes following prolonged blockade of action

potential activity at the mouse neuromuscular junction in vivo (Wang et al., 2011), but it is

unknown whether this form of Rab3A-dependent homeostatic plasticity shares any

characteristics with central synapses. We show here that synaptic scaling of mEPSCs is impaired

in mouse cortical neuron cultures prepared from Rab3A$^{-/-}$ and Rab3A Earlybird mutant mice. To

determine if Rab3A is involved in the well-established homeostatic increase in postsynaptic

AMPA-type receptors (AMPARs), we performed a series of experiments in which

electrophysiological recordings of mEPSCs and confocal imaging of synaptic AMPAR

immunofluorescence were assessed within the same cultures. We found that Rab3A is required

for the increase in synaptic AMPARs following prolonged activity blockade, but the comparison

of mEPSC amplitude and synaptic AMPARs in the same cultures revealed that mEPSC amplitude

cannot solely be determined by postsynaptic AMPAR levels. Finally, we demonstrate that

Rab3A is acting in neurons because selective loss of Rab3A in astrocytes did not disrupt

homeostatic plasticity, whereas selective loss in neurons strongly reduced the homeostatic

increase in mEPSC amplitudes. Taken together with the results at the neuromuscular junction,

we propose that Rab3A is a presynaptic homeostatic regulator that controls quantal size on

both sides of the synapse.

     Introduction

One of the most studied phenomena triggered by prolonged activity blockade is the

increase in amplitudes of miniature excitatory postsynaptic currents (mEPSCs), first

demonstrated in cultures of dissociated cortical neurons (Turrigiano et al., 1998) and spinal

cord neurons (O'Brien et al., 1998). This compensatory response is now termed homeostatic

synaptic plasticity (Turrigiano and Nelson, 2004; Pozo and Goda, 2010) and has been shown to

be dysregulated in several neurodevelopment, psychiatric, and neurodegenerative disorders.

One of the initial studies of homeostatic synaptic plasticity demonstrated that AMPAergic

mEPSCs were uniformly increased after prolonged activity blockade (Turrigiano et al., 1998),

and the authors named this process synaptic scaling. An accompanying increase in AMPA-type

glutamate receptors (AMPARs) was observed in both of the early studies of homeostatic

synaptic plasticity (O'Brien et al., 1998; Turrigiano et al., 1998), and has been confirmed many

times (a non-exhaustive list includes (Ju et al., 2004; Thiagarajan et al., 2005; Shepherd et al.,

2006; Stellwagen and Malenka, 2006; Hou et al., 2008; Gainey et al., 2009; Soden and Chen,

2010; Correa et al., 2012; Altimimi and Stellwagen, 2013; Letellier et al., 2014; Xu and Pozzo-

Miller, 2017).

Homeostatic synaptic plasticity is becoming increasingly implicated in both pathological

brain conditions, for example, epilepsy, neuropsychiatric disorders, Huntington's, and alcohol

use disorder (Trasande and Ramirez, 2007; Fernandes and Carvalho, 2016; Wang et al., 2017;

Lovinger and Abrahao, 2018; Smith-Dijak et al., 2019; Lignani et al., 2020; Suzuki et al., 2021;

Kavalali and Monteggia, 2023), and essential normal functions such as sleep (Tononi and Cirelli,

2014; Diering et al., 2017; Torrado Pacheco et al., 2021), so a molecular understanding of the

process is an extremely important next step. The homeostatic synaptic plasticity field has

identified several proteins required for synaptic scaling of mEPSC amplitudes, the majority of

which are involved in the regulation of AMPAR levels (Shepherd et al., 2006; Seeburg and

Sheng, 2008; Gao et al., 2010; Anggono et al., 2011; Beique et al., 2011; Diering et al., 2014;

Gainey et al., 2015; Tan et al., 2015; Pastuzyn and Shepherd, 2017; Sanderson et al., 2018).

In our previous work studying the increase in miniature endplate currents (mEPC)

following prolonged in vivo activity-block of synaptic transmission at the mouse neuromuscular

junction (NMJ), we were surprised to find no evidence of changes in acetylcholine receptor

(AChR) levels (Wang et al., 2005). This result led us to search for presynaptic molecules that

might homeostatically regulate mEPC amplitude, perhaps via the presynaptic quantum. In

previous studies in chromaffin cells, we identified the small GTPase Rab3A, a synaptic vesicle

protein, as a regulator of synaptic vesicle fusion pore opening (Wang et al., 2008), so we

examined whether deletion of Rab3A (Rab3A$^{-/-}$) might prevent homeostatic upregulation of

mEPC amplitude. The results were clear: in the Rab3A$^{-/-}$ mouse, the homeostatic increase in

mEPC amplitude was strongly reduced, and was completely abolished in mice expressing a

point mutation in Rab3A, the Earlybird mutant (Rab3A$^{Ebd/Ebd}$) (Wang et al., 2011).

The Rab3A$^{-/-}$ mouse has minimal phenotypic abnormalities, with evoked synaptic

transmission and mEPSCs essentially normal in hippocampal slices (Geppert et al., 1994). At the

Rab3A$^{-/-}$ NMJ, reductions in evoked transmission were detected, but only under conditions of

reduced extracellular calcium (Coleman et al., 2007). The most dramatic effect of loss of Rab3A

is the disruption of a presynaptic form of long-term potentiation (LTP) at the mossy fiber-CA3

synapse (Weisskopf et al., 1994; Castillo et al., 1997). To our knowledge, there is no evidence of

Rab3A involvement in the expression or trafficking of postsynaptic receptors. Our results at the

mammalian NMJ suggest that in addition to its importance in mossy fiber LTP, Rab3A may be

required for the homeostatic plasticity of mEPSC amplitude via a presynaptic mechanism.

In the current work, we explored whether the in vivo findings at the NMJ might also

apply to the more typically studied homeostatic synaptic plasticity of mEPSC amplitude in

dissociated cortical neuron cultures. We report that our findings in vivo at the NMJ were almost

exactly recapitulated in cultures of dissociated cortical neurons: 1. strong reduction of the

homeostatic increase in mEPSC amplitude in the absence of Rab3A; 2. complete abolishment of

the homeostatic increase in mEPSC amplitude in the presence of the Rab3A Earlybird mutant;

and 3. increased mEPSC amplitude in the Rab3A Earlybird mutant prior to activity blockade.

However, in contrast to the unchanged AChR levels at the mammalian NMJ, there was a

modest increase in levels of the GluA2-type AMPARs at cortical synapses after activity-

blockade, which appeared to be disrupted in cultures prepared from Rab3A$^{-/-}$ mice.

Importantly, when compared within the same cultures, GluA2 receptor levels did not always

parallel mEPSC amplitudes. We also determined that Rab3A must be present in neurons, but

not astrocytes, for full expression of homeostatic plasticity. The NMJ and cortical culture data

taken together strongly suggest neuronal Rab3A is important for both postsynaptic receptor

upregulation and a second mechanism affecting mEPSC amplitude, likely the amount of

transmitter released by a single vesicle.

Materials and Methods

Animals. Rab3A$^{+/-}$ heterozygous mice were bred and genotyped as previously described

(Kapfhamer et al., 2002; Wang et al., 2008). Rab3A$^{Ebd/Ebd}$ mice were identified in an EU-

mutagenesis screen of C57BL/6J mice, and after a cross to C3H/HeJ, were backcrossed for 3

generations to C57BL/6J (Kapfhamer et al., 2002). Rab3A$^{+/Ebd}$ heterozygous mice were bred at

Wright State University and genotyped in a two-step procedure: 1. a PCR reaction with RabF1

and Dcaps3R as primers; and 2. a digestion with enzyme Bsp1286I (New England Biolabs) that

distinguishes the Earlybird mutant by its different base-pair products. Rab3A$^{+/-}$ mice were

backcrossed with Rab3A$^{+/+}$ mice from the Earlybird heterozygous colony for 11 generations in

an attempt to establish a single wild type strain, but differences in mEPSC amplitude and

adrenal chromaffin cell calcium currents persisted, likely due to genes that are close to the

Rab3A site, resulting in two wild type strains: 1. Rab3A$^{+/+}$ from the Rab3A$^{+/-}$ colony, and 2.

Rab3A$^{+/+}$ from the Rab3A$^{+/Ebd}$ colony.

Primary Culture of Mouse Cortical Neurons. Primary dissociated cultures of mixed neuronal and

astrocyte populations were prepared as previously described (Hanes et al., 2020). Briefly,

postnatal day 0-2 (P0-P2) Rab3A$^{+/+}$, Rab3A$^{-/-}$ or Rab3A$^{Ebd/Ebd}$ neonates were euthanized by rapid

decapitation, as approved by the Wright State University Institutional Animal Care and Use

Committee, and brains were quickly removed. Each culture was prepared from the cortices

harvested from two animals; neonates were not sexed. Cortices were collected in chilled

Neurobasal-A media (Gibco) with osmolarity adjusted to 270 mOsm and supplemented with 40

U/ml DNAse I (ThermoFisher Scientific). The tissues were digested with papain (Worthington

Biochemical) at 20 U/ml at 37°C for 20 minutes followed by trituration with a sterile, fire-

polished Pasteur pipette, then filtered through a 100 μm cell strainer, and centrifuged at 1100

rpm for 2 minutes. After discarding the supernatant, the pellet was resuspended in room

temperature Neurobasal-A media (270 mOsm), supplemented with 5% fetal bovine serum for

astrocyte growth, and 2% B-27 supplement to promote neuronal growth (Gibco), L-glutamine,

and gentamicin (ThermoFisher Scientific). Neurons were counted and plated at 0.15 * 10^6

cells/coverslip onto 12 mm coverslips pre-coated with poly-L-lysine (BioCoat, Corning). The

culture media for the first day (0 DIV) was the same as the above Neurobasal-A media

supplemented with FBS, B-27, L-glutamine, and gentamicin, and was switched after 24 hours (1

DIV) to media consisting of Neurobasal-A (270 mOsm), 2% B-27, and L-glutamine without FBS to

avoid its toxic effects on neuronal viability and health (Stellwagen and Malenka, 2006). Half of

the media was changed twice weekly and experiments were performed at 13-14 DIV. Two days

prior to experiments, tetrodotoxin (TTX) (500 nM; Tocris), a potent Na^+ channel blocker, was

added to some cultures to chronically silence all network activity and induce homeostatic

synaptic plasticity mechanisms, while untreated sister cultures served as controls. Cultures

prepared from mutant mice were compared with cultures from wild-type mice from their

respective colonies. Note that the cultures comprising the Rab3A$^{+/+}$ data here are a subset of

the data previously published in Hanes et al., 2020, and therefore the plots in Figure 1 are not

identical to those in the previously published work. This smaller data set was restricted to the

time period over which cultures were prepared from Rab3A$^{-/-}$ mice.

Preparation of Astrocyte Feeder Layers. Astrocyte feeder layers were prepared from the

cortices of P0-P2 Rab3A$^{+/+}$ or Rab3A$^{-/-}$ mouse pups as described previously (Stellwagen and

Malenka, 2006). Briefly, cortices were dissected and cells were dissociated as described above.

Cell suspensions of mixed neuronal and astrocyte populations were plated onto glass coverslips

pre-coated with poly-L-lysine in Dulbecco's Modified Eagle Media (ThermoFisher Scientific)

supplemented with 5% FBS (to promote astrocyte proliferation and to kill neurons), L-

glutamine, and gentamicin, and maintained in an incubator at 37°C, 5% CO_2; cultures were

maintained in this manner for up to 1 month to generate purely astrocytic cultures (all neurons

typically died off by 7 DIV). Culture media was replaced after 24 hours, and subsequent media

changes were made twice weekly, replacing half of the culture media with fresh media. Feeder

layers were not used for neuronal seeding until all native neurons were gone and astrocytes

approached 100% confluency (visually inspected).

Plating of Neurons on Glial Feeder Layers. Cortical neurons were obtained as described above.

The cell pellet obtained was resuspended in Neurobasal-A (osmolarity adjusted to 270 mOsm)

containing B27 (2%, to promote neuronal growth), L-glutamine, and 5-fluorodeoxyuridine (FdU,

a mitotic inhibitor; Sigma). Addition of FdU was used to prevent astrocyte proliferation and

contamination of the feeder layer with new astrocytes, promoting only neuronal growth on the

feeder layers (FdU-containing media was used for the maintenance of these cultures and all

subsequent media changes). Astrocyte culture media was removed from the feeder layer

cultures, and the neuronal cell suspension was plated onto the astrocyte feeder cultures. The

culture strategy used to distinguish the relative roles of neuronal and astrocytic Rab3A is

outlined in Figure 9. At 1 DIV, all of the culture media was removed and replaced with fresh

Neurobasal-A media containing FdU described above, and half of the media was replaced twice

per week for all subsequent media changes. Cultures were maintained in a 37°C, 5% CO_2

incubator for 13-14 DIV.

Whole-Cell Voltage Clamp to Record mEPSCs. At 13-14 DIV, mEPSCs from TTX-treated and

untreated sister cultures of Rab3A$^{+/+}$ or Rab3A$^{-/-}$ neurons from the Rab3A$^{+/-}$ colony, or Rab3A$^{+/+}$

or Rab3A$^{Ebd/Ebd}$ neurons from the Rab3A$^{+/Ebd}$ colony, were recorded via whole-cell voltage clamp

to assess the role of Rab3A in homeostatic synaptic plasticity. Recordings were taken from

pyramidal neurons, which were identified visually by a prominent apical dendrite; images were

taken of all cells recorded from. Cells were continuously perfused with a solution consisting of

(in mM): NaCl (115), KCl (5), CaCl$_2$ (2.5), MgCl$_2$ (1.3), dextrose (23), sucrose (26), HEPES (4.2),

pH = 7.2 (Stellwagen and Malenka, 2006). On the day of recording, the osmolarity of the media

from the cultures was measured (normally 285 – 295 mOsm) and the perfusate osmolarity was

adjusted to match the culture osmolarity, to protect against osmotic shock to the neurons. To

isolate glutamatergic mEPSCs, TTX (500 nM) and picrotoxin (50 μM) were included in the

perfusion solution to block action potentials and GABAergic currents, respectively. The NMDA

receptor antagonist, APV, was not included in the perfusion solution because all mEPSCs were

blocked by CNQX and picrotoxin, demonstrating no APV-sensitive mEPSCs were present (data

not shown). Patch electrodes (3.5 – 5 MΩ) were filled with an internal solution containing (in

mM): K-gluconate (128), NaCl (10), EGTA (1), CaCl$_2$ (0.132), MgCl$_2$ (2), HEPES (10), pH = 7.2.

Osmolarity was adjusted to 10 mOsm less than the perfusion solution osmolarity. Neurons

were clamped at a voltage of -60 mV using an Axopatch 200B patch-clamp (Axon Instruments),

recorded from for 2 – 5 minutes, and data were collected with Clampex 10.0/10.2 (Axon

Instruments). The antagonist of Ca^{2+}-permeable AMPA receptors (including GluA1 but not

GluA2), N-naphthyl acetylspermine (NASPM, 20 μM; Tocris), was applied during recordings in a

subset of experiments. Because NASPM is an open channel blocker, it was applied with a

depolarizing high K$^+$ solution (25 mM KCl, 95 mM NaCl). Baseline recordings were performed for

2 minutes in our standard perfusate, then were suspended while NASPM + High K$^+$ solution was

applied for 45 seconds, followed by a NASPM only solution for 5 minutes, after which recording

was recommenced for 5 minutes (because we found in pilot experiments that frequency was

reduced following NASPM application).

Data Analysis and Statistics of mEPSCs. Miniature excitatory postsynaptic currents were

manually selected using Mini Analysis software (Synaptosoft) to identify mEPSCs. The program

threshold was set at 3 pA but the smallest mEPSC selected was 4.01 pA. Records were filtered

at 2 kHz using a low-pass Butterworth filter prior to selection. For computing mEPSC mean in

each experimental condition, individual cell means were pooled across multiple cultures and

compared with the non-parametric Kruskal-Wallis test, with n = the number of cells and the

overall means presented as ± SEM. For cumulative probability distribution functions (CDFs) of

mEPSC amplitude, 30 quantiles were computed for each cell and pooled across cultures (see

(Hanes et al., 2020)), and a Kolmogorov-Smirnov test (KS test) was used to test for significant

differences, with n = the number of mEPSC quantiles. The rank order plots were created by

computing a matched number of quantiles from the two experimental conditions (usually

control (CON) and TTX). We used an algorithm to identify the product closest to, but above, 24

quantiles for a data set. For example, if there were 13 CON cells and 12 TTX cells, we computed

24 quantiles for each of the 13 cells and 26 quantiles for each of the 12 cells, for an equal total

number of 312 quantiles for both data sets. The quantiles were sorted from smallest to largest,

TTX amplitudes plotted vs. CON amplitudes, and the relationship fit with a linear regression

function with the intercept term allowed to vary. The ratio plots were created by taking the

ratio of TTX/CON and plotting as a function of the CON amplitude quantiles.

Immunocytochemistry, microscopy, and data analysis. Primary cultures of mouse cortical

neurons were grown for 13-14 DIV. Antibodies to GluA2 (mouse ab against N-terminal, EMD

Millipore) were added directly to live cultures at 1:40 dilution, and incubated at 37 °C in a CO_2

incubator for 45 minutes. Cultures were rinsed 3 times with PBS/5% donkey serum before being

fixed with 4% paraformaldehyde. After 3 rinses in PBS/5% donkey serum, cultures were

incubated in CY3-labeled donkey-anti-mouse secondary antibodies for 1 hour at room

temperature, rinsed in PBS/5% donkey serum, permeabilized with 0.2% saponin, and incubated

in chick anti-MAP2 (1:2500, AbCAM) and rabbit anti-VGLUT1 (1:4000, Synaptic Systems) for 1

hour at room temperature in PBS/5% donkey serum. After rinsing with PBS/5% donkey serum,

coverslips were incubated with 488-anti chick and CY5-anti rabbit secondary antibodies for 1

hour at room temperature, rinsed, blotted to remove excess liquid, and inverted on a drop of

Vectashield (Vector Labs). Coverslips were sealed with nail polish and stored at 4 °C for < 1

week before imaging. All secondary antibodies were from Jackson Immunoresearch and were

used at 1:225 dilution.

Coverslips were viewed on a Fluoview FV1000 laser scanning confocal microscope with a

60x oil immersion, 1.35 NA objective. Once a pyramidal neuron was identified, Fluoview 2.1

software was used to zoom in on the primary dendrite (5X) and confocal sections were taken

every 0.5 μm. Images were analyzed offline with ImagePro 6 (Cybernetics). The composite

image was used to locate synaptic sites containing both VGLUT1 and GluA2 immunoreactivity in

close apposition to each other and to the primary dendrite or a secondary branch. An area of

interest (AOI) was manually drawn around the receptor cluster in the confocal section in which

it was the brightest. The AOIs for a dendrite were saved in a single file; the AOI number and the

confocal section it was associated with were noted for later retrieval. For quantification, AOIs

were loaded, an individual AOI was called up on the appropriate section, and the

count/measurement tool used to apply a threshold (400 or 450; identical for CON and TTX

coverslips but different for different cultures); pixels within the cluster that were above the

threshold were automatically outlined, and size, average intensity, and integral of the outlined

region reported. For quantile sampling, only dendrites with a minimum of 6 synaptic sites were

included. For creating CDFs, 30 quantiles were computed for each dendrite. For the rank order

and ratio plots, we matched the total quantiles for CON and TTX GluA2 receptor characteristics

as was done above for mEPSC amplitudes.

<u>Results</u>

We previously reported that mixed cultures of cortical neurons and astrocytes prepared

from postnatal day 0-2 mouse pups responded to a block of action potential-mediated activity

by a 48 hour TTX treatment with an increase in mEPSC amplitude (Hanes et al., 2020). Here, we

asked the question whether cortical cultures prepared from mice lacking the small GTPase

Rab3A, or, expressing a point mutation of Rab3A, Rab3A Earlybird, have an altered homeostatic

plasticity response to a loss of network activity. To obtain Rab3A$^{-/-}$ and Rab3A$^{Ebd/Ebd}$

homozygotes, we established two mouse colonies of heterozygous breeders with cultures

prepared from pups derived from a final breeding pair of homozygotes. Although we

backcrossed Rab3A$^{+/-}$ with Rab3A$^{+/Ebd}$ for 11 generations, clear differences in mEPSC amplitudes

in untreated cultures (see below) and in calcium current amplitudes in adrenal chromaffin cells

(unpublished obs.) remained. Therefore, throughout this study we keep the two strains

separate and there are two Rab3A$^{+/+}$ or 'wild type' phenotypes.

Example current traces of spontaneously occurring mEPSCs recorded from pyramidal

neurons in untreated (CON) 13-14 DIV cortical cultures and sister cultures treated with 500 nM

TTX for 48 hours prepared from wild type animals in the Rab3A$^{+/-}$ colony are shown in Figure

1A. Average mEPSC waveforms from the same recordings are shown in Figure 1B. The mean

mEPSC amplitudes for 30 control and 23 TTX-treated neurons are displayed in the box and

whisker plot in Figure 1C; after activity blockade the average mEPSC amplitude increased from

13.9 ± 0.7 pA to 18.2 ± 0.9 pA (p = 4.58 * 10^{-4}, Kruskal-Wallis test). Example current traces of

mEPSCs and average mEPSC waveforms are shown in Figures 1D and E, respectively, for cortical

cultures prepared from Rab3A$^{-/-}$ mice. In contrast to the behavior of neurons in cultures

prepared from the wild type strain, the increase in mEPSC amplitudes in Rab3A$^{-/-}$ neurons after

activity blockade was dramatically reduced, and the average mEPSC amplitude was not

significantly increased, for 25 untreated cells and 26 TTX-treated cells (Figure 1F, 13.6 ± 0.1 vs.

14.3 ± 0.6, p = 0.318, Kruskal-Wallis test). To further examine homeostatic plasticity in the

presence and absence of Rab3A, we computed 30 quantiles for the mEPSC amplitude

distribution of each neuron (Hanes et al., 2020), pooled the quantiles, and plotted the data as

cumulative distribution functions (CDFs) for CON and TTX. For mEPSC amplitudes from cultures

prepared from Rab3A$^{+/+}$ mice, the difference between CDFs was highly significant (Figure 1Gi,

test statistic D = 0.172, p = 1.62 * 10^{-10}, Kolmogorov Smirnov (KS) test). In contrast, the increase

in mEPSC amplitude after activity blockade was dramatically reduced in cortical cultures from

Rab3A$^{-/-}$ mice (Figure 1Hi). A KS test for the CDFs was significant, but the test statistic much

smaller (0.070) and the p value much larger (0.042) than for the wild type strain.

290   It was originally proposed that loss of activity produced a uniform multiplicative

increase in mEPSC amplitude across the entire distribution of mEPSCs (Turrigiano et al., 1998).

This uniform scaling would be expected to preserve original differences in synaptic weights

resulting from other forms of plasticity (Turrigiano, 1999; Turrigiano and Nelson, 2004). In

addition, uniform scaling suggested a cell-wide mechanism (in fact, a culture-wide mechanism)

to identically modify all synapses. However, we recently showed that in mouse cortical

neurons, rat cortical neurons, and mouse hippocampal neurons, scaling is non-uniform, a

phenomenon we called "divergent scaling;" the multiplicative factor is smallest (close to 1) for

small mEPSC amplitudes and increases to ~1.4 for larger mEPSC amplitudes (Hanes et al., 2020).

In subsequent reviewing of the current literature (Koesters et al., 2022), we concluded that the

majority of previous studies of homeostatic plasticity of mEPSC amplitude following activity

blockade also show divergent scaling, most notably the study that originally defined synaptic

scaling (Turrigiano et al., 1998). Divergent scaling is not obvious on the standard plot used to

demonstrate synaptic scaling, the rank-order plot (Turrigiano et al., 1998), but is only apparent

once the ratio of TTX mEPSC amplitude/CON mEPSC amplitude is computed and plotted as a

function of CON mEPSC amplitude (Hanes et al., 2020). To compare between uniform and

divergent scaling in the presence and absence of Rab3A, we created both rank order plots and

ratio plots for Rab3A$^{+/+}$ data. The rank order plot for Rab3A$^{+/+}$ neurons (Figure 1Gii) shows that

treatment with TTX appears to cause a uniform multiplication of control mEPSC amplitudes

based on the linear fit to the data, which had a slope of 1.43 (R^2 = 0.982). However, in the ratio

plot for mEPSCs in Rab3A$^{+/+}$ cultures, the scaling was clearly divergent with the smallest mEPSC

amplitudes having the smallest ratios (at 1) and the ratio increasing with mEPSC amplitude to a

maximum of 1.51 (Figure 1Giii). In the absence of Rab3A, the rank ordered mEPSC data follow

the line of identity and the slope of the linear fit was 0.97 (R^2 = 0.985), suggesting complete

absence of homeostatic plasticity (Figure 1Hii). However, the more sensitive ratio plot reveals a

residual divergent scaling to a peak ratio of 1.13 (Figure 1Hiii). We previously observed a

residual homeostatic effect of activity blockade on mEPCs at the mouse NMJ in the absence of

Rab3A (Wang et al., 2011) suggesting that there may be compensatory or redundant

mechanisms after deletion of Rab3A.

We next examined whether expression of a single point mutant of Rab3A (Rab3A$^{Ebd/Ebd}$)

would abolish synaptic scaling in the mouse cortical cultures as we previously showed for

mEPCs at the mouse NMJ (Wang et al., 2011). Example current traces of mEPSCs, average

mEPSC waveforms, and box and whisker plots are shown in Figures 2A and B, respectively, for

cortical cultures prepared from wild type mice in the Rab3A$^{+/Ebd}$ colony. Treatment with TTX for

48 hours leads to a significant increase in the average mEPSC amplitude of 23 TTX-treated cells

when compared to 20 untreated cells (Figure 2C, CON, 11.0 ± 0.6 pA; TTX, 15.0 ± 1.3 pA, p =

0.02, Kruskal-Wallis test). We note here that while the two strains respond very similarly to

activity blockade, the mean mEPSC amplitude in untreated cultures was significantly different in

the two wild type strains, 13.9 ± 0.7 pA, wild type from Rab3A$^{+/-}$ colony; 11.0 ± 0.6 pA, wild type

from Rab3A$^{+/Ebd}$ colony (p = 0.004, Kruskal-Wallis test).

We found a complete disruption of homeostatic plasticity in cortical cultures prepared

from Rab3A$^{Ebd/Ebd}$ mice, as can been seen in viewing example mEPSC traces and average mEPSC

waveforms (Figures 2D and E, respectively). The lack of TTX effect was confirmed in a

comparison of mEPSC amplitude means for 21 untreated and 22 TTX-treated cells (Figure 2F,

CON, 15.1 ± 1.0 pA vs. TTX, 14.6 ± 1.1 pA, p = 0.81, Kruskal-Wallis test). For cultures prepared

from the wild type strain of the Earlybird heterozygote colony, the CDFs of the pooled quantiles

from control and TTX-treated cells were significantly different (Figure 2Gi, D = 0.177, p value =

2.88 * 10^{-9}, KS test), but for cultures prepared from Rab3A$^{Ebd/Ebd}$ mice, the CDF of mEPSC

amplitudes from TTX treated cells was shifted to the *left*, indicating a slight *reduction* after

activity blockade, which did not reach statistical significance (Figure 2Hi; D = 0.067, p = 0.097,

KS test). The rank ordered data indicate that the increased slope after TTX observed in the

wildtype data (Figure 2Gii, slope 1.73, R^2 = 0.993) was completely abolished in the Earlybird

data (Figure 2Hii, slope 1.03, R^2 = 0.989). Finally, in the wild type data, the ratio of TTX mEPSC

amplitude/CON mEPSC amplitude increases from 1.08 for the smallest amplitudes to over 1.60

for the largest amplitudes (Figure 2Giii), whereas for mEPSC amplitudes from Rab3A$^{Ebd/Ebd}$

cultures, the ratio of TTX/CON actually falls below 1 (minimum 0.9, Figure 2 Hiii).

Our results show that the homeostatic increase of mEPSC amplitude after activity

blockade is disrupted in both Rab3A$^{-/-}$ and the Rab3A$^{Ebd/Ebd}$ cortical neurons, strongly supporting

a crucial role of functioning Rab3A in the synaptic scaling process. However, it is important to

note that the disruption differs for Rab3A$^{-/-}$ and Rab3A$^{Ebd/Ebd}$. In the Rab3A$^{-/-}$ data set, mEPSCs

from untreated cultures were indistinguishable from mEPSCs from Rab3A$^{+/+}$ untreated cultures,

demonstrating loss of Rab3A has no impact on basal activity mEPSC amplitudes, but the

increase in mEPSC amplitudes after activity blockade was strongly diminished in Rab3A$^{-/-}$

neurons. In the Rab3A$^{Ebd/Ebd}$ data set, mEPSC amplitudes from untreated cultures were

significantly larger than those of untreated cultures from wild type mice, as can be seen in

Figure 3 for the CDFs (Figure 3A, D = 0.245; p = 1.52 * 10^{-16}, KS test) and the box and whisker

plot (Figure 3A Inset, Rab3A$^{+/+}$, 11.0 ± 0.7 pA, vs. Rab3A$^{Ebd/Ebd}$, 15.1 ± 1.0 pA p = 0.0027). The

linear fit of the rank ordered data of Rab3A$^{Ebd/Ebd}$ vs. wildtype mEPSCs had a slope of 1.63

(Figure 3B). In the ratio plot (Figure 3C), scaling was more uniform than in any other data set we

have examined so far, with values starting at 1.4, declining to 1.2, then rising again to 1.4 for

the majority of the data, instead of beginning near 1 and rising to 1.5, as happened for

homeostatic plasticity (Figure 1Giii). Given this distinct behavior, it cannot be stated that the

presence of mutated Rab3A causes the identical effect on mEPSC amplitudes as that of activity

blockade. It may be that the much longer time in the presence of the Rab3A Earlybird mutant

compared to the 48 hour TTX treatment leads to a stable increase in the smallest mEPSCs.

Alternatively, these are two distinct mechanisms of mEPSC amplitude augmentation that

occlude each other. In any case, the increase in mEPSC amplitude in cultures from Rab3A$^{Ebd/Ebd}$

mice is consistent with the increase in mEPC amplitude we observed at the Rab3A$^{Ebd/Ebd}$ NMJ

(Wang et al., 2011).

The small GTPase Rab3A is generally thought to function presynaptically to regulate

synaptic vesicle trafficking, possibly in an activity-dependent manner (Castillo et al., 1997;

Lonart et al., 1998; Leenders et al., 2001; Schluter et al., 2006; Coleman and Bykhovskaia, 2009;

Tian et al., 2012). In contrast, homeostatic plasticity of mEPSC amplitude has been attributed to

an increase in postsynaptic receptors on the surface of the dendrite (O'Brien et al., 1998;

Turrigiano et al., 1998). Is Rab3A required for the increase in surface AMPA-type glutamate

receptors that has been confirmed by multiple studies (see Introduction)? It should be noted

that at the NMJ in vivo we could find no evidence for an increase in AChRs after TTX block of

the sciatic nerve in vivo (Wang et al., 2005). The type of AMPA receptor that has been shown to

be increased, calcium-permeable GluA1-containing AMPA receptor or calcium-impermeable

GluA2-containing AMPA receptor, appears to depend on the experimental manipulation. For

example, block of activity by APV (NMDA antagonist) combined with TTX (inhibitor of Na

channels) induces an increase in mEPSC amplitude in dissociated hippocampal cultures that is

completely reversed by acute application of the Ca-permeable receptor-specific inhibitor

NASPM (Sutton et al., 2006), indicating the entirety of the homeostatic effect is due to those

receptors, likely GluA1. In contrast, the increase in mEPSC amplitude in mouse hippocampal

slice cultures induced by TTX alone was not affected by another GluA1-specific inhibitor,

philanthotoxin, suggesting mediation by either Ca-impermeable receptors or a presynaptic

effect on quantal size (Soden and Chen, 2010); see also, (Dubes et al., 2022). Because we used

TTX alone to block network activity, we expected that NASPM would not reverse the TTX-

induced increase in mEPSC amplitude in our mouse cortical cultures, and we found that this

was indeed the case. Figure 4A shows that the TTX-induced increase in mean mEPSC amplitude

was nearly identical in a set of 11 CON and 11 TTX-treated cells before and after NASPM

treatment (before NASPM, CON 12.9 ± 3.5 pA; TTX, 17.5 ± 3.1 pA, p = 0.009; after NASPM, CON

11.9 ± 2.6 pA; TTX 16.1 ± 3.5 pA, p = 0.006, Kruskal-Wallis test). Application of NASPM caused a

modest decrease in mEPSC amplitude in both untreated and TTX-treated cultures (Figure 4B,

CON, before NASPM, 12.9 ± 3.5 pA; after, 11.9 ± 2.6 pA, p = 0.08; TTX, before NASPM, 17.5 ± 3.1

pA; after, 16.1 ± 3.5 pA, p = 0.08, paired t test), and also decreased mEPSC frequency (Figure

4C, CON, before NASPM, 1.84 sec^{-1}; after NASPM, 1.56 sec^{-1}; p = 0.003, paired t-test; TTX,

before NASPM, 4.40 ± 3.51 mEPSCs sec^{-1}; after NASPM, 2.68 ± 2.25 mEPSCs sec^{-1}, p = 0.02,

paired t-test), indicating that effective concentrations of NASPM were reached and mEPSCs due

to AMPA receptors composed entirely of Ca-permeable subunits were abolished.

Having established that GluA1 receptors are not contributing to the homeostatic

increase in mEPSC amplitude, we turned to immunohistochemistry and confocal imaging to

assess whether GluA2 receptor expression was increased in our wild type mouse cortical

cultures following 48 hour treatment with TTX. Since mEPSCs necessarily report synaptic levels

of receptors, we used VGLUT1-positivity to identify synapses on pyramidal primary apical

dendrites labeled with MAP-2 immunofluorescence. Figure 5 shows 6 pairs of VGLUT1- and

GluA2-immunofluorescent clusters (white frames) along each of two primary dendrites, one in

an untreated cortical culture (CON, left), the other in a culture treated with TTX for 48 hours

(TTX, right), both from Rab3A$^{+/+}$ mice. A dendrite typically required ~10 confocal sections to be

fully captured, and the total number of synaptic pairs for all the sections imaged on a dendrite

was usually < 20, so this is an atypically high number of pairs within a single section; these

particular dendrites and sections were selected for illustration purposes. In addition to the

synaptic pairs, we observed many GluA2-immunoreactive clusters not associated with VGLUT1

immunoreactivity; those along the dendrites may be extrasynaptic receptors, and those outside

the dendrites may be on astrocytes (Fan et al., 1999). There are also GluA2 immunoreactive

clusters that are close to VGLUT1 immunoreactivity but are not located along any apparent

MAP2-positive neurite, suggesting axon-axonal contacts, although VGLUT1 has also been

detected in astrocytes (Ormel et al., 2012). Only sites that contained both VGLUT1 and GluA2

immunoreactivity close to the primary MAP2-postive dendrite or a secondary branch were

selected for analyses.

Variability in the magnitude of the homeostatic response from culture to culture is

averaged out in physiological experiments by the pooling of data from many cells across many

cultures. To reduce the necessity for many cultures, we chose to pair experiments in the same

cultures by recording mEPSCs from one set of coverslips, and processing another set of

coverslips from the same culture for immunohistochemistry. We completed this matched

paradigm of physiology and immunohistochemistry, which to our knowledge has never been

done before, for 3 cultures prepared from Rab3A$^{+/+}$ mice and 3 cultures prepared from Rab3A$^{-/-}$

mice. We present the results for mEPSC data and imaging data pooled across the 3 experiments

in Figure 6 (Rab3A$^{+/+}$), Figure 7 (Rab3A$^{-/-}$), and Table 1. Levels of GluA2 immunoreactivity at

synaptic sites were quantified by the size of the GluA2-positive receptor cluster and the average

intensity value of the receptor cluster. We analyzed the imaging data the same way we

analyzed mEPSC data, sampling 30 quantiles from each dendrite's data set, and calculating the

means, pooling quantiles across dendrites to create CDFs, and sorting quantiles from smallest

to largest to produce rank order and ratio plots.

In the data pooled from the 3 matched experiments, neurons from Rab3A$^{+/+}$ cultures

showed a significant increase in mean mEPSC amplitudes following activity blockade (Figure 6A

inset, CON, 13.7 ± 4.5 pA, n = 23; TTX, 16.4 ± 4.3 pA, n = 24; p = 0.016, Kruskal-Wallis test); a

significant shift of the TTX CDF to larger mEPSC amplitude values (Figure 6A, D = 0.162, p = 1.42

* 10^{-8}, KS test); a slope of 1.11 in the rank order plot (Figure 6B, R^2 = 0.98); and a mean ratio of

the 50th to 75th percentile of 1.24 in the ratio plot (Figure 6C). These data indicate that the

homeostatic response averaged across the 3 cultures was very similar to, but slightly smaller

than, that of the previous data set presented in Figure 1. The means for size and intensity of

GluA2 receptor clusters, while showing trends to higher values after activity blockade, were not

significantly different (Figure 6D inset, size, CON, 0.97 ± 0.38 μm^2; TTX, 1.15 ± 0.59 μm^2, p =

0.44; Figure 6G inset, intensity, CON, 673 ± 90, TTX, 687 ± 72, p = 0.25, Kruskal-Wallis test; Table

1). However, when comparing the CDFs, where the sample size is much greater (equal to

usually close to 30 quantiles * number of cells), the shifts to larger values for the TTX CDF did

reach statistical significance for GluA2 receptor cluster size (Figure 6D, D = 0.089, p = 0.002, KS

test) and intensity (Figure 6G, D = 0.120, p = 6.73 * 10^{-6}, KS test). In the rank order plot for

GluA2 receptor cluster sizes (Figure 6E), the slope was actually larger than that for mEPSC

amplitude (1.33), but for intensity (Figure 6H), the slope value was below 1 (0.90). The weaker

effects on synaptic GluA2 receptor levels relative to mEPSC amplitude suggest the possibility

that GluA2 receptors are not the sole determining factor in the increased mEPSC amplitude

following activity blockade. In addition, the plot of the ratio of TTX GluA2 receptor cluster

size/control size as a function of ranked control values showed a surprising deviation from the

divergent scaling we have observed for mEPSC amplitude (compare Figure 6C to Figure 6F). The

smallest GluA2 receptor clusters showed a TTX/CON ratio as high as 3, but the mean ratio

across the 50th to 75th percentile settled down to 1.11 within < 20 samples. This finding of larger

ratios for the smallest GluA2 receptor clusters is supported by a previous study that followed

the same fluorescently labeled postsynaptic sites over time following activity blockade. Wang

and colleagues found that the biggest increases after activity blockade occurred at the

smallest/dimmest synaptic sites (Wang et al., 2019). A similar inverse relationship was reported

for changes in individual fluorescently labeled postsynaptic sites followed over time under

normal activity conditions (Minerbi et al., 2009; Statman et al., 2014). The mismatch in ratio

values between the smallest mEPSC amplitudes and smallest GluA2 receptor cluster sizes may

be due to the inability of physiological assays to detect the mEPSCs coming from the smallest

synapses.

We proceeded to determine the homeostatic responses of mEPSC amplitude and GluA2

receptor levels in 3 cultures prepared from Rab3A$^{-/-}$ mice. As shown for the data set in Figure 1,

mean mEPSC amplitude was not increased following activity blockade in the data pooled from

this new set of 3 Rab3A$^{-/-}$ cultures (Figure 7A inset, CON 14.9 ± 3.8 pA, n = 21; TTX, 14.0 ± 4.0

pA, n = 19; p = 0.34, Kruskal-Wallis test). Notably, the TTX CDF was shifted to *smaller* mEPSC

amplitude values (Figure 7A, D = 0.072, p = 0.085, KS test), the slope of the rank order plot was

below 1 (Figure 7B, slope 0.93, R^2 = 0.997), and the mean ratio across the 50th to 75th percentile

was below 1 (Figure 7C, 0.96). Thus, the disruption of homeostatic plasticity of mEPSC

amplitude shown with the previous data sets was recapitulated in this data set. For

immunohistochemistry results from the same Rab3A$^{-/-}$ cultures, we found that mean GluA2

receptor cluster size was unchanged following activity blockade (Figure 7D inset, CON, 0.93 ±

0.27 μm^2, TTX, 0.91 ± 0.28 μm^2, p = 0.74, Kruskal-Wallis test), the TTX CDF was not significantly

shifted (Figure 7D, D = 0.045, p = 0.31, KS test), the slope on the rank order plot was 1.01

(Figure 7E), and the ratio hovered at or below 1 except for a small group of ratios at the high

end of the control values (Figure 7F). For intensity, mean values were not increased after

activity blockade (Figure 7G inset, CON, 766 ± 68, TTX, 776 ± 79, p = 0.47, Kruskal-Wallis test),

the CDFs were not significantly different (Figure 7G, D = 0.060, p = 0.080), the slope on the rank

order plot was 1.09 (Figure 7H), and the mean ratio value across the 50^{th} to 75^{th} percentile was

1.02 (Figure 7I). Taken together, these results indicate that the modest increases in GluA2

receptor cluster size and intensity following activity blockade observed in wild-type cultures do

not occur in the absence of Rab3A.

As noted above, the magnitude of the homeostatic effect on mEPSC amplitude

appeared to be more robust than that on receptor levels in cultures from Rab3A$^{+/+}$ mice.

Further evidence that there is not a one-to-one correspondence in the homeostatic response of

mEPSC amplitudes and GluA2 receptor levels was apparent when the individual experiments'

data were compared. The electrophysiology experiments were technically difficult because in

order to evaluate whether a particular culture displayed homeostatic plasticity of mEPSC

amplitude, we set a minimum requirement of recording from 6 cells per condition, meaning a

successful experiment required at least 12 (6 CON and 6 TTX) usable recordings (i.e. low holding

current, stable baseline, low noise, etc.) in one day. Ultimately, our sample size ranged from 6

to 10 cells per condition per culture. Figures 8A-C show the CDFs and ratio plots for 3 individual

Rab3A$^{+/+}$ cultures. Rank order plots are not included here because the ratio plot is more

sensitive than the rank order plot when determining effect magnitude. We found that the

mEPSC amplitude effects differed from the GluA2 cluster size effects at both the broad level

(increases vs decreases), and in the specific ways in which the effect was non-uniform. Broadly,

although two of the three Rab3A$^{+/+}$ experiments show a homeostatic increase in the TTX CDF

and a ratio of TTX/CON > 1 for both mEPSC amplitudes (Figures 8Ai,iii and 8Bi,iii) and synaptic

GluA2 receptor cluster sizes (Figures 8Aii,iv and 8Bii,iv), Culture #3 shows an increase in mEPSC

amplitude but a *decrease* in the GluA2 receptor cluster size, based on both the shifts in CDFs

(mEPSC, Figure 8Ci, vs GluA2 receptor cluster size, Figure 8Cii) and the majority of ratio values

(mEPSC, Figure 8Ciii, vs. GluA2 receptor cluster size, Figure 8Civ). Similarly, we found an obvious

mismatch between the mEPSC amplitude data and the GluA2 cluster size data for one of the

three Rab3A$^{-/-}$ experiments. For Culture #3, the mEPSC TTX CDF shows a shift to the left (Figure

8Fi), and the ratios were below 1 for control mEPSC amplitudes < 20 pA and above 1 for control

mEPSC amplitudes > 20 pA (Figure 8Fiii). In the same culture, GluA2 receptor cluster size CDF

shows a shift to the right after activity blockade (Figure 8Fii), and in the ratio plot, values were

at or above 1 throughout the range of control sizes (Figure 8Fiv).

Perhaps more striking than the broad mismatches, which could be attributed to the

small sample sizes, is the complete lack of correspondence of TTX/CON ratios in the mEPSC

amplitudes compared to GluA2 cluster sizes in the same cultures. In cultures from Rab3A$^{+/+}$

mice, mEPSC amplitudes show divergent scaling similar to what we have previously reported,

with the smallest CON amplitudes showing the smallest ratios, and the ratio increasing

monotonically to a plateau, or, a peak followed by a gradual decline, as CON mEPSC amplitude

becomes very large. In contrast, the very smallest GluA2 cluster sizes in Rab3A$^{+/+}$ cultures have

the largest increase, with ratios as high as 3 in the two cultures that showed a broad overall

increase. The ratios decline dramatically to values around 1.5, but then remain substantially

above 1 throughout the entire range of data. Interestingly, the dramatic increase in the size of

the smallest GluA2 clusters is absent in the two cultures from Rab3A$^{-/-}$ mice that broadly

showed no increase, suggesting this aspect is also dependent on Rab3A. It is not even possible

to cut off the ratios at a certain point, to address a threshold effect such as GluA2 cluster sizes

below a certain value not being measurable electrophysiologically, and get the ratios for the

remaining data to match up. Taken together, these data indicate that variation in homeostatic

effects on GluA2 cluster size is not responsible for the variation in homeostatic effects on

mEPSC amplitude.

In summary, our matched mEPSC and receptor cluster results indicate that: 1. activity

blockade resulted in an increase in mEPSC amplitudes and synaptic GluA2 receptors; 2. loss of

Rab3A disrupted both the increase in mEPSC amplitudes and the increase in GluA2 receptor

levels; and 3. The homeostatic effects on mEPSC amplitudes and the GluA2 cluster sizes within

the same cultures differed at both the broad level, and in the specific way the ratios were non-

uniform, making it difficult to conclude modulation of GluA2 cluster size is the sole determinant

of homeostatic increases in mEPSC amplitude. Although it remains a possibility that the

disparities in homeostatic effects on mEPSC amplitudes and GluA2 cluster size TTX/CON ratios

might arise due to a sampling error, given the small number of cells sampled in each individual

experiment ($\leq$ 10), it seems unlikely that differences arising from sampling error alone would be

so consistent. However, only when we are able to more easily sample larger numbers of cells,

or it becomes possible to measure mEPSCs and receptor levels in the same cells, will we be able

to resolve this question.

We have demonstrated here for the first time that the synaptic vesicle protein Rab3A

influences the homeostatic regulation of postsynaptic GluA2 receptors. What are possible ways

that Rab3A could exert a postsynaptic action? It has previously been shown that exogenous

addition of TNFα to hippocampal cultures causes an increase in surface expression of GluA1

receptors (although not GluA2 receptors), and that the homeostatic increase in mEPSC

amplitudes is abolished in cultures prepared from the TNFα deletion mouse (Stellwagen et al.,

2005; Stellwagen and Malenka, 2006). Furthermore, neurons from TNF$\alpha^{+/+}$ mice plated on

astrocytic feeder layers derived from TNF$\alpha^{-/-}$ mice fail to show the increase in mEPSC amplitude

after TTX treatment, indicating that the TNFα inducing the receptor increases following activity

blockade comes from the astrocytes (Stellwagen and Malenka, 2006). Rab3A has been detected

in astrocytes (Maienschein et al., 1999; Hong et al., 2016), so to determine whether Rab3A is

acting via regulating TNFα release from astrocytes, we performed experiments similar to those

of Stellwagen and Malenka (2006) (schema illustrated in Figure 9, left side). We compared the

effect of activity blockade on mEPSC amplitudes recorded from cortical neurons from Rab3A$^{+/+}$

mice plated on Rab3A$^{+/+}$ astrocytic feeder layers (Figure 9A); neurons from Rab3A$^{+/+}$ mice

plated on Rab3A$^{-/-}$ astrocytes (Figure 9B), and neurons from Rab3A$^{-/-}$ mice plated on Rab3A$^{+/+}$

astrocytes (Figure 9C). If Rab3A is required for TNFα release from astrocytes, then Rab3A$^{+/+}$

neurons plated on Rab3A$^{-/-}$ astrocytes should not show a homeostatic increase in mEPSC

amplitude after treatment with TTX, and any cultures with Rab3A$^{+/+}$ astrocytes should have a

normal homeostatic response. We found the opposite result: mEPSC amplitudes increased

dramatically in cultures where Rab3A was present in neurons (Figures 9A and B), but much

more modestly increased in cultures where Rab3A was present only in astrocytes (Figure 9C).

Interestingly, the homeostatic effects appear to be larger in the feeder layer cultures,

compared to the effects observed in our previous neuronal/astrocyte mixed cultures prepared

directly onto poly-L-lysine coated coverslips, but the ratio plots clearly show scaling was

divergent under both plating conditions (compare Figures 1Giii with Figures 9Aiii and Biii). We

noticed that in the astrocyte feeder layer cultures, and in the matched mEPSC amplitude and

GluA2 receptor measurement experiments, the mean mEPSC amplitude in the untreated

cultures prepared from Rab3A$^{-/-}$ mice was slightly, but not significantly, larger, compared to the

mean mEPSC amplitude for untreated cultures prepared from Rab3A$^{+/+}$ mice in the same

experiments (astrocyte feeder layer, Rab3A$^{+/+}$ neurons on Rab2A$^{+/+}$ astrocytes, 13.3 ± 0.5 pA,

Rab3A$^{-/-}$ on Rab3A$^{+/+}$ astrocytes, 15.2 ± 1.1 pA; matched mEPSC and receptor experiments,

Rab3A$^{+/+}$ 13.7 ± 4.5 pA, Rab3A$^{-/-}$ 14.9 ± 3.8 pA. It is therefore possible that loss of Rab3A, like

expression of the Rab3A Earlybird mutant, affects basal mEPSC amplitude, albeit to a lesser

extent. However, it could also be that these differences reflect random variation in mEPSC

amplitude from culture to culture.

In summary, in the absence of Rab3A from astrocytes, the divergent scaling of mEPSC

amplitude following activity blockade was completely normal, whereas in the absence of Rab3A

in neurons, scaling was greatly diminished. This result makes it highly unlikely that Rab3A is

required for the release of TNFα, or another factor from astrocytes, that induces a homeostatic

upregulation of postsynaptic receptors and thereby increases mEPSC amplitude following TTX

treatment. Neuronal Rab3A appears to mediate the homeostatic increase in mEPSC amplitude

following activity blockade.

Discussion

We found that homeostatic synaptic plasticity of mEPSC amplitude in dissociated mixed

cultures of mouse cortical neurons and astrocytes behaved remarkably similar to the mouse

NMJ in response to loss of Rab3A function: scaling up of mEPSC amplitude following prolonged

network silencing by TTX was strongly diminished in cultures from Rab3A[-/-], and in cultures

from Rab3A[Ebd/Ebd] mice, basal mEPSC amplitude was increased compared to that of wild-type

cultures and was not further modified following 48-hour treatment with TTX. These results

suggest that normal function of the presynaptic vesicle protein Rab3A is required for the

homeostatic scaling up of mEPSC amplitude in cortical cultures and at the NMJ in vivo.

Rab3A not likely to regulate receptor trafficking

We demonstrated that an increase in synaptic GluA2 receptors accompanies the

increase in mEPSC amplitude in cultures of dissociated mouse cortical neurons after 48-hour

TTX treatment. By examining the same cultures with both mEPSC and immunofluorescence

measurements, we found: 1. the effect on GluA2 receptor levels, whether measured as the size

of a synaptic cluster or the density of receptors in a cluster, was less robust than the effect on

mEPSC amplitudes; 2. the homeostatic effect on mEPSC amplitudes and GluA2 receptors

substantially differed from each other both broadly (mEPSC amplitudes could increase while

GluA2 cluster sizes decreased, and vice versa), and in the details of non-uniform scaling, one

example being that the smallest mEPSC amplitudes showed the least effects but the smallest

GluA2 clusters showed the greatest effects. Interestingly, there are other examples where the

effect of activity blockade by either TTX alone (Hu et al., 2010), TTX and AP5 (Letellier et al.,

2014), or DNQX (Blackman et al., 2012) on AMPAR levels did not match that of mEPSC

amplitudes, although in these cases, the increases in receptors were larger than the increases

of mEPSC amplitudes. The striking differences between TTX effects on AMPARs and mEPSC

amplitudes indicates that there may be another factor contributing to mEPSC amplitude

besides levels of AMPAR expression. One possibility is alterations in the type of AMPA receptor

expressed, since GluA1 homomers have much greater conductance than GluA2-containing

receptors (Oh and Derkach, 2005; Benke and Traynelis, 2019). Increases in mEPSC amplitudes

not accompanied by increases in receptor numbers have been attributed to such a switch (Hou

et al., 2015; Silva et al., 2019; Dubes et al., 2022).

In addition to finding a disparity between synaptic GluA2 receptor and mEPSC responses

to prolonged activity blockade, there are two other reasons we do not think Rab3A impacts

homeostatic synaptic plasticity solely through postsynaptic regulation of GluA2 receptor

trafficking to the plasma membrane. First, prolonged activity blockade with TTX at the NMJ led

to an increase in mEPC amplitude that was not accompanied by an increase in AChR levels, yet

it was dependent on Rab3A (Wang et al., 2011). Second, for Rab3A to modulate AMPAR

trafficking, it must be located in the postsynaptic dendrite. Multiple presynaptic molecules,

such as SNARE proteins, synaptotagmins, and NSF have been identified in the postsynaptic

compartment (Lledo et al., 1998; Nishimune et al., 1998; Osten et al., 1998; Song et al., 1998;

Araki et al., 2010; Kennedy et al., 2010; Suh et al., 2010; Jurado et al., 2013; Hussain and

Davanger, 2015; Gu et al., 2016; Wu et al., 2017), but to our knowledge, Rab3A has not.

Rab3A may regulate quantal size presynaptically

Based on our findings, the lack of postsynaptic Rab3A expression and the known

presynaptic expression of Rab3A, Rab3A may regulate mEPSC amplitude presynaptically by

modulating: a) the amount of transmitter transported into the vesicle, b) the size of the vesicle,

or c) fusion pore characteristics. mRNA for the glutamate transporter, VGLUT1, was increased in

rat cortical cultures after 48-hr treatment with TTX (De Gois et al., 2005), and VGLUT1 relative

to synapsin I in rat hippocampal cultures was increased after 48-hr treatment with AP-5 (Wilson

et al., 2005). This activity-dependent regulation of VGLUT1 was recently corroborated in a

proteome study (Dorrbaum et al., 2020), and it has also been shown that mEPSC amplitudes are

increased after overexpression of VGLUT at the Drosophila NMJ (Daniels et al., 2004) and in rat

hippocampal cultures (Wilson et al., 2005). These data suggest that increases in VGLUT1 after

activity-blockade could contribute to the homeostatic increase in mEPSC amplitude. Although

there is no direct evidence for an interaction between VGLUT1 and Rab3A, Rab3A levels are

reduced 50% in hippocampal extracts from mice lacking VGLUT1 (Fremeau et al., 2004). A

Rab3A-dependent modulation of presynaptic quantal size would explain the homeostatic

effects on mEPSC amplitudes that are not matched by changes in receptor levels.

Interestingly, Rab3A plays a role in regulating vesicle size. In adrenal chromaffin cells

from mice lacking all 4 Rab3 isoforms (A, B, C and D) or mice heterozygous for Rab3A and

lacking the B, C, and D isoforms (Rab3A$^{+/-}$BCD$^{-/-}$) with only 1 copy of Rab3A, there was a 20%

increase in large dense core vesicle (DCV) diameter that did not reach significance (p = 0.11)

(Schonn et al., 2010). Similarly, in the exocrine pancreas and parotid gland of the Rab3D$^{-/-}$, large

DCV diameter was increased by 22% (Riedel et al., 2002). This magnitude of change in diameter

would produce a greater than 80% increase in volume. Also, an increase in volume has been

shown to increase miniature endplate junctional current amplitude at the Drosophila NMJ

(Karunanithi et al., 2002). There have not been any reports that small synaptic vesicle size is

increased in Rab3$^{-/-}$ mice, but small synaptic vesicle size is regulated by activity: vesicle size

increased in retinotectal terminals within the optic tectum following enucleation in pigeon

(Cuenod et al., 1970), with similar results in cat and monkey (Lloret and Saavedra, 1975), in rat

caudate nucleus following cortical ablation (Kawana et al., 1971), and in rat ventral spinal cord

following immobilization (Cheresharov et al., 1978).

Finally, we previously showed that at the mouse NMJ, both loss of Rab3A and activity

blockade by TTX caused an increase in the frequency of abnormally slow rising and/or

prolonged duration mEPCs (Wang et al., 2011). We further demonstrated that the loss of Rab3A

increased the frequency of very small amplitude fusion pore feet in mouse chromaffin cells

(Wang et al., 2008). Taken together, these data suggested that Rab3A acts to prevent abnormal

fusion pore release events. However, at the NMJ and in chromaffin cells, the kinetics (rise time,

decay, half width) of the vast majority of release events were unchanged in the absence of

Rab3A, making it unlikely that the shift in the distribution of mEPSC amplitudes is mediated by a

Rab3A-dependent change in fusion pore kinetics.

Neuronal, not astrocytic, Rab3A may be required for the homeostatic release of a signaling

molecule

Our results support the surprising possibility that presynaptic Rab3A regulates

postsynaptic AMPARs, rather than occurring through a direct effect on the trafficking of

AMPARs by Rab3A in the dendrite. Astrocytic release of TNFα was shown to mediate the

increase in AMPARs in prolonged activity-blocked hippocampal cultures (Stellwagen and

Malenka, 2006), but we found that loss of Rab3A in astrocytes does not disrupt the increase in

mEPSC after activity blockade. Further, we previously showed that the homeostatic increase in

NMJ mEPC amplitude was completely normal in the absence of TNFα (Wang et al., 2011).

Another possible way Rab3A might mediate regulation of postsynaptic AMPARs is

through the presynaptic release of a signaling molecule, such as brain-derived neurotrophic

factor (BDNF), that acts anterogradely to alter postsynaptic AMPAR levels. Addition of

exogenous BDNF to neuronal cultures prevents the increase in mEPSC amplitude following

activity blockade (Rutherford et al., 1998; Benevento et al., 2016) but see (Smith-Dijak et al.,

2019), and BDNF mRNA is reduced after activity blockade in vitro (Benevento et al., 2016;

Miyasaka and Yamamoto, 2021) and in vivo (Castren et al., 1992). Also, after reduction of

secreted BDNF in culture media via the BDNF scavengers TrkB-FC or TrkB-IgG, mEPSC

amplitudes are increased (Rutherford et al., 1998; Benevento et al., 2016; Smith-Dijak et al.,

2019). Rab3B, a Rab3 family member with strong homology to Rab3A but expressed more

highly in inhibitory nerve terminals (Tsetsenis et al., 2011), is modified epigenetically following

TTX treatment by the same DNA methylation pathway as that for BDNF (Benevento et al.,

2016). In addition, reduction of Rab3A in astrocytes decreases astrocytic BDNF release (Hong et

al., 2016), and treatment of cultures with BDNF can increase Rab3A levels (Takei et al., 1997;

Thakker-Varia et al., 2001) but see (Shinoda et al., 2014). Taken together, these results suggest

an interplay between BDNF and Rab3A in the activity-dependent regulation of AMPA receptors.

Rab3A is required for divergent scaling

We have established that homeostatic synaptic plasticity in mouse cortical cultures, rat

cortical cultures, and mouse hippocampal cultures demonstrates divergent, rather than

uniform, scaling (Hanes et al., 2020; Koesters et al., 2022). The deviation from uniformity

suggests that individual synapses may be independently regulated, and do not require a global,

cell-wide signal. Our conclusion is supported by multiple studies demonstrating local activity-

dependent regulation of mEPSC amplitudes at subsets of synapses (Ju et al., 2004; Sutton et al.,

2006; Hou et al., 2008; Beique et al., 2011; Hou et al., 2011; Letellier et al., 2014). It is further

supported by the recent finding that in hippocampal cultures treated with TTX, AMPA receptors

increase selectively at synaptopodin-positive sites, which are already larger and contain higher

amounts of AMPA receptors (Dubes et al., 2022). The authors provide evidence that the

process depends on the microRNA miR-124: miR-124 is reduced after activity blockade; and

overexpression of miR-124 blocks the TTX-induced increase in AMPA receptors and mEPSC

amplitudes. Increases in miR-124 have also been linked to decreases in mEPC amplitudes at the

NMJs of the slow channel syndrome mouse (mSCS) in which prolongation of ACh currents

causes excess excitation (Zhu et al., 2013). Zhu and colleagues provide evidence for a

postsynaptic calpain-Cdk5-nNOS pathway that is activated by increased calcium levels, with the

NO feeding back retrogradely to regulate miR-124, which in turn directly interacts with Rab3A

to affect quantal content (Zhu et al., 2013), but it is also possible that Rab3A is contributing to

the mEPSC amplitude changes in mSCS. In sum, these studies suggest miR-124 could tie

together presynaptic changes (Rab3A, transmitter released from a single vesicle) and

postsynaptic changes (synaptopodin, AMPA receptors) in divergent homeostatic synaptic

plasticity.

A model for the role of Rab3A in homeostatic plasticity of mEPSC amplitude

We propose a model for Rab3A function in homeostatic synaptic plasticity of mEPSC

amplitude in which Rab3A contributes to presynaptic quantal size through the amount of

transmitter released by fusion of a single vesicle, and to postsynaptic quantal size through the

number of postsynaptic receptors. Normally, Rab3A cycles between GTP and GDP-bound forms

("State 1," indicated by red tag in Figure 10A) and promotes synaptic vesicle mobilization and

fusion, at the same time maintaining vesicle size/transmitter content within a narrow range. In

this normally cycling Rab3A state, an anterograde signal dependent on Rab3A maintains GluA2

receptors at their normal, restricted level. Activity blockade leads to a buildup of an alternate

form of Rab3A, possibly one that is unable to cycle through its GTP and GDP-bound forms

("State 2," indicated by gray tag in Figure 10B). Without the quality control mechanism

conferred by State 1 Rab3A, vesicle size/transmitter content skews to higher values, and the

anterograde signal is abnormal, leading to GluA2 receptors levels rising outside normal limits.

Not shown here, the Earlybird point mutation may permanently be in "State 2," as the mutation

is in the guanine nucleotide binding region (Kapfhamer), which would lead to increased

transmitter released from single vesicles and increased receptor levels.

We have demonstrated an essential role of the presynaptic vesicle protein Rab3A in the

homeostatic increase in mEPSC amplitude and GluA2 AMPA receptor levels at synaptic sites in

mouse dissociated cortical cultures. To our knowledge, this is the first evidence of a presynaptic

protein being implicated in the homeostatic regulation of mEPSC amplitude and AMPARs in

neuronal cultures. The current results extend our previous findings that showed the

homeostatic increase in mEPC amplitudes at the mouse NMJ in vivo depends on Rab3A. Under

conditions where postsynaptic receptors are close to being saturated by the amount of

transmitter in a single vesicle (possibly in cortical cultures but not at the NMJ), it may be

necessary to increase both the amount of transmitter released and the number or function of

postsynaptic receptors in order to observe a homeostatic increase in the physiological signal,

mEPSC amplitude. Our results suggest that in cortical cultures, neuronal Rab3A is a key player

regulating the homeostatic increase in synaptic strength on both sides of the synapse.

References

Altimimi HF, Stellwagen D (2013) Persistent synaptic scaling independent of AMPA receptor subunit
composition. J Neurosci 33:11763-11767.
Anggono V, Clem RL, Huganir RL (2011) PICK1 loss of function occludes homeostatic synaptic scaling. J
Neurosci 31:2188-2196.
Araki Y, Lin DT, Huganir RL (2010) Plasma membrane insertion of the AMPA receptor GluA2 subunit is
regulated by NSF binding and Q/R editing of the ion pore. Proc Natl Acad Sci U S A 107:11080-
11085.
Beique JC, Na Y, Kuhl D, Worley PF, Huganir RL (2011) Arc-dependent synapse-specific homeostatic
plasticity. Proc Natl Acad Sci U S A 108:816-821.
Benevento M, Iacono G, Selten M, Ba W, Oudakker A, Frega M, Keller J, Mancini R, Lewerissa E, Kleefstra
T, Stunnenberg HG, Zhou H, van Bokhoven H, Nadif Kasri N (2016) Histone Methylation by the
Kleefstra Syndrome Protein EHMT1 Mediates Homeostatic Synaptic Scaling. Neuron 91:341-355.
Benke T, Traynelis SF (2019) AMPA-Type Glutamate Receptor Conductance Changes and Plasticity: Still a
Lot of Noise. Neurochem Res 44:539-548.
Blackman MP, Djukic B, Nelson SB, Turrigiano GG (2012) A critical and cell-autonomous role for MeCP2
in synaptic scaling up. J Neurosci 32:13529-13536.
Castillo PE, Janz R, Sudhof TC, Tzounopoulos T, Malenka RC, Nicoll RA (1997) Rab3A is essential for
mossy fibre long-term potentiation in the hippocampus. Nature 388:590-593.
Castren E, Zafra F, Thoenen H, Lindholm D (1992) Light regulates expression of brain-derived
neurotrophic factor mRNA in rat visual cortex. Proc Natl Acad Sci U S A 89:9444-9448.
Cheresharov L, Ovtscharoff W, Manolov S (1978) Ultrastructural changes in the anterior horn synapses
of rat spinal cord under different locomotor conditions. J Neural Transm 42:9-21.
Coleman WL, Bykhovskaia M (2009) Rab3a-mediated vesicle recruitment regulates short-term plasticity
at the mouse diaphragm synapse. Mol Cell Neurosci 41:286-296.
Coleman WL, Bill CA, Bykhovskaia M (2007) Rab3a deletion reduces vesicle docking and transmitter
release at the mouse diaphragm synapse. Neuroscience 148:1-6.
Correa SA, Hunter CJ, Palygin O, Wauters SC, Martin KJ, McKenzie C, McKelvey K, Morris RG, Pankratov
Y, Arthur JS, Frenguelli BG (2012) MSK1 regulates homeostatic and experience-dependent
synaptic plasticity. J Neurosci 32:13039-13051.
Cuenod M, Sandri C, Akert K (1970) Enlarged synaptic vesicles as an early sign of secondary
degeneration in the optic nerve terminals of the pigeon. J Cell Sci 6:605-613.

Daniels RW, Collins CA, Gelfand MV, Dant J, Brooks ES, Krantz DE, DiAntonio A (2004) Increased
expression of the Drosophila vesicular glutamate transporter leads to excess glutamate release
and a compensatory decrease in quantal content. J Neurosci 24:10466-10474.
De Gois S, Schafer MK, Defamie N, Chen C, Ricci A, Weihe E, Varoqui H, Erickson JD (2005) Homeostatic
scaling of vesicular glutamate and GABA transporter expression in rat neocortical circuits. J
Neurosci 25:7121-7133.
Diering GH, Gustina AS, Huganir RL (2014) PKA-GluA1 coupling via AKAP5 controls AMPA receptor
phosphorylation and cell-surface targeting during bidirectional homeostatic plasticity. Neuron
84:790-805.
Diering GH, Nirujogi RS, Roth RH, Worley PF, Pandey A, Huganir RL (2017) Homer1a drives homeostatic
scaling-down of excitatory synapses during sleep. Science 355:511-515.
Dorrbaum AR, Alvarez-Castelao B, Nassim-Assir B, Langer JD, Schuman EM (2020) Proteome dynamics
during homeostatic scaling in cultured neurons. Elife 9.
Dubes S, Soula A, Benquet S, Tessier B, Poujol C, Favereaux A, Thoumine O, Letellier M (2022) miR-124-
dependent tagging of synapses by synaptopodin enables input-specific homeostatic plasticity.
EMBO J 41:e109012.
Fan D, Grooms SY, Araneda RC, Johnson AB, Dobrenis K, Kessler JA, Zukin RS (1999) AMPA receptor
protein expression and function in astrocytes cultured from hippocampus. J Neurosci Res
57:557-571.
Fernandes D, Carvalho AL (2016) Mechanisms of homeostatic plasticity in the excitatory synapse. J
Neurochem 139:973-996.
Fremeau RT, Jr., Kam K, Qureshi T, Johnson J, Copenhagen DR, Storm-Mathisen J, Chaudhry FA, Nicoll
RA, Edwards RH (2004) Vesicular glutamate transporters 1 and 2 target to functionally distinct
synaptic release sites. Science 304:1815-1819.
Gainey MA, Hurvitz-Wolff JR, Lambo ME, Turrigiano GG (2009) Synaptic scaling requires the GluR2
subunit of the AMPA receptor. J Neurosci 29:6479-6489.
Gainey MA, Tatavarty V, Nahmani M, Lin H, Turrigiano GG (2015) Activity-dependent synaptic GRIP1
accumulation drives synaptic scaling up in response to action potential blockade. Proc Natl Acad
Sci U S A 112:E3590-3599.
Gao M, Sossa K, Song L, Errington L, Cummings L, Hwang H, Kuhl D, Worley P, Lee HK (2010) A specific
requirement of Arc/Arg3.1 for visual experience-induced homeostatic synaptic plasticity in
mouse primary visual cortex. J Neurosci 30:7168-7178.
Geppert M, Bolshakov VY, Siegelbaum SA, Takei K, De Camilli P, Hammer RE, Sudhof TC (1994) The role
of Rab3A in neurotransmitter release. Nature 369:493-497.
Gu Y, Chiu SL, Liu B, Wu PH, Delannoy M, Lin DT, Wirtz D, Huganir RL (2016) Differential vesicular sorting
of AMPA and GABAA receptors. Proc Natl Acad Sci U S A 113:E922-931.
Hanes AL, Koesters AG, Fong MF, Altimimi HF, Stellwagen D, Wenner P, Engisch KL (2020) Divergent
Synaptic Scaling of Miniature EPSCs following Activity Blockade in Dissociated Neuronal Cultures.
J Neurosci 40:4090-4102.
Hong Y, Zhao T, Li XJ, Li S (2016) Mutant Huntingtin Impairs BDNF Release from Astrocytes by Disrupting
Conversion of Rab3a-GTP into Rab3a-GDP. J Neurosci 36:8790-8801.
Hou Q, Gilbert J, Man HY (2011) Homeostatic regulation of AMPA receptor trafficking and degradation
by light-controlled single-synaptic activation. Neuron 72:806-818.
Hou Q, Zhang D, Jarzylo L, Huganir RL, Man HY (2008) Homeostatic regulation of AMPA receptor
expression at single hippocampal synapses. Proc Natl Acad Sci U S A 105:775-780.
Hou Q, Ruan H, Gilbert J, Wang G, Ma Q, Yao WD, Man HY (2015) MicroRNA miR124 is required for the
expression of homeostatic synaptic plasticity. Nat Commun 6:10045.

Hu JH, Park JM, Park S, Xiao B, Dehoff MH, Kim S, Hayashi T, Schwarz MK, Huganir RL, Seeburg PH,
Linden DJ, Worley PF (2010) Homeostatic scaling requires group I mGluR activation mediated by
Homer1a. Neuron 68:1128-1142.
Hussain S, Davanger S (2015) Postsynaptic VAMP/Synaptobrevin Facilitates Differential Vesicle
Trafficking of GluA1 and GluA2 AMPA Receptor Subunits. PLoS One 10:e0140868.
Ju W, Morishita W, Tsui J, Gaietta G, Deerinck TJ, Adams SR, Garner CC, Tsien RY, Ellisman MH, Malenka
RC (2004) Activity-dependent regulation of dendritic synthesis and trafficking of AMPA
receptors. Nat Neurosci 7:244-253.
Jurado S, Goswami D, Zhang Y, Molina AJ, Sudhof TC, Malenka RC (2013) LTP requires a unique
postsynaptic SNARE fusion machinery. Neuron 77:542-558.
Kapfhamer D, Valladares O, Sun Y, Nolan PM, Rux JJ, Arnold SE, Veasey SC, Bucan M (2002) Mutations in
Rab3a alter circadian period and homeostatic response to sleep loss in the mouse. Nat Genet
32:290-295.
Karunanithi S, Marin L, Wong K, Atwood HL (2002) Quantal size and variation determined by vesicle size
in normal and mutant Drosophila glutamatergic synapses. J Neurosci 22:10267-10276.
Kavalali ET, Monteggia LM (2023) Rapid homeostatic plasticity and neuropsychiatric therapeutics.
Neuropsychopharmacology 48:54-60.
Kawana E, Akert K, Bruppacher H (1971) Enlargement of synaptic vesicles as an early sign of terminal
degneration in the rat caudate nucleus. J Comp Neurol 142:297-307.
Kennedy MJ, Davison IG, Robinson CG, Ehlers MD (2010) Syntaxin-4 defines a domain for activity-
dependent exocytosis in dendritic spines. Cell 141:524-535.
Koesters AG, Rich MM, Engisch KL (2022) Diverging from the Norm: Reevaluating What Miniature
Excitatory Postsynaptic Currents Tell Us about Homeostatic Synaptic Plasticity.
Neuroscientist:10738584221112336.
Leenders AG, Lopes da Silva FH, Ghijsen WE, Verhage M (2001) Rab3a is involved in transport of synaptic
vesicles to the active zone in mouse brain nerve terminals. Mol Biol Cell 12:3095-3102.
Letellier M, Elramah S, Mondin M, Soula A, Penn A, Choquet D, Landry M, Thoumine O, Favereaux A
(2014) miR-92a regulates expression of synaptic GluA1-containing AMPA receptors during
homeostatic scaling. Nat Neurosci 17:1040-1042.
Lignani G, Baldelli P, Marra V (2020) Homeostatic Plasticity in Epilepsy. Front Cell Neurosci 14:197.
Lledo PM, Zhang X, Sudhof TC, Malenka RC, Nicoll RA (1998) Postsynaptic membrane fusion and long-
term potentiation. Science 279:399-403.
Lloret IL, Saavedra JP (1975) Enlargement of synaptic vesicles in degenerating nerve endings: a
comparison between cat and monkey. J Neurocytol 4:1-6.
Lonart G, Janz R, Johnson KM, Sudhof TC (1998) Mechanism of action of rab3A in mossy fiber LTP.
Neuron 21:1141-1150.
Lovinger DM, Abrahao KP (2018) Synaptic plasticity mechanisms common to learning and alcohol use
disorder. Learn Mem 25:425-434.
Maienschein V, Marxen M, Volknandt W, Zimmermann H (1999) A plethora of presynaptic proteins
associated with ATP-storing organelles in cultured astrocytes. Glia 26:233-244.
Minerbi A, Kahana R, Goldfeld L, Kaufman M, Marom S, Ziv NE (2009) Long-term relationships between
synaptic tenacity, synaptic remodeling, and network activity. PLoS Biol 7:e1000136.
Miyasaka Y, Yamamoto N (2021) Neuronal Activity Patterns Regulate Brain-Derived Neurotrophic Factor
Expression in Cortical Cells via Neuronal Circuits. Front Neurosci 15:699583.
Nishimune A, Isaac JT, Molnar E, Noel J, Nash SR, Tagaya M, Collingridge GL, Nakanishi S, Henley JM
(1998) NSF binding to GluR2 regulates synaptic transmission. Neuron 21:87-97.
O'Brien RJ, Kamboj S, Ehlers MD, Rosen KR, Fischbach GD, Huganir RL (1998) Activity-dependent
modulation of synaptic AMPA receptor accumulation. Neuron 21:1067-1078.

Oh MC, Derkach VA (2005) Dominant role of the GluR2 subunit in regulation of AMPA receptors by
CaMKII. Nat Neurosci 8:853-854.
Ormel L, Stensrud MJ, Bergersen LH, Gundersen V (2012) VGLUT1 is localized in astrocytic processes in
several brain regions. Glia 60:229-238.
Osten P, Srivastava S, Inman GJ, Vilim FS, Khatri L, Lee LM, States BA, Einheber S, Milner TA, Hanson PI,
Ziff EB (1998) The AMPA receptor GluR2 C terminus can mediate a reversible, ATP-dependent
interaction with NSF and alpha- and beta-SNAPs. Neuron 21:99-110.
Pastuzyn ED, Shepherd JD (2017) Activity-Dependent Arc Expression and Homeostatic Synaptic Plasticity
Are Altered in Neurons from a Mouse Model of Angelman Syndrome. Front Mol Neurosci
10:234.
Pozo K, Goda Y (2010) Unraveling mechanisms of homeostatic synaptic plasticity. Neuron 66:337-351.
Riedel D, Antonin W, Fernandez-Chacon R, Alvarez de Toledo G, Jo T, Geppert M, Valentijn JA, Valentijn
K, Jamieson JD, Sudhof TC, Jahn R (2002) Rab3D is not required for exocrine exocytosis but for
maintenance of normally sized secretory granules. Mol Cell Biol 22:6487-6497.
Rutherford LC, Nelson SB, Turrigiano GG (1998) BDNF has opposite effects on the quantal amplitude of
pyramidal neuron and interneuron excitatory synapses. Neuron 21:521-530.
Sanderson JL, Scott JD, Dell'Acqua ML (2018) Control of Homeostatic Synaptic Plasticity by AKAP-
Anchored Kinase and Phosphatase Regulation of Ca(2+)-Permeable AMPA Receptors. J Neurosci
38:2863-2876.
Schluter OM, Basu J, Sudhof TC, Rosenmund C (2006) Rab3 superprimes synaptic vesicles for release:
implications for short-term synaptic plasticity. J Neurosci 26:1239-1246.
Schonn JS, van Weering JR, Mohrmann R, Schluter OM, Sudhof TC, de Wit H, Verhage M, Sorensen JB
(2010) Rab3 proteins involved in vesicle biogenesis and priming in embryonic mouse chromaffin
cells. Traffic 11:1415-1428.
Seeburg DP, Sheng M (2008) Activity-induced Polo-like kinase 2 is required for homeostatic plasticity of
hippocampal neurons during epileptiform activity. J Neurosci 28:6583-6591.
Shepherd JD, Rumbaugh G, Wu J, Chowdhury S, Plath N, Kuhl D, Huganir RL, Worley PF (2006)
Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. Neuron 52:475-484.
Shinoda Y, Ahmed S, Ramachandran B, Bharat V, Brockelt D, Altas B, Dean C (2014) BDNF enhances
spontaneous and activity-dependent neurotransmitter release at excitatory terminals but not at
inhibitory terminals in hippocampal neurons. Front Synaptic Neurosci 6:27.
Silva MM, Rodrigues B, Fernandes J, Santos SD, Carreto L, Santos MAS, Pinheiro P, Carvalho AL (2019)
MicroRNA-186-5p controls GluA2 surface expression and synaptic scaling in hippocampal
neurons. Proc Natl Acad Sci U S A 116:5727-5736.
Smith-Dijak AI, Nassrallah WB, Zhang LYJ, Geva M, Hayden MR, Raymond LA (2019) Impairment and
Restoration of Homeostatic Plasticity in Cultured Cortical Neurons From a Mouse Model of
Huntington Disease. Front Cell Neurosci 13:209.
Soden ME, Chen L (2010) Fragile X protein FMRP is required for homeostatic plasticity and regulation of
synaptic strength by retinoic acid. J Neurosci 30:16910-16921.
Song I, Kamboj S, Xia J, Dong H, Liao D, Huganir RL (1998) Interaction of the N-ethylmaleimide-sensitive
factor with AMPA receptors. Neuron 21:393-400.
Statman A, Kaufman M, Minerbi A, Ziv NE, Brenner N (2014) Synaptic size dynamics as an effectively
stochastic process. PLoS Comput Biol 10:e1003846.
Stellwagen D, Malenka RC (2006) Synaptic scaling mediated by glial TNF-alpha. Nature 440:1054-1059.
Stellwagen D, Beattie EC, Seo JY, Malenka RC (2005) Differential regulation of AMPA receptor and GABA
receptor trafficking by tumor necrosis factor-alpha. J Neurosci 25:3219-3228.
Suh YH, Terashima A, Petralia RS, Wenthold RJ, Isaac JT, Roche KW, Roche PA (2010) A neuronal role for
SNAP-23 in postsynaptic glutamate receptor trafficking. Nat Neurosci 13:338-343.

Sutton MA, Ito HT, Cressy P, Kempf C, Woo JC, Schuman EM (2006) Miniature neurotransmission
stabilizes synaptic function via tonic suppression of local dendritic protein synthesis. Cell
125:785-799.
Suzuki K, Kim JW, Nosyreva E, Kavalali ET, Monteggia LM (2021) Convergence of distinct signaling
pathways on synaptic scaling to trigger rapid antidepressant action. Cell Rep 37:109918.
Takei N, Sasaoka K, Inoue K, Takahashi M, Endo Y, Hatanaka H (1997) Brain-derived neurotrophic factor
increases the stimulation-evoked release of glutamate and the levels of exocytosis-associated
proteins in cultured cortical neurons from embryonic rats. J Neurochem 68:370-375.
Tan HL, Queenan BN, Huganir RL (2015) GRIP1 is required for homeostatic regulation of AMPAR
trafficking. Proc Natl Acad Sci U S A 112:10026-10031.
Thakker-Varia S, Alder J, Crozier RA, Plummer MR, Black IB (2001) Rab3A is required for brain-derived
neurotrophic factor-induced synaptic plasticity: transcriptional analysis at the population and
single-cell levels. J Neurosci 21:6782-6790.
Thiagarajan TC, Lindskog M, Tsien RW (2005) Adaptation to synaptic inactivity in hippocampal neurons.
Neuron 47:725-737.
Tian M, Xu CS, Montpetit R, Kramer RH (2012) Rab3A mediates vesicle delivery at photoreceptor ribbon
synapses. J Neurosci 32:6931-6936.
Tononi G, Cirelli C (2014) Sleep and the price of plasticity: from synaptic and cellular homeostasis to
memory consolidation and integration. Neuron 81:12-34.
Torrado Pacheco A, Bottorff J, Gao Y, Turrigiano GG (2021) Sleep Promotes Downward Firing Rate
Homeostasis. Neuron 109:530-544 e536.
Trasande CA, Ramirez JM (2007) Activity deprivation leads to seizures in hippocampal slice cultures: is
epilepsy the consequence of homeostatic plasticity? J Clin Neurophysiol 24:154-164.
Tsetsenis T, Younts TJ, Chiu CQ, Kaeser PS, Castillo PE, Sudhof TC (2011) Rab3B protein is required for
long-term depression of hippocampal inhibitory synapses and for normal reversal learning. Proc
Natl Acad Sci U S A 108:14300-14305.
Turrigiano GG (1999) Homeostatic plasticity in neuronal networks: the more things change, the more
they stay the same. Trends Neurosci 22:221-227.
Turrigiano GG, Nelson SB (2004) Homeostatic plasticity in the developing nervous system. Nat Rev
Neurosci 5:97-107.
Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB (1998) Activity-dependent scaling of
quantal amplitude in neocortical neurons. Nature 391:892-896.
Wang G, Zhong J, Guttieres D, Man HY (2019) Non-scaling regulation of AMPA receptors in homeostatic
synaptic plasticity. Neuropharmacology 158:107700.
Wang JKT, Langfelder P, Horvath S, Palazzolo MJ (2017) Exosomes and Homeostatic Synaptic Plasticity
Are Linked to Each other and to Huntington's, Parkinson's, and Other Neurodegenerative
Diseases by Database-Enabled Analyses of Comprehensively Curated Datasets. Front Neurosci
11:149.
Wang X, Thiagarajan R, Wang Q, Tewolde T, Rich MM, Engisch KL (2008) Regulation of quantal shape by
Rab3A: evidence for a fusion pore-dependent mechanism. J Physiol 586:3949-3962.
Wang X, Wang Q, Yang S, Bucan M, Rich MM, Engisch KL (2011) Impaired activity-dependent plasticity of
quantal amplitude at the neuromuscular junction of Rab3A deletion and Rab3A earlybird mutant
mice. J Neurosci 31:3580-3588.
Wang X, Li Y, Engisch KL, Nakanishi ST, Dodson SE, Miller GW, Cope TC, Pinter MJ, Rich MM (2005)
Activity-dependent presynaptic regulation of quantal size at the mammalian neuromuscular
junction in vivo. J Neurosci 25:343-351.
Weisskopf MG, Castillo PE, Zalutsky RA, Nicoll RA (1994) Mediation of hippocampal mossy fiber long-
term potentiation by cyclic AMP. Science 265:1878-1882.

Wilson NR, Kang J, Hueske EV, Leung T, Varoqui H, Murnick JG, Erickson JD, Liu G (2005) Presynaptic
regulation of quantal size by the vesicular glutamate transporter VGLUT1. J Neurosci 25:6221-
6234.
Wu D, Bacaj T, Morishita W, Goswami D, Arendt KL, Xu W, Chen L, Malenka RC, Sudhof TC (2017)
Postsynaptic synaptotagmins mediate AMPA receptor exocytosis during LTP. Nature 544:316-
321.
Xu X, Pozzo-Miller L (2017) EEA1 restores homeostatic synaptic plasticity in hippocampal neurons from
Rett syndrome mice. J Physiol 595:5699-5712.
Zhu H, Bhattacharyya B, Lin H, Gomez CM (2013) Skeletal muscle calpain acts through nitric oxide and
neural miRNAs to regulate acetylcholine release in motor nerve terminals. J Neurosci 33:7308-
7324.

986
987   FIGURE 1

Figure 1. Block of activity with 48 hours TTX treatment increased mEPSC amplitudes in cortical
cultures from wild type mice but not Rab3A$^{-/-}$ mice. A) 10 second current traces recorded at -60
mV in pyramidal cortical neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in
cultures prepared from Rab3A$^{+/+}$ mice from the Rab3A$^{+/-}$ colony. B) Average traces for the
recordings shown in (A). C) Box and whisker plots for average mEPSC amplitudes from 30 CON
and 23 TTX cells (Rab3A$^{+/+}$, Rab3A$^{+/-}$ colony; mean mEPSC amplitudes, CON, 13.9 ± 0.7 pA; TTX,
18.2 ± 0.9 pA, p = 4.58 * 10^{-4}, Kruskal-Wallis test). D) Current traces recorded in pyramidal
cortical neurons from a CON and TTX-treated neuron in cultures prepared from Rab3A$^{-/-}$ mice.
E) Average traces for the recordings shown in (D). F) Box and whisker plots for average mEPSC
amplitudes from 25 CON and 26 TTX cells, Rab3A$^{-/-}$ data (Rab3A$^{-/-}$, mean mEPSC amplitudes,
CON, 13.6 ± 0.1; TTX, 14.3 ± 0.6, p = 0.318, Kruskal-Wallis test). G) Analysis of pooled mEPSC
amplitudes for cultures from Rab3A$^{+/+}$ mice of the Rab3A$^{+/-}$ colony (i) Cumulative distribution
functions (CDFs) of quantiles of mEPSC amplitudes pooled from CON (black curve) and TTX-
treated cultures (red curve) (Kolmogorov Smirnov (KS) test, test statistic (D) = 0.149, p = 8.53 *
10^{-13}. (ii) Quantiles pooled from CON and TTX-treated cells were ranked from smallest to
largest, plotted against each other, and fit with linear regression functions. Gray line, identity;
red dashed line, linear regression fit. Inset, full range of the data. (iii) The same ranked mEPSC
quantiles were paired and the ratio of CON/TTX computed for each pair, then plotted as a
function of the CON mEPSC for the pair. Dashed line at ratio = 1.0 (no plasticity). H) Analysis of
pooled mEPSCs amplitudes for cultures from Rab3A$^{-/-}$ mice. (i) CDFs of quantiles pooled from
CON (black curve) and TTX-treated cultures (red curve) (KS test, D = 0.070, p = 0.042). (ii)
Quantiles pooled from CON and TTX-treated cells were ranked from smallest to largest, plotted
against each other, and fit with a linear regression function. Gray lines, identity; red dashed
lines, linear regression fits. Inset: full range of the data. (iii) The same ranked mEPSC quantiles
were paired and the ratio of CON/TTX computed for each pair, then plotted as a function of the
CON mEPSC for the pair. Dashed line at ratio = 1 (no plasticity).

1016

FIGURE 2

1018

Figure 2. Block of activity with 48 hours TTX treatment increased mEPSC amplitudes in cortical
cultures from wild type mice but not Rab3A$^{Ebd/Ebd}$ mice, which had an increased mEPSC
amplitude at baseline. A) 10 second current traces recorded at -60 mV in pyramidal cortical
neurons from an untreated ("CON") and TTX-treated ("TTX") neuron in cultures prepared from
Rab3A$^{+/+}$ mice from the Rab3A$^{+/Ebd}$ colony. B) Average traces for the recordings shown in (A). C)
Box and whisker plots for average mEPSC amplitudes from 20 CON and 23 TTX cells (Rab3A$^{+/+}$,
Rab3A$^{+/Ebd}$ colony, mean mEPSCs amplitudes, CON, 11.0 ± 0.6 pA; TTX, 15.0 ± 1.3 pA, p = 0.02,
Kruskal-Wallis test). D) Current traces recorded in pyramidal-shaped cortical neurons from a
CON and TTX-treated neuron in cultures prepared from Rab3A$^{Ebd/Ebd}$ mice. E) Average traces for
the recordings shown in (D). F) Box and whisker plots of average mEPSC amplitudes from 21
CON and 22 TTX cells (Rab3A$^{Ebd/Ebd}$, mean mEPSC amplitudes, CON, 15.1 ± 1.0 pA; TTX, 14.6 ±
1.1 pA, p = 0.81, Kruskal-Wallis test). G) Analysis of pooled mEPSC amplitudes for cultures from
Rab3A$^{+/+}$ mice of the Rab3A$^{+/Ebd}$ colony. (i) CDFs of quantiles pooled from CON (black curve) and
TTX-treated cultures (red curve) (KS test, D = 0.177, p value = 2.88 * 10^{-9}). (ii) Quantiles pooled
from CON and TTX-treated cells were ranked from smallest to largest, plotted against each
other, and fit with linear regression function. Gray line, identity; red dashed line, linear
regression fit. Inset: full range of the data. (iii) The same ranked mEPSC quantiles were paired
and the ratio of CON/TTX computed for each pair, then plotted as a function of the CON mEPSC
for the pair. Dashed line at ratio = 1 (no plasticity). H) Analysis of pooled mEPSC amplitudes for
cultures from Rab3A$^{Ebd/Ebd}$ mice. (i) CDFs of quantiles pooled from CON (black curve) and TTX-
treated cultures (red curve). (KS test, D = 0.067, p value = 0.097 (TTX mEPSCs were slightly
reduced)). (ii) Quantiles pooled from CON and TTX-treated cells were ranked from smallest to
largest, plotted against each other, and fit with a linear regression function. Gray line, identity;
red dashed line, linear regression fit. Inset: full range of the data. (iii) The same ranked mEPSC
quantiles were paired and the ratio of CON/TTX computed for each pair, then plotted as a
function of the CON mEPSC for the pair. Dashed line at ratio = 1 (no plasticity).

1045

1046

1047   FIGURE 3

Figure 3. The increase in mEPSC amplitudes in neurons from cultures prepared from
Rab3A$^{Ebd/Ebd}$ mice, relative to those from Rab3A$^{+/+}$ mice in the same strain, showed less
divergent scaling than that of the activity-block induced increase in mEPSC amplitudes typically
observed in cultures from Rab3A$^{+/+}$ mice. A) Cumulative distribution function of quantiles
pooled from Rab3A$^{+/+}$ (black curve) and Rab3A$^{Ebd/Ebd}$ cultures (red curve), D = 0.245; p = 1.52 *
10^{-16}, KS test. *Inset*: box and whisker plot of average mEPSC amplitudes from 20 Rab3A$^{+/+}$ and
21 Rab3A$^{Ebd/Ebd}$ cells, mean mEPSC amplitudes, Rab3A$^{+/+}$, 11.0 ± 0.7 pA; Rab3A$^{Ebd/Ebd}$, 15.1 ± 1.0
pA, p = 0.0027, Kruskal-Wallis test. B) Quantiles pooled from Rab3A$^{+/+}$ and Rab3A$^{Ebd/Ebd}$ cells
were ranked from smallest to largest, plotted against each other, and fit with linear regression
functions. Gray line, identity; red dashed line, linear regression fit. C) The same ranked mEPSC
quantiles were paired and the ratio of Rab3A$^{+/+}$/Rab3A$^{Ebd/Ebd}$ computed for each pair, then
plotted as a function of the Rab3A$^{+/+}$ mEPSC for the pair. Dashed line at ratio = 1.0 (no
difference).

1061

1062

1063

1064

1065

1066

1067   FIGURE 4

1068

Figure 4. Ca^{2+}-permeable, NASPM-sensitive glutamate receptors did not mediate the TTX-
induced increase in mEPSC amplitude in cultures of dissociated mouse cortical neurons. A)
mEPSC amplitudes for 11 mouse cortical neurons from untreated cultures ("CON") and 11
neurons from TTX treated cultures ("TTX") were recorded before and after application of 20 μM
NASPM. mEPSC amplitudes were still significantly larger in TTX-treated cultures compared to
untreated control cultures after NASPM treatment (TTX effect pre-NASPM, p = 0.009; TTX effect
post-NASPM, p = 0.006, Kruskal-Wallis non-parametric test). B) Acute application of NASPM had
a modest effect on mEPSC amplitude in cells from CON or TTX cultures that did not reach
statistical significance (CON, pre-NASPM, 12.9 ± 1.1 pA, post-NASPM, 11.9 ± 0.8 pA, p = 0.08;
TTX, pre-NASPM, 17.5 ± 0.9 pA, post-NASPM, 16.1 ± 1.0 pA, p = 0.08; student's paired t-test). C)
Acute application of NASPM consistently reduced mEPSC frequency in cells from untreated and
cells from TTX-treated cultures (CON, pre-NASPM, 1.84 ± 1.82 mEPSCs sec^{-1}, post-NASPM, 1.56
± 1.74 mEPSCs sec^{-1}; p = 0.003; TTX, pre-NASPM, 4.40 ± 3.51 mEPSCs sec^{-1}, post-NASPM, 2.68 ±
2.25 mEPSCs sec^{-1}, p = 0.02; student's paired t-test).

1083

1084

1085

1086

1089

1090   FIGURE 5

1091

Figure 5. Confocal images of GluA2 and VGLUT1 immunofluorescence on MAP2-positive
primary dendrites of pyramidal-shaped neurons in high density dissociated cortical cultures
prepared from Rab3A[+/+] mice. Shown are 5X confocal sections of two primary dendrites, one
from an untreated culture (CON) and the other from a culture treated with TTX (TTX).
Postsynaptic GluA2-positive receptor clusters were identified by their proximity to VGLUT1
immunoreactive presynaptic terminals within the same confocal section. GluA2-VGLUT1
synaptic pairs, which were of maximal brightness in these confocal slices, are indicated by white
trapezoids. It can be seen that substantial numbers of GluA2-positive clusters located on the
dendrites are unopposed by VGLUT1-positive terminals, and that some GluA2-positive and
VGLUT1-positive clusters are found away from any MAP2 positivity, suggesting localization to
axons or astrocytes. Scale bar = 10 μm.

1103

1104

1105   FIGURE 6

Figure 6. GluA2 receptor cluster size and average intensity increased following activity blockade
in parallel with mEPSC amplitudes recorded in the same cortical cultures prepared from
Rab3A$^{+/+}$ mice, but the effects on receptor levels were less robust and differed from mEPSC
amplitudes in characteristics of scaling. A) Cumulative distribution functions (CDFs) of mEPSC
amplitudes recorded from untreated ("CON") and TTX-treated cells ("TTX") pooled from 3
cultures were significantly different (KS test, D = 0.162, p = 1.42 * 10^{-8}, CON, n = 690, TTX, n =
720). *Inset*, mean mEPSC amplitudes for 23 CON cells and 24 TTX cells were significantly
different (CON, 13.7 ± 4.5 pA; TTX, 16.4 ± 4.3 pA, p = 0.016, Kruskal-Wallis test). B) 24 mEPSC
amplitude quantiles from each of 23 untreated cells were sorted from smallest to largest and
plotted vs. 23 mEPSC amplitude quantiles from each of 24 TTX-treated cells, and the resulting
relationship fit with a linear regression equation, Y = 1.106x + 1.22 (R^2 = 0.984). *Inset*, full range
of data used for fit. C) Sorted data from (B) were used to calculate the ratio of TTX mEPSC
amplitude/CON mEPSC amplitude and the ratios plotted vs. CON mEPSC amplitude. D) CDFs of
size of GluA2 receptor clusters imaged from primary dendrites in the same untreated cultures
and TTX-treated cultures as in (A) – (C) were significantly different (KS test, D = 0.089, p = 0.002,
CON, n = 870, TTX, n = 870). *Inset*, mean GluA2 cluster size for 29 CON dendrites and 29 TTX
dendrites were not significantly different (CON, 0.97 ± 0.38 μm^2; TTX, 1.15 ± 0.59 μm^2, p = 0.44,
Kruskal-Wallis test). E) 29 GluA2 cluster size quantiles from each of 29 untreated dendrites
were sorted from smallest to largest and plotted vs. 29 cluster size quantiles from each of 29
TTX-treated dendrites, and the resulting relationship fit with a linear regression equation, Y =
1.328x -0.136 (R^2 = 0.98). *Inset*, full range of data used for fit. F) Sorted data from (E) were used
to calculate the ratio of TTX GluA2 cluster size/CON GluA2 cluster size and the ratios plotted vs.
CON GluA2 cluster size. G) CDFs of intensity of GluA2 receptor clusters imaged from primary
dendrites in untreated cultures and TTX-treated cultures were significantly different (KS test, D
= 0.120, p = 6.73 * 10^{-6}, CON, n = 870, TTX, n = 870). *Inset*, mean GluA2 cluster intensity for 29
CON dendrites and 29 TTX dendrites were not significantly different (CON, 673 ± 90 a.u.; TTX,
687 ± 72 a.u., p = 0.25, Kruskal-Wallis test). H) 29 GluA2 cluster intensity quantiles from 29
untreated dendrites were sorted from smallest to largest and plotted vs. 29 cluster intensity
quantiles from 29 TTX-treated dendrites, and the resulting relationship fit with a linear
regression equation, Y = 0.899x + 82 (R^2 = 0.987). *Inset*, full range of data used for fit. I) Sorted
data from (H) were used to calculate the ratio of TTX cluster intensity/CON cluster intensity and
plotted vs. CON cluster intensity.

1138

FIGURE 7

Figure 7. GluA2 receptor cluster size, average intensity, and mEPSC amplitudes recorded in the
same cortical cultures prepared from Rab3A$^{-/-}$ mice were not increased following activity
blockade. A) Cumulative distribution functions (CDFs) of mEPSC amplitudes recorded from
untreated ("CON") and TTX-treated cells ("TTX") pooled from 3 Rab3A$^{-/-}$ cultures were not
significantly different (KS test, D = 0.072, p = 0.085, CON, n = 630, TTX, n = 570). *Inset*, mean
mEPSC amplitudes for 21 CON cells and 19 TTX cells were not significantly different (CON, 14.9
± 3.8 pA; TTX, 14.0 ± 4.0 pA, p = 0.34, Kruskal-Wallis test). B) 19 mEPSC quantiles from each of
21 untreated cells were sorted from smallest to largest and plotted vs. 21 mEPSC quantiles
from each of 19 TTX-treated cells, and the resulting relationship fit with a linear regression
equation, Y = 0.93x + 0.13 (R^2 = 0.997). *Inset*, full range of data used for fit. C) Sorted data from
(B) were used to calculate the ratio of TTX mEPSC amplitude/CON mEPSC amplitude and
plotted vs. CON mEPSC amplitude. D) CDFs of GluA2 receptor cluster size imaged from primary
dendrites in the same untreated cultures and TTX-treated cultures as in (A) – (C) were not
significantly different (KS test, D = 0.045, p = 0.31, CON, n = 900, TTX, n = 870). *Inset*, mean
GluA2 cluster size for 30 CON dendrites and 29 TTX dendrites were not significantly different
(CON, 0.93 ± 0.27 μm^2; TTX, 0.91 ± 0.28 μm^2, p = 0.74, Kruskal-Wallis test). E) 29 GluA2 cluster
size quantiles from each of 30 untreated dendrites were sorted from smallest to largest and
plotted vs. 30 cluster size quantiles from each of 29 TTX-treated dendrites, and the resulting
relationship fit with a linear regression equation, Y = 1.01x -0.03 (R^2 = 0.993). *Inset*, full range of
data used for fit. F) Sorted data from (E) were used to calculate the ratio of TTX GluA2 cluster
size/CON GluA2 cluster size and the ratios plotted vs. CON GluA2 cluster size. G) CDFs of GluA2
receptor cluster average intensity imaged from primary dendrites in untreated cultures and
TTX-treated cultures were not significantly different (KS test, D = 0.0.060, p = 0.08, CON, n =
900, TTX, n = 870). *Inset*, mean GluA2 cluster intensity for 30 CON dendrites and 29 TTX
dendrites were not significantly different (CON, 766 ± 68 a.u.; TTX, 776 ± 79 a.u., p = 0.47,
Kruskal-Wallis test). H) 29 GluA2 cluster intensity quantiles from 30 untreated dendrites were
sorted from smallest to largest and plotted vs. 30 cluster intensity quantiles from 29 TTX-
treated dendrites, and the resulting relationship fit with a linear regression equation, Y = 1.09x -
60 (R^2 = 0.993). *Inset*, full range of data used for fit. I) Sorted data from (H) were used to
calculate the ratio of TTX cluster intensity/CON cluster intensity and the ratios plotted vs. CON
cluster intensity.

1172

1173

FIGURE 8

Figure 8. mEPSC amplitudes and GluA2 receptor cluster sizes did not always change in parallel

after treatment with TTX in cultures from Rab3A$^{+/+}$ and Rab3A$^{-/-}$ mice. A) **Rab3A$^{+/+}$ Culture #1**:

mEPSC amplitudes and GluA2 receptor cluster sizes increased in parallel after treatment in TTX.

i) CDFs of mEPSC amplitudes of cells from untreated and TTX treated dishes were significantly

different (KS test, D = 0.16, p = 0.010; CON, n = 180, TTX, n = 210). ii) CDFs of GluA2 receptor

cluster sizes of dendrites from untreated and TTX treated dishes were significantly different (KS

test, D = 0.17, p = 3.79 * 10^{-4}; CON, n = 300, TTX, n = 300). iii) 35 mEPSC amplitude quantiles

from each of 6 untreated cells and 30 quantiles from each of 7 TTX-treated cells were sorted

from smallest to largest and used to calculate the ratio of TTX mEPSC amplitude/CON mEPSC

amplitude, which was plotted vs. CON mEPSC amplitude. iv) 30 GluA2 receptor cluster size

quantiles from each of 10 untreated dendrites and 30 GluA2 receptor cluster size quantiles

from each of 10 TTX-treated dendrites were used to calculate the ratio of TTX GluA2 cluster

size/CON GluA2 cluster size, and plotted vs. CON GluA2 cluster size. *Inset*, full range of data. B)

Rab3A$^{+/+}$ Culture #2: mEPSC amplitudes and GluA2 receptor cluster sizes increased in parallel

after treatment in TTX. i) CDFs of mEPSC amplitudes of cells from untreated and TTX treated

dishes were significantly different (KS test, D = 0.21, p = 1.06 * 10^{-4}; CON, n = 210, TTX, n = 240).

ii) CDFs of GluA2 receptor cluster sizes of dendrites from untreated and TTX treated dishes

were significantly different (KS test, D = 0.21, p = 5.39 * 10^{-6}; CON, n = 270, TTX, n = 300). iii) 32

mEPSC amplitude quantiles from each of 7 untreated cells, and 28 mEPSC amplitude quantiles

from each of 8 TTX treated cells were sorted from smallest to largest and were used to

calculate the ratio of TTX mEPSC amplitude/CON mEPSC amplitude, and plotted vs. CON mEPSC

amplitude. iv) 30 GluA2 receptor cluster size quantiles from each of 9 untreated dendrites and

27 GluA2 receptor cluster size quantiles from each of 10 TTX-treated dendrites were sorted

from smallest to largest and used to calculate the ratio of TTX GluA2 cluster/CON GluA2 cluster

size, and plotted vs. CON GluA2 cluster size. *Inset*, full range of data. C) **Rab3A$^{+/+}$ Culture #3**:

mEPSC amplitudes increased, but GluA2 receptor cluster sizes decreased, after treatment in

TTX. i) CDFs of mEPSC amplitudes of cells from untreated and TTX treated dishes were

significantly different (KS test, D = 0.15, p = 0.002; CON, n = 300, TTX, n = 270). ii) CDFs of GluA2

receptor cluster sizes of dendrites from untreated and TTX treated dishes were significantly

different, but the TTX CDF is shifted to the *left* (KS test, D = 0.16, p = 0.001; CON, n = 300, TTX, n

= 270) iii) 27 mEPSC amplitude quantiles from each of 10 untreated cells, and 30 mEPSC

amplitude quantiles from each of 9 TTX-treated cells, were sorted from smallest to largest, used

to calculate the ratio of TTX mEPSC amplitude/ CON mEPSC amplitude, and plotted vs. CON

mEPSC amplitude. **iv)** 27 GluA2 receptor cluster size quantiles from each of 10 untreated

dendrites and 30 GluA2 receptor cluster sizes from each of 9 TTX-treated dendrites were sorted

from smallest to largest, used to calculate the ratio of TTX GluA2 receptor cluster size/CON

GluA2 receptor cluster size, and plotted vs. CON GluA2 receptor cluster size. D) **Rab3A$^{-/-}$ Culture**

#1 (note, this culture is distinct from that of Rab3A$^{+/+}$ Culture #1). Neither mEPSC amplitudes,

nor GluA2 receptor cluster size, increased following TTX treatment. i) CDFs of mEPSC

amplitudes of cells from untreated and TTX-treated cultures were not significantly different (KS

test, D = 0.05, p = 0.90; CON, n = 210, TTX, n = 180). ii) CDFs of GluA2 receptor cluster sizes of

dendrites from untreated and TTX (Figure 8 continued) treated dishes were not significantly

different, although the TTX CDF is shifted slightly to the *left* (KS test, D = 0.10, p = 0.08; CON, n =

300, TTX, n = 300). iii) 30 mEPSC amplitude quantiles from each of 7 untreated cells and 35

mEPSC quantiles from each of 6 TTX-treated cells were sorted from smallest to largest, used to
calculate the ratio TTX mEPSC amplitude/CON mEPSC amplitude, and plotted vs. CON mEPSC
amplitude. iv) 30 GluA2 receptor cluster size quantiles from each of 10 untreated dendrites and
10 TTX-treated dendrites were sorted from smallest to largest, used to calculate the ratio TTX
cluster size/CON cluster size, and plotted vs. CON cluster size. E) **Rab3A$^{-/-}$ Culture #2** (distinct
from Rab3A$^{+/+}$ Culture #2) mEPSC amplitudes and GluA2 receptor cluster sizes decreased after
TTX treatment. i) TTX mEPSC amplitudes CDF is shifted slightly to the *left* compared to the CON
mEPSC amplitudes CDF but the difference is not significant (KS test, D = 0.08, p = 0.28, CON, n =
240, TTX, n = 240). ii) TTX GluA2 receptor cluster sizes CDF is significantly shifted to the *left*
compared to CON GluA2 receptor cluster sizes CDF (KS test, D = 0.13, p = 0.01; CON, n = 300,
TTX, n = 300). iii) 24 mEPSC amplitude quantiles from each of 8 untreated cells and 8 TTX-
treated cells were sorted from smallest to largest, used to calculate the ratio TTX mEPSC
amplitude/CON mEPSC amplitude, and plotted vs. CON mEPSC amplitude. iv) 27 GluA2 receptor
cluster size quantiles from each of 10 untreated dendrites, and 30 GluA2 receptor cluster size
quantiles from each of 9 TTX treated dendrites, were sorted from smallest to largest, used to
calculate the ratio TTX GluA2 cluster size/CON cluster size, and plotted vs. CON cluster size.
Both mEPSC amplitude and GluA2 receptor cluster size ratios were below 1 across the majority
of data. F) **Rab3A$^{-/-}$ Culture #3** (distinct from Rab3A$^{+/+}$ Culture #3). mEPSC amplitudes were
decreased following TTX treatment, but GluA2 receptor cluster sizes were increased. i) TTX CDF
of mEPSC amplitudes was significantly shifted to the left of CON CDF (KS test, D = 0.21, p = 4.85
* 10^{-4}, CON, n = 180, TTX, n = 180). ii) TTX CDF of GluA2 receptor cluster sizes was significantly
shifted to the right of CON CDF (KS test, D = 0.15, p = 0.003, CON, n = 300, TTX, n = 300). iii) 30
mEPSC amplitude quantiles from each of 6 untreated and 6 TTX-treated cells were sorted from
smallest to largest, used to calculate the ratio TTX mEPSC amplitude/CON mEPSC amplitude,
and plotted vs. CON mEPSC amplitude. iv) 30 GluA2 receptor cluster size quantiles from each of
10 untreated dendrites and 10 TTX-treated dendrites were sorted from smallest to largest, used
to calculate the ratio TTX GluA2 cluster size/CON GluA2 cluster size, and plotted vs. CON CluA2
cluster size.

1247
1248

                                    FIGURE 9

Figure 9. Rab3A in neurons, not astrocytes, was required for full homeostatic plasticity after
activity blockade. A-C, Data from dissociated cortical neurons plated on an astrocyte feeder
layer, each prepared separately from the type of mice depicted in the schema at left: A)
Neurons from Rab3A[+/+] mice plated on astrocytes from Rab3A[+/+] mice; i) CDFs for mEPSC
amplitudes from untreated (CON, black curve) and TTX-treated (TTX, red curve) cultures (KS
test, test statistic, D = 0.141, p = 2.74 * 10^{-5}, n = 510 CON, 600 TTX samples; *inset*, means, CON,
13.3 ± 0.5 pA, n = 17 cells, TTX, 16.7 ± 1.2 pA, n = 20 cells, p = 0.03, Kruskal Wallis test; ii) rank
order plot and linear regression fit, slope = 1.89, intercept = -8.51, R^2 = 0.97; iii) ratio plot of TTX
mEPSC amplitude/CON amplitude vs. CON amplitude; B) Neurons from Rab3A[+/+] mice plated on
astrocytes from Rab3A[-/-] mice; i) CDFs of mEPSC amplitudes, CON, black curve; TTX, red curve
(KS test, test statistic 0.273, p = 2.98 * 10^{-11}), n = 330 CON, 330 TTX samples; *inset*, means, CON,
13.3 ± 1.0 pA, n = 11 cells, TTX, 18.8 ± 1.4 pA, n = 11 cells, p = 0.005, Kruskal-Wallis test; ii) rank
order plot and linear regression fit, slope = 1.76, intercept = -4.64, R^2 = 0.98; iii) ratio plot of TTX
mEPSC amplitude/TTX mEPSC amplitude vs. CON mEPSC amplitude; C) Neurons from Rab3A[-/-]
neurons plated on astrocytes from Rab3A[+/+] mice; i) CDFs of mEPSC amplitudes, CON, black
curve, TTX, red curve (KS test, test statistic 0.126, p = 0.005), n = 420 CON, 330 TTX samples;
inset, means, CON, 15.2 ± 1.1 pA, n = 14 cells, TTX, 16.9 ± 0.7 pA, n = 11 cells, p = 0.125, Kruskal-

Wallis test;; ii) rank order plot and linear regression fit; slope = 1.01, intercept = 1.53, R^2 = 0.97;
iii) ratio plot, TTX mEPSC amplitude/CON mEPSC amplitude vs. CON mEPSC amplitude. Gray
line, identity; red dashed line, linear regression fit. Insets in rank order plots depict full range of
data.

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285 FIGURE 10

1286

Figure 10. Rab3A is required for increased mEPSC amplitude after activity blockade via 2
mechanisms: 1. increased amount of transmitter released from a single vesicle and 2. increased
postsynaptic receptors. *Left*: in the presence of normal activity, Rab3A is in its control state
("State 1," red), maintaining mEPSC amplitude within a limited range by regulating either
vesicle size or transmitter content, or both, and by regulating the number of postsynaptic
receptors, likely through release of an anterograde signal. *Right*: after activity is blocked, Rab3A
accumulates in an alternate functional state ("State 2," gray) resulting in quantal size expanding
to larger ranges because of increased vesicle size or transmitter content and because of
increased numbers of receptors no longer being under the normal control of the anterograde
signal.

1297

1298

	Rab3A[+/+]		Rab3A[-/-]	
mEPSC amplitude (pA)	**CON**	**TTX**	**CON**	**TTX**
mean	13.7 ± 4.5 (n=23)	16.4 ± 4.3 (n=24)	14.9 ± 3.8 (n=21)	14.0 ± 4.0 (n=19)
KW p value	p = 0.016		p = 0.34	
KS comparison of CDF	N CON (n=690), N TTX (n=720) D = 0.162 p = 1.42 * 10^{-8}		N CON (n=630), N TTX (n=570) D = 0.072 p= 0.085	
Rank order linear regression (slope, intercept, R^2)	Slope, 1.11 Intercept, 1.22 R^2, 0.984		Slope, 0.93 Intercept, 0.13 R^2, 0.997	
Mean ratio TTX/CON 50-75[th] quantile	1.24		0.957	
GluA2 receptor cluster size (μm^2)	**CON**	**TTX**	**CON**	**TTX**
mean	0.97 ± 0.38 (n=29)	1.15 ± 0.59 (n=29)	0.93 ± 0.27 (n=30)	0.91 ± 0.28 (n=29)
KW p value	p = 0.44		p = 0.74	
KS comparison of CDF	N CON (n=870), N TTX (n=870) D = 0.089 p = 0.002		N CON (n=900), N TTX (n=870) D = 0.045 p = 0.31	
Rank order linear regression (slope, intercept, R^2)	Slope, 1.33 Intercept, -0.13 R^2, 0.98		Slope, 1.01 Intercept, -0.03 R^2, 0.993	
Mean ratio TTX/CON 50-75[th] quantile	1.11		0.976	
GluA2 receptor cluster intensity (arb. units)	**CON**	**TTX**	**CON**	**TTX**
mean	673 ± 90 (n=29)	687 ± 72 (n=29)	766 ± 68 (n=30)	776 ± 79 (n=29)
KW p value	p = 0.25		p = 0.47	
KS comparison of CDF	N CON (n=870), N TTX (n=870) D = 0.120 p = 6.73 * 10^{-6}		N CON (n=900), N TTX (n=870) D = 0.060 p = 0.08	
Rank order linear regression (slope, intercept, R^2)	Slope, 0.90 Intercept, 82 R^2, 0.987		Slope, 1.09 Intercept, -60 R^2, 0.993	
Mean ratio TTX/CON 50-75[th] quantile	1.02		1.02	

1299

1300   TABLE 1

1301   Table 1. Comparison of effect of activity blockade with TTX on mEPSC amplitude and GluA2 receptor

1302   cluster characteristics. Voltage clamp recordings of mEPSCs and confocal imaging of GluA2 receptor

1303   immunofluorescence at VGLU1-positive synapses were performed on coverslips from the same 3

1304   Rab3A$^{+/+}$ cultures (left) and 3 Rab3A$^{-/-}$ cultures (right). p values are not corrected for multiple

1305   comparisons.