Title: **Evolutionary engineering a larger porin using a loop-to-hairpin mechanism**

Authors: Rik Dhar[1], Alexander M. Bowman[1], Brunojoel Hatungimana[1], and Joanna SG Slusky[1,2*]

Affiliations: [1]Department of Molecular Biosciences, The University of Kansas, Lawrence KS 66045

[2] Computational Biology Program, The University of Kansas, Lawrence KS 66047

Corresponding author information: slusky@ku.edu

Abstract

In protein evolution, diversification is generally driven by genetic duplication. The hallmarks of this mechanism are visible in the repeating topology of various proteins. In outer membrane β-barrels, duplication is visible with β-hairpins as the repeating unit of the barrel. In contrast to the overall use of duplication in diversification, a computational study hypothesized evolutionary mechanisms other than hairpin duplications leading to increases in the number of strands in outer membrane β-barrels. Specifically, the topology of some 16- and 18-stranded β-barrels appear to have evolved through a loop to β-hairpin transition. Here we test this novel evolutionary mechanism by creating a chimeric protein from an 18-stranded β-barrel and an evolutionarily related 16-stranded β-barrel. The chimeric combination of the two was created by replacing loop L3 of the 16-stranded barrel with the sequentially matched transmembrane β-hairpin region of the 18-stranded barrel. We find the resulting chimeric protein is stable and has characteristics of increased strand number. This study provides the first experimental evidence supporting the evolution through a loop to β-hairpin transition.

Highlights

- We find evidence supporting a novel diversification mechanism in membrane β-barrels

- The mechanism is the conversion of an extracellular loop to transmembrane β-hairpin

- A chimeric protein modeling this mechanism folds stably in the membrane

- The chimera has more β-structure and a larger pore, consistent with a loop-to-hairpin transition

Keywords

Outer membrane proteins, β-barrel, loop-to-hairpin, molecular evolution, Neisserial porin, barrel enlargement, PorB, *Neisseria gonorrhoeae*, MOMP, *Campylobacter jejuni*

Abbreviations

OMP – Outer membrane proteins

MOMP – Major outer membrane protein

SDS-PAGE – Sodium dodecyl sulfate-polyacrylamide gel electrophoresis

CD – circular dichroism

SEC – size exclusion chromatography

MALDI-TOF – matrix-assisted laser desorption/ionization-time of flight

pLDDT – predicted local distance difference test (pLDDT)

PMSF – phenylmethylsulfonyl fluoride

Introduction

Outer membrane proteins (OMPs) are found in the outer membrane of Gram-negative bacteria, mitochondria, and the chloroplast [1, 2]. Unlike inner membrane proteins which consist of membrane-spanning alpha-helices, OMPs generally consist of membrane-spanning anti-parallel β-barrels, connected sequentially with shorter turns in the inter-membrane side and longer loops on the outward side [3-5]. The loops of OMPs can have various functions including structural interaction with the lipopolysaccharides [6], channel gating [7, 8], protease activity [9], interaction with target protein [10], and inter-barrel interactions [11].

OMPs differ from each other by the number of strands forming their transmembrane β-barrel domain, which ranges from 8 to 36 strands [12, 13]. Similar to divergent protein evolution more generally [14-22], OMPs primarily diversified through genetic duplication of a primordial peptide specifically an ancestral β-hairpin peptide [23]. As a result, the β-hairpin motif forms the repeating unit in OMPs, which is evident from internal sequence similarities [25-27]. Analysis of these internal sequence repeats in structurally resolved OMPs revealed an overview of the OMP evolutionary pathway of OMPs tracing the relationship between β-barrels classes of different strand numbers [21].

In addition to duplication events, further diversification occurred by mutation, deletion, and insertion followed by further gene duplication [20, 24] creating a relationship web among most outer membrane proteins. Among the most thought-provoking of these relationships was that transmembrane strands are not only related to each other, but some are also related to extracellular loops. The presence of sequence alignment between β-hairpins of 18-stranded β-barrels and the loops of 16-stranded β-barrels led to the hypothesis that extracellular loops are sometimes evolutionarily converted into transmembrane β-hairpins. This mechanism might be particularly surprising because it is a non-duplication event causing a repeat protein topology.

Moreover, soluble and transmembrane regions are not generally thought to be interchangeable. However, since outer membrane proteins and soluble proteins have similar hydrophobicities and are just "inside out" [28], this hypothesized mechanism may in fact occur in nature.

Due to the importance of the duplication evolutionary mechanism in OMP diversification, many have worked to engineer OMPs using this principle [19, 29-32]. However, incorporating additional strands into β-barrels is challenging, as evidenced by the unsuccessful integration of additional strands in OmpG [33] and the subsequent loss of channel conductance in LamB after the insertion of additional β-hairpin [32]. Another laboratory approach for strand accretion involves creating chimeras from two different β-barrel [19, 32, 34, 35]. Again this is also challenging as not all combinations result in a stable protein [35, 36] nor do they result in oligomeric proteins [19, 34]. The interaction of neighboring residues [37], the sensitivity of residues to mutations [38] and the role of hydrogen bonding in the backbone [3], can frustrate design attempts involving strand insertion.

OMP engineering is most likely to be successful when natively adjacent strands are inserted, such as the first and the last strand, in 8 to 16-stranded OmpX engineering [19] or when strands are replaced with corresponding ones in the OmpF and OmpG chimera [34]. These approaches preserve the interaction between conserved residues, ensuring the stability of the resulting protein. Because of this, usage of evolutionary relationships is known to be more likely to yield success in the difficult endeavor of enlarging OMP β-barrels.

Here we use the hypothesized loop-to-hairpin mode of evolution to engineer an 18-stranded β-barrel from a 16-stranded β-barrel. We selected a protein pair from our previous study that shows evidence for strand accretion through loop-to-hairpin transition [21]. This protein pair consists of the 16-stranded β-barrel PorB from *Neisseria Gonorrhoeae* (PDBID 4AUI) [39]. and 18-stranded β-barrel MOMP from *Campylobacter jejuni* (PDBID 5LDV) [40]. Loop L3 of PorB is its longest loop which lines the lumen of the barrel domain [39]. This loop was found to align with a β-hairpin of

MOMP. We engineer a chimeric protein by replacing loop L3 of PorB with the aligned region of MOMP. Despite being created from proteins from different species with only 21.7% sequence identity, the chimeric protein stably expresses, folds, and inserts into the outer membrane. Using secondary structure estimation from circular dichroism and pore size estimates from single channel conductance, the chimeric protein most likely forms an 18-stranded barrel.

Results

Chimeric protein folds in the outer membrane

To engineer an 18-stranded barrel from a 16-stranded barrel based on loop-to-hairpin evolution we created a chimeric protein from PorB and MOMP. This chimeric protein, which we call $PorB_{18}$, was created by replacing the loop L3 of wild-type PorB ($PorB_{WT}$) with the sequentially aligned region of MOMP (Fig. 1A and B). To evaluate the stability of the chimeric protein, we tested its production in the outer membrane. For this, the N-terminal signal sequence from the *Escherichia coli* OMP OmpA was added to the N-terminal of $PorB_{WT}$ and $PorB_{18}$ to target the proteins to the outer membrane [41, 42]. For expression, we used an *E. coli* strain that lacks the periplasmic protease DegP, as neisserial porins are difficult to express recombinantly in *E. coli* outer membrane [43, 44]. We lysogenized this ΔDegP strain with λDE3 which allows for the use of the pET expression system for the 'leaky' expression of $PorB_{WT}$ and $PorB_{18}$ [45].

A

```
Int Num 2780
E-value 2.8E-5
4aui_A 000  AMADVTLYGTIKAGVETSRSVAHHGAQADRVKTATEIADLGSKIGFKGQEDLGNGLKAIWQLEQKAYVSG--------TDTGWGNRQSFIGLKG--GFGK  090
5ldv_A 000  AIKDVDVSGVLRYRYDTGNFDKNFL-NNSNLNNSK--QDHKYRAQVNFSAAIADNFKAFVQFDYNAVDGGTGVDNATNAEKGLFVRQLYLTYTNEDVATS  097

4aui_A 091  VRVGRLNN--ILKDT----GNFNP-------WEGKSYY-SGLSNIAQ--PEE--------------------R-HVS----VRYDSP-EFAGFSGSVQ  146
5ldv_A 098  VIAGKQQLNIIWTDNGVDGLVGTGVKVVNNSIDGLTLAAFAVDSFMAAEQGSDLLGQSTYVGNGKNNNDSFKLDSIGNLYGAAAVGSYDLAGGQFNPQLW  197

4aui_A 147  YVPNDNSGKNRSESYHAGFNYKND---GFFVQYAGSYKRHNYT--------TEKHQVHRLVGGYDHDALYASVAVQQQDAKL--------T-WRD----D  222
5ldv_A 198  LAYWD----QVAFFYAVDAAYSTTIFDGINWTLEGAYLGNSLDSELDDKKTYANGNLFALKGSIEVNGWDASLGGLYYGDKEKASTVVIEDQGNLGSLLA  293

4aui_A 223  N-------------SHNSQTEVATTVAYRFG-NVTPRVSYAHGFKGSVYD-ADHDNTYDQVVVGAEYDFSKRTSALVSAGWLQKGKGAE---KFVATVGG  304
5ldv_A 294  GEEIFYTTGSRLNGDTGRNIFGYVTGGYTFNETVRVGADFVYGGTKTEATTHLGGGKKLEAVARVDYKYSPKLNFSAFYSYVNLDQGVNTNESADHSTVR  393

4aui_A 305  VGLRHKF  311
5ldv_A 394  LQALYKF  400
```

B

PorB (4aui)
16 stranded

MOMP (5ldv)
18 stranded

PorB₁₈

Figure 1: PorB$_{18}$, a chimera of PorB$_{WT}$ and MOMP: (A) Sequence alignment between 16-stranded PorB (PDB ID 4aui [39], green text) and 18-stranded MOMP (PDB ID 5ldv [46], brown text). (Adapted from Franklin [47]) The highlighted colors represent the β-strands as identified from the crystal structures: the N-terminal strand is in magenta, the C-terminal strand is in red, and the intermediate strands are highlighted in a rainbow gradient. Red/blue text indicates the region of the loop/hairpin alignment **(B)** Construction of the chimeric protein PorB$_{18}$. The majority

of loop L3 (red) from PorB$_{WT}$ (green) has been removed, and the aligned peptide sequence (blue) of MOMP has been inserted (brown) in L3's position.

Western blot of different subcellular fractions of *E. coli* showed that PorB$_{WT}$ and PorB$_{18}$ are expressed and are trafficked to the outer membrane (Fig. 2A). To assess the proper folding and insertion of the recombinant proteins in the outer membrane, we carried out an *in vivo* Proteinase K digestion. Proteinase K is a serine protease that predominantly cleaves the peptide bond adjacent to the carboxylic group of an aliphatic or aromatic residue [48]. Proteinase K assay has been used in the study of membrane proteins to discover exposed or disordered regions [49] and to evaluate OMP insertion in the outer membrane [50]. *E. coli* expressing the recombinant proteins were permeabilized and treated with Proteinase K for 30 minutes on ice. Western blot of these Proteinase K treated cells indicated both PorB$_{WT}$ and PorB$_{18}$ were resistant to Proteinase K digestion (Fig. 2B). We used TolC as a control for this assay as it has a 5kDa disordered region in the periplasm that is susceptible to Proteinase K digestion [51]. PorB$_{WT}$, PorB$_{18}$, and TolC were found to have a low level of degradation products, which can be attributed to the endogenously expressed OmpT protease in the *E. coli* strain used for expression. Resistance to Proteinase K digestion indicates that both PorB$_{WT}$ and chimeric PorB$_{18}$ insert and fold into the outer membrane.

Figure 2: Expression of PorB$_{18}$ in E. coli outer membrane: (A, B) Western blots with the nitrocellulose membrane stained with Ponceau S and subsequently immunoblotted against various antibodies. **(A)** Western blot of different fractions of *E. coli* displaying the localization of the his-tagged recombinant proteins PorB$_{WT}$ and PorB$_{18}$ in the outer membrane. SurA, a periplasmic protein, indicates the localization of periplasmic proteins. Whole-cell fractions (WC), soluble (S), inner membrane and trapped periplasmic protein (IM/P), and outer membrane (OM) **(B)** Western blot displaying the susceptibility of proteins to Proteinase K under different conditions. His-tagged PorB$_{WT}$ (blue asterisks) and PorB$_{18}$ (green asterisks) show resistance to Proteinase K when not boiled but are partially susceptible when heat denatured. TolC (red asterisks) is used as a positive control as it shows a 5 kDa cleavage in the presence of proteinase K resulting in a 45 kDa cleavage product (red arrows) [51].

Chimeric protein has increased β-sheet content

To estimate the secondary structure of the chimeric protein PorB$_{18}$ we measured and analyzed the circular dichroism (CD) spectra of these proteins. Because a higher concentration of protein was required for CD analysis, the recombinant proteins were expressed as inclusion bodies, purified, and refolded into LDAO detergent micelles [52]. Expression as inclusion bodies provides a much higher yield compared to expression in the outer membrane. The refolded proteins were separated by size exclusion chromatography (SEC) to isolate the folded trimeric fractions. The eluents were treated with a reversible crosslinker DSP (Fig. 3A) to verify the quaternary structure. Crosslinking and subsequent visualization by SDS-PAGE indicated that both the recombinant proteins PorB$_{WT}$ and PorB$_{18}$ trimerized (Fig. 3B). It is therefore likely that the inserted region from MOMP in the chimeric protein PorB$_{18}$ does not destabilize the quaternary structure.

CD spectra of PorB$_{WT}$ and PorB$_{18}$ revealed both proteins have β-sheet-like spectra with minima at 218 nm (Fig. 3C). PorB$_{18}$ spectra had higher β-sheet content than the PorB$_{WT}$ as determined

by JASCO multivariate SSE program (Fig. 3E). This supports that the inserted region is adding extra β-strands to the structure. The proteins exhibited similar thermal stability (T_m for $PorB_{WT}$ = 69 °C, T_m for $PorB_{18}$ = 68 °C) as determined by tracking temperature-dependent unfolding through CD absorption measurement at 218 nm (Fig.3D).

Figure 3: Analysis of in vitro $PorB_{WT}$ and $PorB_{18}$: (A) Reversible crosslinker DSP. **(B)** SDS-PAGE displaying DSP crosslinked purified proteins, with samples either untreated or treated with the reducer, β-mercaptoethanol to reverse crosslinking. Bands corresponding to $PorB_{WT}$ (green) and $PorB_{18}$ (blue). monomers are marked with (∗), dimer with (⁂), and trimer with (⁂⁑). **(C)** CD spectra of the proteins with error bars indicated by vertical lines. **(D)** Normalized CD spectra at 218 nm show the thermal unfolding of $PorB_{WT}$ (green) and $PorB_{18}$ (blue). The lines represent the

regression curve with a sigmoidal fit. Dotted vertical lines indicate the T_m. **(E)** Secondary structure estimation from the CD spectra.

Chimeric β-strands are buried

To assess if the extra β-strands are exposed or buried, $PorB_{WT}$ and $PorB_{18}$ folded in detergent micelles were subjected to Proteinase K for a longer duration. Proteins in detergent micelle are likely to be more susceptible to Proteinase K digestion as they are more exposed than in the outer membrane, where the LPS shields them. We found $PorB_{18}$ to be more susceptible to Proteinase K than $PorB_{WT}$ (Fig. 4A – B) as it is fully degraded after 60 minutes and $PorB_{WT}$ is not. To evaluate which regions of the proteins are exposed and are susceptible to Proteinase K, we analyzed the cleavage products with matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry (Fig. 4C & D).

Analysis of the cleavage products by MALDI-TOF mass spectrometry indicated one major cleavage product for $PorB_{WT}$ (Fig. 4E) at 34,364.9 Da. This product is difficult to differentiate from the uncleaved protein band on SDS-PAGE gel at this figure size but can be more clearly seen in a larger version (Fig. 4C). The peak at 28,887.5 Da is likely Proteinase K with a known molecular weight of 28.9 kDa. In contrast to the single cleavage product from $PorB_{WT}$, four major cleavage products are formed for $PorB_{18}$. (Fig. 4F). The additional peak at 20.2 kDa evident in 0 – 5 minute time points is the doubly charged uncleaved protein.

All possible Proteinase K cleavage sites were predicted using Expasy PeptideCutter [53] (Fig. S1). Using the predicted cleavage sites and the peaks from the mass spectra, the likely Proteinase K digest products were calculated (Fig. S2A & B). Despite having multiple Proteinase K cleavage sites, none of the calculated products start or end in the MOMP β-strands region (Fig 4G & S2B). Therefore, the inserted β-strands are likely to be buried in the structure.

Figure 4: Proteinase K digestion of in vitro folded PorB₁₈. SDS-PAGE of in vitro folded proteins **(A)** PorB_WT and **(B)** PorB₁₈, treated with Proteinase K for different durations. An enlarged view of the lanes from 5 min to 60 min is shown in panel **(C)** for PorB_WT and **(D)** for PorB₁₈. The black arrow marks the band that appears as a peak in MALDI-TOF. The purple arrow marks the Proteinase K band. MALDI-TOF of **(E)** PorB_WT (green) and **(F)** PorB₁₈ (blue) treated with Proteinase K for different durations. The numbers above the graph are the mass-by-charge ratio

of the peaks. Peaks indicated by black asterisks are doubly charged peaks of full-length PorB$_{18}$ seen at 40513.1. Peaks indicated by purple numbers are Proteinase K which has a molecular weight of 28.9 kDa. **(G)** The total number of proteinase K (PK) susceptible sites in PorB$_{18}$ predicted through Expasy PeptideCutter [53] is for each structural region (plain bars) is compared with all the possible cleavage sites calculated from the MALDI-TOF result (striped bars). The region of the added hairpin is highlighted in pink.

Chimeric strands increase the pore radius

Although the CD spectra show that the inserted region in PorB$_{18}$ is forming β-strands and the MALDI-TOF analysis of Proteinase K indicates that those strands are buried in the structure, neither demonstrates that these extra β-strands are becoming part of the β-barrel. Through ColabFold [54] we used AlphaFold-multimer [55, 56] to predict the structure of the chimeric protein (Fig 5A and B). The top two models both predict the inserted region to be buried within the structure, but one predicts the β-strands to be inside the lumen of the β-barrel forming a 16-stranded β-barrel (Fig 5A) while the second predicts the β-strands to be part of the β-barrel forming an 18-stranded β-barrel (Fig 5B). Notably, the confidence of the prediction as measured by the predicted local distance difference test (pLDDT) is very low for both predictions, However, the prediction confidence was higher when it predicted the structure to be 18-stranded (Fig. 5C). The theoretical CD spectra for the AlphaFold models where estimated using PDB2CD web server [57] (Fig. 5D). The 18-stranded PorB$_{18}$ model was estimated to be 61.7% β-sheet, and 16-stranded model was estimated to be 57.6% β-sheet. The theoretical CD for the PDB structure of PorB$_{WT}$ (PDB ID 4aui) was calculated to be 55.0% β-sheet.

Figure 5: AlphaFold2 prediction of PorB₁₈. (A and B) Top two ranked predicted structures of PorB₁₈: **(A)** 18-stranded prediction of PorB₁₈, **(B)** 16-stranded prediction of PorB₁₈. **(C)** Plots showing pLDDT scores per residue for each of the two predicted structures. The shaded grey area shows the insert region. **(D)** The predicted β-sheet percentage calculated from the experimental CD data of PorBWT and PorB18 is compared with the predicted β-sheet percentage from theoretical CD spectra of the PDB structure of PorBWT and the AlphaFold models calculated using the PDB2CD server [57]. **(E − G)** Channel prediction using CAVER 3.03 for 16-stranded

PorB$_{18}$ **(E)**, 18-stranded PorB$_{18}$ **(F)**, and PorB$_{WT}$ **(G)**. The number next to the cartoon is the bottleneck radius.

To verify which of the predicted structures is more likely, we investigated the pore size of the chimeric protein PorB$_{18}$. We computationally mapped out the pores with CAVER [58] for each of the FastRelaxed [59-61] predicted structures. The bottleneck radius for the predicted 16-stranded PorB$_{18}$ is 1.55 Å (Fig. 5E) and while for the 18-stranded PorB$_{18}$ is 3.86 Å (Fig. 5F). This is slightly larger than the bottleneck radius of 3.74 Å for PorB$_{WT}$ (Fig 5G).

To verify the pore size experimentally, we measured the channel conductance of PorB$_{WT}$ and PorB$_{18}$. The channel conductance of PorB$_{WT}$ was measured at 4.1 nS (Fig.6A). The experimental conductance for the chimeric PorB$_{18}$ was measured to be 4.4 nS (Fig. 6A). This is slightly more than PorB$_{WT}$ (Fig 6B).

The conductance of PorB$_{WT}$ shows frequent closing-opening of its channel when compared to PorB$_{18}$ (Fig 6A). This frequent channel closure is likely due to the presence of the N-terminal His-tag, which in PorB$_{WT}$ leads to brief obstructions in the pore opening. We confirmed the absence of channel closure in PorB$_{WT}$ when the N-terminal His-tag was absent (Fig. S3). The absence of channel closure in PorB$_{18}$, even with a His-tag present, possibly indicates that either the opening of the channel for PorB$_{18}$ is wide enough to be not affected by the His-tag or that the His-Tag is in conformation that does not interact with the channel. Despite the artifactual noisiness of the His-tagged PorB$_{WT}$, we compare His-tagged constructs as we could not purify PorB$_{18}$ without the His-tag.

The theoretical conductance PorB$_{18}$ AlphaFold models were calculated using equation 1 and by solving the unknown parameters based on the experimental conductance of PorB$_{WT}$ without the His-Tag (Fig 6C). The 18-stranded model has a theoretical conductance of 4.35 nS whereas the 16-stranded model has a theoretical conductance of 0.79 nS. The experimental conductance of

PorB$_{18}$ is much closer to the theoretical conductance of the 18-stranded AlphaFold model and much larger than that of the 16-stranded model (Fig 6C). This shows that the inserted region is not obstructing the pore and is likely integrating into the barrel to form a larger pore size.

To rule out the possibility that the higher conductance of PorB$_{18}$ is not from multiple PorB$_{18}$ trimers but rather from a single trimeric PorB$_{18}$, we measured the conductance at a higher voltage. The high voltage causes the pores of the porins to close and open at a much higher frequency. This causes multiple channels to close simultaneously, resulting in different levels of conductance. The distribution of conductance can provide insight into the number of individual porins present [34]. Both PorB$_{WT}$ and PorB$_{18}$ showed four levels of conductance providing evidence that the observed conductance is from a trimer (Fig 6D). The highest level of conductance for PorB$_{WT}$ is seen to be lower than 4.1nS because the frequency with which all three channels are open at the same time is very low, shifting the distribution of the conductance downward.

Figure 6: Single channel conductance of in vitro folded PorB₁₈ (A) Single channel conductance of PorB₁₈ (blue) compared to PorB_WT (green). **(B)** Box and whisker plot comparing the conductance of PorB_WT (green) and PorB₁₈(blue). ****: p-value <= 10⁻⁴ **(C)** Graph shows comparison of experimental single-channel conductance of PorB_WT with His-Tag (green) and without His-tag (light green), PorB₁₈ (blue), theoretical shingle-channel conductance of AlphaFold 18-stranded model (magenta) and 16-stranded model (cyan) of PorB₁₈. **(D)** Channel conductance

at 100 mV showing channel closing for individual subunits of PorB$_{WT}$ (green) and PorB$_{18}$ (blue).*Shortened loop constructs collapse*

To evaluate the importance of the long loops, shorter loop constructs for PorB$_{WT}$ and PorB$_{18}$ were created with their long loops shortened (loop 3 for PorB$_{WT}$ the MOMP loop 4 or PorB$_{18}$) (Fig. 7A). We called these shorter loop constructs, PorB$_{SL}$ and PorB$_{18SL}$. These were produced in *E. coli* with an outer membrane targeting signal sequence. Western blot of the whole cells showed that PorB$_{SL}$ does not express well indicating that loop L3 might be required for stable expression of PorB$_{WT}$ (Fig.7B). *In vitro* expression and folding showed PorB$_{SL}$ to be dimer rather than trimer (Fig. 7C). The shortened loop may be disrupting the interaction between loop1 and loop4 at the barrel-barrel interface (Fig. 7D).

While PorB$_{18SL}$ was able to express fold in *E. coli* outer membrane, it was found to be susceptible to Proteinase K (Fig. 7E). Single channel conductance of PorB$_{18SL}$ inserted into lipid bilayer membrane showed that its conductance is an order of magnitude smaller than PorB$_{18}$ (Fig. 7F). PorB$_{18SL}$ is likely collapsing without the long loop. This has been observed for engineered barrels without long loops [62].

CD spectra for the short loop constructs show a primarily β-sheet structure (Fig. S4A). The thermal stability of the short loop construct was comparable to PorB$_{WT}$ measured through CD absorption at 218 nm (Fig. S4B). Interestingly, the small rise in CD absorption at 218 nm seen between 20 °C and 60 °C for PorB$_{WT}$ (Fig 3D) is absent in PorB$_{SL}$ (Fig. S4B). This suggests that loop L3 undergoes temperature-dependent structural change. Secondary structure estimates from the CD spectra indicate a higher β-strand percentage for both PorB$_{SL}$ and PorB$_{18SL}$ (Fig. S4C) compared to the PorB$_{WT}$ (Fig. 3E), as expected due to the removal of the loops.

Figure 7: Loop shortening in PorB$_{WT}$ and Por$_{18}$ leads to pore collapse. (A) Top sequence alignment shows the short loop construct of PorB$_{WT}$ with L3 deletion named PorB$_{SL}$, compared with PorB$_{WT}$. The bottom three sequences compare the short loop construct of PorB$_{18}$ with L4 deletion named PorB$_{18SL}$, compared with PorB$_{WT}$. and PorB$_{18}$. Strand sequences from PorB$_{WT}$ are in black and loops are in green. Strand and loop sequences taken from MOMP are in blue and red respectively. **(B)** Western blot showing expression levels of PorB$_{SL}$ and PorB$_{18SL}$ in whole cells. **(C)** DSP crosslinking of the peak fractions from SEC. Monomeric bands are marked with (∗)

and dimers with (⚛). **(D)** Western blot showing PorB$_{18}$ is susceptible to Proteinase K digestion. **(E)** Cartoon of PorB$_{WT}$ trimer showing loops L1 to L4 (PDB ID 4aui) [39]. **(F)** Single channel conductance of PorB$_{18SL}$

Discussion

Here, we engineer a larger pored β-barrel from PorB by inserting domains from another β-barrel protein from a different species (MOMP). Despite having a large section modified the chimeric protein PorB$_{18}$ is found to be stable likely due to the evolutionary relation between PorB and MOMP [21]. Outer membrane β-barrels have been engineered into larger-sized barrels through duplication of existing strands [19, 29-32]. These methods mimic the gene duplication of topology accretion in repeat proteins which are generally considered to be the predominant way of protein evolution. OMP evolution with a loop-to-hairpin mode of strand accretion has not been explored experimentally before. Our results with PorB$_{18}$ support that this mode of strand accretion can also be used to engineer larger barrels and was likely to have been used by evolution.

Biophysical assays indicate that the chimeric PorB$_{18}$ has more β-strand content and the strands are integrating into the β-barrel resulting in a larger pore size. While we did not have an experimental structure for the chimeric protein PorB$_{18}$, the increase in single-channel conductance when compared to PorB$_{WT}$ closely matches the increase in β-structure and increase in predicted pore size for the 18-stranded AlphaFold prediction for PorB$_{18}$. This supports that the chimeric protein is 18-stranded compared to the 16-stranded wild type, enlarging the β-barrel.

While we tested the loop-to-β-hairpin evolution between porins PorB and MOMP, there are other evolutionarily related pairs that could also be tested to determine how widely this method can be used.

In the case of PorB we observed that the lumen-lining long loop is necessary for the stability of the protein. Deletion of loop L3 prevented PorB from trimerizing. We hypothesize that the truncated loop is preventing interaction between loop L2 and L4 of neighboring monomers, destabilizing the trimer. This can be further explored with mutational studies to verify the importance of this interaction for the trimerization of PorB$_{WT}$.

The fact that an extracellular loop can easily be converted into a transmembrane β-hairpin brings OMP folding questions to the fore. Future studies may clarify if the barrel folds synchronously or if the original 16 strands fold first and then the newer strands are added. Moreover, it remains possible that some loops are transiently hairpins, or some hairpins are transiently loops. Such a transition within a protein has been found within soluble proteins [63] though has not yet been found in membrane proteins. In membrane β-barrels such a transition would bring previously unknown plasticity to the barrel radius allowing for the passage of larger solutes or acting as a gating mechanism.

Conclusion

Here we show the first experimental evidence of the evolutionarily unusual topology accretion through the loop-to-hairpin transition in OMBB evolution using a PorB chimeric protein. This provides proof for one such protein evolutionary pair and opens up the avenue to explore more of such relationships in outer membrane β barrels as well as other protein structures.

Materials and Methods:

Cloning and in vivo expression:

Plasmids for recombinant protein expression were constructed using mega primer insertion PCR [64] using the Plasmid Champion pET303/CT-His vector (Invitrogen). The double-stranded DNA

inserts (Integrated DNA Technologies) consist of the codon-optimized signal peptide from *E. coli* OmpA protein, a 6x Histidine tag, a TEV protease cleavage site, the recombinant protein, two stop codons, and two 50-nucleotide homologous regions from the vector flanking on each side. NEB Turbo *E. coli* strain (New England Biolabs) was used for cloning and plasmid propagation.

E. coli ΔDegP strain (KS476 strain acquired from CGSC)[65] was modified by integration of λDE3 prophage using the Novagen λDE3 Lysogenization Kit (MilliporeSigma) to allow for the use of T7 polymerase-based plasmid vectors. The plasmid vector for the recombinant protein was transformed on LB agar plates with carbenicillin as the selection antibiotic. A single colony was resuspended in 2xYT media with antibiotics grown overnight at 37 °C at 250 RPM shaking incubator. Protein expression was achieved in these overnight cultures through the leaky expression of the pET303 vector (without any inducing agent).

Outer membrane fractionation

Overnight cultures were collected by centrifugation at 3000 g for 10 min at 4 °C. The cell pellets were then resuspended in lysis buffer (50 mM Tris, pH 7.4, 200 mM NaCl, 10 mM MgCl$_2$, 5 mM Benzamidine, 0.5 mM PMSF, 5 µg/mL DNase). Resuspended cells were freeze-thawed thrice by freezing them at -80 °C and rapidly thawing at 37 °C. Resuspended cells were then lysed using a sonicator (Qsonica Q500 Sonicator) submerged in an ice bath. Sonication was carried out at 40% amplitude with a pulse of 2 sec on and 4 sec off for a total 'on time' of 5 min. The outer membrane was fractionated as previously described [66] with some modifications. Specifically, the inner membrane solubilization buffer was modified to 20 mM Sodium Phosphate, pH 7.4, 5 mM Benzamidine, and 2% N-Lauroylsarcosine. The outer membrane solubilization buffer was modified to 20 mM NaPi pH 6.5, 100 mM NaCl, 5 mM Benzamidine, 2 µM Lysozyme, and 1% LDAO. The outer membrane was solubilized for 30 min at room temperature and then separated

from non-soluble fractions by centrifugation at 15,600 g for 30 min. Fractions were analyzed by western blot.

In vivo Proteinase K assay

The *in vivo* Proteinase K assay was conducted as previously described [50] with some modifications. In brief, 5 mL of overnight cultures expressing the recombinant protein were collected by centrifugation (3000 g, 5 min, 4 °C). The cells were permeabilized by resuspending the cell pellets in the permeabilization buffer (33 mM Tris pH 8.0, 40% sucrose, 2 µM lysozyme, and 2 mM EDTA) and incubating on ice for 20 min. Half of the cell suspensions were treated with 3.5 µM Proteinase K (Sigma-Aldrich) and half were left untreated as a control. The digestion was stopped with 1 mM PMSF. The cell suspensions were precipitated with 10% TCA. The precipitates were collected by centrifugation at 20,000 g for 5 min. The precipitates were washed once with 100% acetone and twice with 50 mM Tris pH 8.0, and 200 mM NaCl. Precipitates were mixed with 2x Laemmli sample buffer (BioRad) and boiled for 10 min at 98 °C to prepare them for western blot.

Crosslinking assay

Protein eluates from SEC were crosslinked at 2.5 µM concentration with 250 µM dithiobis succinimidyl propionate (DSP) (Thermo Scientific) at room temperature for 30 min. The reactions were quenched with 50 mM final concentration of Tris-HCl pH 7.5 for 15 min. Aliquot from each reaction was mixed with 2x Laemmli SDS sample buffer (Bio-Rad) for SDS-PAGE analysis.

SDS-PAGE and western blot

Purified proteins, cell fractions, or whole cells were mixed with 2x Laemmli sample buffer (BioRad) and loaded onto a 4-20% polyacrylamide gel (Bio-Rad). Electrophoresis was carried out at 300

V, for 15 min. Gels were stained with ReadyBlue Coomassie protein gel stain (MilliporeSigma) and imaged on Gel Doc EZ imaging system (BioRad)

For the Western blot, after the electrophoresis step, gels were sandwiched with 0.2 µm nitrocellulose blotting membrane (BioRad), and the transfer was carried out at 100 V for 30 min on ice. The transfer buffer was prepared with 10x Tris/Glycine buffer (BioRad) and 10% Ethanol. To verify the transfer, blots were stained with 0.2% (w/v) Ponceau S in 5% glacial acetic acid. Blots were washed thrice in TBST (20 mM Tris, pH 7.5, 150 mM NaCl, 0.1% Tween 20) and blocked with TBST with 1% gelatin from cold water fish skin (MilliporeSigma). The following antibodies were diluted as per manufacturer recommendation with TBST with 1% gelatin: THE Anti-His tag monoclonal mouse antibody (GenScript), anti-*E. coli* SurA polyclonal rabbit antibody (Cusabio), anti-*E. coli* BamA polyclonal rabbit antibody (Cusabio), anti-*E. coli* GroEL polyclonal rabbit antibody (Enzo Life Sciences), IRDye 800CW goat anti-mouse IgG secondary antibody (LI-COR Biosciences), IRDye 680RD donkey anti-rabbit IgG secondary antibody (LI-COR Biosciences). anti-*E. coli* TolC was purified from rabbit sera and diluted to 1:2500 with 1% gelatin in TBST. All incubations were carried out at 37 °C for 30 min, followed by three 10 min rinses with TBST at 37 °C. Blots were imaged with LI-COR Odyssey infrared imager at 700 nm and 800 nm.

Protein expression, refolding, and purification

Recombinant proteins were expressed in *E. coli* BL21(DE3) as inclusion bodies by removing the OmpA signal sequence. Protein expression was induced at a culture density of 0.6 OD600 with 1 mM IPTG. Cells were collected by centrifugation at 4000 g for 30 min. Inclusion bodies were harvested by lysing the cell suspension in ice-cold lysis buffer (50 mM Tris pH 8, 200 mM NaCl, 1 mM EDTA, 1 mM phenylmethylsulfonyl fluoride (PMSF)) with 2% TritonX-100. Lysis was carried out by sonication at 50% amplitude, with a pulse of 2 sec on and 2 sec off for a total 'on time' of

5 min per litter of cells, submerged in an ice bath. Cells were washed thrice by centrifugation at 11000 g for 10 min, followed by resuspension in ice-cold lysis buffer without TritonX-100.

Inclusion bodies of his-tagged protein were solubilized in binding buffer (20 mM NaPi, pH 6.5, 100 mM NaCl, 20 mM imidazole, 7.2 M urea). Solubilized inclusion bodies were purified by metal affinity chromatography using a precharged Ni Sepharose 6 Fast Flow prepacked column (Cytiva) on an ÄKTA pure FPLC system (Cytiva). Protein was eluted with a gradient of 0 – 50 % elution buffer (20 mM NaPi, pH 6.5, 100 mM NaCl, 500 mM imidazole, 7.2 M urea). Protein eluent was concentrated using 10 kDa Amicon Ultra-15 centrifugal filter units (MilliporeSigma) to a maximum concentration of 20 mg/mL for $PorB_{WT}$ and 7 mg/mL for $PorB_{18}$. The protein concentrate was refolded by mixing it with refolding buffer (10% LDAO, 20 mM NaPi pH 6.5, 100 mM NaCl, 1 mM EDTA) at a 1:2 ratio [52]. Freshly folded protein was separated from aggregates using size exclusion chromatography using a prepacked Superdex 200 10/300 GL column (Cytiva) equilibrated with running buffer (0.1% LDAO, 20 mM NaPi pH 6.5, 100 mM NaCl, 1 mM EDTA).

Circular dichroism

Samples for circular dichroism (CD) were prepared by buffer exchanging the purified protein into CD buffer (0.1% LDAO, 20 mM NaPi pH 6.5, 100 mM NaF) using a Superdex 200 10/300 GL column (Cytiva). The eluent was concentrated using 30 kDa Amicon Ultra-0.5 centrifugal filter devices (MilliporeSigma) to a concentration of 1 mg/mL. Protein samples were loaded onto a 0.1 mm pathlength Suprasil quartz cuvette (Hellma) for CD measurement.CD spectra were collected using a JASCO J-815-150S CD spectrometer with the following parameters: wavelength range of 260-175 nm, speed of 100 nm/min, data pitch of 0.5 nm, D.I.T of 2 sec, bandwidth of 1 nm, and 7 accumulations for each sample. The secondary structure was estimated using JASCO's multivariate secondary structure estimation program. Thermal melt curves were generated by

using the spectrophotometer with a JASCO PTC-423 Peltier controller accessory. CD spectra were collected at different temperatures starting from 20 °C and up to 90 °C at 5 °C increments.

Single channel conductance

The single-channel setup includes a BC-535 bilayer clamp amplifier from Warner Instruments, National Instruments' PCI-6221 DAQ card for data acquisition, and National Instruments' BNC 2110 digitizer. Data was recorded using WinEDR software (J. Dempster, University of Strathclyde, UK). The bilayer chambers consist of Warner instruments BCH-M13 derlin chamber and CD13A-200 derlin cuvette with 200 μm aperture. 1.4 mL 1 M KCl was added to each chamber as well as the electrode chambers. The electrode chambers were connected to the main chambers using 3 M KCl, 2% agar bridges. The cis chamber was connected to the input and a trans chamber was connected to the ground. The bilayer was formed by 'painting' a solution of 10 mg/mL 1,2-Diphytanoyl-sn-glycero-3-phosphocoline in n-decane over the aperture. Bilayer formation was confirmed by observing an increase in membrane capacitance to 80 pF to 110 pF. 2 μL of 1 mg/mL protein sample was added to the cis chamber near the aperture. A voltage of 100 mV was applied for incorporating the protein into the bilayer. The theoretical conductance was calculated using the following previously derived equation[67]:

$$d = \frac{\rho g}{\pi}\left(\frac{\pi}{2} + \sqrt{\frac{\pi^2}{4} + \frac{4\pi l}{\rho g}}\right)$$

(1)

Where, d is the diameter of the pore, g is the conductance of the pore, ρ is the resistivity of the buffer, and l is the length of the pore.

In vitro Proteinase K and mass spectrometry

For *in vitro* Proteinase K digests, purified proteins at a concentration of 2 µM were treated with 0.1 µM of Proteinase K and placed on ice. PMSF was added at 1 mM concentration to stop the digestion at certain time points. Samples were mixed with 2x Laemmli sample buffer (BioRad) for SDS-PAGE analysis.

Proteinase K treated samples were analyzed using matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry. Samples were made acidic by adding TFA to 0.1% final concentration and then desalted using a ZipTip with C_4 resin (MilliporeSigma). The desalted protein was eluted with solvent TA30 (30:70 [v/v] acetonitrile/0.1% TFA in water) and mixed with HCCA matrix solution (supersaturated HCCA solution in solvent TA30) at a 1:1 ratio. 1 µL of the protein matrix mixture was deposited on a 96-spot, polished, stainless steel target plates (Bruker Daltonics) and dried. For calibration, Protein Standard II (Bruker Daltonics) was prepared with the HCCA matrix solution and 1 µL was deposited next to the protein sample spots. The protein mass to charge spectra were acquired on a MicroFlex LT MALDI-TOF system (Bruker Daltonics) with 1000-2000 shots using the instrument's 337 nm nitrogen laser.

Structure prediction and pore mapping

To predict the structure of $PorB_{18}$, we used AlphaFold v2.3.1 [55, 56] through ColabFold v1.5.2 [54], using MOMP as a template (See supplementary text for configuration). The top two predictions with the highest pLDDT were relaxed in RosettaScripts [61] FastRelax protocol [59, 60]. The relaxed structure with the lowest Rosetta score was used for selecting for mapping its pore using CAVER 3.03 PyMol plugin [58]. The starting point for the tunnel was selected on the periplasmic side of the protein. The correct path was selected visually for the bottleneck radius of the pore.

Data analysis and graphs

Single-channel data analysis and Proteinase K digest product calculations were done using Python 3.10.7 and the Pandas 1.3.4 library [68]. Graphs for these were generated using Python libraries Seaborn v0.11.2 [69] and Matplotlib v3.5.1 library [70]. Chromatography and MALDI-TOF data were graphed using GraphPad Prism 9.5.1 (GraphPad Software, San Diego, California USA, www.graphpad.com).

Acknowledgments

We thank Ayotunde Paul Ikujuni, Jimmy Budiardjo, Ryan Feehan, and Daniel Montezano for their helpful insights and discussion. We also gratefully acknowledge NIGMS awards DP2GM128201 and P20GM113117, and P20GM103418 and NSF 2226804 to JSGS.

Reference

[1] Roumia AF, Theodoropoulou MC, Tsirigos KD, Nielsen H, Bagos PG. Landscape of Eukaryotic Transmembrane Beta Barrel Proteins. J Proteome Res. 2020;19:1209-21.

[2] Paschen SA, Waizenegger T, Stan T, Preuss M, Cyrklaff M, Hell K, et al. Evolutionary conservation of biogenesis of beta-barrel membrane proteins. Nature. 2003;426:862-6.

[3] Koebnik R, Locher KP, Van Gelder P. Structure and function of bacterial outer membrane proteins: barrels in a nutshell. Mol Microbiol. 2000;37:239-53.

[4] Schulz GE. The structure of bacterial outer membrane proteins. Biochim Biophys Acta. 2002;1565:308-17.

[5] Franklin MW, Slusky JSG. Tight Turns of Outer Membrane Proteins: An Analysis of Sequence, Structure, and Hydrogen Bonding. J Mol Biol. 2018;430:3251-65.

[6] Piggot TJ, Holdbrook DA, Khalid S. Conformational dynamics and membrane interactions of the E. coli outer membrane protein FecA: a molecular dynamics simulation study. Biochim Biophys Acta. 2013;1828:284-93.

[7] Van Gelder P, Saint N, van Boxtel R, Rosenbusch JP, Tommassen J. Pore functioning of outer membrane protein PhoE of Escherichia coli: mutagenesis of the constriction loop L3. Protein Eng. 1997;10:699-706.

[8] Liu NZ, Delcour AH. The spontaneous gating activity of OmpC porin is affected by mutations of a putative hydrogen bond network or of a salt bridge between the L3 loop and the barrel. Protein Engineering. 1998;11:797-802.

[9] Kramer RA, Zandwijken D, Egmond MR, Dekker N. In vitro folding, purification and characterization of Escherichia coli outer membrane protease ompT. Eur J Biochem. 2000;267:885-93.

[10] Leonard-Rivera M, Misra R. Conserved residues of the putative L6 loop of Escherichia coli BamA play a critical role in the assembly of β-barrel outer membrane proteins, including that of BamA itself. Journal of bacteriology. 2012;194:4662-8.

[11] Phale PS, Philippsen A, Kiefhaber T, Koebnik R, Phale VP, Schirmer T, et al. Stability of trimeric OmpF porin: the contributions of the latching loop L2. Biochemistry. 1998;37:15663-70.

[12] Roumia AF, Tsirigos KD, Theodoropoulou MC, Tamposis IA, Hamodrakas SJ, Bagos PG. OMPdb: A Global Hub of Beta-Barrel Outer Membrane Proteins. Front Bioinform. 2021;1:646581.

[13] Horne JE, Brockwell DJ, Radford SE. Role of the lipid bilayer in outer membrane protein folding in Gram-negative bacteria. J Biol Chem. 2020;295:10340-67.

[14] McLachlan AD. Repeating sequences and gene duplication in proteins. J Mol Biol. 1972;64:417-37.

[15] Barker WC, Ketcham LK, Dayhoff MO. A comprehensive examination of protein sequences for evidence of internal gene duplication. J Mol Evol. 1978;10:265-81.

[16] McLachlan A. Gene duplication and the origin of repetitive protein structures. Cold Spring Harbor symposia on quantitative biology: Cold Spring Harbor Laboratory Press; 1987. p. 411-20.

[17] Stephens S. Possible significance of duplication in evolution. Advances in genetics: Elsevier; 1951. p. 247-65.

[18] Zhang JZ. Evolution by gene duplication: an update. Trends in Ecology & Evolution. 2003;18:292-8.

[19] Arnold T, Poynor M, Nussberger S, Lupas AN, Linke D. Gene duplication of the eight-stranded beta-barrel OmpX produces a functional pore: a scenario for the evolution of transmembrane beta-barrels. J Mol Biol. 2007;366:1174-84.

[20] Magadum S, Banerjee U, Murugan P, Gangapur D, Ravikesavan R. Gene duplication as a major force in evolution. Journal of genetics. 2013;92:155-61.

[21] Franklin MW, Nepomnyachyi S, Feehan R, Ben-Tal N, Kolodny R, Slusky JS. Evolutionary pathways of repeat protein topology in bacterial outer membrane proteins. Elife. 2018;7.

[22] Dhar R, Slusky JS. Outer membrane protein evolution. Curr Opin Struct Biol. 2021;68:122-8.

[23] Alva V, Söding J, Lupas AN. A vocabulary of ancient peptides at the origin of folded proteins. Elife. 2015;4:e09410.

[24] Söding J, Lupas AN. More than the sum of their parts: on the evolution of proteins from peptides. Bioessays. 2003;25:837-46.

[25] Neuwald AF, Liu JS, Lawrence CE. Gibbs motif sampling: detection of bacterial outer membrane protein repeats. Protein science. 1995;4:1618-32.

[26] Remmert M, Biegert A, Linke D, Lupas AN, Söding J. Evolution of outer membrane β-barrels from an ancestral ββ hairpin. Molecular biology and evolution. 2010;27:1348-58.

[27] Reddy BL, Saier Jr MH. Properties and phylogeny of 76 families of bacterial and eukaryotic organellar outer membrane pore-forming proteins. PloS one. 2016;11.

[28] Dhar R, Feehan R, Slusky JS. Membrane barrels are taller, fatter, inside-out soluble barrels. The Journal of Physical Chemistry B. 2021;125:3622-8.

[29] Krewinkel M, Dworeck T, Fioroni M. Engineering of an E. coli outer membrane protein FhuA with increased channel diameter. J Nanobiotechnology. 2011;9:33.

[30] Liu Z, Ghai I, Winterhalter M, Schwaneberg U. Engineering enhanced pore sizes using FhuA Δ1-160 from E. coli outer membrane as template. ACS sensors. 2017;2:1619-26.

[31] Srivastava SR, Mahalakshmi R. Evolutionary selection of a 19-stranded mitochondrial beta-barrel scaffold bears structural and functional significance. J Biol Chem. 2020;295:14653-65.

[32] Charbit A, Andersen C, Wang J, Schiffler B, Michel V, Benz R, et al. In vivo and in vitro studies of transmembrane beta-strand deletion, insertion or substitution mutants of the Escherichia coli K-12 maltoporin. Mol Microbiol. 2000;35:777-90.

[33] Korkmaz F, van Pee K, Yildiz O. IR-spectroscopic characterization of an elongated OmpG mutant. Arch Biochem Biophys. 2015;576:73-9.

[34] Lin M, Zhang G, Fahie M, Morgan LK, Chen M, Keiderling TA, et al. Engineering a Novel Porin OmpGF Via Strand Replacement from Computational Analysis of Sequence Motif. Biochim Biophys Acta Biomembr. 2017;1859:1180-9.

[35] Chaturvedi D, Mahalakshmi R. Folding Determinants of Transmembrane beta-Barrels Using Engineered OMP Chimeras. Biochemistry. 2018;57:1987-96.

[36] Zeth K, Thein M. Porins in prokaryotes and eukaryotes: common themes and variations. Biochem J. 2010;431:13-22.

[37] Jackups Jr R, Liang J. Interstrand pairing patterns in β-barrel membrane proteins: the positive-outside rule, aromatic rescue, and strand registration prediction. Journal of molecular biology. 2005;354:979-93.

[38] Stapleton JA, Whitehead TA, Nanda V. Computational redesign of the lipid-facing surface of the outer membrane protein OmpA. Proc Natl Acad Sci U S A. 2015;112:9632-7.

[39] Zeth K, Kozjak-Pavlovic V, Faulstich M, Fraunholz M, Hurwitz R, Kepp O, et al. Structure and function of the PorB porin from disseminating Neisseria gonorrhoeae. Biochem J. 2013;449:631-42.

[40] Ferrara LGM, Wallat GD, Moynie L, Dhanasekar NN, Aliouane S, Acosta-Gutierrez S, et al. MOMP from Campylobacter jejuni Is a Trimer of 18-Stranded beta-Barrel Monomers with a Ca(2+) Ion Bound at the Constriction Zone. J Mol Biol. 2016;428:4528-43.

[41] Freudl R, Schwarz H, Degen M, Henning U. The signal sequence suffices to direct export of outer membrane protein OmpA of Escherichia coli K-12. J Bacteriol. 1987;169:66-71.

[42] Macintyre S, Freudl R, Eschbach ML, Henning U. An Artificial Hydrophobic Sequence Functions as Either an Anchor or a Signal Sequence at Only One of 2 Positions within the Escherichia-Coli Outer-Membrane Protein-Ompa. Journal of Biological Chemistry. 1988;263:19053-9.

[43] Wolff K, Stern A. The class 3 outer membrane protein (PorB) of Neisseria meningitidis: gene sequence and homology to the gonococcal porin PIA. FEMS Microbiol Lett. 1991;67:179-85.

[44] Carbonetti NH, Sparling PF. Molecular cloning and characterization of the structural gene for protein I, the major outer membrane protein of Neisseria gonorrhoeae. Proc Natl Acad Sci U S A. 1987;84:9084-8.

[45] Zhang Z, Kuipers G, Niemiec L, Baumgarten T, Slotboom DJ, de Gier JW, et al. High-level production of membrane proteins in E. coli BL21(DE3) by omitting the inducer IPTG. Microb Cell Fact. 2015;14:142.

[46] Ferrara LG, Wallat GD, Moynié L, Dhanasekar NN, Aliouane S, Acosta-Gutiérrez S, et al. MOMP from Campylobacter jejuni is a trimer of 18-stranded β-barrel monomers with a Ca2+ ion bound at the constriction zone. Journal of molecular biology. 2016;428:4528-43.

[47] Franklin MW. Drawing Insight from Structure to Guide Design [Doctoral Dissertation]. Ann Arbor: University of Kansas; 2019.

[48] Ebeling W, Hennrich N, Klockow M, Metz H, Orth HD, Lang H. Proteinase K from Tritirachium album Limber. Eur J Biochem. 1974;47:91-7.

[49] Sweeney PJ, Walker JM. Proteinase K (EC 3.4.21.14). Methods Mol Biol. 1993;16:305-11.

[50] Wang X, Peterson JH, Bernstein HD. Bacterial Outer Membrane Proteins Are Targeted to the Bam Complex by Two Parallel Mechanisms. mBio. 2021;12:e00597-21.

[51] Werner J, Augustus AM, Misra R. Assembly of TolC, a structurally unique and multifunctional outer membrane protein of Escherichia coli K-12. J Bacteriol. 2003;185:6540-7.

[52] Tanabe M, Nimigean CM, Iverson TM. Structural basis for solute transport, nucleotide regulation, and immunological recognition of Neisseria meningitidis PorB. Proc Natl Acad Sci U S A. 2010;107:6811-6.

[53] Gasteiger E, Hoogland C, Gattiker A, Wilkins MR, Appel RD, Bairoch A. Protein identification and analysis tools on the ExPASy server. The proteomics protocols handbook. 2005:571-607.

[54] Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making protein folding accessible to all. Nature Methods. 2022;19:679-+.

[55] Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, et al. Protein complex prediction with AlphaFold-Multimer. BioRxiv. 2021:2021.10. 04.463034.

[56] Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate protein structure prediction with AlphaFold. Nature. 2021;596:583-9.

[57] Mavridis L, Janes RW. PDB2CD: a web-based application for the generation of circular dichroism spectra from protein atomic coordinates. Bioinformatics. 2017;33:56-63.

[58] Pavelka A, Sebestova E, Kozlikova B, Brezovsky J, Sochor J, Damborsky J. CAVER: algorithms for analyzing dynamics of tunnels in macromolecules. IEEE/ACM transactions on computational biology and bioinformatics. 2015;13:505-17.

[59] Khatib F, Cooper S, Tyka MD, Xu K, Makedon I, Popovic Z, et al. Algorithm discovery by protein folding game players. Proc Natl Acad Sci U S A. 2011;108:18949-53.

[60] Maguire JB, Haddox HK, Strickland D, Halabiya SF, Coventry B, Griffin JR, et al. Perturbing the energy landscape for improved packing during computational protein design. Proteins. 2021;89:436-49.

[61] Fleishman SJ, Leaver-Fay A, Corn JE, Strauch EM, Khare SD, Koga N, et al. RosettaScripts: a scripting language interface to the Rosetta macromolecular modeling suite. PLoS One. 2011;6:e20161.

[62] Zhang Z, Ryoo D, Balusek C, Acharya A, Rydmark MO, Linke D, et al. Inward-facing glycine residues create sharp turns in beta-barrel membrane proteins. Biochim Biophys Acta Biomembr. 2021;1863:183662.

[63] Lou J, Zhu C. Flow induces loop-to-β-hairpin transition on the β-switch of platelet glycoprotein Ibα. Proceedings of the National Academy of Sciences. 2008;105:13847-52.

[64] Barik S. Site-directed mutagenesis in vitro by megaprimer PCR. Methods Mol Biol. 1996;57:203-15.

[65] Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol. 2006;2:2006 0008.

[66] Leo JC, Oberhettinger P, Linke D. Assessing the Outer Membrane Insertion and Folding of Multimeric Transmembrane beta-Barrel Proteins. Methods Mol Biol. 2015;1329:157-67.

[67] Cruickshank CC, Minchin RF, Le Dain AC, Martinac B. Estimation of the pore size of the large-conductance mechanosensitive ion channel of Escherichia coli. Biophys J. 1997;73:1925-31.

[68] The pandas development team. Pandas. v1.2.0 ed: Zenodo; 2020.

[69] Michael Waskom and the seaborn development team. Seaborn. v0.11.1 ed: Zenodo; 2020.

[70] Hunter JD. Matplotlib: A 2D graphics environment. Computing in Science & Engineering. 2007;9:90-5.