The paradox of extremely fast evolution driven by genetic drift in multi-copy gene systems

Xiaopei Wang[1], Yongsen Ruan[1], Lingjie Zhang[1], Xiangnyu Chen[1], Zongkun Shi[1], Haiyu Wang[1], Bingjie Chen[1], Miles Tracy[1], Liying Huang[1], Chung-I Wu[1,*] and Haijun Wen[1,*]

[1]State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, China

[*] Corresponding authors: wenhj5@mail.sysu.edu.cn (H. Wen); ciwu@uchicago.edu (C.I. Wu)

ABSTRACT

Multi-copy gene systems that evolve within, as well as between, individuals are common. They include viruses, mitochondrial DNAs, multi-gene families etc. The paradox is that neutral evolution in two stages should be far slower than single-copy systems but the opposite is often true, thus leading to the suggestion of natural selection. We now apply the new Generalized Haldane (GH) model to quantify genetic drift in the mammalian ribosomal RNA genes (or rDNAs). On average, rDNAs have $C \sim 150$ - 300 copies. A neutral mutation in rDNA should take $4NC^*$ generations to become fixed (N, the population size; C^*, the effective copy number). While $C > C^* \gg 1$ is expected, the observed fixation time in mouse and human is $< 4N$, hence the paradox of $C^* < 1$. Genetic drift thus appears as much as 100 times stronger for rRNA genes as for single-copy genes. The large increases in genetic drift are driven by a host of molecular mechanisms such as gene conversion and unequal crossover. Although each mechanism of drift has been extremely difficult to quantify, the GH model permits the estimation of their total effects on genetic drift. In conclusion, the GH model can be generally applicable to multi-copy gene systems without being burdened by tracking the diverse molecular mechanisms individually.

Introduction

In this study, we focus on multi-copy gene systems, where the evolution takes place in two stages: both within (stage I) and between individuals (stage II). Multi-copy gene systems include viruses, transposons, mitochondria and multi-gene families (Alexandrov, et al. 2001; Szitenberg, et al. 2016; Xu, et al. 2019; Ruan, et al. 2021). Given the extra stage of within-host fixation, the neutral evolutionary rate of multi-copy systems should be much slower than in single-copy systems. However, the rapid evolution of multi-copy systems has been extensively documented (Charlesworth, et al. 1994; Eickbush and Eickbush 2007; Jurka, et al. 2007; Hou, et al. 2023) but the reason is unclear.

The speed of neutral evolution is the basis for determining how fast or slow all types of molecular evolution take place. Neutral evolution is driven by random transmission, gene conversion, stochastic replication etc., which collectively constitute genetic drift. Hence, genetic drift is the fundamental force of molecular evolution. All other evolutionary forces, such as selection, mutation and migration, may be of greater biological interest, but inferences are possible only when genetic drift is fully accounted for. In the companion study, we show that the standard Wright-Fisher (WF) model may often under-account genetic

drift, thus leading to the over-estimation of selection.

The companion paper (Ruan, et al. 2024) re-introduces and generalizes the Haldane model to account for all
random forces in molecular evolution. This generalized Haldane model is referred to as the GH model. The
Haldane (as well as GH) model is based on the branching process. In haploids, each individual produces K
progeny with the mean and variance of $E(K)$ and $V(K)$. Genetic drift is primarily $V(K)$ as there would be no
drift if $V(K) = 0$. Gene frequency change in the population is then scaled by N (population size), expressed
as $V(K)/N$(Chen, et al. 2017). In diploids, K would be the number of progenies to whom the gene copy of
interest is transmitted. (The adjustments between haploidy and multi-ploidy are straightforward).

An important element missing in the WF models is the generation of N within the model. All WF models
including subsequent modifications have to be supplied N's externally. The Haldane model is capable of
generating $N' = N E(K)$ where N' is the population size of the next generation. However, the original
Haldane (Haldane 1927; Haldane 1932) model reset $N' = N$ thus depriving the branching process an
important advantage. The GH model permits $E(K)$ to depend on N and is thus capable of generating and
regulating N as well as tracking gene frequencies (see (Ruan, et al. 2024)).

Furthermore, in the WF model, gene frequency is governed by $1/N$ (or $1/2N$ in diploids) because K would
follow the Poisson distribution whereby $V(K) = E(K)$. As $E(K)$ is generally ~1 in evolutionary timescale,
$V(K)$ would also be ~ 1. In this backdrop, many "modified WF" models have been developed (Der, et al.
2011), most of them permitting $V(K) \neq E(K)$ (Karlin and McGregor 1964; Chia and Watterson 1969;
Cannings 1974). Nevertheless, paradoxes encountered by the standard WF model apply to these modified
WF models as well, because all WF models share the key feature of being supplied N externally (see below
and (Ruan, et al. 2024)). One of the paradoxes, first noted in (Chen, et al. 2017) is genetic drift during tumor
growth whereby drift appears to become stronger as N increases (Wu, et al. 2016; Chen, Wu, et al. 2022;
Zhai, et al. 2022). This trend is in stark opposite to the central tenet of the WF models.

A paradox requiring dedicated efforts to analyze concerns multi-copy gene systems, which are as diverse as
viral epidemics (Huang, et al. 2021), transposons (Szitenberg, et al. 2016), mitochondrial DNAs (Xu, et al.
2019), satellite DNAs (Cabot, et al. 1993; Alexandrov, et al. 2001), and ribosomal RNA genes (van Sluis and
McStay 2019; Hori, et al. 2021). In COVID-19, the inability of the WF models to track both within- and
between-host evolution simultaneously is a main reason for much confusion about the origin, spread and
driving forces of SARS-CoV-2 (Ruan, et al. 2021; Deng, et al. 2022; Guan and Zhong 2022; Pan, Liu, et al.
2022; Ruan, Wen, et al. 2022; Zhou, et al. 2022; Hou, et al. 2023).

We now apply the GH model to multi-copy gene systems, using ribosomal RNA genes (rDNAs) as an
example. In such systems, the molecular mechanisms driving neutral evolution within individuals are
diverse. It has been difficult to model each mechanism individually, let alone in combination for rDNAs
(Nagylaki 1983; Ohta and Dover 1983; Ohta and Dover 1984), viruses (Ruan, et al. 2021; Ruan, Hou, et al.
2022) and mitochondria DNA (Rand 2001; Xu, et al. 2019). Since our interest is to filter out the total effects
of genetic drift in order to study the more interesting forces such as selection, mutation etc, the GH model is
applied for this purpose. The details of each molecular mechanism will be missed but genetic drift, unlike
selection, is really noises in biological evolution.

While the WF models have led to speculations of pervasive natural selection in multi-copy gene systems
(Dover 1982; Arkhipova 2018; Chen, Yang, et al. 2022; Pan, Zhang, et al. 2022; Wang, et al. 2022), the GH
model may reverse the view. For rDNAs, the GH model will show that neutrality is often a simpler
explanation. If and when the model rejects the neutrality explanation, the rejection would also lead to the

more convincing proof of selection,

Results

PART I presents a best-known multi-gene system of the rRNA genes. PART II (Theory) consolidates aspects

of the Haldane model applied to polymorphism within species as well as divergence between species. In

PART III (Data analyses), we apply the theory to rDNA evolution in mice and apes (human and

chimpanzee).

PART I - The biology of rRNA gene clusters

The ribosomal RNA genes (rDNAs) are multi-copy gene clusters (Bowman, et al. 2020) that are arrayed as

tandem repeats on multiple chromosomes as shown in Fig. 1A (Guillén, et al. 2004; Cazaux, et al. 2011). In

humans, the copy number can vary from 60 to 1600 per individual (mean, 315; SD, 104; median, 301)

(Parks, et al. 2018). For each haploid genome, $C \sim 150$ on average in humans and $C \sim 110$ in mice (Parks, et

al. 2018). In humans, the five rRNA clusters are located on the short arm of the five acrocentric

chromosomes (Smirnov, et al. 2021). Such an arrangement permits crossovers between chromosomes

without perturbing the rest of the genomes. In *Mus*, the rDNAs are all located in the pericentromeric or sub-

telomeric region, on the long arms of telocentric chromosomes (Cazaux, et al. 2011; Potapova and Gerton

2019). Thus, unequal crossovers between non-homologous chromosomes may involve centromeres while

other genic regions are also minimally perturbed.

Fig. 1A-D here

Each copy of rRNA gene has a functional and non-functional part as shown in Fig. 1B. The "functional"

regions of rDNA, 18S, 5.8S, and 28S rDNA, are believed to be under strong negative selection, resulting in

a slow evolution rate in animals (Salim and Gerton 2019). In contrast, the transcribed spacer (ETS and ITS)

and the intergenic spacer (IGS) show significant sequence variation even among closely related species

(Eickbush and Eickbush 2007). Clearly, these "non-functional" sequences are less constrained by negative

selection. In this study of genetic drift, we focus on the non-functional parts. Data on the evolution of the

functional parts will be provided only for comparisons.

The pseudo-population of ribosomal DNA copies within each individual

While a human haploid with 200 rRNA genes may appear to have 200 loci, the concept of "gene loci"

cannot be applied to the rRNA gene clusters. This is because DNA sequences can spread from one copy to

others on the same chromosome via replication slippage. They can also spread among copies on different

chromosomes via gene conversion and unequal crossovers (Nagylaki 1983; Ohta and Dover 1983; Stults, et

al. 2008; Smirnov, et al. 2021). Replication slippage and unequal crossovers would also alter the copy

number of rRNA genes. These mechanisms will be referred to collectively as the homogenization process.

Copies of the cluster on the same chromosome are known to be nearly identical in sequences (Hori, et al.

2021; Nurk, et al. 2022). Previous research has also provided extensive evidence for genetic exchanges

between chromosomes (Krystal, et al. 1981; Arnheim, et al. 1982; van Sluis, et al. 2019).

In short, rRNA gene copies in an individual can be treated as a pseudo-population of gene copies. Such a

pseudo-population is not Mendelian, but its genetic drift can be analyzed using the branching process (see

below). The pseudo-population corresponds to the "chromosome community" proposed recently

(Guarracino, et al. 2023). As seen in Fig. 1C, the five short arms harbor a shared pool of rRNA genes that

can be exchanged among them. Fig. 1D presents the possible molecular mechanisms of genetic drift within

individuals whereby mutations may spread, segregate or disappear among copies. Hence, rRNA gene diversity or polymorphism refers to the variation across all rRNA copies, as these genes exist as paralogs rather than orthologs. This diversity can be assessed at both individual and population levels according to the multi-copy nature of rRNA genes.

PART II - Theory

1. The Haldane model of genetics drift applied to multi-copy gene systems

The Haldane model of genetic drift based on the branching process is intuitively appealing. In the model, each copy of the gene leaves K copies in a time interval with the mean and variance of $E(K)$ and $V(K)$. If $V(K) = 0$, there is no gene frequency change and no genetic drift. In the standard WF model, $V(K) = E(K)$ whereas $V(K)$ is decoupled from $E(K)$ in the Haldane model; the latter thus being more flexible. [In the companion paper, we discuss the modified WF models with $V(K) \neq E(K)$ which, nevertheless, do not resolve the paradoxes generated from population sampling.]

Below, we compare the strength of genetic drift in rRNA genes vs. that of single-copy genes using the Haldane model (Ruan, et al. 2024). We shall use * to designate the equivalent symbols for rRNA genes; for example, $E(K)$ vs. $E^*(K)$. Both are set to 1, such that the total number of copies in the long run remains constant.

For simplicity, we let $V(K) = 1$ for single-copy genes. (If we permit $V(K) \neq 1$, the analyses will involve the ratio of $V^*(K)$ and $V(K)$ to reach the same conclusion but with unnecessary complexities.) For rRNA genes, $V^*(K) \geq 1$ may generally be true because K for rDNA mutations are affected by a host of homogenization factors including replication slippage, unequal cross-over, gene conversion and other related mechanisms not operating on single-copy genes. Hence,

$$N^* = \frac{NC}{V^*(K)} = N\left[\frac{C}{V^*(K)}\right] = NC^* \qquad Eq.\,(1)$$

where N^* and N are the effective population size of rDNAs and single-copy genes, C is the average number of rRNA genes in an individual, and $V^*(K)$ reflects the homogenization process on rRNA genes (Fig. 1D). Thus,

$$C^* = C/V^*(K)$$

represents the effective copy number of rRNA genes in the population, determining the level of genetic diversity relative to single-copy genes. Since C is in the hundreds and $V^*(K)$ is expected to be > 1, the relationship of $1 << C^* \leq C$ is hypothesized. Fig. 1D is a simple illustration that the homogenizing process may enhance $V^*(K)$ substantially over the WF model.

In short, genetic drift of rRNA genes would be equivalent to single-copy genes in a population of size NC^* (or N^*). Since $C^* >> 1$ is hypothesized, genetic drift for rRNA genes is expected to be slower than for single-copy genes.

2. rDNA polymorphism within species

A standard measure of genetic drift is the level of heterozygosity (H). At the mutation-selection equilibrium

$$H_{equi} = \frac{2N_e\mu}{2N_e\mu + 1}$$

where μ is the mutation rate of the entire gene and N_e is the effective population size. In this study, $N_e = N$ for single-copy gene and $N_e = C^*N$ for rRNA genes. The empirical measure of nucleotide diversity H is given by

$$H = \frac{\sum_{i=1}^{L} 2p_i(1-p_i)}{L} \qquad Eq.\,(2)$$

where L is the gene length (for each copy of rRNA gene, $L \sim 43kb$) and p_i is the variant frequency at the i-th site.

We calculate H of rRNA genes at three levels – within-individual, within-species and then, within total samples (H_I, H_S and H_T, respectively). H_S and H_T are standard population genetic measures (Crow and Kimura 1970; Hartl, et al. 1997). In calculating H_S, all sequences in the species are used, regardless of the source individuals. A similar procedure is applied to H_T. The H_I statistic is adopted for multi-copy gene systems for measuring within-individual polymorphism. Note that copies within each individual are treated as a pseudo-population (see Fig. 1 and text above). With multiple individuals, H_I is averaged over them.

Given the three levels of heterozygosity, there are two levels of differentiation. First, F_{IS} is the differentiation among individuals within the species, defined by

$$F_{IS} = [H_S - H_I]/H_S$$

F_{IS} is hence the proportion of genetic diversity in the species that is found only between individuals. We will later show $F_{IS} \sim 0.05$ in human rDNA (Table 2), meaning 95% of rDNA diversity is found within individuals.

Second, F_{ST} is the differentiation between species within the total species complex, defined as

$$F_{ST} = [H_T - H_S]/H_T$$

F_{ST} is the proportion of genetic diversity in the total data that is found only between species. Between mouse species, F_{ST} distribution is close to 1 (Fig. 4C), indicating a large genetic distance between species relative to within-species polymorphisms.

3. rDNA divergence between species

Whereas the level of genetic diversity is a function of the effective population size, the rate of divergence between species, in its basic form, is not. The rate of neutral molecular evolution (λ), although driven by mutation and genetic drift, is generally shown by Eq. (3) (Crow and Kimura 1970; Hartl, et al. 1997; Li 1997):

$$\lambda = N\mu \frac{1}{N} = \mu \qquad Eq.\,(3)$$

Note that the factor of $1/N$ in Eq. (3) indicates the fixation probability of a new mutation. For rDNA mutations, fixation must occur in two stages – fixation within individuals and then among individuals in the

population. (We note again that new mutations can be fixed via homogenization in an individual, effectively forming a pseudo-population for rRNA genes.) Due to the cancelation of N in Eq. (3), the evolutionary rate of rRNA genes in the long run should be the same as single-copy genes.

Fig. 2 here

Eq. (3) is valid in the long-term evolution. For shorter-term evolution, it is necessary to factor in the fixation time (Fig. 2), T_f, which is the time between the emergence of a mutation and its fixation. If we study two species with a divergent time (T_d) equal to or smaller than T_f, then few mutations of recent emergence would have been fixed as species diverge.

Note that T_d is about 6 million years (Myrs) between human and chimpanzee while T_f (as measured by coalescence) is 0.6 - 1 Myrs in humans. Mutations of single-copy genes would not get fixed during the more recent T_f, as indicated in Figure 2. Thus, the realized substitution rate may be 1/6 to 1/10 lower than the theoretical value. In comparison, T_f^* of rRNA mutations should be at least 4.8 - 8 Myrs based on the C^* estimate (see PATR III). Thus, the substitution rate could be at least 80% lower than calculated for single-copy genes.

Our own theoretical derivation has shown that the fixation time for rRNA genes is close to $4N^*$ as is the case for single-copy genes at $\sim 4N$. If $V^*(K)$ is sufficiently larger than $V(K)$, it is in fact possible for $N^* < N$ such that genetic drift is stronger for rRNA genes than for single-copy genes and $T_f^* < T_f$. Therefore, T_{f1} can represent the T_f^* of mutations in rRNA genes in Fig. 2. This is interesting because, if the homogenization is powerful enough, rRNA genes would have an effective copy number of $C^* < 1$. A short T_f^*, which can be obtained by large $V^*(K)$, would lead to a smaller T_f^*/T_d than T_f/T_d, and thus a higher substitution rate than single-copy genes, particularly in short-term evolution. However, even T_f^* approaches 0, the substitution rate can exceed that of single-copy genes but is still limited by fixation probability. As noted above, the rapidly fixed mutations may not be well represented in the polymorphic data but can accumulate in species divergence.

PART III - Data Analyses

Before presenting the main analyses, we provide some empirical observations on the rapid homogenization of rRNA genes within individuals (Stage I). These observations are needed for PART III that comprises i) the analyses of rDNA polymorphisms within species in mouse and human; and ii) the analysis of the divergence between species.

Empirical measurements of homogenization within cells

In an accompanying study (Wang, et al. unpublished data), the evolutionary rate of neutral rRNA variants within cells is measured. Here, genetic drift operates via the homogenizing mechanisms that include gene conversion, unequal crossover and replication slippage. In the literature, measurements of neutral rRNA evolution are usually based on comparisons among individuals. Therefore, the Mendelian mechanisms of chromosome segregation and assortment would also shuffle variants among individuals. Segregation and assortment would confound the measurements of homogenization effects within individuals.

In one experiment, the homogenization effects in rDNAs are measured in cultured cell lines over 6 months of evolution. For in vivo homogenization, we analyze the evolution of rRNA genes within solid tumors. We estimate the rate at which rRNA variants spread among copies within the same cells, which undergo an asexual

process. The measurements suggest that, in the absence of recombination and chromosome assortment, the fixation time of new rRNA mutations within cells would take only 1 - 3 kyrs (thousand years). Since a new mutation in single-copy genes would take 300 - 600 kyrs to be fixed in human populations, the speed of genetic drift in Stage I evolution is orders faster than in Stage II. Therefore, despite having several hundred copies of rRNA genes per genome, the speed of genetic drift in rRNA genes may not be that much slower than in single-copy genes. This postulate will be tested below.

1. rDNA polymorphism within species

1) Polymorphism in mice

For rRNA genes, H_I of 10 individuals ranges from 0.0056 to 0.0067 while H_S is 0.0073 (Table 1). Thus, $F_{IS} = [H_S - H_I]/H_S$ for mice is 0.14, which means 86% of variation is within each individual. In other words, even one single randomly chosen individual would yield 85% of the diversity of the whole species. Hence, the estimated H_S should be robust as it is not affected much by the sampling.

H_S for *M. m. domesticus* single-copy genes is roughly 1.40 per kb genome-wide (Geraldes, et al. 2008) while H_S for rRNA genes is 7.25 per kb (Table 1), 5.2 times larger. In other words, $C^* = N^*/N \sim 5.2$. If we use the polymorphism data, it is as if rDNA array has a population size 5.2 times larger than single-copy genes. Although the actual copy number on each haploid is $\sim$ 110, these copies do not segregate like single-copy genes and we should not expect N^* to be 100 times larger than N. The H_S results confirm the prediction that rRNA genes should be more polymorphic than single-copy genes.

Based on the polymorphism result, one might infer slower drift for rDNAs than for single-copy genes. However, the results from the divergence data in later sections will reveal the under-estimation of drift strength from polymorphism data. Such data would miss variants that have a fast drift process driven by, for example, gene conversion. Strength of genetic drift should therefore be measured by the long-term fixation rate.

Tables 1- 2 and Fig. 3 here

2) Polymorphism in human

F_{IS} for rDNA among 8 human individuals is 0.059 (Table 2), much smaller than 0.142 in *M. m. domesticus* mice, indicating minimal genetic differences across human individuals and high level of genetic identity in rDNAs between homologous chromosomes among human population. Consisted with low F_{IS}, Fig. 3 shows strong correlation of the polymorphic site frequency of rDNA transcribed region among each pair of individuals from three continents (2 Asians, 2 Europeans and 2 Africans). Correlation of polymorphic sites in IGS region is shown in Supplementary Fig. 1. The results suggest that the genetic variation due to the sampling of chromosomes during sexual reproduction (e.g., segregation and assortment) is substantially attenuated by the effects of homogenization process within individual. Like those in mice, the pattern indicates that intra-species polymorphism is mainly preserved within individuals. The observed H_I of humans for rDNAs is 0.0064 to 0.0077 and the H_S is 0.0072 (Table 2). Research has shown that heterozygosity for the human genome is about 0.00088 (Zhao, et al. 2000), meaning the effective copy number of rDNAs is roughly, or C^* $\sim$ 8. This reduction in effective copy number from 150 to 8 indicates strong genetic drift due to homogenization

force.

2. rDNA divergence between species

We now consider the evolution of rRNA genes between species by analyzing the rate of fixation (or near
fixation) of mutations. Polymorphic variants are filtered out in the calculation. Note that Eq. (3) shows that
the mutation rate, μ, determines the long-term evolutionary rate, λ. Since we will compare the λ values
between rDNA and single-copy genes, we first need to compare their mutation rates by analyzing their long-
term divergence, which is hardly influenced by fixation time. As shown in Table S1, λ falls in the range of
50-60 (differences per Kb) for single-copy genes and $40 - 70$ for the non-functional parts of rRNA genes.
The data thus suggest that rRNA and single-copy genes are comparable in mutation rate. Differences
between their λ values will have to be explained by other means.

1) Between mouse species - Genetic drift as the sole driving force of the rapid divergence

We now use the F_{ST} statistic to delineate fixation and polymorphism. The polymorphism in *M. m. domesticus*
is compared with two outgroup species, *M. spretus* and *M. m. castaneus*, respectively. There are hence two
depths in the phylogeny with two T_d's, as shown in Fig. 4A (Rudra, et al. 2016; Kumar, et al. 2022). There is
a fourth species, *M. m. musculus* (shown in grey in Fig. 4A), which yields very similar results as *M. m.*
domesticus in these two comparisons. These additional analyses are shown in Supplement Table S2- S3.

The F_{IS} values of polymorphic sites in 3 outbred mice are primarily below 0.2 and rarely above 0.8 in Fig. 4B,
indicating the low genetic differentiation in rDNAs within these 3 *M. m. domesticus*. While the F_{IS} distribution
of 10 mice from Table 1, including 7 inbred and 3 outbred mouse strains, exhibits a noticeable right skewness,
but does not exceed 0.8. This suggests that inbreeding to a certain extent limits the process of homogenization
and enhances population differentiation. In comparison, the distribution of the F_{ST} of variant sites between *M.*
m. domesticus and *Mus spretus* has a large peak near $F_{ST} = 1$. This peak in Fig. 4C represents species
divergence not seen within populations (i.e., F_{IS}). We use $F_{ST} = 0.8$ as a cutoff for divergence sites between
the two species. Roughly, when a mutant is > 0.95 in frequency in one species and < 0.05 in the other, F_{ST}
would be > 0.80.

We first compare the divergence between *M. m. domesticus* and *M. m. castaneus* whereby T_d has been
estimated to be less than 0.5 Myrs (Fujiwara, et al. 2022). In comparison, between *Mus m. domesticus* and
Mus spretus, T_d is close to 3 Myrs (Rudra, et al. 2016). As noted above, the reduction in the divergence rate
relative to that of Eq. (3) is proportional to T_f/T_d (for single copy genes) or T_f^*/T_d (for rRNA genes). As T_f and
T_f^* are both from *M. m. domesticus* and T_d is 6 times larger in comparison with *M. spretus*, we expect the
results to be quite different between the two sets of species comparisons.

Although T_f and T_f^* estimates are less reliable than T_d estimates, both comparisons use the same T_f and T_f^*
from *M. m. domesticus*. Hence, the results should be qualitatively unbiased. For a demonstration, we shall use
the estimates of T_f (i.e., the coalescence time) at 0.2 Myrs for single-copy genes by using an average nucleotide
diversity of 0.0014 and the mutation rate of 5.7×10^{-9} per base pair per generation (Geraldes, et al. 2008; Phifer-
Rixey, et al. 2020). Based on the estimated C^* above, we obtain T_f^* for rDNA mutations at 5×0.2 Myrs, or 1
Myrs. While some have estimated T_f to be close to 0.4 Myrs (Fujiwara, et al. 2022), we aim to show that the
pattern of reduction in rRNA divergence is true even with the conservative estimates.

Between *M. m. domesticus* and *M. m. castaneus*, the reduction in substitution rate for single copy gene should

be ~ 40% (T_f/T_d = 0.2/0.5), and the reduction for rRNA genes should be 100% (T_f^*/T_d = 1/0.5 > 1). Table 3 on

their DNA sequence divergence shows that rRNA genes are indeed far less divergent than single-copy genes.

In fact, only a small fraction of rDNA mutations is expected to be fixed as T_f^* for rDNA at 1 Myrs is 2 times

larger than the divergence time, T_d. We should note again that the non-negligible fixation of rRNA mutations

suggests that C^* at 5 is perhaps an over-estimate.

Tables 3, 4, 5 and Fig. 4 here

Between *Mus m. domesticus* and *Mus spretus*, the reduction in actual substitution rate from theoretical limit

for single-copy genes should be 6.7% (T_f/T_d = 0.2/3) and, for rRNA genes, should be 33% (T_f^*/T_d = 1/3). The

evolutionary rate (i.e. the fixation rate) of IGS region is lower than single-copy genes, 0.01 in IGS and 0.021

in genome-wide (Table 4), as one would expect. However, ETS and ITS regions have evolved at a surprising

rate that is 12% higher than single-copy genes. Note that the reduction in C^*, even to the lowest limit of C^*

=1, would only elevate the rate of fixation in rRNA genes to a parity with single-copy genes. From Eq. (1),

the explanation would be that $V^*(K)$ has to be very large, such that C^* is < 1. With such rapid homogenization,

the fixation time approaches 0 and the substitution rate in rRNA genes can indeed reach the theoretical limit

of Eq. (3). In such a scenario, the substitution rate in ETS and ITS, compared to single-copy genes in mice,

may increase by 7%, $T_f/(T_d - T_f)$ = 0.2/(3-0.2). If we use T_f ~ 0.4 Myrs in an alternative estimation, the increase

can be up to 15%.

In conclusion, the high rate of fixation in ETS and ITS may be due to very frequent gene conversions that

reduce C^* to be less than 1. In contrast, IGS may have undergone fewer gene conversions and its long-term

C^* is slightly larger than 1. Indeed, the heterozygosity in IGS region, at about 2-fold higher than that of ETS

and ITS regions (8‰ for IGS, 5‰ for ETS and 3‰ for ITS), supports this interpretation.

2) Between Human and Chimpanzee - Positive selection in addition to rapid drift in rDNA divergence

Like the data of mouse studies, the polymorphism of rDNAs in humans would suggest a slower short-term

evolution rate. The same caveat is that C^* estimated from the polymorphism data would have missed those

rapidly fixed variants. Hence, the long-term C^* obtained from species divergence might be much smaller than

8.

Our results show that the evolutionary rate of rRNA genes between human and chimpanzee is substantially

higher than that of other single-copy genes (Table 5). Especially, 5'ETS region shows a 100% rate

acceleration, at 22.7‰ vs. 11‰ genome-wide. Even after removing CpG sites, their fixation rate still

reaches 22.4‰. In this case, even if C^* <<1, the extremely rapid fixation will only increase the substitution

rate by $T_f/(T_d - T_f)$ by 11%, compared to single-copy genes. Thus, the much accelerated evolution of rRNA

genes between humans and chimpanzees cannot be entirely attributed to genetic drift. In the next and last

section, we will test if selection is operating on rRNA genes by examining the pattern of gene conversion.

3) Positive selection for rRNA mutations in apes, but not in mice – Evidence from gene conversion patterns

For gene conversion, we examine the patterns of AT-to-GC vs. GC-to-AT changes. While it has been

reported that gene conversion would favor AT-to-GC over GC-to-AT conversion (Jeffreys and Neumann

2002; Meunier and Duret 2004) at the site level, we are interested at the gene level by summing up all

conversions across sites. We designate the proportion of AT-to-GC conversion as *f* and the reciprocal, GC-to-

AT, as *g*. Both *f* and *g* represent the proportion of fixed mutations between species (see Methods). So

defined, *f* and *g* are influenced by the molecular mechanisms as well as natural selection. The latter may

favor a higher or lower GC ratio at the genic level between species. As the selective pressure is distributed

over the length of the gene, each site may experience rather weak pressure.

Let p be the proportion of AT sites and q be the proportion of GC sites in the gene. The flux of AT-to-GC
would be pf and the flux in reverse, GC-to-AT, would be qg. At equilibrium, $pf = qg$. Given f and g, the ratio
of p and q would eventually reach $p/q = g/f$. We now determine if the fluxes are in equilibrium ($pf = qg$). If
they are not, the genic GC ratio is likely under selection and is moving to a different equilibrium.

In these genic analyses, we first analyzed the human lineage (Brown and Jiricny 1989; Galtier and Duret
2007). Using chimpanzees and gorillas as the outgroups, we identified the derived variants that became
nearly fixed in humans with frequency > 0.8 (Table 6). The chi-square test shows that the GC variants had a
significantly higher fixation probability compared to AT. In addition, this pattern is also found in
chimpanzees (p < 0.001). In *M. m. domesticus* (Table 6), the chi-square test reveals no difference in the
fixation probability between GC and AT (p = 0.957). Further details can be found in Supplementary Figure
2. Overall, a higher fixation probability of the GC variants is found in human and chimpanzee, whereas this
bias is not observed in mice.

364   **Tables 6-7 here**

Based on Table 6, we could calculate the value of p, q, f and g (see Table 7). Shown in the last row of Table
7, the (pf)/(qg) ratio is much larger than 1 in both the human and chimpanzee lineages. Notably, the ratio in
mouse is not significantly different from 1. Combining Tables 4 and 7, we conclude that the slight
acceleration of fixation in mice can be accounted for by genetic drift, due to gene conversion among rRNA
gene copies. In contrast, the different fluxes corroborate the interpretations of Table 5 that selection is
operating in both humans and chimpanzees.

Discussion

The Haldane model is an "individual-output" model of genetic drift (Chen, et al. 2017). Hence, it does not
demand the population to follow the rules of WF populations. Indeed, it would be difficult to define WF
populations in any multi-copy gene systems. The GH model is also sufficiently flexible for studying various
stochastic forces other than the sampling errors that together drive genetic drift. In the companion study
(Ruan, et al. 2024), we address the ecological forces of genetic drift and, in this study, we analyze the neutral
evolution of rRNA genes. Both examples are amenable to the analysis by the Haldane model, but not by the
WF model.

In multi-copy systems, there are several mechanisms of homogenization within individuals. For rRNA genes,
whether on the same or different chromosomes (Gonzalez and Sylvester 2001; van Sluis and McStay 2019),
the predominant mechanism of homogenization mechanism are gene conversion and unequal crossover. In the
process of exchanging DNA sections in meiosis, gene conversions are an order of magnitude more common
than crossover (Cole, et al. 2012; Williams, et al. 2015).

There have been many rigorous analyses that confront the homogenizing mechanisms directly. These studies
(Smith 1974; Ohta 1976; Dover 1982; Nagylaki 1983; Ohta and Dover 1983) modeled gene conversion and
unequal cross-over head on. Unfortunately, on top of the complexities of such models, the key parameter
values are rarely obtainable. In the branching process, all these complexities are wrapped into $V^*(K)$ for
formulating the evolutionary rate. In such a formulation, the *collective* strength of these various forces may

indeed be measurable, as shown in this study.

By compressing all stochastic elements into $V(K)$, or $V^*(K)$, we will be able to filter out random factors and
focus on forces of greater biological interest such as selection. Interestingly, while rDNA evolution in mice
can be fully accounted for by genetic drift alone, its evolution in the great apes must have been driven by
selection along with genetic drift. The contrast shows that the GH model only tries to delineate genetic drift
more accurately than has been done by the WF model.

The branching process is a model for general processes. Hence, it can be used to track genetic drift in systems
with two stages of evolution - within- and between-individuals, even though TEs, viruses or rRNA genes are
very different biological entities. We use the rRNA genes to convey this point. Multi-copy genes, like rDNA,
are under rapid genetic drift via the homogenization process. The drift is strong enough to reduce the copy
number in the population from $\sim 150N$ to $< N$. A fraction of mutations in multi-copy genes may have been
fixed by drift almost instantaneously in the evolutionary time scale. This acceleration is seen in mice but would
have been interpreted to be due to positive selection by the convention. Interestingly, while positive selection
may not be necessary to explain the mice data, it is indeed evident in human and chimpanzee, as the
evolutionary rate of rRNA genes exceeds the limit of the strong drift.

In multi-copy systems, the copy number (designated C) is often in the hundreds or thousands per individual.
Nevertheless, $C = 2$ as in all diploids is also a multi-copy gene system as the two copies may often evolve
interactively via gene conversion or segregation distortion (Wu, et al. 1988; McDermott and Noor 2010). The
WF models have been noted to be inadequate even in diploid systems (Charlesworth 2009; Chen, et al. 2017).
After all, the WF model is essentially a haloid model with $2N$ copies in the population.

In conclusion, the GH model is far more general than the WF model as the two companion studies clearly
demonstrate. Its $E(K)$ parameter would define N's while $V(K)$ would track random changes in gene
frequency. It can quantify genetic drift in multi-gene systems including viruses such as SARS-CoV-2. The
chaotic analyses of COVID-19 have shown why an accurate account of genetic drift is indispensable (Deng,
et al. 2022; Pan, Liu, et al. 2022; Ruan, Wen, et al. 2022; Hou, et al. 2023).

Materials and Methods

Data Collection.

We collected high-coverage whole-genome sequencing data for our study. The genome sequences of human
(n = 8), chimpanzee (n = 1) and gorilla (n = 1) were sourced from National Center for Biotechnology
Information (NCBI) (Supplementary Table 3). Human individuals were drawn from diverse geographical
origins encompassing three continents (4 Asians, 2 Europeans and 2 Africans) and Asia CRC was the normal
tissue of Case 1 patient from (Chen, Wu, et al. 2022). Genomic sequences of mice (n = 13) were sourced
from the Wellcome Sanger Institute's Mouse Genome Project (MGP) (Keane, et al. 2011). Although some
artificial selection has been performed on laboratory mouse strains (Yang, et al. 2011), the WSB/EiJ,
ZALENDE/Ei and LEWES/EiJ strains were derived from wild populations. Incorporating these wild-
derived laboratory strains, along with other inbred strains, a cohort of 10 mice was utilized to approximate
the population representation of *M. m. domesticus*. Furthermore, the low F_{IS} of 0.14 for rDNA in *M. m.*

domesticus found in this study suggests that each mouse covers 86% of the population's genetic diversity,
thereby mitigating concerns about potential sampling biases. Accessions and the detailed information of
samples used in this study are listed in the Supplementary Table 3 and Table 4.

Variant allele frequency.

Following adapter trimming and the removal of low-quality sequences, these whole-genome sequencing
data of apes and mice were mapped against respective reference sequences: the human rDNA reference
sequence (Human ribosomal DNA complete repeating unit, GenBank: U13369.1) and the mouse rDNA
reference sequence (*Mus musculus* ribosomal DNA, complete repeating unit, GenBank: BK000964.3).
Alignment was performed using Burrows-Wheeler-Alignment Tool v0.7.17 (Li and Durbin 2009) with
default parameters. All mapping and analysis are performed among individual copies of rRNA genes. The
pipelines in variant calling are similar to the ones used in the literature (Ma, et al. 2022; Sun, et al. 2022).
Each individual was considered as a psedo-population of rRNA genes and the diversity of rRNA genes was
calculated using this psedo-population of rRNA genes. To determine variant frequency within individual,
variants were called from multiple alignment files using bcftools (Danecek, et al. 2021) to ensure the
inclusion of all polymorphic sites that appeared in at least one sample and to maintain consistent processing
steps. Per-sample variant calling results were generated in a VCF file using the following settings: 'bcftools
mpileup --redo-BAQ --max-depth 50000 --per-sample-mF --annotate' and 'bcftools call -mv'. Our analysis
specifically focused on single nucleotide variants (SNVs) with only two alleles, then calculated the
frequency of major allele, which is less susceptible to technical errors, to estimate the nucleotide diversity
using Eq. (2), Variant information of interest per sample was extracted from the VCF files using the
command 'bcftools view -V indels' and 'bcftools query –f
'%CHROM\t%POS\t%REF\t%ALT\t[\t%GT]\t[\t%DP]\t[\t%AD]\n'\'. The AD (total high-quality bases of
allelic depths) was used to obtain the number of reference-supporting reads (n.ref) and alternative-
supporting reads (n.alt) in each sample. Sites with a depth < 10 in any sample were filtered out, resulting in
an average of the minimum depths above 3000 for each remaining site across all samples. This filtering step
enhances the robustness of variant frequency estimation, where the variant frequency was approximated by
the ratio n.alt/(n.ref + n.alt). This process allowed us to identify all polymorphic sites present in the samples
and the variant frequency within each individual. Then the population-level variant frequency was computed
by averaging variant frequencies across all individuals.

Identification of Divergence Sites.

Polymorphism represents the transient phase preceding divergence. With variant frequency within
individual, within species, and between species, we obtained the F_{IS} and F_{ST} values for each site. Within the
M. m. domesticus population, 779 polymorphic sites were identified. Among these, 744 sites (95.5%)
exhibited an F_{IS} below 0.4 and rarely above 0.8 in Fig. 4B. In the comparison between *M. m. domesticus* and
Mus spretus, 1579 variant sites were found, among which 453 sites displayed an F_{ST} above 0.8, indicating
swift fixation of mutations during species divergence.

F_{ST} analysis between human and chimpanzee was conducted for each human individual, summarized in
Table 5. We identified a range from 672 to 705 sites with F_{ST} values above 0.8 across individuals, depicting
robust divergence sites. Considering the high mutation rate in CpG sites (Ehrlich and Wang 1981; Sved and
Bird 1990) and predominantly GC content in rDNA (*e.g.* 58% GC in human), we further estimated the
evolutionary rate at non-CpG sites during the interspecies divergence. To achieve this, mutations in CpG
sites were manually removed by excluding all sites containing CpG in one species and TpG or CpA in the
other; the reverse was similarly discarded. Additionally, the count of non-CpG sites within the mapping

length, where site depth exceeded 10, was performed by Samtools (Danecek, et al. 2021) with the settings 'samtools mpileup -Q 15 -q 20'. As a result, the evolutionary rate of rDNA in non-CpG sites was ascertained.

For assessing diversity and divergence across gene segments, we used 'samtools faidx' to partition variants into a total of 8 regions within rRNA genes, including 5'ETS, ITS1, ITS2, 3'ETS, IGS, 18S, 5.8S, and 28S, aligning them with corresponding reference sequences for further analysis. The functional parts (18S, 5.8S, and 28S) were subject to strong negative selection, exhibiting minimal substitutions during species divergence as expected. This observation is primarily used for comparison with the non-functional parts and reflects that the non-functional parts are less constrained by negative selection.

Genome-wide Divergence Estimation.

To assess the genome-wide divergence between 4 mouse strain species, we downloaded their toplevel reference genomes from Ensembl genome browser (GenBank Assembly ID: GCA_001624865.1, GCA_001624775.1, GCA_001624445.1, and GCA_001624835.1). Then we used Mash tools (Ondov, et al. 2016) to estimate divergence across the entire genome (mainly single-copy genes) with 'mash sketch' and 'mash dist'. Additionally, the effective copy number of rRNA genes, denoted as C^*, can be estimated by calculating the ratio of population diversity observed in rDNA to that observed in single-copy genes.

Estimation of Site Conversion.

To estimate site conversions, whether accidental or directional, it was essential to identify new mutant alleles in each lineage after divergence. New mutations were defined as derived alleles that differed from ancestral alleles shared by two outgroup species, where the ancestral state also exhibited high identity among copies. By focusing on new mutations with low initial frequency, we minimized the influence of their initial frequency on fixation probability and fixation time. Specifically, variants shared between chimpanzee and gorilla, humans and gorilla, *M. m. castaneus* and *M. spretus*, each with a frequency greater than 0.8, were considered as the ancestral for human, chimpanzee, and *M. m. domesticus*, respectively.

Derived variants were then categorized into two groups: (nearly) fixed (frequency > 0.8) and not fixed mutations in the lineages of humans, chimpanzees and *M. m. domesticus* (Table 6). The frequency threshold of >0.8 was chosen to balance the need for a sufficient number of sites to calculate of (f/g), and to ensure their reliability. We also applied a more stringent threshold of >0.9, which yielded similar results.

In this study, six types of mutations were tabulated, representing ancestral-to-derived as depicted in Supplementary Fig. 2. For example, A-to-G represented the both A-to-G and T-to-C types of mutations. The C-to-G (or G-to-C) and A-to-T (or T-to-A) types of mutations were excluded in the subsequent analysis.

Specifically, f represents the proportion of fixed mutations where an A or T nucleotide has been converted to a G or C nucleotide. The numerator for f is the number of fixed mutations from A-to-G, T-to-C, T-to-G, or A-to-C. .Since the fixed sites accounted for less than 1% of the non-functional length of rDNA, the denominator is the total number of A or T sites in the rDNA sequence of the species lineage.

Similarly, g is defined as the proportion of fixed mutations where a G or C nucleotide has been converted to an A or T nucleotide. The numerator for g is the number of fixed mutations from G-to-A, C-to-T, C-to-A, or G-to-T. The denominator is the total number of G or C sites in the rDNA sequence of the species lineage.

The consensus rDNA sequences for the species lineage were generated by Samtools consensus (Danecek, et al. 2021) from the bam file after alignment. The following command was used:

'samtools consensus -@ 20 -a -d 10 --show-ins no --show-del yes input_sorted.bam output.fa'.

The alternative hypotheses of GC-biased mutation process (Wolfe, et al. 1989; Francino and Ochman 1999) alone can be rejected in this study. According to the prediction of the mutational mechanism hypotheses, AT

or GC variants should have equal fixation probabilities. We quantified the nearly fixed number of AT-to-GC and GC-to-AT types of mutations and conducted a chi-square test to assess their fixation probabilities. Notably, we found the G (or C) variant had a significant higher fixation probability than A (or T) at site level in apes, but not in mice. In order to test whether there is an equilibrium state at the genic level in these three species, we computed the $(pf)/(qg)$ ratio in Table 7, and a significant deviation of the ratio from 1 would imply biased genic conversion.

Data Availability

No new data were generated in this study. The genomic data used in this study are available from National Center for Biotechnology Information (NCBI) (https://www.ncbi.nlm.nih.gov/) and the Mouse Genomes Project (https://www.sanger.ac.uk/data/mouse-genomes-project/). The specific accession numbers recorded in Supplementary Table S4 and S5.

Acknowledgments

We are grateful for the helpful comments from many colleagues on the Chat Group "Cancer - The new evolving species", in particular, Weiwei Zhai, Yong Zhang, GuoDong Wang of CAS and Jianrong Yang of SYSU. This work was supported by the National Natural Science Foundation of China (32150006, 32293193/32293190 to C.I.W., 32200493 to Y.R., and 82341092 to HJ Wen.), the National Key Research and Development Projects of the Ministry of Science and Technology of China (2021YFC2301300, 2021YFC0863400), and Guangdong Key Research and Development Program (No. 2022B1111030001).

References

Alexandrov I, Kazakov A, Tumeneva I, Shepelev V, Yurov Y. 2001. Alpha-satellite DNA of primates: old and new families. Chromosoma 110:253-266.

Arkhipova IR. 2018. Neutral Theory, Transposable Elements, and Eukaryotic Genome Evolution. Molecular Biology and Evolution 35:1332-1337.

Arnheim N, Treco D, Taylor B, Eicher EM. 1982. Distribution of ribosomal gene length variants among mouse chromosomes. Proc Natl Acad Sci U S A 79:4677-4680.

Bowman JC, Petrov AS, Frenkel-Pinter M, Penev PI, Williams LD. 2020. Root of the Tree: The Significance, Evolution, and Origins of the Ribosome. Chemical Reviews 120:4848-4878.

Brown T, Jiricny J. 1989. Repair of base-base mismatches in simian and human cells. Genome / National Research Council Canada = Génome / Conseil national de recherches Canada 31:578-583.

Cabot EL, Doshi P, Wu ML, Wu CI. 1993. Population genetics of tandem repeats in centromeric heterochromatin: unequal crossing over and chromosomal divergence at the Responder locus of Drosophila melanogaster. Genetics 135:477-487.

Cannings C. 1974. The latent roots of certain Markov chains arising in genetics: A new approach, I. Haploid models. Advances in Applied Probability 6:260-290.

Cazaux B, Catalan J, Veyrunes F, Douzery EJP, Britton-Davidian J. 2011. Are ribosomal DNA clusters rearrangement hotspots? A case study in the genus Mus (Rodentia, Muridae). BMC Evolutionary

559   Biology 11:124.

560 Charlesworth B. 2009. Effective population size and patterns of molecular evolution and variation. Nature

561   Reviews Genetics 10:195-205.

562 Charlesworth B, Sniegowski P, Stephan W. 1994. The evolutionary dynamics of repetitive DNA in

563   eukaryotes. Nature 371:215-220.

564 Chen BJ, Wu XR, Ruan YS, Zhang YL, Cai QC, Zapata L, Wu CI, Lan P, Wen HJ. 2022. Very large hidden

565   genetic diversity in one single tumor: evidence for tumors-in-tumor. National Science Review

566   9:nwac250.

567 Chen QP, Yang H, Feng X, Chen QJ, Shi SH, Wu CI, He ZW. 2022. Two decades of suspect evidence for

568   adaptive molecular evolution-negative selection confounding positive-selection signals. National

569   Science Review 9:nwab217.

570 Chen Y, Tong D, Wu CI. 2017. A New Formulation of Random Genetic Drift and Its Application to the

571   Evolution of Cell Populations. Mol Biol Evol 34:2057-2064.

572 Chia AB, Watterson GA. 1969. Demographic effects on the rate of genetic evolution I. constant size

573   populations with two genotypes. Journal of Applied Probability 6:231-248.

574 Cole F, Kauppi L, Lange J, Roig I, Wang R, Keeney S, Jasin M. 2012. Homeostatic control of recombination

575   is implemented progressively in mouse meiosis. Nature Cell Biology 14:424-430.

576 Crow JF, Kimura M. 1970. An Introduction to Population Genetics Theory: Blackburn Press.

577 Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA,

578   Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. Gigascience 10.

579 Deng S, Xing K, He X. 2022. Mutation signatures inform the natural host of SARS-CoV-2. National Science

580   Review 9:nwab220.

581 Der R, Epstein CL, Plotkin JB. 2011. Generalized population models and the nature of genetic drift.

582   Theoretical Population Biology 80:80-99.

583 Dover G. 1982. Molecular drive: a cohesive mode of species evolution. Nature 299:111-117.

584 Ehrlich M, Wang RY. 1981. 5-Methylcytosine in eukaryotic DNA. Science 212:1350-1357.

585 Eickbush TH, Eickbush DG. 2007. Finely orchestrated movements: evolution of the ribosomal RNA genes.

586   Genetics 175:477-485.

587 Francino MP, Ochman H. 1999. Isochores result from mutation not selection. Nature 400:30-31.

588 Fujiwara K, Kawai Y, Takada T, Shiroishi T, Saitou N, Suzuki H, Osada N. 2022. Insights into Mus

589   musculus Population Structure across Eurasia Revealed by Whole-Genome Analysis. Genome Biol

590   Evol 14.

591 Galtier N, Duret L. 2007. Adaptation or biased gene conversion? Extending the null hypothesis of molecular

592   evolution. Trends in Genetics 23:273-277.

593 Geraldes A, Basset P, Gibson B, Smith KL, Harr B, Yu HT, Bulatova N, Ziv Y, Nachman MW. 2008.

594   Inferring the history of speciation in house mice from autosomal, X-linked, Y-linked and mitochondrial

595   genes. Mol Ecol 17:5349-5363.

596 Gibbons JG, Branco AT, Godinho SA, Yu S, Lemos B. 2015. Concerted copy number variation balances

597   ribosomal DNA dosage in human and mouse genomes.201416878.

598 Gonzalez IL, Sylvester JE. 2001. Human rDNA: Evolutionary Patterns within the Genes and Tandem Arrays

599   Derived from Multiple Chromosomes. Genomics 73:255-263.

600 Guan W-j, Zhong N-s. 2022. Strategies for reopening in the forthcoming COVID-19 era in China. National

601   Science Review 9:nwac054.

602 Guarracino A, Buonaiuto S, de Lima LG, Potapova T, Rhie A, Koren S, Rubinstein B, Fischer C, Abel HJ,

603   Antonacci-Fulton LL, et al. 2023. Recombination between heterologous human acrocentric

604   chromosomes. Nature 617:335-343.

605 Guillén AKZ, Hirai Y, Tanoue T, Hirai H. 2004. Transcriptional repression mechanisms of nucleolus

organizer regions (NORs) in humans and chimpanzees. Chromosome Research 12:225-237.

Haldane JBS. 1932. The Causes of Evolution. London: Longmans, Green and Company.

Haldane JBS. 1927. A mathematical theory of natural and artificial selection, part V: selection and mutation: Cambridge University Press.

Hartl DL, Clark AG, Clark AG. 1997. Principles of population genetics: Sinauer associates Sunderland.

Hori Y, Shimamoto A, Kobayashi T. 2021. The human ribosomal DNA array is composed of highly homogenized tandem clusters. Genome Res 31:1971-1982.

Hou M, Shi JR, Gong ZK, Wen HJ, Lan Y, Deng XZ, Fan QH, Li JJ, Jiang ML, Tang XP, et al. 2023. Intra- vs. Interhost Evolution of SARS-CoV-2 Driven by Uncorrelated Selection-The Evolution Thwarted. Molecular Biology and Evolution 40:msad204.

Huang J, Liu X, Zhang L, Zhao Y, Wang D, Gao J, Lian X, Liu C. 2021. The oscillation-outbreaks characteristic of the COVID-19 pandemic. National Science Review 8:nwab100.

Jeffreys AJ, Neumann R. 2002. Reciprocal crossover asymmetry and meiotic drive in a human recombination hot spot. Nat Genet 31:267-271.

Jurka J, Kapitonov VV, Kohany O, Jurka MV. 2007. Repetitive sequences in complex genomes: structure and evolution. Annu Rev Genomics Hum Genet 8:241-259.

Karlin S, McGregor J. 1964. Direct Product Branching Processes and Related Markov Chains. Proceedings of the National Academy of Sciences 51:598-602.

Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, Heger A, Agam A, Slater G, Goodson M, et al. 2011. Mouse genomic variation and its effect on phenotypes and gene regulation. Nature 477:289-294.

Krystal M, D'Eustachio P, Ruddle FH, Arnheim N. 1981. Human nucleolus organizers on nonhomologous chromosomes can share the same ribosomal gene variants. Proceedings of the National Academy of Sciences of the United States of America 78:5744-5748.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges SB. 2022. TimeTree 5: An Expanded Resource for Species Divergence Times. Mol Biol Evol 39.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754-1760.

Li W-H. 1997. Molecular evolution. Sunderland, Mass.: Sinauer Associates.

Ma Y, Mao X, Wang J, Zhang L, Jiang Y, Geng Y, Ma T, Cai L, Huang S, Hollingsworth P, et al. 2022. Pervasive hybridization during evolutionary radiation of Rhododendron subgenus Hymenanthes in mountains of southwest China. National Science Review 9:nwac276.

McDermott SR, Noor MAF. 2010. The role of meiotic drive in hybrid male sterility. Philosophical Transactions of the Royal Society B-Biological Sciences 365:1265-1272.

Meunier J, Duret L. 2004. Recombination drives the evolution of GC-content in the human genome. Molecular Biology and Evolution 21:984-990.

Nagylaki T. 1983. Evolution of a large population under gene conversion. Proc Natl Acad Sci U S A 80:5941-5945.

Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger MR, Altemose N, Uralsky L, Gershman A, et al. 2022. The complete sequence of a human genome. Science 376:44-53.

Ohta T. 1976. Simple model for treating evolution of multigene families. Nature 263:74-76.

Ohta T, Dover GA. 1984. The Cohesive Population Genetics of Molecular Drive. Genetics 108:501-521.

Ohta T, Dover GA. 1983. Population genetics of multigene families that are dispersed into two or more chromosomes. Proc Natl Acad Sci U S A 80:4079-4083.

Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM. 2016. Mash: fast genome and metagenome distance estimation using MinHash. Genome Biology 17:132.

Pan Y, Liu P, Wang F, Wu P, Cheng F, Jin X, Xu S. 2022. Lineage-specific positive selection on ACE2

contributes to the genetic susceptibility of COVID-19. National Science Review 9:nwac118.

Pan Y, Zhang C, Lu Y, Ning Z, Lu D, Gao Y, Zhao X, Yang Y, Guan Y, Mamatyusupu D, et al. 2022. Genomic diversity and post-admixture adaptation in the Uyghurs. National Science Review 9:nwab124.

Parks MM, Kurylo CM, Dass RA, Bojmar L, Lyden D, Vincent CT, Blanchard SC. 2018. Variant ribosomal RNA alleles are conserved and exhibit tissue-specific expression. Science Advances 4.

Phifer-Rixey M, Harr B, Hey J. 2020. Further resolution of the house mouse (Mus musculus) phylogeny by integration over isolation-with-migration histories. BMC Evol Biol 20:120.

Potapova T, Gerton J. 2019. Ribosomal DNA and the nucleolus in the context of genome organization. Chromosome Research 27.

Rand DM. 2001. The units of selection on mitochondrial DNA. Annual Review of Ecology and Systematics 32:415-448.

Ruan Y, Hou M, Tang X, He X, Lu X, Lu J, Wu CI, Wen H. 2022. The Runaway Evolution of SARS-CoV-2 Leading to the Highly Evolved Delta Strain. Mol Biol Evol 39.

Ruan Y, Wang X, Hou M, Diao W, Tracy ME, Xu S, Liufu Z, Wen H, Wu C-I. 2024. Resolving Paradoxes in Molecular Evolution: The Integrated WF-Haldane (WFH) Model of Genetic Drift. bioRxiv:2024.2002.2019.581083.

Ruan Y, Wen H, He X, Wu CI. 2021. A theoretical exploration of the origin and early evolution of a pandemic. Sci Bull (Beijing) 66:1022-1029.

Ruan Y, Wen H, Hou M, He Z, Lu X, Xue Y, He X, Zhang YP, Wu CI. 2022. The twin-beginnings of COVID-19 in Asia and Europe-one prevails quickly. Natl Sci Rev 9:nwab223.

Rudra M, Chatterjee B, Bahadur M. 2016. Phylogenetic relationship and time of divergence of Mus terricolor with reference to other Mus species. J Genet 95:399-409.

Salim D, Gerton JL. 2019. Ribosomal DNA instability and genome adaptability. Chromosome Research 27:73-87.

Smirnov E, Chmúrčiaková N, Liška F, Bažantová P, Cmarko D. 2021. Variability of Human rDNA. Cells 10.

Smith GP. 1974. Unequal Crossover and the Evolution of Multigene Families.   38:507-513.

Stults DM, Killen MW, Pierce HH, Pierce AJ. 2008. Genomic architecture and inheritance of human ribosomal RNA gene clusters. Genome Res 18:13-18.

Sun N, Yang L, Tian F, Zeng H, He Z, Zhao K, Wang C, Meng M, Feng C, Fang C, et al. 2022. Sympatric or micro-allopatric speciation in a glacial lake? Genomic islands support neither. National Science Review 9:nwac291.

Sved J, Bird AJPotNAoS. 1990. The expected equilibrium of the CpG dinucleotide in vertebrate genomes under a mutation model.   87:4692-4696.

Szitenberg A, Cha S, Opperman CH, Bird DM, Blaxter ML, Lunt DH. 2016. Genetic Drift, Not Life History or RNAi, Determine Long-Term Evolution of Transposable Elements. Genome Biology and Evolution 8:2964-2978.

Tatsumoto S, Go Y, Fukuta K, Noguchi H, Hayakawa T, Tomonaga M, Hirai H, Matsuzawa T, Agata K, Fujiyama A. 2017. Direct estimation of de novo mutation rates in a chimpanzee parent-offspring trio by ultra-deep whole genome sequencing. Sci Rep 7:13561.

van Sluis M, Gailín M, McCarter JGW, Mangan H, Grob A, McStay B. 2019. Human NORs, comprising rDNA arrays and functionally conserved distal elements, are located within dynamic chromosomal regions. Genes Dev 33:1688-1701.

van Sluis M, McStay B. 2019. Nucleolar DNA Double-Strand Break Responses Underpinning rDNA Genomic Stability. Trends in Genetics 35:743-753.

Varki A, Altheide TK. 2005. Comparing the human and chimpanzee genomes: searching for needles in a haystack. Genome Res 15:1746-1758.

Wang X, He Z, Guo Z, Yang M, Xu S, Chen Q, Shao S, Li S, Zhong C, Duke NC, et al. 2022. Extensive

gene flow in secondary sympatry after allopatric speciation. National Science Review 9:nwac280.

Williams AL, Genovese G, Dyer T, Altemose N, Truax K, Jun G, Patterson N, Myers SR, Curran JE,

Duggirala R, et al. 2015. Non-crossover gene conversions show strong GC bias and unexpected

clustering in humans. Elife 4.

Wolfe KH, Sharp PM, Li WH. 1989. Mutation rates differ among regions of the mammalian genome. Nature

337:283-285.

Wu C-I, Lyttle TW, Wu M-L, Lin G-F. 1988. Association between a satellite DNA sequence and the

responder of segregation distorter in D. melanogaster. Cell 54:179-189.

Wu CI, Wang HY, Ling S, Lu X. 2016. The Ecology and Evolution of Cancer: The Ultra-Microevolutionary

Process. Annu Rev Genet 50:347-369.

Xu J, Nuno K, Litzenburger UM, Qi YY, Corces MR, Majeti R, Chang HY. 2019. Single-cell lineage tracing

by endogenous mutations enriched in transposase accessible mitochondrial DNA. Elife 8.

Yang H, Wang JR, Didion JP, Buus RJ, Bell TA, Welsh CE, Bonhomme F, Yu AH-T, Nachman MW, Pialek

J, et al. 2011. Subspecific origin and haplotype diversity in the laboratory mouse. Nat Genet 43:648-

655.

Zhai W, Lai H, Kaya NA, Chen J, Yang H, Lu B, Lim JQ, Ma S, Chew SC, Chua KP, et al. 2022. Dynamic

phenotypic heterogeneity and the evolution of multiple RNA subtypes in hepatocellular carcinoma: the

PLANET study. National Science Review 9:nwab192.

Zhao Z, Jin L, Fu Y-X, Ramsay M, Jenkins T, Leskinen E, Pamilo P, Trexler M, Patthy L, Jorde LB, et al.

2000. Worldwide DNA sequence variation in a 10-kilobase noncoding region on human chromosome

22. Proceedings of the National Academy of Sciences 97:11354-11358.

Zhou T, Shi T, Li A, Zhu L, Zhao X, Mao N, Qin W, Bi H, Yang M, Dai M, et al. 2022. A third dose of

inactivated SARS-CoV-2 vaccine induces robust antibody responses in people with inadequate

response to two-dose vaccination. National Science Review 9:nwac066.

Figures and Tables

Figure 1. The "chromosome community of rDNAs on five acrocentric chromosomes. (A) The genomic locations of rDNA tandem repeats in human (Gibbons, et al. 2015) and mouse (Cazaux, et al. 2011). rDNAs are located on the short arms (human), or the proximal end of the long arms (mouse), of the chromosome. Either way, inter-chromosomal exchanges are permissible. (B) The organization of rDNA repeat unit. IGS (intergenic spacer) is not transcribed. Among the transcribed regions, 18S, 5.8S and 28S segments are in the mature rRNA while ETS (external transcribed spacer) and ITS (internal transcribed spacer) are excluded. (C) The pseudo-population of rRNA genes is shown by the "chromosomes community" map (Guarracino, et al. 2023), which indicates the divergence distance among chromosome segments. The large circle encompasses rDNAs from all 5 chromosomes. It shows the concerted evolution among rRNA genes from all chromosomes, which thus resemble members of a (pseudo-)population. The slightly smaller thin circle, from the analysis of this study, shows that the rDNA gene pool from each individual captures approximately 95% of the total diversity of human population. (D) A simple illustration that shows the transmissions of two new mutations (#1 and #2 in red letter). Mutation 1 experiences replication slippage, gene conversion and unequal crossover and grows to 9 copies ($K = 9$) after transmission. Mutation 2 emerges and disappears ($K = 0$). This shows how $V(K)$ may be augmented by the homogenization process.

Figure 2. Fixation of mutations at two levels of species divergence, (T_{d1}) and (T_{d2}). (T_{f1}) and (T_{f2}) are mutations with a shorter and longer fixation time, respectively for single-copy and multi-copy genes. Note that mutations with a longer T_f would show a lower fixation rate in short-term evolution.

Figure 3. Correlation of variant frequencies between human individuals. The pairwise correlation of variant site frequency in the transcribed region of rDNAs among 6 individuals (2 Asians, 2 Europeans, and 2 Africans). The high correlations suggest that the diversity in each individual can well capture the population diversity. Each color represents a region of rDNA. The diagonal plots present the variant frequency distribution. The upper right section summarizes the Pearson correlation coefficients derived from the mirror symmetric plots in the bottom left. The analysis excluded the 18S, 3'ETS, and 5.8S regions due to the limited polymorphic sites. The result for IGS region is presented in Supplementary Figure S1.

Figure 4. Levels of polymorphism and divergence in mice. (A) Phylogeny of *Mus musculus* and *Mus*
spretus mice. The divergence times are obtained from http://timetree.org/. The line segment labeled 0.5
represents 0.5 Myrs. (B) F_{IS} distribution within *M. m. domesticus*. The distribution of F_{IS} for polymorphic
sites in 3 outbred mouse strains or 10 mouse strains (including 7 inbred mice) in Table 1 (Inset) is shown.
(C) F_{ST} distribution between *M. m. domesticus* and *Mus spretus*. Note that the *F* values rise above 0.8 only in
(C).

Table 1. rRNA gene nucleotide diversity in the 10 *M. m. domesticus* strains of a global collection.

Mouse strains	H_I (‰) - diversity within individuals	
	Functional parts (18S 5.8S and 28S)	Non-functional parts (ETS, ITS and IGS)
WSB/EiJ	0.36	6.45
ZALENDE/Ei	0.44	6.29
LEWES/EiJ	0.28	5.56
BALB/cJ	0.19	6.51
C57BL/6NJ	0.20	6.30
ST/bJ	0.24	6.73
NZW/LacJ	0.22	4.96
FVB/NJ	0.35	6.47
DBA/1J	0.22	6.44
C3H/HeJ	0.20	6.50
Averaged H_I diversity across individuals	0.27	**6.22**
H_S **for rDNA in** *M. m. domesticus*	0.34	7.25
H_S **for single-copy genes in** *M. m. domesticus*	---	1.40[*]

WSB/EiJ, ZALENDE/Ei and LEWES/EiJ are outbred strains, and the rest are inbred strains.

Sources: Wellcome Sanger Institute's Mouse Genome Project (MGP).

[*]Data from (Geraldes, et al. 2008).

Table 2. rRNA gene nucleotide diversity in the 8 humans of a global collection.

Human Individuals	H_I (‰) - diversity within individuals	
	Functional parts (18S 5.8S and 28S)	Non-functional parts (ETS, ITS and IGS)
Asia CRC	0.53	7.69
Asia 1590	0.28	7.33
Europe 4929	0.42	6.53
Europe 1383	0.32	6.74
Africa 7733	0.36	6.65
Africa 9161	0.34	6.55
Asia F9551	0.87	6.37
Asia M9552	0.88	6.58
Averaged diversity H_I across individuals	0.50	6.81
H_S for rDNA in human population	0.68	7.24
H_S for single-copy genes in human population	---	0.88[*]

8 humans are from three continents (4 Asians, 2 Europeans and 2 Africans).

Source: National Center for Biotechnology Information (NCBI).

[*] Data from (Zhao, et al. 2000).

Table 3. Divergence in rRNA genes between *M. m. domesticus* and *M. m. castaneus*.

	Length	mapping length (L) (w/o CpG sites)	Divergent site (D) (w/o CpG sites)	D/(L/1000) (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	6757	6755 (5321)	4(4)	0.6(0.8)
Non-Functional parts				
5' ETS	4007	4006 (3165)	9(7)	2.2(2.2)
ITS1+ITS2	2088	2088 (1530)	11(7)	5.3(4.6)
3'ETS	551	539 (371)	1(1)	1.9(2.7)
IGS	31902	26101 (25461)	45(36)	1.7(1.4)
Genome-wide				
Single-copy genes	-	-	-	9.5

  Note the larger divergence of single-copy genes (9.5) than for rRNA genes (1.7 - 5.3)

Table 4. Divergence in rRNA genes between *M. m. domesticus* and *Mus spretus*.

	Length	mapping length L (w/o CpG sites)	Divergent site D (w/o CpG sites)	D/(L/1000) (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	6757	6757(5321)	12(10)	1.8(1.9)
Non-Functional parts				
5' ETS	4007	4005(3165)	100(68)	25.0(21.5)
ITS1+ITS2	2088	2077(1530)	54(38)	26.0(24.8)
3'ETS	551	551(371)	17(13)	30.9(35.0)
IGS	31902	25516(24796)	270(241)	10.6(9.7)
Genome-wide				
Single-copy genes	-	-	-	21

796 Note the divergence of single-copy genes (21) is often smaller than rRNA genes (25.0 - 30.9)

797 The divergence between *M. m. musculus* and the two outgroups species of *M. m. castaneus* and *Mus spretus*

798 are listed in Supplementary Table S2 – S3.

Table 5. Divergence in rRNA genes between Human and Chimpanzee.

	Length	mapping length L[*] (w/o CpG sites)	Divergent site D[†] (w/o CpG sites)	D/(L/1000) [‡] (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	7063	7036(5496)	13±0.53 (10±0.35)	1.85±0.08 (1.81±0.06)
Non-Functional parts				
5' ETS	3656	3608(2446)	82±1.07 (55±0.53)	22.73±0.30 (22.4±0.22)
ITS1+ITS2	2250	2234(1430)	24±3.34 (12+2.31)	10.74±1.49 (8.39±1.62)
3'ETS	345	345(215)	7±0.53 (7±0.53)	20.29±1.55 (32.56±2.47)
IGS	29685	29452(26650)	544±11.80 (410±8.01)	18.47±0.40 (15.38±0.30)
Genome-wide				
Single-copy genes	3.2billion	-	35million	11(Varki and Altheide 2005)

803 Note the divergence of single-copy genes (= 11) is usually smaller than rRNA genes.

804 Human data are from Table 2 and chimpanzee genome sequencing data are from (Tatsumoto, et al. 2017).

805 [*] The mapping length between two species; [†] Divergent sites ($F_{ST} > 0.8$) are shown as the mean ± SD; [‡]

806 Divergent sites per kilobase are shown as the mean ± SD. SD, standard deviation.

Table 6. The A/T to G/C and G/C to A/T changes in apes and mouse.

	Direction of changes	Not fixed mutation counts	Fixed mutation counts	Chi-square test
Human	A/T to G/C	163	223	P = 1.559e-15
	G/C to A/T	159	48	
Chimpanzee	A/T to G/C	289	210	P = 0.000469
	G/C to A/T	201	83	
M. m. domesticus	A/T to G/C	99	10	P = 0.9568
	G/C to A/T	156	14	

810 New mutants with a frequency >0.8 within an individual are considered as (nearly) fixed, except for

811 humans, where the frequency was averaged over 8 individuals in the Table 2. The X-squared values for each

812 species are as follows: 63.556 for human, 12.236 for chimpanzee, and 0.0029283 for *M. m. domesticus*.

817 **Table 7.** The parameter values of p, q, f and g in the evolution between A/T and G/C.

Species	**Human**			**Chimpanzee**			***M. m. domesticus***		
Sequence segments	Non-functional	Total rDNA	Whole genome	Non-functional	Total rDNA	Whole genome	Non-functional	Total rDNA	Whole genome
p	0.43	0.42		0.43	0.42		0.54	0.51	
q	0.57	0.58	0.41	0.57	0.58	0.42	0.46	0.49	0.42
f	0.0142	0.0125		0.0136	0.012		0.0005	0.0004	
g	0.0023	0.0019		0.0041	0.0034		0.0007	0.0006	
$(pf)/(qg)$	4.66	4.76		2.5	2.56		0.84	0.69	

819 The proportion of AT sites and GC sites are represented by p and q respectively. The proportion of AT-to-GC

820 and GC-to-AT conversions (shown in Table 6) are represented by f and g. When $(pf)/(qg) = 1$, the evolution

821 between A/T and G/C is in equilibrium.