TITLE PAGE

TITLE:

A Generalized approach to genetic drift and its applications -

I. An example of the evolution of ribosomal RNA genes

Xiaopei Wang[1], Yongsen Ruan[1], Lingjie Zhang[1], Xiangnyu Chen[1], Zongkun Shi[1], Haijun Wen[1,*], Chung-I Wu[1,*]

[1]State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, China

[*] Corresponding authors: wenhj5@mail.sysu.edu.cn; m18819268566@163.com

RUNNING TITLE:

genetic drift and rRNA genes

ABSTRACT

Genetic drift should be any random forces that change gene frequency stochastically. While the conventions including the WF (Wright-Fisher) model are about sampling errors between generations, the branching process (BC) has been known to be a more general framework. In addition to connecting ecology and population genetics, BC can address genetic drift in systems of multi-copy sequences that include transposons, viruses and multi-copy genes. Here, we apply this general concept to rRNA genes, which have a median of 300 copies per human individual (designated $C \sim 150$ copies for haploid). By using BC, we determine the strength of genetic drift in rRNA genes to be ~ 20 times higher than single-copy genes in human and mouse polymorphisms. The large increases in drift, likely due to the homogenizing forces (such as unbiased gene conversion) within individuals, reduce Ne* to < 10Ne, despite $C \sim 150$ (Ne* and Ne being the effective population sizes of rRNA genes and single-copy genes respectively). Significantly, when the divergence between species is analyzed, some variants appear to experience extremely strong drift such that Ne* becomes smaller than Ne, as if C < 1. Therefore, the rapid divergence of rDNAs between mouse species can be easily accounted for by strong drift. Positive selection needs to be invoked only when the strong drift can be convincingly ruled out, as is between human and chimpanzee. In conclusion, the branching process reveals many stochastic forces in molecular evolution that are mistakenly attributed to selection in the conventional models.

INTRODUCTION

Genetic drift is defined as the change in gene frequency due to random forces, independent of deterministic influences such as selection or migration. The simplest of these forces is random sampling form the population of size N as formulated in the Wright-Fisher (WF) model (Crow and Kimura 1970; Kimura 1983; Hartl et al. 1997; Li 1997). The Moran model (Moran 1958; Moran and Oliphant 1997), an alternative that also defines genetic drift solely in sampling terms, can usually be scaled to correspond to the WF model. Other definitions of genetic drift, such as the Ornstein-Uhlenbeck model, are about non-specific stochastic noises in long-term evolution, unconnected to genetic variation (Szitenberg et al. 2016).

Deviations from the standard WF model are usually corrected by Ne (effective population size) in place of N. For example, the fluctuations in N can be absorbed into their harmonic mean (Crow and Kimura 1970). However, there are many other stochastic forces that elude the WF model. For example, in many systems of evolution, multiple copies of the same gene, or genomes, compete within individuals before entering the conventional competitions among individuals. We shall refer to the evolution within and between individuals as Stage I and Stage II, respectively. A general model of genetic drift should incorporate both stages, a task not suited to the WF model. In Discussion, we will further address the fundamental limitations of WF.

A common multi-copy system is viral evolution, illustrated vividly by the recent pandemic caused by SARS-CoV-2 (Ruan et al. 2021a; Ruan et al. 2021b; Ruan et al. 2022; Ruan et al. 2023). Another example is systems of large number of identical genes within the same genome such as transposons (Szitenberg et al. 2016), satellite DNAs(Wu et al. 1989; Cabot et al. 1993; Alexandrov et al. 2001), and ribosomal RNA genes (van Sluis and McStay 2019; Hori et al. 2021). In these examples, the copy number (designated C) is in the hundreds up to millions per individual. (Most multi-gene families are excluded from this consideration as their members are not functionally identical.)

A mutation that exists in some of these copies, even completely neutral in fitness, would fluctuate in frequency within individuals. Such random changes in mutation frequency are also genetic drift but these changes are not tractable by the binomial sampling. Each system would require a specific model for the genetic drift in Stage I. For rRNA genes, the forces include unequal cross-over, replication slippage and gene

conversion (Nagylaki 1983; Ohta and Dover 1983; Ohta and Gillespie 1996; Smirnov et al. 2021). For transposons, the proliferation may be even more stochastic as the host genome often suppresses the transposition of members of the same family (Wu et al. 1989; Wilkinson et al. 2021). Furthermore, the stochasticity in viral evolution is extremely high even in Stage II as only 5-6 travelers have been estimated to start the COVID-19 epidemics in each country or region (Ruan et al. 2021a).

In short, a general theoretical framework for genetic drift is needed. Its absence has often led to speculations of adaptation by natural selection when stochasticity could be a simpler explanation (Cabot et al. 1993; Alexandrov et al. 2001; Lynch and Conery 2003; Lynch 2007; Ruan et al. 2021a). A solution may be the branching process that has been applied to stochastic processes in population genetics (Crow and Kimura 1970). Recently, Chen et al. (Chen et al. 2017) have proposed the branching process as a more inclusive approach to the stochastic changes in gene frequency. In this study, we choose the rRNA genes as an example for studying the general process of genetic drift. Hence, we introduce the relevant properties of rRNA genes below.

The "chromosome community" of ribosomal RNA genes

The ribosomal RNA genes (rDNAs) are perhaps the best known multi-copy gene clusters (Bowman et al. 2020). In mammals, they are arrayed as tandem repeats on multiple chromosomes as shown in Fig. 1A (Guillén et al. 2004; Cazaux et al. 2011). In humans, the copy number can vary from 60 to 1600 per individual (mean, 315; SD, 104; median, 301) (Parks et al. 2018). We shall use C = 150 to designate the median number per haploid in humans (2C = 300). In mouse, C ~ 110 (Parks et al. 2018).

In humans, the five rRNA clusters are located on the short arm of the five acrocentric chromosomes (Smirnov et al. 2021). Such an arrangement permits crossovers between chromosomes without perturbing the rest of the genomes. In *Mus*, the rDNAs are all located in the pericentromeric or subtelomeric region, on the long arms of telocentric chromosomes (Cazaux et al. 2011; Potapova and Gerton 2019). Thus, unequal crossovers between non-homologous chromosomes may involve centromeres while other genic regions are also minimally perturbed.

Fig. 1. The "chromosome community of rDNAs on five acrocentric chromosomes
(A) The genomic locations of rDNA tandem repeats in human (Gibbons et al. 2015) and mouse (Cazaux et al. 2011).
rDNAs are located on the short arms (human), or the proximal end of the long arms (mouse), of the acrocentric
chromosome. Either way, inter-chromosomal exchanges are permissible. (B) The organization of rDNA repeat unit.
IGS (intergenic spacer) is not transcribed. Among the transcribed regions, 18S, 5.8S and 28S segments are in the
mature rRNA while ETS (external transcribed spacer) and ITS (internal transcribed spacer) are excluded. (C) The
"chromosomes community" map modified from Fig. 1 of (Guarracino et al. 2023). The map shows the distance of
divergence among chromosome segments. The large circle encompasses rDNAs from all 5 chromosomes with color
dots indicating the chromosome origin. It is clear that rDNAs experience concerted evolution regardless of their
genomic locations. The slightly smaller thin circle, from the analysis of this study, shows that the rDNA gene pool
from each individual captures approximately 95% of the total diversity of human population. (D) A simple illustration
that shows the transmissions of two new mutations (#1 and #2 in red letter). Mutation 1 experiences replication
slippage, gene conversion and unequal crossover and grows to 9 copies (k=9) after transmission. Mutation 2 emerges
and disappears (k = 0). This shows how V(k) may be augmented by the homogenization process.

Each copy of rRNA gene has functional and non-functional parts as shown in Fig. 1B. The "functional"
regions of rDNA, 18S, 5.8S, and 28S rDNA, are believed to be under strong negative selection, resulting in
a slow evolution rate (Salim and Gerton 2019). In contrast, the transcribed spacer (ETS and ITS) and the
intergenic spacer (IGS) show significant sequence variation even among closely related species (Eickbush
and Eickbush 2007). This suggests that these "non-functional" regions are less constrained by negative
selection. In this study of genetic drift, we focus on the non-functional parts. Data on the evolution of the
functional parts will be provided only for comparisons.

While a human individual with 300 rRNA genes may appear to have 300 loci, the concept of distinct
"gene loci" cannot be applied to the rRNA gene clusters. This is because DNA sequences can spread from
one copy to others on the same chromosome via replication slippage. They can also spread among all other
copies on the same or different chromosomes via gene conversion and unequal crossovers (Nagylaki 1983;
Ohta and Dover 1983; Stults et al. 2008; Smirnov et al. 2021). Replication slippage and unequal crossovers
would also alter the copy number of rRNA genes, thus accounting for the copy number variation among
individuals. These mechanisms will be referred to collectively as the homogenization process.

Copies of the cluster on the same chromosome are widely known to be nearly identical in sequences
(Hori et al. 2021; Nurk et al. 2022). Previous research has also provided extensive evidence that rDNA
clusters among nonhomologous chromosomes share the same variants, thus confirming genetic exchange
between chromosomes (Krystal et al. 1981; Arnheim et al. 1982; van Sluis et al. 2019). For example, Nurk
et al. (Nurk et al. 2022) report a 5-kb segment that covers mostly non-functional sites showing a 98.7%
identity among the short arms of the five acrocentric chromosomes. Additionally, variants are often found in
all human rDNAs but not in other great apes, suggesting rapid fixation (Arnheim et al. 1980). The rapid
fixation is unexpected because, for a new rRNA mutation to be fixed, it has to be fixed among all ~ 300
copies in all individuals in the population. In this process, both rapid drift and positive selection can
contribute to these fixations. However, the main difference between them is that genetic drift only shorten
the fixation time of mutations without changing their fixation probability, whereas selection can alter both.
As a result, rapid drift can only accelerate the evolution rate up neutral molecular evolutionary rate (Eq. (3)
below), while positive selection is not constrained by this limit.

With the extensive exchanges via the homogenization process, copies of rRNA genes in an individual
can be treated as a sub-population of gene copies. Such sub-populations do not have to be in the sense of a
Mendelian population when using the branching process to track genetic drift (see Discussion and Chen et
al. 2017). (Guarracino et al. 2023) recently proposed the view of "chromosome community" for the short

arms of the 5 acrocentric chromosomes, where rRNA genes are located. As seen in Fig. 1C, these five short arms harbor a pool of rRNA genes that do not have strong associations with genomic locations. Fig. 1C further illustrates that each human individual's rRNA gene pool can capture ~ 95% of the total diversity of the human populations (see Results).

Our goal is to propose a framework for studying the genetic drift in systems like rRNAs with multiple nearly identical copies within each individual. While we choose rRNA genes, the analysis of genetic drift in systems with two stages of evolution, within as well as between individuals, should be general.

RESULTS

PART I (Theory) of Results consolidate all relevant theories. The level of polymorphism within species is separately considered from the level of divergence between species. In PART II (Data analyses), we apply the theory to the study of rDNA evolution in mice and apes (human and chimpanzee). The comparisons between these two mammalian taxa are most interesting as the driving forces have evolved distinct dynamics in the two lineages.

Empirical measurements of homogenization in vitro and in vivo

The homogenizing mechanisms such as gene conversion, unequal crossover and replication slippage drive the rapid genetic drift in rDNAs. Direct measurements in the literature are usually based on the comparisons among individuals Therefore, the Mendelian mechanisms of chromosome segregation and assortment would also shuffle variants among individuals. Segregation and assortment would confound the measurements of homogenization effects within individuals.

In a separated study, we measure the homogenization effects in rDNAs using cultured cell lines over an extended period of time. The measurements show the homogenization effects to be sufficiently strong to account for the long-term acceleration of genetic drift reported in this study. For the homogenization effects in vivo, we analyze somatic evolution in solid tumors. taking advantage of the single-cell origin of each solid tumor. The divergence among samples obtained from different sites of the same tumor also reveals very genetic drift of rDNA mutations. These direct empirical measurements of homogenization will shed further insight into the general process of genetic drift.

PART I - Theory

1. Generalized genetics drift

In its basic form, genetic drift is the change in gene frequency due strictly to stochastic elements that include forces beyond sampling fluctuations. Therefore, an intuitively appealing approach, alternative to the WF model, is the branching process whereby each copy of the gene leaves k copies in a time interval. Given $E(k)$ and $V(k)$ (for the mean and variance of k), genetic drift is simply $V(k)$. If $V(k) = 0$, there is no gene frequency change and no genetic drift.

In the WF model, $V(k) = E(k)$ because the distribution of k would be Poisson. However, $V(k) = E(k)$ is untenable in most species (Chen et al. 2017). For example, $E(k)$ is the same between the two sexes but $V(k)$ is usually different; hence, $V(k) = E(k)$ is unlikely. In the branching process, genetic drift is a much more flexible process as $V(k)$ is decoupled from $E(k)$. In Discussion, we will review these issues further.

Below, we compare the strength of genetic drift in rRNA genes vs. that of single-copy genes using the branching process. $E(k)$ is set to 1 (i.e., N being near the carrying capacity of the population) so that we can focus on $V(k)$. For single-copy genes, the effective population size that determines genetic drift would be $Ne = N/V(k)$ (Kimura and Crow 1963). We shall use * to designate the equivalent symbols for rRNA genes.

Hence,

$$Ne^* = N^*/V^*(k) = C^*N / V^*(k) \qquad \text{Eq. (1)}$$

where N* is the number of rRNA genes in the entire population and, therefore, N* = C* N. For rRNA genes, C* >> 1 but < 150, C* being the effective copy number of rRNA genes. (C =1 for single copy genes.) The value of C* is uncertain because each haploid in humans, on average, has ~ 150 copies that do not segregate but assort among chromosomes (Parks et al. 2018). V*(k) for rRNA genes is likely to be much larger than V(k) because k for rRNA mutations may be affected by a host of factors including replication slippage, unequal cross-over, gene conversion and other related mechanisms not operating on single copy genes. We thus designate Vh = V*(k)/V(k), as the homogenizing factor of rRNA genes.

Fig. 1D is a simple illustration that the homogenizing process may enhance V(k) substantially over the conventional WF model as each mechanism amplifies the variance in K. Importantly, the homogenization process is considered stochastic rather than deterministic because it affects both the mutants and wild type equally, as expected of genetic drift.

We then designate R* = (1/Ne*)/(1/Ne) as the strength of genetic drift of rRNA genes, relative to that of single copy genes. R* is measurable, as will be done below.

$$R^* = (1/Ne^*)/(1/Ne) = [V^*(k)/V(k)]/C^* = Vh/C^* \qquad \text{Eq. (2)}$$

In short, R* is the ratio of two factors. A larger Vh speeds up the drift strength of rRNA genes whereas a larger C* slows down the process of fixation by drift. It would be most interesting if R* > 1 as it would mean that the effective population size of rRNA genes is smaller than single-copy genes. In this case, the homogenization force should be strong enough to overcome the large copy number of rRNA genes (~ 150) during the fixation of new mutations.

2. rDNA polymorphism within species

For rDNAs, there are 3 levels of diversity: H_I, H_S and H_T. They represent, respectively, the heterozygosity within individual (I), within species (S) and in the total data set (T, which often comprises two species or more). Briefly, H = 2p(1-p) if there are two variants with the frequency of p and 1-p. In an equilibrium haploid population,

$$H_S = 2N_eu /[2N_eu + 1]$$

Given u, H_S is a measure of genetic drift within a population. For rRNA genes, H_S is expected to be larger than that single-copy genes as N* > N. With the three levels of heterozygosity for multi-copy genes, there are two levels of differentiation. First, F_{IS} is the differentiation among individuals within the species, defined by

$$F_{IS} = [H_S - H_I]/H_S \quad (1)$$

Second, F_{ST} is the differentiation between populations (or species) in the context of the total diversity, defined as

$$F_{ST} = [H_T - H_S]/H_T \quad (2)$$

The two F statistics will inform about the distribution of nucleotide diversity among individuals, or between

species. H_T is the expected heterozygosity when there is no subdivision between species and H_s is the observed heterozygosity. Therefore, F_{ST} will be equal to 0 when there is no subdivision and will be positive as long as there is variation in allele frequency across subdivisions.

3. rDNA divergence between species

Whereas the level of genetic diversity is a function of the effective population size, the rate of divergence between species, in its basic form, is not. The rate of neutral molecular evolution is (λ), although driven by mutation and genetic drift, is generally shown by Eq. (3) below (Crow and Kimura 1970; Hartl et al. 1997; Li 1997):

$$\lambda = N \, u \, [1/N] = u \qquad\qquad \text{Eq. (3).}$$

Hence, we might view the evolution of rRNA genes in the long run to be the same as single-copy genes. However, Eq. (3) is valid only in the long-term evolution. For shorter-term evolution, it is necessary to factor in the fixation time (Fig. 2), T_f, which is the time between the emergence of a mutation and its fixation. If we study two species with a divergent time (T_d) equal to or smaller than T_f, then few mutations of recent emergence would have been fixed as species diverge.

Note that T_d is about 6 million years (Myrs) between human and chimpanzee while T_f (as measured by coalescence) is 0.6 - 1 Myrs in humans. Mutations of single-copy genes would not get fixed during the more recent T_f, as indicated in Figure 2. Thus, λ may be 1/6 to 1/10 lower than the theoretical value. In comparison, T_f of rRNA mutations should be at least 4.8 - 8 Myrs based on the C* estimate. Thus, λ may be at least 80% lower than calculated for single-copy genes.

Fig. 2. Fixation of mutations at two levels of species divergence, (T_{d1}) and (T_{d2}). (T_{f1}) and (T_{f2}) are mutations with a shorter and longer fixation time, respectively for single-copy and multi-copy genes. Note that mutations with a longer T_f would show a lower fixation rate in short-term evolution.

Our own theoretical derivation has shown that the fixation time for rRNA genes is close to 4Ne* as is the case for single-copy genes at ~ 4Ne. If V*(k) is sufficiently larger than V(k), it is in fact possible for Ne* < Ne such that genetic drift is stronger for rRNA genes than for single-copy genes and T_f* < T_f. Therefore, T_{f1} can represent the T_f* of mutations in rRNA genes in Fig. 2. This is interesting because, if the homogenization is powerful enough, rRNA genes would have an effective copy number of C* < 1. A short T_f*, which can be obtained by large V*(k), would lead to a small T_f*/T_d and thus a higher λ,   particularly in

short-term evolution. However, even T_f^* approaches 0, λ can exceed that of single-copy genes but still limited by fixation probability. As noted above, the rapidly fixed mutations may not be captured in the polymorphic data but would appear in species divergence.

PART II - Data Analyses

The theory of PART I is now used on rDNA data in PART II. As in PART I, we shall first analyze rDNA polymorphisms within species, in mouse and human separately. On the basis of polymorphism results, we shall study the divergence between species.

1. rDNA polymorphism within species

1) Polymorphism in mice

For rRNA genes, H_I of 10 individuals ranges from 0.0050 to 0.0067 while H_S is 0.0073 (Table 1). Thus, $F_{IS} = [H_S - H_I]/H_S$ for mice is 0.14, which means 86% of variation is within each individual. In other words, even one single randomly chosen individual would yield 85% of the diversity of the whole species. Hence, the estimated H_S should be robust as it is not affected much by the sampling.

Table 1. rRNA gene diversity in the 10 *M. m. domesticus* strains of a global collection.

Mouse strains	H_I (‰) - diversity within individuals	
	Functional parts (18S 5.8S and 28S)	Non-functional parts (ETS, ITS and IGS)
WSB/EiJ	0.36	6.45
ZALENDE/Ei	0.44	6.29
LEWES/EiJ	0.28	5.56
BALB/cJ	0.19	6.51
C57BL/6NJ	0.20	6.30
ST/bJ	0.24	6.73
NZW/LacJ	0.22	4.96
FVB/NJ	0.35	6.47
DBA/1J	0.22	6.44
C3H/HeJ	0.20	6.50
Averaged diversity across individuals	0.27	**6.22**
H_S for rDNA in *M. m. domesticus*	0.34	7.25
H_S for single-copy genes in *M. m. domesticus*	---	1.40[a]

WSB/EiJ, ZALENDE/Ei and LEWES/EiJ are outbred strains, and the rest are inbred strains.
Sources: Wellcome Sanger Institute's Mouse Genome Project (MGP).
[a] Data from (Geraldes et al. 2008).

H_S for *M. m. domesticus* single-copy genes is roughly 1.40 per kb genome-wide (Geraldes et al. 2008) while H_S for rRNA genes is 7.25 per kb, 5.2 times larger. In other words, $C^* = N^*/N \sim 5.2$. If we use the polymorphism data, it is as if rDNA array has a population size 5.2 times larger than single-copy genes. Although the actual copy number on each haploid is ~ 110, these copies do not segregate like single-copy genes and we should not expect N^* to be 100 times larger than N. The H_S results confirm the prediction that rRNA genes should be more polymorphic than single-copy genes.

Based on the polymorphism result, one might infer slower drift for rDNAs than for single-copy genes. However, the results from the divergence data in later sections will reveal the under-estimation of drift strength from polymorphism data. Such data would miss variants that have a fast drift process driven by, for example, gene conversion. Strength of genetic drift should therefore be measured by the long-term fixation rate.

2) Polymorphism in human

F_{IS} for rDNA among 8 human individuals is 0.058 (Table 2), much smaller than 0.142 in *M. m. domesticus* mice, suggesting a major role of stage I evolution in humans. Consisted with low F_{IS}, Fig. 3

Table 2. rRNA gene diversity in the 8 humans of a global collection.

	H_I (‰) – diversity within individuals	
	Functional parts (18S 5.8S and 28S)	**Non-functional parts (ETS, ITS and IGS)**
Asia CRC	0.53	7.69
Asia 1590	0.28	7.33
Europe 4929	0.42	6.53
Europe 1383	0.32	6.74
Africa 7733	0.36	6.65
Africa 9161	0.34	6.55
Asia F9551	0.87	6.37
Asia M9552	0.88	6.58
Averaged diversity across individuals	0.50	6.81
H_S for rDNA in human population	0.68	7.24
H_S for single-copy genes in human population	---	0.88[a]

8 humans are from three continents (4 Asians, 2 Europeans and 2 Africans).

Source: National Center for Biotechnology Information (NCBI).

[a] Data from (Zhao et al. 2000).

shows strong correlation of the polymorphic site frequency of rDNA transcribed region among each pair of individuals from three continents (2 Asians, 2 Europeans and 2 Africans). Correlation of polymorphic sites in IGS region is shown in Supplementary Fig. S1. Like those in mice, the pattern indicates that intra-species

polymorphism is mainly preserved within individuals. In addition, the 6 panels on the diagonal line given in Fig. 3 demonstrated that the allele frequency spectrum is highly concentrated at very low frequency within individuals, indicating rapid homogenization of rDNA mutations among different copies even different chromosomes. This homogenization can lead to the elimination of the genetic variation of this region. The observed H_I of humans for rDNAs is 0.0064 to 0.0077 and the H_S is 0.0072 (Table 2). Research has shown that heterozygosity for the human genome is about 0.00088 (Zhao et al. 2000), meaning the effective copy number of rDNAs is roughly, or $C^* \sim 8$.

Fig. 3. Correlation of variant frequencies between individuals.

The pairwise correlation of variant site frequency in the transcribed region of rDNAs among 6 individuals (2 Asians, 2 Europeans, and 2 Africans). The high correlations suggest that the diversity in each individual can well capture the population diversity. Each color represents a region of rDNA. The diagonal plots present the variant frequency distribution. The upper right section summarizes the Pearson correlation coefficients derived from the mirror symmetric plots in the bottom left. The analysis excluded the 18S, 3'ETS, and 5.8S regions due to the limited polymorphic sites. The result for IGS region is presented in Supplementary Figure S1.

2. rDNA divergence between species

   We now consider the evolution of rRNA genes between species by analyzing the rate of fixation (or near fixation) of mutations. Polymorphic variants are filtered out in the calculation.

1) Between mouse species - Genetic drift as the sole driving force of the rapid divergence

   We now use the F_{ST} statistic to delineate fixation and polymorphism. The polymorphism in *M. m. domesticus* is compared with two outgroup species, *M. spretus* and *M. m. castaneus*, respectively. There are hence two depths in the phylogeny with two T_d's, as shown in Fig. 4A (Rudra et al. 2016; Kumar et al. 2022). There is a fourth species, *M. m. musculus* (shown in grey in Fig. 4A)*,* which yields very similar

results as *M. m. domesticus* in these two comparisons. These additional analyses are shown in Supplement.

Fig. 4. Levels of polymorphism and divergence in mice. (A) Phylogeny of *Mus musculus* and *Mus spretus* mice. The divergence times are obtained from http://timetree.org/. The line segment labelled 0.5 represents 0.5 Myrs. (B) F_{IS} distribution within *M. m. domesticus*. The distribution of F_{IS} for polymorphic sites in 3 outbred mouse strains or 10 mouse strains (including 7 inbred mice) in Table 1 (Inset) as shown. (C) F_{ST} distribution between *M. m. domesticus* and *Mus spretus*. Note that the F values rise above 0.8 only in (C).

The F_{IS} values of polymorphic sites in 3 outbred mice are primarily below 0.2 and rarely above 0.8 in Fig. 4B, indicating the low genetic differentiation in rDNAs within these 3 *M. m. domesticus*. While the F_{IS} distribution of 10 mice from Table 1, including 7 inbred and 3 outbred mouse strains, exhibits a noticeable right skewness, but does not exceed 0.8. This suggests that inbreeding to a certain extent limits the process of homogenization and enhances population differentiation. In comparison, the distribution of the F_{ST} of variant sites between *M. m. domesticus* and *Mus spretus* has a large peak near $F_{ST} = 1$. This peak in Fig. 4C represents species divergence not seen within populations (i.e., F_{IS}). We use $F_{ST} = 0.8$ as a cutoff for divergence sites between the two species. Roughly, when a mutant is > 0.95 in frequency in one species and < 0.05 in the other, F_{ST} would be > 0.80.

We first compare the divergence between *M. m. domesticus* and *M. m. castaneus* whereby T_d has been estimated to be less than 0.5 Myrs (Fujiwara et al. 2022). In comparison, between *Mus m. domesticus* and *Mus spretus*, T_d is close to 3 Myrs (Rudra et al. 2016). As noted above, the reduction in the divergence rate relative to that of Eq. (3) is proportional to T_f/T_d (for single copy genes) or T_f^*/T_d (for rRNA genes). As T_f and T_f^* are both from *M. m. domesticus* and T_d is 6 times larger in comparison with *M. spretus*, we expect the results to be quite different between the two sets of species comparisons.

Although T_f and T_f^* estimates are less reliable than T_d estimates, both comparisons use the same T_f and T_f^* from *M. m. domesticus*. Hence, the results should be qualitatively unbiased. For a demonstration, we shall use the estimates of T_f (i.e., the coalescence time) at 0.2 Myrs for single-copy genes by using an average nucleotide diversity of 0.0014 and the mutation rate of 5.7×10^{-9} per base pair per generation (Geraldes et al. 2008; Phifer-Rixey et al. 2020). Based on the estimated C* above, we obtain T_f^* for rDNA mutations at 5 x 0.2 Myrs, or 1 Myrs. While some have estimated T_f to be close to 0.4 Myrs (Fujiwara et al.

2022), we aim to show that the pattern of reduction in rRNA divergence is true even with the conservative estimates.

Between $M.\ m.\ domesticus$ and $M.\ m.\ castaneus$, the reduction in substitution rate for single copy gene should be $\sim 40\%$ ($T_f/T_d = 0.2/0.5$), and the reduction for rRNA genes should be 100% ($T_f/T_d = 1/0.5 > 1$). Table 3 on their DNA sequence divergence shows that rRNA genes are indeed far less divergent than single-copy genes. In fact, only a small fraction of rRNA mutations is expected to be fixed as T_f^* for rDNA at 1 Myrs is 2 times larger than the divergence time, T_d. We should note again that the non-negligible fixation of rRNA mutations suggests that C* at 5 is perhaps an over-estimate.

Table 3. Divergence in rRNA genes between $M.\ m.\ domesticus$ and $M.\ m.\ castaneus$.

	Length	mapping length L (w/o CpG sites)	Divergent site D (w/o CpG sites)	D/(L/1000) (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	6757	6755 (5321)	4(4)	0.6(0.8)
Non-Functional parts				
5' ETS	4007	4006 (3165)	9(7)	2.2(2.2)
ITS1+ITS2	2088	2088 (1530)	11(7)	5.3(4.6)
3'ETS	551	539 (371)	1(1)	1.9(2.7)
IGS	31902	26101 (25461)	45(36)	1.7(1.4)
Genome-wide				
Single-copy genes	-	-	-	9.5

Note the larger divergence of single-copy genes (9.5) than for rRNA genes (1.7 - 5.3)

Table 4. Divergence in rRNA genes between $M.\ m.\ domesticus$ and $Mus\ spretus$.

	Length	mapping length L (w/o CpG sites)	Divergent site D (w/o CpG sites)	D/(L/1000) (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	6757	6757(5321)	12(10)	1.8(1.9)
Non-Functional parts				
5' ETS	4007	4005(3165)	100(68)	25.0(21.5)
ITS1+ITS2	2088	2077(1530)	54(38)	26.0(24.8)
3'ETS	551	551(371)	17(13)	30.9(35.0)
IGS	31902	25516(24796)	270(241)	10.6(9.7)
Genome-wide				
Single-copy genes	-	-	-	21

Note the divergence of single-copy genes (21) is often smaller than rRNA genes (25.0 - 30.9)

The divergence between $M.\ m.\ musculus$ and the two outgroups species of $M.\ m.\ castaneus$ and $Mus\ spretus$ are listed in Supplementary Table S1 - S2.

Between $Mus\ m.\ domesticus$ and $Mus\ spretus$, the reduction in actual evolution rate from theoretical

limit for single-copy genes should be 6.7% ($T_f/T_d = 0.2/3$) and, for rRNA genes, should be 33% ($T_f^*/T_d = 1/3$). The evolutionary rate (i.e. the fixation rate) of IGS region is lower than single-copy genes, 0.01 in IGS and 0.021 in genome-wide (Table 4), as one would expect. However, ETS and ITS regions have evolved at a surprising rate that is 12% higher than single-copy genes. Note that the reduction in C*, even to the lowest limit of C* =1, would only elevate the rate of fixation in rRNA genes to a parity with single-copy genes. From Eq. 2, the explanation would be that the homogenization factor V*(k)/V(k) has to be very large, such that C* is < 1. With such rapid homogenization, the fixation time approaches 0 and the evolutionary rate in rRNA genes can indeed reach the theoretical limit of Eq. (3). In such a scenario, the substitution rate in ETS and ITS, compared to single-copy genes in mice, may increase by 7%, $T_f/(T_d - T_f) = 0.2/(3-0.2)$. If we use T_f ~ 0.4 Myrs in an alternative estimation, the increase can be up to 15%.

In conclusion, the high rate of fixation in ETS and ITS may be due to very frequent gene conversions that reduce C* to be less than 1. In contrast, IGS may have undergone fewer gene conversions and its long-term C* is slightly larger than 1. Indeed, the heterozygosity in IGS region, at about 2-fold higher than that of ETS and ITS regions (8‰ for IGS, 5‰ for ETS and 3‰ for ITS), supports this interpretation.

2) Between Human and Chimpanzee - Positive selection in addition to rapid drift in rDNA divergence

Like the data of mouse studies, the polymorphism of rDNAs in humans would suggest a slower long-term evolution rate. The same caveat is that C* estimated from the polymorphism data would have missed those rapidly fixed variants. Hence, the long-term C* obtained from species divergence might be much smaller than 8.

Our results show that the evolutionary rate of rRNA genes between human and chimpanzee is substantially higher than that of other single-copy genes (Table 5). Especially, 5'ETS region shows a 100% rate acceleration, at 22.7‰ vs. 11‰ genome-wide. Even after removing CpG sites that are considered to have a ten-fold higher mutation rate, their fixation rate still reaches 22.4‰. In this case, even if C* <<1, the extremely rapid fixation will only increase the substitution rate by $T_f / (T_d - T_f)$ by 11%, compared to single-copy genes. Thus, the much accelerated evolution of rRNA genes between humans and chimpanzees cannot be entirely attributed to genetic drift. Other forces, such as biased gene conversion, must have contributed to this acceleration as well.

Table 5. Divergence in rRNA genes between Human and Chimpanzee.

	Length	mapping length L[a] (w/o CpG sites)	Divergent site D[b] (w/o CpG sites)	D/(L/1000)[c] (w/o CpG sites)
Functional parts				
18S+5.8S + 28S	7063	7036(5496)	13±0.53 (10±0.35)	1.85±0.08 (1.81±0.06)
Non-Functional parts				
5' ETS	3656	3608(2446)	82±1.07 (55±0.53)	22.73±0.30 (22.4±0.22)
ITS1+ITS2	2250	2234(1430)	24±3.34 (12+2.31)	10.74±1.49 (8.39±1.62)
3'ETS	345	345(215)	7±0.53 (7±0.53)	20.29±1.55 (32.56±2.47)
IGS	29685	29452(26650)	544±11.80 (410±8.01)	18.47±0.40 (15.38±0.30)
Genome-wide				
Single-copy genes	3.2billion	-	35million	11(Varki, A.2005,Genome Res)

Note the divergence of single-copy genes (11) is usually smaller than rRNA genes.

Human data are from Table 2 and chimpanzee genome sequencing data are from (Tatsumoto et al. 2017).

3) Biased gene conversion - Positive selection for rRNA mutations in apes, but not in mice

As stated, gene conversion should increase V(k) and speed up genetic drift. Conversion is commonly observed in highly recombinogenic regions, especially in repetitive sequences. If gene conversion is unbiased, the effect on the rate of divergence between species would be quite modest unless $T_d <= T_f$. This is because unbiased conversion would only reduce the fixation time, T_f, as demonstrated in Fig. 2.

In this context, it is necessary to define unbiased gene conversion as it has different meanings at the site level vs. at the gene level. At the site level, when an AT site is paired with a GC site, it has been commonly reported that gene conversion would favor AT-to-GC over GC-to-AT conversion (Jeffreys and Neumann 2002; Meunier and Duret 2004). We first designate the proportion of AT-to-GC conversion as f and the reciprocal, GC-to-AT, as g. Given f ≠ g, this bias is true at the site level. However, at the genic level, the conversion may in fact be unbiased for the following reason. Let p be the proportion of AT sites and q be the proportion of GC sites in the gene. The flux of AT-to-GC would be pf and the flux in reverse, GC-to-AT, would be qg. At equilibrium, pf = qg. Given f and g, the ratio of p and q would eventually reach p/q = g/f. Thus, gene conversion at the genic level would be unbiased when the AT/GC ratio is near the equilibrium. In other words, there is no selection driving either AT-to-GC or GC-to-AT changes.

We now show that *genic* conversion is indeed unbiased (or nearly unbiased) between mouse species. In contrast, between human and chimpanzee, the fluxes are unequal. It appears that selection has favored the AT-to-GC changes in the last 6 million years that separate the two species.

In these genic analyses, we first determine whether there is biased conversion in the human lineage (Brown and Jiricny 1989; Galtier and Duret 2007). Using the same variant found in chimpanzees and gorillas as the ancestral state, we identified the derived variants that became nearly fixed in humans with frequency > 0.8 (Table 6). The chi-square test shows that the GC variants had a significantly higher fixation probability compared to AT. In addition, this pattern is also found in chimpanzees (p < 0.001). Between *M. m. domesticus* and *M. m. castaneus* (Table 6), the chi-square test reveals no difference in the fixation probability between GC and AT (p = 0.957). Further details can be found in Supplementary Figure S2. Overall, a higher fixation probability of the GC variants is found in human and chimpanzee, whereas this bias is not observed in mice.

Table 6. The A/T to G/C and G/C to A/T changes in apes and mouse.

	Direction of changes	Not fixed	Fixed	Chi-square test
Human	A/T to G/C	163	223	P < 0.001
	G/C to A/T	159	48	
Chimpanzee	A/T to G/C	289	210	P < 0.001
	G/C to A/T	201	83	
M. m. domesticus	A/T to G/C	99	10	P = 0.957
	G/C to A/T	156	14	

Based on Table 6, we could calculate the value of p, q, f and g as shown in Table 7. In particular, the pf/qg ratio is much larger than 1 in both the human and chimpanzee lineages. In contrast, the ratio in mouse is not significantly different from 1. The ratio for total rDNAs in mouse is lower than 1, but that is likely due to the influence of selective constraints on the functional parts of these genes.

Table 7. The parameter values of p, q, f and g in the evolution between A/T and G/C.

	Human			Chimpanzee			*M. m. domesticus*		
	Non-functional	Total rDNA	Whole genome	Non-functional	Total rDNA	Whole genome	Non-functional	Total rDNA	Whole genome
p	0.43	0.42		0.43	0.42		0.54	0.51	
q	0.57	0.58	0.41	0.57	0.58	0.42	0.46	0.49	0.42
f	0.0142	0.0125		0.0136	0.012		0.0005	0.0004	
g	0.0023	0.0019		0.0041	0.0034		0.0007	0.0006	
(pf)/(qg)	4.66	4.76		2.5	2.56		0.84	0.69	

The proportion of AT sites and GC sites are represented by p and q respectively. The proportion of AT-to-GC and GC-to-AT conversions (shown in Table 6) are represented by f and g. When $(pf)/(qg) = 1$, the evolution between A/T and G/C is in equilibrium.

Discussion

To conclude the study, we first discuss the specific implications of the evolution of rDNAs as well as other multi-copy gene systems. These discussions pertain to PART II of Results. We then discuss the broader stochastic forces that together constitute genetic drift, as presented in PART I of Results.

On the evolution of rRNA gene clusters and multi-copy gene systems

In multi-copy systems, there are several mechanisms of homogenization. For rRNA genes, whether on the same or different chromosomes (Gonzalez and Sylvester 2001; van Sluis and McStay 2019), the predominant mechanism of homogenization is recombination that may lead to gene conversion or unequal crossover. In the process of exchanging DNA sections in meiosis, gene conversions are an order of magnitude more common than crossover (Cole et al. 2012; Williams et al. 2015). It is not clear how large a role is played by replication slippage which affects copes of the same cluster.

There have been many rigorous analyses that confront the homogenizing mechanisms directly. These studies (Smith 1974; Ohta 1976; Dover 1982; Nagylaki 1983; Ohta and Dover 1983) modeled gene conversion and unequal cross-over head on. Unfortunately, on top of the complexities of such models, the key parameter values are rarely obtainable. In the branching process, all these complexities are wrapped into V(k) for formulating the evolutionary rate. In such a formulation, the *collective* strength of these various forces may indeed be measurable, as shown in this study.

The branching process is a model of general processes. Hence, it can be used to track genetic drift in systems with two stages of evolution - within- and between-individuals, even though TEs, viruses or rRNA genes are very different biological entities. We use the rRNA genes to convey this point. Multi-copy genes, like rDNA, are under rapid genetic drift via the homogenization process. The drift is strong enough to reduce the copy number in the population from ~ 150N to < N. A fraction of mutations in multi-copy genes may have been fixed by drift almost instantaneously in the evolutionary time scale. This acceleration is seen in mice but would have been interpreted to be due to positive selection by the convention. Interestingly, while positive selection may not be necessary to explain the mice data, it is indeed evident in human and chimpanzee, as the evolutionary rate of rRNA genes exceeds the limit of the strong drift.

On the broad stochastic forces governing gene frequencies and the models of genetic drift

The branching process is an "individual-output" model of genetic drift (Chen et al. 2017). Hence, it does not demand the population to follow the rules of Mendelian populations. In contrast, populations in the WF

model have to be defined but can rarely correspond to any actual populations. An investigator often cannot determine the boundary of the studied population - whether at the study site, within the geographical region, or across the species' range. Although the WF model uses the concept of "effective population size" (Ne) to be connected to the actual populations, there are multiple Ne's depending on the subjects of interest ranging from gene frequency, inbreeding, sex ratio, family size, population subdivision, temporal fluctuation and so on (Kimura and Crow 1963; Kimura and Crow 1964; Crow and Kimura 1970).

An adequate model of genetic drift should have at least two parameters, E(k) and V(k), which are unlikely to be equal in magnitude in any populations. The random sampling of the WF model constrains V(k) to be equal to E(k), which can be captured by a single parameter, Ne. Thus, the WF model appears to have one parameter too few. As a result, Ne's are cumbersome fixes and it would be more straightforward to start the model with more than one parameter.

Most importantly, the population size, N, is extrinsically imposed in the conventional approaches including either the WF or Moran model (Hartl et al. 1997; Moran and Oliphant 1997). In the branching process, the population size is intrinsic to the model. It is ironic that this aspect has been a major criticism of the branching process as it would result in either population extinction or unrealistic population growth (Chen et al. 2017). The modification of the branching process by making E(k) a function of N should overcome this problem. For example, E(k) > 1 when N is below the carrying capacity (Ck) and E(k) < 1 when N is above Ck.

To regulate N, (Chen et al. 2017) adopted the logistic population growth with changing E(k)'s. When a population starts from a level far below Ck, Chen et al. obtain the unorthodox result that, as N increases, the strength of genetic drift increases, rather than decreases as is generally thought. Indeed, the WF model implicitly assumes that populations are at, or near, Ck. This is the basis of thinking that the strength of genetic drift decreases when N increases. When the exponential growth is incorporated into the WF model (Durrett 2013; Ling et al. 2015), it in fact assumes that Ck is growing exponentially and N is always near Ck. This discussion shows why the regulation of N needs to be in the model itself, rather than imposed on the model externally.

By incorporating the population size regulation, the branching process would be a model of population genetics that is connected to ecology. For that reason, it is also sufficiently flexible to study various other stochastic forces that together drive genetic drift. The evolution of rRNA genes, as shown here, is one such system among many other examples.

Methods

Data Collection

We collect whole-genome sequencing data with high coverage level. The genome sequence data of human, chimpanzee and gorilla are downloaded from National Center for Biotechnology Information (NCBI). Human individuals come from three continents (4 Asians, 2 Europeans and 2 Africans) and Asia CRC is the normal tissue of Case 1 patient from (Chen et al. 2022). Mice genomic sequences are downloaded from Wellcome Sanger Institute's Mouse Genome Project (MGP). Although some artificial selection has been performed on laboratory mouse strains (Yang et al. 2011), the WSB/EiJ, ZALENDE/Ei and LEWES/EiJ strains are derived from wild populations. By incorporating these wild-derived laboratory strains along with other inbred strains, a group of 10 mice can roughly represent the population of *M. m. domestics*. In addition, the low F_{IS} of 0.14 for rDNA in *M. m. domestics* suggests that each mouse covers 86% of population's genetic diversity, thus mitigating concerns about potential sampling biases. Accessions and the detailed information of samples used in this study are list in Supplementary Table S4 and Table S5.

Data Processing

After cutting adapters and filtering out of low-quality sequences, these whole-genome sequencing data of apes and mice were mapped against the human rDNA reference sequence (Human ribosomal DNA complete repeating unit, GenBank: U13369.1) and mouse rDNA reference sequence (*Mus musculus* ribosomal DNA, complete repeating unit, GenBank: BK000964.3), respectively, using Burrows-Wheeler-Alignment Tool v0.7.17 (Li and Durbin 2009) with default parameters. Each individual was considered as a pool of rRNA genes, so there were three levels of diversity. To determine the variant frequency within individual, we used bcftools (Danecek et al. 2021) to call variants from multiple alignment files and obtained per-sample allele depths with following settings 'bcftools mpileup --redo-BAQ --max-depth 50000 --per-sample-mF --annotate FORMAT/AD,FORMAT/ADF,FORMAT/ADR,FORMAT/DP,INFO/AD,INFO/ADF,INFO/ADR' and 'bcftools call -mv'. The sites with depth <10 were filtered out in the following analysis, and the average of the minimum depths for each remaining site across all samples is above 3000, which ensure a more robust estimation of the variant frequency. In this process, we can identity all the polymorphic sites present in these samples and the variant frequency within each individual. Then population variant frequency is calculated by the average frequency of variant across all individuals. In all cases, we only focus on single nucleotide variants (SNVs) and discard other types of mutations.

Identification of divergence sites.

Polymorphism is the transient phase of divergence. With variant frequency within individual, within species and between species, we obtained the F_{IS} and F_{ST} for every site. Among the *M. m. domestics* population, 779 polymorphic sites were identified, with 744 sites (95.5%) having an F_{IS} below 0.4 and rarely above 0.8 in Fig. 4B. Between *M. m. domesticus* and *Mus spretus*, 1579 variant sites were found while among them, 453 sites had an F_{ST} above 0.8, indicating the occurrence of rapid fixed mutations during species divergence. We also do the F_{ST} analysis between human and chimpanzee for each human individual and summarized the results in Table 5. We identified a range from 672 to 705 sites with F_{ST} values above 0.8, showing small variance across individuals. Due to the high mutation rate in CpG sites and predominantly GC content in rDNA, we further estimated the evolutionary rate at non-CpG sites during the comparison of the divergence between species. The mutations in CpG sites were manually removed by excluding all sites that contained CpG in one species and TpG or CpA in the other species; the reverse is also removed. We also counted the number of non-CpG sites within the mapping length where site depth exceeded 10 by Samtools (Danecek et al. 2021) with settings 'samtools mpileup -Q 15 -q 20'. Then we obtained the evolution rate of rDNA in non-CpG sites.

In the estimation of diversity and divergence in different parts of gene, we used 'samtools faidx' to partition variants into a total of 8 regions within rRNA genes, including 5'ETS, ITS1, ITS2, 3'ETS, IGS, 18S, 5.8S, and 28S, by aligning them to the corresponding reference sequences for further analysis. The functional parts (18S, 5.8S, and 28S) are subject to strong negative selection, resulting in minimal substitutions during species divergence as expected. This observation is primarily used for comparison with the non-functional parts and reflects that the non-functional parts are less constrained by negative selection.

Genome wide divergence estimation

To assess the genome-wide divergence between 4 mouse strain species, we downloaded their toplevel reference genomes from Ensembl genome browser (GenBank Assembly ID: GCA_001624865.1, GCA_001624775.1, GCA_001624445.1, and GCA_001624835.1). Then we used Mash tools (Ondov et al. 2016) to estimate divergence across the entire genome (mainly single-copy genes) with 'mash sketch' and 'mash dist'. The effective copy number of rRNA genes, C*, can be estimated by calculating the ratio of population heterozygosity observed in rDNA to that in single-copy genes.

Estimation of site conversion

By utilizing the identical variant found in chimpanzees and gorillas, humans and gorillas, *M. m. castaneus* and *M. spretus* as the ancestral state, we identified the derived variants that became nearly fixed in the lineages of human, chimpanzee and *M. m. domesticus* lineage with frequency > 0.8. In this study, we counted six types of mutations, from ancestral to derived as depicted in Supplementary Fig. S2. For example, A-to-G represented the both A-to-G and T-to-C types of mutations. The C-to-G (or G-to-C) and A-to-T (or T-to-A) types of mutations were excluded in the subsequent analysis. The alternative hypotheses of biased mutation pattern (Wolfe et al. 1989; Francino and Ochman 1999) alone can be rejected in this study. According to the prediction of the mutational mechanism hypotheses is that AT or GC variants should have equal fixation probability. We counted the nearly fixed (frequency > 0.8) number of AT-to-GC and GC-to-AT types of mutations and conducted a chi-square test to examine the fixation probabilities of AT and GC. As a result, we found the GC variant had significant higher fixation probability than AT at site level in apes but not in mice. Since the fixed sites accounted for less than 1% of non-functional length of rDNA, we calculated the AT-to-GC conversion rate, f, by dividing the number of fixed A or T sites by the number of GC base pairs, and the GC-to-AT conversion rate, g, can be calculated using the same approach. In order to test whether there is an equilibrium state at the genic level in these three species, we computed the (pf)/(qg) ratio in Table 7, and a significant deviation of the ratio from 1 can suggest biased genic conversion.

DATA ACCESS

We have no newly generated data.

COMPETING INTEREST STATEMENT

We declare no competing interest.

ACKNOWLEDGMENTS

We are grateful for the helpful comments from many colleagues on the Chat Group "Cancer - The new evolving species", in particular, Weiwei Zhai, Yong Zhang, GuoDong Wang of CAS and Jianrong Yang of SYSU.

REFERENCES

Alexandrov I, Kazakov A, Tumeneva I, Shepelev V, Yurov Y. 2001. Alpha-satellite DNA of primates: old and new families. *Chromosoma* **110**: 253-266.

Arnheim N, Krystal M, Schmickel R, Wilson G, Ryder O, Zimmer E. 1980. Molecular evidence for genetic exchanges among ribosomal genes on nonhomologous chromosomes in man and apes. *Proc Natl Acad Sci U S A* **77**: 7323-7327.

Arnheim N, Treco D, Taylor B, Eicher EM. 1982. Distribution of ribosomal gene length variants among mouse chromosomes. *Proc Natl Acad Sci U S A* **79**: 4677-4680.

Bowman JC, Petrov AS, Frenkel-Pinter M, Penev PI, Williams LD. 2020. Root of the Tree: The Significance, Evolution, and Origins of the Ribosome. *Chemical Reviews* **120**: 4848-4878.

Brown T, Jiricny J. 1989. Repair of base-base mismatches in simian and human cells. *Genome / National Research Council Canada = Génome / Conseil national de recherches Canada* **31**: 578-583.

Cabot EL, Doshi P, Wu ML, Wu CI. 1993. Population genetics of tandem repeats in centromeric heterochromatin: unequal crossing over and chromosomal divergence at the Responder locus of Drosophila melanogaster. *Genetics* **135**: 477-487.

Cazaux B, Catalan J, Veyrunes F, Douzery EJP, Britton-Davidian J. 2011. Are ribosomal DNA clusters rearrangement hotspots? A case study in the genus Mus (Rodentia, Muridae). *BMC Evolutionary Biology* **11**: 124.

Chen B, Wu X, Ruan Y, Zhang Y, Cai Q, Zapata L, Wu CI, Lan P, Wen H. 2022. Very large hidden genetic diversity in one single tumor: evidence for tumors-in-tumor. *Natl Sci Rev* **9**: nwac250.

Chen Y, Tong D, Wu CI. 2017. A New Formulation of Random Genetic Drift and Its Application to the Evolution of Cell Populations. *Mol Biol Evol* **34**: 2057-2064.

Cole F, Kauppi L, Lange J, Roig I, Wang R, Keeney S, Jasin M. 2012. Homeostatic control of recombination is implemented progressively in mouse meiosis. *Nat Cell Biol* **14**: 424-430.

Crow J, Kimura MJNY, US A. 1970. An Introduction to Population Genetics Theory Harper & Row.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM et al. 2021. Twelve years of SAMtools and BCFtools. *Gigascience* **10**.

Dover G. 1982. Molecular drive: a cohesive mode of species evolution. *Nature* **299**: 111-117.

Durrett R. 2013. POPULATION GENETICS OF NEUTRAL MUTATIONS IN EXPONENTIALLY GROWING CANCER CELL POPULATIONS. *The annals of applied probability : an official journal of the Institute of Mathematical Statistics* **23**: 230-250.

Eickbush TH, Eickbush DG. 2007. Finely orchestrated movements: evolution of the ribosomal RNA genes. *Genetics* **175**: 477-485.

Francino MP, Ochman H. 1999. Isochores result from mutation not selection. *Nature* **400**: 30-31.

Fujiwara K, Kawai Y, Takada T, Shiroishi T, Saitou N, Suzuki H, Osada N. 2022. Insights into Mus musculus Population Structure across Eurasia Revealed by Whole-Genome Analysis. *Genome Biol Evol* **14**.

Galtier N, Duret L. 2007. Adaptation or biased gene conversion? Extending the null hypothesis of molecular evolution. *Trends Genet* **23**: 273-277.

Geraldes A, Basset P, Gibson B, Smith KL, Harr B, Yu HT, Bulatova N, Ziv Y, Nachman MW. 2008. Inferring the history of speciation in house mice from autosomal, X-linked, Y-linked and mitochondrial genes. *Mol Ecol* **17**: 5349-5363.

Gibbons JG, Branco AT, Godinho SA, Yu S, Lemos B. 2015. Concerted copy number variation balances ribosomal DNA dosage in human and mouse genomes. doi:10.1073/pnas.1416878112 %J Proceedings of the National Academy of Sciences: 201416878.

Gonzalez IL, Sylvester JE. 2001. Human rDNA: Evolutionary Patterns within the Genes and Tandem Arrays Derived from Multiple Chromosomes. *Genomics* **73**: 255-263.

Guarracino A Buonaiuto S de Lima LG Potapova T Rhie A Koren S Rubinstein B Fischer C Abel HJ Antonacci-Fulton LL et al. 2023. Recombination between heterologous human acrocentric chromosomes. *Nature* **617**: 335-343.

Guillén AKZ, Hirai Y, Tanoue T, Hirai H. 2004. Transcription repression mechanisms of nucleolus organizer regions (NORs) in humans and chimpanzees. *Chromosome Research* **12**: 225-237.

Hartl DL, Clark AG, Clark AG. 1997. *Principles of population genetics*. Sinauer associates Sunderland.

Hori Y, Shimamoto A, Kobayashi T. 2021. The human ribosomal DNA array is composed of highly homogenized tandem clusters. *Genome Res* **31**: 1971-1982.

Jeffreys AJ, Neumann R. 2002. Reciprocal crossover asymmetry and meiotic drive in a human recombination hot spot. *Nature genetics* **31**: 267-271.

Kimura M. 1983. *The Neutral Theory of Molecular Evolution*. Cambridge University Press, Cambridge.

Kimura M, Crow JF. 1963. THE MEASUREMENT OF EFFECTIVE POPULATION NUMBER. *Evolution; international journal of organic evolution* **17**: 279-288.

Kimura M, Crow JF. 1964. THE NUMBER OF ALLELES THAT CAN BE MAINTAINED IN A FINITE POPULATION. *Genetics* **49**: 725-738.

Krystal M, D'Eustachio P, Ruddle FH, Arnheim N. 1981. Human nucleolus organizers on nonhomologous chromosomes can share the same ribosomal gene variants. *P Natl Acad Sci USA* **78**: 5744-5748.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges SB. 2022. TimeTree 5: An Expanded Resource for Species Divergence Times. *Mol Biol Evol* **39**.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)* **25**: 1754-1760.

Li W-H. 1997. *Molecular evolution*. Sinauer Associates, Sunderland, Mass.

Ling S, Hu Z, Yang Z, Yang F, Li Y, Lin P, Chen K, Dong L, Cao L, Tao Y et al. 2015. Extremely high genetic diversity in a single tumor points to prevalence of non-Darwinian cell evolution. *Proc Natl Acad Sci U S A* **112**: E6496-6505.

Lynch M. 2007. The frailty of adaptive hypotheses for the origins of organismal complexity. *Proc Natl Acad Sci U S A* **104 Suppl 1**: 8597-8604.

Lynch M, Conery JS. 2003. The Origins of Genome Complexity. *Science* **302**: 1401-1404.

Meunier J, Duret L. 2004. Recombination drives the evolution of GC-content in the human genome. *Molecular Biology and Evolution* **21**: 984-990.

Moran PAP. 1958. Random processes in genetics. *Mathematical Proceedings of the Cambridge Philosophical Society* **54**: 60-71.

Moran PAP, Oliphant MLE. 1997. A general theory of the distribution of gene frequencies - I. Overlapping generations. *Proceedings of the Royal Society of London Series B - Biological Sciences* **149**: 102-112.

Nagylaki T. 1983. Evolution of a large population under gene conversion. *Proc Natl Acad Sci U S A* **80**: 5941-5945.

Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger MR, Altemose N, Uralsky L, Gershman A et al. 2022. The complete sequence of a human genome. *Science* **376**: 44-53.

Ohta T. 1976. Simple model for treating evolution of multigene families. *Nature* **263**: 74-76.

Ohta T, Dover GA. 1983. Population genetics of multigene families that are dispersed into two or more chromosomes. *Proc Natl Acad Sci U S A* **80**: 4079-4083.

Ohta T, Gillespie JH. 1996. Development of Neutral and Nearly Neutral Theories. *Theoretical Population Biology* **49**: 128-142.

Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM. 2016. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol* **17**: 132.

Parks MM, Kurylo CM, Dass RA, Bojmar L, Lyden D, Vincent CT, Blanchard SC. 2018. Variant ribosomal RNA alleles are conserved and exhibit tissue-specific expression. *Sci Adv* **4**.

Phifer-Rixey M, Harr B, Hey J. 2020. Further resolution of the house mouse (Mus musculus) phylogeny by integration over isolation-with-migration histories. *BMC Evol Biol* **20**: 120.

Potapova T, Gerton J. 2019. Ribosomal DNA and the nucleolus in the context of genome organization. *Chromosome Research* **27**.

Ruan Y, Luo Z, Tang X, Li G, Wen H, He X, Lu X, Lu J, Wu CI. 2021a. On the founder effect in COVID-19 outbreaks: how many infected travelers may have started them all? *Natl Sci Rev* **8**: nwaa246.

Ruan Y, Wen H, He X, Wu CI. 2021b. A theoretical exploration of the origin and early evolution of a pandemic. *Sci Bull (Beijing)* **66**: 1022-1029.

Ruan Y, Wen H, Hou M, He Z, Lu X, Xue Y, He X, Zhang YP, Wu CI. 2022. The twin-beginnings of COVID-19 in Asia and Europe-one prevails quickly. *Natl Sci Rev* **9**: nwab223.

Ruan Y, Wen H, Hou M, Zhai W, Xu S, Lu X. 2023. On the epicenter of COVID-19 and the origin of the pandemic strain. *National*

Science Review **10**: nwac286.

Rudra M, Chatterjee B, Bahadur M. 2016. Phylogenetic relationship and time of divergence of Mus terricolor with reference to other Mus species. *Journal of genetics* **95**: 399-409.

Salim D, Gerton JL. 2019. Ribosomal DNA instability and genome adaptability. *Chromosome Research* **27**: 73-87.

Smirnov E, Chmúrčiaková N, Liška F, Bažantová P, Cmarko D. 2021. Variability of Human rDNA. *Cells-Basel* **10**.

Smith GP. 1974. Unequal Crossover and the Evolution of Multigene Families.   **38**: 507-513.

Stults DM, Killen MW, Pierce HH, Pierce AJ. 2008. Genomic architecture and inheritance of human ribosomal RNA gene clusters. *Genome Res* **18**: 13-18.

Szitenberg A, Cha S, Opperman CH, Bird DM, Blaxter ML, Lunt DH. 2016. Genetic Drift, Not Life History or RNAi, Determine Long-Term Evolution of Transposable Elements. *Genome Biol Evol* **8**: 2964-2978.

Tatsumoto S, Go Y, Fukuta K, Noguchi H, Hayakawa T, Tomonaga M, Hirai H, Matsuzawa T, Agata K, Fujiyama A. 2017. Direct estimation of de novo mutation rates in a chimpanzee parent-offspring trio by ultra-deep whole genome sequencing. *Sci Rep* **7**: 13561.

van Sluis M, Gailín M, McCarter JGW, Mangan H, Grob A, McStay B. 2019. Human NORs, comprising rDNA arrays and functionally conserved distal elements, are located within dynamic chromosomal regions. *Genes Dev* **33**: 1688-1701.

van Sluis M, McStay B. 2019. Nucleolar DNA Double-Strand Break Responses Underpinning rDNA Genomic Stability. *Trends Genet* **35**: 743-753.

Wilkinson GS, Adams DM, Haghani A, Lu AT, Zoller J, Breeze CE, Arnold BD, Ball HC, Carter GG, Cooper LN et al. 2021. DNA methylation predicts age and provides insight into exceptional longevity of bats. *Nature Communications* **12**: 1615.

Williams AL, Genovese G, Dyer T, Altemose N, Truax K, Jun G, Patterson N, Myers SR, Curran JE, Duggirala R et al. 2015. Non-crossover gene conversions show strong GC bias and unexpected clustering in humans. *Elife* **4**.

Wolfe KH, Sharp PM, Li WH. 1989. Mutation rates differ among regions of the mammalian genome. *Nature* **337**: 283-285.

Wu C-I, True JR, Johnson N. 1989. Fitness reduction associated with the deletion of a satellite DNA array. *Nature* **341**: 248-251.

Yang H, Wang JR, Didion JP, Buus RJ, Bell TA, Welsh CE, Bonhomme F, Yu AH-T, Nachman MW, Pialek J et al. 2011. Subspecific origin and haplotype diversity in the laboratory mouse. *Nature genetics* **43**: 648-655.

Zhao Z, Jin L, Fu Y-X, Ramsay M, Jenkins T, Leskinen E, Pamilo P, Trexler M, Patthy L, Jorde LB et al. 2000. Worldwide DNA sequence variation in a 10-kilobase noncoding region on human chromosome 22. *Proceedings of the National Academy of Sciences* **97**: 11354-11358.