The influence of prenatal alcohol exposure and maternal diet on offspring DNA methylation: a cross-species study

Mitchell Bestry[1], Alexander N. Larcombe[2,3], Nina Kresoje[4], Emily K Chivers[2], Chloe Bakker[2], James P Fitzpatrick[5], Elizabeth J Elliott[6,7], Jeffrey M Craig[8,9,10], Evelyne Muggli[9,10], Jane Halliday[10,11], Delyse Hutchinson[11,12,13,14], Sam Buckberry[4,15,16], Ryan Lister[15,16], Martyn Symons[1,17], David Martino[18].

1. *Telethon Kids Institute, The University of Western Australia, Perth, Australia.*

2. *Respiratory Environmental Health, Wal-yan Respiratory Research Centre, Telethon Kids Institute, Nedlands, Western Australia, Australia.*

3. *Occupation, Environment and Safety, School of Population Health, Curtin University, Perth, Western Australia.*

4. *Telethon Kids Institute, 15 Hospital Ave, Nedlands, Western Australia, Australia.*

5. *School of Psychological Sciences, University of Western Australia, Perth, Australia.*

6. *University of Sydney, Faculty of Medicine and Health, Specialty of Child and Adolescent Health, Sydney, NSW 2006, Australia.*

7. *Sydney Children's Hospitals Network (Westmead) and Kids Research, Sydney, NSW 2145, Australia.*

8. *Deakin University, IMPACT – the Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Geelong, Australia.*

9. *Murdoch Children's Research Institute, Royal Children's Hospital, Flemington Road, Parkville, Victoria 3052, Australia.*

10. *Department of Paediatrics, University of Melbourne, Royal Children's Hospital, Flemington Road, Parkville, Victoria 3052, Australia.*

11. *Public Health Genetics, Murdoch Children's Research Institute, Parkville, VIC,*

Australia.

12. *Deakin University, School of Psychology, Faculty of Health, Geelong, Australia.*

13. *Murdoch Children's Research Institute, Centre for Adolescent Health, Royal*

Children's Hospital, Melbourne, Australia.

14. *University New South Wales, National Drug and Alcohol Research Centre, Sydney,*

Australia.

15. *Harry Perkins Institute of Medical Research, QEII Medical Centre and Centre for*

Medical Research, The University of Western Australia, Perth, WA 6009, Australia.

16. *ARC Centre of Excellence in Plant Energy Biology, School of Molecular Sciences,*

The University of Western Australia, Perth, WA 6009, Australia.

17. *National Drug Research Institute, enAble Institute, Curtin University.*

18. *Wal-yan Respiratory Research Centre, Telethon Kids Institute, Nedlands, Western*

Australia, Australia.

Abstract

Prenatal alcohol exposure (PAE) is associated with changes to offspring DNA methylation

and adverse neurodevelopmental outcomes. Prior studies have utilised candidate gene or

microarray approaches resulting in biased representation of the methylome. We employed an

established experimental murine model of moderate PAE until gestational days 8-10, with

genome-wide DNA methylation sequencing of neonatal brain and liver. Although global

DNA methylation levels were generally preserved, we detected 78 alcohol-sensitive regions

in neonatal brain, and 759 alcohol-sensitive regions in neonatal liver ($p < 0.05$ and delta beta

> 0.05). Affected coding regions were significantly enriched in genes involved in

neurodevelopmental pathways, but behavioural outcomes were unaltered in adult littermates.

Seven alcohol-sensitive regions identified in mice were statistically replicated in a human

cohort of adolescents with fetal alcohol spectrum disorder (FASD) and known to be linked

with traits related to facial morphology, intelligence, educational attainment, autism,

schizophrenia, ageing and haematological disorders in the GWAS catalogue. The high methyl

donor diet (HMD) across pregnancy partially protected against the effects of alcohol on DNA

methylation in offspring tissues. Collectively this study provides a comprehensive evaluation

of PAE on offspring DNA methylation and suggests prenatal dietary intervention may be a

warranted mitigation strategy.

Introduction

Prenatal alcohol exposure (PAE) is the most common preventable cause of

neurodevelopmental impairments in children (1). Maternal alcohol consumed during

pregnancy passes through the placenta and acts as a teratogen in fetal tissues. PAE can cause

Fetal Alcohol Spectrum Disorder (FASD), a diagnostic term encompassing the spectrum of

physical, cognitive, behavioural and neurodevelopmental abnormalities that can occur in

offspring as a result of PAE (2, 3). Although PAE is necessary to cause FASD, outcomes

depend on multiple environmental and genetic factors including the dose and timing of

alcohol use during pregnancy, maternal diet, smoking, stress, and potentially other factors

that collectively influence fetal gene regulation and development (4-6).

DNA methylation is an epigenetic mechanism of gene regulation involving the

covalent modification of DNA cytosine bases, mostly within the cytosine-guanine sequence

(CpG) of genomic DNA. This is facilitated by a family of methyltransferase enzymes that

catalyse the addition of methyl groups to cytosines during one-carbon metabolism.

Methylation of DNA can alter chromatin density and influence patterns of gene expression in a tissue-specific and developmentally appropriate manner. Such modifications are mitotically stable throughout development. As DNA methylation is one of the control mechanisms for gene activity, alterations to this system by PAE could potentially disrupt fetal development and may be the cause of some of the difficulties experienced by people with FASD (7, 8). Previous studies on human participants (9-11) and animals (12-14) have implicated PAE as a mechanism of fetal DNA methylation disruption, either globally (11-13), and/or at specific gene regions (10, 13, 14) with limited replication of DNA methylation sites between studies (15). There has been much interest in the utility of DNA methylation as a potential biomarker for PAE, although results have been inconsistent (15). A mechanistic explanation for the effect of alcohol on DNA methylation is still lacking, but evidence from rodent (16) and human studies (17) suggests that PAE can either directly inhibit DNA methyltransferase enzymes or disrupt one-carbon metabolism via inhibition of bioavailability of dietary methyl donors, such as folate and choline to the fetus. Indeed several clinical trials have explored the potential for dietary choline to attenuate the cognitive deficits caused by PAE in affected individuals (18, 19), or when administered during pregnancy (20, 21), with results suggesting a high methyl donor diet (HMD) could mitigate the adverse effects of PAE on various behavioural outcomes.

There are many limitations when studying PAE in humans such as a reliance on observational methodologies and self-reported alcohol consumption. In addition, it is difficult to measure and control for multiple factors that may influence fetal DNA methylation regulation, including diet and smoking. Thus, we employed an established experimental model of PAE combined with validation studies in human cohorts. We adopted the design of a murine study used by Kaminen-Ahola et al. (22) who performed site-specific DNA methylation sequencing on the Agouti allele in C57BL/6J mice exposed to moderate PAE

during the first eight days of pregnancy. This model is largely based on early PAE during the

equivalent of the first trimester in humans when neurulation occurs and reflects a typical

situation in which women may consume alcohol until pregnancy recognition (23). We

employed whole genome bisulfite sequencing (WGBS) of fetal brain and liver to gain an

unbiased view of the methylation state of all CpG dinucleotides while factoring the effect of

tissue type on DNA methylation (24). Our WGBS approach also enabled sufficient

sequencing coverage in the mouse genome to examine for alcohol-sensitive regions and the

effects of a dietary intervention. We also assessed adult offspring mice for several

behavioural outcomes that are known to be affected by PAE to determine if early, moderate

PAE had any effects on these outcomes in adulthood, and if any of these effects were

mitigated by the dietary intervention. The aims of this study were to: (1) determine if

moderate PAE affected several behavioural outcomes in adult offspring mice; (2) identify

candidate DNA methylation biomarkers for moderate PAE during the first trimester of

pregnancy in a murine model; (3) investigate if HMD across pregnancy can mitigate any

effects of PAE on offspring DNA methylation or behavioural outcomes; and (4) validate

candidate biomarkers from the murine model within human cohorts for PAE or FASD.

Results

Comparison of prenatal characteristics across treatment groups

To investigate the effects of PAE and HMD on offspring DNA methylation and behavioural

outcomes, we employed a murine model with four treatment groups (Figure 1). The trajectory

of weight gain during pregnancy was similar across all treatment groups with some evidence

of more rapid weight gain in the HMD groups in the last 2 days (Linear mixed effects

regression model; H_2O-HMD: -2.282 ± 0.918 g, P = 0.0177; PAE-HMD: -1.656 ± 0.814 g, P

= 0.0493; figure 2a), although the total amount of weight gained between days 1 and 17-19

was not significantly different between treatment groups by linear mixed effects regression

(Figures S1a-b). The total amount of liquid consumed over the course of pregnancy was

significantly lower in HMD dams by unpaired t-test (Figures 2b and S1c). There was no

significant difference in the average litter size (6.525 ± 0.297 pups, figure 2c) and pup sex

ratios (Figure 2d) between treatment groups by unpaired t-test.

Effects of PAE and HMD on behavioural outcomes in adult mice

Behavioural testing was carried out on adult mice from each study group. There was no

evidence that PAE had a significant effect on any of behavioural outcomes tested (Figure S3).

Mice exposed to HMD exhibited greater locomotor activity, in terms of distance travelled

(Figure 3).

Effects of PAE on DNA methylation in newborn mice tissue.

Whole-genome bisulfite sequencing (WGBS) of brain and liver samples from 16 newborn

pups equally matched by treatment group and sex was performed to determine the effects of

PAE on fetal DNA methylation. Global levels of DNA methylation stratified across different

genomic features were preserved across treatment conditions, with no evidence of differences

in average methylation content (Figure S2).

To investigate region-specific effects of PAE on newborn DNA methylation, we conducted

genome-wide testing on newborn NC mice. Comparison of methylation profiles between

PAE and non-PAE groups within NC mice revealed 78 differentially methylated regions

(DMRs) in the brain from 19,113,529 CpG sites tested after coverage filtering and 759 DMRs

in the liver ($P < 0.05$ and delta > 0.05) from 19,335,801 CpG sites tested after coverage

filtering (Figure 4a-b). These regions were annotated to nearby genes using *annotatr* and are

provided in Tables S1-2. Two of the DMRs overlapped in mouse brain and liver (tissue

agnostic), but the remainder were tissue specific. Among these tissue agnostic regions was

the *Impact* gene on chromosome 18, which was hypomethylated in PAE+NC mice compared

to H_2O+NC mice in both the brain and liver (Figure 4c-d). The other tissue agnostic region

was within 5kb downstream of *Bmf*, and was hypermethylated in brain and liver tissue of

PAE+NC mice.

In both tissues, hypomethylation with PAE in NC mice was more frequently observed in

DMRs (52.6% of DMRs in brain, 93.5% of DMRs in liver hypomethylated with PAE)

compared to hypermethylation. The three genes associated with DMRs in both brain and the

liver tissues were the Autism Susceptibility Gene 2 (*Auts2*), Androglobin (*Adgb*), and RNA

Binding Protein Fox 1 (*Rbfox1*) genes (Table 1).

In both brain and liver tissues, DMRs were enriched in non-coding intergenic and open sea

regions and relatively underrepresented in coding and CpG island regions (Figure 4e-f).

Using open chromatin assay and histone modification datasets from the ENCODE project, we

found overwhelming enrichment ($p < 0.05$) of DMRs in open chromatin regions (ATAC-

seq), enhancer regions (H3K4me1), and active gene promoter regions (H3K27ac), in mouse

fetal forebrain tissue and fetal liver (Table 2).

Gene ontology enrichment analysis of liver DMRs that did localise to genes showed

enrichment in ten pathways, with neuron projection being the most significant (Figure 4g,

Tables S3-4).

HMD mitigates the effects of PAE on DNA methylation

To determine whether administration of a HMD throughout pregnancy could mitigate the

effects of PAE on offspring DNA methylation, we examined alcohol sensitive DMRs in the

HMD mice. Compared to H_2O+NC mice, PAE+HMD mice exhibited methylation differences

in only 12/78 (15%) brain (Table S7), and 124/759 (16%) liver (Table S8) DMRs, suggesting

the effects were predominantly mitigated. The overall difference in DNA methylation with

PAE within brain and liver DMRs in the HMD mice were significantly lower than in the NC

group mice (Figure 5a-d), with differences in DNA methylation with PAE more than 25%

lower in some DMRs in the HMD mice compared to NC mice (Figure 5e-f).

Replication studies in Human PAE and FASD case-control cohorts

We undertook validation studies in human cohorts to address the generalizability of our

murine model of PAE. Only 36 of the 78 (46.2%) brain DMRs, and 294 of the 759 (38.8%)

liver DMRs, had homologous regions in the human genome and were able to be tested.

In the Australian AQUA cohort, we tested a total of 1,898 CpG sites that corresponded to

mouse DMRs but did not find evidence of differential methylation (FDR < 0.05).

In the Lussier et al. cohort, differential testing of 2,048 CpG sites that were homologous to

mouse DMRs identified statistical evidence of 47 DMR associations with FASD status (FDR

P < 0.05). In models adjusted for participant age, sex, array number, and cell counts, 16/47

DMRs remained significant, seven of which had the same direction of effect observed in the

murine model (Table 3). Visual comparison of methylation changes across these seven

DMRs revealed striking differences in effect sizes between people with FASD and mice

(Figure 6). We investigated the list of genes implicated by association with the 7 replicated

DMRs using the GWAS and EWAS catalogue. Virtually all the genes of interest have been

associated with haematological traits (*GADD45A, KLF1, PEX12, AP2B1, DNASE2*) and

ageing phenotypes (*GADD45A, DNASE2, ZNF823, A2BP1*). GWAS studies have linked

variations in these genes with traits related to facial morphology (*GADD45A* (46)),

educational attainment (*AP2B1* (47)*,* intelligence (*RP9* (48)), autism and schizophrenia

(*ZNF823* (49)). The *Impact* gene, which contained a DMR in the brain and liver of newborn

mice, did not contain any DMRs in any of the human cohorts (Figure 7).

Candidate Gene Analysis of previously defined alcohol sensitive regions.

There were 21 CpG sites (FDR<0.05) identified in the brain from 15,132 CpG sites tested,

including two sites in the *Mest* (*Peg1*) gene and 19 sites in *Kcnq1* (*KvDMR1*) (Table S9).

There were nine FDR-significant CpG sites identified in the liver out of 15,382 CpG sites

tested, all of which were in *Peg3* (Table S10). All FDR-significant CpG sites identified in

both tissues were hypomethylated in mice with PAE.

Discussion

In this study we report on the effects in mice of maternal PAE on offspring DNA

methylation, using a whole-genome bisulfite sequencing approach, which provides a

comprehensive assessment of the mammalian DNA methylome. We found that moderate

early (first trimester) PAE was associated with site-specific differences to DNA methylation

in newborn pups but did not affect behavioural outcomes in adult mice. Global levels of DNA

methylation were not significantly different with PAE, and effects were characterized

predominantly by a loss of methylation (hypomethylation), mostly at non-coding regions of

the genome. Research evidence remains inconclusive about whether alcohol's effects on

DNA methylation are stochastic, or deterministic, with varying reports of loss or gain of

methylation in different tissues (15). We observed large differences in the effects of PAE on

DNA methylation between brain and liver, only detecting two genomic regions and four

genes that were similarly affected in both tissues from different germ layers. The latter

finding suggests alcohol can perturb methylation events as early in pregnancy as during

gastrulation, although on balance, our data were more supportive of a stochastic rather than a

deterministic influence of alcohol on DNA methylation.

Although our murine model did not result in any significant differences in behavioural

outcomes due to PAE, we identified site-specific differences in DNA methylation associated

with early moderate PAE. Timing, dose, and other tissue-related factors all likely influence

the extent to which genome-regulation is perturbed and manifests as differences in DNA

methylation. It is possible that more severe PAE might induce more consistent changes across

tissues, whereas sustained exposure would conceivably induce tissue-specific methylation

changes relating to the organ's metabolic processes.

We also found that DMRs were enriched in non-coding regions of the genome with

regulatory potential suggesting alcohol may have broad effects on genome regulation.

Despite this, the candidate gene analysis found reproducible associations in our model with

previously published reports of hypomethylation within *Peg3* and *KvDMR1* from South

African children with fetal alcohol syndrome (10). Both these genes are methylated in a

parent-of-origin specific manner, suggesting that alcohol affects imprinting processes,

although results from other studies are inconsistent (50, 51). We also replicated the alcohol-

sensitive DMRs we found in the murine model in a human cohort of adolescents with FASD

at genes with known associations with traits indicative of neurodevelopmental disorders

(Table 3). That these effects were detectable in different tissues despite species-related

differences is remarkable, and suggests that at least some methylation changes are stable.

Collectively, our data are encouraging for biomarker studies and aid in the prioritisation of

associations for future follow-up, particularly in relation to diagnosis of FASD. For example,

three genes are zinc finger proteins (*RP9*, *PEX12*, and *ZNF823*) that play an important role in

fetal gene regulation. Notably, *PEX12* is associated with Zellweger syndrome, which is a rare

peroxisome biogenesis disorder (the most severe variant of Peroxisome biogenesis disorder

spectrum), characterized by neuronal migration defects in the brain, dysmorphic craniofacial

features, profound hypotonia, neonatal seizures, and liver dysfunction (52).

Another key finding from this study was that HMD mitigated some of the effects of PAE on

DNA methylation. Preclinical studies of choline supplementation in rodent models have

similarly reported attenuation of memory and behavioural deficits associated with PAE (53,

54), and mitigating effects on DNA methylation (55). These data have been largely consistent

and collectively support the notion that alcohol induced perturbation of epigenetic regulation

may occur, at least in part, through disruption of the one-carbon metabolism. The most

encouraging aspect of this relates to the potential utility for evidence-informed

recommendations for dietary advice or supplementation, particularly in population groups

with limited access to antenatal care or healthy food choices.

This study provides a comprehensive assessment of the effects of PAE on DNA methylation

using a well-established murine model of early moderate exposure. Our study design also

utilized existing human data sets to explore the generalizability of the alcohol sensitive

DMRs we found, which were replicated at seven human genes. Despite this, the level of PAE

was insufficient to induce recognizable behavioural changes in adult littermates, and we did

not assess markers of brain or liver physiology. We therefore had limited ability to infer

pathogenic mechanisms from the methylation changes observed. We were also unable to

determine the contribution of specific cell types within tissues to the methylation changes

observed. In conclusion, we have identified novel targets with screening or diagnostic

potential that warrant further study. Additionally, our findings add to the body of evidence

that suggests a potential role for a dietary intervention to protect from the adverse effects of

PAE.

Materials and Methods

Murine subjects and housing

To study the effects of PAE on offspring DNA methylation processes, we adapted a murine

model study design that has previously reported epigenetic changes at the A^{vy} locus in Agouti

mice (22) (Figure 1). This study received animal ethics approval from the Telethon Kids

Institute Animal Ethics Committee (Approval Number: 344). Sixty nulliparous C57BL/6JArc

female mice aged ~8 weeks were mated with equivalent stud male mice. Pregnant dams were

randomly assigned to one of four treatment groups (n = 15 dams per group) that varied based

on composition of the drinking water and chow given to the dams:

i.   PAE-NC (Prenatal Alcohol Exposure-Normal Chow): 10% (v/v) ethanol in non-

acidified water *ad libitum* from 10 days prior to mating until gestational days (GD) 8-

10. This is intended to replicate moderate PAE during the first trimester of pregnancy

in humans. Dams received non-acidified reverse osmosis water for the remainder

pregnancy and NC (Rat and Mouse Cubes, Speciality Feeds, Glen Forrest, Australia)

throughout pregnancy.

ii.   PAE-HMD (Prenatal Alcohol Exposure-High Methyl Donor diet): 10% (v/v) ethanol

in non-acidified water *ad libitum* from 10 days prior to mating until GD8-10 and non-

acidified reverse osmosis water for remainder of pregnancy. Isocaloric high methyl

donor (HMD) chow consisting of 20 mg/kg folate and 4,970 mg/kg choline

throughout pregnancy (Speciality Feeds, Glen Forrest, Australia).

iii. H_20-NC (Water-Normal Chow): non-acidified water and NC throughout pregnancy.

iv. H_20-HMD (Water-High Methyl Donor diet): non-acidified water and HMD chow

throughout pregnancy.

Whole-genome bisulfite sequencing of newborn mouse tissues

Two male and two female pups from each treatment group (n=16 total) were euthanised by

intraperitoneal injection with ketamine and xylazine on the day of birth for WGBS of their

brain and liver tissues. Mouse tissue samples were stored at -80ºC. Pups were selected from

average sized litters for WGBS to minimize variability in PAE caused by differences in litter

sizes. Remaining littermates grew until adulthood for behavioural testing. Ten milligrams of

tissue were extracted from each liver and brain sample. Total nucleic acid was extracted from

the tissues using the Chemagic 360 instrument (PerkinElmer) and quantified with Qubit DNA

High Sensitivity Kit (Catalogue Number: Q32854, Thermo Scientific). 100 ng of genomic

DNA was spiked with 0.5 ng of unmethylated lambda DNA (Catalogue Number: D1521,

Promega) to assess the bisulfite conversion efficiency. Each sample was digested with 2 µl

RNase A (Invitrogen) at 37^{o}C for 20 minutes to remove RNA. 100ng of genomic DNA from

each sample was sheared using a Covaris M220 (300bp settings, Covaris). Libraries were

prepared using the Lucigen NxSeq AmpFREE Low DNA Library Kit (Catalogue Number:

14000-1, Lucigen), according to the manufacturer's instructions. Nextflex bisulfite-seq

barcodes (Catalgoue Number: Nova-511913, PerkinElmer) were used as the adapters with

incubation at 25^{o}C for 30 minutes. The libraries were bisulfite converted using the Zymo EZ

DNA Methylation-Gold Kit (Catalogue Number: D5005, Zymo Research) and PCR

amplified using the KAPA HiFi Uracil PCR Kit (Catalogue Number: ROC-07959052001,

Kapa Biosystems). The final libraries were assessed with the Agilent 2200 Tapestation

System using D1000 Kit (Catalogue Number:). WGBS was performed by Genomics WA

sequencing core on a NovaSeq 6000 (Illumina) using 2x150bp chemistry on an S4 flow cell.

The bisulfite conversion rate in each tissue sample was at least 99%. The overall mean

coverage in each sample was 9.69x (range: 6.51-12.12x). Raw fastq files were mapped to the

mm10 mouse reference genome with BSseeker 2 (version 2.1.8) (25) and CG-maptools

(version number 0.1.2) (26) using a custom bioinformatics pipeline.

Behavioural testing in adult mice

Littermates who were not sacrificed were reared on normal chow and drinking water *ad*

libitum until adulthood (~8 weeks after birth) when they underwent behavioural testing for

five neurodevelopmental functions including locomotor activity, anxiety, spatial recognition,

memory, motor coordination and balance. Behavioural tests included the open field test

(locomotor activity, anxiety) (27), object recognition test (locomotor activity, spatial

recognition) (28), object in place test (locomotor activity, spatial recognition) (29), elevated

plus maze test (locomotor activity, anxiety) (30), and rotarod test (motor coordination,

balance) (31).

Validation studies in human cohorts

Validation studies were undertaken in human cohorts with existing genome-wide DNA

methylation data sets and matching PAE data. These are described below:

1. The Asking Questions About Alcohol in Pregnancy (AQUA) cohort (32) is an

observational Australian cohort of 1,570 mother-child pairs. Timing, dose, and

frequency of alcohol consumption data were collected once in each trimester using

detailed questionnaires that were designed with consumer input to encourage honest

and accurate reporting (33). Genome-wide DNA methylation measurements were

available for 147 newborns from buccal swabs assayed on the

IlluminaHumanMethylation450k array. Newborns were classified according to the

level of maternal alcohol use during pregnancy as abstainers (51 subjects), moderate

discontinued (40 subjects), moderate sustained (49 subjects), or high sustained (7

subjects) (34).

2. The Lussier et al. cohort (9) is a cohort study for clinical diagnosis of FASD in

Canadian children. The cohort (GSE80261) initially contained 216 children (112

FASD and 104 controls) from the NeuroDevNet FASD cohort. Genome-wide DNA

methylation measures were taken from buccal swabs assayed on the

IlluminaHumanMethylation450k array. We performed initial quality filtering of the

data and included only Caucasians to avoid ethnic confounding (ethnicity chi-squared

$P = 8.28*10^{-14}$ before filtering). This reduced the sample size to 118 individuals (30

FASD and 88 controls).

3. The Cobben et al. cohort (35) is a cohort study for clinical diagnosis of FASD in

Canadian children. We assessed the combined discovery (GSE112987) and

replication (GSE113012) cohorts, which contained a total of 138 children (92 FASD

and 64 controls). Genome-wide DNA methylation measures were taken from blood

samples assayed on the IlluminaHumanMethylation450k array.

For the AQUA and Cobben et al. datasets, raw .iDAT files were pre-processed using the

minfi package (36) from the Bioconductor project (http://www.bioconductor.org) in the R

statistical environment (http://cran.r-project.org/, version 4.2.2). Sample quality was assessed

using control probes on the array. Between-array normalization was performed using the

stratified quantile method to correct for Type 1 and Type 2 probe bias. Probes exhibiting a *P*-

detection call rate of >0.01 in one or more samples were removed prior to analysis. Probes

containing SNPs at the single base extension site, or at the CpG assay site were removed, as

were probes measuring non-CpG loci (32,445 probes). Probes reported to have off-target

effects in McCartney et al. (37) were also removed. The pre-processed Lussier et al. cohort

dataset was obtained directly from the Gene Expression Omnibus database using the

getGenomicRatioSetFromGEO command from *minfi*.

Mouse DMRs were converted into human equivalent regions using an mm10 to hg19 genome

conversion with the liftover tool in the UCSC Genome Browser (38). A minimum 0.1 ratio of

bases that must remap was specified as recommended for liftover between regions from

different species and multiple output regions were allowed.

Statistical analysis

Dam characteristics and pup behavioural testing results were generally assessed using

unpaired t-tests comparing each treatment group to the baseline control group that was given

non-acidified reverse osmosis water and normal chow throughout pregnancy. The exceptions

were the trajectories of liquid consumption and weight gain across pregnancy, which were

assessed using a quadratic mixed effects model and the trajectory of chow consumption

across pregnancy which was assessed using a linear mixed effects model.

We converted CGmap files from each mouse sample into CG methylation bsseq objects, and

then combined the bsseq objects from each sample. We filtered the data to include only the

autosomes and then combined reads from mice in the same treatment group using the

collapseBSseq function, to maximise coverage prior to differential methylation analysis. The

X chromosome was excluded as we were searching for a PAE biomarker that was

independent of sex. CpG sites with an aggregated coverage below 10x in each tissue type

were removed prior to modelling to ensure there was sufficient coverage in all assessed CpG

sites. This retained 94.9% of CpG sites in the brain and 93.8% of CpG sites in the liver.

Differentially methylated regions (DMRs) were identified within tissues using a Bayesian

hierarchical model comparing exposed and unexposed groups using the Wald test with

smoothing, implemented in the R package *DSS* (39). We declared DMRs as those with P-

value < 0.05 based on the p-values of each individual CpG site in the DMR, and an effect

size (delta) > 0.05. For validation studies in humans, differential testing of mouse DMRs was

carried out using the R package *DMRcate* (40) for each dataset and DMRs were declared as

minimum smoothed false-discovery rate (FDR) < 0.05. Gene ontology analysis was carried

out using Gene Set Enrichment Analysis (41), which assessed brain and liver DMRs for

transcription start sites and associated biological processes.

Brain and liver DMRs were tested for enrichment within ENCODE Project data sets (42) by

an overlap permutation test with 100 permutations using the *regioneR* package. The

ENCODE Project data sets that were assessed included ENCFF845WSI, ENCFF764NTQ,

ENCFF937JHP, ENCFF269TLO, ENCFF676TSV and ENCFF290MLR. DMRs were then

tested for enrichment within specific genic and CpG regions of the mouse genome, compared

to a randomly generated set of regions in the mouse genome generated with *resampleRegions*

in *regioneR*, with equivalent means and standard deviations.

We compiled a set of key genes and genomic regions identified in previous mammalian PAE

studies for site-specific testing based on our prior systematic review of the literature, which

identified 37 candidate genes ((15), Tables S5-6). The brain and liver datasets were filtered to

candidate gene regions and differential testing was then performed across the entire coding

sequence, separately in the brain and liver of the NC mice using the *callDML* feature in DSS.

Competing Interest Statement

399 The authors declare no competing interests.

Acknowledgements

401 We wish to acknowledge the assistance of Dr Jahnvi Pflueger who provided training on

402 preparation of whole-genome bisulfite sequencing libraries. We wish to acknowledge the

403 financial contribution of the Centre for Research Excellence in FASD who supported the

404 murine experiments.

Author Contributions

DM, MS, AL, RL, DH, EM, JC, JH, JF, SB, MB, EE contributed to the study design and

funding application. EC, AL performed the mouse work including administering the mating,

treatments, measurements, monitoring and extracting biological samples. EC, AL, MB

performed the mouse behavioural testing. CB analysed the videos from the mouse

behavioural testing. MB prepared the whole-genome bisulfite sequencing libraries and

performed the data analyses. NK assisted with preparing whole-genome bisulfite sequencing

libraries. SB, DM provided advice and support on the data analysis. JH, EM contributed

human datasets for reproducibility analysis. MB, DM drafted the manuscript. MS, AL, RL,

DH, EM, JC, JH, JF, SB, EE, NK contributed to the development and editing of the

manuscript.

References

1.   Popova S, Charness ME, Burd L, Crawford A, Hoyme HE, Mukherjee RAS, et al. Fetal alcohol
spectrum disorders. Nat Rev Dis Primers. 2023;9(1):11.
2.   Bower C, Elliott EJ, Zimmet M, Doorey J, Wilkins A, Russell V, et al. Australian guide to the
diagnosis of foetal alcohol spectrum disorder: A summary. J Paediatr Child Health. 2017;53(10):1021-
3.
3.   Popova S, Lange S, Shield K, Mihic A, Chudley AE, Mukherjee RAS, et al. Comorbidity of fetal
alcohol spectrum disorder: a systematic review and meta-analysis. Lancet. 2016;387(10022):978-87.
4.   Jacobson JL, Akkaya-Hocagil T, Ryan LM, Dodge NC, Richardson GA, Olson HC, et al. Effects of
prenatal alcohol exposure on cognitive and behavioral development: Findings from a hierarchical
meta-analysis of data from six prospective longitudinal U.S. cohorts. Alcohol Clin Exp Res.
2021;45(10):2040-58.
5.   Sambo D, Goldman D. Genetic Influences on Fetal Alcohol Spectrum Disorder. Genes (Basel).
2023;14(1).
6.   Chung DD, Pinson MR, Bhenderu LS, Lai MS, Patel RA, Miranda RC. Toxic and Teratogenic
Effects of Prenatal Alcohol Exposure on Fetal Development, Adolescence, and Adulthood. Int J Mol
Sci. 2021;22(16).
7.   Jin Z, Liu Y. DNA methylation in human diseases. Genes Dis. 2018;5(1):1-8.
8.   Fransquet PD, Hutchinson D, Olsson CA, Wilson J, Allsop S, Najman J, et al. Perinatal
maternal alcohol consumption and methylation of the dopamine receptor DRD4 in the offspring: the
Triple B study. Environ Epigenet. 2016;2(4):dvw023.
9.   Lussier AA, Morin AM, MacIsaac JL, Salmon J, Weinberg J, Reynolds JN, et al. DNA
methylation as a predictor of fetal alcohol spectrum disorder. Clinical Epigenetics. 2018;10.
10.   Masemola ML, van der Merwe L, Lombard Z, Viljoen D, Ramsay M. Reduced DNA
methylation at the PEG3 DMR and KvDMR1 loci in children exposed to alcohol in utero: a South
African Fetal Alcohol Syndrome cohort study. Front Genet. 2015;6:85.
11.   Jarmasz JS, Basalah D, Del Bigio MR, Stirton H, Davie JR, Clarren SK, et al. Global DNA
Methylation and Histone Posttranslational Modifications in Human and Nonhuman Primate Brain in
Association with Prenatal Alcohol Exposure. Alcoholism: Clinical & Experimental Research.
2019;43(6):1145-62.
12.   Chen YY, Ozturk NC, Zhou FC. DNA Methylation Program in Developing Hippocampus and Its
Alteration by Alcohol. PLoS One. 2013;8(3).
13.   Abbott CW, Rohac DJ, Bottom RT, Patadia S, Huffman KJ. Prenatal Ethanol Exposure and
Neocortical Development: A Transgenerational Model of FASD. Cereb Cortex. 2018;28(8):2908-21.
14.   Downing C, Johnson TE, Larson C, Leakey TI, Siegfried RN, Rafferty TM, et al. Subtle
decreases in DNA methylation and gene expression at the mouse Igf2 locus following prenatal
alcohol exposure: effects of a methyl-supplemented diet. Alcohol. 2011;45(1):65-71.

15.   Bestry M, Symons M, Larcombe A, Muggli E, Craig JM, Hutchinson D, et al. Association of
prenatal alcohol exposure with offspring DNA methylation in mammals: a systematic review of the
evidence. Clin Epigenetics. 2022;14(1):12.
16.   Chen YL, Yang SS, Peng HC, Hsieh YC, Chen JR, Yang SC. Folate and vitamin B12 improved
alcohol-induced hyperhomocysteinemia in rats. Nutrition. 2011;27(10):1034-9.
17.   Hutson JR, Stade B, Lehotay DC, Collier CP, Kapur BM. Folic acid transport to the human
fetus is decreased in pregnancies with chronic alcohol exposure. PLoS One. 2012;7(5):e38057.
18.   Nguyen TT, Risbud RD, Mattson SN, Chambers CD, Thomas JD. Randomized, double-blind,
placebo-controlled clinical trial of choline supplementation in school-aged children with fetal alcohol
spectrum disorders. Am J Clin Nutr. 2016;104(6):1683-92.
19.   Wozniak JR, Fink BA, Fuglestad AJ, Eckerle JK, Boys CJ, Sandness KE, et al. Four-year follow-
up of a randomized controlled trial of choline for neurodevelopment in fetal alcohol spectrum
disorder. J Neurodev Disord. 2020;12(1):9.
20.   Jacobson SW, Carter RC, Molteno CD, Stanton ME, Herbert JS, Lindinger NM, et al. Efficacy of
Maternal Choline Supplementation During Pregnancy in Mitigating Adverse Effects of Prenatal
Alcohol Exposure on Growth and Cognitive Function: A Randomized, Double-Blind, Placebo-
Controlled Clinical Trial. Alcohol Clin Exp Res. 2018;42(7):1327-41.
21.   Thomas JD, Abou EJ, Dominguez HD. Prenatal choline supplementation mitigates the
adverse effects of prenatal alcohol exposure on development in rats. Neurotoxicol Teratol.
2009;31(5):303-11.
22.   Kaminen-Ahola N, Ahola A, Maga M, Mallitt KA, Fahey P, Cox TC, et al. Maternal Ethanol
Consumption Alters the Epigenotype and the Phenotype of Offspring in a Mouse Model. Plos
Genetics. 2010;6(1).
23.   Muggli E, O'Leary C, Donath S, Orsini F, Forster D, Anderson PJ, et al. "Did you ever drink
more?" A detailed description of pregnant women's drinking patterns. BMC Public Health.
2016;16:683.
24.   Loke YJ, Muggli E, Saffery R, Ryan J, Lewis S, Elliott EJ, et al. Sex- and tissue-specific effects of
binge-level prenatal alcohol consumption on DNA methylation at birth. Epigenomics.
2021;13(24):1921-38.
25.   Guo W, Fiziev P, Yan W, Cokus S, Sun X, Zhang MQ, et al. BS-Seeker2: a versatile aligning
pipeline for bisulfite sequencing data. BMC Genomics. 2013;14:774.
26.   Guo W, Zhu P, Pellegrini M, Zhang MQ, Wang X, Ni Z. CGmapTools improves the precision of
heterozygous SNV calls and supports allele-specific methylation detection and visualization in
bisulfite-sequencing data. Bioinformatics. 2018;34(3):381-7.
27.   Seibenhener ML, Wooten MC. Use of the Open Field Maze to measure locomotor and
anxiety-like behavior in mice. J Vis Exp. 2015(96):e52434.
28.   Lueptow LM. Novel Object Recognition Test for the Investigation of Learning and Memory in
Mice. J Vis Exp. 2017(126).
29.   Murai T, Okuda S, Tanaka T, Ohta H. Characteristics of object location memory in mice:
Behavioral and pharmacological studies. Physiol Behav. 2007;90(1):116-24.
30.   Komada M, Takao K, Miyakawa T. Elevated plus maze for mice. J Vis Exp. 2008(22).
31.   Deacon RM. Measuring motor coordination in mice. J Vis Exp. 2013(75):e2609.
32.   Muggli E, Halliday J, Elliott EJ, Penington A, Thompson D, Spittle AJ, et al. Cohort profile:
early school years follow-up of the Asking Questions about Alcohol in Pregnancy Longitudinal Study
in Melbourne, Australia (AQUA at 6). BMJ Open. 2022;12(1):e054706.
33.   Muggli E, Cook B, O'Leary C, Forster D, Halliday J. Increasing accurate self-report in surveys
of pregnancy alcohol use. Midwifery. 2015;31(3):e23-8.
34.   Muggli E, Hearps S, Halliday J, Elliott EJ, Penington A, Thompson DK, et al. A data driven
approach to identify trajectories of prenatal alcohol consumption in an Australian population-based
cohort of pregnant women. Sci Rep. 2022;12(1):4353.

35.   Cobben JM, Krzyzewska IM, Venema A, Mul AN, Polstra A, Postma AV, et al. DNA
methylation abundantly associates with fetal alcohol spectrum disorder and its subphenotypes.
Epigenomics. 2019;11(7):767-85.
36.   Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al. Minfi: a
flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation
microarrays. Bioinformatics. 2014;30(10):1363-9.
37.   McCartney DL, Walker RM, Morris SW, McIntosh AM, Porteous DJ, Evans KL. Identification of
polymorphic and off-target probe binding sites on the Illumina Infinium MethylationEPIC BeadChip.
Genom Data. 2016;9:22-4.
38.   Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, et al. The human genome
browser at UCSC. Genome Res. 2002;12(6):996-1006.
39.   Wu H, Xu T, Feng H, Chen L, Li B, Yao B, et al. Detection of differentially methylated regions
from whole-genome bisulfite sequencing data without replicates. Nucleic Acids Res.
2015;43(21):e141.
40.   Peters TJ, Buckley MJ, Chen Y, Smyth GK, Goodnow CC, Clark SJ. Calling differentially
methylated regions from whole genome bisulphite sequencing with DMRcate. Nucleic Acids Res.
2021;49(19):e109.
41.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.
Proc Natl Acad Sci U S A. 2005;102(43):15545-50.
42.   Luo Y, Hitz BC, Gabdank I, Hilton JA, Kagda MS, Lam B, et al. New developments on the
Encyclopedia of DNA Elements (ENCODE) data portal. Nucleic Acids Res. 2020;48(D1):D882-D9.
43.   Roffé M, Hajj GN, Azevedo HF, Alves VS, Castilho BA. IMPACT is a developmentally regulated
protein in neurons that opposes the eukaryotic initiation factor 2α kinase GCN2 in the modulation of
neurite outgrowth. J Biol Chem. 2013;288(15):10860-9.
44.   Yamashiro K, Hori K, Lai ESK, Aoki R, Shimaoka K, Arimura N, et al. AUTS2 Governs Cerebellar
Development, Purkinje Cell Maturation, Motor Function and Social Communication. iScience.
2020;23(12):101820.
45.   Fernàndez-Castillo N, Gan G, van Donkelaar MMJ, Vaht M, Weber H, Retz W, et al. RBFOX1,
encoding a splicing regulator, is a candidate gene for aggressive behavior. Eur
Neuropsychopharmacol. 2020;30:44-55.
46.   Indencleef K, Hoskens H, Lee MK, White JD, Liu C, Eller RJ, et al. The Intersection of the
Genetic Architectures of Orofacial Clefts and Normal Facial Variation. Front Genet. 2021;12:626403.
47.   Okbay A, Wu Y, Wang N, Jayashankar H, Bennett M, Nehzati SM, et al. Polygenic prediction
of educational attainment within and between families from genome-wide association analyses in 3
million individuals. Nat Genet. 2022;54(4):437-49.
48.   Davies G, Lam M, Harris SE, Trampush JW, Luciano M, Hill WD, et al. Author Correction:
Study of 300,486 individuals identifies 148 independent genetic loci influencing general cognitive
function. Nat Commun. 2019;10(1):2068.
49.   Consortium ASDWGoTPG. Meta-analysis of GWAS of over 16,000 individuals with autism
spectrum disorder highlights a novel locus at 10q24.32 and a significant overlap with schizophrenia.
Mol Autism. 2017;8:21.
50.   Marjonen H, Toivonen M, Lahti L, Kaminen-Ahola N. Early prenatal alcohol exposure alters
imprinted gene expression in placenta and embryo in a mouse model. PLoS One.
2018;13(5):e0197461.
51.   Stouder C, Somm E, Paoloni-Giacobino A. Prenatal exposure to ethanol: a specific effect on
the H19 gene in sperm. Reprod Toxicol. 2011;31(4):507-12.
52.   Konkoľová J, Petrovič R, Chandoga J, Halasová E, Jungová P, Böhmer D. A novel mutation in
the PEX12 gene causing a peroxisomal biogenesis disorder. Mol Biol Rep. 2015;42(9):1359-63.

53.   Thomas JD, Biane JS, O'Bryan KA, O'Neill TM, Dominguez HD. Choline supplementation
following third-trimester-equivalent alcohol exposure attenuates behavioral alterations in rats.
Behav Neurosci. 2007;121(1):120-30.
54.   Thomas JD, La Fiette MH, Quinn VR, Riley EP. Neonatal choline supplementation ameliorates
the effects of prenatal alcohol exposure on a discrimination learning task in rats. Neurotoxicol
Teratol. 2000;22(5):703-11.
55.   Otero NK, Thomas JD, Saski CA, Xia X, Kelly SJ. Choline supplementation and DNA
methylation in the hippocampus and prefrontal cortex of rats exposed to alcohol during
development. Alcohol Clin Exp Res. 2012;36(10):1701-9.

Table 1: Table of DMRs identified in the intronic regions of genes that contained DMRs in

both the brain and liver. Δmeth indicates the percentage change in average methylation level

within the DMR with PAE compared to non-PAE mice.

Gene	Tissue	Intronic DMR	Δmeth	P-value
Auts2	Brain	chr5:131510296-131510465	-23.5%	<0.05
Auts2	Liver	chr5:131621828-131621999	-22.5%	<0.05
Adgb	Brain	chr10:10455557-10455883	-25.0%	<0.05
Adgb	Liver	chr10:10353338-10353613	-25.9%	<0.05
Rbfox1	Brain	chr16:6813039-6813217	-24.3%	<0.05
Rbfox1	Liver	chr16:6781985-6782330	-22.6%	<0.05

574   Table 2: Number and percentage of brain and liver DMRs that overlap with tissue-specific

575   atacseq, H3K4me1 and H3K27ac regions obtained at 0 days postnatal from the ENCODE

576   database. P-values for permutation testing using a randomisation strategy.

Assay type	Brain DMRs	Brain randomised regions	Liver DMRs	Liver randomised regions
ATAC-seq	21/78 (26.92%) P = 0.01	1/78 (1.28%) P = 0.16	53/759 (6.98%) P = 0.01	22/759 (2.90%) P = 0.31
H3K4me1	4/78 (5.13%) P = 0.03	2/78 (2.56%) P = 0.18	38/759 (5.01%) P = 0.05	35/759 (4.61%) P = 0.32
H3K27ac	9/78 (11.54%) P = 0.01	2/78 (2.56%) P = 0.74	48/759 (6.32%) P = 0.01	19/759 (2.50%) P = 0.26

587 Table 3: DMRs identified in the murine model that were validated in the Lussier et al. human

588 case-control cohort for a clinical diagnosis of FASD. The upper section describes properties

589 of Lussier et al. human DMRs. The lower section describes properties of the equivalent

590 murine model DMRs.

DMR	organism	tissue	chr	start	end	width	no. cpgs	FDR	meandiff	gene
1	human	buccal	1	68151571	68152310	740	5	0.028636	-0.00497	*GADD45A*
2	human	buccal	19	13000782	13002357	1576	11	0.000197	-0.00203	*GCDH*
3	human	buccal	7	33148815	33149316	502	11	0.001149	-0.00011	*RP9*
4	human	buccal	17	33905444	33905888	445	14	0.000171	-0.00359	*AP2B1, PEX12*
5	human	buccal	17	27181503	27182342	840	11	0.018536	-0.00246	*ERAL1, FAM222B*
6	human	buccal	19	12992181	12992479	299	9	0.037431	-0.00179	*CTD-2265O21.7, DNASE2*
7	human	buccal	19	11849531	11850013	483	9	0.022724	-0.00244	*ZNF823*
1	mouse	liver	6	67034885	67035082	197	4	<0.05	-0.220833	*E230016M11Rik*
2	mouse	liver	8	84901298	84901544	246	5	<0.05	-0.234457	*Klf1*
3	mouse	liver	9	22453836	22453893	57	5	<0.05	-0.226427	*Rp9*
4	mouse	brain	14	21403570	21403622	52	4	<0.05	-0.234193	*Adk*
5	mouse	liver	11	78069463	78070002	539	9	<0.05	-0.255864	*Mir144, Mir451a*
6	mouse	liver	11	78072079	78072313	234	4	<0.05	-0.215227	*Mir144, Mir451a*
7	mouse	liver	2	177091927	177092945	1018	5	<0.05	-0.224354	Intergenic

Figure 1. Overview of PAE model.

A schematic representation of the experiment design is shown in figure 1. Fifteen dams were

allocated to each treatment group. Prenatal alcohol exposure (PAE) mice were exposed to

ethanol (10% v/v in non-acidified reverse osmosis drinking water *ad libitum*) from one week

before pregnancy to gestational days 8-10 and the remaining mice received water (H_2O). The

PAE and H_2O groups received either normal chow (NC) or a high methyl donor (HMD) diet

(NC containing 20 mg/kg folate and 4970 mg/kg choline) from one week before pregnancy

until birth.

Figure 2. PAE and HMD had varied effects on dam characteristics.

(a) Dam weight progression was significantly affected by HMD but not PAE by quadratic mixed-effects model without interaction (b) Trajectory of liquid consumption across pregnancy was affected by PAE and HMD by quadratic mixed effects model. PAE and HMD significantly interacted with trimester of pregnancy. (c) litter size and (d) pup sex ratios were not significantly associated with PAE or HMD by unpaired t-test or ANOVA. All line and bar plots show raw means and SEM. NC = normal chow, HMD = high methyl diet, PAE = prenatal alcohol exposure. Comparisons show p-value by unpaired t-test compared to the H_2O-NC baseline treatment group.

Figure 3. HMD was associated with increased locomotor activity.

HMD was associated with increased locomotor activity compared to NC, indicated by significantly greater total distance travelled in the (a) open field test (sex P = 0.0568), (b) object recognition test (sex P = 0.1519), (c) elevated plus maze test (sex P = 0.2043), and (d) object in place test (sex P = 0.1639) by unpaired t-test. Bars show mean and standard deviation. Each point represents one mouse. NC = normal chow, HMD = high methyl diet, PAE = prenatal alcohol exposure. Time interval for each mouse was (a-c) 300 seconds and (d) 180 seconds. The p-value (Sex P) represents the statistical significance of the association between sex and the behavioral outcome. It was calculated using a two-way ANOVA that included PAE status and HMD status as factors, without considering any interactions.

Figure 4. PAE was associated with site-specific differences in offspring DNA methylation.

The majority of DMRs were hypomethylated with PAE in (a) brain and (b) liver of mice given normal chow. Each point represents one DMR. Point colour indicates change in DNA methylation with PAE. PAE was also associated with hypomethylation in the DMRs identified in the promoter of the *Impact* gene in (c) brain and (d) liver, within NC mice. Each plot represents a separate treatment group. Each blue vertical line indicates a CpG site, with the height and corresponding left y-axis indicating the methylation ratio. The grey line and corresponding right y-axis indicate coverage at each CpG site. The black horizontal dotted line indicates 40% methylation for comparison purposes. The x-axis indicates the base position on chromosome 18, with the pink shaded area highlighting the DMR. DMR plots include 200 base pair flanking regions on each side of the DMR. DMRs identified in (e) brain and (f) liver were enriched in intergenic and inter-CpG regions, whilst being underrepresented in CpG and gene regions. The bar plot compares the number of WGBS DMRs in red to a set of equivalent randomly generated regions in blue. (g) Gene ontology analysis of liver DMRs shows enrichment within neuronal cellular components and biological processes. BP/red point = biological process, CC/blue point = Cellular component. X-axis of point indicates FDR of ontology. Size of point indicates number of overlapping genes with ontology. There were insufficient number of DMRs identified in the brain for a gene ontology analysis.

Figure 5. HMD partially mitigated effects of PAE on offspring DNA methylation.

Average DNA methylation effect sizes above 30% with PAE were observed in some (a) brain, and (b) liver DMRs in NC mice. Mean absolute difference in methylation with PAE is reduced within the HMD mice in (c) brain, and (d) liver. Each point represents one DMR. Point colour indicates change in DNA methylation with PAE. Points with a high number of CpGs and methylation difference are annotated with associated gene if located within a genic region. HMD was associated with (e) hypermethylation in the DMR identified proximal to *Lamb1* on chromosome 12 in brain and (f) hypomethylation in the DMR identified proximal to *Socs5* on chromosome 17 in liver. Each plot represents a separate treatment group. Each blue vertical line indicates a CpG site, with the height and corresponding left y-axis indicating the methylation ratio. The grey line and corresponding right y-axis indicate coverage at each CpG site. The black horizontal line indicates (e) 40% and (f) 80% methylation for comparison purposes. The x-axis indicates the base position on the chromosome, with the pink shaded area highlighting the DMR. DMR plots include 200 base pair flanking regions on each side of the DMR.

Figure 6. Seven PAE DMRs identified in the murine model were successfully replicated in the Lussier et al. human FASD cohort.

Examples of two PAE DMRs that were significantly hypomethylated with a clinical diagnosis of FASD in the Lussier et al. cohort (a and c), while their mouse liftover DMR was also significantly hypomethylated with PAE in the murine model experiment (b and d).

Figure 7. Methylation in the *Impact* gene and surrounding regions was not significantly affected by PAE or FASD in human cohorts.

The *Impact* gene, which contained DMRs significantly associated with PAE in the mouse brain and liver, was not significantly associated with (a) PAE in the AQuA cohort, or a clinical diagnosis of FASD in the (b) Cobben et al. cohort, or (c) Lussier et al. cohort.