Full title

Disease-attenuated pneumococcal biosynthesis gene mutants invade the mucosal

epithelium and induce innate immunity

Short title

Epithelial-pneumococcal interactions

Caroline M. Weight[1][*][±], Gabriele Pollara[1], Modupeh Betts[1], Roberta Ragazzini[2], Elisa Ramos-

Sevillano[3], Jesús Reiné[4], Matthew Whelan[1], José Afonso Guerra-Assunção[1], Michael Connor[5],

Paola Bonfanti[1,2], Clare Jolly[1], Mahdad Noursadeghi[1], Daniela M. Ferreira[4], Jeremy S. Brown[3],

Robert S. Heyderman[1][±]

[1]Research Department of Infection, Division of Infection and Immunity, UCL London, UK

[2]Francis Crick Institute, London, UK

[3]Department of Respiratory Medicine, UCL, London, UK

[4]Oxford Vaccine Group, University of Oxford, Oxford and Clinical Sciences, Liverpool School of

Tropical Medicine, Liverpool, UK

[5]Chromatin and Infection, Institut Pasteur, France

*Current address Biomedical and Life Sciences, Faculty of Health and Medicine, Lancaster

University, UK

±Corresponding Authors: Dr Caroline M. Weight and Professor Robert S. Heyderman

Email: c.weight@lancaster.ac.uk and r.heyderman@ucl.ac.uk

Word Count:

Introduction, results, discussion, methods = 3500

ABSTRACT

Nasopharyngeal colonisation by *Streptococcus pneumoniae* is characterised by bacterial adherence to epithelial cells, microinvasion and innate immune activation. Previously, we have shown two serotype 6B *S. pneumoniae* mutant strains affecting bacterial metabolism (Δ*proABC/pia* and Δ*fhs/pia*) colonise humans and mice, but in a murine disease model do not cause invasive infection. Here, we explore whether *S. pneumoniae* epithelial microinvasion and the induction of innate immune responses persist despite disease attenuation.

We show that under serum stress, these biosynthesis gene mutations had a broad but different impact on pneumococcal virulence gene expression, oxidative stress regulation, and purine and carbohydrate metabolism genes. However, although these mutations did not attenuate microinvasion in human challenge and epithelial models, there was less transmigration of Detroit 562 nasopharyngeal epithelial cells by the mutants compared to WT. Cellular reorganisation of primary human airway epithelium varied considerably between strains. Compared to WT, infection of Detroit 562 epithelial cells by the Δ*fhs/piaA* strain, but not the Δ*proABC/piaA* strain was less pro-inflammatory, induced less caspase 8 production, and were associated with increased pneumococcal hydrogen peroxide and reduced pneumolysin secretion.

These findings suggest that the observed differences in microinvasion and the epithelial response were driven by the differential expression of multiple bacterial virulence and metabolic pathways, rather than single genes or pathways of genes. These data highlight the complex impact of single gene mutations on bacterial virulence and suggest that the virulence determinants of pneumococcal epithelial colonisation, microinvasion and innate immunity are not necessarily directly linked to disease.

Author Summary

Streptococcus pneumoniae (the pneumococcus) commonly colonises the back of the human nose, and is a leading cause of pneumonia, meningitis, and sepsis. During colonisation, the pneumococcus adheres to the cells in the nose, invades these cells (so-called microinvasion), and activates them.

Colonisation is a pre-requisite for disease, however, since disease is largely a dead end for *S. pneumoniae*, it remains unclear whether these processes are directly linked to disease progression. We have previously shown that if we introduce gene mutations into *S. pneumoniae* that affect key metabolic pathways, these bacteria retain their ability to colonize human and animal models without causing disease. We now show that these mutants retain their ability to microinvade epithelial cells in human and mouse models, and some may still cause inflammation, but are less able to pass through the epithelial barrier. However, although the attenuation of disease may be explained by the broad-ranging impact of these mutations on pneumococcal virulence, oxidative stress, and metabolism, they are not driven by a single determinant. Our findings suggest that pneumococcal microinvasion and immune activation are not necessarily pre-cursors to disease progression. This supports the idea that *S. pneumoniae* adapts and evolves to promote colonisation and ultimately transmission rather than cause disease.

Keywords

Biosynthesis genes, host-pathogen interactions, Innate immunity, *Streptococcus pneumoniae*, airway epithelium, colonisation, disease

Graphical Abstract

S. pneumoniae colonisation is characterised by mucus association, epithelial adherence, microcolony formation and microinvasion – where the pneumococcus invades the epithelial barrier without causing disease. Although mutations in *S. pneumoniae* biosynthesis genes (Δ*proABC* and Δ*fhs*) attenuate disease in a murine model, they do not attenuate microinvasion in either experimental human pneumococcal challenge (EHPC), *ex vivo* or *in vitro* epithelial cells. Transmigration of the epithelial barrier is attenuated. These mutations show strain-dependent effects on both the epithelial and bacterial responses to infection. Factors such as epithelial cellular reorganisation, inflammation and caspase 8 activity alongside pneumococcal metabolic adaptation, virulence factor expression and response to stress are important components of these processes.

INTRODUCTION

Streptococcus pneumoniae is a common commensal of the URT, yet translocation of the pneumococcus to the lungs, blood or the brain leads to life-threatening disease [1]. Among children under five, *S. pneumoniae* is the leading cause of pneumonia-related deaths globally [2].

We have previously shown that *S. pneumoniae* colonisation is characterised by epithelial surface microcolony formation and invasion, termed microinvasion [3]. Pneumococcal colonization elicits an innate immune response that may facilitate clearance while also promoting bacterial transmission [3, 4]. Beyond the capsule, numerous virulence factors have evolved to optimise colonization and transmission, rather to cause invasive disease. We therefore hypothesize that epithelial microinvasion and the innate immune response to colonization are not necessarily linked to disease causation.

To test this, we have used two *S. pneumoniae* serotype 6B biosynthesis gene mutants (Δ*proABC* and Δ*fhs*), which were designed to induce protective immunity through nasopharyngeal colonization without causing disease [5]. The *proABC* operon encodes enzymes for proline biosynthesis, while *fhs* encodes a tetrahydrofolate reductase involved in one-carbon metabolism, both essential for pneumococcal growth and virulence [6-8]. In murine models, these mutants colonize the nasopharynx but fail to cause invasive disease [5, 8]. In EHPC, Δ*proABC* and Δ*fhs* mutants carrying an additional *piaA* deletion (disrupting a key iron uptake transporter, introduced to prevent reversion) retained the ability to colonise and elicit IgG-mediated immunity [9, 10], [10]. These isogenic mutants thus provide a controlled system to investigate epithelial microinvasion and innate immune activation independently of capsular serotype.

Here, we show that although the Δ*proABC/piaA* and Δ*fhs/piaA* mutations attenuate disease through broad-ranging impacts on pneumococcal virulence, oxidative stress, and metabolism, and reduced epithelial transmigration, they retained the capacity to colonize and invade epithelial cells. The Δ*fhs* but not the Δ*proABC* mutant was less pro-inflammatory. Hence pneumococcal microinvasion and immune activation are not necessarily pre-cursors to disease progression, highlighting the complex impact of biosynthesis gene mutations on pneumococcal virulence, with differential effects on epithelial colonisation and disease.

RESULTS

Deletion of the *proABC* and *fhs* genes have broad-ranging differential effects on the *S. pneumoniae* virulence gene expression and metabolic gene pathways when under stress

We have previously shown that Δfhs and ΔproABC showed considerable derangement in global gene transcription when cultured in human serum compared to growth under optimal nutrient conditions [8]. Metabolic analysis suggested that in sera, Δ*fhs* had an impaired stringent response, and both mutants were under increased oxidative stress and had altered lipid profiles [8]. The Δ*proABC* mutation resulted in the accumulation of glycolytic pathway and peptidoglycan synthesis intermediates. To further understand links between disease attenuation, epithelial colonisation and the innate immune response, we extended the transcriptomic analysis of the mutants' responses to serum stress [8]. This revealed upregulation of virulence factors such as *ply*, *nanA*, *psaA,* those regulating oxidative stress like *SpxB*, *lctO* and *adhE*, and genes involved in purine and carbohydrate metabolism (Supplemental Information Table 1). Phosphotransferase systems, amino nucleotide sugar, fructose and mannose and metabolic pathways were the most significantly enriched pathways in the Δ*proABC* mutant (Table 1). In contrast, competence, purine, pyruvate, metabolic pathways, amino nucleotide sugar pathways and secondary metabolites were the most significantly enriched pathways in the Δ*fhs* mutant under stress (Table 1).

Table 1. Differentially expressed pathways from single isogenic mutants in human serum compared to WT.

Pathway	GeneRatio	P-adjust	qvalue
Δ*fhs* vs. WT			
Fatty acid metabolism/biosynthesis	10/57	1.08E-09	6.49E-10
Biosynthesis of secondary metabolites	26/57	2.09E-04	1.26E-04
Purine metabolism	10/57	1.34E-03	8.06E-04
Quorum sensing	11/57	3.20E-03	1.92E-03
Pantothenate and CoA biosynthesis	5/57	3.59E-03	2.16E-03
beta-Lactam resistance	5/57	7.59E-03	4.57E-03
Propanoate metabolism	4/57	1.51E-02	9.09E-03
2-Oxocarboxylic acid metabolism	4/57	1.51E-02	9.09E-03
Metabolic pathways	40/57	1.94E-02	1.16E-02
Amino sugar and nucleotide sugar metabolism	7/57	4.22E-02	2.54E-02
Pyruvate metabolism	5/57	4.31E-02	2.59E-02
Δ*proABC* vs. WT			
Fatty acid metabolism/biosynthesis	10/71	1.17E-08	8.94E-09
Galactose metabolism	12/71	1.25E-04	2.47E-08
Phosphotransferase system (PTS)	13/71	2.54E-03	1.93E-03
Amino sugar and nucleotide sugar metabolism	9/71	2.89E-02	2.20E-02

Table 1. Differentially expressed pathways from single isogenic mutants in human serum compared to WT. 55 genes were selected within competence pathways, amino acid

metabolism, virulence factors, oxidative stress pathways and other known genes important

for colonisation, for assessment of changes in gene expression following serum stress. *q-*

value measures the proportion of false positives incurred (the false discovery rate) when

that particular test is significant.

***S. pneumoniae* 6B disease-attenuated mutants colonise and microinvade nasal epithelium in**

EHPC

We investigated whether the Δ*proABC/piaA* or Δ*fhs/piaA* mutations affected epithelial association

and microinvasion (Fig 1A) using nasal curette samples from EHPC of healthy volunteers [11].

Visualisation of mucosal cells by confocal microscopy showed colonisation by the WT and mutant

pneumococcal strains at day 6 post challenge (Fig 1B): 4/11 volunteers challenged with the WT,

5/11 volunteers with the Δ*proABC/piaA, and* 4/5 volunteers with the Δ*fhs/piaA*. Microinvasion

(intracellular bacteria) was seen in 2/11 volunteers with the WT, 5/11 volunteers with the

Δ*proABC/piaA,* and 3/5 volunteers with the Δ*fhs/piaA* (Fig 1C). Together, these data show that

compared to WT, there was no attenuation of colonisation or microinvasion with either the

Δ*proABC/piaA* or Δ*fhs/piaA* strains in EHPC.

Colonisation and microinvasion of human airway epithelium by the disease-attenuated

mutants is associated with cellular reorganisation and preservation of barrier function

To explore whether disease-attenuation influences epithelial cellular organisation, we infected

normal human bronchial epithelial airway (NHBE-A) cells with these pneumococcal strains. We

observed epithelial association and microinvasion with all three strains (Fig 2A, top). TEM

demonstrated that the intracellular bacteria for all three strains were likely enclosed within vacuoles

(Fig 2A, bottom). WT and mutant strains co-localised with Junctional Adhesion Molecule A (JAM-A,

Fig 2B, arrow), with no change in JAM-A expression. Quantification by image analysis demonstrated

differential effects on the expression of other junctional proteins with increased intensity of Zonula

Occludens 1 (ZO-1) at the junctional membrane after infection with WT (Fig 2C), and increased

intensity of β catenin staining infected with either Δ*proABC/piaA or Δfhs/piaA*, compared to cells exposed to these strains but not infected (Fig 2D), suggesting cellular reorganisation following pneumococcal exposure.

NHBE cells cultured at an air-liquid interface differentiate into a pseudostratified columnar ciliated epithelium, consisting of goblet, Clara, basal and ciliated cells (Fig 3A). Following infection, strains were observed amongst cilia (acetylated tubulin positive cells, Fig 3B, Supplemental Information Fig 1) and the Δ*proABC/piaA* co-localised with mucus (mucin5ac positive, Fig 3C, arrows). Infection with WT and Δ*proABC/piaA* but not Δ*fhs/piaA* resulted in an increased abundance of secreted mucus globules compared to uninfected cells (Fig 3A). This appeared due to secretion of preformed mucus rather than de-novo production, as expression of Mucin5AC was unaffected (Fig 3C). Infection with the Δ*proABC/piaA* strain increased expression of acetylated tubulin within pneumococcal-associated cells (Fig 3B), known to be associated with increased microtubule conformation and elasticity [12, 13]. In contrast, only infection with Δ*fhs/piaA* caused increased expression of uteroglobin, a multifunctional epithelial secreted protein with anti-inflammatory and anti-chemotactic properties, and a marker for non-ciliated Clara cells (Fig 3D).

Δ*fhs/piaA* but not the Δ*proABC/piaA* mutations attenuate the pro-inflammatory epithelial innate response to pneumococcal colonisation and microinvasion

We have shown that Detroit 562 epithelial nasopharyngeal cells are a good model for responses to *S. pneumoniae* [3]. Indeed, infection of Detroit 562 cells showed that WT, Δ*proABC/piaA* and Δ*fhs/piaA* strains formed colonies on the epithelial surface (Fig 4A), with limited differences between these strains (Fig 4B, top left). Microinvasion of Detroit 562 cells was observed with all strains (Fig 4B, top right) and showed no differences in intracellular viability (Fig 4B, bottom left) or replication (Supplemental Information Fig 2A-F). In line with attenuation of disease, the Δ*proABC/piaA* and Δ*fhs/piaA* were impaired in their ability to transmigrate across the epithelium (Fig 4B, bottom right). Importantly, the single Δp*iaA* mutation in the *S. pneumoniae* serotype 6B did not affect epithelial

association, microinvasion and transmigration, linking the differences seen to the Δ*fhs* and Δ*proABC*

mutations (Fig 4B). Epithelial β catenin expression was also affected following incubation with the

Δ*proABC/piaA* and Δ*fhs/piaA* strains, compared to the WT in pneumococcal-associated and non-

associated cells. (Fig 4C).

We further assessed links between colonisation, microinvasion and epithelial cell innate immunity

using RNAseq. Compared to non-infected cells (Fig 5A), all strains upregulated 78 core epithelial

genes related to cytokine signalling in immune system, NLR signalling and GPCR ligand binding.

193 genes were upregulated in response to WT and Δ*proABC/piaA*, characterised by Rhodopsin-

like receptors, GPCR ligand binding, cytokine signalling in Immune system, signalling by interleukins

and Toll Receptor cascades. Infection with Δ*fhs/piaA* resulted in a markedly different transcriptomic

profile with modest upregulation of 254 genes, characterised by antigen processing, proteasomal

pathways, GTP hydrolysis, and intracellular stress responses (Fig 5A). Overall, the epithelial genes

that were most upregulated following infection with WT or Δ*proABC/piaA* were similar and contrasted

to the reduced gene expression profile in response to Δ*fhs/piaA* despite high levels of cell invasion

(Fig 5B). Transcription factors, cellular kinases and cytokines (Fig 5C, 5D) predicted to regulate the

differentially expressed genes showed the greatest difference in profile following infection with

Δ*fhs/piaA* compared to infection with 6B WT and Δ*proABC/piaA*. Δ*fhs/piaA* was associated with

diminished NFκB and MAPK activation and instead induced a more intrinsic and epithelial repair

response that included TBK1, Raf1, JAK1 kinases and IL-1 and IL-33 cytokine signalling. These data

therefore suggest that attenuation of disease-causing potential is not dependent on a reduction in

the epithelial innate immune response to colonisation.

Enhanced epithelial caspase 8 activity following infection with wild-type and Δ*proABC/piaA*

strains

To further explore the link between disease attenuation and the epithelial innate immune response to colonisation, we assessed differential caspase activation. The NLRP3 inflammasome drives programmed epithelial cell death via caspase 1, caspase 3/7 and caspase 8 activity [14].

Compared to non-infected cells, all strains increased epithelial caspase activity 6-hour post infection, although this was only significant following infection with $\Delta proABC/piaA$ (Fig 6A, dark grey bars). Incubating infected cell cultures with either a caspase 1 inhibitor (YVAD-CHO) or caspase 3/7 inhibitor (Z-VAD-FMK), showed that increased caspase activation following infection with WT and $\Delta fhs/piaA$ was not due to caspase 1 activity (Fig 6B)Compared to non-infected cells, caspase 3/7 accounted for 3% and 15% of the caspase activity induced following infection with WT and $\Delta fhs/piaA$ respectively (Fig 6A). In contrast, $\Delta proABC/piaA$ strain did not induce caspase 3/7 activity. Caspase 8 activation was significantly increased following epithelial infection with 6B WT and $\Delta proABC/piaA$ strain (Fig 6C).

fhs gene mutation is associated with increased pneumococcal hydrogen peroxide secretion and decreased pneumolysin activity

S. pneumoniae SpxB and LctO generate hydrogen peroxide (H_2O_2) which suppresses ENaC-α transcription and inhibits the Na+–K+ pump, an intracellular signal to increase mitochondrial ROS production [15, 16]. We postulated that differences in cellular reorganisation, the innate inflammatory response and caspase 8 activation seen between $\Delta proABC/piaA$ and $\Delta fhs/piaA$ mutants could be explained by differential expression of pneumococcal H_2O_2. After three and six hours, we detected significantly higher levels of H_2O_2 in $\Delta fhs/piaA$ cultures compared to 6B WT and $\Delta proABC/piaA$ strains (Fig 7A). Furthermore, this was maintained during epithelial cell contact (Fig. 7B).

Pneumolysin can both activate and suppress the host innate immune response and mediates the haemolytic activity characteristic of the pneumococcus [17-19]. H_2O_2 increases the release of

pneumolysin but paradoxically also negatively impairs its haemolytic activity [20]. In line with this

paradoxical effect the higher production of H_2O_2 by the $\Delta fhs/piaA$ strain, was associated with

increased *ply* gene expression but reduced red blood cell lysis compared to 6B WT and

$\Delta proABC/PiaA$ (Fig. 7C).

DISCUSSION

We show that when using isogenic serotype 6B biosynthesis pathway mutant strains, attenuated for

systemic disease but not for carriage, epithelial microinvasion is preserved [5, 8, 21]. Epithelial

barrier transmigration was significantly less with the isogenic strains with the $\Delta fhs/piaA$ and

$\Delta proABC/piaA$ mutations compared to WT *S. pneumoniae* 6B. Importantly, while epithelial

microinvasion, cellular reorganization, the innate epithelial immune response, and the propensity to

cause invasive disease following pneumococcal colonisation are all strain-dependent, no single

determinant explained these differences.

In our primary epithelial cell infection model [22], consisting of Clara cells (involved in secretory,

xenobiotic metabolic, and progenitor cell functions), goblet cells (mucus production), ciliated cells

(small particle transport), infection with WT and mutant strains resulted in epithelial cellular re-

organisation involving β catenin, acetylated tubulin and uteroglobin. However, strain differences

seen on the impact on junctional or cytoskeletal protein organisation or mucus production did not

explain the disease attenuation seen with $\Delta fhs/piaA$ and $\Delta proABC/piaA$.

Epithelial infection with WT and $\Delta proABC/piaA$ induced innate inflammatory pathway gene

expression [3], including cytokine/chemokines, interleukins, Toll Like Receptor and NFκB signalling,

alongside apoptotic transcripts such as FOXO3 and TP53. Notably, some epithelial genes were

upregulated exclusively during WT infection, which could be related to its greater disease-causing

potential. These were involved in RNA degradation, O-linked glycosylation, wnt signalling, DNA

sensing, RIG-I pathways all of which play a role in the immune response and signal transduction. In

contrast, the Δ*fhs/piaA* strain activated a distinct pro-survival and adaptation profile, upregulating

genes such as JUN, SMARCA4, NFAT5, KLF6 and TCF4 which are involved in homeostatic

signalling and chromatin remodelling [23, 24], and suggesting a more tolerant epithelial response to

this strain. Ultimately, these capsule-independent differences in pneumococcal-epithelial

interactions may shape the differences seen in adaptive immune response identified using EHPC to

colonisation with the WT, Δ*proABC/piaA* or Δ*fhs/piaA* strains [11, 25].

Importantly, while all strains adhered and microinvade the epithelium, WT serotype 6B infection led

to pneumococcal transmigration, a more pro-inflammatory innate immune transcriptomic profile and

increased caspase 8 activity, indicative of a robust cellular response to *S. pneumoniae*, with

Δ*proABC/piaA* and Δ*fhs/piaA* exhibiting attenuation of these responses. This suggests that

pneumococcal sensing by the epithelium is affected by disruption of *S. pneumoniae* metabolism [18,

26]. Pneumolysin is a key driver of epithelial inflammatory responses and increased acetylated

tubulin and microtubule stabilisation [13] [3]. Under stress conditions, the Δ*proABC* and Δ*fhs*

mutations upregulated multiple metabolic pathways, presumably as a compensatory mechanism,

and also genes encoding several pneumococcal virulence factor genes (Graphical Abstract

Summary) including pneumolysin and H_2O_2. Based on these findings, we propose that differential

expression of pneumolysin, and H_2O_2 by the WT and mutant strains may contribute to the distinct

patterns of epithelial responses that we observed, with the potential to influence both bacterial

clearance and transmission [1]. Whether the distinct epithelial innate immune response to Δ*fhs/piaA*

infection is due to diminished NFκB and MAPK activation (potentially linked to extracellular receptor

engagement), reduced caspase 8 activity mediated by enhanced pneumococcal H_2O_2 release,

decreased pneumolysin activity, or a compensatory metabolic response to the stress, remains to be

determined. As seen in 'professional' intracellular bacterial species such as *Listeria* and *Salmonella*,

the host response to microinvasion might also promote pneumococcal transmission [27, 28].

Whether the less pro-inflammatory Δ*fhs/piaA* is less transmissible than the more inflammatory WT

is unknown, which could be explored using the EHPC model [29].

In conclusion, these data further support our proposed paradigm that epithelial microinvasion by *S.*

pneumoniae is frequent and does not necessarily lead to disease. Microinvasion may aid

transmission through immune evasion and localised inflammation and, transmigration across the

epithelium may represent the first step towards transition to invasive disease. Our findings also

highlight the broad-ranging effects of single gene mutations in pneumococcal biosynthesis

pathways, which may explain the complexity of differences between seemingly closely related

strains. The interplay between *S. pneumoniae* and the host epithelium is modulated by key mediators

such as hydrogen peroxide and pneumolysin, which can have both enhancing and inhibitory effects

on epithelial immune responses depending on the infection context. However, strain differences in

subsequent epithelial cell microinvasion, cellular reorganisation and inflammatory responses appear

driven by the differential expression of multiple bacterial virulence and metabolic pathways, rather

than any single determinant.

METHODS AND PROTOCOLS

Bacteria

The *Streptococcus pneumoniae* serotype 6B (strain BHN 418 [30]) was a kind gift from Prof. Birgitta

Henriques Normark (Karolinska Institute). Δ*proABC/piaA* and Δ*fhs/piaA* mutant strains were

generated using overlap extension PCR as described by Ramos-Sevillano *et al* [5]. Aliquot stocks

frozen at 0.3 O.D.$_{600nm}$ were defrosted, centrifuged at 8000g for eight minutes and resuspended in

1% Foetal Bovine Serum (FBS, Gibco) in MEM media (Gibco) for infections. Starting inoculums were

not significantly different across strains for all assays.

Bacterial RNA-seq sample collection, sequencing and reads

Pneumococcal strains were grown in THY to mid-log phase and pellets resuspended in THY and undiluted human serum for 60mins. The culture was centrifuged and stored in RNA Protect at -70°C before extracting total RNA [8]. Total RNA from each library was ribo-depleted and sent for Illumina Next Seq Sequencing Pathogen Genomics Unit, UCL, as detailed in Supplemental Information.

Differential gene expression analyses

The generated count matrix was imported into R-studio (R v3.4.2) and normalization of counts and differential gene expression analysis was performed by DESeq2 [31]. DESeq2 normalized libraries were regularized log transformed for visualizing heatmaps and clustering. Differential gene expression was performed on raw counts. Fold change shrinkage was implemented with the apeglm package within DESeq2. Genes with a log2 fold change > 1.5 and false discovery rate (FDR) of < 0.05 were categorized as differentially expressed. KEGG pathway enrichment and module analysis were performed on differentially regulated genes using over representation analysis (ORA) implemented by clusterProfiler [32].

Experimental Human Pneumococcal Carriage Model (EHPC)

A full explanation of experimental design and selection criteria has been previously described [3] and detailed in Supplemental Information.

Normal Human Epithelial Cells

Normal Human Bronchial/Tracheal Epithelial Cells (NHBE-A, ATCC® PCS-300-010™) were plated at 6×10^3 cells/cm^2 on a layer of sub-lethally γ-irradiated (60 Gy) mouse embryonic fibroblasts 3T3-J2 cells (kind donation from Howard Green, Harvard Medical School, Boston [33]) with keratinocyte culture medium cFAD (3:1 DMEM (Gibco) to Ham F-12 Nut Mix (Gibco), 10% FBS (HyClone, Sigma), 1% Penicillin-Streptomycin (100X, Sigma), 0.4 μg/mL Hydrocortisone (Calbiochem, USA), 5 μg/ml Insulin (various), 10-10 Cholera Toxin (Sigma) and 2x10-9 Triodothyronine (Sigma)). Epithelial cells

were stimulated with 10 ng/mL human epidermal growth factor (hEGF, PeproTech, USA) on days 3

and 5. Cultures were grown at 37°C 6% CO_2. For NHBE-A differentiation in Air Liquid Interface

culture, cells were plated at $6x10^4/cm^2$ on 1µm pore PET cell culture inserts (Greiner). All chambers

were supplemented with cFAD medium. After 48 hours, medium was removed from the top chamber.

Lower chamber media was replaced with PneumaCult™ differentiation Medium (Stemcell

Technologies) every 2 days for 3 weeks. Antibiotics were removed from media and cells were

washed 24 hours before infection experiments.

Human Epithelial Cell Lines

Human pharyngeal carcinoma Detroit 562 epithelial cells (ATCC_CCL-138) were grown in 10% FCS

in alpha MEM media (Gibco). For each set of experiments, cells were used within 5 passages. Cells

consistently tested PCR negative for mycoplasma (Cambridge Biosciences).

Pneumococcal-epithelial cell co-culture

For association and internalisation assays, confluent monolayers of Detroit 562 cells cultured on 12

well plates (Corning), were infected(1 cell to 10 pneumococci) for three hours in 1% FCS MEM or

six hours for HNBE-A cells in 1% Pneumocult media without supplements [3]. Cells washed three

times in $HBSS^{+/+}$ before incubation in 1% saponin for 10 minutes at 37°C. Bacterial dilutions were

plated on horse blood agar for 16 hours and colony forming units counted. For internalisation

quantification, 200µg/ml gentamicin was added one hour before incubation with saponin.

For transmigration assays, confluent monolayers of Detroit 562 cells cultured on 3µm pore, PET

Transwell Inserts (ThermoFisher) [3] were infected for three hours. 50µl of basal media was

repeatedly removed to measure pneumococcal load by counting CFU/well.

Confocal Microscopy

Epithelial monolayers on transwell membranes were infected then fixed in 4% PFA (Pierce, Methanol

Free) as previously described and detailed in Supplemental Information [3]. Cells were processed

for image analysis using an inverted Zeiss LSM (either 700 or 880). Z stacks were recorded at 1μm

intervals at either 40x oil or 63x oil objectives. Automated image analysis was carried out using

Cellprofiler [34].

Electron Microscopy

Preparation of Detroit 562 cells and NHBE-A cells was performed as previously described and details

can be found in Supplemental Information [3].

Pneumolysin activity

Adapted from Kirkham *et al* [35]. Final concentration of 2% horse blood (EO Labs) solution was

prepared in phenol free alpha MEM (Life Technologies) and added to a 96 U-bottom plate (Corning).

2.5×10^6 bacteria were added to 96 well. Saponin was used as a positive control for lysis. Solutions

were incubated for 30 minutes at 37°C before centrifugation at 1000g for one minute. Solution was

transferred into a new 96 well plate and absorbance recorded at 540nm on a plate reader.

Hydrogen Peroxide Assay

Detroit 562 cells were cultured in white 96 well plates (Corning). Cells were incubated with H_2O_2

substrate and infected for up to six hours according to manufacturers' instructions (ROS-Glo™ H_2O_2

Assay, Promega). Luminescence was read using a GloMax -Multi Detection System plate reader

(Promega).

Caspase assay

Detroit 562 cells were cultured in white 96 well plates (Corning). Cells were infected for up to six

hours. Primary cell supernatant was transferred to a white 96 well plate. Samples were incubated

with Caspase-Glo® 1 Reagent, Caspase-Glo® 1 YVAD-CHO Reagent, Caspase-Glo® Z-VAD-FMK

Reagent or Caspase-Glo® 8 Reagent, for one hour and according to manufacturers' instructions

(Promega). Luminescence was read using a GloMax-Multi Detection System plate reader

(Promega).

RNA samples and sequencing (RNA-Seq)

Confluent monolayers of Detroit 562 epithelial cells were infected for four hours before collection in RNALater. RNA was extracted using a Qiagen RNAEasy micro kit and treated for DNA removal using a Qiagen Turbo DNA-free kit. RNA quality was assessed and quantified using a BioAnalyser (Agilent 2100). Samples were sequenced using Illumina Nextseq 500/550 High Output 75 cycle kit giving 15-20 million 41bp paired-end reads per sample (Pathogens Genomic Unit, UCL).

Statistics

Conditions within each experiment were performed in duplicate or triplicate, and each experiment was performed independently more than three times, unless stated otherwise. Error bars represent SEM, unless stated otherwise. GraphPad Prism Version 7 was used for parametric (t-tests or ANOVA) or non-parametric (Mann-Whitney or Kruskal-Wallis tests) analysis, which was based on the Shapiro-Wilk normality test. Ad hoc tests were performed using Tukey's (parametric data) or Dunn's (non-parametric data) multiple comparisons test. P values <0.05 were considered significant.

Data and code availability

Source data are provided as a Source Data file. RNAseq data from the Detroit 562 cell infections is publicly accessible through Zenodo with Accession (DOI:10.5281/zenodo.7997789).

ACKNOWLEDGEMENTS

This study was funded by CMW from a Postdoctoral Innovation Award from UCL, a Pump-Priming Award and Enhancement Award from the Human Infection Challenge Network for Vaccine Development (HIC-VAC (funded by the GCRF Networks in Vaccines Research and Development) which is co-funded by the MRC and BBSRC). This UK funded award is part of the EDCTP2 programme supported by the European Union. CMW and RSH are supported by the Medical

Research Council (grant MR/T016329/1). The authors would like to thank Prof. Peter Hermans

(Radboud University Medical Centre) for donating the 6B pneumococcal strain originally from Prof.

Birgitta Henriques-Normark (Karolinska Institute) and obtained directly from DF at LSTM. EHPC

sample collection was supported by the Medical Research Council (grant MR/M011569/1), Bill and

Melinda Gates Foundation (grant OPP1117728) awarded to DF and the National Institute for Health

Research (NIHR) Local Comprehensive Research Network. The authors also wish to thank the

EHPC Clinical Team at LSTM and all the volunteers. Confocal imaging facilities at LSTM were

funded by a Wellcome Trust Multi-User Equipment Grant (104936/Z/14/Z). Electron microscopy

processing and image capture was performed by Dr. Elizabeth Slavik-Smith in the Biosciences EM

facility at UCL. MC is supported by a Springboard to Independence grant (AirwayStasis) from the

French Government's Investissement d'Avenir program, the Laboratoire d'Excellence "Integrative

Biology of Emerging Infectious Diseases" (ANR-10-LABX-62-IBEID) and the Chromatin and infection

unit headed by Melanie A. Hamon. RNA-Seq library preparation was undertaken at UCL through the

UCL/UCLH Biomedical Research Centre and MRC funded Pathogen Genomics Unit (PGU,

G0900950). All RNAseq data processing was completed by the PGU. MN is supported by the

Wellcome Trust and by NIHR Biomedical Research Centre Funding to University College Hospitals

NHS Foundation Trust and University College London. RR is supported by Marie Skłodowska-Curie

Individual Fellowships 896014. PB is supported by the European Research Council (ERC-Stg No.

639429), the Rosetrees Trust (M362-F1; M553), the CF Trust (SRC006;SRC020), the London

Advanced Therapies - Research England (C2N-AT.006), the MRC Confidence in Concept scheme

(Award: MC_PC_17180), the Innovate UK (Smart grant n.10005465), the Duchenne Parent Project

(DPP) and the NIHR GOSH BRC. MW is supported by a 223065 Wellcome Investigator Award to

Clare Jolly, UCL. ERS and JSB are supported by MRC grant R/N02687X/1 and Wellcome grant

221803/Z/20/Z. JSB, MN, and RSH acknowledge funding from the Department of Health's National

Institute of Health and Care Research Biomedical Research Centre funding to UCL and UCLH. RSH

is a NIHR Senior Investigator.

AUTHOR CONTRIBUTIONS: CMW conceived, funded, designed and conducted the study, performed the experiments, acquired samples and processed data, and wrote the first draft of the manuscript. GP analysed and interpreted the RNAseq data. MB analysed and interpreted the bacterial transcriptomic data. RR cultured and differentiated primary epithelial cells. ERS generated the 6B mutant strains and bacterial transcriptomic data. JR collected and processed EHPC samples. JAGA processed the RNA-Seq data. MW ran the image analysis. MC provided technical advice. PB and CJ provided investigator support. MN provided RNA-Seq analysis support. DF provided investigator support and EHPC clinical samples. JSB generated the 6B mutant strains, interpreted the data and revised the manuscript. RSH provided investigator support, designed and interpreted the data, revised the manuscript. All authors commented on and approved the manuscript.

COMPETING INTERESTS: The authors declare no competing interests.

REFERENCES

1. Weiser JN, Ferreira DM, Paton JC. Streptococcus pneumoniae: transmission, colonization and invasion. Nat Rev Microbiol **2018**; 16:355-67.

2. O'Brien KL, Wolfson LJ, Watt JP, et al. Burden of disease caused by Streptococcus pneumoniae in children younger than 5 years: global estimates. Lancet **2009**; 374:893-902.

3. Weight CM, Venturini C, Pojar S, et al. Microinvasion by Streptococcus pneumoniae induces epithelial innate immunity during colonisation at the human mucosal surface. Nat Commun **2019**; 10:3060.

4. Zafar MA, Hamaguchi S, Zangari T, Cammer M, Weiser JN. Capsule Type and Amount Affect Shedding and Transmission of Streptococcus pneumoniae. MBio **2017**; 8:e00989-17.

5. Ramos-Sevillano E, Ercoli G, Felgner P, et al. Preclinical Development of Virulence-attenuated Streptococcus pneumoniae Strains Able to Enhance Protective Immunity against Pneumococcal Infection. Am J Respir Crit Care Med **2021**; 203:1037-41.

6. Brill J, Hoffmann T, Bleisteiner M, Bremer E. Osmotically controlled synthesis of the compatible solute proline is critical for cellular defense of Bacillus subtilis against high osmolarity. J Bacteriol **2011**; 193:5335-46.

7. Crowley PJ, Gutierrez JA, Hillman JD, Bleiweis AS. Genetic and physiologic analysis of a formyl-tetrahydrofolate synthetase mutant of Streptococcus mutans. J Bacteriol **1997**; 179:1563-72.

8. Ramos-Sevillano E, Ercoli G, Betts M, et al. Essential role of proline synthesis and the one-carbon metabolism pathways for systemic virulence of Streptococcus pneumoniae. mBio **2024**; 15:e0175824.

9. Brown JS, Gilliland SM, Holden DW. A Streptococcus pneumoniae pathogenicity island encoding an ABC transporter involved in iron uptake and virulence. Mol Microbiol **2001**; 40:572-85.

10. Hill H, Mitsi E, Nikolaou E, et al. A Randomized Controlled Clinical Trial of Nasal Immunization with Live Virulence Attenuated Streptococcus pneumoniae Strains Using Human Infection Challenge. Am J Respir Crit Care Med **2023**; 208:868-78.

11. Ferreira DM, Neill DR, Bangert M, et al. Controlled human infection and rechallenge with Streptococcus pneumoniae reveals the protective efficacy of carriage in healthy adults. Am J Respir Crit Care Med **2013**; 187:855-64.

12. Eshun-Wilson L, Zhang R, Portran D, et al. Effects of alpha-tubulin acetylation on microtubule structure and stability. Proc Natl Acad Sci U S A **2019**; 116:10366-71.

13. Iliev AI, Djannatian JR, Opazo F, et al. Rapid microtubule bundling and stabilization by the Streptococcus pneumoniae neurotoxin pneumolysin in a cholesterol-dependent, non-lytic and Src-kinase dependent manner inhibits intracellular trafficking. Mol Microbiol **2009**; 71:461-77.

14. Surabhi S, Jachmann LH, Shumba P, Burchhardt G, Hammerschmidt S, Siemens N. Hydrogen Peroxide Is Crucial for NLRP3 Inflammasome-Mediated IL-1beta Production and Cell Death in Pneumococcal Infections of Bronchial Epithelial Cells. J Innate Immun **2022**; 14:192-206.

15. Erttmann SF, Gekara NO. Hydrogen peroxide release by bacteria suppresses inflammasome-dependent innate immunity. Nat Commun **2019**; 10:3493.

16. Alibayov B, Scasny A, Khan F, et al. Oxidative Reactions Catalyzed by Hydrogen Peroxide Produced by Streptococcus pneumoniae and Other Streptococci Cause the Release and Degradation of Heme from Hemoglobin. Infect Immun **2022**; 90:e0047122.

17. Zafar MA, Wang Y, Hamaguchi S, Weiser JN. Host-to-Host Transmission of Streptococcus pneumoniae Is Driven by Its Inflammatory Toxin, Pneumolysin. Cell Host Microbe **2017**; 21:73-83.

18. Periselneris J, Turner CT, Ercoli G, et al. Pneumolysin suppresses the initial macrophage pro-inflammatory response to Streptococcus pneumoniae. Immunology **2022**; 167:413-27.

19. Terra VS, Plumptre CD, Wall EC, Brown JS, Wren BW. Construction of a pneumolysin deficient mutant in streptococcus pneumoniae serotype 1 strain 519/43 and phenotypic characterisation. Microb Pathog **2020**; 141:103999.

20. Bazant J, Ott B, Hudel M, Hain T, Lucas R, Mraheil MA. Impact of Endogenous Pneumococcal Hydrogen Peroxide on the Activity and Release of Pneumolysin. Toxins (Basel) **2023**; 15.

21. Ramos-Sevillano E, Ercoli G, Guerra-Assuncao JA, et al. Protective Effect of Nasal Colonisation with cps/piaA and cps/proABCStreptococcus pneumoniae Strains against Recolonisation and Invasive Infection. Vaccines (Basel) **2021**; 9.

22. Travaglini KJ, Nabhan AN, Penland L, et al. A molecular cell atlas of the human lung from single-cell RNA sequencing. Nature **2020**; 587:619-25.

23. Herta T, Bhattacharyya A, Bollensdorf C, et al. DNA-release by Streptococcus pneumoniae autolysin LytA induced Krueppel-like factor 4 expression in macrophages. Sci Rep **2018**; 8:5723.

24. Barutcu AR, Lajoie BR, Fritz AJ, et al. SMARCA4 regulates gene expression and higher-order chromatin structure in proliferating mammary epithelial cells. Genome Res **2016**; 26:1188-201.

25. Wilson R, Cohen JM, Jose RJ, de Vogel C, Baxendale H, Brown JS. Protection against Streptococcus pneumoniae lung infection after nasopharyngeal colonization requires both humoral and cellular immune responses. Mucosal Immunol **2015**; 8:627-39.

26. Periselneris J, Ercoli G, Pollard T, et al. Relative Contributions of Extracellular and Internalized Bacteria to Early Macrophage Proinflammatory Responses to Streptococcus pneumoniae. mBio **2019**; 10.

27. Galan JE. Salmonella Typhimurium and inflammation: a pathogen-centric affair. Nat Rev Microbiol **2021**; 19:716-25.

28. Li G, Wang G, Li M, et al. Morin inhibits Listeria monocytogenes virulence in vivo and in vitro by targeting listeriolysin O and inflammation. BMC Microbiol **2020**; 20:112.

29. Connor V, German E, Pojar S, et al. Hands are vehicles for transmission of Streptococcus pneumoniae in novel controlled human infection study. Eur Respir J **2018**; 52:1800599.

30. Browall S, Norman M, Tangrot J, et al. Intraclonal variations among Streptococcus pneumoniae isolates influence the likelihood of invasive disease in children. J Infect Dis **2014**; 209:377-88.

31. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol **2014**; 15:550.

32. Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological themes among gene clusters. OMICS **2012**; 16:284-7.

33. Rheinwald JG, Green H. Serial cultivation of strains of human epidermal keratinocytes: the formation of keratinizing colonies from single cells. Cell **1975**; 6:331-43.

549   34. Carpenter AE, Jones TR, Lamprecht MR, et al. CellProfiler: image analysis software for identifying and
550   quantifying cell phenotypes. Genome Biol **2006**; 7:R100.
551   35. Kirkham LA, Kerr AR, Douce GR, et al. Construction and immunological characterization of a novel
552   nontoxic protective pneumolysin mutant for use in future pneumococcal vaccines. Infect Immun **2006**;
553   74:586-93.

FIGURE LEGENDS

Fig 1. Pneumococcal – epithelial interactions in the EHPC model. (A) Schematic showing that

during colonisation, pneumococci adhere to the epithelial cell surface (association), internalise inside

epithelial cells (microinvasion). Mucus – lime; pneumococci – purple; intracellular vacuole – dark

green; epithelial cells – yellow; nucleus – brown; cell junctions – pink. (B) Representative microscopy

images showing pneumococcal-epithelial interactions by the 6B WT, Δ*proABC/piaA* or Δ*fhs/piaA*

mutants. Surface carbohydrates (red), nuclei (DAPI, blue), pneumococci (green). (C) Table

summarising the microscopy and microbiology counts for each inoculated strain 6 days post

challenge per volunteer. Adherent represents pneumococcal surface association and intracellular

represents pneumococci that have microinvaded (inside) epithelial cells.

Fig 2. Impact of pneumococcal infection on NHBE-A cell junctional protein expression and

barrier integrity. NHBE-A Cells were cultured at ALI and infected with pneumococcal strains for

6hrs. (A) Top: Representative confocal microscopy images from 4 independent experiments with

replicates, showing pneumococcal (green) association with the epithelium (red, surface

carbohydrates; nuclei blue) at 6hr post infection. Bottom: Representative transmission electron

microscopy images from 2 independent experiments, showing pneumococcal microinvasion

examples of each strain (arrows), enclosed within a vacuole after 4hr infection. Representative

confocal microscopy images from four independent experiments with replicates showing

pneumococcal (green) interactions with junctional proteins (red); JAM-A (B), ZO-1 (C) and β catenin

(D) in primary HBE-A cells after a 6hr exposure. Nuclei stained with DAPI in blue. Co-localisation of

junctional proteins and pneumococci appear yellow, as highlighted by arrows. Integrated intensity of

β catenin in pneumococcal negative (not pneumococcal associated) or positive (pneumococcal

associated) cells. Non infected v pneumococcal negative cells **** p <0.0001 (NI v Δ*proABC/piaA*),

** p = 0.0026 (NI v Δ*fhs/piaA*); non-infected v pneumococcal positive cells; **** p <0.0001 (NI v

Δ*proABC/piaA* and NI v Δ*fhs/piaA*); Kruskal-Wallis, n = 4 independent experiments.

Fig 3. Pneumococcal – epithelial interactions with NHBE-A. Cells were cultured at ALI and infected with pneumococcal strains for 6hrs. (A) Representative scanning electron microscopy images from 2 independent experiments, showing the surface of the epithelium. Representative examples of differentiated epithelia (basal cells, Clara cells, goblet cells, ciliated cells) are shown in yellow text. Colonising pneumococci are highlighted by arrows. Representative confocal microscopy images from 4 independent experiments with replicates for (B) Ciliated cells (acetylated tubulin, red), (C) Goblet cells (mucin5ac, red, (arrows indicate co-localisation of mucin5ac with pneumococci)), (D) Clara cells (uteroglobin, red) after 6hr infection. Pneumococci (green), nuclei (DAPI, blue), colocalization of red and green appears as yellow. Integrated intensity of acetylated tubulin (B) or Muc5ac (C) in pneumococcal negative or positive cells from two or more images. Acetylated tubulin, neg p = 0.2677, pos p = 0.017, ANOVA. Mucin5ac, neg p = 0.6852, pos p = 0.8510, N = 3 or 4 independent experiments. (D) Integrated intensity of uteroglobin normalised to non-infected cells (left, p = 0.0115 non-infected v infected, Kruskal-Wallis) and percentage of Clara positive cells following infection (right, p = 0.0003 non-infected v strains, p = 0.0036 6B v mutant strains, Kruskal-Wallis). N = 3 independent experiments.

Fig 4. **Strain association, microinvasion and transmigration of Detroit 562 cells.** Cells were infected with *S. pneumoniae* for three hours and CFU counts recorded. (A) Representative confocal microscopy images from 6 independent experiments with replicates, showing pneumococcal (green) interactions with surface carbohydrates (red). (B, top left) Association of pneumococcal strains to epithelial cells. p = 0.0009 (Kruskal-Wallis), n = 6 with replicates. (B, top right) Internalisation of strains into cells, following a gentamicin protection assay. p = 0.0219 (Kruskal-Wallis) N = 6 with replicates. (B, bottom left) Pneumococcal viability inside epithelial cells following a gentamicin protection assay. n = 3 or 4 independent experiments with replicates. Counts were normalised to microinvasion counts for each strain after 3 hours infection and compared to 6B WT; +1hr p = 0.6517, +2hr p = 0.1377, +3hr p = 0.1183 (ANOVA). (B, bottom right) Transmigration of pneumococcal strains across confluent monolayers on transwell inserts. CFUs were recorded from the basal

chamber. N = 6 independent experiments with replicates. *p* < 0.001 (Kruskal-Wallis). (C) Representative confocal microscopy images from 6 independent experiments with replicates, showing pneumococcal (green) interactions with β catenin (red). Nuclear DAPI stain (blue).

Fig 5. Detroit 562 cell transcriptomic responses to pneumococcal strains. Epithelial cells were infected for four hours with pneumococcal strains and RNA extracted and processed for RNAseq. 6B n = 11, PP n = 17, FP n = 14, non-infected n = 12, from 6 independent experiments. (A) All significantly log2 fold change upregulated genes according to strain compared to non-infected cells. Significance was defined as DEGs with a log2 fold change (logFC) of > 1.5 and false discovery rate (FDR) of < 0.05. (B) Comparison of the top 20 most upregulated genes for each strain, compared to non-infected cells. Upstream regulator analysis for (C) transcription factors and (D) kinases.

Fig 6. Effects of pneumococcal infection on caspase activity in Detroit 562 cells. Caspase activity was assessed by luminescence, following infection with pneumococcal strains for 6 hours. (A) Caspase 3/7 activity was determined following incubation with the specific inhibitor Z-VAD-FMK. N = 4 with replicates. In comparison to non-infected cells, caspase 3/7 *p* = 0.0022; caspase 3/7 + Z-VAD-FMK inhibitor *p* = 0.0005. In comparison to 6B WT, caspase 3/7 *p* = 0.0215; caspase 3/7 + Z-VAD-FMK inhibitor *p* = 0.0012 (Kruskal-Wallis). (B) Caspase 1 activity was determined following incubation with the specific inhibitor YVAD-CHO. Negative signals indicate caspase activity other than caspase 1. N = 6 with replicates. In comparison to non-infected cells, *p* = 0.39; caspase+YVAD-CHO inhibitor *p* = 0.3604 and, in comparison to 6B WT, *p* = 0.6226; caspase+YVAD-CHO inhibitor *p* = 0.5386 (Kruskal-Wallis). (C) Caspase 8 activity was assessed following infection with pneumococcal strains. N = 3 with replicates. In comparison to non-infected cells, *p* = 0.0015 (ANOVA) and, in comparison to 6B WT, *p* = 0.0172 (ANOVA).

Fig 7. Hydrogen peroxide secretion and pneumolysin activity from pneumococcal strains in the absence and presence of Detroit 562 cells. (A) Fold change of hydrogen peroxide production

from the mutants compared to WT, normalised to the CFU counts. 3hr p = 0.0087 (ANOVA), 6hr p

= 0.0209 (Kruskal-Wallis). (B) Fold change of hydrogen peroxide production from the mutants

compared to WT, normalised to the CFU counts, following infection of Detroit 562 cells. Media was

replaced after 1hr infection and cells incubated for a further 5hrs. N = 3 independent experiments

with replicates. Compared to 6B WT, p = 0.0034 (ANOVA). (C) Pneumolysin activity was assessed

using a red blood cell lysis assay. Strains were incubated with red blood cells for 30 minutes and

absorbance measured. N = 4 experiments with replicates. Pneumococcal strain haemolytic activity

is calculated in comparison to 100% lysis by 0.0625% saponin. p <0.0001 compared to 6B WT

(ANOVA).

Figure 1. Pneumococcal – epithelial interactions in the EHPC model

A

B

C

6B	MICROSCOPY		MICROBIOLOGY
	Adherent	Intracellular	
#1	0	0	0
#2	2	0	520
#3	0	0	0
#4	1	0	0
#5	0	0	9
#6	1	2	223
#7	0	0	5725
#8	39	4	1110
#9	0	0	0
#10	0	0	767
#11	0	0	0

ΔproABC/piaA	MICROSCOPY		MICROBIOLOGY
	Adherent	Intracellular	
#1	0	0	0
#2	0	0	0
#3	0	0	0
#4	5	3	0
#5	3	2	0
#6	8	2	5
#7	0	0	0
#8	0	2	0
#9	0	2	1
#10	0	0	0
#11	0	0	0

Δfhs/piaA	MICROSCOPY		MICROBIOLOGY
	Adherent	Intracellular	
#1	2	0	21856
#2	1	3	0
#3	3	3	16
#4	51	54	0
#5	0	0	520325

Figure 2. Impact of pneumococcal infection on NHBE-A cell junctional protein expression and barrier integrity.

Figure 3. Pneumococcal – epithelial interactions with primary human epithelial cells.

Figure 4. Strain association, microinvasion and transmigration of Detroit 562 cells.

Figure 5. Detroit 562 cell transcriptomic responses to pneumococcal strains.

A

B

log2foldchange	6B	ΔproABC/piaA	Δfhs/piaA
Upregulated genes			
AC256236.1	32.9	25.5	0.0
AC022149.1	30.2	27.6	0.0
Z84488.2	29.1	21.6	0.0
MUC21	25.8	29.0	0.0
VASN	20.9	0.0	0.0
BRD2	18.4	24.7	0.0
AC012254.2	18.2	13.8	0.0
CTAGE4	18.1	0.0	0.0
MICE	17.1	20.3	0.0
AC107075.1	10.1	0.0	0.0
CSNK2B	0.0	20.2	0.0
MTRNR2L6	0.0	19.4	15.8
LSM2	0.0	17.1	0.0
ALOX5	0.0	0.0	6.9
AC006064.6	0.0	0.0	3.4
TAP2	0.0	0.0	3.2
CYP1A1	0.0	0.0	1.7
FOSB	0.0	0.0	1.4
IL1B	0.0	0.0	1.3
CXCL8	0.0	0.0	1.1
CXCL2	0.0	0.0	1.1
CYP1B1	0.0	0.0	1.0

C

Transcription factors

D

Kinases

Figure 6. Effects of pneumococcal infection on caspase activity in Detroit 562 cells.

Figure 7. Hydrogen peroxide secretion and pneumolysin activity from pneumococcal strains in the absence and presence of Detroit 562 cells.

Graphical abstract