Full title

Epithelial microinvasion and immunity elicited by disease-attenuated pneumococcal

biosynthesis gene mutants

Running title

Epithelial microinvasion, immunity and disease

Caroline M. Weight[1*], Gabriele Pollara[1], Modupeh Betts[1], Roberta Ragazzini[2], Elisa Ramos-

Sevillano[3], Jesús Reiné[4], Matthew Whelan[1], José Afonso Guerra-Assunção[1], Michael Connor[5],

Paola Bonfanti[1,2], Clare Jolly[1], Mahdad Noursadeghi[1], Daniela M. Ferreira[4], Jeremy S. Brown[3],

Robert S. Heyderman[1]

[1]Research Department of Infection, Division of Infection and Immunity, UCL London, UK

[2]Francis Crick Institute, London, UK

[3]Department of Respiratory Medicine, UCL, London, UK

[4]Oxford Vaccine Group, University of Oxford, Oxford and Clinical Sciences, Liverpool School of

Tropical Medicine, Liverpool, UK

[5]Chromatin and Infection, Institut Pasteur, France

*Current address Biomedical and Life Sciences, Faculty of Health and Medicine, Lancaster

University, UK

Corresponding Authors: Dr Caroline M. Weight and Professor Robert S. Heyderman

Email: c.weight@lancaster.ac.uk and r.heyderman@ucl.ac.uk

ABSTRACT

Nasopharyngeal colonisation by *Streptococcus pneumoniae* is characterised by adherence to

epithelial cells, microinvasion and an innate immune response. Here, we explore the hypothesis that

pneumococcal virulence has primarily evolved to promote these colonisation processes and

subsequent transmission, rather than to cause disease. We have used a *S. pneumoniae* serotype

6B strain, alongside two biosynthesis gene mutants (Δ*proABC* and Δ*fhs* with an additional Δ*piaA*

mutation) known to colonise murine and human models without causing disease. We show that these

mutations do not attenuate microinvasion in either experimental human pneumococcal challenge or

in vitro in primary and immortalised epithelial cells. However, interactions and subsequent cellular

reorganisation by primary human airway epithelial and Detroit 562 nasopharyngeal cells were strain

dependent, and the inflammatory epithelial innate response and caspase 8 activity were attenuated

by the Δ*fhs/piaA* but not the Δ*proABC/piaA* mutation. Growth of the Δ*fhs/piaA* mutant was associated

with increased pneumococcal hydrogen peroxide secretion and reduced pneumolysin activity when

compared to the WT and Δ*proABC/piaA* mutant strains. Analysis of the bacterial transcriptome under

stress conditions revealed that the mutations resulted in broad, differential effects on pneumococcal

virulence and metabolic gene expression suggestive of compensatory changes that preserve

microinvasion. The findings that cellular reorganisation, the innate immune response, and the ability

to cause invasive disease are not necessarily interconnected emphasise the complexity of bacterial-

host interactions at the mucosal surface. Colonisation outcome cannot be fully explained by single

or functionally grouped virulence or metabolic factors.

IMPORTANCE

Colonization of the nasopharynx by *Streptococcus pneumoniae* is a precursor to both disease and

bacterial transmission. This study challenges the assumption that virulence factors have primarily

evolved to cause disease, proposing instead that these pathogenic factors more aligned with

facilitating bacterial carriage and spread. Using a pneumococcal clinical isolate strain and two

biosynthesis gene mutants in experimental human pneumococcal challenge and epithelial cell line

models, we demonstrate that epithelial microinvasion can occur independently of disease causation. Moreover, we uncover a complex relationship between bacterial adaptation and epithelial immune response, with differential effects on epithelial cell reorganization, inflammation, and bacterial growth dynamics. Our findings underscore our contention that pneumococcal colonisation and disease outcomes cannot be predicted by individual gene profiles alone. This phenotyping approach offers the opportunity to effectively unravel *S. pneumoniae* pathogenesis and its adaptation to the nasopharyngeal niche.

Keywords

Biosynthesis genes, host-pathogen interactions, Innate immunity, *Streptococcus pneumoniae*, airway epithelium

Graphical Abstract

S. pneumoniae colonisation is characterised by mucus association, epithelial adherence and microinvasion – where the pneumococcus invades the epithelial barrier without causing disease. Microcolonies form at the epithelial surface and pneumococcal transmigration can occur between epithelial cells. Although mutations in *S. pneumoniae* biosynthesis genes (Δ*proABC* and Δ*fhs*) attenuate disease in a murine model, they do not attenuate microinvasion in either experimental human pneumococcal challenge (EHPC) or *in vitro* in primary and immortalised epithelial cells. These mutations show strain-dependent effects on both the epithelial and bacterial responses to infection. While not directly co-correlated, factors such as epithelial cellular reorganisation, inflammation and caspase 8 activity alongside pneumococcal metabolic adaptation, virulence factor expression and response to stress are key determinants of the outcome of colonisation in the nasopharyngeal niche.

INTRODUCTION

Adaptation to the host environment is crucial for successful colonisation of the upper respiratory tract (URT) by a range of human pathogens, including bacteria, viruses and fungi. This includes the ability to withstand different ambient temperatures, overcome physical barriers such as mucus, evade epithelial soluble mediators such as antimicrobial peptides, compete with neighbouring polymicrobial communities and compensate for nutrient restriction (1). *Streptococcus pneumoniae* (the pneumococcus) is a common commensal of this URT niche (1). However, translocation of the pneumococcus from the nasopharynx to the lungs, blood or the brain results in life-threatening disease (1). Indeed, among children under five years, *S. pneumoniae* is the leading cause of pneumonia deaths globally (2).

Using experimental human pneumococcal challenge (EHPC) of healthy volunteers and well-characterised epithelial cell models, we have previously shown that colonisation by *S. pneumoniae* is characterised by epithelial surface microcolony formation and invasion which we have called microinvasion (i.e. invasion of the epithelial barrier without disease) (3). The pneumococcal capsule is a major virulence determinant, influencing carriage, transmission from person to person, and invasive disease potential (4). Indeed, we have shown that *S. pneumoniae* epithelial microinvasion also varies markedly between strains expressing different capsular serotypes, with a serotype 4 strain invading Detroit 562 nasopharyngeal cell monolayers more readily than serotype 6B and 23F strains, and inducing more marked pro-inflammatory epithelial transcriptomic and cytokine responses (3).

We and others have previously proposed that although the epithelial innate immune response that results from pneumococcal colonisation may facilitate clearance, this highly variable pathobiont (>100 serotypes; >600 global pneumococcal sequence clusters (5)) tunes the host response to enhance transmission(3, 4). In addition to capsule, these processes have been shown to be

modulated by a broad array of subcapsular and cytoplasmic virulence factors (6), and more recently,

by bacterial carbohydrate, nucleotide, fatty acid and purine/ pyrimidine metabolism (7-11).

To explore the hypothesis that pneumococcal virulence has primarily evolved to promote

colonisation and subsequent transmission, but not necessarily disease, we have used a unique set

of *S. pneumoniae* serotype 6B Δ*proABC* and Δ*fhs* biosynthesis gene mutants developed to induce

protective immunity without causing disease (12). *proABC* and *fhs* were chosen as targets through

the analysis of previously published transcriptome and transposon mutational screens (11, 13, 14).

The *proABC* operon, consisting of genes predicted to encode a γ-glutamyl phosphate reductase

(*proA*), a γ-glutamyl kinase (*proB*) and a pyrroline-5-carboxylate reductase (*proC*), and *fhs* predicted

to encode a folate tetrahydrofolate reductase involved in one carbon metabolism pathways have

been implicated in metabolic adaptation to growth (15-17), and are highly transcribed in mouse

models of pneumococcal infection (16, 17). To further avoid reversion to a disease-causing

phenotype during EHPC, these mutants also contain a deletion of *piaA* which encodes the lipoprotein

component of the dominant pneumococcal iron uptake ABC transporter (18, 19). We have previously

shown that in a murine model, when compared to the *S. pneumoniae* serotype 6B wild-type parent

strain (WT, BHN418), the Δ*proABC/piaA* and Δ*fhs/piaA* mutants colonise the nasopharynx as

effectively. However, despite expressing the same capsule as the wild type 6B strain, neither the

Δ*proABC/piaA* and Δ*fhs/piaA* 6B strains causes invasive disease (12) (17). We have subsequently

shown that this attenuated phenotype is at least in part due to impaired growth in serum and is

associated with increased oxidative stress and altered lipid profiles (17). In EHPC, we have also

shown that both Δ*proABC/piaA* and Δ*fhs/piaA* retain the ability to colonise the human host, and can

induce IgG-mediated immunity (20). Therefore, these otherwise isogenic strains allow us to compare

the pneumococcal-epithelial interactions of disease-causing and non-disease-causing phenotypes,

while avoiding the confounding of differences in capsular serotype.

Here, we reveal that although the *S. pneumoniae* Δ*proABC/piaA* and Δ*fhs/piaA* metabolic function mutations result in disease attenuation, the ability of the mutant strains to colonise and cause microinvasion in both EHPC and *in vitro* epithelial infection models is preserved. Crucially, we also show that despite similar patterns of epithelial cellular engagement, disruption of these key pneumococcal metabolic pathways has differential effects on cellular reorganisation and innate responses, which we propose influences niche adaptation during colonisation.

RESULTS

***S. pneumoniae* 6B with Δ*proABC* or Δ*fhs* mutants colonise in EHPC with evidence of epithelial microinvasion**

We have defined microinvasion as the process by which *S. pneumoniae* invades the mucosal epithelium, inducing an innate immune response, without necessarily causing disease (Fig 1A). Nasal curette samples from EHPC healthy volunteers inoculated with the *S. pneumoniae* serotype 6B WT or isogenic mutant strains *(Δ*proABC/piaA* or Δ*fhs/piaA*)* (21) were collected to investigate the extent of epithelial association and microinvasion. A positive carrier was defined as colonisation at any time point (day 2 or day 6 post-inoculation), as detected by microbiology culture (3). Visualisation of the mucosal cells by confocal microscopy showed colonisation by the WT and mutant pneumococcal strains (Fig 1B). At day 6 post challenge, compared to WT, there was no attenuation of microinvasion with either the Δ*proABC/piaA* or Δ*fhs/piaA* strains, with a trend for greater microinvasion (Fig 1C) with a higher microinvasive index (percentage of internalised bacteria / associated bacteria; Supplemental Information Table 1).

***S. pneumoniae* colonisation and microinvasion of primary human airway epithelium leads to cellular reorganisation with retention of barrier function**

To further explore the impact of microinvasion on epithelial biology in a complex environment, we cultured *S. pneumoniae* serotype 6B WT and the isogenic biosynthesis mutants with primary normal

human bronchial epithelial airway (NHBE-A) cells in air-liquid interface. We observed epithelial cell association and microinvasion with all three strains (Fig2A, top). TEM demonstrated that intracellular bacteria for all three strains were enclosed within vacuoles (Fig 2A, bottom). As seen previously with several different pneumococcal strains (3), the 6B WT and the mutant strains co-localised with Junctional Adhesion Molecule A (JAM-A, Supplemental Information Fig 1A, arrow), with no change in overall JAM-A expression. However, using confocal microscopy and automated image analysis, for cells infected with the 6B WT strain, we did observe an increase in the intensity of Zonula Occludens 1 (ZO-1) staining at the junctional membrane (Supplementary Fig 1B), and a significant increase in the intensity of β catenin staining for cells infected with either the Δ*proABC/piaA or* Δ*fhs/piaA* strain (Fig 2B, 2C). These changes in junctional protein expression most likely occurred as a result of cellular reorganisation, as de-novo protein production would have been insufficient over the 6-hour experimental infection period. As measured by transepithelial electrical resistance, barrier integrity was retained over the infection period, (Fig 2D).

When cultured at an air-liquid interface, NHBE cells differentiate into a typical pseudostratified columnar ciliated epithelium, consisting of goblet, Clara, basal and ciliated cells (Fig 3A). Although there was no predilection by *S. pneumoniae* for particular epithelial cell-types, pneumococci were observed amongst cilia (acetylated tubulin positive cells, Fig 3B and Supplemental Information Fig 2) and co-localised with mucus (mucin5ac positive, Fig 3C, arrows). Strikingly, while the cellular composition of the epithelial cultures exposed to the Δ*fhs/piaA* strain were comparable in appearance to non-infected cells by SEM, exposure to the 6B WT and Δ*proABC/piaA* strains resulted in an increased abundance in secreted mucus globules compared to non-infected cells (Fig 3A). This appeared to be due to differential cellular secretion of preformed mucus rather than de-novo production as these differences were not reflected in an increase in the expression of the gel-forming glycoprotein Mucin5AC (Fig 3C and 3E, right graph). Following infection by the Δ*proABC/piaA* strain we found a significant increase in expression of acetylated tubulin (Fig 3B and 3E, left graph), which has been shown to be associated with increased microtubule conformation and elasticity, and occurs by post-translational modification (22, 23). In contrast, following exposure to Δ*fhs/piaA* strain, we

saw an increased expression of the anti-inflammatory protein, uteroglobin (UG), which is a multifunctional epithelial secreted protein with anti-inflammatory and anti-chemotactic properties, and a marker for non-ciliated Clara cells (Fig 3D and 3F, left graph). These changes in UG expression again most likely occurred as a result of cellular reorganisation, or secretion of pre-formed protein, as de-novo protein production would have been insufficient to detect over the 6-hour experimental infection period.

ΔproABC/piaA and Δfhs/piaA mutations of *S. pneumoniae* serotype 6B do not affect microinvasion of Detroit 562 epithelial cells

Detroit 562 epithelial nasopharyngeal cells enable the analysis of these host-pathogen interactions under multiple conditions, which are not feasible with the primary cell lines. We have previously shown that the pattern of Detroit 562 epithelial nasopharyngeal cell microinvasion, and subsequent innate immune response to *S. pneumoniae* 6B infection *in vitro* is similar to that seen in the EPHC *in vivo(3)*. We therefore then used Detroit 562 cells to further characterise colonisation by the WT, ΔproABC/piaA and/or Δfhs/piaA pneumococcal strains. We found that although all the strains formed colonies on the epithelial surface (Fig 4A), the isogenic mutants associated less compared to 6B WT (Fig 4B). Nonetheless, microinvasion was observed with all the strains (Fig 4C), with a significantly elevated micro-invasive index for the ΔproABC/piaA and Δfhs/piaA strains compared to WT (Fig 4D). Differences in microinvasion were not due to differences in intracellular viability (Fig 4E) or replication (Supplemental Information Fig 3A-F). All of the single and double biosynthesis genes mutants (ΔproABC, Δfhs, ΔproABC/piaA, Δfhs/piaA) were impaired in their ability to transmigrate across the epithelium (Fig 4F). Importantly, we show that a single ΔpiaA mutation in the *S. pneumoniae* serotype 6B did not affect epithelial association, microinvasion and transmigration (Fig 4B, 4C, 4F).

Consistent with our previous findings for serotype 6B(3), Detroit 562 epithelial barrier function was not compromised after three hours of infection; indeed, there was a tightening of the barrier evidenced by reduced FITC dextran diffusion after pneumococcal exposure (Fig 4G). Pneumococcal

infection had no effect on Junctional Adhesion Molecule A (JAM-A), Zonula-Occludens 1 (ZO-1),

Claudin 1 or Claudin 4 expression (Supplemental Fig 4). However, we did observe a significant

increase in epithelial β catenin expression following incubation with the ΔproABC/piaA and Δfhs/piaA

strains, compared to the WT (Fig 4H, 4I).

Δfhs/piaA but not the ΔproABC/piaA mutations attenuate the pro-inflammatory epithelial

innate response to pneumococcal colonisation and microinvasion

We have previously shown that the Detroit 562 cell innate immune response to *S. pneumoniae* is

strain dependent (3). Having found that serotype 6B biosynthesis gene mutants that are attenuated

for disease retain their capacity for microinvasion, we have further assessed the link between

colonisation, microinvasion and epithelial cell innate immunity by RNAseq. Compared to non-

infected cells (Fig 5A), all strains upregulated 78 core epithelial genes related to cytokine signalling

in immune system, NLR signalling and GPCR ligand binding. 193 genes were upregulated by both

the 6B WT and ΔproABC/piaA strains, characterised by Rhodopsin-like receptors, GPCR ligand

binding, cytokine signalling in Immune system, signalling by interleukins and Toll Receptor cascades.

Infection with the Δfhs/piaA strain resulted in a markedly different transcriptomic profile with modest

upregulation of 254 genes, characterised by antigen processing, proteasomal pathways, GTP

hydrolysis, and intracellular stress responses (Fig 5A). Overall, the epithelial genes that were most

upregulated following infection with 6B WT or ΔproABC/piaA strain were similar and contrasted to

the reduced gene expression profile seen with the Δfhs/piaA strain despite high levels of cell invasion

(Fig 5B). Since both mutant strains contained the *piaA* mutation, this could not account for the

differences seen in these responses. Together, these data demonstrate that microinvasion, the

innate epithelial response and disease are not necessarily linked.

Transcription factors, cellular kinases and cytokines (Fig 5C, 5D) predicted to regulate the

differentially expressed genes, showed the greatest difference in profile following infection with

Δfhs/piaA strain compared to infection with 6B WT and ΔproABC/piaA strains. Δfhs/piaA was

predicted to show diminished NFκB and MAPK activation and was associated with a more intrinsic

and epithelial repair response that included TBK1, Raf1, JAK1 kinases and IL-1 and IL-33 cytokine

signalling.

Enhanced epithelial caspase 8 activity following pneumococcal infection with the wild-type

and ∆proABC/piaA strains

The NLRP3 inflammasome drives programmed epithelial cell death via caspase 1, caspase 3/7 and

caspase 8 activity (24). Caspase activity, primarily regulated post-translationally, enables a rapid

response to cellular stress and pathogen-mediated activation (25). In health, caspase regulation is

thought to be important for mucosal integrity (26). During bacterial pneumonia, caspase activation is

thought to have an important pro-inflammatory role, possibly triggered by pneumolysin (27, 28). We

have therefore explored whether the different patterns of microinvasion and epithelial reorganisation

seen with the WT 6B, and mutant strains are associated with differential caspase activation.

Compared to non-infected cells, infection with all strains led up to a 1.5-fold increase in epithelial

caspase activity 6-hour post infection (Fig 6A, dark grey bars). To test which caspases were

activated, we incubated infected cell cultures with either a caspase 1 inhibitor (YVAD-CHO), a

caspase 3/7 inhibitor (Z-VAD-FMK), or specifically measured caspase 8 activity. We found that the

increase was not due to caspase 1 activity (Fig 6B). Compared to non-infected cells, caspase 3/7

accounted for 3% and 15% of the caspase activity induced following infection with 6B WT and

∆*fhs/piaA* strain respectively (Fig 6A). In contrast, ∆*proABC/piaA* strain did not induce caspase 3/7

activity. Significant caspase 8 activation was seen following infection with 6B WT and ∆*proABC/piaA*

strain, but not with the ∆*fhs/piaA* strain (Fig 6C).

Together, these data suggest differential caspase induction following infection with the different

strains, linking the proinflammatory profile of the 6B WT and the ∆*proABC/piaA* with caspase 8

activation, which we postulate may regulate pneumococcal survival and transmission during
colonisation in the nasopharynx.

269 Deletion of the *proABC* and *fhs* genes have broad-ranging differential effects on the *S.*
270 *pneumoniae* virulence gene expression and metabolic gene pathways when under stress

We have previously found marked transcriptional differences between the WT and both mutants in
response to serum stress (used to represent the physiological stress found during human infection),
with an over three-fold increase in the number of genes upregulated by the mutant strains compared
to WT (17). Further analysis revealed that these genes include well-characterised pneumococcal
virulence factors such as *ply*, *nanA*, *psaA,* those regulating oxidative stress such as *SpxB*, *lctO* and
adhE, and genes involved in purine and carbohydrate metabolism (Supplemental Information Table
2). Phosphotransferase systems, amino nucleotide sugar, fructose and mannose and metabolic
pathways were the most significantly enriched pathways in the Δ*proABC* mutant under serum stress
(Table 1). In contrast, competence, purine, pyruvate, metabolic pathways, amino nucleotide sugar
pathways and secondary metabolites were the most significantly enriched pathways in the Δ*fhs*
mutant under stress (Table 1). Together these data suggest that deletion of the *proABC* and *fhs*
genes have widespread but differential effects on *S. pneumoniae* virulence and metabolism which
may explain then complex phenotypes seen following epithelial infection.

285 *fhs* gene mutation in *S. pneumoniae* 6B is associated with increased pneumococcal hydrogen
286 peroxide secretion and decreased pneumolysin activity

S. pneumoniae SpxB and LctO generate hydrogen peroxide (H_2O_2) which has been shown to
suppress ENaC-α transcription and inhibit the Na+–K+ pump, acting as an intracellular signal to
increase mitochondrial production of ROS (29, 30). We therefore postulated that differential
expression of pneumococcal H_2O_2 may modulate epithelial functional responses during

nasopharyngeal colonisation by *S. pneumoniae* and explain the differences in the epithelial innate

immune response to the 6B WT, Δ*proABC/piaA* and Δ*fhs/piaA* strains. To assess this, we measured

H_2O_2 concentration using a luminescence-based bioassay, normalising against bacterial CFU. After

three and six hours, we detected significantly higher levels of H_2O_2 in Δ*fhs/piaA* cultures compared

to 6B WT and Δ*proABC/piaA* strains (Fig 7A). Furthermore, this was not ameliorated by epithelial

cell contact (Fig. 7B).

The pore forming pneumococcal toxin pneumolysin can both activate and suppress the host innate

immune response and mediates the haemolytic activity characteristic of the pneumococcus (31-33).

H_2O_2 can increase the release of pneumolysin but paradoxically also negatively impairs its

haemolytic activity (34). Compatible with the latter and the higher production of H_2O_2 by the Δ*fhs*

/piaA strain, despite having increased expression of *ply*, red blood cell lysis was reduced for Δ*fhs*

strains compared to 6B WT and Δ*proABC* (Fig. 7C).

Differential pneumococcal H_2O_2 release may therefore explain at least in part, the effects of the *S.*

pneumoniae serotype 6B WT and Δ*proABC /piaA* compared to the Δ*fhs /piaA* strain on the epithelial

response to colonisation and microinvasion. Importantly, in the 6-hour timeframe assessed, the

increased H_2O_2 release the Δ*fhs /piaA* strain was not associated with epithelial cellular toxicity or

loss of barrier function, but was associated with increased intracellular epithelial stress response, as

detected by RNAseq.

DISCUSSION

There is considerable variation in the propensity of different *S. pneumoniae* strains to colonise the

human host, cause invasive disease and transmit to others (1). During colonisation of the

nasopharyngeal niche, the pneumococcus was previously thought to be confined to the mucus layer

and the epithelial surface (1). However, using EHPC, we have previously reported that

pneumococcal microinvasion occurs in healthy volunteers (3), and we now show that even when

using isogenic serotype 6B pneumococcal biosynthesis pathway mutant strains that are attenuated

for systemic disease but not for carriage, this microinvasion is preserved (12). We found that

pneumococcal colonisation and microinvasion of primary human airway epithelium led to cellular

reorganisation and mucus production while retaining barrier function, although the pattern of this

reorganisation varied by strain. Furthermore, the pro-inflammatory innate response and caspase 8

activity associated with epithelial infection by *S. pneumoniae* serotype 6B were attenuated by the

Δ*fhs/piaA* but not by the Δ*proABC/piaA* mutation. Under stress conditions, along with presumably

compensatory changes in the expression of multiple metabolic pathways, the Δ*proABC* and Δ*fhs*

mutations resulted in the upregulation of several pneumococcal virulence factor genes (Graphical

Abstract Summary). Notably, the *fhs* gene mutation in *S. pneumoniae* serotype 6B was associated

with increased pneumococcal hydrogen peroxide secretion and decreased pneumolysin activity.

Together these findings suggest that despite being attenuated for disease, the metabolic changes

and the increased expression of pneumococcal virulence genes in response to stress may explain

the preservation of epithelial microinvasion by these mutants. However, although epithelial

microinvasion, cellular reorganisation, innate immune response, and the propensity to cause

invasive disease are strain dependent, these properties do not appear to be directly co-correlated.

The complexity of the interaction between *S. pneumoniae* and the mucosal epithelium was

highlighted through infection of our well-characterised primary epithelial cell model(35). The model

consisting of Clara cells (involved in secretory, xenobiotic metabolic, and progenitor cell functions),

goblet cells (mucus production), ciliated cells (responsible for mucus and small particle transport),

exhibited considerable epithelial cellular re-organisation involving β catenin, acetylated tubulin and

uteroglobin in response to pneumococcal infection. In a rabbit pneumococcal meningitis model,

increased acetylated tubulin and microtubule stabilisation was attributed to pneumolysin activity(23).

Previously, using a pneumolysin deficient mutant of *S. pneumoniae* serotype 4 (TIGR4), we have

shown that pneumolysin is a key driver of the epithelial inflammatory response to pneumococcal

infection (3). We therefore speculate that the differential expression of pneumolysin and H_2O_2 by the

WT and mutant strains may contribute to the differing patterns of epithelial reorganisation, and goblet

cell mucus release that we observed, and may impact on both clearance and transmission (1).

Epithelial infection with the 6B WT and Δ*proABC/piaA* strains induced innate inflammatory pathway

gene expression typical of pneumococcal infection (3), including cytokine/chemokines, interleukins,

Toll Like Receptor and NFκB signalling, alongside apoptotic transcripts such as FOXO3 and TP53.

We also identified epithelial genes upregulated only during WT infection, potentially related to a

greater disease-causing potential. These genes are involved in RNA degradation, O-linked

glycosylation, wnt signalling, DNA sensing, RIG-I pathways and signal transduction. In contrast, the

Δ*fhs/piaA* strain activated a pro-survival and adaptation profile, upregulating genes such as JUN,

SMARCA4, NFAT5, KLF6 and TCF4, which are involved in homeostatic signalling and chromatin

remodelling (36, 37), suggestive of a more tolerant response. Ultimately, these capsule-independent

differences in pneumococcal-epithelial interactions may shape the adaptive immune response to

pneumococcal colonisation (21, 38), which could then further influence the outcomes of both

colonisation and disease. How these findings translate to differences in the duration of

nasopharyngeal carriage, invasiveness, transmission or clearance remains to be fully determined.

However, during EHPC, Δ*fhs/piaA* infection was associated with increased density at day 6 post-

inoculation compared the other strains (20).

Importantly, while WT serotype 6B infection led to epithelial cell reorganisation (β catenin localisation

to the junctional membrane and mucus secretion), pneumococcal transmigration, a more pro-

inflammatory innate immune transcriptomic profile and increased caspase 8 activity, all indicative of

a robust cellular response to *S. pneumoniae*, the Δ*fhs/piaA* but not Δ*proABC/piaA* strains was

attenuated for many of these characteristics. This suggests that similar to the macrophage response

to *S. pneumoniae*, intracellular signalling by pneumococci is not the sole determinant of the epithelial

response to infection (32, 39). The *proABC* operon consists of three genes involved in proline

biosynthesis (ProABC)(40). *fhs* catalyses the synthesis of N10 formyl tetrahydrofolate, essential for

carbon metabolism and purine nucleotide biosynthesis in various bacterial species (16, 41),

especially under hypoxic conditions (41, 42). It is increasingly recognised that bacterial virulence and

metabolism are intimately linked (43). Whether the distinct epithelial innate immune response to

Δ*fhs/piaA* infection is due to diminished epithelial NFκB and MAPK activation (potentially linked to

extracellular receptor engagement), reduced caspase 8 activity mediated by enhanced

pneumococcal H_2O_2 release, decreased pneumolysin activity, or a compensatory metabolic

response to the stress remains to be determined.

S. pneumoniae is highly adapted to colonise the human host and transmit from person to person.

Invasive pneumococcal disease, particularly in its most severe forms, is a dead end for the bacteria.

Our observation that the induction of inflammation/ innate immunity is not necessarily linked to

disease potential supports our proposition that microinvasion during colonisation by the

pneumococcus may facilitate immune evasion. As is typical of 'professional' intracellular bacterial

species such as *Listeria* and *Salmonella*, the resulting host response might also promote

transmission (44, 45). Whether the less pro-inflammatory Δ*fhs/piaA* is less transmissible than the

more inflammatory WT remains to be determined but this could be further explored using EHPC

(46).

In conclusion, these data further support our proposed paradigm that epithelial microinvasion by *S.*

pneumoniae in healthy people is frequent and usually does not lead to disease. Our findings highlight

the broad-ranging effects of even single gene mutations in pneumococcal biosynthesis pathways,

which may partially explain the complexity of differences between seemingly closely related strains.

However, strain differences in subsequent epithelial cell microinvasion, cellular reorganisation and

inflammatory responses are likely driven by differential expression of a range of bacterial virulence

and metabolic pathways that are not necessarily co-correlated with disease-causing potential. As

suggested by our bacterial transcriptomic data, no single pneumococcal virulence factor or group of

factors fully explains the differences seen between the strains under stress. The complexity of these

bacterial-host cell interactions is modulated by mediators such as hydrogen peroxide and

pneumolysin, which can have both enhancing and inhibitory effects on the epithelial cell response to

infection in different contexts. *S. pneumoniae* in constantly evolving, with new variants emerging

both due to stochastic events and under ecological pressure from numerous factors including

vaccines and antibiotic exposure (47, 48) (49). Extensive phenotyping of emerging, genotypically

related pneumococcal strains offers the opportunity to more effectively unravel *S. pneumoniae*

pathogenesis and its adaptation to the nasopharyngeal niche.

METHODS AND PROTOCOLS

Bacteria

The *Streptococcus pneumoniae* serotype 6B (strain BHN 418(50)) was a kind gift from Prof. Birgitta

Henriques Normark (Karolinska Institute). Δ*proABC/piaA* and Δ*fhs/piaA* mutant strains were

generated using overlap extension PCR as described by Ramos-Sevillano *et al*(12). Stocks of

bacteria were stored at -80°C when bacterial growth in Todd Hewitt Broth (THY) was 0.3 O.D.$_{600nm}$.

Aliquots were defrosted, centrifuged at 8000g for eight minutes and resuspended in 1% Fetal Bovine

Serum (FBS, Gibco) in MEM media (Gibco) for infections. Approximately 10 bacteria were added for

every epithelial cell. Colony forming units (CFU) were counted by serial dilutions on horse blood agar

plates (EO Labs). Starting inoculums were not significantly different across strains for all assays.

Bacterial RNA-seq sample collection, sequencing and reads

Mutant pneumococcal strains (three biological replicates for each mutant) and 6B WT (three

biological replicates) were grown in THY to mid log phase and the pellets resuspended in THY and

undiluted human serum for 60mins. The culture was then pelleted, and RNA stored in RNA Protect

at -70°C before extracting total RNA (17). Total RNA from each library was ribo-depleted and sent

for Illumina Next Seq Sequencing Pathogen Genomics Unit (PGU), UCL. The quality of raw FASTQ

reads was checked by FastQC v0.11.5, Babraham Bioinformatics, UK (51) and visualised using

multiQC v1.9 (52). Reads were trimmed using Trimmomatic 0.39 (53). Trimmed reads were also

checked by FastQC and multiQC. Trimmed reads were then mapped to KEGG annotated 6B

genome sequence (670-6B, Accession: CP002176.1) using bowtie2 with default settings(54).

Conversion into BAM files was performed using SAMtools (55). Mapped reads were visualized in

the Integrated Genome Browser (IGB) (56). FeatureCounts v2.0.0 was used to summarize read

counts for each annotated feature in the respective reference genomes in multimapping mode (-M)

(57).

Differential gene expression analyses

The generated count matrix was imported into R-studio (R v3.4.2) and normalization of counts and

differential gene expression analysis was performed by DESeq2 (58). DESeq2 normalized libraries

were regularized log transformed for visualizing heatmaps and clustering. Differential gene

expression was performed on raw counts. Fold change shrinkage was implemented with the apeglm

package within DESeq2. Genes with a log2 fold change > 1.5 and false discovery rate (FDR) of <

0.05 were categorized as differentially expressed. KEGG pathway enrichment and module analysis

were performed on differentially regulated genes using over representation analysis (ORA)

implemented by clusterProfiler (59).

Experimental Human Pneumococcal Carriage Model (EHPC)

A full explanation of experimental design and selection criteria has been previously described (3).

Following written informed consent, healthy non-smoking adults between the ages of 18 – 59 were

inoculated with 80,000 CFU/nostril live strains of *S. pneumoniae*. Mucosal cells from the inferior

turbinate using a plastic Rhino-probe™ (curettage, Arlington Scientific) were collected prior to

inoculation, and days 2 and 6 post-inoculation. Samples were processed for confocal microscopy

analysis. Nasal wash samples were collected for CFU count analysis. There were no adverse events.

Ethical approval was given by NHS Research and Ethics Committee (REC)/Liverpool School of

Tropical Medicine (LSTM) REC, reference number: 18/NW/0481 and Human Tissue Authority

licensing number 12548.

Primary Human Epithelial Cells

Normal Human Primary Bronchial/Tracheal Epithelial Cells (NHBE-A, ATCC® PCS-300-010™) were

plated at 6×10^3 cells/cm^2 on a layer of sub-lethally γ-irradiated (60 Gy) mouse embryonic fibroblasts

3T3-J2 cells (kind donation from Howard Green, Harvard Medical School, Boston[60]) with

keratinocyte culture medium cFAD (3:1 DMEM (Gibco) to Ham F-12 Nut Mix (Gibco), 10% FBS

(HyClone, Sigma), 1% Penicillin-Streptomycin (100X, Sigma), 0.4 μg/mL Hydrocortisone

(Calbiochem, USA), 5 μg/ml Insulin (various), 10-10 Cholera Toxin (Sigma) and 2x10-9

Triodothyronine (Sigma)). Epithelial cells were stimulated with 10 ng/mL human epidermal growth

factor (hEGF, PeproTech, USA) on days 3 and 5. Cultures were grown at 37°C 6% CO_2 and sub-

confluent cultures were trypsinised with 0.25% Trypsin-EDTA (Sigma). For NHBE-A differentiation

in Air Liquid Interface (ALI) culture, cells were plated at a density of 6×10^4/cm^2 on 1μm pore PET cell

culture inserts (12 wells, Greiner). The insert was supplemented with cFAD medium on the top and

bottom. After 48 hours, medium was removed from the top of the culture (exposure) and the culture

was fed with fresh PneumaCult™ differentiation Medium (Stemcell Technologies) from the bottom

of the well every 2 days for 3 weeks, until cell ciliation was observed. Antibiotics were removed from

the medium and cells washed 24 hours before infection experiments.

Human Epithelial Cell Lines

Human pharyngeal carcinoma Detroit 562 epithelial cells (ATCC_CCL-138) were grown in 10% FCS

in alpha MEM media (Gibco). For each set of experiments, cells were used within 5 passages. Cells

consistently tested PCR negative for mycoplasma (Cambridge Biosciences).

Pneumococcal-epithelial cell co-culture

For association and internalisation assays, confluent monolayers of Detroit 562 cells (typically day 7

- 8 post plating) cultured on 12 well plates (Corning), were exposed to *S. pneumoniae* (1 cell : 10

pneumococci) for three hours in 1% FCS alpha MEM (six-hour infection for HNBE-A cells in 1%

Pneumocult media without supplements) (3). Cells washed three times in HBSS$^{+/+}$ before incubation

in 1% saponin for 10 minutes at 37°C to lyse cells. Bacterial dilutions were plated on horse blood

agar for 16 hours and colony forming units counted. For internalisation quantification, 200µg/ml

gentamicin was added to the cells for one hour before incubation with saponin.

For transmigration assays, Detroit 562 cells were plated onto 3µm pore, PET Transwell Inserts

(ThermoFisher) typically for 10 days (3). Media was changed to 1% FCS (250µl apical chamber, 1ml

basal chamber) one hour prior to infection experiments. 1mg/ml FITC-dextran (Sigma Aldrich) was

added to the apical chamber of selected inserts for permeability assays. Transepithelial Electrical

Resistance (TEER) was measured before and after infection using an EVOM2 (World Precision

Instruments). Bacteria were added to the cells for three hours during of which 50µl of basal media

was repeatedly removed to measure bacterial load by counting CFU/well. Permeability was recorded

using a FLUOstar Omega (BMG Labtech) at 488nm.

Confocal Microscopy

Sample processing was performed as previously described(3). Briefly, mucosal cells from the EHPC

model were placed directly into 4% PFA and cytospun onto microscope slides. NHBE-A cells and

Detroit 562 cells cultured on transwell membranes were fixed in 4% PFA (Pierce, Methanol Free).

Cells were permeabilised with 0.2% Triton X-100 for 10 minutes, blocked for one hour in blocking

buffer (3% goat serum and 3% BSA in PBS (Sigma)) and incubated with pneumococcal antisera

Pool Q (raised against the polysaccharide capsule, SSI Diagnostica) for one hour. Wheat germ

agglutinin conjugated to Rhodamine (Vector Labs) was added along with goat-anti rabbit IgG

conjugated to Alexa Fluor 488 for 45 minutes in blocking buffer. Primary (used at 1:100) antibodies

used for epithelial cell proteins can be found in Supplemental Information Table 3. Secondary (used

at 1:500) antibodies were either goat anti-mouse or goat anti-rabbit – Alexa Fluor 555 or 546,

ThermoFisher. Of note, for epithelial protein antibodies raised in rabbit (Uteroglobin), no

pneumococcal staining was possible and therefore pneumococci were not visible on the images.

DAPI was added for 5 minutes and cells were mounted with Aqua PolyMount (VWR International)

and a coverslip on the microslide. Pneumococci were recorded by manual scanning of the whole

cytospin for the EHPC samples. More than three fields of view were recorded for each bacterial

strain for Detroit 562 cell infections, using inverted Zeiss LSM (either 700 or 880) confocal

microscopes. Images were processed with Zeiss LSM Image Browser. Z stacks were recorded at

1μm intervals at either 40x oil or 63x oil objectives. Automated image analysis was carried out using

Cellprofiler (61).

Electron Microscopy

Preparation of Detroit 562 cells and NHBE-A cells was identical, and previously described(3). Briefly,

cells cultured on transwells were fixed with 2% paraformaldehyde and 1.5% glutaraldehyde in 0.1 M

cacodylate buffer and post-fixed in 1% OsO_4 / 1.5% $K4Fe(CN)6$ in 0.1 M cacodylate buffer pH7.3.

Samples were dehydrated in graded ethanol-water series and infiltrated with Agar 100 resin mix.

Ultra-thin sections were cut at 70-80 nm using a diamond knife on a Reichert ultracut microtome.

Sections were collected on 300 mesh copper grids and stained with lead citrate. For transmission

electron microscopy the samples were hardened at 60°C for 48 hours. The coverslip was removed,

and a representative area was selected. Ultra-thin sections were cut at 70-80 nm using a diamond

knife on a Reichert ultra-cut S microtome. Sections were collected on copper grids and stained with

lead citrate. Samples were viewed with a Joel 1010 transition electron microscope and the images

recorded using a Gatan Orius CCD camera.

Pneumolysin activity

Adapted from Kirkham et al(62). Briefly, final concentration of 2% horse blood (EO Labs) solution

was prepared in phenol free alpha MEM (Life Technologies) and 100 μl added to wells of a 96 U-

bottom plate (Corning). 100 μl of bacterial solutions (containing ~2.5×10^6 bacteria) were suspended

in phenol free alpha MEM were added to each 96 well. Saponin was used as a positive control for

lysis. Solutions were incubated for 30 minutes at 37°C before centrifugation at 1000g for one minute.

100 µl was transferred into a new 96 well plate and absorbance recorded at 540nm on a plate reader.

Hydrogen Peroxide Assay

Detroit 562 cells were cultured in white 96 well plates (Corning). NHBE-A cells were cultured in 12

well plates in Pneumocult media. Cells were incubated with H_2O_2 substrate and infected with strains

of *S. pneumoniae* for up to six hours according to manufacturers' instructions (ROS-Glo™ H_2O_2

Assay, Promega). ROS-Glo™ Detection solution was added to the cells and incubated for 20

minutes. For primary cells, supernatant was then transferred to a white 96 well plate. Luminescence

was read using a GloMax -Multi Detection System plate reader (Promega).

Caspase assay

Detroit 562 cells were cultured in white 96 well plates (Corning). NHBE-A cells were cultured in 12

well plates in Pneumocult media. Cells were infected with strains of *S. pneumoniae* for up to six

hours. Primary cell supernatant was transferred to a white 96 well plate. Samples were incubated

with Caspase-Glo® 1 Reagent, Caspase-Glo® 1 YVAD-CHO Reagent, Caspase-Glo® Z-VAD-FMK

Reagent or Caspase-Glo® 8 Reagent, for one hour and according to manufacturers' instructions

(Promega). Luminescence was read using a GloMax-Multi Detection System plate reader

(Promega).

RNA samples and sequencing (RNA-Seq)

RNA-Seq data was collected as previously described (3). Confluent monolayers of Detroit 562

epithelial cells were infected with S. pneumoniae strains for four hours before collection in RNALater.

RNA was extracted using a Qiagen RNAEasy micro kit and treated for DNA removal using a Qiagen

Turbo DNA-free kit. RNA quality was assessed and quantified using a BioAnalyser (Agilent 2100).

Samples were sequenced using Illumina Nextseq 500/550 High Output 75 cycle kit giving 15-20

million 41bp paired-end reads per sample (Pathogens Genomic Unit, UCL).

Statistics

Each condition within each experiment was performed in duplicate or triplicate, and each experiment

was performed independently more than three times, unless stated otherwise. Error bars represent

SEM, unless stated otherwise. GraphPad Prism Version 7 was used for parametric (t-tests or

ANOVA) or non-parametric (Mann-Whitney or Kruskal-Wallis tests) analysis, which was based on

the Shapiro-Wilk normality test. Ad hoc tests were performed using Tukey's (for parametric data) or

Dunn's (for non-parametric data) multiple comparisons test. P values less than 0.05 were considered

significant.

Data and code availability

Source data are provided as a Source Data file. RNAseq data from the Detroit 562 cell infections is

publicly accessible through Zenodo with Accession (DOI:10.5281/zenodo.7997789).

ACKNOWLEDGEMENTS

This study was funded by CMW from a Postdoctoral Innovation Award from UCL, a Pump-Priming

Award and Enhancement Award from the Human Infection Challenge Network for Vaccine

Development (HIC-VAC (funded by the GCRF Networks in Vaccines Research and Development)

which is co-funded by the MRC and BBSRC). This UK funded award is part of the EDCTP2

programme supported by the European Union. CMW and RSH are supported by the Medical

Research Council (grant MR/T016329/1). The authors would like to thank Prof. Peter Hermans

(Radboud University Medical Centre) for donating the 6B pneumococcal strain originally from Prof.

Birgitta Henriques-Normark (Karolinska Institute) and obtained directly from DF at LSTM. EHPC

sample collection was supported by the Medical Research Council (grant MR/M011569/1), Bill and

Melinda Gates Foundation (grant OPP1117728) awarded to DF and the National Institute for Health

Research (NIHR) Local Comprehensive Research Network. The authors also wish to thank the

EHPC Clinical Team at LSTM and all the volunteers. Confocal imaging facilities at LSTM were

funded by a Wellcome Trust Multi-User Equipment Grant (104936/Z/14/Z). Electron microscopy

processing and image capture was performed by Dr. Elizabeth Slavik-Smith in the Biosciences EM

facility at UCL. MC is supported by a Springboard to Independence grant (AirwayStasis) from the

French Government's Investissement d'Avenir program, the Laboratoire d'Excellence "Integrative

Biology of Emerging Infectious Diseases" (ANR-10-LABX-62-IBEID) and the Chromatin and infection

unit headed by Melanie A. Hamon. RNA-Seq library preparation was undertaken at UCL through the

UCL/UCLH Biomedical Research Centre and MRC funded Pathogen Genomics Unit (PGU,

G0900950). All RNAseq data processing was completed by the PGU. MN is supported by the

Wellcome Trust and by NIHR Biomedical Research Centre Funding to University College Hospitals

NHS Foundation Trust and University College London. RR is supported by Marie Skłodowska-Curie

Individual Fellowships 896014. PB is supported by the European Research Council (ERC-Stg No.

639429), the Rosetrees Trust (M362-F1; M553), the CF Trust (SRC006;SRC020), the London

Advanced Therapies - Research England (C2N-AT.006), the MRC Confidence in Concept scheme

(Award: MC_PC_17180), the Innovate UK (Smart grant n.10005465), the Duchenne Parent Project

(DPP) and the NIHR GOSH BRC. MW is supported by a 223065 Wellcome Investigator Award to

Clare Jolly, UCL. ERS and JSB are supported by MRC grant R/N02687X/1 and Wellcome grant

221803/Z/20/Z. JSB is partially supported by the Department of Health's National Institute of Health

and Care Research Biomedical Research Centre funding to UCLH. RSH is a NIHR Senior

Investigator.

AUTHOR CONTRIBUTIONS: CMW conceived, funded, designed and conducted the study,

performed the experiments, acquired samples and processed data, and wrote the first draft of the

manuscript. GP analysed and interpreted the RNAseq data. MB analysed and interpreted the

bacterial transcriptomic data. RR cultured and differentiated primary epithelial cells. ERS generated

the 6B mutant strains and bacterial transcriptomic data. JR collected and processed EHPC samples.

JAGA processed the RNA-Seq data. MW ran the image analysis. MC provided technical advice. PB

and CJ provided investigator support. MN provided RNA-Seq analysis support. DF provided

investigator support and EHPC clinical samples. JSB generated the 6B mutant strains, interpreted

the data and revised the manuscript. RSH provided investigator support, designed and interpreted

the data, revised the manuscript. All authors commented on and approved the manuscript.

COMPETING INTERESTS: The authors declare no competing interests.

REFERENCES

1.　Weiser JN, Ferreira DM, Paton JC. 2018. Streptococcus pneumoniae: transmission, colonization and
invasion. Nat Rev Microbiol 16:355-367.
2.　O'Brien KL, Wolfson LJ, Watt JP, Henkle E, Deloria-Knoll M, McCall N, Lee E, Mulholland K, Levine OS,
Cherian T, Hib, Pneumococcal Global Burden of Disease Study T. 2009. Burden of disease caused by
Streptococcus pneumoniae in children younger than 5 years: global estimates. Lancet 374:893-902.
3.　Weight CM, Venturini C, Pojar S, Jochems SP, Reine J, Nikolaou E, Solorzano C, Noursadeghi M, Brown
JS, Ferreira DM, Heyderman RS. 2019. Microinvasion by Streptococcus pneumoniae induces
epithelial innate immunity during colonisation at the human mucosal surface. Nat Commun 10:3060.
4.　Zafar MA, Hamaguchi S, Zangari T, Cammer M, Weiser JN. 2017. Capsule Type and Amount Affect
Shedding and Transmission of Streptococcus pneumoniae. mBio 8.
5.　Gladstone RA, Lo SW, Lees JA, Croucher NJ, van Tonder AJ, Corander J, Page AJ, Marttinen P, Bentley
LJ, Ochoa TJ, Ho PL, du Plessis M, Cornick JE, Kwambana-Adams B, Benisty R, Nzenze SA, Madhi SA,
Hawkins PA, Everett DB, Antonio M, Dagan R, Klugman KP, von Gottberg A, McGee L, Breiman RF,
Bentley SD, Global Pneumococcal Sequencing C. 2019. International genomic definition of
pneumococcal lineages, to contextualise disease, antibiotic resistance and vaccine impact.
EBioMedicine 43:338-346.
6.　Amin-Chowdhury Z, Collins S, Sheppard C, Litt D, Fry NK, Andrews N, Ladhani SN. 2020.
Characteristics of Invasive Pneumococcal Disease Caused by Emerging Serotypes After the
Introduction of the 13-Valent Pneumococcal Conjugate Vaccine in England: A Prospective
Observational Cohort Study, 2014-2018. Clin Infect Dis 71:e235-e243.
7.　Echlin H, Frank M, Rock C, Rosch JW. 2020. Role of the pyruvate metabolic network on carbohydrate
metabolism and virulence in Streptococcus pneumoniae. Mol Microbiol 114:536-552.
8.　Hartel T, Eylert E, Schulz C, Petruschka L, Gierok P, Grubmuller S, Lalk M, Eisenreich W,
Hammerschmidt S. 2012. Characterization of central carbon metabolism of Streptococcus
pneumoniae by isotopologue profiling. J Biol Chem 287:4260-74.
9.　Henriques-Normark B, Tuomanen EI. 2013. The pneumococcus: epidemiology, microbiology, and
pathogenesis. Cold Spring Harb Perspect Med 3.
10.　Kohler S, Voss F, Gomez Mejia A, Brown JS, Hammerschmidt S. 2016. Pneumococcal lipoproteins
involved in bacterial fitness, virulence, and immune evasion. FEBS Lett 590:3820-3839.
11.　van Opijnen T, Camilli A. 2012. A fine scale phenotype-genotype virulence map of a bacterial
pathogen. Genome Res 22:2541-51.
12.　Ramos-Sevillano E, Ercoli G, Felgner P, Ramiro de Assis R, Nakajima R, Goldblatt D, Heyderman RS,
Gordon SB, Ferreira DM, Brown JS. 2020. Preclinical Development of Virulence Attenuated
Streptococcus pneumoniae Strains Able to Enhance Protective Immunity Against Pneumococcal
Infection. Am J Respir Crit Care Med doi:10.1164/rccm.202011-4161LE.
13.　Ogunniyi AD, Mahdi LK, Trappetti C, Verhoeven N, Mermans D, Van der Hoek MB, Plumptre CD, Paton
JC. 2012. Identification of genes that contribute to the pathogenesis of invasive pneumococcal
disease by in vivo transcriptomic analysis. Infect Immun 80:3268-78.

14. Mahdi LK, Wang H, Van der Hoek MB, Paton JC, Ogunniyi AD. 2012. Identification of a novel
pneumococcal vaccine antigen preferentially expressed during meningitis in mice. J Clin Invest
122:2208-20.
15. Brill J, Hoffmann T, Bleisteiner M, Bremer E. 2011. Osmotically controlled synthesis of the compatible
solute proline is critical for cellular defense of Bacillus subtilis against high osmolarity. J Bacteriol
193:5335-46.
16. Crowley PJ, Gutierrez JA, Hillman JD, Bleiweis AS. 1997. Genetic and physiologic analysis of a formyl-
tetrahydrofolate synthetase mutant of Streptococcus mutans. J Bacteriol 179:1563-72.
17. Ramos-Sevillano E. EG, Guerra-Assunção J.F., Betts M., Partridge F., Fernandes V.E., Wall E., Gordon
S.B., Ferreira D.M., Heyderman R.S., Brown J.S. 2023. Essential role of proline synthesis and the one-
carbon metabolism pathways for systemic virulence of Streptococcus pneumoniae. Biorxiv
doi:https://doi.org/10.1101/2023.08.03.550501.
18. Brown JS, Gilliland SM, Holden DW. 2001. A Streptococcus pneumoniae pathogenicity island
encoding an ABC transporter involved in iron uptake and virulence. Mol Microbiol 40:572-85.
19. Hill H, Mitsi E, Nikolaou E, Blizard A, Pojar S, Howard A, Hyder-Wright A, Devin J, Reine J, Robinson R,
Solorzano C, Jochems SP, Kenny-Nyazika T, Ramos-Sevillano E, Weight CM, Myerscough C,
McLenaghan D, Morton B, Gibbons E, Farrar M, Randles V, Burhan H, Chen T, Shandling AD, Campo
JJ, Heyderman RS, Gordon SB, Brown JS, Collins AM, Ferreira DM. 2023. A Randomised Controlled
Trial of Nasal Immunisation with Live Virulence Attenuated Streptococcus pneumoniae Strains Using
Human Infection Challenge. Am J Respir Crit Care Med doi:10.1164/rccm.202302-0222OC.
20. Hill H ME, Nikolaou E, Blizard A, Pojar S, Howard A, Hyder-Wright A, Devin J, Reiné J, Robinson R,
Solórzano C, Jochems S, Kenny-Nyazika T, Ramos-Sevillano E, Weight CM, Myerscough C, McLennan
D, MortonB , Gibbons E, Farrar M, Randles V, Burhan H, Chen T, Shandling AD, Campo JJ, Heyderman
RS, Gordon SB, Brown JS, Collins AM, Ferreira DM. 2023. A Randomised Controlled Trial of Nasal
Immunisation with Live Virulence Attenuated Streptococcus pneumoniae Strains using Human
Infection Challenge. Medrxiv doi:https://doi.org/10.1101/2023.04.14.23288224.
21. Ferreira DM, Neill DR, Bangert M, Gritzfeld JF, Green N, Wright AK, Pennington SH, Bricio-Moreno L,
Moreno AT, Miyaji EN, Wright AD, Collins AM, Goldblatt D, Kadioglu A, Gordon SB. 2013. Controlled
human infection and rechallenge with Streptococcus pneumoniae reveals the protective efficacy of
carriage in healthy adults. Am J Respir Crit Care Med 187:855-64.
22. Eshun-Wilson L, Zhang R, Portran D, Nachury MV, Toso DB, Lohr T, Vendruscolo M, Bonomi M, Fraser
JS, Nogales E. 2019. Effects of alpha-tubulin acetylation on microtubule structure and stability. Proc
Natl Acad Sci U S A 116:10366-10371.
23. Iliev AI, Djannatian JR, Opazo F, Gerber J, Nau R, Mitchell TJ, Wouters FS. 2009. Rapid microtubule
bundling and stabilization by the Streptococcus pneumoniae neurotoxin pneumolysin in a
cholesterol-dependent, non-lytic and Src-kinase dependent manner inhibits intracellular trafficking.
Mol Microbiol 71:461-77.
24. Surabhi S, Jachmann LH, Shumba P, Burchhardt G, Hammerschmidt S, Siemens N. 2022. Hydrogen
Peroxide Is Crucial for NLRP3 Inflammasome-Mediated IL-1beta Production and Cell Death in
Pneumococcal Infections of Bronchial Epithelial Cells. J Innate Immun 14:192-206.
25. Maekawa T, Kashkar H, Coll NS. 2023. Dying in self-defence: a comparative overview of immunogenic
cell death signalling in animals and plants. Cell Death Differ 30:258-268.
26. Gunther C, Buchen B, He GW, Hornef M, Torow N, Neumann H, Wittkopf N, Martini E, Basic M, Bleich
A, Watson AJ, Neurath MF, Becker C. 2015. Caspase-8 controls the gut response to microbial
challenges by Tnf-alpha-dependent and independent pathways. Gut 64:601-10.
27. Gonzalez-Juarbe N, Bradley KM, Shenoy AT, Gilley RP, Reyes LF, Hinojosa CA, Restrepo MI, Dube PH,
Bergman MA, Orihuela CJ. 2017. Pore-forming toxin-mediated ion dysregulation leads to death
receptor-independent necroptosis of lung epithelial cells during bacterial pneumonia. Cell Death
Differ 24:917-928.
28. Gonzalez-Juarbe N, Bradley KM, Riegler AN, Reyes LF, Brissac T, Park SS, Restrepo MI, Orihuela CJ.
2018. Bacterial Pore-Forming Toxins Promote the Activation of Caspases in Parallel to Necroptosis to
Enhance Alarmin Release and Inflammation During Pneumonia. Sci Rep 8:5846.

29. Erttmann SF, Gekara NO. 2019. Hydrogen peroxide release by bacteria suppresses inflammasome-
dependent innate immunity. Nat Commun 10:3493.
30. Alibayov B, Scasny A, Khan F, Creel A, Smith P, Vidal AGJ, Fitisemanu FM, Padilla-Benavides T, Weiser
JN, Vidal JE. 2022. Oxidative Reactions Catalyzed by Hydrogen Peroxide Produced by Streptococcus
pneumoniae and Other Streptococci Cause the Release and Degradation of Heme from Hemoglobin.
Infect Immun 90:e0047122.
31. Zafar MA, Wang Y, Hamaguchi S, Weiser JN. 2017. Host-to-Host Transmission of Streptococcus
pneumoniae Is Driven by Its Inflammatory Toxin, Pneumolysin. Cell Host Microbe 21:73-83.
32. Periselneris J, Turner CT, Ercoli G, Szylar G, Weight CM, Thurston T, Whelan M, Tomlinson G,
Noursadeghi M, Brown J. 2022. Pneumolysin suppresses the initial macrophage pro-inflammatory
response to Streptococcus pneumoniae. Immunology 167:413-427.
33. Terra VS, Plumptre CD, Wall EC, Brown JS, Wren BW. 2020. Construction of a pneumolysin deficient
mutant in streptococcus pneumoniae serotype 1 strain 519/43 and phenotypic characterisation.
Microb Pathog 141:103999.
34. Bazant J, Ott B, Hudel M, Hain T, Lucas R, Mraheil MA. 2023. Impact of Endogenous Pneumococcal
Hydrogen Peroxide on the Activity and Release of Pneumolysin. Toxins (Basel) 15.
35. Travaglini KJ, Nabhan AN, Penland L, Sinha R, Gillich A, Sit RV, Chang S, Conley SD, Mori Y, Seita J,
Berry GJ, Shrager JB, Metzger RJ, Kuo CS, Neff N, Weissman IL, Quake SR, Krasnow MA. 2020. A
molecular cell atlas of the human lung from single-cell RNA sequencing. Nature 587:619-625.
36. Herta T, Bhattacharyya A, Bollensdorf C, Kabus C, Garcia P, Suttorp N, Hippenstiel S, Zahlten J. 2018.
DNA-release by Streptococcus pneumoniae autolysin LytA induced Krueppel-like factor 4 expression
in macrophages. Sci Rep 8:5723.
37. Barutcu AR, Lajoie BR, Fritz AJ, McCord RP, Nickerson JA, van Wijnen AJ, Lian JB, Stein JL, Dekker J,
Stein GS, Imbalzano AN. 2016. SMARCA4 regulates gene expression and higher-order chromatin
structure in proliferating mammary epithelial cells. Genome Res 26:1188-201.
38. Wilson R, Cohen JM, Jose RJ, de Vogel C, Baxendale H, Brown JS. 2015. Protection against
Streptococcus pneumoniae lung infection after nasopharyngeal colonization requires both humoral
and cellular immune responses. Mucosal Immunol 8:627-39.
39. Periselneris J, Ercoli G, Pollard T, Chimalapati S, Camberlein E, Szylar G, Hyams C, Tomlinson G,
Petersen FC, Floto RA, Noursadeghi M, Brown JS. 2019. Relative Contributions of Extracellular and
Internalized Bacteria to Early Macrophage Proinflammatory Responses to Streptococcus
pneumoniae. mBio 10.
40. Belitsky BR, Brill J, Bremer E, Sonenshein AL. 2001. Multiple genes for the last step of proline
biosynthesis in Bacillus subtilis. J Bacteriol 183:4389-92.
41. Sah S, Aluri S, Rex K, Varshney U. 2015. One-carbon metabolic pathway rewiring in Escherichia coli
reveals an evolutionary advantage of 10-formyltetrahydrofolate synthetase (Fhs) in survival under
hypoxia. J Bacteriol 197:717-26.
42. Aluri S, Rex K, Varshney U. 2015. Simultaneous presence of fhs and purT genes is disadvantageous
for the fitness of Escherichia coli growth. FEMS Microbiol Lett 362.
43. Obolski U, Swarthout TD, Kalizang'oma A, Mwalukomo TS, Chan JM, Weight CM, Brown C, Cave R,
Cornick J, Kamng'ona AW, Msefula J, Ercoli G, Brown JS, Lourenco J, Maiden MC, French N, Gupta S,
Heyderman RS. 2023. The metabolic, virulence and antimicrobial resistance profiles of colonising
Streptococcus pneumoniae shift after PCV13 introduction in urban Malawi. Nat Commun 14:7477.
44. Galan JE. 2021. Salmonella Typhimurium and inflammation: a pathogen-centric affair. Nat Rev
Microbiol 19:716-725.
45. Li G, Wang G, Li M, Li L, Liu H, Sun M, Wen Z. 2020. Morin inhibits Listeria monocytogenes virulence
in vivo and in vitro by targeting listeriolysin O and inflammation. BMC Microbiol 20:112.
46. Connor V, German E, Pojar S, Mitsi E, Hales C, Nikolaou E, Hyder-Wright A, Adler H, Zaidi S, Hill H,
Jochems SP, Burhan H, French N, Tobery T, Rylance J, Ferreira DM. 2018. Hands are vehicles for
transmission of Streptococcus pneumoniae in novel controlled human infection study. Eur Respir J
52.
47. Lourenco J, Daon Y, Gori A, Obolski U. 2021. Pneumococcal Competition Modulates Antibiotic
Resistance in the Pre-Vaccination Era: A Modelling Study. Vaccines (Basel) 9.

48. Watkins ER, Kalizang'Oma A, Gori A, Gupta S, Heyderman RS. 2022. Factors affecting antimicrobial
resistance in Streptococcus pneumoniae following vaccination introduction. Trends Microbiol
30:1135-1145.
49. Chaguza C, Senghore M, Bojang E, Gladstone RA, Lo SW, Tientcheu PE, Bancroft RE, Worwui A, Foster-
Nyarko E, Ceesay F, Okoi C, McGee L, Klugman KP, Breiman RF, Barer MR, Adegbola RA, Antonio M,
Bentley SD, Kwambana-Adams BA. 2020. Within-host microevolution of Streptococcus pneumoniae
is rapid and adaptive during natural colonisation. Nat Commun 11:3442.
50. Browall S, Norman M, Tangrot J, Galanis I, Sjostrom K, Dagerhamn J, Hellberg C, Pathak A, Spadafina
T, Sandgren A, Battig P, Franzen O, Andersson B, Ortqvist A, Normark S, Henriques-Normark B. 2014.
Intraclonal variations among Streptococcus pneumoniae isolates influence the likelihood of invasive
disease in children. J Infect Dis 209:377-88.
51. S. A. 2010. FastQC: a quality control tool for high throughput sequence data.
52. Ewels P, Magnusson M, Lundin S, Kaller M. 2016. MultiQC: summarize analysis results for multiple
tools and samples in a single report. Bioinformatics 32:3047-8.
53. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics 30:2114-20.
54. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9:357-9.
55. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome
Project Data Processing S. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics
25:2078-9.
56. Nicol JW, Helt GA, Blanchard SG, Jr., Raja A, Loraine AE. 2009. The Integrated Genome Browser: free
software for distribution and exploration of genome-scale datasets. Bioinformatics 25:2730-1.
57. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. Bioinformatics 30:923-30.
58. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. Genome Biol 15:550.
59. Yu G, Wang LG, Han Y, He QY. 2012. clusterProfiler: an R package for comparing biological themes
among gene clusters. OMICS 16:284-7.
60. Rheinwald JG, Green H. 1975. Serial cultivation of strains of human epidermal keratinocytes: the
formation of keratinizing colonies from single cells. Cell 6:331-43.
61. Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, Guertin DA, Chang JH, Lindquist
RA, Moffat J, Golland P, Sabatini DM. 2006. CellProfiler: image analysis software for identifying and
quantifying cell phenotypes. Genome Biol 7:R100.
62. Kirkham LA, Kerr AR, Douce GR, Paterson GK, Dilts DA, Liu DF, Mitchell TJ. 2006. Construction and
immunological characterization of a novel nontoxic protective pneumolysin mutant for use in future
pneumococcal vaccines. Infect Immun 74:586-93.

FIGURE LEGENDS

Table 1. Differentially expressed pathways from single isogenic mutants in human

serum compared to WT. 55 genes were selected within competence pathways, amino acid

metabolism, virulence factors, oxidative stress pathways and other known genes important

for colonisation, for assessment of changes in gene expression following serum stress. *q-*

value measures the proportion of false positives incurred (the false discovery rate) when

that particular test is significant.

Fig 1. Pneumococcal – epithelial interactions in the EHPC model. (A) Schematic showing that

during colonisation, pneumococci adhere to the epithelial cell surface (association), internalise inside

epithelial cells (microinvasion). Mucus – lime; pneumococci – purple; intracellular vacuole – dark

green; epithelial cells – yellow; nucleus – brown; cell junctions – pink. (B) Representative microscopy

images showing 6B association to the epithelium and microinvasion by the $\Delta proABC/piaA$ or

$\Delta fhs/piaA$ mutants. Surface carbohydrates (red), nuclei (DAPI, blue), pneumococci (green). (C)

Table summarising the percentage of bacteria that were found by microscopy, on the surface

(association) or inside (microinvasion) cells in the EHPC model. Mucosal curettage samples were

collected at day 6 post-pneumococcal challenge. Number of volunteers (n) in each group is indicated

in brackets, along with the total numbers of pneumococci (Spn) counted.

Fig 2. Impact of pneumococcal infection on NHBE-A cell junctional protein expression and

barrier integrity. NHBE-A Cells were cultured at ALI and infected with pneumococcal strains for

6hrs. (A) Top: Representative confocal microscopy images from 4 independent experiments with

replicates, showing pneumococcal (green) association with the epithelium (red, surface

carbohydrates; nuclei blue) at 6hr post infection. Bottom: Representative transmission electron

microscopy images from 2 independent experiments, showing pneumococcal microinvasion

examples of each strain (arrows), enclosed within a vacuole after 4hr infection. (B) Representative

confocal microscopy images from 4 independent experiments with replicates, showing

pneumococcal (green) interactions with junctional proteins (red) of β catenin. Colocalization of red

and green appears yellow. Nuclei (blue). (C) Integrated intensity of β catenin in pneumococcal

negative (not pneumococcal associated) or positive (pneumococcal associated) cells. Non infected

v pneumococcal negative cells **** $p <0.0001$ (NI v $\Delta proABC/piaA$), ** $p = 0.0026$ (NI v $\Delta fhs/piaA$);

non-infected v pneumococcal positive cells; **** $p <0.0001$ (NI v $\Delta proABC/piaA$ and NI v $\Delta fhs/piaA$);

Kruskal-Wallis, n = 4 independent experiments. (D) Transepithelial electrical resistance (TEER) was

recorded pre- and post- 6hr infection to assess barrier integrity. N = 3 independent experiments with

replicates. Average TEER pre-infection was 466Ω.cm^2 (430-484 Ω.cm^2, p = 0.1458 ANOVA) and

average TEER post-infection was 545Ω.cm^2 (527-567 Ω.cm^2. p = 0.04744 ANOVA). Change in

TEER comparing non-infected cells to infected cells p = 0.6592, Kruskal-Wallis. Error bars represent

SEM.

Fig 3. Pneumococcal – epithelial interactions with primary human epithelial cells. Cells were

cultured at ALI and infected with pneumococcal strains for 6hrs. (A) Representative scanning

electron microscopy images from 2 independent experiments, showing the surface of the epithelium.

Representative examples of differentiated epithelia (basal cells, Clara cells, goblet cells, ciliated

cells) are shown in yellow text. Colonising pneumococci are highlighted by arrows. Representative

confocal microscopy images from 4 independent experiments with replicates for (B) Ciliated cells

(acetylated tubulin, red), (C) Goblet cells (mucin5ac, red, (arrows indicate co-localisation of

mucin5ac with pneumococci)), (D) Clara cells (uteroglobin, red) after 6hr infection. Pneumococci

(green), nuclei (DAPI, blue), colocalization of red and green appears as yellow. (E) Integrated

intensity of acetylated tubulin and Muc5ac in pneumococcal negative or positive cells from two or

more images. Acetylated tubulin, neg p = 0.2677, pos p = 0.017; Mucin5ac, neg p = 0.6852, pos p =

0.8510, ANOVA. N = 3 or 4 independent experiments. (F) Integrated intensity of uteroglobin

normalised to non-infected cells (left, p = 0.0115 non-infected v infected, Kruskal-Wallis) and

percentage of Clara positive cells following infection (right, p = 0.0003 non-infected v strains, p =

0.0036 6B v mutant strains, Kruskal-Wallis). N = 3 independent experiments.

Fig 4. **Strain association, microinvasion and transmigration of Detroit 562 cells.** Cells were

infected with *S. pneumoniae* for three hours and CFU counts recorded. (A) Representative confocal

microscopy images from 6 independent experiments with replicates, showing pneumococcal (green)

interactions with surface carbohydrates (red). (B) Association of pneumococcal strains to epithelial

cells. p = 0.0009 (Kruskal-Wallis), n = 6 with replicates. (C) Internalisation of strains into cells,

following a gentamicin protection assay. $p = 0.0219$ (Kruskal-Wallis) N = 6 with replicates. (D)

Microinvasive index is defined as the percentage of internalised bacteria over all associated bacteria.

$p = 0.071$ (Kruskal-Wallis) N = 6 with replicates. (E) Pneumococcal viability inside epithelial cells

following a gentamicin protection assay. n = 3 or 4 independent experiments with replicates. Counts

were normalised to microinvasion counts for each strain after 3 hours infection and compared to 6B

WT; +1hr $p = 0.6517$, +2hr $p = 0.1377$, +3hr $p = 0.1183$ (ANOVA). (F) Transmigration of

pneumococcal strains across confluent monolayers on transwell inserts. CFUs were recorded from

the basal chamber. N = 6 independent experiments with replicates. $p < 0.001$ (Kruskal-Wallis). (G)

Epithelial cell permeability was assessed via leak to 4kDa FITC-Dextran into the basal chamber. N

= 4 with replicates. $p < 0.0001$ compared to non-infected cells, and $p = 0.0689$ between 6B WT and

mutant strains (Kruskal-Wallis). (H) Representative confocal microscopy images from 6 independent

experiments with replicates, showing pneumococcal (green) interactions with β catenin (red).

Nuclear DAPI stain (blue). (I) Image quantification of integrated intensity of β catenin normalised to

non-infected cells, following infection with the different strains. N = 5 independent experiments. Non

infected v pneumococcal negative cells **** $p < 0.0001$ (NI v Δ*fhs/piaA*); non-infected v

pneumococcal positive cells; **** $p < 0.0001$ (NI v 6B), * $p = 0.0183$ (NI v Δ*proABC/piaA*), * $p = 0.0149$

(NI v Δ*fhs/piaA*); Kruskal-Wallis.

Fig 5. Detroit 562 cell transcriptomic responses to pneumococcal strains. Epithelial cells were

infected for four hours with pneumococcal strains and RNA extracted and processed for RNAseq.

6B n = 11, PP n = 17, FP n = 14, non-infected n = 12, from 6 independent experiments. (A) All

significantly log2 fold change upregulated genes according to strain compared to non-infected cells.

Significance was defined as DEGs with a log2 fold change (logFC) of > 1.5 and false discovery rate

(FDR) of < 0.05. (B) Comparison of the top 20 most upregulated genes for each strain, compared to

non-infected cells. Upstream regulator analysis for (C) transcription factors and (D) kinases.

Fig 6. Effects of pneumococcal infection on caspase activity in Detroit 562 cells. Caspase

activity was assessed by luminescence, following infection with pneumococcal strains for 6 hours.

(A) Caspase 3/7 activity was determined following incubation with the specific inhibitor Z-VAD-FMK.

N = 4 with replicates. In comparison to non-infected cells, caspase 3/7 p = 0.0022; caspase 3/7 + Z-

VAD-FMK inhibitor p = 0.0005. In comparison to 6B WT, caspase 3/7 p = 0.0215; caspase 3/7 + Z-

VAD-FMK inhibitor p = 0.0012 (Kruskal-Wallis). (B) Caspase 1 activity was determined following

incubation with the specific inhibitor YVAD-CHO. Negative signals indicate caspase activity other

than caspase 1. N = 6 with replicates. In comparison to non-infected cells, p = 0.39; caspase+YVAD-

CHO inhibitor p = 0.3604 and, in comparison to 6B WT, p = 0.6226; caspase+YVAD-CHO inhibitor

p = 0.5386 (Kruskal-Wallis). (C) Caspase 8 activity was assessed following infection with

pneumococcal strains. N = 3 with replicates. In comparison to non-infected cells, p = 0.0015

(ANOVA) and, in comparison to 6B WT, p = 0.0172 (ANOVA).

Fig 7. Hydrogen peroxide secretion and pneumolysin activity from pneumococcal strains in

the absence and presence of Detroit 562 cells. A) Fold change of hydrogen peroxide production

from the mutants compared to WT, normalised to the CFU counts. 3hr p = 0.0087 (ANOVA), 6hr p

= 0.0209 (Kruskal-Wallis). B) Fold change of hydrogen peroxide production from the mutants

compared to WT, normalised to the CFU counts, following infection of Detroit 562 cells. Media was

replaced after 1hr infection and cells incubated for a further 5hrs. N = 3 independent experiments

with replicates. Compared to 6B WT, p = 0.0034 (ANOVA). C) Pneumolysin activity was assessed

using a red blood cell lysis assay. Strains were incubated with red blood cells for 30 minutes and

absorbance measured. N = 4 experiments with replicates. Pneumococcal strain haemolytic activity

is calculated in comparison to 100% lysis by 0.0625% saponin. p <0.0001 compared to 6B WT

(ANOVA).

Table 1. Differentially expressed pathways from single isogenic mutants in human serum compared to WT.

Pathway	GeneRatio	P-adjust	qvalue
Δ*fhs* vs. WT			
Fatty acid metabolism/biosynthesis	10/57	1.08E-09	6.49E-10
Biosynthesis of secondary metabolites	26/57	2.09E-04	1.26E-04
Purine metabolism	10/57	1.34E-03	8.06E-04
Quorum sensing	11/57	3.20E-03	1.92E-03
Pantothenate and CoA biosynthesis	5/57	3.59E-03	2.16E-03
beta-Lactam resistance	5/57	7.59E-03	4.57E-03
Propanoate metabolism	4/57	1.51E-02	9.09E-03
2-Oxocarboxylic acid metabolism	4/57	1.51E-02	9.09E-03
Metabolic pathways	40/57	1.94E-02	1.16E-02
Amino sugar and nucleotide sugar metabolism	7/57	4.22E-02	2.54E-02
Pyruvate metabolism	5/57	4.31E-02	2.59E-02
Δ*proABC* vs. WT			
Fatty acid metabolism/biosynthesis	10/71	1.17E-08	8.94E-09
Galactose metabolism	12/71	1.25E-04	2.47E-08
Phosphotransferase system (PTS)	13/71	2.54E-03	1.93E-03
Amino sugar and nucleotide sugar metabolism	9/71	2.89E-02	2.20E-02

Figure 1. Pneumococcal – epithelial interactions in the EHPC model

A

Pneumococci

Association

Micro-invasion

Transmigration

B

6B | ΔproABC/piaA | Δfhs/piaA

C

	Day 6 post-challenge	
	Association (%)	Micro-invasion (%)
6B	87.8 (n11, 43spn)	12.2 (n11, 6spn)
ΔproABC/piaA	59.3 (n11, 16spn)	40.7 (n11, 11spn)
Δfhs/piaA	48.7 (n5, 57spn)	51.3 (n5, 60spn)

Figure 2. Impact of pneumococcal infection on NHBE-A cell junctional protein expression and barrier integrity.

Figure 3. Pneumococcal – epithelial interactions with primary human epithelial cells.

Figure 4. Strain association, microinvasion and transmigration of Detroit 562 cells.

Figure 5. Detroit 562 cell transcriptomic responses to pneumococcal strains.

A

6B WT v not infected   ΔproABC/piaA v not infected

81   193   38

78   13

Δfhs/piaA v not infected

B

log2foldchange	6B	ΔproABC/piaA	Δfhs/piaA
Upregulated genes			
AC256236.1	32.9	25.5	0.0
AC022149.1	30.2	27.6	0.0
Z84488.2	29.1	21.6	0.0
MUC21	25.8	29.0	0.0
VASN	20.9	0.0	0.0
BRD2	18.4	24.7	0.0
AC012254.2	18.2	13.8	0.0
CTAGE4	18.1	0.0	0.0
MICE	17.1	20.3	0.0
AC107075.1	10.1	0.0	0.0
CSNK2B	0.0	20.2	0.0
MTRNR2L6	0.0	19.4	15.8
LSM2	0.0	17.1	0.0
ALOX5	0.0	0.0	6.9
AC006064.6	0.0	0.0	3.4
TAP2	0.0	0.0	3.2
CYP1A1	0.0	0.0	1.7
FOSB	0.0	0.0	1.4
IL1B	0.0	0.0	1.3
CXCL8	0.0	0.0	1.1
CXCL2	0.0	0.0	1.1
CYP1B1	0.0	0.0	1.0

C

Transcription factors

0.00   2.01   7.00

CREB1
TP53
HIF1A
NUPR1
RELA
STAT3
NFKBIA
FOXO3
FOXO1
TGIF1
EZH2
JUN
SMARCA4
NFKB1
NFAT5
KLF6
TCF4

D

Kinases

0.00   0.73   4.00

MAPK1
PRKCA
SRC
TBK1
IKBKB
MAP3K8
MAPK8
EIF2AK3
RIPK1
MAP3K14
ERBB2
PRKCD
FGFR1
RAF1
JAK1
CHUK
MAPK7
PRKCE
MAPK3

Figure 6. Effects of pneumococcal infection on caspase activity in Detroit 562 cells.

Figure 7. Hydrogen peroxide secretion and pneumolysin activity from pneumococcal strains in the absence and presence of Detroit 562 cells.

Graphical abstract