POLARIS is a copper-binding peptide required for ethylene signalling control in *Arabidopsis*

Anna J. Mudge[1†], Saher Mehdi[1†‡], Will Michaels[1,2†], Beatriz Orosa-Puente[1‡‡], Weiran Shen[1], Charlie Tomlinson[2‡‡‡], Wenbin Wei[1], Claudia Hoppen[3], Buket Uzun[3], Dipan Roy[1], Flora M. Hetherington[1‡‡‡‡], Jennifer F. Topping[1], Ari Sadanandom[1], Georg Groth[3], Nigel J. Robinson[1,2*], Keith Lindsey[1*]

Affiliations

[1]*Department of Biosciences, Durham University, Durham DH1 3LE, UK*

[2]*Department of Chemistry, Durham University, Durham DH1 3LE, UK*

[3]*Institute of Biochemical Plant Physiology, Heinrich Heine University Du☐sseldorf, D-40204 Du☐sseldorf, Germany*

[†]These authors contributed equally to this work

[‡]Present address: wellOwise Research and Development, IIITD Innovation & Incubation Centre, New Delhi, Delhi - 110020, India.

[‡‡]Present address: School of Biological Sciences, University of Edinburgh, Edinburgh EH9 3FF, UK; and Centro Singular de Investigación en Química Biolóxica e Materiais Moleculares (CIQUS) and Departamento de Química Orgánica, 15782, Universidade de Santiago de Compostela, Spain

[‡‡‡]Present address: Department of Biology, University of York, Heslington, York YO10 5DD, U.K.

[‡‡‡‡]Present address: University of Bath, Claverton Down, Bath BA2 7AY, U.K.

*Joint corresponding authors: keith.lindsey@durham.ac.uk; nigel.robinson@durham.ac.uk

Running title:

POLARIS peptide regulates ethylene receptor function

ABSTRACT

Ethylene signalling represents one of the classic hormonal pathways in plants, with diverse roles in development and stress responses. The dimeric ethylene receptor localizes to the endoplasmic reticulum (ER) and contains Cu(I) ions essential for ethylene binding and signal transduction. We previously discovered that mutants in the *Arabidopsis* gene *POLARIS* (*PLS*), encoding a 36 amino acid peptide, exhibit enhanced ethylene signalling responses, suggestive of reduced receptor activity, but the role and activity of the peptide in this signalling cascade has not been defined. Here we report PLS binds copper as a 1:2 thiol-dependent $Cu(I):PLS_2$ complex, with an affinity of 3.79 ($\pm$1.5) $\times 10^{19}$ M^{-2}, via two cysteine residues also found in the related species *Camelina sativa*. These residues are also essential for biological function. This affinity precludes PLS as a cytosolic Cu chaperone. We demonstrate that PLS localizes to endomembranes and interacts with the transmembrane domain of receptor protein ETR1. PLS-ETR1 binding is increased in the presence of copper, and this interaction provides a Cu-dependent mechanism for mediating a repression of ethylene responses. *PLS* transcription is up-regulated by auxin and down-regulated by ethylene, and so PLS-ETR1 interactions also provide a mechanism to modulate ethylene responses in high auxin tissues.

Keywords:

Plant hormone signalling, Arabidopsis, Ethylene, Protein metallation, Hormone receptor

INTRODUCTION

Ethylene is used by plants as a gaseous hormone to regulate many aspects of development and responses to biotic and abiotic stresses (Johnson and Ecker, 1998). It is perceived by a family of receptors that, in *Arabidopsis*, comprises 5 members, ETR1 (ETHYLENE RESPONSE 1), ERS1 (ETHYLENE RESPONSE SENSOR 1), ERS2, ETR2, and EIN4 (ETHYLENE-INSENSITIVE 4) (Chang et al, 1993; Hua et al., 1995, 1998; Sakai et al., 1998) located on the endoplasmic reticulum (ER) (Chen et al., 2002; Grefen et al., 2008). The receptors are related to bacterial two-component systems (Chen et al., 2002), form dimers through disulphide bonding at the N-terminal hydrophobic domains (Schaller et al., 1995; Hall et al., 2000) and contain Cu(I) ions bound to residues Cys65 and His69, essential for ethylene binding and signal transduction (Rodriguez et al., 1999; McDaniel and Binder, 2012). In the absence of ethylene these receptors activate a negative regulator CTR1 (CONSTITUTIVE TRIPLE RESPONSE 1), which is a mitogen-activated protein kinase kinase kinase (MAPKKK), so preventing ethylene responses (Chang, 2003; Gao et al., 2003). Mechanisms by which receptor activity is regulated are not fully understood.

Introduction of copper to the ER and ethylene receptor requires the RAN1 (RESPONSIVE TO ANTAGONIST1) protein. This is a predicted copper-transporting P-type ATPase homologous to the yeast Ccc2p and to human Menkes and Wilson disease proteins (Hirayama et al., 1999). Strong loss-of-function mutants of *RAN1* in *Arabidopsis* (e.g. *ran1-3*, *ran1-4*) exhibit an enhanced ethylene signalling response (Binder et al., 2010), consistent with a loss of receptor function, and similar to higher order loss-of-function receptor mutants, which also show an ethylene hypersignalling phenotype (Qu et al., 2007). Mechanisms of copper homeostasis at ETR1 are unknown, as is true for other compartmentalized cuproteins supplied with copper, for example via Ccc2p, Menkes or Wilson ATPases. RAN1 in Arabidopsis localizes to endomembrane systems, including *trans*-Golgi and ER compartments, and is necessary for both the biogenesis of ethylene receptors and copper homeostasis - loss of function *ran1* mutants suggest copper is required for both ethylene binding and receptor function (Binder et al., 2010). RAN1 can interact directly with ETR1 and the copper chaperones ANTIOXIDANT1 (ATX1) and COPPER CHAPERONE (CCH), suggestive of a transfer of copper between proteins to deliver it to the ethylene

receptor as part of the receptor biogenesis pathway at the ER (Hoppen et al., 2019).

Our understanding of receptor function is still incomplete, however. For
example, how is Cu(I) delivery from RAN1 to ETR1 mediated? How does Cu(I)
influence receptor conformation and function? Are there other Cu(I)-binding
components involved? Is the receptor regulated in a tissue-specific way or in response
to the hormonal environment in a tissue? Is this process part of the crosstalk
mechanism with other hormone signaling pathways? It is well established that
ethylene signaling interacts with and is impacted by other hormonal pathways but does
this influence the receptor metalation state and with developmental consequences?

We have previously shown that the loss-of-function *polaris* (*pls*) mutant has
some phenotypic similarities to *ran1* loss-of-function alleles and to *ctr1*, exhibiting a
triple response phenotype (short hypocotyl and root, exaggerated apical hook, radial
expansion) in the dark in the absence of ethylene (Chilley et al., 2006), and a short
root in light-grown seedlings, consistent with its known expression in the root meristem
in light-grown seedlings (Figure 1A). Transgenic complementation of the mutant by the
PLS gene (AT4G39403), which encodes a 36 amino acids peptide, suppresses the
mutant phenotype (Casson et al., 2002). The *pls* mutant phenotype is rescued by the
gain-of-function ethylene resistant mutation *etr1-1* and pharmacological inhibition of
ethylene signalling by silver ions (Chilley et al., 2006). Ethylene gas production in the
pls mutant is at wild type levels, indicating the peptide plays a role in ethylene
signalling rather than biosynthesis (Chilley et al., 2006). *PLS* transgenic
overexpressers (PLSOx seedlings) in contrast exhibit a suppression of the triple
response phenotype when grown in the presence of the ethylene precursor ACC,
similar to the gain of function *etr1-1* mutant, but this suppression is incomplete (PLSOx
seedlings show some response to ACC; Casson et al., 2002; Chilley et al., 2006). PLS
overexpression also partially suppresses the *ctr1* mutant phenotype, indicating that the
PLS peptide acts upstream of CTR1 (Chilley et al., 2006).

In this paper we show that the POLARIS peptide is required for correct
ethylene responses, binds copper via two cysteine residues essential for biological
function, but is unlikely to act as a metallochaperone intermediary; however it co-
localises, and forms a copper-adduct with, ETR1. We suggest this provides cell type
control and a crosstalk regulatory mechanism for ethylene responses through a role in

its physical interaction with the ethylene receptor.

RESULTS

The POLARIS peptide is a negative regulator of ethylene responses

The PLS peptide is translated from a low-abundance transcript that, in light-grown
seedlings, is most strongly expressed in the embryo and silique, seedling root tip and
leaf vascular tissues (Figure S1), a pattern reflected in promoter-reporter expression
patterns (Casson et al., 2002). To further investigate the *pls* molecular phenotype, we
carried out an RNA-seq analysis on loss-of-function *pls* mutant and PLSOx transgenic
overexpressor seedlings to identify differentially regulated genes, with mutant and
overexpressor each compared with wild type seedlings of the same ecotype as
controls (Fig. 1B,C). The *pls* mutant expresses no full length PLS coding sequence
due to a T-DNA insertion in the coding region of *PLS* and disrupting function (Casson
et al., 2002), while the PLSOx seedlings express significantly higher levels of *PLS*
transcript compared to wild type (log$_2$ fold 10.8, *padj* value = 2.02 x 10^{-33}; Tables S1,
S2). 836 genes were significantly up-regulated and 292 down-regulated in the *pls*
mutant compared to wild type control seedlings (padj value <0.05) (Table S1). 1487
genes were significantly up-regulated and 1281 down-regulated in PLSOx seedlings
compared to wild type (padj value <0.05; Table S2). Gene Ontology (GO) analysis of
genes upregulated in *pls* mutant seedlings compared to wild type showed significant
enrichment of genes associated with responses to hormone signalling, biotic and
abiotic defence responses, and cell death (Table S3; Figure S2A).

Out of 307 GO:0009723 (response to ethylene) genes, 25 were significantly
up-regulated and 7 down-regulated in the *pls* mutant compared with the wild type, with
40 up-regulated and 18 down-regulated in the PLSOx seedlings (Figure 1C; Table S3,
S4, S5 and S6; Mudge et al., 2024), indicating a requirement for control over *PLS*
expression levels for correct ethylene responses. While GO:0009723 (response to
ethylene) is significantly enriched in genes up-regulated in the *pls* mutant compared to
wild type (*FDR = 0.00062*; Table S3), a large number of up-regulated genes in *pls* are
significantly associated with immunity, response to pathogens and hypersensitive
response (Table S3 and Figure S2A). Down-regulated genes in *pls* were significantly
enriched in GO categories of hormone biosynthetic processes (GO:0042446, *FDR =*

0.038), hormone metabolic process (GO:0042445, *FDR = 0.0096*) and regulation of
hormone levels (GO:0010817, *FDR = 0.0013*) (Table S4, Figure S2B; Mudge et al.,
2024), consistent with our previous studies that described PLS-dependent crosstalk
between ethylene, auxin and cytokinin signalling (Liu et al., 2010, 2013; Moore et al.,
2015). In PLSOx overexpressors, both up- and down-regulated genes similarly
included those associated with GO terms hormone response, stress response and
immune response, and genes associated with photosynthesis are downregulated in
PLSOx seedlings, consistent with repressed photosynthetic activity in root tips (Table
S5 and S6, Figure S2C, D; Casson et al., 2002). These data are consistent with a role
for PLS in a number of ethylene responses and other related signalling, stress and
developmental processes.

To understand better the relationship between PLS peptide structure and
function between species, we carried out hydroponic feeding experiments using
synthetic versions of PLS peptide from both *Arabidopsis* and its close relative
Camelina sativa. C. sativa contains a gene with partial sequence identity to the
Arabidopsis PLS gene, encoding a predicted peptide sequence that is 22 amino acids
long and identical to the N-terminal 22 amino acids of the *Arabidopsis* PLS except for
a phenylalanine to serine substitution at position nine (Figure 2A). We synthesized full-
length PLS peptide, PLS(FL) and truncated versions from both *Arabidopsis* and *C.*
sativa (Figure 1A) and supplied the peptides to *Arabidopsis pls* mutant seedlings
hydroponically. The full-length peptides from both *Arabidopsis* and *C. sativa* and the
N-terminal 22 amino acids sequence of the *Arabidopsis* peptide (N1) were each able
to rescue the short primary root length of the *Arabidopsis pls* mutant (Figure 2B),
similar to transgenic overexpression and genetic complementation using the wild type
PLS coding sequence (Casson et al., 2002; Chilley et al., 2006). The mean primary
root length of wild type (C24) seedlings did not change across all peptide treatments,
while supply of *pls* seedlings with 50 nM PLS(FL) and N1 peptides showed a rescue of
root growth (Figure 2C), and *t*-tests showed there was no significant difference
between the rescue effects of PLS(FL) and PLS(N1); *P* > 0.1, n = 25). However,
neither a 9 amino acids sequence (N2, Figure 2C) from the N-terminus, nor C-terminal
sequences of 14 (C1) or 24 (C2) amino acids from *Arabidopsis* PLS were able to
rescue the mutant (Figure 2C); each of these shorter peptides lacks one of the two

Cys residues found in the functional longer length peptides PLS(FL) and PLS(N1). A

fluorescent tagged (5-carboxyfluorescein, 5-FAM) version of the *Arabidopsis* N-

terminal 22 amino acids sequence of the peptide (N1) is taken up by the roots, and

also rescues the mutant root phenotype (Figures S3A, B).

PLS localizes to endomembranes including the endoplasmic reticulum

Since genetic studies suggest that PLS acts close to the ethylene receptor (Chilley et

al., 2006), we hypothesized that it should localize to the same subcellular

compartment as ETR1. The ethylene receptor in *Arabidopsis* is localized

predominantly to the ER (Chang, 2003), and a *proPLS::PLS:GFP*-generated fusion

protein (PLS:GFP) was used to investigate sub-cellular localization. Of five

independent transformants with a single-copy insertion of the *proPLS::PLS:GFP* gene

in a *pls* mutant background, four were found to complement fully the *pls* mutant

(Figure S4), similar to the wild type cDNA (Casson et al., 2002), demonstrating

functionality of the gene fusion. The PLS:GFP was found to co-localize in transgenic

plants with the ER marker dye ER-Tracker red^{TM}, which binds sulphonylurea receptors

of ATP-sensitive channels on the ER (Thermofisher; Figures 3A-C). It also co-localizes

with an ER lumen-targeted red fluorescent protein RFP-HDEL (Lee et al., 2013)

(Figures 3G-I). These observations suggest PLS:GFP locates both on the ER

membrane and in the lumen. PLS:GFP also appears to localize to the nucleus and

cytoplasm (Figure 3). As controls, free GFP protein expressed under the control of the

PLS promoter is not co-localized to the ER (Figures 3D-F) and, as expected, the Golgi

marker SH:GFP does not co-localize with ER Tracker (Figures 3M-O). *Trans*-Golgi-

localized SULFOTRANSFERASE1 (ST1) mCherry (Bauer and Papenbrock, 2002)

visualization shows PLS:GFP does not localize to the Golgi (Figures 2J-L). To further

clarify the side of the ER membrane on which PLS localizes, transient expression of

redox-sensitive GFP (roGFP2) fusions of PLS were carried out. The different excitation

properties of roGFP2 in an oxidizing (ER lumen) or reducing environment (cytosol)

allows discrimination of the precise location of PLS fused to RoGFP2. Ratiometric

analysis and comparison with proteins of known localization (i.e. cytosolic RoGFP2,

ER luminal RoGFP2, cytosolic v-SNARE SEC22-GFP, ER luminal RoGFP2-SEC22;

Brach et al., 2009) revealed that PLS as either an N- or C-terminal roGFP2 fusion

resides at the cytosolic side of the ER (as well as to other vesicular compartments;

Figure 3P). However, there may be some translocation of PLS:GFP into the ER lumen

as indicated by co-localization with RFP:HDEL.

PLS interacts with the ethylene receptor protein ETR1

We hypothesized that PLS plays a role in receptor function and investigated whether

this involved direct interaction with the receptor complex. Preliminary experiments

using yeast 2-hybrid analysis suggested that PLS interacts with ETR1 (Figure S5).

Confirmation of the physical interaction between PLS and ETR1 in plants came from

co-immunoprecipitation (Co-IP) analysis. *Agrobacterium* containing a plasmid

encoding PLS linked to a C-terminal GFP and ETR1 with a C-terminal HA tag was

infiltrated into *Nicotiana benthamiana* leaves for transient expression. After 3 d,

interaction was confirmed by western blotting after Co-IP with either anti-GFP beads

(showing PLS pulls down ETR1) or anti-HA beads (showing ETR1 pulls down PLS)

(Figure 4A). GFP-only controls did not show binding with ETR1, demonstrating the

interaction is dependent on the presence of the PLS peptide. The addition of 0.5 µM

copper sulphate to the protein extract used for Co-IP experiments stabilized the PLS-

ETR1 interaction. The presence of copper ions resulted in almost 3-fold more

PLS:GFP detected upon pulldowns with ETR1-HA, or conversely of ETR1-HA pulled

down with PLS-GFP, compared to the same assay in the presence of the metal

chelator 2 mM EDTA (Figures 4A, B).

To investigate the specificity of PLS binding, synthetic full length PLS peptide

PLS(FL) was introduced into infiltrated *N. benthamiana* leaves 30 min before tissue

harvest in the presence of both copper to maximise PLS-ETR1 interaction and the

proteasome inhibitor MG-132 to prevent protein degradation. The addition of 25 nM

synthetic PLS caused a ca. 80% reduction in PLS-GFP binding to ETR1-HA (Figures

4C, D), suggesting that the synthetic PLS peptide competes for ETR1 binding,

showing the specificity of PLS for ETR1. The anti-GFP beads bound two sizes of PLS-

GFP protein (Figure 3E), both of which were larger than a GFP-only control,

suggesting that the PLS peptide undergoes cleavage, a change in conformation, post-

translational modification or incomplete reduction of Cys residues on some PLS. When

using ETR1-HA to pull down PLS-GFP, only the larger peptide was present (Figures

4E, F), suggesting that ETR1 binds a longer version of the PLS peptide.

To pinpoint the interaction site at the receptor in more detail, *in vitro* binding

studies were performed with purified receptor variants and PLS by microscale

thermophoresis (MST; Figure 4G). Binding of PLS was observed only with receptor

variants containing the N-terminal transmembrane domain (TMD). In contrast, no

binding was detected with ETR1 lacking this domain (ETR1[306-738]). The TMD harbors

the ethylene and copper binding region (Schott-Verdugo et al., 2019).

PLS binds Cu(I)

Cysteine residues are common metal-ligand binding residues in low molecular weight

copper-handling peptides, and predictions of PLS structure suggests a single a-helix

plus unstructured region with two cysteines ($CX_{10}C$ arrangement where X is any amino

acid), with some analogy to copper-metallochaperones such as Cox17 or other CX_9C

twin proteins (Robinson and Winge, 2010). In view of both structural considerations

and the copper-dependency of ETR1 we determined whether the two cysteine

residues (C6 and C17) in PLS play a functional role, by analysing both *pls* mutant

complementation and in copper binding by PLS variants. A mutated *Arabidopsis* full-

length peptide in which both cysteines were replaced with serines, PLS(FL C6S,

C17S), was non-functional in hydroponic root feeding assays, failing to rescue the

short primary root phenotype of the *pls* mutant (Figure 5A). Furthermore, as indicated

above, the 9 amino acids sequence N2 and C-terminal sequences C1 and C2 from

Arabidopsis PLS, which each contain only one cysteine residue, were unable to

rescue the mutant (Figure 2C). These results indicate the requirement of the two

cysteine residues for biological activity, measured as primary root growth. Interestingly,

RNA-seq data showed that nineteen out of 287 genes associated with response to

metal ion (GO:0010038) were also significantly down-regulated in the *pls* mutant

(Table S4, Figure S2B, enrichment *FDR = 0.000057*; *20*).

To determine a possible role for PLS and the cysteine residues in binding Cu(I),

synthesised PLS peptide was titrated with Cu(I) ions under strictly anaerobic conditions

and monitored for copper-dependent spectral features (Figures 5B-D). Both UV-vis

(putative metal-to-ligand charge-transfer) and fluorescence (putative tyrosine solvent

access) spectra of PLS changed as a function of Cu(I) ion concentration, consistent with

complex-formation (Figure 5C). We titrated PLS peptide against bicinchoninic acid (BCA; Figure 5B), which is a chromophore that binds Cu(I) (Xiao et al., 2011; Figures 5D and S6). The rationale was to determine whether PLS could compete Cu(I) from BCA, indicative of binding by PLS. PLS peptide was titrated with Cu(I) in the presence of 87 µM BCA. PLS peptide (44.8 µM) increased the concentration of Cu(I) required to saturate BCA increased by ca. 20 µM (Figure 5B), and the initial gradient is shallower than in the control of BCA and Cu(I) but lacking PLS. Thus PLS withholds Cu(I) from BCA, implying an affinity for Cu(I) (K_{Cu} PLS) tighter than K_{Cu} of BCA, and these data also provide support for a 2:1 stoichiometry of binding to the tightest site. Titration experiments also show that although synthetic PLS(FL) withholds Cu(I) from BCA (Figure S7), a mutant PLS (FL C6S, C17S) does not (Fig. S8), confirming a role for those cysteine residues in Cu(I) binding. As an additional and complementary strategy, to eliminate potential solubility problems of PLS in determining accurately the stoichiometry and affinity for Cu(I) (these studies require complete solubility for precise affinity quantification), we also tested a PLS fusion to maltose binding protein (MBP-PLS) which was found to retain solubility when titrated with Cu(I) ions under strictly anaerobic conditions. MBP-PLS (14 µM) was found to withhold ca. 7 µM Cu(I) from BCA, again indicative of a 2:1 PLS:Cu(I) stoichiometry, and tight binding is cysteine-dependent (Figure 5E). A β_2 affinity of 3.79 (±1.5) x10^{19} M^{-2} was determined by competition against an excess of BCA, and the fit significantly departs from simulations 10x tighter or weaker (Figures 5F, S9).

Metal binding in biology is challenging to predict because the formation of metal-protein complexes is a combined function of metal affinity for a given protein and metal availability, which would need to be known for Cu(I) in the *Arabidopsis* cytosol in the case of PLS (assuming the same orientation of binding residues as with the GFP tag; Figure 3P). Cu(I) occupancy of the cytosolic copper chaperone ANTIOXIDANT PROTEIN 1 (ATX1) tracks with fluctuations in available cytosolic Cu(I) such that its affinity approximates to the mid-point of the range of Cu(I) availabilities within this eukaryotic compartment (Yu et al., 2017; Morgan et al., 2019). *Arabidopsis* ATX1 was therefore expressed and purified to determine a 1:1 ATX1:Cu(I) stoichiometry and affinity $K_{D\ Cu(I)}$ of 5.47 (±0.6) x10^{-18} M (Figure 5G inset, Figure S10). Figure 5F reveals that the cytosolic concentration of PLS would need to exceed (improbable) millimolar

concentrations for Cu(I)-dependent homodimers to form at this cytosolic Cu(I)

availability (mathematical models and equations shown in Supplementary text). It is

thus unlikely that the Cu(I):PLS$_2$ complex alone delivers Cu(I) to, or retrieves it from,

the interacting cuproprotein ETR1. Cu(I)-dependent PLS-ETR1 heterodimeric

complexes are the more likely functional species, and we conclude that PLS binding

alters ETR1 conformation or ethylene availability to regulate receptor activity.

DISCUSSION

We present evidence of a role for the PLS peptide as a new Cu(I)-binding peptide that,

based on its physical interactions with the receptor, and its ethylene signalling function

as determined by genetic and physiological experiments, acts as a regulatory

component of the ethylene signalling pathway. Peptides can act as ligands for receptor

kinases to regulate signalling pathways, such as in plant immunity (Campos et al.,

2018), development (Willoughby and Nimchuk, 2021) or in response to abiotic

stresses (Kim et al., 2021). We propose that the PLS peptide is a new component of

the regulatory mechanism for ethylene receptor function linked to its copper-binding

activity (Rodriguez et al., 1999). Loss of copper binding through mutation of the two

cysteines in PLS ablates its biological activity (Figures 2B, C; Figure 5; Figure S8).

The *pls* mutant phenotype has some similarities (enhanced ethylene responses) to

strong *ran1* mutant alleles in which copper delivery to the receptor is compromised

(Binder et al., 2010), and shares (statistically significant) overlap in DEGs with the *ctr1*

mutant, whereby 11.9% of the *ctr1* DEGs are shared with *pls* (each compared to the

respective wild types; Figure S11).

Many plant peptides involved in signalling are processed from longer pre-

proteins (Olsson et al., 2019). In contrast, PLS is translated from an open reading

frame to produce a functional 36 amino acids peptide; the peptide appears to be

cleaved or otherwise modified in the cell (Figure 4E). It is possible that the three

arginines (amino acids 10-12) may represent a cleavage site (Duckert et al., 2004;

Capraro et al., 2015) which would produce the observed shorter GFP fusion protein

(ca. 30 kD) detected in anti-GFP co-IP blots (Fig. 4E). The size of this PLS-GFP

cleavage product suggests that this peptide fragment represents GFP (ca. 27 kDa)

plus a 3 kDa fragment of the C-terminal region of PLS. Such cleavage would likely

inactivate PLS function as we demonstrate that shorter N- and C-terminal fragments (each with only one cysteine residue) are not biologically active (Figure 2C). While PLS:GFP localizes to several subcellular compartments (though not the *trans*-Golgi), some localizes to the ER where the ethylene receptor is found (Figure 3). Furthermore PLS can interact directly with the receptor protein ETR1 (and specifically with the N-terminal copper-binding transmembrane domain of ETR1).

Previously there has been proposed a model for copper delivery into the cell via the COPT family of transporters, which are located at the root plasma membrane (Chen et al., 2022), to RAN1 via ATX1 and/or CCH (Li et al., 2017; Hoppen et al., 2019). Arabidopsis has three known copper chaperones, ATX1, CCH and a copper chaperone for superoxide dismutase CCS (Casareno et al., 1998; Chu et al., 2005; Puig et al., 2007). These are required for copper homeostasis, and both ATX1 and CCH have been shown to interact with RAN1 (Li et al., 2017), providing a link with ethylene signalling (Andrés-Colás et al., 2005; Puig et al., 2007). Chaperones are required for the transport of reactive copper to the correct compartment via these chaperones, avoiding cytotoxicity, following import into the cell; and into the xylem from the cytosol via HMA5, which is structurally similar to RAN1 (Williams and Mills, 2005; Burkhead et al., 2009). Loss of function mutants *cch* and *atx1* have no abnormal phenotype when grown under standard conditions, but the *atx1* and the *atx1 cch* double mutant, though not the *cch* single mutant, are hypersensitive to exogenous copper (Shin et al., 2012). ATX1 and CCH functions have likely diverged, a view supported by their differential transcriptional regulation by copper and different cell type specificities (Shin et al., 2012). The *ran1* mutant is however not copper-hypersensitive, while *hma5* is; and the *pls* mutant is, like *ran1*, not copper hypersensitive (Figure S12), indicative of a role that is distinct from that of ATX1 or CCH. Given its predicted low concentration in the cell and 2:1 stoichiometry with Cu(I) (Figures 5F and G, S9 and 10), we conclude PLS is unlikely to act as a cytosolic copper chaperone.

A lack of copper delivery in the strong *ran1-3* and *ran1-4* null alleles causes constitutive ethylene responses (Woeste and Kieber, 2000), producing a triple response phenotype similar to the *pls* mutant (Casson et al., 2002). These *ran1* mutants fail to bind ethylene due to the lack of copper at the ethylene binding site

(Binder et al., 2010). The decrease in ethylene binding in the *ran1-3* and *ran1-4*
mutants is not due to reduced levels of the ETR1 protein (Binder et al., 2010), and
regulation of the ethylene receptor ETR1 is generally regarded to be independent from
ETR1 transcription or degradation (Hua et al., 1998). Weaker *ran1-1* and *ran1-2*
alleles only show an ethylene response phenotype if they are grown in the presence of
copper chelators (Binder et al., 2010). Therefore, defective RAN1 results in defective
copper delivery to wild type receptors, generating non-functional receptors which
cannot interact with CTR1, resulting in downstream ethylene responses.

Intriguingly, the *ran1-3* and *ran1-4* mutants with constitutive ethylene
responses are distinct from the ethylene insensitive *etr1-1* mutant, which also cannot
bind copper. Therefore, both the *etr1-1* mutant receptor and the strong *ran1* mutant
alleles have reduced ethylene binding capability but the *etr1-1* mutation, which is gain-
of-function, generates seedling phenotypes that are very different from seedlings
carrying the ethylene hypersensitive wild type receptors that simply lack copper due to
defective delivery (in *ran1-3* and *ran1-4).* A receptor carrying the *etr1-1* mutation is
maintained in its 'active' state and constitutively inhibits downstream ethylene
responses, whilst in the *ran1* mutants ETR1 remains in the 'inactive' state, promoting
ethylene responses (Binder et al., 2010). Those authors therefore proposed that
copper has more than one role in the regulation of ETR1 and ethylene signalling: it is
crucial for binding ethylene molecules but it is also needed for the process of signal
transduction by the receptor protein, by an unknown mechanism, to regulate
downstream ethylene responses.

We propose that PLS represents a new component of this pathway, providing
some tissue-specificity to ethylene receptor activity. Given the low probability that PLS
would act as a cytosolic copper chaperone, it more likely regulates ethylene responses
via its demonstrated Cu-dependent ETR1 binding. The RFP-HDEL and Rho-GFP co-
localization experiments indicate the presence of PLS in both the cytosol and ER
lumen (as well as other compartments; Figure 3).

Since the *pls* mutant is rescued by the supply of silver ions (Casson et al.,
2002), and silver is likely delivered to the ethylene receptor via RAN1 (Binder et al.,
2010), this argues against a mechanism in which PLS acts downstream of RAN1 to
metalate ETR1 in the ER lumen. This would also be consistent with the data

suggesting PLS is located predominantly at the cytosolic side of the ER; but it binds to
the transmembrane domain of ETR1. It is also unlikely that PLS removes Cu(I) from
the receptor as this would inactivate the receptor and lead to enhanced ethylene
signalling in the presence of PLS, which is the opposite of what would be expected
from the *pls* mutant phenotype. We therefore favour a model (Fig. 6) in which PLS
binds ETR1 as part of the receptor biogenesis mechanism, with the PLS-ETR1
physical interaction enhanced in the presence of copper (Fig. 4), to regulate receptor
conformation to generate the active form, in which ethylene responses are repressed.
This may be through regulation of Cu(I) homeostasis at the receptor, or via other
structural interactions - for example, PLS could physically block ethylene binding at the
Cu(I) pocket. This question can form the basis of future investigations. If PLS is
required for Cu(I) delivery, it may act redundantly with RAN1, accounting for the
relatively mild ethylene phenotype of the *pls* mutant (compared with e.g. *ctr1*).

Given that *PLS* is expressed in specific tissues, notably root tip and leaf
vascular tissue (Casson et al., 2002), it seems likely that PLS adds a level of
regulation of receptor function in specific tissues, with a focus in this paper on the root
(though developmental defects in the vascular patterning of the leaf are also observed
in the *pls* mutant; Casson et al., 2002). We propose that PLS acts to dampen ethylene
responses at sites of high auxin. *PLS* transcription is strongly up-regulated by auxin,
and its expression profile is similar to that of the auxin reporter DR5 (Sabatini et al.,
1999; Casson et al., 2002). The root tip, where *PLS* is most strongly expressed, is a
site of high auxin concentration, whereby an auxin maximum is formed around the
stem cell niche to regulate both stem cell identity and function (Sabatini et al., 1999).
The vascular tissues are also a site of high auxin (Bishopp et al., 2011). Auxin can
induce ethylene biosynthesis by upregulation of the ethylene biosynthetic enzyme
ACC synthase (Abel et al., 1995), and high ethylene concentrations can lead to both
reduced meristem function and aberrant division of the quiescent centre and cells
within the stem cell niche and columella (Ortega-Martinez et al., 2007). Therefore
ethylene biosynthesis and signalling in the root tip must be tightly regulated to allow
growth control. *PLS* expression is transcriptionally activated by auxin and suppressed
by ethylene. We have shown in previous work that there is crosstalk between PLS,
ethylene and other signalling pathways, notably auxin, cytokinin and ABA in the root

(e.g. Liu et al., 2010, 2013; Moore et al., 2015), and the distinctiveness of *pls*
compared to other ethylene signaling mutants may be linked to its restricted
expression pattern. This network provides a mechanism to suppress growth-inhibitory
ethylene responses in the high auxin environment of the root tip, which is dependent
on the tissue-specific expression of *PLS* (Liu et al., 2010) (Figure S13) and is
mediated through its interaction with the ethylene receptor.

Future structural studies should reveal more about the PLS-ETR1-RAN1-
chaperone interaction and diverse roles of copper ions in ethylene signal transduction,
which represents a new paradigm for the regulation of signaling protein function by
metals.

METHODS

Plant Material

Seedlings of *Arabidopsis thaliana* ecotype either C24 or Col-0, or *pls* mutants or
transgenic *PLS* overexpressers, all from lab stocks, were grown on solid sterile half-
strength MS10 medium on 2.5% Phytagel (Sigma-Aldrich) in 90 mm Petri dishes
(Sarstedt, Leicester, UK) at 21°C, under a 16 hour photoperiod as described (Casson
et al., 2002; Chilley et al., 2006). *Nicotiana benthamiana* plants (lab stocks) were
grown in a controlled environment (21°C, 16 h photoperiod) for transient expression
studies in leaf tissue. For hydroponic feeding studies, *Arabidopsis* seedlings were
cultured in liquid medium (1 ml/well) in sterile 24-well plates (Sarstedt), essentially as
described (Matsuzaki et al., 2010). For hormone and peptide assays, one seedling
was grown in each well, at 21°C, under a 16 h photoperiod. For peptide feeding
experiments, purified freeze-dried peptide was dissolved in DMSO to create a 500 µM
stock solution. Peptide stock solution was added to liquid ½ MS10 plant media
containing 0.1% DMSO to make a final peptide concentration of 50 or 100 nM (or 10,
25, 50 and 100 nM for dose-dependent assays). For copper treatments, 1 mM $CuSO_4$
solution was filter-sterilized and added to autoclaved liquid ½ MS10 plant media to
create final $CuSO_4$ concentrations of 0, 5, 10, 15, 20, 25, 30, 35, 40, 45 and 50 µM.
The copper chelator bathocuproine disulphonic acid (BCS) was added to liquid

medium to produce final concentrations of 0, 10, 50, 100, 250 and 500 µM. Seedlings

were scanned to create a digital image and root lengths of seedlings were measured

using ImageJ. Statistical analysis was performed using the Real Statistics Resource

Pack software (Release 3.8, www.real-statistics.com) in Excel (Microsoft).

Peptide Synthesis

Peptides were either obtained from Cambridge Research Biochemicals (Billingham,

UK) or synthesized in the laboratory by Fmoc solid phase peptide synthesis (SPPS) on

a CEM Liberty1 single-channel peptide synthesiser equipped with a Discover

microwave unit. Reactions were performed in a 30 ml PTFE reaction vessel with

microwave heating FW: EpiLife Media and agitation by bubbling nitrogen. Peptide

synthesis was carried out using 2-chlorotrityl chloride resin (0.122 mmol g^{-1},

Novabiochem) using Fmoc-protected amino acids. The first amino acid residue at the

C-terminus (histidine) was coupled manually by mixing 76 mg (1 eq.) Fmoc-His(Trt)-

OH, 0.09 ml (4 eq.) *N,N*-diisopropylethylamine (DIPEA), 1 ml dichloromethane (DCM)

and 1 ml dimethylformamide (DMF) until the amino acid powder had dissolved. The

mixture was added to 0.1 mmol resin and stirred gently for 120 min at room

temperature. Resin was washed with 3x DCM/MeOH/DIPEA (17:2:1), 3x DCM, 2x

DMF and 2x DCM. Amino acid coupling reactions were performed using Fmoc-

protected amino acids present in a 5-fold excess (2 M concentration), HOBt (0.5 M

HOBt in DMF, used at the activator position) and DIC (0.8 M in DMSO, used at the

activator base position). For double and triple couplings the reaction vessel was

drained after each coupling cycle and fresh reagents were added. Before each

coupling, a room temperature preactivation period of 1 to 2 h was used. Microwave-

assisted couplings were performed for 10 min at 75°C at 25W power. Cys and His

residues were coupled at low temperature (10 min at room temperature followed by 10

min at 50°C, 25W). Arg residues were double coupled, firstly by 45 minutes at room

temperature plus 5 minutes at 75°C (25W), and second by the standard microwave

conditions above. Fmoc group removal was carried out by two piperidine solution

treatments (20% piperidine in DMF) in succession: 5 minutes then 10 minutes. Peptide

cleavage from resin was carried out in 3 ml 95% TFA in dH$_2$O/TIPS (2.85 ml TFA, 0.15

ml dH$_2$O, 0.15 ml triisopropylsilane). Peptide was dissolved in water with a small

volume of MeCN and lyophilized to produce a powder using a Christ ALPHA 1-2 LD$_{plus}$

freeze dryer.

Preparative High-Performance Liquid Chromatography (HPLC)

Peptide products were analysed and purified by HPLC at 280 nm. 25-50 mg of freeze-

dried peptide sample was dissolved in 1 ml 1:1 H$_2$O:MeCN and injected onto a Speck

and Burke Analytical C18 Column (5.0 µm, 10.0 x 250 mm) attached to a PerkinElmer

(Massachusetts, USA) Series 200 LC Pump and 785A UV/Vis Detector. Separation

was achieved by gradient elution of 10-80% solvent B (solvent A = 0.08% TFA in

water; solvent B = 0.08% TFA in ACN) over 60 min, followed by 80-100% B over 10

min, with a flow rate of 2 ml/min. Selected peptide fractions were lyophilized and a

mass assigned using MALDI-TOF MS. Peptide sequences were identified using

MALDI-TOF MS, using an Autoflex II ToF/ToF mass spectrometer (Bruker Daltonik

GmBH, Germany) equipped with a 337 nm nitrogen laser. MS data was processed

using FlexAnalysis 2.0 (Bruker Daltonik GmBH).

Imaging

CSLM images for propidium iodide and GFP were obtained by a Leica SP5 TCS

confocal microscope as previously described (Rowe et al., 2016). *pPLS::PLS:GFP,*

pPLS::GFP and *p35S::GFP* seedlings were grown for 7 d on Phytagel ½ MS10

medium before ca. 25 mm of the root tip was removed and mounted in dH$_2$O prior to

imaging. The ER marker *p35S::RFP:HDEL* (*24*) (provided by Dr. Pengwei Wang,

Durham University), and the trans-Golgi apparatus marker *pFGC-ST:mCherry* (from

Nottingham Arabidopsis Stock Centre, www.arabidopsis.info) were introduced into

pPLS::PLS:GFP plants by the floral dip method of transformation (Clough and Bent,

1998) using *A. tumefaciens* GV3 101. ER was also localized using ER Tracker™ Red

(Thermofisher). Seven day-old seedlings were stained for 30 min in the dark in liquid

½ MS10 media containing 1 µM ER Tracker™ Red.

RNA Isolation, RNA Sequencing, RT-qPCR

RNA was extracted from 7 day-old seedlings grown on half strength MS10 medium

essentially as described (Thompson et al., 2023). The Illumina HiSeq 2500 System

was used for RNA sequencing of three biological replicate samples, with libraries
prepared using the Illumina TruSeq Stranded Total RNA with Ribo-Zero Plant Sample
Preparation kit (RS-122-2401), essentially as described (Thompson et al., 2023).
Library quality control was carried out again using a TapeStation with D1000
ScreenTape (catalog number 5067-5582). RNA-seq data were aligned to the TAIR10
(EnsemblePlants, release 58) genome sequence with corresponding gtf file using
STAR (v 2.7.11a; Dobin et al., 2013) to obtain read count per gene. Read count per
gene was analysed using DESeq2 v 1.40.2 (Love et al., 2014) to get p values,
adjusted p values, and log2 fold changes. Differentially expressed genes (DEGs) were
identified with the criteria of adjusted p value < 0.05. Gene ontology analysis was
performed using AgriGO (Tian et al., 2017) singular enrichment analysis. Gene
expression heatmaps were generated from log2 normalized counts using pheatmap
(v1.0.12, https://cran.r-project.org/web/packages/pheatmap/index.html) with row
scaling. RNA-seq data were deposited in GEO (https://www.ncbi.nlm.nih.gov/geo/)
with accession number GSE256166 (Mudge et al., 2024).

For RT-qPCR, RNA was extracted from 7 day-old seedlings (3 biological
replicates, 20 mg of tissue) as described (Thompson et al., 2023). Samples were
checked for the presence of genomic DNA by PCR with *ACTIN2* primers ACT2
forward and reverse. Primer sequences were determined using Primer-BLAST
(https://www.ncbi.nlm.nih.gov/tools/primer-blast/). Primers are listed in Table S7.

Protein-protein interaction studies
Yeast 2-hybrid
The GAL4 two-hybrid phagemid vector system was used to detect protein-protein
interactions *in vivo* in yeast, using the reporter genes β-galactosidase *(lacZ)* and
histidine (*HIS3*) in the YRG-2 yeast strain, essentially as described (Zhong et al.,
2008**)**. DNA that encodes the target (ETR1) and bait (PLS) was inserted into the pAD-
GAL4-2.1 A and pBD-GAL4 Cam phagemid vectors respectively and expressed as
hybrid protein. The hybrid proteins were then assayed for protein-protein interaction.

DNA encoding the target and bait proteins were prepared by PCR
amplification using primer designed the specifically for the DNA encoding the target
(ETR1) and bait (PLS). Each set of primer contained specific endonucleases on the

ends of primer corresponding to the endonucleases in the MCS of pAD-GAL4-2.1 A

and pBD-GAL4 Cam phagemid vectors. The DNA construct of the target (ETR1) and

bait (PLS) with specific restriction sites on the ends was then transformed into the

TOPO 2.1 vector to check the sequence of the amplified DNA by sequencing with M13

forward (CTG GCC GTC GTT TTA C) and M13 reverse (CAG GAA ACA GCT ATG

AC). The two vectors, pAD-GAL4-2.1 and pBD-GAL4 Cam were digested using

specific restriction endonucleases and dephosphorylated prior to ligating the insert

DNA. The DNA encoding the target (ETR1) and bait (PLS) was then ligated into the

same reading frame as the GAL4 AD of the pAD-GAL4-2.1 phagemid vector and the

GAL4 BD of the pBD-GAL4 Cam phagemid vector.

The following primers were used for PCR amplification:

ETR1:

Forward primer GAA TCC ATG GAA GTC TGC AAT GTT A (Eco RI on 5' end)

Reverse primer GTC GAC TTA CAT GCC CTC GTA CA (Sal I on 5'end)

PLS:

Forward primer CTG GAG ATG AAA CCC AGA CTT TGT (Xho I on 5' end)

Reverse primer GTC GAC ATG GAT TTT AAA AAG TTT (Sal I on 5' end)

The pGAL4 control plasmid was used alone to verify that induction of the *lacZ*

and *HIS3* genes has occurred and that the gene products are detectable in the assay

used. The pLamin C control plasmid was used in pairwise combination with the pAD-

WT control plasmid or with the pAD-MUT control plasmid to verify that the *lacZ* and

HIS3 genes are not induced as the protein expressed by each of these pairs do not

interact *in vivo*.

Control plasmids were transformed into the YRG-2 strain prior to the initial

transformation of the bait and the target plasmids and used separately or in pairwise

combination in the transformation of the YRG-2 yeast strain. Yeast competent cells

were cotransformed with the bait and target plasmids by sequential transformation.

Gene cloning for co-immunoprecipitation

To investigate the interaction between the PLS peptide and the ethylene receptor

ETR1, two DNA constructs were created by Gateway cloning. The 105-nucleotide *PLS*

gene (without the stop codon) was inserted into the pEarlyGate103 (pEG103)

destination vector, containing the *p35S* promoter and a C-terminal GFP tag, producing
a vector containing the *p35S::PLS:GFP* DNA. The ETR1 cDNA was inserted into the
pEarlyGate301 (pEG301) vector to create a *p35S::ETR1:HA* construct, producing an
ETR1 protein with a C-terminal HA tag.

Infiltration into *Nicotiana benthamiana*

The transient expression of constructs in *Nicotiana benthamiana* (tobacco) leaves was
as previously described (Voinnet et al., 2007). Experiments were replicated up to 5
times. Competent *Agrobacterium tumefaciens* GV3101 cells were transformed with the
desired plasmid containing the gene of interest and each injected with a syringe. The
plants were approximately 7 to 10 weeks old; the chosen leaves were healthy and of
length 3-6 cm, and 3 to 4 leaves were infiltrated with each construct.

Protein extraction and PLS/ETR1 co-immunoprecipitation

Total protein was extracted from the infiltrated leaves of *N. benthamiana* plants 3 d
after infiltration for co-immunoprecipitation experiments to investigate the interaction
between PLS and ETR1 proteins, essentially as described previously (Srivastava et
al., 2020). For competition assays, 5 nM or 25 nM full-length PLS peptide was also
infiltrated in the presence of 50 µM MG-132 (a proteasome inhibitor) 30 min prior to
tissue freezing. ChromoTek (Planegg, Germany) anti-GFP beads were used to
immunoprecipitate the PLS-GFP protein, and Sigma-Aldrich (St. Louis, USA) anti-HA beads
for the HA-tagged ETR1.

SDS-PAGE was used to separate protein fragments. The complexed proteins
from the pull-down assay were analysed on 10-12% acrylamide gels. Membranes
were incubated with primary antibody for 2.5 h (GFP, Abcam, Cambridge, UK: rabbit,
1:10000; HA (Roche), rat, 1:3000; Rubisco large subunit (Agrisera), rabbit, 1:10000).
Excess primary antibody was then removed by washing three times in 2x TBST (150
mM NaCl, 10 mM Tris, 0.1% (v/v) Tween 20, pH 7.4) for 2 m, 5 min and 10 min, and
subsequently incubated for 1 h with the ECL peroxidase-labelled anti-rabbit or anti-rat
IgG secondary antibody, diluted 1:20000 in TBST. Excess secondary antibody was
removed again by washing three times in 1x TBST, as with the primary antibody. In
order to visualize the probed blot, the membrane was incubated with ECL Western

Blotting Detection Reagent immediately prior to imaging. The horseradish peroxidase

conjugated to the secondary antibody was detected by using X-ray film.

Estimation of synthetic PLS concentration

Freeze-dried synthetic PLS peptide (Cambridge Research Biochemicals, Billingham)

was dissolved in DMSO. An aliquot was added to aqueous buffer (10 mM HEPES

pH7, 20 mM NaCl, 80 mM KCl) and absorbance at 280 nm was recorded.

Concentration was estimated from the absorbance and the ProtParam estimated

extinction coefficient of 2,980 $M^{-1} cm^{-1}$. Concurrent with this a sample was submitted

for quantitative amino acid analysis (Abingdon Health Laboratory Services). From this

analysis a conversion factor of 2.27 was generated, which was applied to

concentrations determined by A280 nm.

ATX1 purification

E. coli BL21(DE3) containing pETatx1 was used to overexpress the wild type *ATX1*

gene from *Arabidopsis thaliana* (optimised for expression in *E. coli*, Novoprolabs).

Harvested cells were collected and frozen at -20 ^{o}C overnight then defrosted,

resuspended in 20 mM HEPES (pH 7.0), 10 mM EDTA, 100 mM NaCl, 10 mM DTT.

Cells were sonicated (Bandelin Sonoplus), supernatant separated by size exclusion

chromatography (GE Healthcare, HiLoad 26.600 Superdex 75 pg) metal-free buffer

lacking EDTA. Fractions containing ATX1 were incubated overnight and pooled before

transfer into an anaerobic chamber (Belle Technology) via desalting column where

reductant was removed. ATX1 was quantified by combination of Bradford assay and

Ellman's reagent to ensure the fully reduced state of the protein. Samples were also

analysed for metal content by ICP-MS to ensure > 95% apo-ATX1.

MBP-PLS/mutant purification

A fusion of PLS to MBP was created using the NEBExpress MBP Fusion and

Purification System. Two complementary oligonucleotide primers encoding PLS

(optimised for expression in *E. coli*) were annealed and inserted into the pMal-c5x

plasmid at XmnI and SalI insertion sites.

The three mutants MBP-PLS(C6S), MBP-PLS(C17S) and MBP-
PLS(C6S/C17S) were created by site-directed mutagenesis (QuikChange II, Agilent).
E. coli NEB Express containing the pMal plasmid with the correct MBP-PLS mutant
was used to overexpress each protein. Harvested cells were resuspended in 20 mM
Tris-HCl (pH 7.4), 200 mM NaCl, 1 mM EDTA and frozen at -20 ^{o}C overnight. Cells
were defrosted in cold H_2O, sonicated, purified by ammonium sulphate precipitation
(where MBP-PLS precipitates >60% saturation), separated on MBP-trap (GE
Healthcare) and eluted using buffer containing 10 mM maltose. MBP-PLS-containing
fractions were pooled and concentrated using centrifugal concentrator (Corning, Spin-
X UF 30 KDa) and buffer exchanged by desalting column into metal-free 20 mM
HEPES (pH 7.0), 50 mM NaCl buffer in an anaerobic chamber (Belle Technology).
Mutants containing thiols were quantified Ellman's assay and MBP-PLS(C6S/C17S),
which lacks all thiols, was quantified by Bradford assay alone. Samples were also
analysed for metal content by ICP-MS to ensure >95% apo-protein.
676

Anaerobic spectroscopic analysis of Cu(I) complexes
All Cu(I) titration experiments were carried out in an anaerobic chamber (Belle
Technology) using metal-free CHELEX-treated, degassed buffers. For experiments
titrating Cu(I), aqueous $CuSO_4$ stock was quantified in advance by ICP-MS and diluted
to working concentrations. The reductant NH_2OH was included at final concentration of
1 mM to retain Cu(I) in its reduced state. Proteins were diluted in buffer to the final
concentration specified in each titration in air-tight quartz cuvettes (Helma), and after
addition of probe to the concentration specified, titrated with $CuSO_4$. After each
addition, solutions were thoroughly mixed and absorbance spectra recorded using a
Lambda 35 UV/Vis spectrophotometer (Perkin Elmer). Titration isotherm data was
fitted using simulated affinity curves using DynaFit (Kuzmič, 2009).
688

Interaction studies of PLS with copper transporter ETR1 by Microscale
Thermophoresis
Fluorescently labelled ETR1 truncation mutants were added to a dilution series of PLS
in 50 mM HEPES, 150 mM NaCl, 0.015 % (wv) FosCholine 16 (pH 7.6) or 50 mM Tris,

300 mM NaCl, 0.015 % (w/v) FosCholine 16 (pH 7.6). Dissociation constants were

calculated using GraphPad Prism 5.

Determination of dissociation constants for the PLS-ETR1 interaction

Full-length ETR1 and truncation mutants were purified and labelled as described in

(Milić et al., 2018). 94 µM PLS were diluted serially in 50 mM Tris and 300 mM NaCl

(pH 7.6). Fluorescently labelled receptor was added at a final concentration of 50 nM.

Thermophoretic behaviour was measured in premium capillaries at 50 % LED and 50

% MST power. In case of a binding event, data were fitted using GraphPad Prism 5.

Ratiometric analysis of roGFP2 fusion proteins

N- and C-terminal roGFP2 fusion proteins of PLS were generated by Gateway cloning.

Infiltration and transient expression of roGFP2 fusions and control proteins were

carried out as described (Brach et al., 2009). Image acquisition and data analysis were

carried out as described by Hoppen et al. (2019). A minimum of 10 leaf optical

sections were imaged and used for ratiometric analysis of the redox sensitive

excitation properties of roGFP2.

SUPPLEMENTAL INFORMATION

All study data are included in the article and/or Supplemental Information are available

at *Molecular Plant Online*. RNA-seq and root assay data are available at GEO

accession GSE256166:

FUNDING

The authors acknowledge financial support from: UK Biotechnology and Biological
Sciences Research Council BB/E006531/1, BBS/B/0773X, BB/J014516/1 (KL),
BB/V006002/1, BB/M011186/1 (NJR); Deutsche Forschungsgemeinschaft (German
Research Foundation) 267205415 – SFB 1208 project B06 (GG). We thank Prof.
Steven Cobb (Durham University Department of Chemistry) for advice on peptide
synthesis, and Dr. Andrew Foster (Durham University Department of Chemistry) for
preliminary peptide-Cu interaction analysis.

AUTHOR CONTRIBUTIONS

734 KL and NJR initiated the project. KL, JFT, NJR, AS and GG designed and supervised
735 aspects of the project. AJM, SM, WM, BO-P, WS, WW, CT, DR, FMH, CH and BU
736 carried out the experimental work and prepared the Figures. KL, NJR and GG drafted
737 the early version of the manuscript, and all authors reviewed and edited the
738 manuscript.

ACKNOWLEDGEMENTS

741 The authors declare no competing interests.

REFERENCES

Abel, S., Nguyen, M., Chow, W., Theologis, A. (1995). *ASC4*, a primary indoleacetic
acid-responsive gene encoding 1-amino-cyclopropane-1-carboxylate synthase in
Arabidopsis thaliana. J. Biol. Chem. **270**: 19093–19099.

Andrés-Colá,s N., Sancenón, V., Rodríguez-Navarro, S., Mayo, S., Thiele, D.J.,
Ecker, J.R., Puig, S., Peñarrubia, L. (2006). The Arabidopsis heavy metal P-type
ATPase HMA5 interacts with metallochaperones and functions in copper
detoxification of roots. Plant J. **45**: 225–236.

Bauer, M., Papenbrock, J. (2002). Identification and characterization of single-domain
thiosulfate sulfurtransferases from *Arabidopsis thaliana*. FEBS Lett. **532**: 427-
431.

Binder, B.M., Rodriguez, F.I., Bleecker, A.B. (2010). The copper transporter RAN1 is
essential for biogenesis of ethylene receptors in Arabidopsis. J. Biol. Chem. **285**:
37263-37270.

Bishopp, A., Help, H., El-Showk, S., Weijers, D., Scheres, B., Friml, J., Benkova, E.,
Mahonen, A.P., Helariutta, Y. (2011). A mutually inhibitory interaction between
auxin and cytokinin specifies vascular pattern in roots. Curr. Biol. **21**: 917-926.

Brach, T., Soyk, S., Müller, C., Hinz, G., Hell, R., Brandizzi, F., Meyer, A.J . (2009).
Non-invasive topology of membrane proteins in the secretory pathway. Plant J.
57: 534-541.

Burkhead, J.L., Reynolds, K.A., Abdel-Ghany, S.E., Cohu, C.M., Pilon, M. (2009).
Copper homeostasis. New Phytol. **182**: 799–816.

Campos, M.L., de Souza, C.M., de Oliviera, K.B.S., Dias, S.C., Franco, O.L. (2018).
The role of antimicrobial peptides in plant immunity. J. Exp. Bot. **69**: 4997-5011.

Capraro, J., Sessa, F., Magni, C., Scarafoni, A., Maffioloi, E., Tedeschi, G., Croy,
R.D.D., Duranti, M. (2015). Proteolytic cleavage at twin arginine residues affects
structural and functional transitions of Lupin seed 11S storage globulin. PLOS
ONE **10**: e0117406.

Casareno, R.L., Waggoner, D., Gitlin, J.D. (1998). The copper chaperone CCS directly
interacts with copper/zinc superoxide dismutase. J. Biol. Chem. **273**: 23625–
23628.

Casson, S.A., Chilley, .PM., Topping, J.F., Evans, I.M., Souter, M.A., Lindsey, K.
(2002). The *POLARIS* gene of Arabidopsis encodes a predicted peptide required
for correct root growth and leaf vascular patterning. Plant Cell **14**: 1705-1721.

Chang, C.R. (2003). Ethylene signaling: the MAPK module has finally landed. Trends
Plant Sci. **8**: 365-368.

Chang, C., Kwok, S.F., Bleecker, A.B., Meyerowitz, E.M. (1993). Arabidopsis ethylene-
response gene *ETR1* - similarity of product to 2-component regulators. Science
262: 539-544.

Chen, G., Li, J., Han, H., Du, R., Wang, X. (2022). Physiological and molecular
mechanisms of plant responses to copper stress. Int. J. Mol. Sci. **23**: 12950.

Chen, Y.F., Randlett, M.D., Findell, J.L., Schaller, G.E. (2002). Localization of the
ethylene receptor ETR1 to the endoplasmic reticulum of Arabidopsis. J. Biol.
Chem. **277**: 19861-19866.

Chilley, P.M., Casson, S.A., Tarkowski, P., Hawkins, N., Wang, K.L., Hussey, P.J.,
Beale, M., Ecker, J.R., Sandberg, G.K., Lindsey, K. (2006). The POLARIS

789   peptide of Arabidopsis regulates auxin transport and root growth via effects on
790   ethylene signaling. Plant Cell **18**: 3058-3072.

791 Chu, C.C., Lee, W.C., Guo, W.Y., Pan, S.M., Li, L.J., Chen, H.M., Jinn, T.L. (2005). A
792   copper chaperone for superoxide dismutase that confers three types of
793   copper/zinc superoxide dismutase activity in Arabidopsis. Plant Physiol. **139**:
794   425–436.

795 Clough, S.J., Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-
796   mediated transformation of *Arabidopsis thaliana*. Plant J. **6**: 735-743.

797 Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
798   Chaisson, M., Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
799   Bioinformatics **29**: 15-21.

800 Duckert, P., Brunak, S., Blom, N. (2004). Prediction of preprotein convertase cleavage
801   sites. Prot. Engin. Design Select. **17**: 107-112.

802 Gao, Z.Y., Chen, Y.-F., Randlett, M.D., Zhao, X.-C., Findell, J.L., Kieber, J.J., Schaller,
803   G.E. (2003). Localization of the Raf-like kinase CTR1 to the endoplasmic
804   reticulum of Arabidopsis through participation in ethylene receptor signaling
805   complexes. J. Biol. Chem. **278**: 34725-34732.

806 Grefen, C., Staedele, K., Ruzicka, K., Obrdlik, P., Harter, K., Horak, J. (2008).
807   Subcellular localization and *in vivo* interactions of the *Arabidopsis thaliana*
808   ethylene receptor family members. Molec. Plant. **1**: 308-320.

809 Hall, A.E., Findell, J.L., Schaller, G.E., Sisler, E.C., Bleecker, A.B. (2000). Ethylene
810   perception by the ERS1 protein in Arabidopsis. Plant Physiol. **123**: 1449-1457.

811 Hirayama, T., Kieber, J.J., Hirayama, N., Kogan, M., Guzman, P., Nourizadek, S.,
812   Alonso, J.M., Dailey, W.P., Dancis, A., Ecker, J.R. (1999). RESPONSIVE-TO-
813   ANTAGONIST1, a Menkes/Wilson disease-related copper transporter, is required
814   for ethylene signaling in Arabidopsis. Cell **97**: 383-393.

815 Hoppen, C., Müller, L., Hänsch, S., Uzun, B., Milic, D., Meyer, A.J., Weidtkamp-Peters,
816   S., Groth, G. (2019). Soluble and membrane-bound protein carrier mediate
817   direct copper transport to the ethylene receptor family. Sci. Rep. **9**: 10715.

818 Hua, J., Chang, C., Su,n Q., Meyerowitz, E.M. (1995). Ethylene insensitivity conferred
819   by Arabidopsis *ERS* gene. Science **269**: 1712-1714.

Hua, J., Sakai, H., Nourizadeh, S., Chen, Q.G., Bleecker, A.B., Ecker, J.R.,
Meyerowitz, E.M. (1998). EIN4 and ERS2 are members of the putative ethylene
receptor gene family in Arabidopsis. Plant Cell **10**: 1321-1332.

Johnson, P.R., Ecker, J.R. (1998). The ethylene gas signal transduction pathway: A
molecular perspective. Ann. Rev. Genet. **32**: 227-254.

Kim, J.S., Jeon, B.W., Kim, J. (2021). Signaling peptides regulating abiotic stresses in
plants. Front. Plant Sci. **12**: 704490.

Kuzmič, P. (2009). DynaFit—a software package for enzymology. Meth. Enzymol. **467**:
247–280.

Lee, H., Sparkes, I., Gattolin, S., Dzimitrowicz, N., Roberts, L.M., Hawes, C., Frigerio,
L. (2013). An Arabidopsis reticulon and the atlastin homologue *RHD3-like2* act
together in shaping the tubular endoplasmic reticulum. New Phytol. **197**: 481-
489.

Li, W., Lacey, R.F., Ye, Y., Lu, J., Yeh, K.-C., Xiao, Y., Li, L., Wen, C.-K., Binder, B.M.,
Zhao, Y. (2017). Triplin, a small molecule, reveals copper ion transport in
ethylene signaling from ATX1 to RAN1. PLOS Genet. **13**: e1006703.

Liu, J., Mehdi, S., Topping, J,. Tarkowski, P., Lindsey, K. (2010). Modelling and
experimental analysis of hormonal crosstalk in Arabidopsis. Molec. Systems Biol.
6: 373.

Liu, J., Mehdi, S., Topping, J., Friml, J., Lindsey, K. (2013). Interaction of PLS and PIN
and hormonal crosstalk in Arabidopsis root development. Front. Plant Sci. **4**: 75.

Love, M.I., Huber, W., Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. **15**: 550.

Matsuzaki, Y., Ogawa-Ohnishi, M., Mori, A., Matsubayashi, Y. (2010). Secreted
peptide signals required for maintenance of root stem cell niche in Arabidopsis.
Science **329**: 1065-1067.

McDaniel, B.K., Binder, B.M. (2012). ETHYLENE RECEPTOR 1 (ETR1) is sufficient
and has the predominant role in mediating inhibition of ethylene responses by
silver in *Arabidopsis thaliana*. J. Biol. Chem. **287**: 26094-26103.

Milić, D., Dick, M., Mulnaes, D., Pfleger, C., Kinnen, A., Gohlke, H., Groth, G. (2018).
Recognition motif and mechanism of ripening inhibitory peptides in plant
hormone receptor ETR1. Sci. Rep. **8**: 1–12.

Moore, S., Zhang, X., Mudge, A., Rowe, J.H., Topping, J.F., Liu, J., Lindsey, K. (2015).
Spatiotemporal modelling of hormonal crosstalk explains the level and patterning
of hormones and gene expression in *Arabidopsis thaliana* wildtype and mutant
roots. New Phytol. **207**: 1110-1122.

Morgan, M.T., Bourassa, D., Harankhedkar, S., McCallum, A.M., Zlatic, S.A., Calvo,
J.S,. Meloni, G., Faundez, V., Fahrni, C.J. (2019). Ratiometric two-photon
microscopy reveals attomolar copper buffering in normal and Menkes mutant
cells. Proc. Natl. Acad. Sci. USA **116**: 12167–12172.

Mudge, A.J., Mehdi, S., Michaels, W., Orosa-Puente, B., Shen, W., Tomlinson, C.,
Wei, W., Hoppen, C., Uzun, B., Roy, D., Hetherington, F.M., Topping, J.F.,
Sadanandom, A., Groth, G., Robinson, N.J., Lindsey, K. (2024). Data from:
POLARIS is a copper-binding peptide required for ethylene signalling control in
Arabidopsis. GEO accession
GSE256166: https://eur01.safelinks.protection.outlook.com/?url=https%3A%2F
%2Fwww.ncbi.nlm.nih.gov%2Fgeo%2Fquery%2Facc.cgi%3Facc%3DGSE25616
6&data=05%7C02%7Ckeith.lindsey%40durham.ac.uk%7C9637f7022a2240cfb10
608dc324244cd%7C7250d88b4b684529be44d59a2d8a6f94%7C0%7C0%7C63
8440507397042151%7CUnknown%7CTWFpbGZsb3d8eyJWIjoiMC4wLjAwMDAi
LCJQIjoiV2luMzIiLCJBTiI6Ik1haWwiLCJXVCI6Mn0%3D%7C0%7C%7C%7C&sd
ata=U4JHCoviYbdEbF76Tws8Ridq7W%2B5Hq2qBVN%2BFjIysgw%3D&reserve
d=0

Olsson, V., Joos, L., Zhu, S., Gevaert, K., Butenko, M., De Smet, I. (2019). Look
closely, the beautiful may be small: Precursor-derived peptides in plants. Ann.
Rev. Plant Biol. **70**: 153-186.

Ortega-Martinez, O., Pernas, M., Carol, R.J., Dolan, L. (2007). Ethylene modulates
stem cell division in the *Arabidopsis thaliana* root. Science **317**: 507- 510.

Puig, S., Mira, H., Dorcey, E., Sancenón, V., Andrés-Colás, N., Garcia-Molina, A.,
Burkhead, J.L., Gogolin, K.A., Abdel-Ghany, S.E., Thiele, D.J., Ecker, J.R., Pilon,
M., Penarrubia, L. (2007). Higher plants possess two different types of ATX1-like
copper chaperones. Biochem. Biophys. Res. Commun. **354**: 385–390.

Qu, X., Hall, B.P., Gao, Z.Y., Schaller, G.E. (2007). A strong constitutive ethylene-
response phenotype conferred on Arabidopsis plants containing null mutations in
the ethylene receptors *ETR1* and *ERS1*. BMC Plant Biol. **7**: 3.

Robinson, N.J., Winge, D.R. (2010). Copper metallochaperones. Ann. Rev. Biochem.
79: 537-562.

Rodriguez, F.I., Esch, J.J., Hall, A.E., Binder, B.M., Schaller, G.E., Bleecker, A.B.
(1999). A copper cofactor for the ethylene receptor ETR1 from Arabidopsis.
Science **283**: 996-998.

Rowe, J.H., Topping, J.F., Liu, J., Lindsey, K. (2016). Abscisic acid regulates root
growth under osmotic stress conditions via an interacting hormonal network with
cytokinin, ethylene and auxin. New Phytol. **211**: 225-239.

Sabatini, S., Beis. D., Wolkenfelt, H., Murfett, J., Guilfoyle, T., Malamy, J., Benfey, P.,
Leyser, O., Bechtold, N., Weisbeek, P., Scheres, B. (1999). An auxin-dependent
distal organizer of pattern and polarity in the Arabidopsis root. Cell **99**: 463-472.

Sakai, H., Hua, J., Chen, Q.G., Chang, C., Medrano, L.J., Bleecker, A.B., Meyerowitz,
E.M. (1998). *ETR2* is an *ETR1*-like gene involved in ethylene signaling in
Arabidopsis. Proc. Natl. Acad. Sci. **USA** 95: 5812-5817.

Schaller, G.E., Ladd, A.N., Lanahan, M.B., Spanbauer, J.M., Bleecker, A.B. (1995).
The ethylene response mediator ETR1 from Arabidopsis forms a disulfide-linked
dimer. J. Biol. Chem. **270**: 12526-12530.

Schott-Verdugo, S., Müller, L., Classen, E., Gohlke, H., Groth, G. (2019). Structural
model of the ETR1 ethylene receptor transmembrane sensor domain. Sci. Rep.
9: 8869.

Shin, L.-J,. Lo, J.-C., Yeh, K.-C. (2012). Copper chaperone antioxidant protein1 is
essential for copper homeostasis. Plant Physiol. **159**: 1099-1110.

Srivastava, M., Srivastava, A.J., Orosa-Puente, B., Campanaro, A., Zhang, C.,
Sadanandom, A. (2020). SUMO conjugation to BZR1 enables brassinosteroid
signaling to integrate environmental cues to shape plant growth. Curr. Biol. **30**:
1410-1423.

Thompson, H.L., Shen, W., Matus, R., Kakkar, M., Jones, C., Dolan, D., Grellscheid,
S., Yang, X., Zhang, N., Mozaffari-Jovin, S., Chen, C., Zhang, X., Topping, J.F.,
Lindsey, K. (2023). MERISTEM-DEFECTIVE regulates the balance between

stemness and differentiation in the root meristem through RNA splicing control.

Development **150**: dev201476.

Tian, T., Liu, Y., Yan, H., You, Q., Yi, X., Du, Z., Xu, W., Su, Z. (2017). agriGO v2.0: a

GO analysis toolkit from the agricultural community, 2017 update. Nucl. Acids

Res. **45**: W122-W129.

Yu, C.H., Yang, N., Bothe, J., Tonelli, M., Nokhrin, S., Dolgova, N.V., Braiterman, L.,

Lutsenko, S., Dmitriev, O.Y. (2017). The metal chaperone Atox1 regulates the

activity of the human copper transporter ATP7B by modulating domain dynamics.

J. Biol. Chem. **292**: 18169.

Voinnet, O., Rivas, S., Mestre, P., Baulcombe, D. (2003). An enhanced transient

expression system in plants based on suppression of gene silencing by the p19

protein of tomato bushy stunt virus. Plant J. **33**: 949-956.

Williams, L.E., Mills, R.F. (2005). P(1B)-ATPases: an ancient family of transition metal

pumps with diverse functions in plants. Trends Plant Sci. **10**: 491–502.

Willoughby, A.C., Nimchuk, Z.L. (2021). WOX going on: CLE peptides in plant

development. Curr. Opin. Plant Biol. **63**: 102056.

Xiao, Z., Brose, J., Schimo, S., Ackland, S.M., La Fontaine, S., Wedd, A.G. (2011).

Unification of the copper(I) binding affinities of the metallo-chaperones Atx1,

Atox1, and related proteins: detection probes and affinity standards. J. Biol.

Chem. **286**: 11047-11055.

Woest, K.E., Kieber, J.J. (2000) A strong loss-of-function mutation in RAN1 results in

constitutive activation of the ethylene response pathway as well as a rosette-

lethal phenotype. Plant Cell **12**: 443-455.

Zhong, S., Lin, Z., Grierson, D. (2008). Tomato ethylene receptor-CTR interactions:

visualization of NEVER-RIPE interactions with multiple CTRs at the endoplasmic

reticulum. J. Exp. Bot. **59**: 965-972.

Figure Legends

Figure 1. The PLS peptide is required for ethylene control of seedling growth.

(**A**). Wild type (left) and *pls* mutant (right); bar = 5 mm. (**B-C**) Heat maps showing
expression levels of 32 ethylene-responsive genes in *pls* (**B**) and of 58 genes in *PLS*
overexpressing (PLSox) seedlings (**C**), compared to wild type levels. Data for 3
biological replicates are shown and are expressed as $\log_2$-fold changes in *pls* mutant
and PLSox seedlings compared wildtype with respective significance levels (*P-adj*
value) provided in Tables S1 and S2. A $P < 0.05$ and $\log_2$-fold of change of ± 0.5 was
chosen to identify differentially expressed genes.

Figure 2 The PLS peptide is structurally and functionally conserved and
complements the *Arabidopsis pls* mutant.

(**A**). Amino acid sequence of the PLS peptide from *Arabidopsis thaliana* (*A.t*) with
Camelina sativa PLS sequence (*C.s*), and synthetic truncations N1, N2, C1 and C2,
indicated by horizontal lines. Two cysteine residues are highlighted in bold. (**B**) Effect
of *Arabidopsis* PLS full length peptide, *A.t.* PLS(FL), and *Camelina* PLS peptide (*C.s.*
PLS) on *Arabidopsis* primary root length. Wildtype (C24; blue bars) and *pls* mutant
seedlings (red bars) were grown hydroponically in the presence (100 nM) or absence
of peptide for 10 d. ***: $P < 0.05$, *t*-test between C24 and *pls*, n = 20-25. (**C**) Effect of
PLS full length and truncated peptides on wild type (blue bars) and *pls* mutant (red
bars) *Arabidopsis* primary root length. Seedlings were grown hydroponically in the
presence of 50 nM peptide for 10 d. C1 = C-terminal 14 amino acids, C2 = C-terminal
24 amino acids, N1 = N-terminal 22 amino acids, N2 = N-terminal 9 amino acids, full
length PLS = 36 amino acids. Error bars show ± 1 standard error, n = 25. ***: $P < 0.05$,
t-test between C24 and *pls*, n = 25.

Figure 3. PLS localizes to the endoplasmic reticulum.

(**A-O**) PLS::PLS:GFP fusion protein (**A, G, J**) colocalizes with endoplasmic reticulum
markers ER Tracker (**B, C, C inset**) and RFP:HDEL (**H, I, I inset**), but free GFP does
not (**D-F**, **M-O**). PLS:GFP staining is seen also in nuclei (n). PLS::PLS:GFP (**J**) does
not co-localize with the *trans*-Golgi markers ST-mCherry SH:GFP (**K, L**). Scale bars =
25 µm (**C, I**), 10 µm (**F, I, O**). Root epidermal cells in the transition zone were imaged.
(**p**) Ratiometric analysis of roGFP2 fusion constructs transiently expressed in *N.*
benthamiana. Comparison of excitation ratios of PLS-roGFP2 and roGFP2-PLS with

control constructs (free roGFP, SEC22 fusions) reveals that PLS localizes to the
cytosolic side of the ER.

Figure 4. PLS interacts with the ethylene receptor ETR1.

(**A**) Co-immunoprecipitation of PLS:GFP by ETR1:HA (upper panel) in leaves of
Nicotiana benthamiana, in the presence and absence of 0.5 µM $CuSO_4$ and EDTA (to
remove Cu). Lower panels show presence of ETR1:HA in extracts using anti-HA
antibody, and protein input blots using anti-GFP and anti-Rubisco. (**B**) Densitometric
scan of immunoblot. (**C**) Competition assay showing a reduced binding between
PLS:GFP and ETR1:HA in the presence of 0, 5 nM or 25 nM PLS peptide, in the
presence of 0.5 µM $CuSO_4$ and 50 µM MG-132, a proteasome inhibitor (upper panel).
Lower panel shows ETR1:HA in extracts using anti-HA antibody. (**D**) Densitometric
scan of immunoblot. (**E**) Co-immunoprecipitation of ETR1:HA (upper panel) in leaves
of *N. benthamiana*, showing the effect of EDTA (to remove Cu) on interaction between
ETR1:HA and PLS:GFP. Lower panels shows presence of ETR1:HA in extracts using
anti-HA antibody, and protein input blots using anti-ETR1-HA and anti-Rubisco. (**F**)
Densitometric scan of immunoblot. α-GFP, anti-GFP antibody; α-HA, anti-HA
antibody. (**G**) Microscale thermophoresis binding curves of different ETR1 truncations
with PLS. Binding of PLS was observed with full-length ETR1 and all C-terminal
truncations but not with ETR1$^{306\text{-}738}$ lacking the N-terminal transmembrane part of the
receptor.

Figure 5. PLS binds copper and interacts with RAN1 and copper chaperones.

(**A**) C6S and C17S in PLS are required for function. Seedlings were grown
hydroponically for 10 d in the presence (100 nM) or absence of peptide (FL or C6S,
C17S). Blue bars are C24, red bars are *pls*. *: $P <0.05$, *t*-test between C24 and *pls*, n =
25. Error bars show ± 1 standard error, n = 14-22. (**B**) BCA absorbance in the
presence of synthetic PLS (44.8 µM, filled symbols) or an equivalent volume (to PLS)
of DMSO (open symbols). Experiments with PLS used 88.3 µM (circles), 84 µM
(triangles) and 88.4 µM (squares) BCA. The control experiment with DMSO used 85.3
µM BCA. (**C**) UV-vis apo subtracted difference spectra of synthetic PLS (19.8 µM)
upon titration with CuCl (top left); binding isotherm of the feature at 242 nm (top right).

Fluorescence spectra of PLS (19.8 µM) upon titration with CuCl (bottom left); binding isotherm of the feature at 306 nm (bottom right). (**D**) Absorption of BCA (17.3 µM) titrated with Cu(I) (representative spectrum, n = 2, full data in Fig. S6). (**E**) Binding isotherms ($A_{358\,nm}$) of BCA (10 µM) in the presence/absence (filled/empty symbols, respectively) of 14 µM MBP-PLS (circles) or MBP-PLS mutants (C6S, C17S and C6S/C17S: triangles, diamonds and squares respectively) titrated with Cu(I) (n = 3, ± SD). Arrow indicates ca. 7 µM withholding of Cu(I) from BCA. (**F**) Binding isotherms ($A_{358\,nm}$) of BCA (50 µM) in the presence/absence of 10 µM MBP-PLS (filled/empty symbols, respectively) titrated with Cu(I). Model (solid line) describes Cu(I)-binding as a 2:1 complex, with β_2 affinity of 3.79 (± 1.5) $\times 10^{19}$ M^{-2}. Dotted lines simulate 10x weaker or tighter affinity (n = 3, ± SD). (**G**) Simulated Cu(I) occupancy (Supplementary Text) as a function of [PLS] using β_2 Cu(I) affinity of 3.79 $\times 10^{19}\,M^{-2}$. Inset, *Arabidopsis* ATX1 (20 µM, filled circles) withholds one Cu(I) equivalent from 20 µM BCA (open circles, BCA-alone) (n = 3, ±SD), with K_D Cu(I) 5.47 $\times 10^{-18}$ M (Fig. S3).

Figure 6. Model of the role of PLS in ethylene signalling.

In this model, we assume that RAN1 and ETR1 are in the same membranous compartment (likely the ER) at least transiently. RAN1 is believed to pump Cu(I) into the ER lumen, and then to ETR1. In air (absence of ethylene; left panel), the receptor complex is active and activates CTR1, which is a negative regulator of ethylene responses. In the presence of ethylene (centre panel), the receptor is in an inactive state, cannot activate CTR1, resulting in ethylene responses. In the Model for PLS function, PLS binds to ETR1 (with binding potentially enhanced by the Cu(I) in the ETR1 transmembrane domain) and affects the receptor conformation, potentially blocking ethylene binding to ETR1 or inducing some other conformational change, leading to active ETR1, active CTR1 and reduced ethylene signalling. This model is consistent with the observed *pls* loss of function mutant showing enhanced ethylene responses.

Figure 1. The PLS peptide is required for ethylene control of seedling growth.

(**A**). Wild type (left) and *pls* mutant (right); bar = 5 mm. (**B-C**) Heat maps showing expression levels of 32 ethylene-responsive genes in *pls* (**B**) and of 58 genes in *PLS* overexpressing (PLSox) seedlings (**C**), compared to wildtype levels. Data for 3 biological replicates are shown, and are expressed as log$_2$-fold changes in *pls* mutant and PLSox seedlings compared wildtype with respective significance levels (*P-adj* value) provided in Tables S1 and S2. A *P* < 0.05 and log$_2$-fold of change of ± 0.5 was chosen to identify differentially expressed genes.

Fig. 1

Figure 2 The PLS peptide is structurally and functionally conserved and complements the *Arabidopsis pls* mutant.
(**A**). Amino acid sequence of the PLS peptide from *Arabidopsis thaliana* (*A.t*) with *Camelina sativa* PLS sequence (*C.s*), and synthetic truncations N1, N2, C1 and C2, indicated by horizontal lines. Two cysteine residues are highlighted in bold. (**B**) Effect of *Arabidopsis* PLS full length peptide, *A.t.* PLS(FL), and *Camelina* PLS peptide (*C.s.* PLS) on *Arabidopsis* primary root length. Wildtype (C24; blue bars) and *pls* mutant seedlings (red bars) were grown hydroponically in the presence (100 nM) or absence of peptide for 10 d. ***: *P* <0.05, *t*-test between C24 and *pls*, n = 20-25. (**C**) Effect of PLS full length and truncated peptides on wild type (blue bars) and *pls* mutant (red bars) *Arabidopsis* primary root length. Seedlings were grown hydroponically in the presence of 50 nM peptide for 10 d. C1 = C-terminal 14 amino acids, C2 = C-terminal 24 amino acids, N1 = N-terminal 22 amino acids, N2 = N-terminal 9 amino acids, full length PLS = 36 amino acids. Error bars show ± 1 standard error, n = 25. ***: *P* <0.05, *t*-test between C24 and *pls*, n = 25.

Fig. 2

Fig. 3

Figure 3. PLS localizes to the endoplasmic reticulum.
(**A-O**) PLS::PLS:GFP fusion protein (**A, G, J**) colocalizes with endoplasmic reticulum markers ER Tracker (**B, C, C inset**) and RFP:HDEL (**H, I, I inset**), but free GFP does not (**D-F**, **M-O**). PLS:GFP staining is seen also in nuclei (n). PLS::PLS:GFP (**J**) does not co-localize with the *trans*-Golgi markers ST-mCherry SH:GFP (**K, L**). Scale bars = 25 μm (**C, L**), 10 μm (**F, I, O**). Root epidermal cells in the transition zone were imaged. (**P**) Ratiometric analysis of roGFP2 fusion constructs transiently expressed in *N. benthamiana*. Comparison of excitation ratios of PLS-roGFP2 and roGFP2-PLS with control constructs (free roGFP, SEC22 fusions) reveals that PLS localizes to the cytosolic side of the ER.

A

GFP PLS-GFP GFP PLS-GFP

| + | + | – | + | EDTA |
| – | – | + | + | Cu |

kDa

35 —
25 —   αGFP   PLS-GFP

100 —
70 —   αHA   ETR1-HA

35 —   αGFP   PLS-GFP
25 —   GFP   } input

Rubisco

B

PLS-GFP / ETR-HA

+ EDTA
– EDTA

GFP PLS-GFP GFP PLS-GFP

C

GFP PLS-GFP GFP PLS-GFP

0	0	25	5	[PLS] (nM)
+	+	+	+	Cu
+	+	+	+	MG-132

KDa

35 —
25 —   αGFP   PLS-GFP

100 —
70 —   αHA   ETR1-HA

D

PLS_GFP / ETR_HA

GFP   PLS C-   PLS_25nM   PLS_5nM

G

PLS + ETR1

Fraction bound

PLS [nM]

PLS_GFP / ETR_HA

GFP   PLS C-   PLS_25nM   PLS_5nM

PLS + ETR1

● ETR1$^{1-788}$ (WT)
  1898 ± 339 nM

■ ETR1$^{1-157}$
  1516 ± 272 nM

△ ETR1$^{1-307}$
  1785 ± 382 nM

Figure 4. PLS interacts with the ethylene receptor ETR1.
(**A**) Co-immunoprecipitation of PLS:GFP by ETR1:HA (upper panel) in leaves of *Nicotiana benthamiana*, in the presence and absence of 0.5 µM $CuSO_4$ and EDTA (to remove Cu). Lower panels show presence of ETR1:HA in extracts using anti-HA antibody, and protein input blots using anti-GFP and anti-Rubisco. (**B**) Densitometric scan of immunoblot. (**C**) Competition assay showing a reduced binding between PLS:GFP and ETR1:HA in the presence of 0, 5 nM or 25 nM PLS peptide, in the presence of 0.5 µM $CuSO_4$ and 50 µM MG-132, a proteasome inhibitor (upper panel). Lower panel shows ETR1:HA in extracts using anti-HA antibody. (**D**) Densitometric scan of immunoblot. (**E**) Co-immunopreciptation of ETR1:HA (upper panel) in leaves of *N. benthamiana*, showing the effect of EDTA (to remove Cu) on interaction between ETR1:HA and PLS:GFP. Lower panels shows presence of ETR1:HA in extracts using anti-HA antibody, and protein input blots using anti-ETR1-HA and anti-Rubisco. (**F**) Densitometric scan of immunoblot. a-GFP, anti-GFP antibody; a-HA, anti-HA antibody. (**G**) Microscale thermophoresis binding curves of different ETR1 truncations with PLS. Binding of PLS was observed with full-length ETR1 and all C-terminal truncations but not with ETR1$^{306-738}$ lacking the N-terminal transmembrane part of the receptor.

Fig 5

Figure 5. PLS binds copper and interacts with RAN1 and copper chaperones.
(**A**) C6S and C17S in PLS are required for function. Seedlings were grown hydroponically for 10 d in the presence (100 nM) or absence of peptide (FL or C6S, C17S). Blue bars are C24, red bars are *pls*. * *P* <0.05, *t*-test between C24 and *pls*, n = 25. Error bars show $\pm$ 1 standard error, n = 14-22. (**B**) BCA absorbance in the presence of synthetic PLS (44.8 µM, filled symbols) or an equivalent volume (to PLS) of DMSO (open symbols). Experiments with PLS used 88.3 µM (circles), 84 µM (triangles) and 88.4 µM (squares) BCA. The control experiment with DMSO used 85.3 µM BCA. (**C**) UV-vis apo subtracted difference spectra of synthetic PLS (19.8 µM) upon titration with CuCl (top left); binding isotherm of the feature at 242 nm (top right). Fluorescence spectra of PLS (19.8 µM) upon titration with CuCl (bottom left); binding isotherm of the feature at 306 nm (bottom right). (**D**) Absorption of BCA (17.3 µM) titrated with Cu(I) (representative spectrum, n = 2, full data in Fig S6). (**E**) Binding isotherms ($A_{358\,nm}$) of BCA (10 µM) in the presence/absence (filled/empty symbols, respectively) of 14 µM MBP-PLS (circles) or MBP-PLS mutants (C6S, C17S and C6S/C17S: triangles, diamonds and squares respectively) titrated with Cu(I) (n = 3, $\pm$SD). Arrow indicates ca. 7 µM withholding of Cu(I) from BCA. (**F**) Binding isotherms ($A_{358\,nm}$) of BCA (50 µM) in the presence/absence of 10 µM MBP-PLS (filled/empty symbols, respectively) titrated with Cu(I). Model (solid line) describes Cu(I)-binding as a 2:1 complex, with β_2 affinity of 3.79 ($\pm$1.5) x10^{19} M^{-2}. Dotted lines simulate 10x weaker or tighter affinity (n = 3, $\pm$SD). (**G**) Simulated Cu(I) occupancy (Supplementary Text) as a function of [PLS] using β_2 Cu(I) affinity of 3.79 x10^{19} M^{-2}. Inset, *Arabidopsis* ATX1 (20 µM, filled circles) withholds one Cu(I) equivalent from 20 µM BCA (open circles, BCA-alone) (n = 3, $\pm$SD), with K_D Cu(I) 5.47 x10^{-18} M (Fig. S3).

In air

In presence of ethylene

Model of PLS action: PLS binds ETR1 to prevent ethylene binding or conformational change

RAN1

Cu^+

Cu^+ Cu^+

Cu^+

ETR1

CTR1

RAN1

Cu^+

C_2H_4

Cu^+ Cu^+

Cu^+

ETR1

CTR1

RAN1

Cu^+

C_2H_4

PLS

Cu^+ Cu^+

Cu^+

ETR1

CTR1

-reduced/no ethylene bound to ETR1
-ETR1 active
-CTR1 active
-reduced ethylene response

-ethylene bound to ETR1
-ETR1 inactive
-CTR1 inactive
-ethylene response

-reduced/no ethylene bound to ETR1
-ETR1 active
-CTR1 active
-reduced ethylene response

Figure 6. Model of the role of PLS in ethylene signalling.

In this model, we assume that RAN1 and ETR1 are in the same membranous compartment (likely the ER) at least transiently. RAN1 is believed to pump Cu(I) into the ER lumen, and then to ETR1. In air (absence of ethylene; left panel), the receptor complex is active and activates CTR1, which is a negative regulator of ethylene responses. In the presence of ethylene (centre panel), the receptor is in an inactive state, cannot activate CTR1, resulting in ethylene responses. In the Model for PLS function, PLS binds to ETR1 (with binding potentially enhanced by the Cu(I) in the ETR1 transmembrane domain) and affects the receptor conformation, potentially blocking ethylene binding to ETR1 or inducing some other conformational change, leading to active ETR1, active CTR1 and reduced ethylene signalling. This model is consistent with the observed *pls* loss of function mutant showing enhanced ethylene responses.