1 **Improved RNA stability estimation through Bayesian modeling reveals most**
2 ***Salmonella* transcripts have sub-minute half-lives**
3

4 Laura Jenniches[1], Charlotte Michaux[2], Linda Popella[2], Sarah Reichardt[1], Jörg Vogel[1,2,3],
5 Alexander J. Westermann[1,2], Lars Barquist[1,3,*]
6
7 [1] Helmholtz Institute for RNA-based Infection Research (HIRI), Helmholtz Centre for Infection
8 Research (HZI), Würzburg, Germany
9 [2] University of Würzburg, Institute of Molecular Infection Biology (IMIB), Würzburg, Germany
10 [3] University of Würzburg, Faculty of Medicine, Würzburg, Germany
11
12 *Correspondence to: lars.barquist@helmholtz-hiri.de

Abstract (238/250 words)
RNA decay is a crucial mechanism for regulating gene expression in response to environmental
stresses. In bacteria, RNA-binding proteins (RBPs) are known to be involved in post-
transcriptional regulation, but their global impact on RNA half-lives has not been extensively
studied. To shed light on the role of the major RBPs ProQ and CspC/E in maintaining RNA
stability, we performed RNA sequencing of *Salmonella enterica* over a time course following
treatment with the transcription initiation inhibitor rifampicin (RIF-seq) in the presence and
absence of these RBPs. We developed a hierarchical Bayesian model that corrects for
confounding factors in rifampicin RNA stability assays and enables us to identify differentially
decaying transcripts transcriptome-wide. Our analysis revealed that the median RNA half-life in
Salmonella in early stationary phase is less than 1 minute, a third of previous estimates. We
found that over half of the 500 most long-lived transcripts are bound by at least one major RBP,
suggesting a general role for RBPs in shaping the transcriptome. Integrating differential stability
estimates with CLIP-seq revealed that approximately 30% of transcripts with ProQ binding sites
and more than 40% with CspC/E binding sites in coding or 3' untranslated regions decay
differentially in the absence of the respective RBP. Analysis of differentially destabilized
transcripts identified a role for ProQ in the oxidative stress response. Our findings provide new
insights into post-transcriptional regulation by ProQ and CspC/E, and the importance of RBPs in
regulating gene expression.

Significance Statement (117/120 words)
Together with transcription and translation, RNA decay is one of the major processes governing
protein production. Here, we have developed a new statistical approach that corrects for
confounding effects when estimating RNA decay rates from RNA-seq in bacteria. Our more
accurate decay rate estimates indicate that *Salmonella* transcripts have half-lives about three
times shorter than previously thought. This approach allowed us to measure the effects of RNA-
binding proteins (RBPs) on decay rates, identifying large cohorts of transcripts with changes in
stability following RBP deletion and conditions where post-transcriptional regulation affects
survival. Our method should lead to a reevaluation of RNA stability estimates across diverse
bacteria and new insights into the role of RBPs in shaping the transcriptome.

44
45
Introduction

47
Rapid adaptation of the proteome to environmental conditions is essential for the survival of

microorganisms. RNA degradation is an important post-transcriptional process directly

influencing protein abundance. The lifetime of bacterial RNA ranges from seconds to an hour (1)

and depends on numerous factors, including transcript identity, genotype and growth condition

(2). RNA-binding proteins (RBPs) in bacteria include structural components of the ribosome and

global post-transcriptional regulators such as Hfq (3, 4) and CsrA (5) which play key roles in

modulating translation and RNA stability in concert with a network of small RNAs (sRNAs) (6,

7). Beyond these model RBPs, recent years have seen the discovery of a menagerie of

bacterial RBPs that bind hundreds or even thousands of transcripts (8–10), though their

functions in shaping the transcriptome remain unclear.

In *Salmonella enterica* serovar Typhimurium (henceforth *Salmonella)*, these recently

identified global RBPs include the less studied FinO-domain containing protein ProQ and the

cold-shock proteins CspC and CspE. While individual deletion of any of these RBPs does not

lead to clear growth phenotypes under standard laboratory conditions (11, 12), mounting

evidence suggests they play important regulatory roles. ProQ has been shown to bind hundreds

of mRNAs and sRNAs (13–15), affecting important biological processes including expression of

virulence factors (12) and formation of antibiotic persisters (16). ProQ has also been shown to

be involved in determining the outcome of regulatory interactions between some sRNAs and

target transcripts (12, 15, 17), though many of the mechanistic details of these interactions

remain unclear (18). CspC and CspE have been shown to play partially redundant roles in

virulence, affecting survival in mice, motility, biofilm formation, and survival of bile stress (11,

19). The molecular details of how these RBPs affect phenotype are not clear, although at least

some of the effects of ProQ and CspC/E are mediated by the direct modulation of mRNA

stability. For instance, CspC/E have been shown to stabilize the mRNA of the bacteriolytic

lipoprotein EcnB by blocking digestion by the endonuclease RNase E (11). ProQ on the other

hand appears to preferentially bind 3′ UTRs where in a few cases it has been shown to protect

transcripts from exonuclease activity (14, 20).

While these results provide hints at the mechanisms by which RBPs regulate target

gene expression, in the absence of transcriptome-wide differential RNA stability measurements

it remains unclear how common regulation through stability modulation is. A classical approach

to study RNA stability is to halt transcription with the transcription initiation inhibitor rifampicin

(21) and monitor RNA decay over time. This approach has been scaled to the whole

transcriptome by combining it with microarrays (22, 23) and high-throughput sequencing (24).

However, the presence of non-linear effects in the resulting time-course data makes inference

of differences in decay rates between experimental conditions difficult.

RNA-seq analysis tools such as limma (25), edgeR (26), and DEseq (27) solve the

problem of accurately estimating dispersion in experiments with many measurements but few

replicates through an empirical Bayes approach (28). In empirical Bayes, information is pooled

across transcripts under the assumption that transcripts with similar concentrations will exhibit

similar biological and technical variation across samples, leading to more robust dispersion

estimates. However, these tools are currently restricted to linear models, which limits the

structure of models that can be implemented and hence the complexity of effects that can be

considered. Recent progress in the optimization of sampling methods has made the

development of fully Bayesian hierarchical models increasingly efficient and accessible. In

particular, the Stan probabilistic programming language (29) separates model description from

sampler implementation, allowing easy development and testing of complex hierarchical

models. This provides a powerful framework for developing analysis methods for sequencing

data that can accommodate complex experimental techniques.

Here, we investigate the effects of the undercharacterized RBPs ProQ and CspC/E on

RNA stability across the entire transcriptome, starting from a fully Bayesian analysis of

rifampicin treatment followed by RNA sequencing (RIF-seq). During model development, we

discovered that accounting for confounding factors in stability assays conducted after rifampicin

treatment dramatically affects the inferred half-life, leading us to substantially revise estimates

for average mRNA half-life in *Salmonella* to less than 1 minute, compared to previous estimates

in the range of 2 to 7 minutes in the closely related species *E. coli* (22–24). We develop a

hypothesis testing procedure for determining differential decay rates that allows us to identify

hundreds of gene transcripts destabilized in the absence of ProQ and CspC/E. We combine our

differential stability estimates with other high-throughput datasets available for *Salmonella* to

further characterize RBP interactions, identifying a role for ProQ in survival of oxidative stress.

We additionally find a substantial population of long-lived transcripts that depend on RBPs for

their stability, illustrating the importance of RBPs in shaping the bacterial transcriptome. Beyond

its utility in investigating RBP interactions, our improved approach to determining transcript half-

life suggests that RNA stability in bacteria has generally been overestimated and will need to be

reassessed in other bacterial species.

Results

A progressive Bayesian analysis revises RNA half-lives

To determine transcriptome-wide half-lives under an infection-relevant condition, we applied RIF-seq to *Salmonella* at early stationary phase (ESP) where host invasion genes are expressed (30). Our RIF-seq workflow for data production and analysis is illustrated in **Figure 1A**: wild-type and isogenic RBP deletion strains were treated with rifampicin, and cellular RNA samples were collected over time to capture RNA decay dynamics. We collected data from eight time points following rifampicin treatment in three (Δ*cspC/E*), six (Δ*proQ*), or nine (wild-type) replicates (see Methods). We included ERCC RNA spike-ins (31) for normalization between samples. Additionally, we developed a center-mean normalization technique to remove batch effects between replicate samples (**Figure S2;** *Methods*). Subsequently, we fitted a Bayesian statistical model to the normalized data using Hamiltonian Monte Carlo with Stan (29).

We employed a progressive Bayesian workflow to arrive at our final model (**Figure 1B**). An advantage of Bayesian analysis is that it allows the modeler to formalize their beliefs about the data generating process and provides a variety of tools for model comparison and selection. In the case of RIF-seq data, the simplest expectation would be that RNA concentrations would exhibit a linear decay on a semilog scale, which could be fit by a simple linear model with gene- and condition-dependent decay rate β. While some of our observations met this expectation (**Figure S3A**), the vast majority of transcripts exhibited more complex dynamics that prevent accurate extraction of decay rates with a linear model (**Figure 1C; S3A**), leading to large unexplained variation at late time points (**Figure S3L**). To account for this, we introduced additional parameters that capture confounding effects in the data. The first confounding effect is a gene-dependent delay parameter γ, which captures the delay commonly observed in RIF-seq data before decay initiates (**Figure 1B&C**, in green). As has been previously described, this is due to ongoing transcription from RNA polymerase already bound to DNA, which rifampicin does not block (24, 32). The ongoing transcription compensates for decay, manifesting as a delayed decay, though our delay parameter may also capture unrelated effects such as the time needed for rifampicin to penetrate and act within cells. To support the relationship between ongoing transcription and the delay parameter, we performed an analysis of elongation times on 60 base sub-genic windows, finding a clear association between the estimated elongation time and distance to annotated transcription start sites (**Figure S3B**). We used this association to infer transcription rates from our data set (see **Figure S3C**, Methods) finding a median transcription rate of 22.2 nt/s (**Figure S3D**), comparable to previous estimates in *E. coli* (24).

The second confounding effect we corrected for was an apparent gene- and condition-
dependent baseline RNA concentration π beyond which no further decay was observed (**Figure**
1B&C, orange). We were initially concerned that this effect may be an artifact of the
pseudocount we used to avoid dividing by zero in our calculations; however, inspection of a
number of decay curves illustrated that the observed baseline was generally well above the
detection threshold (**Figure S3E**, see Methods). We also verified that the half-life of a transcript
is generally constant along an operon (**Figure S3G**). In agreement with previous work (33) we
found a small number of stable subregions which generally corresponded to known sRNAs (e.g.
the FtsO sRNA excised from the *ftsI* mRNA, **Figure S3H**), but since this was not a general
feature of transcripts we excluded this as a source of the observed baseline. To confirm that the
baseline is not a result of our sequencing protocol, we used independent northern blot
quantifications from a rifampicin treatment time course including late time points from previous
studies (13, 14). These quantifications reproduced the observed baseline effect (**Figure**
1D&S3I-J), illustrating that this is a general feature of rifampicin RNA stability assays. Similar
artifacts can be observed in published northern blot quantifications of rifampicin-treated
Salmonella and *E. coli* in both ESP (34, 35) and during exponential growth (36). For wild-type
Salmonella, we find that a median of 2.6% of the initial transcript concentration appears
resistant to decay (**Figure S3K**), and that the exponential decay regime ends at different
timepoints for different transcripts and genotypes (**Figure S6AB&C**). Whether this fraction is
truly resistant to degradation or just degrades much slower than the rest of the transcript
population is unclear. However, the median fraction of baseline RNA increases to 5.7% when
proQ is overexpressed (**Figure S3K**), suggesting that nonspecific RBP-RNA interactions may
play a role in degradation resistance.
To compare the models with and without these two confounding factors, we calculated
the difference in the expected log pointwise predictive density (ELPD), a measure of the
expected predictive accuracy of a model on out-of-sample data, using Pareto-smoothed
importance sampling approximate leave-one-out cross-validation (PSIS-LOO, see Methods)
(37). Comparing the difference in ELPD between a simple linear model, the piecewise linear
model correcting only for extension time, and the full model (henceforth *log-normal* model)
including the baseline correction showed a clear preference for the log-normal model,
particularly at late timepoints (**Figure S3L&M**). Additionally, examination of the fitted variance
unexplained by our decay model, σ_g, illustrated the log-normal model captured the behavior of
late timepoints better than the piecewise model (**Figure S3N-P**). Correcting for confounding
factors has major implications for transcriptome-wide estimates of decay rates: while the linear

and piecewise linear models produced median half-life estimates of 2.7 and 2.5 minutes,

respectively, our final log-normal model estimates a median half-life of 0.9 minutes (**Figure 1E**).

To investigate whether transcripts encoding proteins involved in different cellular

functions systematically differ in their stability, we calculated average half-lives across clusters

of orthologous groups (COG) categories (38) (**Figure 1F; S5**). In agreement with previous work

(22, 23), transcripts for genes involved in energy production and carbohydrate metabolism

tended to be longer lived. We also found many sRNAs to have longer than average half-lives,

with several of the longest-lived also having a connection to metabolism (e.g. GlmZ (39), Spot42

(40), and CyaR (41)). Among the least stable transcripts were those coding for genes involved

in cell division (e.g. *ftsZ*) and DNA replication (e.g. *dnaA*, *dnaN*), suggesting tight control of their

cognate proteins. Taken together, accurately modeling RNA-decay curves led to drastically

reduced transcriptome-wide half-life estimates and allowed us to relate transcript stability to

gene function.

196
Figure 1. Pipeline and model description
(A) RIF-seq workflow: WT and ΔRBP strains are treated with rifampicin, cells are harvested at various
time points and subjected to RNA-seq. Read counts are normalized before the extraction of biologically

relevant parameters with a Bayesian model. Significant differences between strains are identified with
Bayesian p values. (B) A plate diagram of the Bayesian models in this study. The layers indicate which
indices and variables the parameters depend on. The LM is parametrized by the WT decay rate $\beta_{WT,g}$
(purple). In the PLM, the gene-wise elongation time γ_g (green) is added. The LNM adds a baseline
parameter π_{gc} (orange) which corresponds to the fraction of residual RNA ($\pi_{gc} \in [0,0.2]$). The WT decay
rate β_{WT} is a gene-wise parameter that is modeled hierarchically and depends on the hyperparameters
μ_β and σ_β. The difference in decay rate $\Delta\beta_{gc}$ depends on the strain or condition c. The scale parameter
$\sigma_g(t)$ captures variation by our decay model and depends on the time-dependent hyperparameters $\mu_\sigma(t)$
and $\sigma_\sigma(t)$. (C) Representative example of a decay curve in the LNM, illustrating regimes dominated by
the different model parameters. The period of transcription elongation γ is marked in green, the
exponential decay with decay rate β in purple and the constant regime governed by the fraction of
baseline RNA π in orange. (D) Comparison of RNA-seq and model fit with independent northern blot
quantifications for SibC (13). (E) Hyperpriors and median of transcriptome-wide WT half-lives in the three
Bayesian models. (F) Half-life distributions from the log-normal model for transcripts in selected COG
categories. Adjusted p-values (compared to the first COG) were calculated using the Wilcoxon rank sum
test.
216
217
218 **Steady-state abundance does not reflect changes in transcript half-life upon RBP**
219 **deletion**
220
To study the influence of the RBPs ProQ and CspC/E on transcript stability, we applied the log-
normal model to our RIF-seq data for the *proQ* and *cspC/E* deletion strains, as well as a *proQ*
overexpression strain (*proQ++*; see Methods). To prioritize transcripts with changes in stability,
we developed a hypothesis testing procedure based on examination of the posterior distribution
of the change in decay rate from the wild-type (**Figure S4A**) and estimated statistical
significance by calculating Bayesian p-values (**Figure S4B**). Since Bayesian p-value
distributions require calibration (42, 43), we used simulation studies to estimate the false
discovery rate (FDR) (**Figure S4C-E**, see Methods). To evaluate our differential stability
estimates, we examined known targets of ProQ (13, 14) and CspC/E (11) (**Figure 2A-C, S6A**).
For deletion of ProQ we were able to confirm destabilization of the *cspD*, *cspE*, and *ompD*
transcripts (**Figure 2A**), while the *cspC* transcript was hyperstabilized in the *proQ++* background
(**Figure 2B**) in agreement with previous northern analysis (14). Of the mRNAs known to be
stabilized by ProQ, we were only unable to detect destabilization for the *yfiD* transcript.
Similarly, we found the *ecnB* transcript destabilized following *cspC/E* deletion (**Figure 2C**) as
previously reported (11).
For both RBP deletions, we identify hundreds of transcripts with changes in stability at
an FDR of 0.1 (**Figure 2D&E**). Deletion of *cspC/E*, whose role in maintaining transcript stability

is less well explored, led to strong destabilization of a large cohort of transcripts (727), while

only stabilizing 164 (**Figure 2F**). Curiously, we identified more transcripts which were

significantly stabilized (413) than destabilized (276) following *proQ* deletion (**Figure 2G**), which

was unexpected as prior studies have focused on ProQ's stabilizing effect (13, 14).

Nevertheless, stabilized transcripts tended to have much smaller changes in half-life, with a

median change of 0.3 minutes (**Figure S6D**), compared to destabilized transcripts whose half-

lives changed by 0.7 minutes on average. Overexpression of *proQ* restored stability for ~95% of

transcripts identified as destabilized by proQ deletion, but additionally significantly stabilized one

fourth of the transcriptome (**Figure S6E&F**). The majority of the transcripts (> 90%) stabilized

upon *proQ* overexpression were not affected by *proQ* deletion, suggesting these effects are

largely non-specific.

A striking feature of our analysis of both strains was that changes in transcript half-life

are not clearly related to changes in steady-state abundance upon RBP deletion (**Figure 2D-G**).

In the *proQ* deletion strain, less than 10% of destabilized transcripts showed a statistically

significant decrease in steady-state abundance. While this number was higher for the *cspC/E*

deletion (~18%), it was still only a minor fraction of the total number of destabilized transcripts.

This might be explained by altered activity of other regulatory proteins. Deletion of either RBP

led to perturbation of the stability of transcripts encoding major regulatory proteins including the

anti-sigma factor Rsd*,* the transcription termination factors Rho and NusA, the alternative sigma

factor RpoS, the nucleoid-associated HupA/B, and the cAMP receptor protein CRP (**Figure**

S6B). For HupA/B and RpoS, we also observed reduced mRNA abundance in RBP deletion

strains (**Figure S14**). Hence, loss of ProQ or CspC/E likely has complex, and in some cases

indirect, effects on the global transcriptome. This suggests caution should be taken when

deducing direct regulatory interactions from differential expression analysis of RBP-deletion

mutants.

263
Figure 2. Differential analysis of transcript stability in the absence of CspC/CspE/ProQ
(A-C) Decay curves of all known ProQ (*cspD*, *cspE*, *ompD*, *cspC*) or CspC/E (*ecnB*) mRNA targets whose
stability changes were confirmed by our study. See **Figure S6A** for sRNAs. Significant stability changes
are marked with a star. (D) Relative difference in half-life vs. steady state log-fold changes between the
ΔcspCE and the WT strain. (E) Relative difference in half-life vs. steady state log-fold changes between
the ΔproQ and the WT strain. (F) Overlap between stability changes and steady-state log-fold changes in
the ΔcspCE strains. (G) Overlap between stability changes and steady-state log-fold changes in the
ΔproQ strain.

Integrating high-throughput datasets identifies cohorts of mRNAs subject to known RBP

regulatory mechanisms

The location of an RBP binding site within a transcript is often a key determinant of the

mechanism of RBP regulation. To investigate the potential mechanisms underlying the

stabilization activity of ProQ and CspC/E, we integrated our differential stability estimates with

UV crosslinking and immunoprecipitation followed by RNA sequencing (CLIP-seq), which can

localize RBP binding sites within a transcript. For ProQ, we reanalyzed an existing CLIP-seq

dataset (14), identifying 833 peaks indicative of binding (see Methods). We produced new

CLIP-seq datasets for both CspC and CspE and identified 1155 CspC and 861 CspE peaks,

spread across 571 and 462 target transcripts, respectively (**Figure 3A&B;S7A-C**). In total, 717

transcripts are bound by at least one CSP, with 430 CspC peaks directly overlapping with a

CspE peak (**Figure 3B**) supporting the previously reported partial redundancy between these

proteins (11) and similar observations in *E. coli* (44). We saw especially dense clusters of

CspC/E peaks in transcripts encoding genes involved in the TCA cycle, flagellar proteins, and

proteins involved in host invasion associated with the *Salmonella* pathogenicity island 1 (SPI-1)

type three secretion system (**Figure 3A**).

291   We next examined the distribution of RBP binding sites across target transcripts,

beginning with ProQ. As previously reported (14), ProQ binds predominantly at the end of

coding sequences, with half of detected binding sites within 100 nucleotides of the stop codon

(**Figure 3D;S7D**). Amongst those genes with 3′ binding sites, approximately 19% (86

transcripts) were significantly destabilized upon *proQ* deletion (**Table S3**). Besides the known

interaction of ProQ with the *cspE* mRNA, these include transcripts encoding the SPI-1 effectors

SopD and SopE2, involved in host cell invasion (45), and OxyR, a transcription factor involved

in the oxidative stress response (46). The location of these binding sites suggests that ProQ

may protect the 3′ ends of a large cohort of transcript from exoribonucleases attack, as

previously shown for individual model transcripts (14, 20).

301   In contrast to ProQ, CspC and CspE binding sites were spread across coding (CDS)

regions with only slight enrichment in the vicinity of the start and stop codons (**Figure 3C;S7D**).

We identified 177 transcripts with a CspC and/or CspE binding site in the CDS or 5'UTR that

were destabilized upon *cspC/E* deletion (**Table S4**). These included the *ecnB* transcript (**Figure

S10A**), which has previously been shown to bind CspC and CspE *in vitro* and to be protected

from RNase E by CspC/E *in vivo* (11). To further investigate the role of the CSPs in protection

from RNase E cleavage, we combined our stability and CLIP-seq data with a published dataset

mapping RNase E cleavage sites (47). We saw an enrichment of RNase E cleavage sites within

CspC/E CLIP-seq peaks (410/2059 compared to a median of 331/2059 across 100 simulations,

$p \approx 0$, see *Methods*), but the majority of CspC/E binding sites did not directly occlude known

RNase E cleavage sites. Furthermore, the presence of an RNase E cleavage site within a peak

did not appear to influence differential decay rates upon *cspC/E* deletion (**Figure S7E**). This

suggests that rather than directly protecting cleavage sites, CspC/E may interfere with RNase E

scanning (48). This is further supported by the fact that destabilized transcripts have a median

of two CspC/E binding sites, while ligands without stability changes have a median of one

binding site (**Figure S7F**), suggesting multiple CspC/E proteins must bind to create an

obstruction of sufficient size to interfere with RNase E scanning (49).

Figure 3. CspC/E CLIP-seq, comparison with RIF-seq results.
CspC/E CLIP-seq analysis: (A) Number of CspC/E binding sites binned by genomic position for the
positive (outer) and negative (inner) strand. The chromosome is indicated in purple and the three
plasmids in blue, green and yellow. (B) Venn diagram of binding sites, with shared targets defined by an
overlap by at least 12 bases between CspE and CspC sites. (C-D) Metagene plot of transcripts bound by
the respective RBP, ordered by position of CLIP-seq peak relative to the start/stop codon. Target
sequences are colored by the effect of RBP deletion on stability: destabilized (purple), stabilized (yellow),
or no differential decay (non-DD, green).

RBPs play overlapping and complementary roles in infection-relevant pathways

To investigate the physiological consequences of RBP deletion, we identified pathways

enriched in differentially stabilized and differentially expressed transcripts in the *proQ* and

cspC/E deletion strains with the GSEA algorithm (50) (**Figure 4A & S8A**). Intriguingly, we found

a large overlap in enriched gene sets in both deletion backgrounds that may in part reflect

destabilization of the *cspE* mRNA in the *proQ* deletion background (14). On the level of stability

this included responses to extracellular stimulus and oxidative stress, flagellar assembly, and

metabolite transport and utilization pathways including the phosphotransferase system and

glyoxylate and dicarboxylate metabolism. Several of these gene sets were also enriched in

differentially expressed transcripts, though the directions of the changes were often inconsistent

with the observed effects on stability. For instance, genes involved in flagellar assembly were

expressed at lower levels in both deletion strains despite their transcripts being stabilized

(**Figure 4A, S8B & S12**). Some pathways, such as aerobic and anaerobic respiration, showed

consistent changes in expression levels across both strains despite no clear shared enrichment

on the level of stability.

The large overlap in pathways affected at the level of stability and expression between

the two RBP deletion strains led us to investigate the relationship between ProQ- and CspC/E-

mediated regulation. We examined transcripts significantly differentially expressed in both

strains, finding a strong correlation between the steady-state log fold-changes (**Figure 4B**,

r=0.89). The slope of a line fitted to these changes indicated stronger average changes in the

cspC/E deletion; this was particularly clear for genes involved in flagellar assembly and

chemotaxis which exhibited an ~2-fold lower expression in the Δ*cspC/E* background compared

to Δ*proQ*. The lack of a similarly strong correlation for changes in transcript stability (**Figure**

S8C & D) suggested that some of the similarities in changes in steady-state mRNA abundance

between the two deletion strains may be due to indirect regulation, with that in the Δ*proQ*

background possibly mediated by changes in CspE expression.

Despite the large overlaps in mRNA stability and abundance changes, there were a

number of changes specific to each RBP, though these were often in the same pathways. For

instance, deletion of each RBP affected the stability of a discrete set of secreted effectors

involved in host cell invasion (**Figure S9D**). We also identified changes for transcripts involved

in *porin activity* (**Figure S13**), which may be related to ProQ's originally described role in

osmoregulation (51, 52). We found a ProQ CLIP-seq peak in the 5' UTR of the *proP* transcript,

but this did not appear to affect transcript stability or abundance, supporting a potential role in

regulation of translation (53). Both *proQ* and *cspC/E* deletion affected stability and expression of

the transcript for the major porin OmpD (**Figure 2A**), while effects on the transcripts of the well-

characterized osmolarity-responsive OmpF and OmpC appeared to be RBP-specific (**Figure**

S13). Another pathway with changes upon *proQ* and *cspC/E* deletion was in the oxidative stress

response, where we saw a stronger enrichment for destabilized transcripts in the Δ*proQ* strain

(**Figure 4A & S11**). Transcripts destabilized by *proQ* deletion included those encoding for the

oxidative stress regulator OxyR (**Figure S9B & C**), the superoxide dismutase SodB, the

catalase-peroxidase KatG, and the DNA protection during starvation protein Dps; however, few

of these transcripts showed significant differences in mRNA abundance.

To test if destabilization was predictive of phenotype, we exposed the Δ*proQ* and

proQ++ strains to varying concentrations of hydrogen peroxide, including a Δ*oxyR/S* strain as a

control. After exposure to 1.5mM H_2O_2 we saw a survival defect for both Δ*proQ* and *proQ++*

strains intermediate between wild-type survival and that of the Δ*oxyR/S* strain (**Figure S9D**).

However, this defect was concentration dependent: exposure to 2mM H_2O_2 led to a severe

survival defect in the Δ*proQ* strain that could be complemented by *proQ* overexpression (**Figure**

4C), while Δ*oxyR/S* behaved similarly to wild-type. This indicates that the Δ*proQ* survival defect

after exposure to high concentrations of H_2O_2 is independent of any effects ProQ has on the

stability of the *oxyR* transcript and likely depends on the effects of ProQ on other transcripts

involved in the oxidative stress response. The sensitivity of the Δ*proQ* strain to oxidative stress

also shows that changes in transcript stability can be predictive of RBP deletion phenotype,

even without corresponding changes in transcript abundance under standard growth conditions.

387
Figure 4. Integrative analysis of RBP binding and transcript stability
(A) Comparative analysis of pathways enriched in cspCE and proQ RIF-seq data. Pathways enriched in transcripts destabilized or with negative steady-state log-fold changes (min) upon RBP deletion are marked blue. Pathways enriched in stabilized transcripts or positive log-fold changes are marked brown.
(B) Genetic features with significant log-fold changes in both the *proQ* and the *cspC/E* deletion mutant.
(C) Exposure of Δ*proQ*, *proQ*++ and Δ*oxySR Salmonella* strains to 2mM of hydrogen peroxide.

Long-lived transcripts rely on global RBPs for their stability

To further examine the global impact of RBPs in shaping the transcriptome, we investigated the

relationship between our wild-type half-life estimates and RBP binding as determined by CLIP-

seq for five major *Salmonella* RBPs at early stationary phase: ProQ (14), CspC/E (this study),

Hfq and CsrA (54). After sorting transcripts by stability, we saw a clear association between

wild-type half-life and RBP binding, with over half of the 500 most stable transcripts ($t_{1/2} > 2.5$

min) bound by at least one RBP (**Figure 5A & S9E**). While the probability of detecting a CLIP-

seq peak increases with transcript abundance (**Figure S9F&G**), we found no correlation

between transcript abundance and stability (**Figure S3F**) suggesting the relationship between

RBP binding and stability is unlikely to be an artifact of our measurements. Long-lived

transcripts are also more likely to be destabilized upon RBP deletion than shorter-lived ones,

regardless of RBP-binding. Of the 500 most stable transcripts, 32% are significantly destabilized

in the absence of ProQ and 51% in the absence of CspC/E. Investigating the relationship

between transcript half-life and differential half-life upon RBP deletion revealed large changes in

median half-life for stable transcripts (**Figure 5B**), indicating that long-lived transcripts are not

only bound by RBPs but also rely on them for their stability.

Figure 5. Global effect of RBP binding on transcript stability

(A) Fraction of transcripts bound by RBPs. Transcripts were ordered by half-life. The distribution of half-

lives for each individual group can be found in **Fig. S9E**. (B) Relative change in half-life in the Δ*proQ* and

Δ*cspCE* strain for transcripts with a WT half-life of 0-2 min, 2-4 min, and greater than 4 min.

Discussion

Together with transcription and translation, mRNA degradation is one of the fundamental

processes controlling protein production in the cell. Rapid turnover of mRNAs underlies the

ability of bacteria to rapidly adapt to new conditions: as protein production is constrained by the

translational capacity of the available ribosome pool (55), clearance of transcripts encoding for

unneeded proteins is essential to change the composition of the proteome. Previous work

based on RNA-seq and microarray analysis of rifampicin time course data in *E. coli* and

Salmonella has reported average mRNA half-lives in the range of 2 to 7 minutes (22–24, 56–

58). The most similar prior RNA-seq study to our own reported an average half-life of 3.1

minutes across ~1200 transcripts in *E. coli* grown to stationary phase (24), over three times our

estimated average decay rate of 0.9 minutes in *Salmonella* at ESP. This discrepancy appears to

originate from not accounting for the baseline stable RNA concentration, leading to a systematic

underestimation of the decay rate. Interestingly, our estimates are in the range of those derived

from classic experiments that pulse radiolabeled bulk RNA and determined average mRNA half-

life to be ~0.7 minutes in exponentially growing *E. coli* (59), far shorter than any other

subsequent estimates based on high-throughput approaches. This rapid decay may in part

underlie the ability of bacteria to rapidly adapt their transcriptomes, as constant transcription

would be required to maintain mRNA concentrations. Across bacteria, mRNA half-lives have

primarily been determined by rifampicin treatment followed by sequencing or microarray

analysis in those bacteria where transcriptome-wide measurements are available (60) and

appear to contain similar artifacts to those observed here. For instance, biphasic decay curves

suggestive of a stable baseline transcript fraction have also been observed in the slow-growing

Mycobacterium smegmatis (61). Our results thus suggest that mRNA stability has likely been

widely overestimated and that a general reevaluation of bacterial transcript stability is in order.

Our hierarchical Bayesian analysis of RIF-seq data provides a principled framework for

the analysis of RNA turnover, including the determination of differential decay rates after

deletion of an RBP of interest. The flexibility of Bayesian analysis allowed us to account for

nonlinearities due to confounding factors like transcription elongation after rifampicin addition

and RNA baseline concentration, removing substantial biases in our determination of decay

rates. Despite our best efforts, it is likely that there are still some limitations to our analysis. For

instance, our control for false discovery rates means that we have likely missed some genuine

instances of differential decay. Our simulations suggest an 85% sensitivity for the most precisely

measured transcripts, but this falls to ~30% when considering the whole transcriptome (**Figure**

S4D). Other limitations may be due to uncontrollable effects in the data. For example, some

Salmonella promoters have previously been shown to respond specifically to subinhibitory

rifampicin (62), which could introduce some bias to decay rate estimates for affected transcripts

should similar effects occur with the rifampicin concentrations used here. Manual inspection of

our decay curves suggests this is unlikely to be a widespread problem in our data. Similarly, if

RBP deletion leads to modulation of expression of cellular RNases, our individual differential

decay rates may not be reflective of the differential decay induced by simple ablation of an RBP

binding site. Such a bias dependent on the cellular context of rifampicin treatment has

previously been observed for the sRNA RyhB whose stability critically depends on the presence

of its target mRNAs (63).

Regardless of potential biases, high-throughput methods provide at least one major

advantage over classical molecular approaches to RBP characterization: numbers. While our

approach does not definitively demonstrate a causal link between RBP binding and transcript

fate, we are able to provide high-confidence targets for future molecular characterization. Where

previously a small handful of transcripts were known to be stabilized by 3′ binding of ProQ in

Salmonella, we find 86 candidates. Similarly, we expand the number of transcripts known to be

stabilized by CspC/E from two to a predicted cohort of 177. By combining CLIP-seq (14) and

RNase E cleavage profiling (47) with our differential stability data, we have defined cohorts of

transcripts likely subject to particular modes of RBP regulation. Depending on the binding site

within a transcript, up to 44% (CspC/E, CDS) and 32% (ProQ, close to the stop codon) of direct

RBP targets showed altered stability upon deletion of the respective RBP. However, in both

cases transcripts stabilized by RBP binding are outnumbered by those apparently bound, but

unaffected at the level of stability, raising numerous questions about RBP function. How are

transcripts stabilized by RBPs differentiated from those that are not? Do RBP interactions that

do not affect stability perform other functions in the cell? Our analysis suggests CspC/E may

protect some transcripts from RNase E through a roadblock mechanism (48) which could be

investigated further by mapping Rnase E cleavage sites in the presence and absence of CspC

or CspE. (48) Additionally or alternatively, CspC/E targets may be regulated at the level of

translation (64) or antitermination (65) through the manipulation of mRNA secondary structure

as has been shown for the targets of other CSPs. Alternative roles of ProQ remain to be well

defined, but it has been shown to play a role in gene regulation by sRNAs (12, 17). By defining

and partially characterizing RBP targets, our data provides a starting point for the molecular

investigations needed to further define the functions of CspC/E and ProQ.

The degree to which post-transcriptional regulation shapes the bacterial proteome has

long been controversial. Recent work has suggested that, on average, protein concentrations

are primarily determined by promoter on rates with post-transcriptional regulation playing only a

minor role (66). Here in contrast, we have shown that deletion of bacterial RBPs thought to act

primarily at the post-transcriptional level leads to large changes in both RNA stability and

steady-state transcript concentration, and strong phenotypes have been observed for RBP

deletion in a variety of conditions (5, 11, 67). How can these findings be reconciled? Our data

provides at least two potential answers. First, as suggested by the effects of *proQ* and *cspC/E*

deletion on the stability of various global regulators (Figure S6B&S14), modulation of stability or

translation of single transcriptional regulators may ultimately cause phenotypic changes by

indirectly affecting the promoter on rates of a large cohort of transcripts. The lack of correlation

we observe between changes in steady-state RNA levels and differential stability (**Figure

2D&E**) indicates that such indirect effects are widespread. Secondly, our analysis shows that

the majority of RNA half-lives are concentrated at less than 1 minute (**Figure 1E**), and it is

indeed difficult to understand how further destabilization through post-transcriptional regulation

could have strong effects on translation. However, the half-life distribution is long tailed, with

~500 transcripts having half-lives of greater than 2.5 minutes and being preferentially bound by

RBPs (**Figure 4A**). The stability of this population of transcripts is strongly affected by RBP

deletion (**Figure 4B**), further suggesting they may be the major targets of post-transcriptional

regulation.

An accumulating body of work suggests that the post-transcriptional regulatory networks
scaffolded by RBPs are interconnected. At least two Hfq-dependent sRNAs also serve as
sponges for CsrA (68, 69), and RNA-RNA interactome studies have observed a substantial
fraction of shared targets between Hfq and ProQ (15). Regulatory interactions between cold
shock proteins (CSP) have long been observed, with deletion of particular CSPs leading to the
induction of others (11, 70), presumably through undescribed feedback mechanisms. The *cspE*
mRNA has previously been used as a model for understanding the molecular mechanism of
ProQ protection of 3′ ends (14); our results suggest some fraction of the change in steady-state
transcript levels observed in the *proQ* deletion strain may be the result of indirect regulation
through CspE (**Figure 5B**). Additionally, both RBPs affect the stability of mRNAs in similar
pathways (**Figure 5A**), though often by targeting different transcripts, as for the SPI-1 effectors
(**Figure S8B**). We also find effects for both strains on the stability of the CsrA-sponging sRNA
CsrB, with *proQ* and *cspC/E* deletion having opposite effects on half-life (**Figure S10B & C**),
adding a further potential connection between RBP regulatory networks. Our reanalysis of
publicly available CLIP-seq data suggests that a substantial number of mRNAs are targeted by
two or more RBPs (**Figure S10D**). What this apparently dense interconnection between RBP-
mediated regulatory networks means for the cell, and how RBP activity is coordinated to
maintain homeostasis in diverse environmental conditions, is an open question that will likely
take significant conceptual advances to answer.
524

525 **Data Availability**
526 Data deposition: All sequencing data reported in this paper have been deposited in the Gene
527 Expression Omnibus (GEO) database, https://www.ncbi.nlm.nih.gov/geo (SuperSeries no.
528 GSE234010). Transcript annotations and source code for the Stan models have been made
529 available at https://github.com/BarquistLab/RIF-seq_repo
530

Acknowledgements
We thank Joel Belasco, Erik Holmqvist, Susan Gottesman and Anke Sparmann for insightful
comments on the manuscript, and Alexandre Smirnov for providing northern blot quantifications
from (13). This project was funded in part by the Bavarian State Ministry for Science and the
Arts through the research network bayresq.net (to LB, JV).

Methods
538

Media and growth conditions

For all experiments in this study, broth cultures were grown from single colonies overnight at 37 °C in LB medium (5 g/L of yeast extract, 5 g/L of NaCl, and 10 g/L of Tryptone/Peptone ex casein; Roth). Subsequently, cultures were diluted 1:100 in fresh medium, and further grown at 37°C with shaking at 220 rpm to an OD600 of 2.0 (early stationary phase (ESP), a SPI-1 inducing condition (30)).

Bacterial strains and plasmids

Salmonella enterica serovar Typhimurium strain SL1344 (strain JVS-1574 (71)) is considered wild-type (WT). The generation of *proQ* and *cspC/E* deletion strains by lambda red homologous recombination (72) has been previously described (11, 13). For the *proQ++* strain, a strain containing plasmid pZE12-ProQ was used as previously described (12, 13). The complete lists of bacterial strains, plasmids, oligos and antibodies used in this study are provided in SI Appendix, Table S1-4.

Rifampicin assay protocol for sequencing

Wild-type (WT), ΔRBP and RBP++ strains were grown until an OD_{600} of 2.0 in three (WT, *cspCE*) or six (WT, *proQ*, *proQ++*) replicates. The cultures were treated with 500µl/ml of rifampicin (stock solution 50mg/ml resuspended in DMSO). Samples were taken before ($t = 0$min) and after 3, 6, 12, 24 min (proQ) or 2, 4, 8 and 16 min (cspCE) of rifampicin treatment. 2ml were collected for each sample, immediately mixed with 20% vol. stop mix (95% ethanol, 5% phenol) and snap frozen in liquid nitrogen.

Subsequently, the samples were thawed on ice and centrifuged for 20 min at 4500 rpm. Bacterial pellets were resuspended in 600 µl of 0.5 mg/ml of lysozyme in TE buffer pH 8 and transferred into a 2 ml Eppendorf tube. 2.5 µl of 1/10 ERCC spike-ins was added to each sample. Total RNA was then extracted using the hot phenol method. Briefly, 60 µl of 10% w/v SDS was added to the suspension, and the samples were mixed by inversion. Tubes were placed at 64□ for 1-2 min until clearance of the solution, then 66 µl of 3M sodium acetate solution at pH 5.2 was added and tubes were mixed by inversion. 750 µl of phenol (Roti-Aqua phenol #A980.3) was then added to each tube, mixed by inversion and incubated for 6 min at 64□. Tubes were then placed on ice to cool and spun for 15 min at 13 000 rpm, 4□. The resulting aqueous layer was transferred in a 2 ml PLG tube (5PRIME) where 750 µl of chloroform (Roth, #Y015.2) was added. After mixing by inversion, the tubes were spun for 15 min at 13 000 rpm, 4□. The obtained aqueous layer was then collected and precipitated in a 30:1 mix of 100% ethanol: 3M sodium acetate pH 6.5 at -20□ for at least 2 hr. After centrifugation for 30 min, 13 000 rpm, 4□, the pellet was washed with 70% ethanol and the air-dried pellet was resuspended in nuclease-free water. Total RNA was measured by nanodrop, and integrity was checked on TBE agarose gel. 40 µg of RNA in 39.5 µl of nuclease free water were then subjected to DNAse I treatment. Total RNA was denatured for 5 min at 65□ and put back on ice. 5 µl of DNase I (Fermentas), 5 µl of DNase I buffer (Fermentas) and 0.5 µl of Superase In (Thermo Fisher Scientific) were added to the denatured RNA and incubated at 37°C for 30 min. After incubation, 100 µl of nuclease free water was added and each reaction was placed in a PLG tube containing 150 µl of PCI. Tubes were centrifuged for 15 min at 4□, 13 000 rpm. The aqueous phases were collected and precipitated in 30:1 Ethanol/sodium acetate

mix at -20☐ for at least 2 hr. DNase treated pellets were collected by centrifugation (30 min, 4☐,
13 000 rpm) and after 70% ethanol wash, were resuspended in 25 µl nuclease free water.
RNA-seq libraries were prepared by Vertis AG (Freising-Weihenstephan, Germany).
Briefly, the ribodepleted RNA samples were fragmented using ultrasound (4 pulses of 30 s each
at 4°C). Subsequently, an oligonucleotide adapter was ligated to the 3' end of the RNA
molecules. First-strand cDNA synthesis was performed using M-MLV reverse transcriptase and
the 3' adapter as primer. The first-strand cDNA was purified and the 5' Illumina TruSeq
sequencing adapter was ligated to the 3' end of the antisense cDNA. The resulting cDNA was
PCR-amplified to about 10-20 ng/µl using a high fidelity DNA polymerase. cDNA was purified
with the Agencourt AMPure XP kit (Beckman Coulter Genomics) and sequenced on an Illumina
HiSeq2000. Replicate 2 of the 24 minute time point for WT, Δ*proQ*, and *proQ++* was excluded
from subsequent analysis, as rRNA depletion failed.
595
Processing of sequencing reads and mapping of RIF-seq data
The 75 nt RNA-seq reads were demultiplexed and quality control of each sample was
performed with fastQC. Afterwards, Illumina adapters were removed with Cutadapt v4.1, and
STAR (73) was used to align the reads to the SL1344 genome (NCBI accessions: FQ312003.1,
HE654724.1, HE654725.1 and HE654726.1). For all analyses related to annotated genomic
features such as CDSs, tRNAs, and rRNAs, gene annotations from NCBI were used. We use
the same definition of transcriptional units as (54) which is based on the NCBI CDS annotations,
transcription start site annotations (74), and Rho-independent terminator prediction with RNIE
(75). sRNA annotations are based on (13). The ERCC92.fa sequence file for the quantification
of the spike-in was obtained from ThermoScientific. For quantification, htseq-count with default
options was used for counting reads aligning to CDS, sRNA and ERCC spike-ins, while the 60
base sub-genic windows were counted with the option *--nonunique all* to ensure that
overlapping reads are assigned to all overlapping segments. For the 60 base windows, reads
were quantified separately for the positive and the negative strand.
610
Read count normalization of RIF-seq data with ERCC spike-ins
The samples were normalized using a custom implementation of the trimmed mean of M-values
(76) across 30 detected ERCC spike-ins (**Fig. S2B&C**). The cutoff on the M values was set to
0.3 and the cutoff on the A values to 0.05. Only transcripts with more than 10 counts-per-million
(cpm) before normalization in at least three samples in the ProQ assay were retained for further
analysis.
Normalized counts-per-million (cpm) were obtained by adding a pseudo count and then
dividing the read counts $Y_{gcr}(t)$ by the respective library size N_s and normalization factor $n_{f,s}$ of
the sample

$$cpm_{gcr,s} = \frac{\left(Y_{gcr,s} + 0.5\right) \cdot 10^6}{N_s n_{f,s} + 1}$$

620   The Stan models were applied to the natural logarithm of the normalized cpm values $y_{gcr}(t) \equiv$
621    $ln\left(cpm_{gcr}(t)\right).$
622

Removal of batch effects: center-mean normalization

Following spike-in normalization, we observed some clustering by replicate rather than condition within time point groups (**Figure S2D&I**). To account for these batch effects, we developed a center-mean (CM) normalization procedure, which can be applied after a primary normalization, e.g. with spike-ins, and compensates for small variations in the amount of spike-ins added to the individual samples. After the normalization with spike-ins, we calculated a gene-wise mean log-count $y_{gc}(t)$ for every condition and every time-point (see **Figure S2E** for $t = 0$ min). This value was subtracted from the observed value in every sample

$$y_{0,gcr}(t) = y_{gcr}(t) - y_{gc}(t).$$

For every sample s (uniquely defined by condition c, time t, replicate r), we calculated the mean

$$ln(n_{b,s}) = \frac{1}{N_g}\sum_g y_{0,gcr}(t),$$

where $n_{b,s}$ is an additional normalization constant. The batch-corrected cpm values are then given by

$$cpm_{gcr,s} = \frac{(Y_{gcr,s} + 0.5) \cdot 10^6}{N_s n_{f,s} n_{b,s} + 1}.$$

PCA confirmed that the samples separated well by time point and genotype after the CM normalization (**Figure S2F&L**), and boxplots showed an improved alignment of median logcpm values (**Figure S2G,H,J,K**). Before fitting the decay curves with the Bayesian models, we subtracted the mean log-count at $t = 0$ min

$$_{gcr}(t) = y_{gcr}(t) - y_{gc}(t = 0min).$$

Calculation of detection limit

In order to regularize zero counts, we have added a pseudo count of 0.5. The library sizes vary in size around 10 million reads and the normalization constants around 1. This results in an estimated minimum log-count of

$$y_{min} = ln\left(\frac{0.5 \cdot 10^6}{N_s n_{f,s} n_{b,s} + 1}\right) \simeq log\left(\frac{0.5}{10}\right).$$

After subtracting the mean log-count at $t = 0$ min, we can calculate the detection limit for gene g in condition c as

$$_{lim\ ,gc} = y_{min} - y_{gc}(t = 0min).$$

This corresponds to the minimal possible value of the log relative expression.

Differential gene expression analysis

Log-fold changes were calculated using glmQLFit from edgeR (26) with a cutoff of 0.25 on the log-fold changes. Since batch effects were present after TMM normalization (**Fig. S1 A,B&E**), samples were additionally normalized using RUVg (77) (**Fig. S1C&G**). We selected the 800 least varying genes between the ΔRBP and the WT strain. Since the differences between the *proQ++* and the WT strain were larger, we only took the 600 least varying genes between these two conditions. The intersection between these sets is 37 genes which we used as negative control (**Fig. S1F**). The number of factors of unwanted variation k was set to 6. After RUVg normalization, the samples clustered by strain (**Fig. S1C&H**). We selected differentially expressed genes at an FDR of 0.1 (**Fig. S1I-K, Table S2**).

Extraction of RNA half-lives from RIF-seq data

We compared three statistical models, summarized in **Figure 1B**. All models assume that the normalized log counts follow a normal distribution around a condition and gene-dependent mean

$$_{gcr}(t) \sim N\left(\mu_{gc}(t), \sigma_g(t)\right).$$

Normal distributions have been shown to effectively model RNA-seq data (78), and were computationally more efficient in our initial testing than count-based models (see the SI Appendix for a comparison between normal distributions and count-based models). The mean μ_{gc} is parameterized differently in the three statistical models:

Linear model (LM): $\mu_{gc}(t) = -\beta_{gc} \cdot t$

Piecewise linear model (PLM): $\mu_{gc}(t) = -\Theta\left(t - \gamma_g\right) \cdot \beta_{gc} \cdot \left(t - \gamma_g\right)$

Log-normal model (LNM): $\mu_{gc}(t) = ln\left[\pi_{gc} + \left(1 - \pi_{gc}\right)exp\left(-\Theta\left(t - \gamma_g\right) \cdot \beta_{gc} \cdot \left(t - \gamma_g\right)\right)\right]$

where Θ is the Heaviside step function which is 0 for negative arguments and 1 otherwise. The baseline parameter π_{gc} introduced in the LNM corresponds to the fraction of stable RNA for gene g in condition c as compared to steady-state levels at $t = 0$ min. Since we have no prior knowledge of the distribution of the WT decay rates, we impose a hierarchical prior with mean μ_β and width σ_β to "learn" the distribution of the decay rates and to share information across genetic loci. Additionally, we model the gene-wise standard deviation σ_g hierarchically, which shrinks the gene-wise variance towards the mean variance and reduces the effect of outliers. This is similar to the idea of variance shrinkage through empirical Bayes methods implemented in edgeR (26) and limma (79). Since our library sizes do not vary much between replicates within time points (Figure S2M), there were no large differences in measurement precision to account for (78). For other parameters (e.g. difference in decay rate), broad priors were chosen to minimize their influence on posterior estimates. Priors were defined as follows:

WT decay rate $\beta_{g,WT} \sim N\left(\mu_\beta, \sigma_\beta\right)$

Mutant decay rate $\Delta\beta_{gc} = \beta_{gc} - \beta_{g,WT} \sim N(0,0.2)$

Standard deviation $\sigma_g(t) \sim N(\mu_\sigma, \sigma_\sigma)$ and $\sigma_g(t) \geq 0$

Baseline parameter $\pi_{gc} \sim N(0,0.25)$ and $\pi_{gc} \in [0,0.2]$

Hyperparameters $\mu_\beta, \mu_\sigma, \sigma_\beta \sim C(0,1)$, $\sigma_\sigma \sim N(0.3,0.3)$ and $\mu_\beta, \mu_\sigma, \sigma_\beta, \sigma_\sigma \geq 0$

Elongation time $\gamma_g \sim C_{[0,12]}(0,2)$

For $\pi_{gc} = 0$, the LNM is equivalent to the PLM, which converts to the LM as $\gamma_{gc} \to 0$. The statistical models are fitted to the RIF-seq data using the probabilistic programming language Stan (v.2.30.1) (29) with two chains and 1000 MCMC samples each (`method=sample num_samples=1000 num_warmup=1000 adapt delta=0.95 algorithm=hmc engine=nuts max_depth=15`). The statistical model was applied to all four strains (WT, Δ*proQ*, *proQ*++, Δ*cspCE*) at once. The reported parameters (decay rate, half-life, transcription elongation time) correspond to the median of the 2000 MCMC samples. The median of the transcriptome-wide half-lives corresponds to the median of the 2000 MCMC samples of the hyperparameter μ_β. In addition to the 2nd replicate of the time point taken at 24 minutes for the *proQ* experiments, the 1st replicate of the 4 min time point of the Δ*cspCE* mutant was removed

704   from this part of the analysis because it clustered together with the 0 min time point (**Figure S1I**)
705   which strongly influenced differences in decay rate in the Δ*cspCE* mutant.
706
Model comparison using leave-one-out cross validation
For a quantitative comparison of the linear model (LM), the piecewise linear model (PLM) and
the log-normal model (LNM), we estimated the out-of-sample predictive accuracy using leave-
one-out cross validation (LOO-CV) with *Pareto-smoothed importance sampling* (PSIS) (37). The
pointwise log-likelihood `log_lik` was computed in the `generated quantities` block in Stan
during MCMC sampling. We used the `loo()` function from the *loo* R package (version 2.5.1),
which computes the expected log-pointwise predictive density (ELPD) using PSIS.
714
Calculation of transcription velocities
To calculate transcription velocities, we took advantage of ongoing transcription of RNA
polymerase already bound to DNA in the RIF-seq data. We split the genome into 60 base
subgenic windows, and extracted the corresponding elongation times and decay rates using the
log-normal model. We split the dataset into five subsets before running the MCMC sampler (1
chain: `method=sample num_samples=1000 num_warmup=1000 adapt delta=0.95`
`algorithm=hmc engine=nuts max_depth=15`). Subsequently, we verified that the
hierarchical parameters agreed well between the five subsets. The resulting transcription
elongation times γ were combined with operon annotations taken from (54). We fitted a linear
model with y-intercept a_g and slope b_g to the elongation times of operons or individual
transcripts as shown for the mra/fts operons in **Figure S3C,** using the inverse of the 68%
credible intervals of γ_g as obtained from the MCMC samples as weights (example: Fig. 2E). The
transcription velocity v_g is given by the ratio of the window size ($s_{seg} = 60$ nt) and the slope b_g.
Its error was calculated via error propagation $\Delta v_g = \Delta b_g / (s_{seg} \cdot b_g^2)$. We obtain 772 operons
with at least 7 nonzero segments which fulfill the quality criterium $\Delta v_g / v_g < 0.75$.
730
Calculation of Bayesian p-values
The half-lives were calculated from the decay rates $t_{1/2,gc} = ln(2)/\beta_{gc}$. In order to calculate
Bayesian p-values, we tested against the null hypothesis that the difference in half-life $\Delta t_{1/2,c} =$
$t_{1/2,c} - t_{1/2,WT}$ is compatible with zero. There is a limit as to how precisely we measured the WT
half-lives. We determined the minimum of the 90% credible intervals of the WT half-lives
(~0.05). Assuming that we cannot measure a difference in half-life with higher precision than the
WT half-lives, we selected the interval [-0.05, 0.05] as the null hypothesis. The p-value p_{gc} for
gene g in condition c corresponding to the difference in decay rate $\Delta t_{1/2,gc}$ is given by the
fraction $|S_0|/|S|$ of MCMC samples $S = \{s_1, \ldots, s_{2000}\}$ that agrees with the null hypothesis
(**Figure S4A**):
741   For $\Delta t_{1/2,gc} > 0$, the samples $S_0 = \{s \in S | s \leq 0.05\}$ agree with the null hypothesis.
742   For $\Delta t_{1/2,gc} < 0$, the samples $S_0 = \{s \in S | s \geq -0.05\}$ agree with the null hypothesis.
743   We compared the distribution of p-values to the distribution of p-values under the null
744   hypothesis which was obtained by bootstrapping from the distribution of MCMC samples of the
745   WT half-lives and calculating the corresponding p-values (**Figure S4B**).
746

Calibration of posterior predictive p-values

In order to assign a false-discovery rate (FDR) to the p-values, we simulated a dataset with 4000 transcripts, 3 conditions (*WT*, c_1, c_2) with time points 0, 3, 6, 12, 24 and 2 conditions (*WT*, c_3) with time points 0, 2, 4, 8, 16. We drew samples from the following distributions (which we extracted from fitting the LNM to the two RIF-seq data sets), using the definition of the LNM as given above:

$\quad\quad$ Relative log-counts $y_{gcr}(t) \sim N\left(\mu_{gc}(t), \sigma_g(t)\right)$

$\quad\quad$ *WT* decay rate $\beta_{g,WT} \sim N(0.75, 0.3)$

$\quad\quad$ Elongation time $\gamma_g \sim C_{[0,12]}(0,2)$

$\quad\quad$ Standard deviation of log-counts $\sigma_g \sim N(0.35, 0.23)$ and $\sigma_g(t) \geq 0$

$\quad\quad$ Difference in decay rate $\Delta\beta_{gc} = \beta_{gc} - \beta_{g,WT} \sim N(0, 0.08)$

$\quad\quad$ Baseline parameter $\pi_{gc} \sim N(0, 0.05)$

$\quad\quad$ Mean of relative log-counts $\mu_{gc}(t) = ln\left[\pi_{gc} + \left(1 - \pi_{gc}\right)exp\left(-\Theta(t - \gamma_g) \cdot \beta_{gc} \cdot \left(t - \gamma_g\right)\right)\right]$

Then, we fitted the log-normal model to this dataset. Simulated absolute differences in half-life below 0.05 ($\left|\Delta t_{1/2,gc}\right| \leq 0.05$) were assumed to agree with the null hypothesis. The Pearson correlation of 0.86 between simulated and fitted differences in half-life were obtained using the `weightedCorr` function from the *wCorr* package in R with the inverse of the size of the 90% credible intervals of the fitted half-lives as weights (**Figure S4C**). We calculated the posterior predictive p-values for the fitted differences in half-life and varied the p-value cutoff between 0 and 1 with step size 0.01. The corresponding FDR is given by the fraction of transcripts whose simulated difference in half-life agrees with the null hypothesis and the total number of transcripts with a p-value below the cutoff. Subsequently, we fitted a LOESS curve in R (`span=0.2`) to determine the FDR corresponding to any p-value cutoff (**Figure S4E, Table S1**) A p-value of 0.082 corresponds to a FDR of 0.1 which we used as a cutoff for our analysis of differentially decaying transcripts. In addition to controlling the FDR, we verified that at an FDR of 0.1, the log-normal model identifies differentially decaying transcripts with a low simulated standard deviation on log-counts σ_g with high sensitivity (**Figure S4D**). For this, we selected five cutoffs on standard deviation (0.05, 0.1, 0.2, 0.4, 1) and calculated the false positive rate (FPR) and sensitivity for all transcripts below the cutoff.

Supplementary Material

Supplementary Methods, bacterial strains, plasmids, oligos and antibodies used in this study (**Tables S1-S4**) and Supplementary **Figures S1-14** can be found in the SI Appendix. Genome-wide *Salmonella* half-life estimates (**Dataset S1**) and log-fold changes in the absence of the global RBPs ProQ and CspC/E (**Dataset S2**), candidate lists for ProQ and CspC/E protected transcripts (**Dataset S3&4**) and significant CLIP-seq peaks for ProQ, CspC & CspE (**Dataset S5-7**) are provided as XLSX files.

References

1. J. G. Belasco, All things must pass: contrasts and commonalities in eukaryotic and bacterial mRNA decay. *Nat. Rev. Mol. Cell*

Biol. **11**, 467–478 (2010).

2.   M. P. Hui, P. L. Foley, J. G. Belasco, Messenger RNA degradation in bacterial cells. *Annu. Rev. Genet.* **48**, 537–559 (2014).

3.   T. B. Updegrove, A. Zhang, G. Storz, Hfq: the flexible RNA matchmaker. *Curr. Opin. Microbiol.* **30**, 133–138 (2016).

4.   K. Kavita, F. de Mets, S. Gottesman, New aspects of RNA-based regulation by Hfq and its partner sRNAs. *Curr. Opin.*
Microbiol. **42**, 53–61 (2017).

5.   T. Romeo, P. Babitzke, Global Regulation by CsrA and Its RNA Antagonists. *Microbiol Spectr* **6** (2018).

6.   E. Holmqvist, J. Vogel, RNA-binding proteins in bacteria. *Nat. Rev. Microbiol.* (2018) https:/doi.org/10.1038/s41579-018-0049-5.

7.   T. S. Stenum, E. Holmqvist, CsrA enters Hfq's territory: regulation of a base-pairing small RNA. *Mol. Microbiol.* (2021)
https:/doi.org/10.1111/mmi.14785.

8.   T. S. Stenum, *et al.*, RNA interactome capture in Escherichia coli globally identifies RNA-binding proteins. *Nucleic Acids Res.*
(2023) https:/doi.org/10.1093/nar/gkad216.

9.   L.-C. Chu, *et al.*, The RNA-bound proteome of MRSA reveals post-transcriptional roles for helix-turn-helix DNA-binding and
Rossmann-fold proteins. *Nat. Commun.* **13**, 2883 (2022).

10.   E. C. Urdaneta, *et al.*, Purification of cross-linked RNA-protein complexes by phenol-toluol extraction. *Nat. Commun.* **10**, 990
(2019).

11.   C. Michaux, *et al.*, RNA target profiles direct the discovery of virulence functions for the cold-shock proteins CspC and CspE.
Proc. Natl. Acad. Sci. U. S. A., 201620772 (2017).

12.   A. J. Westermann, *et al.*, The Major RNA-Binding Protein ProQ Impacts Virulence Gene Expression in Salmonella enterica
Serovar Typhimurium. *MBio* **10**, e02504–18 (2019).

13.   A. Smirnov, *et al.*, Grad-seq guides the discovery of ProQ as a major small RNA-binding protein. *Proc. Natl. Acad. Sci. U. S. A.*
113, 11591–11596 (2016).

14.   E. Holmqvist, L. Li, T. Bischler, L. Barquist, J. Vogel, Global Maps of ProQ Binding In Vivo Reveal Target Recognition via RNA
Structure and Stability Control at mRNA 3' Ends. *Mol. Cell* **70**, 971–982.e6 (2018).

15.   S. Melamed, P. P. Adams, A. Zhang, H. Zhang, G. Storz, RNA-RNA Interactomes of ProQ and Hfq Reveal Overlapping and
Competing Roles. *Mol. Cell* **77**, 411–425.e7 (2020).

16.   A. Rizvanovic, *et al.*, The RNA-Binding Protein ProQ Promotes Antibiotic Persistence in Salmonella. *MBio*, e0289122 (2022).

17.   A. Smirnov, C. Wang, L. L. Drewry, J. Vogel, Molecular mechanism of mRNA repression intransby a ProQ-dependent small
RNA. *EMBO J.* **36**, 1029–1045 (2017).

18.   E. Holmqvist, S. Berggren, A. Rizvanovic, RNA-binding activity and regulatory functions of the emerging sRNA-binding protein
ProQ. *Biochim. Biophys. Acta Gene Regul. Mech.*, 194596 (2020).

19.   S. Ray, R. Da Costa, S. Thakur, D. Nandi, Salmonella Typhimurium encoded cold shock protein E is essential for motility and
biofilm formation. *Microbiology* **166**, 460–473 (2020).

20.   S. Bauriedl, *et al.*, The minimal meningococcal ProQ protein has an intrinsic capacity for structure-based global RNA
recognition. *Nat. Commun.* **11**, 2823 (2020).

21.   E. A. Campbell, *et al.*, Structural mechanism for rifampicin inhibition of bacterial rna polymerase. *Cell* **104**, 901–912 (2001).

22.   J. A. Bernstein, A. B. Khodursky, P.-H. Lin, S. Lin-Chao, S. N. Cohen, Global analysis of mRNA decay and abundance in
Escherichia coli at single-gene resolution using two-color fluorescent DNA microarrays. *Proc. Natl. Acad. Sci. U. S. A.* **99**,
9697–9702 (2002).

23.   D. W. Selinger, R. M. Saxena, K. J. Cheung, G. M. Church, C. Rosenow, Global RNA half-life analysis in Escherichia coli
reveals positional patterns of transcript degradation. *Genome Res.* **13**, 216–223 (2003).

24.   H. Chen, K. Shiroguchi, H. Ge, X. S. Xie, Genome-wide study of mRNA degradation and transcript elongation in Escherichia
coli. *Mol. Syst. Biol.* **11**, 781 (2015).

25.   M. E. Ritchie, *et al.*, limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Res. **43**, e47 (2015).

26.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential expression analysis of digital
gene expression data. *Bioinformatics* **26**, 139–140 (2010).

27.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome*

Biol. **15**, 550 (2014).

28. I. Lönnstedt, T. Speed, REPLICATED MICROARRAY DATA. *Stat. Sin.* **12**, 31–46 (2002).

29. B. Carpenter, *et al.*, Stan: A probabilistic programming language. *J. Stat. Softw.* **76** (2017).

30. C. A. Lee, S. Falkow, The ability of Salmonella to enter mammalian cells is affected by bacterial growth state. *Proc. Natl. Acad.*
Sci. U. S. A. **87**, 4304–4308 (1990).

31. L. Jiang, *et al.*, Synthetic spike-in standards for RNA-seq experiments. *Genome Res.* **21**, 1543–1551 (2011).

32. R. D. Mosteller, C. Yanofsky, Transcription of the tryptophan operon in Escherichia coli: rifampicin as an inhibitor of initiation. *J.*
Mol. Biol. **48**, 525–531 (1970).

33. D. Dar, R. Sorek, Bacterial Noncoding RNAs Excised from within Protein-Coding Transcripts. *MBio* **9** (2018).

34. G. Matera, *et al.*, Global RNA interactome of Salmonella discovers a 5' UTR sponge for the MicF small RNA that connects
membrane permeability to transport capacity. *Mol. Cell* **82**, 629–644.e4 (2022).

35. S. Z. Abdulla, *et al.*, Small RNAs Activate Salmonella Pathogenicity Island 1 by Modulating mRNA Stability through the hilD
mRNA 3' Untranslated Region. *J. Bacteriol.* **205**, e0033322 (2023).

36. D. Sinha, N. R. De Lay, Target recognition by RNase E RNA-binding domain AR2 drives sRNA decay in the absence of
PNPase. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2208022119 (2022).

37. A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Stat.*
Comput. **27**, 1413–1432 (2017).

38. M. Y. Galperin, *et al.*, COG database update: focus on microbial diversity, model organisms, and widespread pathogens.
Nucleic Acids Res. **49**, D274–D281 (2021).

39. J. H. Urban, J. Vogel, Two seemingly homologous noncoding RNAs act hierarchically to activate glmS mRNA translation. *PLoS*
Biol. **6**, e64 (2008).

40. C. Bækkedal, P. Haugen, The Spot 42 RNA: A regulatory small RNA with roles in the central metabolism. *RNA Biol.* **12**, 1071–
1077 (2015).

41. N. De Lay, S. Gottesman, The Crp-activated small noncoding regulatory RNA CyaR (RyeE) links nutritional status to group
behavior. *J. Bacteriol.* **191**, 461–476 (2009).

42. A. Vehtari, J. Ojanen, A survey of Bayesian predictive methods for model assessment, selection and comparison. *Stat. Surv.* **6**,
142–228 (2012).

43. A. Gelman, Two simple examples for understanding posterior p-values whose distributions are far from uniform. *EJSS* **7**,
2595–2602 (2013).

44. Y. Yair, *et al.*, Cellular RNA Targets of Cold Shock Proteins CspC and CspE and Their Importance for Serum Resistance in
Septicemic Escherichia coli. *mSystems* **7**, e0008622 (2022).

45. M. Raffatellu, *et al.*, SipA, SopA, SopB, SopD, and SopE2 contribute to Salmonella enterica serotype typhimurium invasion of
epithelial cells. *Infect. Immun.* **73**, 146–154 (2005).

46. J. A. Imlay, Transcription Factors That Defend Bacteria Against Reactive Oxygen Species. *Annu. Rev. Microbiol.* **69**, 93–108
(2015).

47. Y. Chao, *et al.*, In Vivo Cleavage Map Illuminates the Central Role of RNase E in Coding and Non-coding RNA Pathways. *Mol.*
Cell **65**, 39–51 (2017).

48. J. Richards, J. G. Belasco, Obstacles to Scanning by RNase E Govern Bacterial mRNA Lifetimes by Hindering Access to Distal
Cleavage Sites. *Mol. Cell* (2019) https:/doi.org/10.1016/j.molcel.2019.01.044.

49. J. Richards, J. G. Belasco, Graded impact of obstacle size on scanning by RNase E. *Nucleic Acids Res.* **51**, 1364–1374 (2023).

50. A. Subramanian, *et al.*, Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 15545–15550 (2005).

51. Jörg Kunte H., Crane Rebecca A., Culham Doreen E., Richmond Deborah, Wood Janet M., Protein ProQ Influences Osmotic
Activation of Compatible Solute Transporter ProP in Escherichia coliK-12. *J. Bacteriol.* **181**, 1537–1543 (1999).

52. C. H. Kerr, D. E. Culham, D. Marom, J. M. Wood, Salinity-dependent impacts of ProQ, Prc, and Spr deficiencies on Escherichia
coli cell structure. *J. Bacteriol.* **196**, 1286–1296 (2014).

53. S. G. Chaulk, *et al.*, ProQ is an RNA chaperone that controls ProP levels in Escherichia coli. *Biochemistry* **50**, 3095–3106

(2011).

54.　E. Holmqvist, *et al.*, Global RNA recognition patterns of post-transcriptional regulators Hfq and CsrA revealed by UV
crosslinking in vivo. *EMBO J.* **35**, 991–1011 (2016).

55.　M. Scott, C. W. Gunderson, E. M. Mateescu, Z. Zhang, T. Hwa, Interdependence of cell growth and gene expression: origins
and consequences. *Science* **330**, 1099–1102 (2010).

56.　T. Esquerré, *et al.*, Dual role of transcription and transcript stability in the regulation of gene expression in Escherichia coli cells
cultured on glucose at different growth rates. *Nucleic Acids Res.* **42**, 2460–2472 (2014).

57.　A. H. Potts, Y. Guo, B. M. M. Ahmer, T. Romeo, Role of CsrA in stress responses and metabolism important for Salmonella
virulence revealed by integrated transcriptomics. *PLoS One* **14**, e0211430 (2019).

58.　Morin Manon, Enjalbert Brice, Ropers Delphine, Girbal Laurence, Cocaign-Bousquet Muriel, Genomewide Stabilization of
mRNA during a "Feast-to-Famine" Growth Transition in Escherichia coli. *mSphere* **5**, e00276–20 (2020).

59.　E. Baracchini, H. Bremer, Determination of synthesis rate and lifetime of bacterial mRNAs. *Anal. Biochem.* **167**, 245–260
(1987).

60.　D. A. Vargas-Blanco, S. S. Shell, Regulation of mRNA Stability During Bacterial Stress Responses. *Front. Microbiol.* **11**, 2111
(2020).

61.　D. A. Vargas-Blanco, Y. Zhou, L. G. Zamalloa, T. Antonelli, S. S. Shell, mRNA Degradation Rates Are Coupled to Metabolic
Status in Mycobacterium smegmatis. *MBio* **10** (2019).

62.　G. Yim, F. de la Cruz, G. B. Spiegelman, J. Davies, Transcription modulation of Salmonella enterica serovar Typhimurium
promoters by sub-MIC levels of rifampin. *J. Bacteriol.* **188**, 7988–7991 (2006).

63.　E. Massé, F. E. Escorcia, S. Gottesman, Coupled degradation of a small regulatory RNA and its mRNA targets in Escherichia
coli. *Genes Dev.* **17**, 2374–2383 (2003).

64.　Y. Zhang, *et al.*, A Stress Response that Monitors and Regulates mRNA Structure Is Central to Cold Shock Adaptation. *Mol.*
Cell (2018) https:/doi.org/10.1016/j.molcel.2018.02.035.

65.　W. Bae, B. Xia, M. Inouye, K. Severinov, Escherichia coli CspA-family RNA chaperones are transcription antiterminators. *Proc.*
Natl. Acad. Sci. U. S. A. **97**, 7784–7789 (2000).

66.　R. Balakrishnan, *et al.*, Principles of gene regulation quantitatively connect DNA to RNA and proteins in bacteria. *Science* **378**,
eabk2066 (2022).

67.　Y. Chao, J. Vogel, The role of Hfq in bacterial pathogens. *Curr. Opin. Microbiol.* **13**, 24–33 (2010).

68.　M. G. Jørgensen, M. K. Thomason, J. Havelund, P. Valentin-Hansen, G. Storz, Dual function of the McaS small RNA in
controlling biofilm formation. *Genes Dev.* **27**, 1132–1145 (2013).

69.　Y.-J. Lai, *et al.*, CsrA regulation via binding to the base-pairing small RNA Spot 42. *Mol. Microbiol.* **117**, 32–53 (2022).

70.　B. Xia, H. Ke, M. Inouye, Acquirement of cold sensitivity by quadruple deletion of the cspA family and its suppression by
PNPase S1 domain in Escherichia coli. *Mol. Microbiol.* **40**, 179–188 (2001).

71.　B. A. Stocker, S. K. Hoiseth, B. P. Smith, Aromatic-dependent "Salmonella sp." as live vaccine in mice and calves. *Dev. Biol.*
Stand. **53**, 47–54 (1983).

72.　K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc.*
Natl. Acad. Sci. U. S. A. **97**, 6640–6645 (2000).

73.　A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

74.　C. Kröger, *et al.*, An infection-relevant transcriptomic compendium for Salmonella enterica Serovar Typhimurium. *Cell Host*
Microbe **14**, 683–695 (2013).

75.　P. P. Gardner, L. Barquist, A. Bateman, E. P. Nawrocki, Z. Weinberg, RNIE: genome-wide prediction of bacterial intrinsic
terminators. *Nucleic Acids Res.* **39**, 5845–5852 (2011).

76.　M. D. Robinson, A. Oshlack, A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol.*
11, R25 (2010).

77.　D. Risso, J. Ngai, T. P. Speed, S. Dudoit, Normalization of RNA-seq data using factor analysis of control genes or samples.
Nat. Biotechnol. **32**, 896–902 (2014).

78.　C. W. Law, Y. Chen, W. Shi, G. K. Smyth, voom: Precision weights unlock linear model analysis tools for RNA-seq read counts.
Genome Biol. **15**, R29 (2014).

79. G. K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat.*
Appl. Genet. Mol. Biol. **3**, Article3 (2004).