Machine learning and phylogenetic models identify predictors of genetic variation in Neotropical

amphibians

RUNNING TITLE: Neotropical amphibian genetic variation

Luis Amador (laamador@unm.edu; https://orcid.org/0000-0003-2638-4068),

Irvin Arroyo-Torres (irtorre5@unm.edu; https://orcid.org/0000-0001-6811-7422),

Lisa N. Barrow (lnbarrow@unm.edu; https://orcid.org/0000-0001-7081-2432) (Corresponding author)

Museum of Southwestern Biology and Department of Biology, University of New Mexico, Albuquerque,

NM, USA, 87131

19
20

Abstract

Aim. Intraspecific genetic variation is key for adaptation and survival in changing environments and is

known to be influenced by many factors, including population size, migration, and life history traits. We

investigated genetic variation within Neotropical amphibian species to provide insights into how natural

history traits, phylogeny, climatic, and geographic characteristics influence intraspecific diversity.

Location. Neotropics.

Taxon. Amphibians.

Methods. We assembled datasets using open-access databases for natural history traits, genetic

sequences, phylogenetic trees, climatic, and geographic data. For each species, we calculated overall

nucleotide diversity (π) and tested for isolation by distance (IBD) and isolation by environment (IBE). We

then identified predictors of π, IBD, and IBE using Random Forest (RF) regression or RF classification.

To incorporate phylogenetic relationships, we fitted phylogenetic generalized linear mixed models

(PGLMMs) to predict π, IBD, and IBE.

Results. We compiled 4,052 mitochondrial DNA sequences from 256 amphibian species (230 frogs and

26 salamanders), georeferencing 2,477 sequences from 176 species that were not linked to occurrence

data. RF regressions and PGLMMs were congruent in identifying range size and precipitation (σ) as the

most important predictors of π. RF classification and PGLMMs identified minimum elevation as an

important predictor of IBD, and maximum latitude and precipitation (σ) as the best predictors of IBE.

Main conclusions. This study unified machine learning and phylogenetic methods and identified

predictors of genetic variation in Neotropical amphibians. This approach was valuable to determine which

predictors were congruent between methods. We found that species with small ranges or living in zones

with less variable precipitation tended to have low genetic diversity. We also showed that Western

Mesoamerica, Andes, and Atlantic Forest biogeographic units harbor high diversity across many species

that should be prioritized for protection. These results could play a key role in the development of

conservation strategies for Neotropical amphibians.

Keywords

Anura, Caudata, elevation, latitude, phylogeny, precipitation, Random Forest, range size.

49 Introduction

Characterizing genetic variation within species and understanding the processes that maintain that

diversity are important goals in evolutionary biology (Ellegren & Galtier, 2016). Genetic variation is

essential for natural populations to develop new traits and adapt to future environmental changes (Stange

et al., 2021), making it a fundamental aspect of biodiversity conservation efforts (DeWoody et al., 2021).

Intraspecific genetic variation includes both the variation within populations and the distribution of spatial

genetic variation between populations. These two components provide insights into demographic history

and evolutionary processes such as changes in effective population size (N_e) or the pattern of genetic

exchange between populations (Paz-Vinas et al., 2018). Therefore, investigating the forces that drive

genetic diversity within species and how this diversity is distributed and influenced spatially, can help

explain the processes underlying the patterns of genetic variation that we observe in nature.

Levels of genetic variation vary greatly in natural populations and among species (see Romiguier et al.,

2014), however, determinants of intraspecific genetic variation remain poorly understood. For decades,

many studies have considered N_e to be the most important evolutionary parameter that has a significant

impact on genetic variation within and between populations (e.g., Frankham, 1996). For example, small,

isolated populations tend to have low genetic variation due to increased genetic drift and reduced gene

flow. Under neutral theory, genetic variation is expected to increase with population size (Kimura, 1983),

due mainly to reduced genetic drift in large populations (Buffalo, 2021). Several empirical studies have

shown results consistent with the neutral theory, in which species with higher population abundances

have higher genetic diversity (e.g., Grundler et al. 2019; Hague and Routman, 2016). However, genetic

diversity does not necessarily correlate with population size in all cases (e.g., Bazin et al. 2006), and

genetic variation levels across species have been observed to be much narrower than their variation in

population size (the so-called Lewontin's paradox; Lewontin, 1974). In nature, genetic variation within

species can be influenced by several additional factors, including environmental and intrinsic

characteristics (e.g., Nevo, 1978).

Distinct geographic, climatic, and life history factors can determine intraspecific genetic variation and

influence the diversification and extinction of populations and species. The range size of a species and the

environmental heterogeneity across the species distribution can influence genetic variation. Species with

larger ranges are expected to have higher genetic diversity, low inbreeding, and reduced genetic drift

because of the direct relationship between geographic range size and population size (see Leffler et al.,

2012). For example, studying nine co-distributed amphibian species, García-Rodriguez et al. (2020) found

that abiotic factors and geographical features affected the genetic diversity of species in Isthmian Central

America. In addition, habitat fragmentation can lead to reduced gene flow and increased genetic

differentiation among populations, ultimately reducing genetic diversity within populations because of

genetic drift (see Dixo et al., 2009). Life history traits also have an impact on genetic variation, providing

a connection between different demographic processes (Duminil et al., 2007). For example, species with

shorter lifespans and higher fecundity may have higher genetic diversity because there are more chances

for mutation and recombination, and to cover the full gradient of environmental pressures experienced by

the species (Romiguier et al. 2014). Body size can be another important predictor of genetic variation

within some taxonomic groups. For example, Brüniche☐Olsen et al. (2019) found that genetic diversity

decreases with increasing body size in Darwin's finches, with the possible explanation that species

abundance is expected to decrease with increasing body size (White et al. 2007). Larger-bodied species

may also have higher dispersal ability resulting in differences in spatial genetic variation, as has been

demonstrated in bees (López-Uribe et al. 2019) and frogs (Paz et al. 2015).

An improved understanding of these factors can provide insights into how genetic variation is shaped in

natural populations. Given the variety of factors and their complex relationships, the approach used to

assess potential predictors of genetic diversity is no less important. One approach is to collect new genetic

data for a set of taxa of interest and analyze linear models of genetic variation with possible predictors

(e.g., Dixo et al., 2009). An alternative approach, which can enable comparisons of many more species, is

to use repurposed data that was previously collected for a different research purpose and that can be

reanalyzed in a common framework. These two approaches converge in macrogenetics (sensu Blanchet et

al., 2017), a field that focuses on the integration of genetic datasets from multiple species at large scales

with environmental datasets to identify drivers of intraspecific genetic variation (Leigh et al., 2021).

Recently, Pelletier and Carstens (2018) applied a machine learning framework to repurposed

georeferenced DNA sequences from more than 8000 species and found that geographic range size and

latitude were the most important predictors of genetic structure. Another approach to address how

landscapes contribute to the evolution of genetic variation is one that quantifies the effects of geography

and ecology using multiple matrix regression with randomization (MMRR; Wang, 2013). Using this

approach, Wieringa et al. (2020) examined the impact of geography and ecology on genetic distance of

several species in the southeastern United States, finding that geographic or environmental distance were

significant factors in most of the species evaluated.

Amphibians have been a preferred study system for several ecological and evolutionary studies because

they exhibit a wide variety of natural history traits (e.g., complex life cycles) and distributional patterns

(e.g., species with limited dispersal capabilities that lead to high levels of genetic differentiation). Many

studies have explored patterns of diversity in amphibians. For example, on a regional scale, a positive

latitudinal gradient of taxonomic, functional, and phylogenetic diversity in the Americas has been

demonstrated in amphibians (Ochoa-Ochoa et al., 2020). Global genetic diversity patterns also appear to

follow a latitudinal gradient in amphibians (Miraldo et al. 2016), but less is known about the predictors of

intraspecific variation. In a study of Nearctic amphibian species, the most important predictors of genetic

diversity were taxonomic family, number of sequences, and for salamander species (N = 98), those at

more northern latitudes had lower genetic diversity (Barrow et al., 2021). In the same region, Schmidt et

al. (2022) analyzed microsatellite data from 19 amphibian species; they found that genetic diversity was

not predicted by the environmental variables they used and that areas with high species richness also had

high genetic structure, but low genetic diversity.

The Neotropical Region includes almost 50% of the world's total amphibian species, which is greater

than any other comparable area on the planet (Menéndez-Guerrero et al., 2020). This exceptionally high

diversity is possibly due to a combination of factors, including the geological history of the region (e.g.,

the formation of the Isthmus of Panama, Andean uplift), climate change, ecological interactions, and

biotic diversification (see Elmer et al., 2013; Antonelli, 2022). For decades, the origins and maintenance

of diversity in this region has been an important focus of research in amphibian ecology and systematics.

For example, Gonzalez-Voyer et al. (2011) found that high-altitude ranges, vascularized ventral skin, and

vegetation type correlated significantly with species richness of New World direct-developing frogs.

Using glassfrogs as a study model, Castroviejo-Fisher et al. (2014) showed how the climatic and

geological history of the Neotropics shaped the diversity of this group, demonstrating that initial uplifts of

the Andes and marine incursions are coincidental with divergence times of most glassfrog clades.

Recently, Tobar-Suárez et al. (2022) found that amphibian species richness in Neotropical cloud forests

increased towards the equator, with frog and caecilian species richness increasing towards lower latitudes,

while salamanders showed the opposite pattern. Despite this active research focus on species-level

diversity, very little is known about the determinants of intraspecific genetic variation in Neotropical

amphibians.

Here, we investigated what factors predict genetic variation within Neotropical frog and salamander

species. We gathered genetic sequences, natural history traits, phylogenetic relationships, and climatic

and geographic information for each species from open-access databases and literature. We estimated

nucleotide diversity (π) and tested for isolation-by-distance (IBD) and isolation-by-environment (IBE)

from repurposed DNA sequences. To identify potential hotspots of intraspecific diversity, we built maps

of nucleotide diversity across the Neotropics. We then applied machine learning and phylogenetic models

to investigate the predictors of π, IBD, and IBE within Neotropical amphibians.

Materials and Methods

DNA sequences and associated geographic coordinates

We obtained sequences of the mitochondrial gene cytochrome-b (Cytb) from open access databases. We

chose Cytb since this was the most abundant gene in Neotropical amphibian studies and it has informative

variation within species (van den Burg et al., 2020; Zeisset and Beebee, 2008). We downloaded

sequences from three sources: phylogatR (Pelletier et al., 2022), ACDC (van den Burg et al., 2020) and

GenBank (National Center for Biotechnology Information). Although these sources include Cytb

sequences for hundreds of species, we chose those species with at least five sequences for further analyses

to more adequately represent genetic variation within species (Barrow et al., 2021). Alignments were

saved in FASTA format and were edited and aligned using AliView v.1.28 (Larsson, 2014) with the

MUSCLE aligner v.3.8.31 (Edgar, 2004) using default settings. Geographic coordinates from each

sequence were obtained from phylogatR and GenBank when available, which corresponded to 1,575

sequences (38.87% of the total sequences) from 80 species (31.25% of the total species). We linked 2,477

additional sequences from 176 species to geographic coordinates using either (1) the GeoNames

(geonames.org) and GEOLocate (geo-locate.org) databases by entering the associated location name of

the sequence to obtain an approximate occurrence for that sequence; or (2) information from manuscripts

(e.g., the original species description, distributional records, or systematic and phylogeographic works;

see Appendix S1).

Metrics of genetic variation

To evaluate genetic diversity within species, we calculated nucleotide diversity (π) from each mtDNA

alignment of species and localities using the function nuc.div() in the *pegas* R package (Paradis, 2010).

To evaluate spatial genetic variation within amphibian species, we calculated isolation-by-distance (IBD)

and isolation-by-environment (IBE) for each species. The 'raw' genetic distance (*gendist*; the proportion

of sites that differ between each pair of sequences) was calculated using the function dist.dna() in the *ape*

R package (Paradis and Schliep, 2019). Topographic distance (*geodist*) was calculated between

coordinates associated with each sequence using the function topoDist() implemented in the *topoDistance*

R package (Wang, 2020). Environmental distance (*envdist*) was calculated based on 19 bioclimatic

variables from the WorldClim database (Hijmans et al., 2005) and tree cover data derived from the ESA

WorldCover database (Zanaga et al., 2021). These layers were retrieved at 30-seconds spatial resolution

using the function landcover() of the *geodata* R package (Hijmans et al., 2023). We conducted Multiple

Matrix Regression with Randomization analyses with the MMRR function in R (Wang, 2013) using the

three distances, *gendist, geodist*, and *envdist*, to determine whether each species showed significant IBD

and IBE. Regression coefficients of geography (IBD, β_D) and ecology (IBE, β_E) and their significance

were calculated after 10,000 permutations. MMRR analyses require datasets with n > 4 and assume that

>3 samples show differences in their distances. Based on this premise we used 194 species for this

analysis. After classifying each species as having significant IBD or not and significant IBE or not, we

used these binary classifications as response variables in subsequent analyses.

Trait and geographic data compilation

We obtained information on traits from the AmphiBIO dataset (Oliveira et al., 2017), AmphibiaWeb

(2022), and specific data from manuscripts (Appendix S1; Table 1). Body size (snout-vent length – SVL),

the type of habitat predominantly used by adults, activity (diurnal or nocturnal), and development mode

(larval or direct development) were included since these traits are involved in the dispersal capacity of

amphibian species and can impact intraspecific genetic differentiation (Hillman et al., 2014). We

collected elevational and latitudinal data (mean, maximum, and minimum) from Rolland et al. (2018),

who generated this information from the Global Biodiversity Information Facility (GBIF), the

International Union for Conservation of Nature (IUCN), and WorldClim (Fick and Hijmans, 2017). From

the latter we also obtained temperature (BIO1 = Annual Mean Temperature) and precipitation (BIO12 =

Annual Precipitation). We also obtained latitudinal and elevational data from specific manuscripts.

Elevational data were also recovered from species accounts in the Amphibian Species of the World 6.1

Online Reference (ASW database; Frost, 2021). The geographical range as a shapefile (.shp) for each

species was downloaded from the IUCN portal. We obtained the shapefile for 24 species (e.g., *Allobates*

hodli, *Bolitoglossa awajun*) not included in IUCN by calculating minimum convex polygons based on the

sequence coordinates and using the function mcp() with the *adehabitatHR* R package (Calenge and

Fortmann-Roe, 2023). When necessary, we corrected the polygon for species recently split and not

included in IUCN, e.g., *Rhinella marina – Rhinella horribilis* (Acevedo et al., 2016). The range area was

calculated in square kilometers (km^2) using the functions shapefile() of the *raster* package and

areaPolygon() of the *geosphere* package (Hijmans, 2021; 2022) in R 4.2.2 (R Core Team, 2022). The

IUCN Red List Category was also recorded for each species (IUCN, 2022). We tested for

multicollinearity of the continuous variables using variance inflation factors (VIF) implemented in the R

package *car* (Fox and Weisberg, 2019).

Geographic framework

To evaluate and visualize how amphibian genetic diversity is distributed in the Neotropical Region, we

divided our study area into 17 biogeographic units following the regionalization proposed by Morrone

(2014) and with modifications from Antonelli et al. (2018), Josse et al. (2003), and Olson et al. (2001)

(Figure 1a). We calculated and mapped intraspecific nucleotide diversity first with a single value of π per

species (mean value of π), and second with a value of π per locality within each species. Each locality

consisted of at least two individuals, and we assigned sequences to the same locality when they shared the

same coordinates. In cases when the sequences were from different geographic coordinates, we combined

sequences based on the distance between points (150 km or less) and shared habitat characteristics (e.g.,

occurring in the same mountain range or river). Although this approach involved arbitrarily choosing a

somewhat coarse resolution, it allowed us to maximize the species and geographic area included to

provide an initial picture of spatial genetic diversity given the data available. We visualized π for three

different resolutions (grid cell size of 150 km^2, 250 km^2, and 350 km^2; Figure 1c, Figure S1b, d) which

corresponds to 3640, 1326, and 673 grid cells, respectively. We also visualized the number of sequences

per grid cell for the three resolutions to present the sampling effort and distribution of sequences

available. All mapping analyses were performed in R. To evaluate the degree of match between the

sampling effort (number of observations) and π, we computed the Moran's I statistic to test for spatial

autocorrelation by using the moran.test() function of the *spdep* package (Bivand, 2022).

Identifying predictors of genetic variation with random forests

We used the random forests machine learning algorithm (Breiman, 2001) to build predictive models and

identify variables (e.g., body size, habitat, elevation, range size; see Table 1) that are important predictors

of π, IBD, and IBE in Neotropical amphibians. RF uses independent variables to create many individual

decision trees (a forest) that act as an ensemble to predict a response. Each decision tree in the random

forest uses a subset of the independent variables and returns a response, and variable importance is

determined based on the increase in model error when that variable is not included (Kabacoff, 2015). RF

regression (for π) and classification (for IBD and IBE) models were created using the randomForest()

function in the *randomForest* R package (Liaw and Wiener, 2002), with 5000 trees and 100 permutations.

We split the data into training (90%) and test (10%) datasets and created RF models with the training

data. We used the tuneRF() function to find the optimal mtry value (number of variables to randomly

sample as candidates at each split), and a new model was built using the best mtry value. We made

predictions on the training and test (unseen data values in the models) datasets that were evaluated with

the Mean Squared Error (MSE) and R-squared (R^2) metrics in the RF regression analysis and with the

confusion matrix in RF classification analyses. Relative importance for each independent variable was

measured and printed using the importance() function in the *randomForest* package, and visualized with

the vip() function of the *vip* R package (Greenwell and Boehmke, 2020) for each model. To evaluate the

effect the number of sequences per species and the number of base pairs (bp) in each alignment might

have on the findings, we also created RF models with two different reduced datasets: (1) a dataset with at

least 10 sequences per species (114 species), and (2) a dataset with at least 400 bp in each species

alignment (197 species).

Testing predictors of genetic variation with phylogenetic comparative methods

Random forests can handle large numbers of variables to build predictive models, but do not explicitly

incorporate phylogenetic relationships (other than taxonomic level as a possible predictor). Therefore, we

also fitted phylogenetic generalized linear mixed models (PGLMMs) to determine relationships between

natural history traits, geographic and climatic variables, and genetic variation of Neotropical amphibians.

We pruned the phylogenetic tree from Jetz and Pyron (2018) to create a phylogeny that includes only the

species in our dataset using the phylo4() function in the *phylobase* R package (Hackathon et al., 2020).

This resource is the most complete phylogeny available (7,238 species), covering ≈83% of the known

amphibian diversity. Of the 256 species in our dataset, 15 species were not included in the phylogeny of

Jetz and Pyron (Appendix S2), therefore for phylogenetic analyses we used a subset consisting of 241

species. We mapped π as a continuous trait using the contMap() function in the *phytools* R package

(Revell, 2012), and the distribution of IBD and IBE were mapped at the tips on the tree. We tested for

phylogenetic signal in π using two metrics, Pagel's λ (lambda; Pagel, 1999) and Blomberg's K (Blomberg

et al., 2003), calculated with the phylosig() function implemented in *phytools*. We tested for phylogenetic

signal in IBD and IBE using the phylo.d() function that calculates the D statistic, a measure of

phylogenetic signal in a binary trait (Fritz and Purvis, 2010), implemented in the R package *caper* (Orme

et al., 2018).

Finally, we used PGLMMs (Ives and Helmus, 2011) implemented in the *MCMCglmm* R package

(Hadfield, 2010) to investigate relationships between a subset of the predictors and three responses: (1) π,

(2) IBD, and (3) IBE, while accounting for phylogeny. For these models, we included species 'random

effects', which account for the variability caused by species-specific effects, and phylogenetic random

effects, which consider the phylogenetic relationship between species, by transforming the phylogeny into

a variance-covariance matrix of relatedness between species (Garamszegi, 2014). For each response

variable (π, IBD, and IBE), we compared 10 models that were generated with different combinations of

predictors based on the results of RF analyses (Table S2). These models included: (1) an intercept-only

null model, (2) a model with both species and phylogenetic random effects and no predictors, (3) a model

with six predictors and no random effect, (4) a model with six predictors and species as a random effect,

(5) a model with six predictors and a phylogenetic random effect, (6) a model with six predictors and

species and phylogenetic random effects, (7) a model with 12 predictors and no random effects, (8) a

model with 12 predictors and a species random effect, (9) a model with 12 predictors and a phylogenetic

random effect, and (10) a model with 12 predictors and with both species and phylogenetic random

effects. For Bayesian model selection, we used the deviance information criterion (DIC).

Results

Data compilation

We compiled 4,052 Cytb mtDNA sequences from 256 Neotropical amphibians (Figure S2), of which only

80 species had associated occurrences (we retrieved occurrences for 50 species from phylogatR and 30

species from GenBank). Occurrences for the sequences of the other 176 species were recovered by us in

the present work by retrieving coordinates from published literature or by georeferencing localities. To

visualize sampling effort across the region, we mapped the number of sequences per grid cell (Figure 1b,

Figure S1a, c). There was weak positive spatial autocorrelation of number of sequences (Moran's I =

0.0569, P-value = 0.019), this suggest that similar number of sequences values are somewhat clustered

together but is not a strong pattern in the region. The total occurrences represented all 17 Neotropical

biogeographic units (Figure S3). The final dataset included 230 frogs in 17 families and 26 salamanders

in family Plethodontidae, three response variables (π, IBD, and IBE), and 22 predictor variables (Table 1,

Table S1). We did not detect correlation among predictor variables in the model (VIF values < 2) except

for the three elevation variables (mean, minimum, and maximum) (Figure S4).

Distribution of genetic variation in the Neotropics

The calculated π ranged from 0 (10 species) to 0.156 (Boquete rocket frog, *Silverstoneia nubicola*, family

Dendrobatidae) with a mean of $\pi = 0.025$ (Table S1). When we visualized nucleotide diversity per

locality, we found higher π values in western Mesoamerica, central Andes, Atlantic Forest, and a few

areas of the Amazon region, compared to their adjacent biogeographic units. On the other hand, the

southern Andes and the southern portion of South America showed low genetic diversity (Figure 1c).

Mean values of π per species were higher in southern Mesoamerica, the Chocó region, northern Andes,

and Atlantic Forest region (Figure S5). Spatial autocorrelation analysis suggests that is likely that π values

are randomly distributed in the geographic space (Moran's I = 0.0662, P-value = 0.009). With respect to

spatial genetic variation, we found that 89 Neotropical amphibian species showed significant IBD, while

105 species did not (Table S1). Species following an IBD pattern were mostly found in Mesoamerica (n =

18), the Antillean Subregion (n = 13), the Atlantic Forest (n = 13), and the Andes region (n = 13). The

Amazon included the most species that did not present IBD (n = 28), followed by the Atlantic Forest and

Andes region (Figure S6). We found that most species did not show IBE (n = 124) and only 70 species

showed significant IBE. Species with IBE were found mostly in the Antillean region (n = 14), Andes (n =

13), and Amazonia (n = 12). The Amazon region also included the most species with no IBE patterns (n =

27), followed by the Atlantic Forest and Mesoamerica (Figure S7)

Important predictors of genetic variation based on random forests

Random forest regressions showed that range size was the most important predictor of π for Neotropical

amphibians (Figure 2a). Precipitation standard deviation (σ) was another important predictor of π,

followed by body size and mean temperature (Figure 2a). The overall variance explained by the RF

regression model shown was 20.31%. Random forest classification found that body size was the most

important predictor of IBD, followed by mean precipitation, IUCN rank, and minimum elevation (Figure

2b). The most important predictor for IBE was maximum latitude, followed by precipitation (σ) and

latitude (mean and minimum) (Figure 2c). The model error for the RF classification model for IBD was

37.21% and for IBE was 40.21%. When we evaluated the effect of the sampling effort (number of

sequences) per species (at least 10 sequences), and the sequence length (number of base pairs, bp) in each

alignment (at least 400 bp) on our results, we found that range size was the best predictor of π for all

models analyzed (Figure S8) as in the model using the complete dataset. For IBD, body size remained one

of the most important predictors for the 'at least 10 sequences' dataset, but not the 'at least 400 bp'

dataset. Instead, mean temperature was the most important predictor of IBD in all the models using

reduced datasets (Figure S9). Mean temperature was also the best predictor of IBE in the 'at least 400 bp'

dataset, while microhabitat was the best predictor for the 'at least 10 sequences' dataset (Figure S10). For

IBE, latitude was still among the most important predictors for both reduced datasets.

Relationships between predictors and genetic variation based on PGLMMs

Phylogenetic generalized linear mixed models using *MCMCglmm* showed that the best fit model for π

was the one with a reduced number of predictors and with both species and phylogenetic random effects

(Table 2). This model indicated that range size and precipitation (σ) predicted π, consistent with our

random forest results. Both range size and precipitation (σ) had positive relationships with π (Figure 2d).

Neotropical amphibian species with larger ranges and living in areas with more variable precipitation

tended to have higher π (Figure 3a-c). For IBD, the best-fit *MCMCglmm* model was the full model with

12 predictors and species and phylogenetic random effects (Table 2). *MCMCglmm* also identified

minimum elevation as a significant predictor of IBD, and their relationship was negative (Figure 2e).

Neotropical amphibians living at higher elevations tended to have no IBD, and those with significant IBD

tended to live at lower elevations (Figure 3d). Similar to RF analyses, precipitation (σ) and maximum

Latitude predicted IBE in PGLMM analyses (Figure 2f, Table 2). Species living at southern latitudes in

the Neotropics tended to have no IBE, contrary to amphibians in northern latitudes that tended to exhibit

IBE (Figure 3e). Species following an IBE pattern mostly occur in areas with lower precipitation (σ)

(Figure 3f). *MCMCglmm* results were verified by checking model diagnostics using trace and density

plots of the MCMC samples.

Testing phylogenetic signal

We found significant phylogenetic signal in π (Pagel's λ = 0.7684, P-value (based on LR test) < 0.0001;

Blomberg's K = 0.1509, P-value (based on 1000 randomizations) = 0.002) (Figure 4; Figure S11a, b). We

found clusters of closely related genera that had dissimilar π values; for example, within Bufonidae,

Atelopus species had low π (average = 0.006, range: 0.001-0.010) while toads of genus *Rhinella* presented

a higher average π (average = 0.020, range: 0.006-0.054). The same pattern was also observed in poison

frogs of the family Dendrobatidae, where *Ameerega* species showed high π (average π = 0.024; range:

0.003-0.059) in contrast to species in the genus *Andinobates* which had very low π (average π = 0.003;

range: 0.000-0.009). We did not find significant phylogenetic signal in IBD or IBE, with values of D that

were greater than 1 and were overdispersed compared to a Brownian threshold model (Estimated D_{IBD} =

1.008, P-value = 0.53; Estimated D_{IBE} = 1.047, P-value = 0.685) (Figure S11c, d).

Discussion

In this study we investigated the predictors of genetic variation (π, IBD, and IBE) in Neotropical

amphibians using repurposed data including mtDNA sequences, natural history traits, and geographic

information gathered from open-access databases. Our analyses revealed that geographic range size,

precipitation, elevation, and latitude were significant predictors of different aspects of genetic variation

within species. Specifically, we found that amphibian species inhabiting smaller ranges and places with

lower variation in precipitation had lower intraspecific π (Figure 3a, b); species living at higher elevations

(e.g., mountain ranges) tended not to exhibit IBD; and species living in southern latitudes tended not to

exhibit IBE (Figure 3d, e).

Geographic range size was the top predictor of π, but did not predict IBD and IBE, i.e., Neotropical

amphibian species with and without IBD or IBE can inhabit small or large geographic ranges. We suspect

that environmental heterogeneity within the geographic range rather than range size may have a greater

influence on IBD and IBE patterns of species in the Neotropics, as has been observed in other vertebrate

studies (e.g., Quiroga-Carmona and D'Elía, 2022). Geographic range size has also emerged as an

important predictor of genetic variation in other taxonomic groups such as squamates (Larkin et al., 2023)

and Darwin's finches (Brüniche☐Olsen et al., 2019). Interestingly, the relationship between genetic

diversity and geographic range is not always straightforward, as evidenced by the case of butterflies

(Mackintosh et al., 2019) and other non-model animal species (Romiguier et al., 2014). These

discrepancies highlight the need for continued study of a variety of taxonomic groups and regions to

better understand the ecological and geographic context of genetic variation. Geographically restricted

species are expected to have less genetic variation, which may also indicate an increased risk of extinction

(Levy et al., 2016). In amphibians, Caviedes-Solis et al. (2020) found that Neotropical treefrogs living in

high elevations were more likely to be classified with threatened status. Our analysis of Neotropical

amphibians from several taxonomic families confirmed expected differences between geographic range

sizes based on IUCN conservation status, with Least Concern (LC) amphibian species occurring in larger

ranges and Data Deficient (DD), Endangered (EN) and Critically Endangered (CR) species occurring in

smaller ranges (Figure S12). Based on our estimates, however, IUCN conservation status did not have as

clear of an association with π (Figure 2a, Figure S12), suggesting that IUCN status may not currently

capture this important parameter for population persistence. Regardless, populations of threatened species

with low π such as harlequin frogs of the genus *Atelopus* should be closely monitored.

Precipitation and temperature, along with associated gradients of elevation, has significant impacts on

genetic variation across animal species (De Kort et al., 2021). Our study identified minimum elevation as

a significant predictor of IBD (and it was also one important predictor of π), and precipitation as one of

the best predictors of π, IBD, and IBE (Figure 2). We observed that species inhabiting higher elevations

did not usually exhibit IBD (see Figure 3d), while those living in lower elevations tended to have both

significant IBD patterns and larger geographic range sizes. The absence of IBD patterns, and in turn low

π, in species living at higher elevations could be attributed to their smaller ranges, leading to smaller

population sizes. Our findings partially agree with those reported by Pelletier and Carstens (2018), who

found that geographic range size, elevation, and latitude predicted IBD in several taxonomic groups. In

our case, latitude (maximum) was the most important predictor of IBE. Species with a significant IBE

pattern occurred at higher latitudes than species without IBE. Although latitude was not a significant

predictor of π, analyzing mean latitude with π and geographical range size revealed that amphibians near

the equator have larger ranges, and diversity increases near the equator and slowly decreases towards

higher latitudes (Figure S13). This pattern of genetic diversity forms a plateau around the equator, a

pattern noted by Pereira (2016) in his perspective about a global study mapping amphibian genetic

diversity by Miraldo et al. (2016).

The nucleotide diversity map of Neotropical amphibians clearly shows areas of high π, such as Chocó,

northern Andes, and Atlantic Forests in South America, and an area of high diversity in the Mesoamerica

region located in Central America. Biogeographic units identified with higher π values are areas that also

have high precipitation rates. Therefore, precipitation could explain genetic variation in the Neotropics;

however, this hypothesis needs to be tested in future studies with more species homogeneously distributed

throughout the region. Other studies have previously shown the important role of annual precipitation in

explaining species richness, phylogenetic diversity, and functional diversity in Neotropical amphibians

(Amador et al., 2019; Ochoa-Ochoa et al., 2019). Differences in research and collecting efforts

throughout the Neotropical region could be complicating the interpretation of our genetic diversity map,

mainly because we do not have an equilibrated sampling and our repurposed dataset has a high

concentration of occurrence points in certain regions (e.g., Amazonia, Atlantic forests or Mesoamerica).

In addition, we were not able to recover more than five CytB sequences (our minimum number of

sequences per species) for any Gymnophiona species, nor were we able to recover genetic information for

centrolenid frogs, one of the most diverse taxa in the Neotropics. We were also unable to recover target

sequences for marsupial frogs of the family Hemiphractidae or for Neotropical microhylid frogs. These

two groups have high species richness but are lacking in available evolutionary studies. To alleviate some

of these sampling issues, future studies should consider comparisons of other mitochondrial and nuclear

genes and ideally standardize the markers sequenced across multiple species.

Comparing our results with those for Nearctic amphibians (Barrow et al. 2021), no differences in average

π within species were evident (Nearctic amphibians: average $\pi = 0.028$, n = 299; Neotropical amphibians:

average $\pi = 0.025$, n = 256). However, the highest values of intraspecific π were found in several

Neotropical species; for example, only three Nearctic species had π values > 0.09 compared to 13

Neotropical species with similar or higher values. This disparity between Neotropical and Nearctic

species could relate to differences in demographic history between regions or could be explained by bias

in taxonomic practices (see Chek et al., 2003). It is possible that the higher genetic variation we observed

in Neotropical amphibians is partially due to taxonomic under-splitting, which could lead to severe

conservation implications for this group. We found very high values of π (e.g., > 0.09) for several species,

suggesting the possibility of cryptic species in our dataset. At least nine of the species with high π values

(Table S3) have been considered as species complexes or cryptic species in previous taxonomic studies

(e.g., *Pristimantis altamazonicus* – Ortega-Andrade et al., 2017; *Anomaloglossus degranvillei* – Vacher et

al., 2017; *Phyllobates lugubris* – Márquez et al., 2020; *Dendropsophus decipiens* – Oliveira et al., 2021).

Higher environmental and climatic heterogeneity may lead to diversification dynamics with higher

speciation and lower extinction rates supporting rapid evolution in the Neotropics (Brown, 2014). Under

this scenario, species would have accumulated faster towards the present mainly due to recent geological

and climatic perturbations (e.g., the elevation of the Andes) (Meseguer et al., 2022). The potential under-

split species in our data set are therefore expected to be young or recently diverged, which is one of the

mechanisms causing cryptic diversity (Fišer et al., 2018). We interpret the results of genetic variation of

these species with caution because they may represent multiple taxa, for example, if we were able to split

cryptic species in separate species, range sizes would be smaller impacting the levels of intraspecific

genetic diversity.

We found that levels of π varied among families, with species in certain families such as Dendrobatidae

and Hylidae having the highest π estimates and Ranidae, Rhinodermatidae, and Telmatobiidae having the

lowest (Figure S14). In contrast, the presence of spatial genetic variation (IBD and IBE) within species

appears to be more randomly distributed throughout the phylogeny of Neotropical amphibians. These

findings highlight the value of employing different methodological frameworks as we did in this study.

With the growing size and complexity of biodiversity datasets, machine learning methods such as random

forests prove valuable in identifying potential predictors, even using both complete and reduced datasets.

Combining these methods with phylogenetically informed models allowed us to gain a deeper

understanding of the relationships between predictors and genetic variation within species. For example,

our RF (regression and classification) and *MCMCglmm* models were consistent in identifying range size

and precipitation as the best predictors of genetic variation (Table 2; Figure 2).

Our study provides valuable insights into the distribution of genetic variation in Neotropical amphibians

and identifies important predictors of intraspecific genetic variation. These findings underscore the

importance of considering both nucleotide diversity and spatial genetic variation in the conservation and

management of Neotropical amphibian populations. For example, the results demonstrate the importance

of preserving forest areas, especially in biogeographic areas where the intraspecific π is very low. This

information is also valuable to assess the conservation status of Neotropical amphibian species and

494 consider the impact of threats these taxa could be facing in the future. The current distribution of genetic

495 variation could play a key role in the development of targeted conservation strategies for amphibian

496 species, particularly considering the diverse life histories observed among Neotropical amphibians. For

497 example, certain species within the genera *Atelopus* or *Telmatobius*, which are highly endangered groups,

498 require specific conservation measures focused on preserving aquatic habitats (e.g., streams, ponds, or

499 lakes). These habitats serve as crucial breeding grounds where frogs lay their eggs in shallow water,

500 making their conservation of paramount importance. To address these and other subjects inherent to

501 amphibian ecology and evolution, we suggest that future work should include more information such as

502 genome-scale data and where possible, add more species within the region and globally.

504 **Acknowledgements**

506 This work was supported by a grant to Lisa Barrow from the U.S. National Science Foundation (DEB-

507 2112946). Thank you to the associate editor and two reviewers for their suggestions that helped to

508 improve this manuscript. We thank the members of the Amphibian & Reptile Biodiversity Lab (The

509 University of New Mexico) for their valuable comments on earlier versions of this manuscript, Bryan

510 Carstens (The Ohio State University) for early access to the phylogatR database, and Jorge "Colo" Ortega

511 for use of the photo of *Boana picturata* in Figure 1.

513 **Conflict of Interest**

514 The authors declare no conflict of interest.

516 **Data Availability**

517 Supplementary table S1 and additional supplementary files (e.g., appendices, fasta sequences used for the

518 analyses) are available on Dryad.

519 https://datadryad.org/stash/share/mAYGsdKODg3I8ouXADLggPVcTkVkU7Nqa1wGPl80jiQ

References

Acevedo, A. A., Lampo, M., & Cipriani, R. (2016). The cane or marine toad, *Rhinella marina* (Anura,

Bufonidae): Two genetically and morphologically distinct species. *Zootaxa*, 4103(6), 574.

https://doi.org/10.11646/zootaxa.4103.6.7

Amador, L., Soto☐Gamboa, M., & Guayasamin, J. M. (2019). Integrating alpha, beta, and phylogenetic

diversity to understand anuran fauna along environmental gradients of tropical forests in western Ecuador.

Ecology and Evolution, 9(19), 11040–11052. https://doi.org/10.1002/ece3.5593

Antonelli, A. (2022). The rise and fall of Neotropical biodiversity. *Botanical Journal of the Linnean*

Society, 199(1), 8–24. https://doi.org/10.1093/botlinnean/boab061

Antonelli, A., Zizka, A., Carvalho, F. A., Scharn, R., Bacon, C. D., Silvestro, D., & Condamine, F. L.

(2018). Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National*

Academy of Sciences, *115*(23), 6034–6039. https://doi.org/10.1073/pnas.1713819115

Barrow, L. N., Masiero da Fonseca, E., Thompson, C. E. P., & Carstens, B. C. (2021). Predicting

amphibian intraspecific diversity with machine learning: Challenges and prospects for integrating traits,

geography, and genetic data. *Molecular Ecology Resources*, *21*(8), 2818–2831.

https://doi.org/10.1111/1755-0998.13303

Bazin, E., Glémin, S., & Galtier, N. (2006). Population Size Does Not Influence Mitochondrial Genetic

Diversity in Animals. *Science*, 312(5773), 570–572. https://doi.org/10.1126/science.1122033

Bivand, R. (2022). R Packages for Analyzing Spatial Data: A Comparative Case Study with Areal Data.

Geographical Analysis, 54(3), 488–518. https://doi.org/10.1111/gean.12319

Blanchet, S., Prunier, J. G. & De Kort, H. (2017), Time to go bigger: Emerging patterns in macrogenetics.

Trends in Genetics, 33(9), 579–580. https://doi.org/10.1016/j.tig.2017.06.007

Blomberg, S. P., Garland, T., & Ives, A. R. (2003). Testing for phylogenetic signal in comparative data:

Behavioral traits are more labile. *Evolution*, 57(4), 717–745. https://doi-

org.libproxy.unm.edu/10.1111/j.0014-3820.2003.tb00285.x

Breiman, L. (2001). Random Forests. *Machine Learning 45*, 5–32.

https://doi.org/10.1023/A:1010933404324

Brown, J. H. (2014). Why are there so many species in the tropics? *Journal of Biogeography*, 41(1), 8–

22. https://doi.org/10.1111/jbi.12228

Brüniche◻Olsen, A., Kellner, K. F., & DeWoody, J. A. (2019). Island area, body size and demographic

history shape genomic diversity in Darwin's finches and related tanagers. *Molecular Ecology*, 28(22),

4914–4925. https://doi.org/10.1111/mec.15266

Buffalo, V. (2021). Quantifying the relationship between genetic diversity and population size suggests

natural selection cannot explain Lewontin's Paradox. *ELife*, 10, e67509.

https://doi.org/10.7554/eLife.67509

Calenge, C., & Fortmann-Roe, S. (2023). _adehabitatHR: Home Range Estimation_. R package version

0.4.21, <https://CRAN.R-project.org/package=adehabitatHR>.

Castroviejo-Fisher, S., Guayasamin, J. M., Gonzalez-Voyer, A., & Vilà, C. (2014). Neotropical

diversification seen through glassfrogs. *Journal of Biogeography*, 41(1), 66–80.

https://doi.org/10.1111/jbi.12208

Caviedes-Solis, I. W., Kim, N., & Leaché, A. D. (2020). Species IUCN threat status level increases with

elevation: A phylogenetic approach for Neotropical tree frog conservation. *Biodiversity and*

Conservation, 29(8), 2515–2537. https://doi.org/10.1007/s10531-020-01986-8

Chek, A. A., Austin, J. D., & Lougheed, S. C. (2003). Why is there a tropical–temperate disparity in the

genetic diversity and taxonomy of species? *Evolutionary Ecology Research*, 5, 69 – 77.

De Kort, H., Prunier, J. G., Ducatez, S., Honnay, O., Baguette, M., Stevens, V. M., & Blanchet, S. (2021).

Life history, climate and biogeography interactively affect worldwide genetic diversity of plant and

animal populations. *Nature Communications*, 12(1), 516. https://doi.org/10.1038/s41467-021-20958-2

DeWoody, J. A., Harder, A. M., Mathur, S., & Willoughby, J. R. (2021). The long□standing significance

of genetic diversity in conservation. *Molecular Ecology*, 30(17), 4147–4154.

https://doi.org/10.1111/mec.16051

Dixo, M., Metzger, J. P., Morgante, J. S., & Zamudio, K. R. (2009). Habitat fragmentation reduces

genetic diversity and connectivity among toad populations in the Brazilian Atlantic Coastal Forest.

Biological Conservation, 142(8), 1560–1569. https://doi.org/10.1016/j.biocon.2008.11.016

Duminil, J., Fineschi, S., Hampe, A., Jordano, P., Salvini, D., Vendramin, G. G., & Petit, R. J. (2007).

Can population genetic structure be predicted from life-history traits?. *The American naturalist*, *169*(5),

662–672. https://doi.org/10.1086/513490

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res. 32(5), 1792-1797. doi:10.1093/nar/gkh340

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, *17*(7),

422–433. https://doi.org/10.1038/nrg.2016.58

Elmer, K. R., Bonett, R. M., Wake, D. B., & Lougheed, S. C. (2013). Early Miocene origin and cryptic

diversification of South American salamanders. *BMC Evolutionary Biology*, 13(1), 59.

https://doi.org/10.1186/1471-2148-13-59

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1□km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology*, 37(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Fišer, C., Robinson, C. T., & Malard, F. (2018). Cryptic species as a window into the paradigm shift of

the species concept. *Molecular Ecology*, 27(3), 613–635. https://doi.org/10.1111/mec.14486

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression*, Third edition. Sage, Thousand

Oaks CA. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Frankham, R. (1996). Relationship of genetic variation to population size in wildlife. *Conservation*

Biology, 10(6), 1500–1508. https://doi.org/10.1046/j.1523-1739.1996.10061500.x

Fritz, S. A., & Purvis, A. (2010). Selectivity in Mammalian Extinction Risk and Threat Types: A New

Measure of Phylogenetic Signal Strength in Binary Traits: Selectivity in Extinction Risk. *Conservation*

Biology, 24(4), 1042–1051. https://doi.org/10.1111/j.1523-1739.2010.01455.x

Frost, D. R. (2021). Amphibian Species of the World: an Online Reference. Version 6.1 (Date of access).

Electronic Database accessible at https://amphibiansoftheworld.amnh.org/index.php. American Museum

of Natural History, New York, USA. Doi.org/10.5531/db.vz.0001

Garamszegi, L. Z. (Ed.). (2014). Modern Phylogenetic Comparative Methods and Their Application in

Evolutionary Biology: Concepts and Practice. Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-

662-43550-2

García-Rodríguez, A., Guarnizo, C. E., Crawford, A. J., Garda, A. A., & Costa, G. C. (2021).

Idiosyncratic responses to drivers of genetic differentiation in the complex landscapes of Isthmian Central

America. *Heredity*, 126(2), 251–265. https://doi.org/10.1038/s41437-020-00376-8

Gonzalez-Voyer, A., Padial, J. M., Castroviejo-Fisher, S., De La Riva, I., & Vilà, C. (2011). Correlates of

species richness in the largest Neotropical amphibian radiation: Correlates of species richness. *Journal of*

Evolutionary Biology, 24(5), 931–942. https://doi.org/10.1111/j.1420-9101.2011.02243.x

Greenwell, B. M., & Boehmke, B. C. (2020). Variable Importance Plots—An Introduction to the vip

Package. *The R Journal*, 12(1), 343--366. https://doi.org/10.32614/RJ-2020-013.

Grundler, M. R., Singhal, S., Cowan, M. A., & Rabosky, D. L. (2019). Is genomic diversity a useful

proxy for census population size? Evidence from a species◻rich community of desert lizards. *Molecular*

Ecology, 28(7), 1664–1674. https://doi.org/10.1111/mec.15042

Hackathon, R., Bolker. B., Butler, M., Cowan, P., de Vienne, D., Eddelbuettel, D., Holder, M., Jombart,

T., Kembel, S., Michonneau, F., Orme, D., O'Meara, B., Paradis, E., Regetz, J. & Zwickl D. (2020).

phylobase: Base package for phylogenetic structures and comparative data. R package version 0.8.10.

https://CRAN.R◻project.org/package=phylobase

Hadfield, J. D. (2010). MCMC Methods for Multi-Response Generalized Linear Mixed Models: The

MCMCglmm R Package. *Journal of Statistical Software*, 33(2), 1–22.

https://doi.org/10.18637/jss.v033.i02

Hague, M. T. J., & Routman, E. J. (2016). Does population size affect genetic diversity? A test with

sympatric lizard species. *Heredity*, 116(1), 92–98. https://doi.org/10.1038/hdy.2015.76

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25(15), 1965–

-1978.

Hijmans, R. J. (2021). geosphere: Spherical Trigonometry. R package version 1.5-14. https://CRAN.R-

project.org/package=geosphere

Hijmans, R. J. (2022). raster: Geographic Data Analysis and Modeling. R package version 3.6-3.

https://CRAN.R-project.org/package=raster

Hijmans, R. J., Barbosa, M., Ghosh, A., & Mandel, A. (2023). _geodata: Download Geographic Data_. R

package version 0.5-9, <https://CRAN.R-project.org/package=geodata>.

Hillman, S. S., Drewes, R. C., Hedrick, M. S., & Hancock, T. V. (2014). Physiological Vagility:

Correlations with Dispersal and Population Genetic Structure of Amphibians. *Physiological and*

Biochemical Zoology, *87*(1), 105–112. https://doi.org/10.1086/671109

IUCN. 2022. The IUCN Red List of Threatened Species. Version 2022-1. https://www.iucnredlist.org.

Accessed on 01 May 2022.

Ives, A.R., & Helmus, M.R. (2011). Generalized linear mixed models for phylogenetic analyses of

community structure. *Ecological Monographs*, 81, 511–525.

Josse, C., Navarro, G., Comer, P., Evans, R., Faber-Langendoen, D., Fellows, M., Kittel, G., Menard, S.,

Pyne, M., Reid, M., Schulz, K., Snow, K., & Teague, J. (2003). Ecological Systems of Latin America and

the Caribbean: A Working Classification of Terrestrial Systems. NatureServe, Arlington, VA.

Kabacoff, R. I. (2015). *R in Action* (2nd ed.). Manning Publications.

Kimura, M. (1983). *The Neutral Theory of Molecular Evolution.* Cambridge: Cambridge University

Press.

Larkin, I. E., Myers, E. A., Carstens, B. C., & Barrow, L. N. (2023). Predictors of genomic diversity

within North American squamates. *Journal of Heredity*, 114(2), 131–142.

https://doi.org/10.1093/jhered/esad001

Larsson, A. (2014). AliView: a fast and lightweight alignment viewer and editor for large data sets.

Bioinformatics, *30*(22): 3276-3278. http://dx.doi.org/10.1093/bioinformatics/btu531

Leffler, E. M., Bullaughey, K., Matute, D. R., Meyer, W. K., Ségurel, L., Venkat, A., Andolfatto, P., &

Przeworski, M. (2012). Revisiting an Old Riddle: What Determines Genetic Diversity Levels within

Species? *PLoS Biology*, 10(9), e1001388. https://doi.org/10.1371/journal.pbio.1001388

Leigh, D. M., Van Rees, C. B., Millette, K. L., Breed, M. F., Schmidt, C., Bertola, L. D., Hand, B. K.,

Hunter, M. E., Jensen, E. L., Kershaw, F., Liggins, L., Luikart, G., Manel, S., Mergeay, J., Miller, J. M.,

Segelbacher, G., Hoban, S., & Paz-Vinas, I. (2021). Opportunities and challenges of macrogenetic

studies. *Nature Reviews Genetics*, 22(12), 791–807. https://doi.org/10.1038/s41576-021-00394-0

Levy, E., Byrne, M., Coates, D. J., Macdonald, B. M., McArthur, S., & van Leeuwen, S. (2016).

Contrasting Influences of Geographic Range and Distribution of Populations on Patterns of Genetic

Diversity in Two Sympatric Pilbara Acacias. *PLOS ONE*, 11(10), e0163995.

https://doi.org/10.1371/journal.pone.0163995

Lewontin, R. (1974). *The Genetic Basis of Evolutionary Change.* Columbia University Press.

Liaw A., & Wiener, M. (2002). Classification and Regression by randomForest. R News 2(3), 18--22.

López☐Uribe, M. M., Jha, S., & Soro, A. (2019). A trait☐based approach to predict population genetic

structure in bees. *Molecular Ecology*, 28(8), 1919–1929. https://doi.org/10.1111/mec.15028

Mackintosh, A., Laetsch, D. R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., & Lohse, K.

(2019). The determinants of genetic diversity in butterflies. *Nature Communications*, *10*(1), 3466.

https://doi.org/10.1038/s41467-019-11308-4

Márquez, R., Linderoth, T. P., Mejía☐Vargas, D., Nielsen, R., Amézquita, A., & Kronforst, M. R. (2020).

Divergence, gene flow, and the origin of leapfrog geographic distributions: The history of colour pattern

variation in *Phyllobates* poison☐dart frogs. *Molecular Ecology*, 29(19), 3702–3719.

https://doi.org/10.1111/mec.15598

Menéndez☐Guerrero, P. A., Green, D. M., & Davies, T. J. (2020). Climate change and the future

restructuring of Neotropical anuran biodiversity. *Ecography*, 43(2), 222–235.

https://doi.org/10.1111/ecog.04510

Meseguer, A. S., Michel, A., Fabre, P.-H., Pérez Escobar, O. A., Chomicki, G., Riina, R., Antonelli, A.,

Antoine, P.-O., Delsuc, F., & Condamine, F. L. (2022). Diversification dynamics in the Neotropics

through time, clades, and biogeographic regions. *ELife*, 11, e74503. https://doi.org/10.7554/eLife.74503

Miraldo, A., Li, S., Borregaard, M. K., Flórez-Rodríguez, A., Gopalakrishnan, S., Rizvanovic, M., Wang,

Z., Rahbek, C., Marske, K. A., & Nogués-Bravo, D. (2016). An Anthropocene map of genetic diversity.

Science, 353(6307), 1532–1535. https://doi.org/10.1126/science.aaf4381

Morrone, J. J. (2014). Biogeographical regionalisation of the Neotropical region. *Zootaxa*, *3782*(1), 1.

https://doi.org/10.11646/zootaxa.3782.1.1

Nevo, E. (1978). Genetic variation in natural populations: Patterns and theory. *Theoretical Population*

Biology, 13(1), 121–177. https://doi.org/10.1016/0040-5809(78)90039-4

Ochoa☐Ochoa, L. M., Mejía☐Domínguez, N. R., Velasco, J. A., Marske, K. A., & Rahbek, C. (2019).

Amphibian functional diversity is related to high annual precipitation and low precipitation seasonality in

the New World. *Global Ecology and Biogeography*, 28(9), 1219–1229. https://doi.org/10.1111/geb.12926

Ochoa-Ochoa, L. M., Mejía-Domínguez, N. R., Velasco, J. A., Dimitrov, D., & Marske, K. A. (2020).

Dimensions of amphibian alpha diversity in the New World. *Journal of Biogeography*, 47, 2293 – 2302.

https://doi.org/10.1111/jbi.13948

Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017). AmphiBIO, a

global database for amphibian ecological traits. *Scientific Data*, *4*(1), 170123.

https://doi.org/10.1038/sdata.2017.123

Oliveira, R. F. de, Magalhães, F. de M., Teixeira, B. F. da V., Moura, G. J. B. de, Porto, C. R.,

Guimarães, F. P. B. B., Giaretta, A. A., & Tinôco, M. S. (2021). A new species of the *Dendropsophus*

decipiens Group (Anura: Hylidae) from Northeastern Brazil. *PLOS ONE*, 16(7), e0248112.

https://doi.org/10.1371/journal.pone.0248112

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C.,

D'amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura,

Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., & Kassem, K. R. (2001). Terrestrial Ecoregions of the

World: A New Map of Life on Earth. *BioScience*, *51*(11), 933. https://doi.org/10.1641/0006-

3568(2001)051[0933:TEOTWA]2.0.CO;2

Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, Pearse W (2018). caper: Comparative

Analyses of Phylogenetics and Evolution in R. R package version 1.0.1 https://CRAN.R-

project.org/package=caper

Ortega-Andrade, H. M., Rojas-Soto, O. R., Espinosa de los Monteros, A., Valencia, J. H., Read, M., &

Ron, S. R. (2017). Revalidation of *Pristimantis brevicrus* (Anura, Craugastoridae) with taxonomic

comments on a widespread Amazonian direct-developing frog. Herpetological Journal 26, 81 – 97.

Pagel, M. (1999). Inferring the historical patterns of biological evolution. *Nature*, 401(6756), 877–884.

https://doi.org/10.1038/44766

Paradis, E. (2010). pegas: an R package for population genetics with an integrated-modular approach.

Bioinformatics 26, 419-420.

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* 35, 526 - 528.

Paz, A., Ibáñez, R., Lips, K. R., & Crawford, A. J. (2015). Testing the role of ecology and life history in

structuring genetic variation across a landscape: A trait-based phylogeographic approach. *Molecular*

Ecology, 24(14), 3723–3737. https://doi.org/10.1111/mec.13275

Paz-Vinas, I., Loot, G., Hermoso, V., Veyssiere, C., Poulet, N., Grenouillet, G., & Blanchet, S. (2018).

Systematic conservation planning for intraspecific genetic diversity. *Proceedings of the Royal Society B*

285, 20172746. http://dx.doi.org/10.1098/rspb.2017.2746

721 Pelletier, T. A., & Carstens, B. C. (2018). Geographical range size and latitude predict population genetic

722 structure in a global survey. *Biology Letters*, 14(1), 20170566. https://doi.org/10.1098/rsbl.2017.0566

723 Pelletier, T. A., Parsons, D. J., Decker, S. K., Crouch, S., Franz, E., Ohrstrom, J., & Carstens, B. C.

724 (2022). PHYLOGATR: Phylogeographic data aggregation and repurposing. *Molecular Ecology Resources*,

725 *22*(8), 2830–2842. https://doi.org/10.1111/1755-0998.13673

726 Pereira, H. M. (2016). A latitudinal gradient for genetic diversity. *Science*, 353(6307), 1494–1495.

727 https://doi.org/10.1126/science.aah6730

728 Quiroga-Carmona, M., & D'Elía, G. (2022). Climate influences the genetic structure and niche

729 differentiation among populations of the olive field mouse *Abrothrix olivacea* (Cricetidae: Abrotrichini).

730 *Scientific Reports*, 12(1), 22395. https://doi.org/10.1038/s41598-022-26937-x

731 R Core Team (2022). R: A language and environment for statistical computing. R Foundation for

732 Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

733 Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other things):

734 *phytools: R package*. *Methods in Ecology and Evolution*, *3*(2), 217–223. https://doi.org/10.1111/j.2041-

735 210X.2011.00169.x

736 Rolland, J., Silvestro, D., Schluter, D., Guisan, A., Broennimann, O., & Salamin, N. (2018). The impact

737 of endothermy on the climatic niche evolution and the distribution of vertebrate diversity. *Nature Ecology*

738 *& Evolution*, *2*(3), 459–464. https://doi.org/10.1038/s41559-017-0451-9

739 Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y., Dernat, R.,

740 Duret, L., Faivre, N., Loire, E., Lourenco, J. M., Nabholz, B., Roux, C., Tsagkogeorga, G., Weber, A. A.-

741 T., Weinert, L. A., Belkhir, K., Bierne, N., … Galtier, N. (2014). Comparative population genomics in

742 animals uncovers the determinants of genetic diversity. *Nature*, 515(7526), 261–263.

743 https://doi.org/10.1038/nature13685

Schmidt, C., Munshi-South, J., Dray, S., & Garroway, C. J. (2022). Determinants of genetic diversity and

species richness of North American amphibians. *Journal of Biogeography, 49*, 2005– 2015.

https://doi.org/10.1111/jbi.14480

Stange, M., Barrett, R. D. H., & Hendry, A. P. (2021). The importance of genomic variation for

biodiversity, ecosystems and people. *Nature Reviews Genetics*, *22*(2), 89–105.

https://doi.org/10.1038/s41576-020-00288-7

Tobar-Suárez, C., Urbina-Cardona, N., Villalobos, F., & Pineda, E. (2022). Amphibian species richness

and endemism in tropical montane cloud forests across the Neotropics. *Biodiversity and Conservation*,

31(1), 295–313. https://doi.org/10.1007/s10531-021-02335-z

Vacher, J.-P., Kok, P. J. R., Rodrigues, M. T., Lima, J. D., Lorenzini, A., Martinez, Q., Fallet, M.,

Courtois, E. A., Blanc, M., Gaucher, P., Dewynter, M., Jairam, R., Ouboter, P., Thébaud, C., & Fouquet,

A. (2017). Cryptic diversity in Amazonian frogs: Integrative taxonomy of the genus *Anomaloglossus*

(Amphibia: Anura: Aromobatidae) reveals a unique case of diversification within the Guiana Shield.

Molecular Phylogenetics and Evolution, 112, 158–173. https://doi.org/10.1016/j.ympev.2017.04.017

van den Burg, M. P., Herrando-Pérez, S., & Vieites, D. R. (2020). ACDC, a global database of amphibian

cytochrome-b sequences using reproducible curation for GenBank records. *Scientific Data*, *7*(1), 268.

https://doi.org/10.1038/s41597-020-00598-9

Wang, I. J. (2013). Examining the full effects of landscape heterogeneity on spatial genetic variation: A

multiple matrix regression approach for quantifying geographic and ecological isolation. *Evolution*,

67(12), 3403–3411. https://doi.org/10.1111/evo.12134

Wang, I. J. (2020). Topographic path analysis for modeling dispersal and functional connectivity:

calculating topographic distances using the topoDistance R package. *Methods in Ecology and Evolution*,

11:265-272.

White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between body size

and abundance in ecology. *Trends in Ecology & Evolution*, 22(6), 323–330.

https://doi.org/10.1016/j.tree.2007.03.007

Wieringa, J. G., Boot, M. R., Dantas-Queiroz, M. V., Duckett, D., Fonseca, E. M., Glon, H., Hamilton,

N., Kong, S., Lanna, F. M., Mattingly, K. Z., Parsons, D. J., Smith, M. L., Stone, B. W., Thompson, C.,

Zuo, L., & Carstens, B. C. (2020). Does habitat stability structure intraspecific genetic diversity? It's

complicated... *Frontiers of Biogeography*, 12(2). https://doi.org/10.21425/F5FBG45377

Jetz, W., & Pyron, R.A. (2018). The interplay of past diversification and evolutionary isolation with

present imperilment across the amphibian tree of life. *Nature Ecology & Evolution*, *2*, 850–858.

https://doi.org/10.1038/s41559-018-0515-5

Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R.,

Wevers, J., Grosu, A., Paccini, A., Vergnaud, S., Cartus, O., Santoro, M., Fritz, S., Georgieva, I., Lesiv,

M., Carter, S., Herold, M., Li, Linlin, Tsendbazar, N.E., Ramoino, F., & Arino, O. (2021). ESA

WorldCover 10 m 2020 v100. doi:10.5281/zenodo.5571936.

Zeisset, I., & Beebee, T. J. C. (2008). Amphibian phylogeography: A model for understanding historical

aspects of species distributions. *Heredity*, 101(2), 109–119. https://doi.org/10.1038/hdy.2008.30

https://doi.org/10.1038/hdy.2008.30

Biosketch

Luis Amador is an evolutionary biologist, currently studying the determinants of global amphibian

genetic/genomic diversity as a postdoctoral fellow in the Amphibian and Reptile Biodiversity Lab

(ARBL) at the University of New Mexico (UNM). Irvin Arroyo-Torres contributed to this work as an

undergraduate researcher and is pursuing graduate studies in ecology and conservation at UNM. Lisa

Barrow is Curator of Amphibians and Reptiles at the Museum of Southwestern Biology and principal

investigator of ARBL at UNM, which focuses on research in evolution, ecology, and conservation of

herpetofauna.

Author contributions: L.A. and L.N.B. conceived the ideas; L.A. and I.A.T. assembled the data; L.A.

analyzed the data; and L.A. and L.N.B. led the writing.

Appendices

Appendix S1 Literature used to retrieve occurrences associated with DNA sequences and geographical

data.

Appendix S2 Species not included in the phylogenetic analyses.

Appendix S3 Supplementary tables.

Appendix S4 Supplementary figures.

 Tables

 Table 1. Variables used in this study to predict nucleotide diversity, isolation-by-distance, and isolation-

 by-environment in Neotropical amphibians. Variables with an asterisk (*) indicate non-biologically

 motivated variables and were not used in all analyses. The rationale for considering each variable is

 provided as an example prediction and is not meant to be an exhaustive explanation.

Variable	Value	Rationale
Habitat	Aquatic, Arboreal, Burrowing, Semi-aquatic, Semi-arboreal, Terrestrial	Some habitats may have more natural barriers (e.g., fully aquatic species would have lower dispersal), leading to genetic differentiation of populations.
Activity	Diurnal, Nocturnal	Diurnal or nocturnal activity could restrict or promote gene flow, leading to genetic variation between populations.
Body size	millimeter (mm)	Larger-bodied species are expected to have higher dispersal ability, facilitating gene flow between populations. Larger-bodied species also may have limits on population abundance.
Development mode	Direct, Larval	Species with larval development may have lower dispersal ability since they are tied to water bodies.
Sampling effort*	number of sequences >5	Sampling effort can be considered as a proxy of abundance (population size). Species with more sequences sampled are expected to have higher genetic variation.
Sequence length*	number of base pairs	Species with longer CytB fragments may have more variable sites sampled.
Elevation (mean, minimum, maximum)	meters above sea level (masl)	Species at higher elevations may experience physical barriers or occur in more isolated populations, increasing genetic differentiation.
Latitude (mean, minimum, maximum)	decimal degrees	Latitudinal gradients in climatic variables can influence patterns of distribution and abundance within species.

Precipitation (mean, σ)	precipitation millimeter (mm)	Amphibian diversity is directly affected by precipitation, e.g, species living in areas with high precipitation may have higher abundances.
Temperature (mean, σ)	average temperature (°C)	Temperature may not have strong influence on Neotropical species due to the relatively stable and mild temperatures of the region.
Range size	area in square kilometers (km2)	Species with larger ranges are expected to have higher genetic variation due to larger population sizes and more chances of dispersal.
Order	Anura, Caudata	Order-level classification may be associated with characteristics (e.g., reproductive mode, body size) that influence genetic diversity.
Family	Anura (17 families), Caudata (Plethodontidae)	Family-level classification may be associated with characteristics (e.g., reproductive mode, distribution) that influence genetic diversity.
Biogeographic Unit	17 units shown in Figure 1a	Biogeographic units are associated with different climates, geologic features, or total area that can affect population sizes and gene flow between populations leading to genetic differentiation.
Conservation Status	NE = Not Evaluated, DD = Data Deficient, LC = Least Concern, NT = Near Threatened, VU = Vulnerable, EN = Endangered, CR = Critically Endangered, EX = Extinct	Threatened species with small population sizes are expected to have low genetic diversity; Least concern species are expected to have large range sizes and associated high genetic diversity.
Land cover (mean)	0/1, 1 indicates forest presence	Species living in forested areas (greater number of trees) are expected to have higher abundances and more genetic variation than species living in deforested areas.

  Table 2. Model comparison with deviance information criterion (DIC) scores from each MCMCglmm

model for nucleotide diversity (π), isolation-by-distance (IBD), and isolation-by-environment (IBE).

Comparisons were made with a major and reduced number of predictors and without predictors; with no

random effects and with species and phylogenetic (phylo) random effects. For each model, four different

runs were performed; we present the average DIC score with the standard deviation (SD).

Model detail						Response variables		
Predictors				**Random effects**		π – DIC (SD)	IBD – DIC (SD)	IBE – DIC (SD)
None	Reduced	Major		Species	Phylo			
X						-1002.391 (0.013)	250.5857 (0.012)	244.9121 (0.009)
X				X	X	-1101.188 (0.399)	35.5618 (16.544)	83.9517 (31.095)
	X					-1041.873 (0.028)	247.7395 (0.054)	245.0202 (0.033)
	X			X		-1043.356 (0.124)	5.3151 (0.497)	8.0073 (0.599)
	X				X	-1051.133 (0.035)	248.1518 (0.020)	245.8697 (0.008)
	X			X	X	**-1125.022 (0.302)**	5.4907 (0.756)	8.6062 (3.430)
		X				-1027.762 (0.039)	253.1265 (0.073)	257.0998 (4.340)
		X		X		-1031.574 (0.219)	4.8898 (0.578)	**4.6792 (0.359)**

				-1032.643	252.0281	259.9906
X			X	(0.036)	(0.032)	(0.037)
X	X	X		-1116.827	**4.6857**	5.1812
				(0.698)	**(0.269)**	(0.501)

 Figures

 Figure 1. (a) Neotropical biogeographic units were used in this study. Regionalization follows Morrone

 (2014), with modifications from Olson et al. (2001), Josse et al. (2003), and Antonelli et al. (2018). (b)

 Number of observations/sequences per grid cell. (c) Amphibian genetic diversity patterns in the

 Neotropics. The map uses equal-area grid cells of 350 km.

 Figure 2. Top ranked predictors according to their variable importance for random forest models in

 explaining (a) π, (b) IBD, and (c) IBE. Effect sizes of π (d), IBD (e), and IBE (f) across predictors in

 MCMCglmm analyses. The associated 95% credible intervals (CIs) do not cross zero for range size and

 precipitation in π, for minimum elevation in IBD, and precipitation and maximum latitude in IBE;

 indicating that these predictors are statistically significant.

 Figure 3. (a) Range size (log2) and mtDNA π for all amphibian species in our dataset; (b) precipitation

 (σ) and π relationship; (c) Range size (log2) and π relationship without outliers, the colors indicate

 precipitation (σ) of occurrences in each species. (d) Minimum elevation for species with and without of

 isolation-by-distance (IBD). (e and f) Maximum latitude and precipitation (σ) for species with and

 without isolation-by-environment (IBE). Each box-whisker plot indicates the median (bold lines), the

 interquartile range (boxes), and dots represent outliers.

 Figure 4. Phylogeny of Neotropical amphibians (downloaded and subset from Jetz and Pyron, 2018) with

 values of CytB π presented as light blue bars, and presence (black dots) or absence (grey dots) of IBD and

 IBE in Neotropical amphibians. Silhouettes of frogs and salamanders were obtained from phylopic.org.

Figure 1. (a) Neotropical biogeographic units used in this study. Regionalization follows Morrone

(2014), with modifications from Olson et al. (2001), Josse et al. (2003), and Antonelli et al. (2018). (b)

Number of observations/sequences per grid cell. (c) Amphibian genetic diversity patterns in the

Neotropics. The map uses equal-area grid cells of 350 km.

Figure 2. Top ranked predictors according to their variable importance for random forest models in explaining (a) π, (b) IBD, and (c) IBE. Effect sizes of π (d), IBD (e), and IBE (f) across predictors in *MCMCglmm* analyses. The associated 95% credible intervals (CIs) do not cross zero for range size and precipitation in π, for minimum elevation in IBD, and precipitation and maximum latitude in IBE; indicating that these predictors are statistically significant.

Figure 3. (a) Range size (log2) and mtDNA π for all amphibian species in our dataset; (b) precipitation

(σ) and π relationship; (c) Range size (log2) and π relationship without outliers, the colors indicate

precipitation (σ) of occurrences in each species. (d) Minimum elevation for species with and without of

isolation-by-distance (IBD). (e and f) Maximum latitude and precipitation (σ) for species with and

without isolation-by-environment (IBE). Each box-whisker plot indicates the median (bold lines), the

interquartile range (boxes), and dots represent outliers.

Figure 4. Phylogeny of Neotropical amphibians (downloaded and subset from Jetz and Pyron, 2018) with

values of CytB π presented as light blue bars, and presence (black dots) or absence (grey dots) of IBD and

IBE in Neotropical amphibians. Silhouettes of frogs and salamanders were obtained from phylopic.org.