1 **Modelling Human Post-Implantation Development via**

2 **Extra-Embryonic Niche Engineering**

4 Joshua Hislop[1-3], Amir Alavi[4,5], Qi Song[4,5], Rayna Schoenberger[1,3], Kamyar Keshavarz F.[1-3], Ryan LeGraw[2,3], Jeremy
5 Velazquez [2,3], Tahere Mokhtari[1-3], Mohammad Nasser Taheri[1-3], Matthew Rytel[2,3], Susana M Chuva de Sousa Lopes[8],
6 Simon Watkins[6,7], Donna Stolz[6,7], Samira Kiani [1-3,9], Berna Sozen[10], Ziv Bar-Joseph[4,5], Mo R. Ebrahimkhani[1-3,9] *
7
8 Affiliations:
9
1.   Department of Bioengineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA
15261, USA
2.   Department of Pathology, Division of Experimental Pathology, School of Medicine, University of
Pittsburgh, Pittsburgh, PA 15213, USA
3.   Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA 15261, USA
4.   Computational Biology Department, School of Computer Science, Carnegie Mellon University, Pittsburgh,
PA, USA
5.   Machine Learning Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA,
USA.
6.   Center for Biologic Imaging, University of Pittsburgh, Pittsburgh, PA, USA.
7.   Department of Cell Biology and Molecular Physiology, University of Pittsburgh, Pittsburgh, PA, USA
8.   Department of Anatomy and Embryology, Leiden University Medical Center, Einthovenweg, 2333 ZC
Leiden, the Netherlands
9.   McGowan Institute for Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA 15219, USA
10.   Department of Genetics, Yale School of Medicine, Yale University, New Haven, CT, 06510, USA.
*Correspondence: mo.ebr@pitt.edu

26
27
28 **Abstract**
29

Implantation of the human embryo commences a critical developmental stage that comprises profound
morphogenetic alteration of embryonic and extra-embryonic tissues, axis formation, and gastrulation
events. Our mechanistic knowledge of this window of human life remains limited due to restricted access
to *in vivo* samples for both technical and ethical reasons. Additionally, human stem cell models of early
post-implantation development with both embryonic and extra-embryonic tissue morphogenesis are
lacking. Here, we present iDiscoid, produced from human induced pluripotent stem cells via an
engineered synthetic gene circuit. iDiscoids exhibit reciprocal co-development of human embryonic tissue
and engineered extra-embryonic niche in a model of human post-implantation. They exhibit unanticipated
self-organization and tissue boundary formation that recapitulates yolk sac-like tissue specification with
extra-embryonic mesoderm and hematopoietic characteristics, the formation of bilaminar disc-like
embryonic morphology, the development of an amniotic-like cavity, and acquisition of an anterior-like
hypoblast pole and posterior-like axis. iDiscoids offer an easy-to-use, high-throughput, reproducible, and
scalable platform to probe multifaceted aspects of human early post-implantation development. Thus,
they have the potential to provide a tractable human model for drug testing, developmental toxicology,
and disease modeling.

45
46

Main Text

Decoding mechanisms of human embryogenesis has tremendous biomedical impact, from treating congenital diseases and infertility, to engineering functional human organs. Immediately after implantation, the embryo and co-developing extra-embryonic tissues are profoundly remodeled and initiate morphological changes central to the success of pregnancy, including formation of the amniotic cavity and emergence of yolk sac hematopoiesis[1,2]. However, hidden within the uterine tissues, the key steps of early post-implantation development in humans largely remain beyond our reach, owing to limited access to this stage for both technical and ethical reasons[3,4]. The early post-implantation stage human embryo, with its flat bilaminar disc structure, is morphologically different from its mouse counterpart. As such it may demand new engineering approaches to be efficiently modeled. *In vitro* models of embryogenesis based on both mouse and human-derived cell types have emerged as instrumental platforms to probe human-specific developmental mechanisms[5-20]. However, they often suffer limitations, such as low efficiency and high technical complexity. Traditionally these studies use cocktails of growth factors at supra-physiological levels and encounter challenges finding common media that support diverse cellular fates. Additionally, human post-implantation embryo models with co-developing embryonic and extra-embryonic layers are still missing, and a full integrated human embryo-like model can encounter ethical challenges, which limits our ability to study integrated morphogenetic events of human embryo after implantation[3,21].

Here, we present a human stem cell-derived embryo model, dubbed "*iDiscoid*", that overcomes several of these limitations. The iDiscoid platform leverages an approach that genetically engineers a population of human induced pluripotent stem cells (hiPSCs) with inducible circuits. Upon activation of these genetic circuits, an extra-embryonic niche is formed that triggers self-organization of the adjacent wild-type hiPSC clusters. Using iDiscoids, we demonstrate development of human bilaminar disc-like structures, specification of anterior-like hypoblast cells, lumenogenesis, formation of amnion-like tissue, symmetry breaking leading to posterior-like axis specification, and yolk sac specification with hematopoietic characteristics. iDiscoids model human embryogenesis at an early stage of implantation when the embryo *in vivo* is at its most inaccessible state.

Engineering co-development of embryonic and extra-embryonic endoderm tissues

We engineered an hiPSC cell line with an inducible gene circuit that expresses human GATA6, a key transcription factor for extra-embryonic endodermal fate (henceforth referred to as iGATA6) **(Fig. 1A, Supplementary Fig. 1A)**. These iGATA6 cells express different levels of GATA6 upon the addition of a small molecule, Doxycycline (Dox), distinguishable by expression of the reporter enhanced green fluorescent protein (EGFP) **(Supplementary Fig. 1B)**. When iGATA6 and wild-type (WT) hiPSCs were co-assembled in 3D, EGFP+ extra-embryonic endoderm cells segregated to the outer layer, similar to what occurs during implantation period *in vivo.* However, the overall organization of these aggregates did not recapitulate the structural format of bilaminar disc and yolk sac tissue. This approach yielded aggregates in which the entire EGFP+ extra-embryonic endoderm is in close contact with wild type epiblast-like cells, missing the part of the yolk sac tissue (parietal) that is formed away from the epiblast cells. Additionally, in these 3D aggregates, we found that WT cells could not consistently induce symmetry breaking to amnion-like and epiblast-like domains **(Supplementary Fig. 1C-D)**. Moreover, *in vitro* reattachment of both synthetic and true human blastocysts to a dish to mimic implantation has shown organizational instability that results in tissue organization that cannot reflect the developmental stages approaching gastrulation[7,16,20,22-24]. Mimicking stable co-development of embryonic and extra-embryonic cell types through post-implantation embryonic stages from an initial 3D state is technically difficult to control.

To address these challenges, we aimed to develop an alternative strategy to generate structures off
the cell culture plate via igniting 2D-to-3D self-organization of hiPSCs. The iGATA6-hiPSCs were mixed with
WT hiPSCs, and the mixed population of cells were seeded onto standard culture plates at a defined ratio
and cell density **(Fig. 1A)**. After treatment with Dox, the iGATA6 cells expectedly upregulate GATA6 and
EGFP and exhibit loss of the pluripotency markers NANOG **(Supplementary Fig. 2A)**. During the first 48
hours, from an initial state of random distribution, induced iGATA6 cells and WT cells proliferate, and WT
cells are organized into disc-shaped clusters confined by iGATA6 cells **(Supplementary Video 1, Fig. 1B,**
Supplementary Fig. 1G). Subsequently, the migration of the iGATA6 cells over the upper surface of these
WT clusters occurs in conjunction with 3D growth within the WT disc **(Supplementary Fig. 2C,**
Supplementary Video 2). These events result in a membrane of extra-embryonic endoderm-like cells that
express GATA4 and SOX17 but lack OCT4 over a multilayer WT structure **(Fig. 1A, 1C-D, Supplementary**
Fig. 2C-D). The process of upward iGATA6 migration is in parallel with the deposition of a laminin
membrane surrounding the WT clusters **(Supplementary Fig. 3A-B)**. The laminin deposition by the iGATA6
cells can subsequently trigger cell polarization within cells of the WT cluster, which promotes the
conversion of cellular rosettes into lumens **(Fig. 1C, Supplementary Fig. 3C, Supplementary Video 3,**
4)[25,26]. This is facilitated by expression of PODXL, an apically expressed antiadhesive surface protein
implicated in embryonic lumen formation, and ZO-1, an apical tight junction protein **(Supplementary Fig.**
3C, D)[27]. This laminin deposition and lumen formation is not observed in WT-only culture in equivalent
conditions **(Supplementary Fig. 3E)**.

We observed that within WT clusters for which an iGATA6 layer was not surrounding the cluster,
lumen formation was not observed **(Fig. 1E)**. The number of clusters with no coverage by an iGATA6 layer
progressively decreases by day 5 as the migration of iGATA6 cells moves forward **(Fig. 1E)**. Ultimately,
following expansion of the lumen, the two cell monolayers of epiblast-like WT (NANOG[+]) and endoderm-
like GATA6[+] cells are positioned on either side of a laminin membrane **(Fig. 1A, 1C, Supplementary Fig.**
3C), forming a cellular arrangement similar to the bilaminar disc in the early post-implantation human
embryo **(Supplementary Fig. 1E)**[28]. Analysis of iDiscoid culture media supported high levels of secreted
AFP and APOA1, two proteins known to be produced by primitive endoderm and yolk sac tissues
(Supplementary Fig. 3F).

In our initial analysis, we observed that for WT clusters at day 5 with higher area and often lower
circularity, cavity formation occurred via multiple lumens **(Supplementary Fig. 4A, B)**. In order to minimize
the occurrence of large WT clusters with multiple lumens and control the cluster size and number, we
tested a range of seeding ratio and densities for iGATA6 and WT cells, with the objective to achieve cluster
area most consistently within a size range corresponding to the E9-E17 human bilaminar disc
(Supplementary Fig. 4C, D) (Carnegie #8004, #7700 in [29], Carnegie #7801 in [30]). To this end, we selected a
seeding ratio of 81:5 iGATA6:WT at a density of 54k cells per cm^2 **(Supplementary Fig. 4C, D**, dotted boxes**)**.
When we applied this optimized seeding density, we could detect 74% of iDiscoids in the expected
physiological size range **(Supplementary Fig. 4E)**. Importantly we also observed the majority (70.9%) of
iDiscoids in this size range produced a single lumen **(Supplementary Fig. 4E)**. Hence, by altering the initial
parameters we could control iDiscoid size, number, circularity, and lumenogenesis.

Single cell RNA transcriptomics reveals embryonic and extra-embryonic fate trajectories

To investigate the cellular identities of the developed population, we performed single cell RNA
sequencing (scRNA-seq) from day 0 to day 5 of iDiscoid cultures **(Supplementary Fig. 5-7)**. Performing an
unsupervised comparison of the transcriptomes with the E16-19 human embryo[31], E6-E14 human embryo

samples[24,32], and the E18-20 cynomolgus embryo[33] revealed the acquisition of distinct post-implantation
cellular fates in identified iDiscoid subpopulations **(Fig. 1F, G, Supplementary Fig. 6A-C)**.

145
Beginning as early as 24 hours (D1) after induction, iDiscoids showed the emergence of cell
populations (D1_G1, G2, G3) displaying transcriptomic similarity to human hypoblast, alongside a
corresponding epiblast-like population (D1_W1) **(Supplementary Fig. 6A-C)**. By 36 hours (D1.5), iGATA6
cells become statistically similar to the human E12 hypoblast lineage in from Xiang et al., and this similarity
strengthens through day 5 of culture **(Supplementary Fig. 6B)**[32]. Comparison with the E16-19 human and
cynomolgus embryo shows significant similarity to yolk sac endoderm and hypoblast appearing at D1 and
strengthening through day 5 (p = 6.00×10^{-54} to 2.84×10^{-93} similarity significance to YS Endoderm at D5)
(Fig 1F, Supplementary Fig. 6A, 6C).

154
On day 2, there is significant statistical similarity to yolk sac endoderm within the iGATA6 population
(Clusters D2_G1, D2_G2, D2_G3, D2_G4; p = 1.3×10^{-17} to 9.5×10^{-69}), with the highest GATA6-expressing
cells (Cluster D2-G5) diverging from this fate towards a more yolk sac mesoderm-like state (p = 6.8×10^{-11})
(Fig. 1F, Supplementary Fig. 6A-C). By day 4, we observed the strengthening of the identities of
subpopulations within the iGATA6 and fate divergence within the WT compartment. The lower-GATA6
clusters (D4_G1 and D4_G2) showed increased expression of key yolk sac marker genes (*GATA4*, *PDGFRA*,
LAMA1, *CUBN*, *AMN*, *NODAL*) and statistical similarity to human yolk sac endoderm **(Fig. 1G,**
Supplementary Fig. 5B, 6A-C, p_{G1} = 1.3×10^{-92}, p_{G2} = 1.8×10^{-63}**)**[34]. A "medium" GATA6 population (D4_G3
and D4_G4) showed lower statistical similarity to either of these fates, but has a strong anterior hypoblast
identity (homologous to mouse Anterior Visceral Endoderm, AVE), with specific upregulation of anterior
markers (*LHX1*, *HHEX*, *FZD5*, *CER1*, *LEFTY1*) **(Supplementary Fig. 5B)**[31,35-37]. The highest-GATA6 clusters,
D4_G5 and D4_G6, demonstrated statistical similarity to human yolk sac mesoderm **(Fig. 1G,**
Supplementary Fig. 6A,B, p_{G5} = 1.7×10^{-22}, p_{G6} = 2.5×10^{-25}**)**.Taken together, this demonstrates a divergence
in yolk sac-associated fates within iGATA6 cells that is correlated with different GATA6 expression levels,
with yolk sac endoderm, anterior hypoblast, and yolk sac mesoderm-like cells stratifying between
populations expressing low, medium, and high average levels of *GATA6* **(Supplementary Fig. 6** top row**)**.
Recently observations in cynomolgus monkey by Zhai et al. suggest a similar stratification of GATA6 levels
in a variety of tissues, notably between yolk sac endoderm (lower GATA6) and mesoderm (higher GATA6)
populations[38].

174
Within the WT clusters, identified via a lack of *GATA6* or *EGFP* transcripts, we observe the
divergence of three separate populations from the epiblast-like compartment. The first and largest cluster
(D4_W1) maintained the similarity to human epiblast, with expression of the pluripotency markers
POU5F1 (OCT4), *SOX2*, and *NANOG* **(Supplementary Fig. 5B,** p_{W1} = 1.15×10^{-49}**)**. The second (D4_W2) WT
cluster had lower statistical similarity to human epiblast (p_{W2} = 3.4×10^{-20} to D16-19 Epi), but stronger
similarity to human primitive streak (p_{W2} = 3.6×10^{-95} to D16 PrS) **(Fig. 1G, Supplementary Fig. 5B, 6A)**. Our
assessment of the third wild type cluster at this timepoint, (D4_W3) showed occurrence of a unique
transcriptomic similarity to human and cynomolgus amnion (p_{W3} = 8.0×10^{-15} and 1.6×10^{-25}, respectively)
(Fig. 1G, Supplementary Fig. 5B, 6A, 6C)[17,39,40]. We did not observe a similarity to the trophoblast lineages
(Supplementary Fig. 6).

185
When we integrated all clusters across different days, iGATA6 and WT lineages occupied distinct
compartments within the projection space. However, we did observe spatial positioning of populations
aligned with their time of sampling **(Supplementary Fig 7A)**. Populations sampled at a different time can
show similar fates (i.e., D3 versus D5 iGATA6 yolk sac endoderm cells) **(Supplementary Fig 7B)**. The top

upregulated genes show distinct patterns across different cell fates, with overlapping genes present
across the similar fates sampled at different time points **(Supplementary Fig 7C)**.
192
193 **Specification of amniotic ectoderm in iDiscoids**
194
Primate amniogenesis is structurally and temporally distinct from the process in mouse,
highlighting the need for human model systems to provide a window into understanding human-specific
developmental dynamics[40]. *In vivo*, following lumenogenesis within the epiblast of the primate post-
implantation embryo, the cells of the epiblast undergo dorsal-ventral (D-V) patterning to separate the
epiblast from amniotic ectoderm[15,17,29]. Our single cell transcriptomic analysis in wild type cells showed a
group of cells with transcriptomic similarity to human and cynomolgus amnion that emerges around day
3 **(Fig. 2A-B, Supplementary Fig. 6A, 7B)**. These cells express amnion markers, including *ISL1*, *TFAP2A* (AP-
2α), and *GATA3*, maintaining expression of OCT4 while substantially lacking expression of *NANOG* **(Fig.**
2A-B)[17,39,40]. Our analysis via immunofluorescent staining corroborated this finding, demonstrating a layer
of ISL1[+]/AP-2α[+] in iDiscoids **(Fig. 2C-E, Supplementary Fig. 8A)**. We observe that these cells are spatially
segregated within the cavitated sac-like structure, with the amnion-like lineage positioning away from the
bilaminar iGATA6/WT disc, forming a membrane against the tissue culture dish. WT cells above within the
bilaminar iGATA6/WT disc display expression of NANOG **(Fig. 2C-E)**.
208
Further immunofluorescence staining for pSMAD1, pSMAD5, and pSMAD8/9 (BMP4 effectors)
revealed BMP4 signal transduction in a ring pattern concentrated at the edges of the iDiscoid **(Fig. 2F)**.
scRNA-seq at day 4 also showed specific upregulation of BMP4 targets and related genes within the
amnion-like population **(Supplementary Fig. 8B)**. Recently, the active role of primate amniotic tissue to
promote BMP4 signaling has been suggested, acting downstream of ISL1[40]. It was also reported that BMP4
can promote the formation of ISL1[+] amniotic tissue in a human embryo model[15,17,39]. Hence, we aimed to
inhibit BMP signaling via treatment with the inhibitor Noggin to investigate the relevance of amnion-like
tissue formation in iDiscoid. We observed effective suppression of SMAD1/5/8 phosphorylation **(Fig. 2G)**,
and a marked decrease in amnion fate acquisition assessed by ISL1-expression **(Fig. 2H-I).** However, the
expression of NANOG remained intact in epiblast-like cells **(Fig. 2H)**. To this end, we showed 98.8% of WT
clusters express the amnion marker ISL1 around day 4 of iDiscoid development that was consistent across
experiments. We show that all (n = 87) but one of the observed WT clusters (n = 88) have ISL1 expression
above the expression levels in the noggin treatment group.
222
223   Collectively, we conclude that iDiscoids show robust amniogenesis with formation of a dorsal-
224 ventral (D-V) axis that is dependent on BMP4 signaling, supporting the utility of iDiscoids for studying this
225 key developmental event in humans.
226
227 **Development of anterior hypoblast poles and posterior axis in iDiscoids**
228
In vivo evidence shows a group of cells with anterior visceral endoderm characteristics in human
and non-human primates that have been termed anterior hypoblast[36]. In mouse, this tissue has been
shown to pattern the specification of the posterior axis via the expression of signaling inhibitors including
CER1. However, common *in vitro* stem cell models of the human embryo (i.e., blastoids) have not yet
captured this specification[7,16,20]. We asked whether iDiscoids can exhibit features associated with anterior-
posterior (A-P) axis formation **(Fig. 3)**. Our scRNA-seq analysis of iGATA6 cells as early as day 2 revealed
a cluster of cells that exhibit markers associated with anterior hypoblast. These cells are positive for *CER1*,
LHX1, *HHEX*, *LEFTY1*, and *LEFTY2*, and continued to present these markers in the following days **(Fig.**
3A)[31,35-37]. Additionally, we also detected the emergence of a distinct cluster of cells at day 3 and 4

presenting *TBXT* (Brachyury), *MIXL1*, *GDF3*, and *EOMES*, key markers associated with development of the
posterior pole in the embryo **(Fig. 3B)**[41].
240
Immuno-staining identified domains of cells expressing the anterior hypoblast-related proteins
(CER1, LHX1, HHEX) in a subset of extra-embryonic cells surrounding the WT clusters **(Fig. 3C)**. Out of 396
of iDiscoids assessed we identified 42.4±4.1% (169) discs with polar CER1-expressing domains on day 4
(Fig. 3C), exhibiting a similar efficiency to existing *ex vivo* human blastocyst cultures[36]. In parallel, we
showed development of TBXT[+] domains in 46.4±9.0% (323 out of 746) of WT clusters on day 4 in iDiscoids,
of which 30.1±6.3% (215) of clusters with these domains displayed asymmetric (polar) expression **(Fig.
3D-E)**. These posterior domains are dependent to local BMP4 signaling, as a small molecule, Noggin, could
eliminate TBXT[+] poles with no effect on development of CER1-expressing cells. This supports a role for
BMP4 from amnion in posterior axis specification in line with past studies within these tissues in iDiscoids
(Supplementary Fig. 9)[42].
251
We then analyzed iDiscoid cultures to evaluate A-P axis positionings. On day 4, we observed that
60.4±5.3% (102 of 169) of iDiscoids with polar configuration in CER1-expressing cells possessed a TBXT-
expressing pole **(Fig. 3F)**. Within iDiscoids with polarity in both CER1 and TBXT we observed two distinct
patterns **(Fig. 3F, G, Supplementary Fig. 10)**. 42.2±3.2% exhibit anti-polar configuration, where the
TBXT[+] pole occupies the opposite radial region as the CER1[+] domain, similar to A-P axis positioning in
mouse at E6.5 **(Fig. 3F-H)**. In a second set of polar iDiscoids, we saw high expression of CER1 in the
proximity of TBXT[+] domains **(Fig. 3F-H, Supplementary Fig. 10A)**. This set of iDiscoids shows CER1[+] and
TBXT[+] poles with a same radial region (syn-polar), in contrast to past reported literature. We also noted
expression of CER1 in a subset of TBXT[+] cells in the posterior domains in some cases **(Supplementary
Fig. 10A, B)** and subsequent scRNA-seq analysis showed the presence of a subpopulation of *CER1[+]/TBXT[+]*
cells in the cluster that represents posterior identity **(Supplementary Fig. 10C)**.
263
Next, we aimed to determine whether CER1 expression adjacent to or within the posterior region
observed in iDiscoids can be identified during *in vivo* embryogenesis. To address this question, we first
examined the recent spatial scRNA-seq data from human and non-human primates. In marmoset
embryos[37], we noted non-polar *CER1* at CS5, followed by very slight polarization away from the TBXT-
expressing posterior at CS6 (subtly anti-polar-like) **(Supplementary Fig. 11A)**. However, in human E16-19
(CS7)[31], we observed higher *CER1* transcript expression in the same caudal embryonic region as the
primitive streak **(Supplementary Fig. 11B)**. Lastly, pseudospatial assessment of the E18-20 cynomolgus
embryo shows *CER1* and *TBXT* expression overlapping within the lower region of the embryonic disc
(hypoblast) and embryonic disc compartments (syn-polar-like) **(Supplementary Fig. 11C)**[43]. These three
datasets show a subset of cells or regions that display expression of *CER1* near or within the posterior
domain. To re-examine this phenotype in mouse, we assessed three independent mouse scRNA-seq
datasets spanning ~E6.5-E8 and could also detect a similar pattern of *Cer1* expression within the anterior
boundary of the primitive streak **(Supplementary Fig. 11D-F)**[44-46].
277
Taken together, our analyses of iDiscoids, in tandem with exploration of available *in vivo* data,
suggest that the observed expression and positioning of TBXT[+] and CER1[+] cells in iDiscoids can be seen *in
vivo* and reveal distinct developmental stages. Anti-polar positioning of CER1 and TBXT domains reflect
positioning of anterior hypoblast-like and posterior domains during early primitive streak formation
around E14; syn-polar positioning and co-expression of CER1 and TBXT is reflective of a more advanced
stages post-E14 where CER1 expression acts either to prevent spreading of posterior domain[47] or
represent early mesendoderm commitment derived from a primitive streak[38]. We propose iDiscoids

provide a model to further study A-P axis biology, the formation of anterior hypoblast-like cells, the
posterior pole and their respective cellular progeny in a human context.
287
288 **Specification of yolk sac mesoderm-like cells with hematopoietic potential**
289
The yolk sac represents a multifunctional extra-embryonic tissue that provides nutritional support
for the embryo while also acting as a hub for early blood production that emerges around E16-18 of
human development and matures thereafter[2,48,49]. Multicellular models of human yolk sac morphogenesis
with subsets of mesoderm-derived cells and *in vivo*-like structural characteristics are missing, imposing a
challenge for the dissection molecular underpinnings of the process. We detected progressive
developmental maturation aligned with yolk sac cellular fates in iGATA6 compartment of iDiscoids **(Fig.**
1G, Supplementary Fig 3F, 6). We further asked if iGATA6 cells in iDiscoids could mimic development of
mesodermal subsets within the developing yolk sac endoderm. Hypergeometric statistical analysis of the
clusters with highest expression of GATA6 on day 4 (D4_G6) and 5 (D5_G3) reveals significant similarity to
human yolk sac mesoderm, hematopoietic lineage progenitors, and hemogenic endothelial cells as
compared with the E16-19 human embryo **(Fig. 4A, Supplementary Fig. 7A)**[31]. Examination of the D5_G3
cluster reveals higher expression of the yolk sac mesoderm marker cell surface marker BST2, as well as of
ECM proteins **(Supplementary Fig. 12A)**[49,50]. Subpopulations of cells within this cluster also exhibit a
higher average expression markers of human yolk sac mesoderm (*CREB3L1*, *NR2F2*, *PLAGL1*, *ANXA1*,
NID2)[31,40,50], markers of endothelial cells (*CD34*, *PECAM1* (CD31), *TEK*, *ICAM1, ETV2*)[31,51,52], and markers of
human hematopoiesis (*GATA1*, *GATA2*, *KLF1*, *KLF2*, *ZEB1*, *ZEB2*, *GYPA* (CD235a), and *GYPB* (CD235b)) **(Fig.**
4B, Supplementary Fig. 12B)[31,49,53].
307
Hematopoietic processes emerge *in vivo* in the form of blood islands within yolk sac tissues. In
iDiscoids, immunofluorescence analysis initially reveals spindle shape cells positive for TAL1 (scl), a key
regulator of blood development[54]. Further analysis shows a subset of these cells were also positive for
CD34 and ERG, markers associated with hematopoietic and endothelial fates or RUNX1, a master
transcription factor of hematopoiesis **(Supplementary Fig. 13A-C)**[54]. Further analysis shows a subset of
these cells were also positive for CD34 and ERG, markers associated with hematopoietic and endothelial
fates or RUNX1, a master transcription factor of hematopoiesis **(Supplementary Fig. 13A-C)**[55,56]. Image
analysis of z confocal sections shows ERG$^+$ cells consistently emerged underneath an iGATA6 layer **(Fig.**
4C, Supplementary Fig. 13D), a morphology that may mimic an initial *in vivo* localization similar to human
yolk sac mesoderm, where this tissue is positioned against ECM in the area underneath the yolk sac
endoderm[55,56]. Observed CD34$^+$/TAL1$^+$ cells were also EGFP$^+$, indicating they develop from an iGATA6-
derived parental population, supporting a hypoblast origin **(Fig. 4C)**. Staining for other endothelial markers
KDR (VEGFR2) and CDH5 also confirmed similar localization in day 5 iDiscoids **(Supplementary Fig. 13E)**.
A colony-forming unit (CFU) assay initiated with CD34$^+$ cells followed by flow cytometry analysis of the
cultures, showed generation of CD45$^+$ cells, erythroid cells (CD71$^+$, CD235a$^-$ (early stage) or CD235a$^+$ (late
stage)) as well as CD11b$^+$ myeloid cells **(Supplementary Fig. 13F)**.
324
Having observed early hematopoietic potential in day 5 iDiscoid, we followed iDiscoids beyond
day 5 to better examine the hematopoietic potential of the system. We observed minimal expansion of
the CD34+ area when the culture was maintained on mTeSR **(Supplementary Fig. 3G)**. Therefore, to
facilitate the differentiation of progenitor cells, we used basal media (IMDM without Dox) after day 5. We
observed the expansion of the CD34$^+$ endothelial-like cell area significantly between day 5 and day 12,
from covering 1.2±0.46% of the culture area on day 5 to covering 14.7±6.0% of the culture area at day 12,
an expansion that was also confirmed by flow cytometry analysis **(Supplementary Fig. 13G, H)**. Around
this time, we observed the emergence of condensed areas containing spherical cells within the yolk sac-

like compartment **(Fig. 4D)**. Immunofluorescence staining of these areas revealed the presence of a CD31[+] endothelial layer surrounding spherical cells expressing Leukosialin (CD43), a marker of hematopoietic progenitors **(Fig. 4E)**[53,57]. Further investigation of these spherical cells found that many expressed the erythropoietic marker genes CD235a and hemoglobin **(Supplementary Fig. 14A, 15A)**[31]. Having observed yolk sac mesodermal and hemogenic progenitors of this cell type primarily correlating within the GATA6-hi clusters in scRNA-seq, we further supplemented iDiscoid cultures at the time of seeding with high-GATA6 expressing cells and observed significantly increased generation of these CD43[+] hematopoietic foci **(Fig. 4F-G, Supplementary Fig. 15B)**, demonstrating predictable and controllable fate augmentation at the time of seeding.

Further investigation of the cell types observed at day 12 revealed cells with markers of yolk sac-derived myeloid lineages, including macrophages (CD33[+]/CX$_3$CR1[+]) and megakaryocytes (CD41[+]/CD42b[+]) **(Supplementary Fig. 14B-C)**[58-60]. Structural image analysis of CD43[+] foci show intricate 3D hierarchical organization, with CD34[+] endothelial vessels packed between FOXA2[+] endodermal cells occupying the top compartment and Desmin[+] mesodermal cells forming the basal layer **(Fig. 4H-J, Supplementary Fig. 16A-B)**. This is in line with the tissue morphology reported *in vivo* in blood islands of the developing human yolk sac at around E19-23 **(Fig. 4K)**[49]. Late time point analysis of day 21 iDiscoid cultures using scRNA-seq and flow cytometry similarly supports emergence of populations aligned with those expected from the human yolk sac[61]. This analysis shows identification of: *CDH5* (VE-cadherin)[+] /*CD34*[+] /*CD31*[+] /*KDR* (VEGFR2)[+] /*CD93* (AA4.1)[+] /*NT5E* (CD73)[−] /*GYPA*[−] hemogenic endothelium-like cells; *GYPA*[+] /*GYPB*[+] /*TRDN* (CD71)[+] /*HBG1*[+] /*HBE1*[+] /*HBZ*[+] erythrocyte-like cells; *CD226*[+] /*CLEC1B*[+] /*PLEK*[+] /*GP1BA* (CD42b)[+] /*GATA1*[+] megakaryocyte-like cells; and *CD33*[+] /*CD68*[+] /*CD45*[+] /*CD53*[+] /*CX3CR1*[+] macrophage-like cells **(Supplementary Fig. 14D-F)**. Flow cytometry analysis of these cultures also corroborated the development of a CD43[+]/CD235a[+]/CD71[+] erythrocyte-like population, a CD43[+]/CD33[+]/CD31[+] macrophage-like population, and a CD43[+]/CD42b[+] megakaryocyte-like population **(Supplementary Fig. 14G)**.

Discussion

iDiscoids, presented here, develop both embryonic and extra-embryonic lineages with several features observed together for the first time in an *in vitro* stem cell-based human developmental model. These features show close proximity to native post-implantation human embryogenesis yet are uniquely observed together in an hiPSC-based system **(Supplementary Fig. 17)**. The iDiscoid platform shows morphogenesis of bilaminar disc-like structures including the formation of an amniotic cavity, the segregation of amnion from pluripotent epiblast-like cells, the generation of anterior hypoblast-like cells and a posterior axis, as well as yolk sac morphogenesis. Our extended follow up revealed morphologically relevant blood islands and the specification of hematopoietic lineages **(Supplementary Fig. 17)**. In our follow up, WT clusters exhibited characteristics mainly aligned with ectodermal fates, which is a subject for future studies and optimization **(Supplementary Fig. 18)**. We also show the utility of the iDiscoid platform to study developmental processes.

From an engineering angle, iDiscoid platform highlights the power of tissue niche engineering and co-differentiation to promote morphogenetic events *in vitro*. It demonstrates that a combination of genetic circuit-based tissue engineering with the formation of tissue-tissue boundaries enables geometric confinement and subsequent self-organization in 3D. Use of genetic circuits can also alleviate the need for complex regimes of growth factors to support heterotypic tissues. The extra-embryonic differentiation of iDiscoids is pre-programmed into undifferentiated hiPSCs via an inducible genetic switch and mixed with WT hiPSCs; hence, the undifferentiated cell mix can be expanded, cryostored, and shipped for use on demand, requiring only 2D culture plates, commercially available medium, and addition of a small

molecule, Dox **(Supplementary Fig. 19)**. The high-throughput format, efficient generation, compatibility with live imaging, and an easy-to-implement protocol enables establishment and utility across different labs. These characteristics, coupled with the potential to be produced from hiPSCs with diverse genetic backgrounds **(Supplementary Fig. 20)**, facilitate the studies on early stages of post-implantation, opening new routes for drug testing, developmental toxicology, and tractable disease modeling with fewer ethical concerns in the human context.

References

Muller, F. & O'Rahilly, R. The primitive streak, the caudal eminence and related structures in staged human embryos. *Cells Tissues Organs* **177**, 2-20, doi:10.1159/000078423 (2004).

Palis, J. & Yoder, M. C. Yolk-sac hematopoiesis: the first blood cells of mouse and man. *Exp Hematol* **29**, 927-936, doi:10.1016/s0301-472x(01)00669-5 (2001).

Clark, A. T. *et al.* Human embryo research, stem cell-derived embryo models and in vitro gametogenesis: Considerations leading to the revised ISSCR guidelines. *Stem Cell Reports* **16**, 1416-1424, doi:https://doi.org/10.1016/j.stemcr.2021.05.008 (2021).

Pera, M. F. Human embryo research and the 14-day rule. *Development* **144**, 1923-1925, doi:10.1242/dev.151191 (2017).

Beccari, L. *et al.* Multi-axial self-organization properties of mouse embryonic stem cells into gastruloids. *Nature* **562**, 272-276, doi:10.1038/s41586-018-0578-0 (2018).

Harrison, S. E., Sozen, B., Christodoulou, N., Kyprianou, C. & Zernicka-Goetz, M. Assembly of embryonic and extraembryonic stem cells to mimic embryogenesis in vitro. *Science* **356**, doi:10.1126/science.aal1810 (2017).

Kagawa, H. *et al.* Human blastoids model blastocyst development and implantation. *Nature* **601**, 600-605, doi:10.1038/s41586-021-04267-8 (2022).

Karzbrun, E. *et al.* Human neural tube morphogenesis in vitro by geometric constraints. *Nature* **599**, 268-272, doi:10.1038/s41586-021-04026-9 (2021).

Li, R. *et al.* Generation of Blastocyst-like Structures from Mouse Embryonic and Adult Cell Cultures. *Cell* **179**, 687-702 e618, doi:10.1016/j.cell.2019.09.029 (2019).

Simunovic, M. *et al.* A 3D model of a human epiblast reveals BMP4-driven symmetry breaking. *Nat Cell Biol* **21**, 900-910, doi:10.1038/s41556-019-0349-7 (2019).

Sozen, B. *et al.* Self-Organization of Mouse Stem Cells into an Extended Potential Blastoid. *Dev Cell* **51**, 698-712 e698, doi:10.1016/j.devcel.2019.11.014 (2019).

Veenvliet, J. V. *et al.* Mouse embryonic stem cells self-organize into trunk-like structures with neural tube and somites. *Science* **370**, doi:10.1126/science.aba4937 (2020).

Warmflash, A., Sorre, B., Etoc, F., Siggia, E. D. & Brivanlou, A. H. A method to recapitulate early embryonic spatial patterning in human embryonic stem cells. *Nat Methods* **11**, 847-854, doi:10.1038/nmeth.3016 (2014).

Yu, L. *et al.* Blastocyst-like structures generated from human pluripotent stem cells. *Nature* **591**, 620-626, doi:10.1038/s41586-021-03356-y (2021).

Zheng, Y. *et al.* Controlled modelling of human epiblast and amnion development using stem cells. *Nature* **573**, 421-425, doi:10.1038/s41586-019-1535-2 (2019).

Liu, X. *et al.* Modelling human blastocysts by reprogramming fibroblasts into iBlastoids. *Nature* **591**, 627-632, doi:10.1038/s41586-021-03372-y (2021).

Shao, Y. *et al.* A pluripotent stem cell-based model for post-implantation human amniotic sac development. *Nature Communications* **8**, 208, doi:10.1038/s41467-017-00236-w (2017).

Mathew, B. *et al.* Mouse ICM Organoids Reveal Three-Dimensional Cell Fate Clustering. *Biophysical Journal* **116**, 127-141, doi:https://doi.org/10.1016/j.bpj.2018.11.011 (2019).

19 Schröter, C., Rué, P., Mackenzie, J. P. & Martinez Arias, A. FGF/MAPK signaling sets the switching
threshold of a bistable circuit controlling cell fate decisions in embryonic stem cells.
Development **142**, 4205-4216, doi:10.1242/dev.127530 (2015).
20 Fan, Y. *et al.* Generation of human blastocyst-like structures from pluripotent stem cells. *Cell*
Discovery **7**, 81, doi:10.1038/s41421-021-00316-8 (2021).
21 Weatherbee, B. A. T., Cui, T. & Zernicka-Goetz, M. Modeling human embryo development with
embryonic and extra-embryonic stem cells. *Dev Biol* **474**, 91-99,
doi:10.1016/j.ydbio.2020.12.010 (2021).
22 Zhao, C. *et al.* Reprogrammed blastoids contain amnion-like cells but not trophectoderm.
bioRxiv, 2021.2005.2007.442980, doi:10.1101/2021.05.07.442980 (2021).
23 Deglincerti, A. *et al.* Self-organization of the in vitro attached human embryo. *Nature* **533**, 251-
254, doi:10.1038/nature17948 (2016).
24 Chhabra, S. & Warmflash, A. BMP-treated human embryonic stem cells transcriptionally
resemble amnion cells in the monkey embryo. *Biol Open* **10**, doi:10.1242/bio.058617 (2021).
25 Bedzhov, I. & Zernicka-Goetz, M. Self-organizing properties of mouse pluripotent cells initiate
morphogenesis upon implantation. *Cell* **156**, 1032-1044, doi:10.1016/j.cell.2014.01.023 (2014).
26 Li, S., Edgar, D., Fassler, R., Wadsworth, W. & Yurchenco, P. D. The role of laminin in embryonic
cell polarization and tissue organization. *Dev Cell* **4**, 613-624, doi:10.1016/s1534-5807(03)00128-
x (2003).
27 Shahbazi, M. N. *et al.* Pluripotent state transitions coordinate morphogenesis in mouse and
human embryos. *Nature* **552**, 239-243, doi:10.1038/nature24675 (2017).
28 O'Rahilly, R., Müller, F. & Washington, C. I. o. *Developmental Stages in Human Embryos:*
Including a Revision of Streeter's "Horizons" and a Survey of the Carnegie Collection. (Carnegie
Institution of Washington, 1987).
29 Hertig, A. T. & Rock, J. Two human ova of the pre-villous stage, having a developmental age of
about 8 and 9 days respectively. *Contrib Embryol* **33**, 169-186 (1949).
30 Heuser, C. H., Rock, J. & Hertig, A. T. Two human embryos showing early stages of the definitive
yolk sac. *Contributions to embryology* **201**, 85-99 (1945).
31 Tyser, R. C. V. *et al.* Single-cell transcriptomic characterization of a gastrulating human embryo.
Nature **600**, 285-289, doi:10.1038/s41586-021-04158-y (2021).
32 Xiang, L. *et al.* A developmental landscape of 3D-cultured human pre-gastrulation embryos.
Nature **577**, 537-542, doi:10.1038/s41586-019-1875-y (2020).
33 Ma, H. *et al.* In vitro culture of cynomolgus monkey embryos beyond early gastrulation. *Science*
366, doi:10.1126/science.aax7890 (2019).
34 Cindrova-Davies, T. *et al.* RNA-seq reveals conservation of function among the yolk sacs of
human, mouse, and chicken. *Proc Natl Acad Sci U S A* **114**, E4753-E4761,
doi:10.1073/pnas.1702560114 (2017).
35 Costello, I. *et al.* Lhx1 functions together with Otx2, Foxa2, and Ldb1 to govern anterior
mesendoderm, node, and midline development. *Genes Dev* **29**, 2108-2122,
doi:10.1101/gad.268979.115 (2015).
36 Mole, M. A. *et al.* A single cell characterisation of human embryogenesis identifies pluripotency
transitions and putative anterior hypoblast centre. *Nat Commun* **12**, 3679, doi:10.1038/s41467-
021-23758-w (2021).
37 Bergmann, S. *et al.* Spatial profiling of early primate gastrulation in utero. *Nature* **609**, 136-143,
doi:10.1038/s41586-022-04953-1 (2022).
38 Zhai, J. *et al.* Primate gastrulation and early organogenesis at single-cell resolution. *Nature* **612**,
732-738, doi:10.1038/s41586-022-05526-y (2022).

39   Shao, Y. *et al.* Self-organized amniogenesis by human pluripotent stem cells in a biomimetic
implantation-like niche. *Nature Materials* **16**, 419-425, doi:10.1038/nmat4829 (2017).
40   Yang, R. *et al.* Amnion signals are essential for mesoderm formation in primates. *Nat Commun*
12, 5126, doi:10.1038/s41467-021-25186-2 (2021).
41   Bardot, E. S. & Hadjantonakis, A.-K. Mouse gastrulation: Coordination of tissue patterning,
specification and diversification of cell fate. *Mechanisms of Development* **163**, 103617,
doi:https://doi.org/10.1016/j.mod.2020.103617 (2020).
42   Chhabra, S., Liu, L., Goh, R., Kong, X. & Warmflash, A. Dissecting the dynamics of signaling events
in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human
gastruloids. *PLoS Biol* **17**, e3000498, doi:10.1371/journal.pbio.3000498 (2019).
43   Cui, G. *et al.* Spatial molecular anatomy of germ layers in the gastrulating cynomolgus monkey
embryo. *Cell Reports* **40**, 111285, doi:https://doi.org/10.1016/j.celrep.2022.111285 (2022).
44   Mittnenzweig, M. *et al.* A single-embryo, single-cell time-resolved model for mouse gastrulation.
Cell **184**, 2825-2842.e2822, doi:10.1016/j.cell.2021.04.004 (2021).
45   Peng, G. *et al.* Spatial Transcriptome for the Molecular Annotation of Lineage Fates and Cell
Identity in Mid-gastrula Mouse Embryo. *Dev Cell* **36**, 681-697, doi:10.1016/j.devcel.2016.02.020
(2016).
46   Pijuan-Sala, B. *et al.* A single-cell molecular map of mouse gastrulation and early organogenesis.
Nature **566**, 490-495, doi:10.1038/s41586-019-0933-9 (2019).
47   Stuckey, D. W., Di Gregorio, A., Clements, M. & Rodriguez, T. A. Correct patterning of the
primitive streak requires the anterior visceral endoderm. *PLoS One* **6**, e17620,
doi:10.1371/journal.pone.0017620 (2011).
48   Nakazawa, F., Alev, C., Jakt, L. M. & Sheng, G. Yolk sac endoderm is the major source of serum
proteins and lipids and is involved in the regulation of vascular integrity in early chick
development. *Dev Dyn* **240**, 2002-2010, doi:10.1002/dvdy.22690 (2011).
49   Ross, C. & Boroviak, T. E. Origin and function of the yolk sac in primate embryogenesis. *Nat*
Commun **11**, 3760, doi:10.1038/s41467-020-17575-w (2020).
50   Pham, T. X. A. *et al.* Modeling human extraembryonic mesoderm cells using naive pluripotent
stem cells. *Cell Stem Cell* **29**, 1346-1365.e1310, doi:https://doi.org/10.1016/j.stem.2022.08.001
(2022).
51   Bruveris, F. F. *et al.* Human yolk sac-like haematopoiesis generates RUNX1-, GFI1- and/or GFI 1B-
dependent blood and SOX17-positive endothelium. *Development* **147**, doi:10.1242/dev.193037
(2020).
52   Koyano-Nakagawa, N. *et al.* Etv2 is expressed in the yolk sac hematopoietic and endothelial
progenitors and regulates Lmo2 gene expression. *Stem Cells* **30**, 1611-1623,
doi:10.1002/stem.1131 (2012).
53   Vanhee, S. *et al.* In vitro human embryonic stem cell hematopoiesis mimics MYB-independent
yolk sac hematopoiesis. *Haematologica* **100**, 157-166, doi:10.3324/haematol.2014.112144
(2015).
54   Vagapova, E. R., Spirin, P. V., Lebedev, T. D. & Prassolov, V. S. The Role of TAL1 in Hematopoiesis
and Leukemogenesis. *Acta Naturae* **10**, 15-23 (2018).
55   Enders, A. C. & King, B. F. in *The Human Yolk Sac and Yolk Sac Tumors* (ed Francisco F. Nogales)
33-47 (Springer Berlin Heidelberg, 1993).
56   Knoth, M. & Larsen, J. F. Ultrastructure of a human implantation site. *Acta Obstet Gynecol Scand*
51, 385-393, doi:10.3109/00016347209156879 (1972).
57   Vodyanik, M. A., Thomson, J. A. & Slukin, II. Leukosialin (CD43) defines hematopoietic
progenitors in human embryonic stem cell differentiation cultures. *Blood* **108**, 2095-2105,
doi:10.1182/blood-2006-02-003327 (2006).

58   Stremmel, C. *et al.* Yolk sac macrophage progenitors traffic to the embryo during defined stages
of development. *Nature Communications* **9**, 75, doi:10.1038/s41467-017-02492-2 (2018).
59   Kappelmayer, J. *et al.* Consecutive Appearance of Coagulation Factor XIII Subunit A in
Macrophages, Megakaryocytes, and Liver Cells During Early Human Development. *Blood* **86**,
2191-2197, doi:https://doi.org/10.1182/blood.V86.6.2191.bloodjournal8662191 (1995).
60   Miyawaki, K. *et al.* Identification of unipotent megakaryocyte progenitors in human
hematopoiesis. *Blood* **129**, 3332-3343, doi:10.1182/blood-2016-09-741611 (2017).
61   Wittamer, V. & Bertrand, J. Y. Yolk sac hematopoiesis: does it contribute to the adult
hematopoietic system? *Cellular and Molecular Life Sciences* **77**, 4081-4091, doi:10.1007/s00018-
020-03527-6 (2020).

533
Ethics regarding development of iDiscoids
This research was approved and performed under the oversight of the University of Pittsburgh
Human Stem Cell Research Oversight Committee. The work that has been reported in this study followed
2016 Guidelines for Stem Cell Research and Clinical Translation as well as 2021 ISSCR Guidelines on Ethical
Standards for Stem Cell Embryo Models[62,63]. To this end, iDiscoids do not involve human embryonic stem
cells and are generated from human iPSCs that are derived from human somatic cells such as fibroblasts.
iDiscoids are attached to a cell culture dish, lacking an extra-embryonic trophectoderm tissue critical for
full integrated embryo development and implantation in the uterine cavity. The yolk sac cavity in iDiscoid
is not closed, and the tissue cannot be harvested for any implantation without substantial disruption of
their structures. TBXT$^+$ posterior domains were observed during the study but were not sustained during
the development of our system. These features collectively restrict the ability of this model to undergo
the full integrated development of a human embryo *in vitro* and/or implant to support further
development of a conceptus *in vivo*.
547

Acknowledgements
This work was supported by a Startup Fund from the Department of Pathology at University of
Pittsburgh. We acknowledge support from the Center for Biological Imaging and the University of
Pittsburgh Flow Cytometry Core facilities. Characterization of hematopoietic cellular subsets was partly
supported by an R01 from the National Heart Lung and Blood Institute (HL141805) to M.R.E.
Development and characterization of the iPSC lines used in these studies were partly supported by an
R01 from the National Institute of Biomedical Imaging and Bioengineering (EB028532) and NSF award #
2134999 to M.R.E. The computational analysis of published embryo datasets was partly supported by
NIH grants 1R01GM122096, OT2OD026682, 1U54AG075931 and 1U24CA268108 to Z.B.J. This work is
also partly supported by the Pittsburgh Liver Research Center (NIHNIDDK P30DK120531), the Cellular
Approaches to Tissue Engineering and Regenerative Medicine Training Grant (T32 EB001026), and NIH
1S10OD019973-01. We would like to thank Katherine Helfrich and Michael Calderon at the Center for
Biological Imaging for their assistance related to imaging. We also thank Alaina Sutherland, Chloe Hislop,
and Raini Porterfield for helpful discussion during this work. Biorender was used to create several of the
schematics shown.

Author Contributions
J.H. and M.R.E. conceived the study; M.R.E., Z.B.J., B.S., and S.K. supervised the project; J.H.,
M.R.E., S.K., A.A., K.K.F., J.V., R.L., D.S., Z.B.J. conceived the methodology for experiments; J.H., R.S., K.K.F.,
J.V., R.L., M.N.T., and M.R. performed iDiscoid experiments; J.H., R.S., K.K.F., T.M., M.N.T., S.W. and D.S.
performed imaging and analysis; J.H., A.A., and Q.S. performed the bioinformatic analysis of scRNA-seq
datasets; R.L. and K.K.F. performed flow cytometry and analysis; J.H., A.A., K.K.F., J.V., R.L., D.S., M.R.E.
performed visualization of the data; M.R.E., S.K., Z.B.J., and J.H. acquired funding for the project; J.H. and

M.R.E. wrote the original draft of the manuscript; J.H., R.S., K.K.F., M.R.E., B.S., S.M.C.S.L., A.A., J.V., S.K.,
and Z.B.J. were involved in editing and input to the final draft of the manuscript.
573
574 **Competing Interests:**
575 The authors declare no conflicts of interest. J.H., S.K., and M.R.E. have filed for IP for the
576 technology presented here.

Figure 1

Figure 1: Engineering co-development of embryonic and extra-embryonic endoderm tissues in iDiscoids

(A) Schematic demonstrating cell mixing and subsequent organization of iGATA6 (green) and WT (strongly orange) hiPSCs after GATA6 induction by Dox.

(B) Immunofluorescence staining demonstrating self-organization of iDiscoids in the cultures. Scale bar = 1000 μm

(C) Live time lapse images of one iDiscoid showing growth of a central lumen from a rosette indicated by an arrow. Scale bar = 50 μm.

(D) Phase and fluorescence images from live day 4 iDiscoid cultures showing developed iDiscoid morphologies with EGFP-expressing iGATA6 around a WT cluster. Scale bar = 100 μm

(E) Quantification of characteristics of WT clusters possessing different iGATA6 coverage and lumen formation characteristics. No islands without coverage were observed possessing a lumen (-/+ = 0%)

(F) Single cell UMAP and hypergeometric statistical comparisons of differentially expressed gene (DEG) lists between day 2 iDiscoid and human and NHP embryo single cell datasets. Labels above heatmaps indicate pre-implantation versus post-implantation embryo sample comparisons; gray bars under labels indicated NHP comparisons.

(G) Single cell UMAP and hypergeometric statistical comparisons of differentially expressed gene (DEG) lists between day 4 iDiscoid and human and NHP embryo single cell datasets. Black boxes highlight high DEG similarity (high P values) to relevant populations of interest.

Figure 2

A Wild-Type

D0
D0.5
D1
D1.5
D2
D3
D4
D5

B Amnion-like

OCT4 (POU5F1) NANOG

ISL1 TFAP2A GATA3

Expression Level
7.5
5.0
2.5

C Day 4
EGFP-GATA6 NANOG ISL1

D
EGFP-GATA6 NANOG ISL1
Z = 0µm   Z = 20µm   Z = 30µm

Bottom ← → Top

E
GATA6+
NANOG+
ISL1+
Cell Culture Dish

Top of Cavity

Bottom of Cavity

F Day 4
EGFP-GATA6 pSMAD1/5/8
F-actin

G D2 100ng/ml Noggin
Day 4
EGFP-GATA6 pSMAD1/5/8
F-actin

H D2 100ng/ml Noggin
Day 4
EGFP-GATA6 ISL1 NANOG
F-actin

I
****
ISL1 Intensity (a.u.)
Ctrl   D2 Noggin

Figure 2: Amniotic-like cavity formation and expansion

(A) Merged UMAP of all WT lineages from iDiscoid labeled by day of development.

(B) The merged WT population showing the compartment expressing markers of amnion. *ISL1*, *TFAP2A* (AP-2$^\alpha$), and *GATA3* are expressed in this area, while *NANOG* is negative and OCT4 is low.

(C) An orthogonal slice of an individual iDiscoid showing top-bottom compartmentalization of ISL1$^+$ (cyan) and NANOG$^+$ (red) cells while iGATA6 cells cover the top (white).

(D) Horizontal z-slices at the indicated distance from the dish of the WT cluster from (C).

(E) Schematic showing the position of each population within a single iDiscoid. Dotted lines indicate the area of slices shown.

(F) Expression patterns of BMP4 effectors (phosphorylated SMAD1, SMAD5, and SMAD8/9) in iDiscoid. Lower images show a lateral slice of the WT disc shown.

(G) Immunofluorescence staining for the BMP4 effectors (phosphorylated SMAD1, SMAD5, and SMAD8/9) in iDiscoid at day 4 after application of the inhibitor Noggin at Day 2 of development. Lower images show a lateral slice of the WT disc shown.

(H) Immunofluorescence staining for ISL1 and NANOG in iDiscoid at day 4 after application of Noggin at Day 2 of development. Lower images show a lateral slice of the WT disc shown.

(I) Scatterplot showing ISL1 expression (D4) intensities within WT clusters; control (Ctrl) iDiscoid versus BMP4 inhibition (Noggin) conditions. n[control] = 87, n[Noggin] = 134, **** $p < 0.0001$

Scale bars = 100 µm. n represents separate biological samples from a single experiment harvested on the corresponding day.

Figure 3

Figure 3: Anterior hypoblast-like domain and posterior axis in iDiscoids

(A) Merged UMAP of all iGATA6 lineages showing the compartments expressing markers of anterior hypoblast. Two separate domains of these markers were observed, one within day 2 iDiscoid cells and one within day 3-5 iDiscoid cells. (UMAP labeled by day can be seen in Supplementary Figure 5)

(B) Merged UMAP of all WT lineages showing the compartment expressing markers of the posterior axis and primitive streak.

(C) Immunofluorescence staining showing a z-slice of an iDiscoid with a HHEX/LHX1/CER1 co-positive domain adjacent to a WT clusters. Dotted line indicates boundaries of the WT cluster in each image, arrow indicates co-positive domain.

(D) Immunofluorescence staining showing a TBXT/MIXL1 co-positive domain within the WT clusters of the iDiscoid.

(E) Examples of WT clusters with polarized TBXT/MIXL1 domains (top) or without co-expression of markers (bottom). Venn diagram shows proportions of cluster types observed in iDiscoid cultures, n = 746 total.

(F) Venn diagram showing the proportion of clusters with a particular TBXT/CER1 polarity type when CER1 confined to one pole is present. n = 169 total.

(G) Representative WT clusters showing TBXT and CER1 expression domains at opposing poles of the cluster (anti-polar, top) or at the same pole of the cluster (syn-polar, bottom).

(H) Diagrams indicating the average radial expression patterns of TBXT in each polarity pattern from WT clusters with a polarized CER1-expressing domain indicated in (F). All diagrams are scaled to the same expression intensity value (1 a.u.). Degrees indicate radial distance around the circularized perimeter of a WT cluster from the CER1 peak shown. Shaded areas indicate region of highest average TBXT polarity corresponding to the polarity types.

Scale bars = 100 μm. n represents separate biological samples from a 3-4 separate experiments harvested on day 4 after iDiscoid induction.

Figure 4

Figure 4: iDiscoid hematopoietic lineages and blood island-like structures

(A) Heatmap showing day 5 iDiscoid scRNA-seq populations compared to hematopoietic populations from the human E16-19 embryo.

(B) Dot plot showing the expression pattern of hematopoietic markers in day 5 iDiscoid scRNA-seq populations.

(C) Immunofluorescence image showing the distribution of cells expressing CD34 and TAL1 (scl) in iDiscoid culture. Cells expressing TAL1 localize between the yolk sac endoderm compartment and the tissue culture dish and form arrangements of spindle-shaped cells. Orthogonal slice shows the position of spindle cells against the dish. Dashed line indicates the position from which the slice was taken.

(D) Live phase image taken at day 12 showing cells spherical cells generated within iDiscoid tissue. Dotted box indicates area of inset.

(E) Immunofluorescence image of day 12 iDiscoid showing the generation of CD43$^+$ spherical cells within CD31$^+$ endothelial cells.

(F) Immunofluorescence image of a day 12 iDiscoid with 25% GATA6-high cells substituted into the normal iDiscoid ratio.

(G) Histogram showing the number of CD43$^+$ cells detected in experiments in which the given percentages of GATA6-hi cells were substituted. n[0%] = 4 experimental replicates, n[10%] = 2 experimental replicates, n[25%] = 2 experimental replicates.

(H) Two slices from a single blood island-like structure demonstrating Desmin$^+$ mesoderm cells and CD34$^+$ endothelial cells localized underneath a FOXA2$^+$ endoderm.

(I) Histogram showing the z-distribution of the indicated markers between the bottom of the dish and the top of the culture. Bimodal distribution of FOXA2 indicates areas of expression outside of blood island-like foci. Distributions are averaged from 9 foci from 3 technical replicates.

(J) 3D reconstruction of one blood island area showing positioning of endoderm (FOXA2), endothelial (CD34), and mesoderm (Desmin) markers

(K) Schematic depicting in vivo embryonic yolk sac blood islands and in vitro iDiscoid yolk sac blood island-like structures.

Scale bars = 100 μm.

Extended Data Figures

Modelling Human Post-Implantation Development via Extra-Embryonic Niche Engineering

Joshua Hislop[1-3], Amir Alavi[4,5], Qi Song[4,5], Rayna Schoenberger[1,3], Kamyar Keshavarz F.[1-3], Ryan LeGraw[2,3], Jeremy Velazquez[2,3], Tahere Mokhtari[1-3], Mohammad Nasser Taheri[1-3], Matthew Rytel[2,3], Susana M Chuva de Sousa Lopes[8], Simon Watkins[6,7], Donna Stolz[6,7], Samira Kiani[1-3,9], Berna Sozen[10], Ziv Bar-Joseph[4,5], Mo R. Ebrahimkhani[1-3,9]*

Affiliations:

1. Department of Bioengineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA 15261, USA
2. Department of Pathology, Division of Experimental Pathology, School of Medicine, University of Pittsburgh, Pittsburgh, PA 15213, USA
3. Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA 15261, USA
4. Computational Biology Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA
5. Machine Learning Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA.
6. Center for Biologic Imaging, University of Pittsburgh, Pittsburgh, PA, USA.
7. Department of Cell Biology and Molecular Physiology, University of Pittsburgh, Pittsburgh, PA, USA
8. Department of Anatomy and Embryology, Leiden University Medical Center, Einthovenweg, 2333 ZC Leiden, the Netherlands
9. McGowan Institute for Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA 15219, USA
10. Department of Genetics, Yale School of Medicine, Yale University, New Haven, CT, 06510, USA.

*Correspondence: mo.ebr@pitt.edu

Extended Data Figure 1

Middle Slice

EGFP-GATA6
ISL1
NANOG

Hoechst

EGFP-GATA6

NANOG

ISL1

A
pConst — rtTA3
dox
pTREt — GATA6 | 2A | EGFP

B

C

D

E

In Vitro

Aggregates in 3D

In Vivo

Cell Culture Media

Self-organization (3D):
Cell sorting
Cavity formation

No dorso-ventral symmetry breaking

iDiscoid is similar to a flattened yolk sac cavity with epiblast interface and amniotic cavity

In Vitro

Derivation of iDiscoid

Cell Culture Media

Cell Culture Media

Self-organization (2D->> 3D):
Cell sorting
Confinement
Migration
Cavity formation

Cell Culture Media

Cell Culture Dish

Cell Culture Dish

Yolk Sac

Bilaminar Disc

Yolk sac tissue away from epiblast

Uterine Wall

Trophoblast

■ Yolk Sac Endoderm   ■ WT/Epiblast   ■ Amniotic Ectoderm

Extended Data Figure 1: 3D iDiscoid cultures.

(A) The gene circuit used to create inducible GATA6-expressing iPSCs. pConst is a constitutively active promoter.

(B) Heterogeneity of EGFP (GATA6) activation in iGATA6 cells, detected via flow cytometry analysis. Higher gene circuit copy numbers leads to higher expression level of EGFP and GATA6.

(C) 3D culture of iGATA4 and WT showing co-expression of the amnion marker ISL1 and the pluripotency marker NANOG spread fully throughout the WT layer without D-V polarization. Middle slice shows the development of a central lumen.

(D) 3D culture of iGATA4 and WT showing expression of the pluripotency marker NANOG throughout the WT layer but lack of ISL1 expression in notable subset of these 3D tissues. These spheres do not exhibit polarization as well.

(E) Schematic depicting iDiscoid development in 3D versus from 2D>>3D in comparison to embryo morphology. iDiscoid is similar to a flattened yolk sac cavity with epiblast interface and amniotic cavity.

Extended Data Figure 2

Extended Data Figure 1: Sorting and symmetry breaking events following GATA6 induction.

(A) Immunofluorescence images of fixed cultures demonstrating cell organization of iGATA6 (green) and WT (NANOG) hiPSCs between day 0 and day 3 after GATA6 induction. PDGFRα rises within the iGATA6 cells as they acquire a more yolk sac endoderm-like morphology.

(B) Time lapse images of a single position within the iGATA6/WT co-culture from day 0 to day 3 after GATA6 induction. Top cropped images correspond to positions within the white boxes in the images below. Scale bar = 200 μm.

(C) Time lapse images showing the initial confinement of red WT cells with green iGATA6 cells, followed by migration of iGATA6 cells over the WT disc prior to D2 after iDiscoid induction.

(D) Z-slices of two representative iDiscoids showing localization of OCT4, GATA4, and SOX17 within the bilaminar disc-like area of iDiscoid culture, as well as development of a central lumen by day 5.

Extended Data Figure 3

Extended Data Figure 3: Lumen development within WT of iDiscoids

(A) Immunofluorescence images showing dynamic of LAMA1 deposition on days 1, 3, and 5 of iDiscoid development after induction. Dotted boxes show the areas of inset in the fourth panels of the day 3 and day 5 images.

(B) Timecourse graphs showing the increase in LAMA1 signal in immunofluorescence images over the first five days of iDiscoid development. Dots represent randomly sampled areas from a single experiment harvested at each day

(C) Immunofluorescence image showing the deposition of laminin around a WT cluster with a central lumen as well as polarization of PODXL .

(D) Immunofluorescence showing horizontal and lateral slices of a representative WT cluster with polarization of PODXL and ZO-1 towards a central lumen.

(E) A representative cluster of WT iPSCs at day 3. No laminin deposition is observable in the vicinity of the cluster.

(F) ELISA comparing secreted AFP and APOA1 detected on D0 and D5 after GATA6 induction with Dox. Individual dots represent biological replicates.

Scale bars = 100µm

Extended Data Figure 4

A

Average Per- WT cluster
Lumen #: 2.8821
n = 331

Lumen Number (y-axis, 0–20)
Area (µm) (x-axis, 0 to 2.5×10⁵)

64%   33.2%   39.8%

B

Lumen Number (y-axis, 0–20)
Circularity (x-axis, 0.0 to 0.8)

EGFP-GATA6 ZO-1 PODXL

C

Disc Area

iGATA6 Seeding Density (per cm²) (y-axis: 34k, 51k, 66k, 85k)
WT Seeding Density (per cm²) (x-axis: 1.3k, 1.9k, 2.5k, 3.1k, 3.8k, 5.7k, 7.5k, 9.4k)

2×10⁵ 1×10⁵ µm²

D

Disc Circularity

(y-axis: 34k, 51k, 66k, 85k)
(x-axis: 1.3k, 1.9k, 2.5k, 3.1k, 3.8k, 5.7k, 7.5k, 9.4k)

Circularity 0.3 0.2

Disc Number

(y-axis: 34k, 51k, 66k, 85k)
(x-axis: 1.3k, 1.9k, 2.5k, 3.1k, 3.8k, 5.7k, 7.5k, 9.4k)

per mm² 10 5 0

E

Average Per- WT cluster
Lumen # (Optimized): 1.064
n = 79

Lumen Number (y-axis, 0–20)
Area (µm) (x-axis, 0 to 2.5×10⁵)

74%   70.9%   89.7%

Extended Data Figure 4: Lumen development within WT of iDiscoids

(A) Distribution of WT cluster areas versus the number of lumens observed in iDiscoids with iGATA6 coverage. Shaded area indicates the areas of the bilaminar disc between E9 and E17 as recorded by Hertig and Rock, 1949 and Heuser et al. 1945; 64% of discs observed have areas that fall into this range of values, and 39.8% have a single lumen. 33.2% fall into both of these categories. n = 331 total.

(B) Distribution of WT cluster circularity versus the number of lumens observed in iDiscoids with iGATA6 coverage, with a representative image of an iDiscoid with the indicated characteristics. n = 331, Scale bar = 100μm

(C) Heatmaps displaying the average area of iDiscoids resulting from different initial seeding densities of iGATA6 and WT. Dotted box shows optimized seeding density used for iDiscoid experiments.

(D) Heatmaps displaying the average circularity and average resulting disc number pre mm^2 of iDiscoids resulting from different initial seeding densities of iGATA6 and WT. Dotted box shows optimized seeding density used for iDiscoid experiments.

(E) Distribution of WT cluster areas versus the number of lumens observed in iDiscoids with iGATA6 coverage when seeded at the optimized ratio indicated in C. Shaded area indicates the areas of the bilaminar disc between E9 and E16-19 as recorded by Hertig and Rock, 1949 and Heuser et al. 1945; 74% of discs that fall into this area range. 89.7% of discs have a single lumen. 70.9% of discs fall into both of these categories. n = 79 total.

Extended Data Figure 5: Merged clustering of D0-D5 iDiscoid single cell populations
(A) Individual UMAP projections and clustering for each time point recorded through day 5 in iDiscoid development.
(B) Violin plots showing a curated set of genes in iDiscoid day 0, day 2, and day 4 clusters.
iDiscoid clusters are ordered by lowest to highest GATA6 expression level. "W" clusters are clusters with putative wild-type lineage; "G" clusters are clusters with putative iGATA6 lineage.

Extended Data Figure 6

Extended Data Figure 6: Hypergeometric statistical comparison of iDiscoid time points to human and NHP embryo data.

(A) Hypergeometric statistical comparison of each iDiscoid day to the annotated human embryo populations from Tyser et al. 2022. Blue dots above each column indicates the relative GATA6 expression level of each population indicated per day.

(B) Hypergeometric statistical comparison of each iDiscoid day to the annotated human embryo populations from Xiang et al. 2020. Scale used is the same as shown in panel A.

(C) Hypergeometric statistical comparison of each iDiscoid day to the annotated cynomolgus embryo populations from Ma et al. 2019. Scale used is the same as shown in panel A.

iDiscoid clusters correspond to those in the individual day-by-day clustering in Extended Data Figure 5 and are ordered from left to right on the x-axis by lowest to highest GATA6 expression level. "W" clusters are clusters with putative wild-type lineage; "G" clusters are clusters with putative iGATA6 lineage. Dotted outlines indicate fate comparisons of the most interest for each cluster.

Abbreviations from other datasets: NNE = Non-neural ectoderm, DE (P) = Definitive endoderm (proliferative), DE (NP) = Definitive endoderm (not proliferative), YS = Yolk sac, PGC = Primordial germ cell, CTB = Cytotrophoblast, STB = Syncytiotrophoblast, EVT = Extravillous trophoblasts, PSA-EPI = Primitive streak anlage in the epiblast, EXMC = Extraembryonic mesoderm cells, E- = Early, L- = Late, Gast = Gastrulating cells, AM = Amnion, VE/YE = Visceral endoderm/Yolk sac endoderm

Extended Data Figure 7

Extended Data Figure 7: Merged clustering of Day 0 – Day 5 iDiscoid single cell populations

(A) UMAP showing the merged iDiscoid clusters annotated by day.

(B) UMAP showing the merged iDiscoid clusters with unsupervised clustering applied. (*repeated figures*: Cluster 14 is the amnion-like cluster shown in Figure 2B; Cluster 6 is the D2 anterior-like cluster shown in Figure 3A; Cluster 2 is the D3-5 anterior-like cluster shown in Figure 3A; and Cluster 15 is the posterior-like cluster shown in Figure 3B.)

(C) Heatmap showing the top 20 genes corresponding to each cluster of the merged day 0 through day 5 dataset.

Extended Data Figure 8: Major signaling interactions between day 5 iDiscoid cell clusters.
(A) Immunofluorescence staining for the amnion markers ISL1 and AP-2α at day 4. Top-down widefield image of a flattened coverslip. Scale Bar = 100 μm
(B) Dot plot of marker genes from the BMP pathway from day 4 iDiscoid scRNA-seq. BMP4 expression and a number of associated genes (boxed in red) are highest in the amnion-like population.

Extended Data Figure 9

Extended Data Figure 9: Effect of BMP4 inhibition on anterior-like and posterior domains in iDiscoids.

(A) Control iDiscoids showing development of TBXT⁺ posterior domains and LHX1⁺ areas expressing CER1.

(B) iDiscoids with 100ng/mL Noggin added at day 2 showing presence of LHX1⁺ areas expressing CER1, but no expression of TBXT⁺ domains. TBXT expression is seen in iGATA6-lineage cells at the periphery of the WT disc.

Scale bars = 100 μm

Extended Data Figure 10

Extended Data Figure 10: Examples of TBXT and CER1 observed in iDiscoids.

(A) A representative WT cluster showing syn-polarity of TBXT and CER1 within the WT cluster and iGATA6 layers, respectively. Filled arrow indicates CER1-expressing cells within the iGATA6 layer; empty arrow indicates CER1 and TBXT co-expressing cells within the WT layer. Z-slices are representative slices from the center of two different WT discs.

(B) Representative WT clusters showing anti-polarity of TBXT and CER1 within the WT cluster and iGATA6 layers, respectively. Filled arrow indicates CER1-expressing cells within the iGATA6 layer; empty arrow indicates CER1 and TBXT co-expressing cells within the WT layer. Z-slice is a representative slice from the center of a WT disc.

(C) Top shows a merged UMAPs of all D0-D5 scRNA-seq data showing the posterior compartment expressing the inhibitor CER1. Dotted boxes show WT lineages. Posterior region was illustrated in Figure 3B. Bottom shows a scatterplot from merged D0-D5 iDiscoid scRNA-seq showing a population of cells derived from D3-D4 iDiscoid that co-express CER1 and TBXT. Axes represent scaled expression levels of each marker; dots represent individual cells colored by day.

Scale Bars = 100 μm

Extended Data Figure 11

Marmoset (Bergmann et al 2022)

A

CS5   CS6

Human (Tyser et al 2021)

B

Cynomolgus Monkey (Cui et al 2022)

C

Mouse (Pijuan-Sala et al 2018)

D

Mouse (Mittnenzweig et al 2021)

E

Mouse (Peng et al 2016)

F

Extended Data Figure 11: Positioning of TBXT and CER1 expression in transcriptomic human, NHP and mouse datasets.

(A) Spatial expression of TBXT and CER1 from the CS5 and CS6 marmoset embryo from Bergmann et al. 2022. In marmoset CS5, CER1 has high and unpolarized expression in the extraembryonic endoderm overlaying the embryonic disc; by CS6, overall expression has decreased, and a very slight polarization away from the putative posterior is observed. From CS5 to CS6, TBXT moves from unregionalized/bipolar expression within the embryonic disc to strong expression at one pole of the embryonic disc.

(B) Regional expression of TBXT and CER1 in the E16-19 (CS7) human embryo from Tyser et al. 2022. Highest expression of TBXT and CER1 is observable in the caudal portion of the embryo, which corresponds to the area of the primitive streak and the hypoblast overlaying this area, as shown in the diagram.

(C) Pseudospatial representation of TBXT and CER1 expression in the cynomolgus E18-E20 embryo from Cui et al. 2022. Diagram to the right shows the microdissected embryo sections with labels corresponding to each identified tissue layer. CER1 expression strongly overlays TBXT expression in the lower embryonic (EM-Low) and middle embryonic (EM-Mid) regional layers.

(D) Expression levels of T and Cer1 from the E6.5-E8.5 mouse embryo. There is a pronounced co-expression within the anterior domain of the primitive streak.

(E) Expression levels of T and Cer1 from an anterior primitive streak metacell from 153 mouse embryos spanning E6.5 – E7.5. Within this representative cell/cell type, T and Cer1 rise together over time at the anterior primitive streak.

(F) Spatial expression levels of T and Cer1 within the ~E6.5-E7.5 mouse embryo. These markers are co-expressed in the area near the anterior tip of the posterior domain. Boxed area in the diagram corresponds highlights this range. Right diagrams show expression patterns of each marker at the index 3 plane in the Y axis of left diagrams.

Extended Data Figure 12: Expression of yolk sac mesoderm and extracellular matrix genes in day 4 iDiscoid scRNA-seq.

(A) Violin plots showing the distribution of ECM genes as well as the yolk sac mesoderm marker gene BST2 in day 5 iDiscoid scRNA-seq clusters.

(B) Dot plot showing a subset of day 5 yolk sac mesoderm and endothelial marker genes. The highest GATA6 population expresses the highest levels of YSM marker genes. The GATA6 expression is shown to the left with different scaling of circle sizes than the marker list to the right.

Extended Data Figure 13: Identification of endothelial/hematopoietic populations.
(A) Scatterplots showing the distribution of markers obtained via image analysis of 3 independent experiments at day 5. Percentages correspond to the fraction of cells recorded with corresponding marker expression levels above the thresholds represented by the dotted lines.
(B) Immunofluorescence image showing cells expressing hematopoietic markers in iDiscoid culture. Cells expressing the hematopoietic marker TAL1 (Scl) localize between the yolk sac endoderm compartment and the tissue culture dish. Scale bar = 100 µm
(C) Immunofluorescence image showing cells expressing RUNX1 and TAL1 positioned against the dish. Scale bar = 100 µm
(D) Image analysis of z-slices from 5 areas in 3 biological replicates. The peak of ERG expression is underneath the peak of EGFP expression representing the iGATA6 endoderm layer. Dotted curves represent SEM calculated at each point.
(E) Immunofluorescence image showing cells expressing VE-cadherin and VEGFR2 positioned against the dish. Arrow indicates likely differentiating endothelial cell from the overlaying iGATA6 layer. Scale bar = 100 µm
(F) Flow cytometry plots showing the presence of hematopoietic markers on day 14 after a CFU assay initiated CD34-expressing cells (enriched from iDiscoids).
(G) Results from image analysis demonstrating the change in area of CD34$^+$ cells between day 5 and day 12 of iDiscoid culture, assessed via analysis of immunofluorescence images. Individual dots represent biological replicates. **: P value=0.0055
(H) Representative flow cytometry plot on day 5 and day 12 of iDiscoid showing expansion of the CD34$^+$ population by day 12 (n=3).

Extended Data Figure 14

Extended Data Figure 14: Hematopoietic cells in iDiscoid.

(A) Erythroid markers (CD235a and Hemoglobin) expressed in a subset of CD43$^+$ round cells in day 12 iDiscoids. Scale bar = 100µm

(B) Cells expressing myeloid (CD31, CD33) and macrophage (CX$_3$CR1) markers in a representative cluster in day 12 iDiscoid. Scale bar = 100µm

(C) Cells expressing megakaryocyte markers in day 12 iDiscoid. Scale bar = 100µm

(D) UMAP showing unbiased clustering of scRNA-seq performed on day 21. Boxed area shows populations expressing hematopoietic marker genes.

(E) Hypergeometric statistical comparison of day 21 iDiscoid populations to the E16-19 human embryo populations from Tyser et al. 2022. The red box shows populations of interest with similarity to in vivo hematopoietic lineages. iDiscoid clusters are grouped by similar fates on Y axis.

(F) Expression of hematopoietic marker genes within the red boxed population in panel D. The four clusters show distinct marker gene expression profiles associated with yolk sac-derived hematopoietic populations.

(G) Flow cytometry analysis on day 21 iDiscoid. Cells were gated for positive CD43 expression prior to gating for the markers shown (n=2, representative sample shown).

0% iGATA6-hi Supplementation

25% iGATA6-hi Supplementation

Extended Data Figure 15: Comparison of iDiscoids with and without supplementation with high iGATA6 expressing (iGATA6-hi) cells.

(A) Expression of CD43+ and Hemoglobin+ cells in day 12 iDiscoid without additional supplementation of GATA6-hi cells. Red color in high Hoechst areas is nonspecific staining. Scale bar = 500 μm. Insets show areas of CD43+ cells. Dotted outlines on right show zoomed in areas shown within the image. Scale bar = 100 μm

(B) Expression of CD43+ and Hemoglobin+ cells in day 12 iDiscoid with supplementation of 25% GATA6-hi cells at initial seeding. A significant expansion of CD43-expressing cells is observable. Scale bar = 500 μm. A representative image of 2-4 biological replicates.

Extended Data Figure 16: Structure of blood island-like associated tissues within iDiscoid.

(A) Z-slices of a day 12 iDiscoid at the indicated height above the bottom focal plane. Expression of the mesodermal marker Desmin is regionalized close to the bottom of the tissue; endothelial marker CD34 is expressed near the bottom and middle of the tissue, with highest expression just above Desmin (z = 6-8); endoderm marker FOXA2 is exclusively expressed above the other tissues. Scale bar = 50 μm

(B) Z-slices around the blood island-like area shown in Figure 4E. Desmin is observed regionalized below the endothelial (CD31[+]) and hematopoietic (CD43[+]) tissues. No stain for endoderm is shown. Scale bar = 100 μm

Extended Data Figure 17

A

Extended Data Figure 17: iDiscoid formation from hiPSCs to model human early post-implantation development *in vitro*.

(A) From an initially mixed state, iDiscoid cells segregate into WT clusters surrounded by iGATA6 cells. These iGATA6 cells migrate laterally to create a bilaminar boundary on top of the WT clusters. These clusters then undergo lumenogenesis, specification of amnion-like cells, and formation of anterior hypoblast-like and posterior domains. In iGATA6-only areas, yolk sac mesoderm specifies and hematopoietic cell specification is observed.

Extended Data Figure 18

Extended Data Figure 18: Investigation of WT fates in day 12 iDiscoid.

(A) Most structures corresponding to former WT clusters at day 12 in iDiscoid have taken on expression of ISL1 and lost expression of the pluripotency markers SOX2 and NANOG. A limited number of cells express SOX2 at the core of former WT clusters, potentially indicating the specification of an ectoderm-like fate in a small number of WT-lineage cells.

(B) A subset of former WT clusters have taken on markers of ectoderm differentiation.

Scale bars = 500 µm

Extended Data Figure 19

A

iGATA6
(Uninduced/
Preprogrammed)   Wild-Type

Parental Cell
Population

Expand/Passage
Freeze   Seed in 2D

+Dox

High-Throughput
System   iDiscoid (3D)

Easy to Use
Reproducible

Developmental Toxicology/Screening
Disease Modeling
Study Human Development

B GFP-GATA6 OCT4 Day 5

Post-cryostorage
WT Cluster # = 601
Avg area = 30630 μm^2 ± 1044
Avg Distance from Nearest = 6085 μm^2 ± 102

C GFP-GATA6 OCT4 Day 5

Passage = 1
WT Cluster # = 415
Avg cluster area = 32540 μm^2 ± 1176
Avg Distance from Nearest = 5765 μm^2 ± 115

GFP-GATA6 OCT4 Day 5

Passage = 15
WT Cluster # = 438
Avg area = 26358 μm^2 ± 855
Avg Distance from Nearest = 5291 μm^2 ± 106

Extended Data Figure 19: Mixed iDiscoid cell lines exhibit consistent cluster formation after repeated passaging and cryostorage prior to induction.

(A) Schematic showing the creation of the iDiscoid parental cell line. iGATA6 cells with heterogeneous copy numbers of the inducible GATA6 circuit are mixed with wild-type. This cell combination is then maintained together or frozen prior to induction for iDiscoid experiments.

(B) iDiscoid morphology and characteristics following cryostorage and defrosting. Scale Bar = 500 µm

(C) Characteristics and morphology of iDiscoid cultures induced immediately after mixing (passage 1) or maintained together and passaged for two months (passage 15). Scale Bars = 500 µm

Extended Data Figure 20

Anti-polar　　**Syn-polar**

Extended Data Figure 20: iDiscoid engineering in a separate iPSC cell line results in the generation all major features of interest

Day 4 immunofluorescence images of iDiscoid engineered using the iPSC cell line PGP9.

(A) A portion of PGP9 iGATA6 cells expressing high levels of GATA6 (EGFP) also express the anterior endoderm marker HHEX near the edge of a WT cluster.

(B) A ring of PODXL expression lines the inside of a cavities formed in PGP9 WT clusters.

(C) ISL1[+] cells specify away from NANOG[+] cells, along the base of a cavity formed in PGP9 WT.

(D) TBXT[+] cells develop polar domains at the edge of WT clusters. Representative islands show that both syn- and anti-polar arrangement of cells are observable.

Scale Bars = 100 μm

Extended Data Figure 21

Extended Data Figure 21: Example of FACS gating strategy.
(Cited in Methods) Cell debris was excluded via an SSC-A vs FSC-A gate; aggregates were excluded by comparing FSC-A and FSC-H; dead cells were gated out using the 7-AAD stain to identify positive cells.

Modelling Human Post-Implantation Development via Extra-Embryonic Niche Engineering

Joshua Hislop[1-3], Amir Alavi[4,5], Qi Song[4,5], Rayna Schoenberger[1,3], Kamyar Keshavarz F.[1-3], Ryan LeGraw[2,3], Jeremy Velazquez[2,3], Tahere Mokhtari[1-3], Mohammad Nasser Taheri[1-3], Matthew Rytel[2,3], Susana M Chuva de Sousa Lopes[8], Simon Watkins[6,7], Donna Stolz[6,7], Samira Kiani[1-3,9], Berna Sozen[10], Ziv Bar-Joseph[4,5], Mo R. Ebrahimkhani[1-3,9]*

Affiliations:

1. Department of Bioengineering, Swanson School of Engineering, University of Pittsburgh, Pittsburgh, PA 15261, USA
2. Department of Pathology, Division of Experimental Pathology, School of Medicine, University of Pittsburgh, Pittsburgh, PA 15213, USA
3. Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA 15261, USA
4. Computational Biology Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA
5. Machine Learning Department, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, USA.
6. Center for Biologic Imaging, University of Pittsburgh, Pittsburgh, PA, USA.
7. Department of Cell Biology and Molecular Physiology, University of Pittsburgh, Pittsburgh, PA, USA
8. Department of Anatomy and Embryology, Leiden University Medical Center, Einthovenweg, 2333 ZC Leiden, the Netherlands
9. McGowan Institute for Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA 15219, USA
10. Department of Genetics, Yale School of Medicine, Yale University, New Haven, CT, 06510, USA.

*Correspondence: mo.ebr@pitt.edu

Supplementary Video 1: Self-organization of hiPSC cells that result in WT discs surrounded by iGATA6 cells from day 0 to day 3

Supplementary Video 2: Migration of iGATA6 cells over the top of the WT cells to form a contiguous membrane
iGATA6 cells (green) cover the top of the WT disc (red) between day 0.5 and day 2.

Supplementary Video 3: Expansion of a lumen from a rosette in a single iDiscoid.

Supplementary Video 4: Fly-through of an iDiscoid with a single lumen
Z-slices of a single fixed iDiscoid moved through from end-to-end.
DAPI: blue, iGATA6: Green, PODXL: Red, F-actin: Gray