Short Title: Role of BST4 in Chlamydomonas

Title: The role of BST4 in the pyrenoid of *Chlamydomonas reinhardtii*

Authors: Liat Adler[1,2,3], Chun Sing Lau[4], Kashif M. Shaikh[5], Kim A. van Maldegem[5],
Alex L. Payne-Dwyer[4,6], Cecile Lefoulon[7], Philipp Girr[4], Nicky Atkinson[1,2], James
Barrett[4], Tom Z. Emrich-Mills[4], Emilija Dukic[5], Michael R. Blatt[7], Mark C. Leake[4,6],
Gilles Peltier[8], Cornelia Spetea[†5], Adrien Burlacot[†,3,9], Alistair J. McCormick[†,1,2], Luke
C. M. Mackinder[†4], Charlotte E. Walker[†4]

[1]Institute of Molecular Plant Sciences, School of Biological Sciences, University of
Edinburgh, EH9 3BF, United Kingdom.
[2]Centre for Engineering Biology, University of Edinburgh, EH9 3BF, United Kingdom.
[3]Department of Plant Biology, The Carnegie Institution for Science, Stanford, CA,
94305 USA
[4]Centre for Novel Agricultural Products (CNAP), Department of Biology, University of
York, Heslington, York YO10 5DD, United Kingdom.
[5]Department of Biological and Environmental Sciences, University of Gothenburg,
Gothenburg 40530, Sweden.
[6]School of Physics, Engineering and Technology, University of York, Heslington, York
YO10 5DD, United Kingdom.
[7]Laboratory of Plant Physiology and Biophysics, Bower Building, University of
Glasgow, Glasgow, United Kingdom.
[8]Aix-Marseille Université, CEA, CNRS, Institut de Biosciences et Biotechnologies Aix-
Marseille, CEA Cadarache, 13108 Saint-Paul-lez-Durance, France
[9]Department of Biology, Stanford University, Stanford, CA, 94305, USA

[†]The author(s) responsible for distribution of materials integral to the findings
presented in this article in accordance with the policy described in the Instructions for
Authors (https://academic.oup.com/plcell/pages/General-Instructions) are:

• Cornelia Spetea, Department of Biological and Environmental Sciences
University of Gothenburg, Gothenburg 40530, Sweden +46-31 786 9332,
cornelia.spetea.wiklund@bioenv.gu.se

- Adrien Burlacot, Carnegie Institution for Science, Stanford, CA, 94305 USA, aburlacot@carnegiescience.edu
- Alistair McCormick, School of Biological Sciences, University of Edinburgh, +44 1316 505316, alistair.mccormick@ed.ac.uk
- Luke Mackinder, Department of Biology, University of York, +44 1904 328984, luke.mackinder@york.ac.uk
- Charlotte Walker, Department of Biology, University of York, +44 1904 328500, charlotte.walker@york.ac.uk

One-sentence summary: In Chlamydomonas, the pyrenoid-localized bestrophin-like protein BST4 is a putative ion channel involved in pH regulation of the thylakoid lumen, possibly by mediating bicarbonate transport.

Funding information: This work was funded by the Biotechnology and Biological Sciences Research Council (BBSRC) (grant number (BB/S015531/1) and Leverhulme Trust (RPG-2017-402). LA acknowledges funding support by EASTBIO DTP (BB/J01446X/1) and the Carnegie Institution for Science. CEW acknowledges funding support from BBSRC Discovery Fellowship (BB/W009587/1). CSL was supported by Bill & Melinda Gates Agricultural Innovations (Investment ID 53197). CS acknowledges grants from the Swedish Research Council (VR 2016-03836 and 2021-03790). KMS was recipient of a postdoctoral fellowship from the Carl Tryggers Foundation (CTS 20:406). A United Kingdom Research and Innovation Future Leaders Fellowship to LCMM (MR/T020679/1), Biotechnology and Biological Sciences Research Council grants (BB/T017589/1, BB/S015337/1, BB/R001014/1) and Engineering and Physical Research Council Grant (EP/W024063/1) with ML and APD. AB acknowledges the support of the Carnegie Institution for Science. The authors would like to thank the University of York Biosciences Technology Facility for confocal microscopy access and support. CL was funded by BBSRC grants BB/P011586/1 and BB/T013508/1 to MRB. PG was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 456013262.

ABSTRACT

In many eukaryotic algae, CO_2 fixation by Rubisco is enhanced by a CO_2-concentrating mechanism, which utilizes a Rubisco-rich organelle called the pyrenoid.

The pyrenoid is traversed by a network of thylakoid-membranes called pyrenoid
tubules, proposed to deliver CO_2. In the model alga *Chlamydomonas reinhardtii*
(**Chlamydomonas**), the pyrenoid tubules have been proposed to be tethered to the
Rubisco matrix by a bestrophin-like transmembrane protein, BST4. Here, we show
that BST4 forms a complex that localizes to the pyrenoid tubules. A Chlamydomonas
mutant impaired in the accumulation of BST4 (*bst4*) formed normal pyrenoid tubules
and heterologous expression of BST4 in *Arabidopsis thaliana* did not lead to the
incorporation of thylakoids into a reconstituted Rubisco condensate. Chlamydomonas
bst4 mutant did not show impaired growth at air level CO_2. By quantifying the non-
photochemical quenching (**NPQ**) of chlorophyll fluorescence, we show that *bst4*
displays a transiently lower thylakoid lumenal pH during dark to light transition
compared to control strains. When acclimated to high light, *bst4* had sustained higher
NPQ and elevated levels of light-induced H_2O_2 production. We conclude that BST4 is
not a tethering protein, but rather is an ion channel involved in lumenal pH regulation
possibly by mediating bicarbonate transport across the pyrenoid tubules.

INTRODUCTION
Maintaining improvement in crop yields to keep pace with the rising demands for food
is becoming increasingly challenging (Horton et al., 2021). Current models predict that
an increase in food supply between 35 and 56% from 2010 to 2050 is required (van
Dijk et al., 2021). A possible solution to overcome this challenge is engineering a
biophysical CO_2-concentrating mechanism (**CCM**) into C3 crop plants, which has been
proposed to improve crop yields by between 8 and 60%, as well as water-use and
nitrogen-use efficiency (Price et al., 2013; McGrath and Long, 2014; Long et al., 2019;
Fei et al., 2022; Wu et al., 2023). The biophysical CCMs in algae typically function by
concentrating CO_2 into a liquid-liquid phase separated microcompartment called a
pyrenoid, which is predominantly made up of a Ribulose-1,5-bisphosphate
carboxylase/oxygenase (**Rubisco)**-rich matrix. This raises the $[CO_2]:[O_2]$ ratio around
the primary CO_2-fixing enzyme Rubisco, which brings Rubisco closer to its maximal
carboxylation rate and minimizes the competing oxygenation reaction.
Chlamydomonas reinhardtii (hereafter **Chlamydomonas**) has the most well
understood pyrenoid-based CCM and has become the blueprint for engineering such
a CCM into C3 plants (Hennacy and Jonikas, 2020; Adler et al., 2022).

An important yet little understood aspect of the Chlamydomonas CCM is the function
and biogenesis of the thylakoid tubule network that traverses the pyrenoid, known as
the pyrenoid tubules. The pyrenoid tubules are continuous with the thylakoid
membrane (which harbors the photosynthetic electron transport chain) (Engel et al.,
2015) and are thought to function as a delivery system for inorganic carbon (**Ci**) to the
Rubisco-rich pyrenoid matrix (Mitra et al., 2005; Raven 1997). In the current model,
bicarbonate (**HCO$_3^-$**) is channelled into the thylakoid lumen by bestrophin-like proteins
1-3 (**BST1-3**) (Mukherjee et al., 2019) and diffuses to the pyrenoid tubules where it is
converted to CO_2 by carbonic anhydrase 3 (**CAH3**) (Karlsson et al., 1998; Mitra et al.,
2005) thanks to a low lumenal pH generated by the photosynthetic electron transport
chain (Burlacot et al., 2022). Traversion of the pyrenoid Rubisco-matrix by tubules is
predicted to be essential for an efficient CCM (Fei et al., 2022). Therefore,
understanding the mechanisms of pyrenoid tubule formation and function will be
crucial for future plant pyrenoid engineering efforts.

The protein **BST4** (bestrophin-like protein 4, also known as Rubisco binding
membrane protein 1, RBMP1; Cre06.g261750) localizes exclusively to the pyrenoid
tubules and has been proposed to function as a tether protein, linking the Rubisco
matrix to the tubules (Meyer et al., 2020). BST4 is a predicted transmembrane protein
and has two Rubisco binding motifs (**RBMs**) on its long, disordered C-terminus (**Fig.**
1A) (He et al., 2020; Meyer et al., 2020). RBMs facilitate the targeting of proteins to
the pyrenoid and are also hypothesized to underpin the assembly of the pyrenoid.
Meyer et al. (2020) proposed that BST4, together with other tether proteins, may
recruit Rubisco to the tubule network.

BST4 also has a well-conserved bestrophin domain similar to those present in the
thylakoid-localized BST1-3 proteins (Mukherjee et al. 2019). Bestrophins primarily act
as anion channels and are found in a wide diversity of organisms, including animals,
plants and fungi. They are best known to have permeability to chloride and HCO_3^- (Qu
and Hartzell, 2008), although some are reportedly permeable to cations (Yang et al.,
2014) and larger organic anions (Roberts et al., 2011). It is possible that BST4
functions as an HCO_3^- channel, like that proposed for BST1-3 (Mukherjee et al. 2019),
but there are currently no data to support this hypothesis.

In this study we aimed to elucidate the role of BST4 in the Chlamydomonas CCM. We
tested BST4 as a thylakoid-Rubisco tethering protein as well as its suitability in
promoting a thylakoid-Rubisco matrix interface in the model land plant Arabidopsis.
We also explored an alternative hypothesis that BST4 functions as an ion channel in
the pyrenoid tubules and discussed the implications this has for our current
understanding of CO_2 fixation by the pyrenoid.

RESULTS AND DISCUSSION
BST4 is a bestrophin-like protein that is localized in the intra-pyrenoid thylakoid
tubules in Chlamydomonas
The amino acid sequence of BST4 has two key unique features compared to the
previously characterized BST1-3 (**Fig. 1**) (Mukherjee et al. 2019). First, BST4 has an
extended disordered C-terminus that contains two RBMs (**Fig. 1A**) (He et al., 2020;
Meyer et al., 2020). Second, BST4 has a phenylalanine residue in the first position of
the putative selection pore, as opposed to valine which is conserved throughout BST1-
3 (Mukherjee et al. 2019), although the residue at this position is variable in other well
characterized bestrophin proteins (*Malus domestica* voltage-dependent Cl⁻ channel 1
(***Md*VCCN1**) and *Homo sapiens Bestrophin 1* (***Hs*BEST1**)) (**Fig.1B, C**). In
corroboration, we found the evolutionary history of BST4 diverges from BST1-3 when
investigated using maximum likelihood phylogenetic analysis. We analysed the full-
length sequence (**Fig. S1**) and a truncated version without the disordered C-terminal,
leaving the bestrophin domain and N-terminal (**Fig. 1D**), both of which found BST4 to
resolve in a distinct clade from BST1-3 but within the wider green algae group.

Figure 1. BST4 is a bestrophin protein that is distinct from BST1-3. A. Schematic of the topology of BST4. BST4 is predicted to have four transmembrane domains and a disordered C-terminus containing two Rubisco Binding Motifs (RBMs) **B.** AlphaFold v2 models of two BST4 bestrophin domains (amino acid residues 53-386 shown for clarity) to show a typical bestrophin channel pore. Pore lining residues are highlighted in red. **C.** AlphaFold v2 structure of the BST4 bestrophin domains alongside the predicted structure of another bestrophin-like protein from *Chlamydomonas reinhardtii* (*Cr*BST1; Alphafold amino acids 51-end shown for clarity), and experimentally determined structures of bestrophins *Malus domestica* voltage-dependent Cl⁻ channel 1 (*Md*VCCN1;7EK1) and *Homo sapiens* Bestrophin 1 (*Hs*BEST1; 8D1I). Residues known to line the channel pore are highlighted. **D.** Phylogenetic analysis of the BST4 bestrophin domain (bold) with the disordered C-terminal removed. The alignment used was trimmed at residue 369. The evolutionary history of BST4 was inferred by using the maximum likelihood method based on the Le and Gascuel substitution model with discrete Gamma distribution (5 categories) and 500 bootstrap replicates. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

As well as being distinct at a sequence level, BST4 also localizes differently from BST1-3 (**Fig. 2**). While BST1-3 localize throughout the thylakoid membrane and are enriched at the pyrenoid periphery (Mukherjee et al., 2019), BST4 localizes to the centre of the pyrenoid in a pattern that resembles the pyrenoid thylakoid tubule system (**Fig. 2A, B**; Meyer et al., 2020). To confirm tubule localization, we generated a dual-tagged line that expressed BST4-mScarlet-I and the Chlamydomonas Rubisco small subunit 1 (*Cr***RBCS1**) fused to Venus. BST4-mScarlet-I was enriched where

*Cr*RBCS1-Venus was depleted (**Fig. 2C, D**), indicating that BST4 is located in the tubules and not the Rubisco-enriched pyrenoid matrix. Previous work suggests that the C-terminal RBMs of BST4 enable the protein to interact with *Cr*RBCS1 (Meyer et al. 2020). We confirmed using a yeast-2-hybrid approach that the C-terminus of BST4 interacts with *Cr*RBCS1 (**Fig. S2A**). We also measured the efficiency of Förster resonance energy transfer (**FRET**) from Venus to mScarlet-I and found that the FRET efficiency was ~35%, supporting the proximity of BST4 to *Cr*RBCS1 (**Fig. S2B**).

Bestrophins typically form pentameric assemblies (Bharill et al., 2014; Hagino et al., 2022). When five chains of BST4 were inputted to AlphaFold a typical bestrophin pentameric structure was predicted (**Fig. 2E**). To test the complex assembly of BST4 *in vivo*, we utilized a Slimfield microscopy molecular tracking method (Plank et al., 2009). For this method, it is important to have only fluorescently tagged BST4 molecules and no native (untagged) BST4 molecules. To find a strain that was impaired in the accumulation of BST4 protein, we screened *bst4* insertional mutants from the CLiP library (Zhang et al., 2014; Li et al., 2019) and confirmed a mutant strain (***bst4***) (**Fig. S3** and methods). We then expressed BST4-mScarlet-I in the *bst4* mutant background (**Fig. S3**). We used Slimfield microscopy to image BST4 in the pyrenoid tubules and subsequent image analysis (detailed in methods) to track individual fluorescent molecules of BST4-mScarlet-I and quantified the number of BST4 monomers per complex. The resulting probability distribution revealed that the most common BST4 complex is made up of five molecules (**Fig. 2F**). Other peaks showed complexes with numbers of molecules divisible by five, which may be multiple pentameric channels grouping together. The interval between the probability peaks was also five (**Fig. 2G**). To further support higher order complex assembly of BST4 we ran purified Chlamydomonas thylakoid membranes on a Blue Native-PAGE gel (**BN-PAGE**) and immunodetected BST4 (**Fig. 2H**). BST4 formed a smear at ~1000 kDa, which is considerably larger than a pentamer (~330kDa). This could be due to higher order assemblies of BST4, pentameric BST4 in a complex with other proteins and/or aberrant migration during BN-PAGE due to influences by complex shape. Collectively, our data support *in vivo* higher order assembly of BST4 potentially as a pentamer.

Figure 2. BST4 forms higher order assemblies in the pyrenoid tubules in Chlamydomonas. A. Diagram of a Chlamydomonas cell with the pyrenoid Rubisco matrix and pyrenoid tubules indicated. **B.** Confocal image of a Chlamydomonas cell expressing BST4-Venus. Scale bar is 2 μm. **C.** Pyrenoid in dual-tagged Chlamydomonas. BST4-mScarlet-I and RBCS-Venus are shown in green and magenta, respectively. Overlap appears white. The yellow line shows the 1D cross-section used for generating the line plot in (D). Scale bar is 1 μm. **D**. Plot of normalized fluorescence intensity values from a 1D cross-section from (C). mScarlet-I and RBCS-Venus are shown in green and magenta, respectively. **E**. AlphaFold Multimer v2 prediction of pentameric BST4 structure. Top structure includes the disordered C-terminus, bottom structure displays amino acid residues 53-386 only for clarity. **F**. BST4-mScarlet-I stoichiometry probability distribution based on single particle tracking and molecular counting in live Chlamydomonas (n=347 tracks) using Slimfield microscopy. **G**. Averaging the intervals between peaks in BST4-mScarlet-I stoichiometry (n=22 intervals) indicates a consistent pentameric unit. **H**. Immunoblot of proteins from *bst4* mutant and wild type (WT) Chlamydomonas thylakoids separated by Blue Native-PAGE.

BST4 localizes to the stroma lamellae thylakoid membrane in Arabidopsis

We used *Arabidopsis thaliana* (**Arabidopsis** hereafter) as a heterologous system to examine if BST4 was able to tether the Rubisco matrix to thylakoid membranes. In previous work, an Arabidopsis Rubisco small subunit (*At*RBCS) double mutant (*1a3b*) was complemented with *Cr*RBCS2, resulting in a line with hybrid Rubisco representing

~50% of the Rubisco pool (**S2$_{Cr}$**) (Izumi et al., 2012; Atkinson et al., 2017). Subsequent
expression of the pyrenoid linker protein Essential PYrenoid Component 1 (**EPYC1**),
in S2$_{Cr}$ resulted in the formation of an EPYC1-hybrid Rubisco condensate or 'proto-
pyrenoid' (Atkinson et al., 2020). The S2$_{Cr}$ line was therefore used as a platform to test
whether BST4 acts as a tether protein.

We initially localized a BST4-mNeon fusion protein following stable expression in S2$_{Cr}$
(**Fig. 3**). BST4-mNeon was observed in chloroplasts, demonstrating that the native
chloroplast signal peptide was compatible with the chloroplast targeting mechanism in
land plants (**Fig. 3A**), as seen previously for other Chlamydomonas proteins (Atkinson
et al., 2016). The fluorescence signal from BST4-mNeon had a sponge-like pattern
that was inversely correlated with the punctate chlorophyll autofluorescence signal that
represents the grana stacks (**Fig. 3B, C**). The sponge-like pattern was similar to
previous observations of autofluorescence originating from Photosystem (**PS**) I
(Hasegawa et al., 2010), which is enriched in the stroma lamellae of thylakoids. We
subsequently generated a stable Arabidopsis transgenic line expressing untagged
BST4 to confirm its location by biochemical fractionation. BST4 was detected in the
thylakoid fraction and not in the stromal fraction (**Fig. 3D**). The thylakoids were then
further fractionated into grana stacks and stroma lamellae sub-fractions. BST4 was
found in the stroma lamellae fraction (**Fig. 3E**), which is consistent with the observed
sponge-like fluorescence pattern.

Figure 3. BST4 assembles as a complex in the stroma lamellae of thylakoids in Arabidopsis. A. Confocal
image of WT protoplasts expressing BST4-mNeon. Scale bar is 5 µm. **B.** Mesophyll chloroplast from S2$_{Cr}$
Arabidopsis expressing BST4-mNeon. BST4mNeon and chlorophyll autofluorescence are shown in green and
magenta, respectively. Overlap appears white. Scale bar is 2 µm. Yellow lines indicate selections for profile plot in

(C). **C.** Plot of normalized fluorescence intensity values from a 1D cross-section (yellow line) through two grana
stacks from (B). mNeon and chlorophyll autofluorescence are shown in green and magenta, respectively. **D.**
Immunoblots of sub-chloroplast fractions isolated from Arabidopsis line S2$_{Cr}$ expressing BST4. RbcL and CP43
were probed for as stromal and thylakoid controls, respectively. **E.** Immunoblots of fractionation thylakoids from
Arabidopsis line S2$_{Cr}$ expressing BST4. CP43 and PsaB were used for grana stack and stroma lamellae controls,
respectively **F.** Trypsin protease protection assay. Intact thylakoids containing BST4 subjected to 0 or 100 μg/ml
trypsin with or without the addition of 1% (v/v) Triton. AtpB and PsbO used as controls for stromal facing (exposed)
and lumen facing (protected), respectively. **G.** Immunoblot of proteins from thylakoids separated by Blue Native-
PAGE from either BST4 stable line or S2$_{Cr}$ background. Abbreviations: Clp, whole chloroplast; CP43, CP43-like
chlorophyll binding protein; RbcL, Rubisco large subunit; PsbO, photosystem II manganese-stabilizing
polypeptide;AtpB, ATP synthase subunit beta; PsaB, photosystem I P700 chlorophyll a apoprotein A2; Str, stromal
fraction; Thy, thylakoid fraction.

As BST4 has two RBMs on its C-terminus, we hypothesized that BST4 should be
orientated with the C-terminus facing the stroma so that the RBMs are available to
interact with *Cr*RBCS. To determine the orientation of BST4, we performed a protease
protection assay on thylakoids isolated from the untagged BST4 transgenic plants.
Our antibody was raised against the C-terminal end of BST4, so it could be used to
assess whether the C-terminus was exposed for degradation in the stroma or if it was
protected in the lumen. We found that the BST4 C-terminus was fully degraded after
a 60 min treatment of trypsin, indicating that it faced the stroma (**Fig. 3F**). There was
some degradation of the lumenal control, the PSII subunit PsbO, which we attribute to
a portion of the thylakoid membrane preparation not being fully intact. However, PsbO
was fully degraded when the membranes were solubilized, indicating that they were
sufficiently intact to differentiate between lumenal- and stromal-facing peptides.
Therefore, BST4 was observed in the expected location and orientation in plant
thylakoid membranes.

Finally, we also tested whether BST4 forms a complex in the Arabidopsis thylakoid
membrane. We subjected thylakoids from the untagged BST4 transgenic plants to BN-
PAGE and detected a single band of ~850 kDa (**Fig. 3G**). Thus, BST4 forms a similar
high order complex in Arabidopsis to that in Chlamydomonas but may be lacking
additional interaction partners present in Chlamydomonas.

BST4 is not sufficient for integration of thylakoids into the Arabidopsis proto-
pyrenoid
To test whether BST4 could facilitate incorporation of thylakoids into hybrid Rubisco
condensates (i.e., proto-pyrenoids) in plants, BST4-mCherry and EPYC1-tGFP were
co-expressed in the S2$_{Cr}$ background (**Fig. 4**). When we expressed EPYC1 alone,
Rubisco condensates formed in chloroplasts as previously described (Atkinson et al.,

2020), and were visible as a ~2 µm wide puncta in the tGFP channel (**Fig. 4A**). When

BST4-mCherry and EPYC1-tGFP were co-expressed, approximately 60% of the

BST4-mCherry fluorescence signal was observed in the condensate region (**Fig. 4B,**

D), with the remaining signal exhibiting the same sponge-like pattern as seen in **Fig.**

3A.

To confirm that the observed co-localization was due to an interaction between BST4

and CrRBCS2, we mutated the first two residues of each core RBM motif of BST4 to

glutamic acid (WR to EE, the new version noted BST4$_{WR/EE}$). Previously, these

substitutions have been reported to disrupt the binding interface between EPYC1 and

CrRBCS2 (He et al., 2020). Using yeast-2-hybrid we confirmed that the interaction

between the C-terminus of BST4 and CrRBCS2 was disrupted by these mutations

(**Fig. S3**). When BST4$_{WR/EE}$-mCherry was expressed with EPYC1-tGFP in S2$_{Cr}$, no

enrichment of the mCherry signal was observed in the Rubisco condensate (**Fig. 4C,**

D). Thus, the RBMs of BST4 were responsible for the enrichment of BST4 in the

vicinity of the Rubisco condensate in Arabidopsis.

However, when condensates were sedimented and analysed by immunoblotting,

BST4-mCherry was not detected in the condensate fraction (**Fig. 4E**). When leaf

samples were subjected to formaldehyde cross-linking prior to sedimentation of the

condensate, a small amount of BST4 was present in the condensate fraction. We

concluded that BST4 was not present in the condensate itself but preferentially

occupied the thylakoid membranes surrounding the CrRBCS2-enriched condensate

(Atkinson et al., 2020), likely due to interactions with CrRBCS2.

Figure 4. BST4 is not sufficient to enable the inclusion of thylakoid membranes in a Rubisco condensate in Arabidopsis and is not required for pyrenoid tubule inclusion in the Chlamydomonas pyrenoid. A. Confocal images showing EPYC1-tGFP in S2$_{Cr}$ Arabidopsis background. Scale bar is 10 µm. **B.** Confocal images showing BST4-mCherry co-expressed with EPYC1-tGFP in S2$_{Cr}$ Arabidopsis background. Scale bar is 10 µm. **C.** Confocal images showing BST4 with mutated RBMs (BST4$_{(WR/EE)}$) fused to mCherry co-expressed with EPYC1-tGFP in the S2$_{Cr}$ Arabidopsis background. **D.** Ratio of mean mCherry fluorescence associated with the Rubisco condensate compares do the rest of chloroplast when mCherry was fused to either BST4 or BST4$_{(WR/EE)}$ (n=22-23). **E.** Immunoblot of sedimented Rubisco condensates from S2$_{Cr}$ Arabidopsis expressing EPYC1-tGFP and BST4-mCherry. Abbreviations: I, input; S, supernatant; P, pellet. Rubisco condensates are enriched in the pelleted fraction. Cross-linked samples were prepared by vacuum infiltrating intact leaves with 1% (v/v) formaldehyde prior to sedimentation. **F.** Transmission electron micrograph of a chloroplast from S2$_{Cr}$ Arabidopsis expressing either EPYC1 alone or EPYC1-tGFP and BST4-mCherry. Rc is the Rubisco condensate. Scale bar is 1 µm. **G.** Transmission electron micrograph of pyrenoid from WT Chlamydomonas or *bst4* mutant. Abbreviations: pt, pyrenoid tubules; m, matrix; s, starch sheath. Scale bar is 250 nm.

Although BST4 partially co-localized with the Rubisco condensate, we found no evidence to suggest that BST4 could facilitate the inclusion of thylakoid membranes. Confocal microscopy was not sufficient to determine if chlorophyll autofluorescence was in the proto-pyrenoids of S2$_{Cr}$ lines expressing BST4-mCherry and EPYC1-tGFP. However, transmission electron microscopy (**TEM**) revealed no visible indication of thylakoid membranes in the condensates, which were structurally similar to

325   condensates in S2$_{Cr}$-EPYC1 lines lacking BST4 (**Fig. 4F**). Thus, BST4 appeared

326   insufficient to enable thylakoid inclusion in the Rubisco condensate in Arabidopsis.

BST4 is not necessary for tubule formation in Chlamydomonas

To investigate whether BST4 was necessary for normal formation of the thylakoid

tubule-structure in Chlamydomonas we compared the structure of pyrenoids of *bst4*

compared to the WT control strain (CMJ030) (Zhang et al., 2014; Li et al., 2019). TEM

images showed that pyrenoids from *bst4* were structurally comparable to the WT

control, including the presence of pyrenoid tubules (**Fig. 4G**). There were also no

differences in pyrenoid size or shape between the two lines when comparing pyrenoids

from 40-50 cells from each genotype (**Fig. S4**). As a result, we conclude that BST4 is

not necessary for the pyrenoid tubule-Rubisco matrix interface in Chlamydomonas.

The C-terminus of BST4 is required for localization to the pyrenoid tubules

Multiple copies of the RBM are sufficient to target proteins to the pyrenoid (Meyer et

al., 2020). Furthermore, we have shown that the BST4 RBMs are required for RBCS

interaction via yeast-2-hybrid (**Fig. S2A)** and that they are required for enrichment with

the proto-pyrenoid in Arabidopsis (**Fig. 4B-D**). To investigate the role of the RBMs in

BST4 localization in Chlamydomonas, we generated a truncated version of BST4

(residues 1 to 386) that lacked the C-terminus containing the two RBMs (BST4$_{\Delta C\text{-term}}$)

and compared localization with full-length BST4 expressed in WT and the *bst4* mutant

(**Fig. 5 and Fig. S5**). BST4$_{\Delta C\text{-term}}$-mScarlet expressed in the *bst4* mutant line did not

localize to the pyrenoid tubules but was found throughout the thylakoid membrane

(**Fig. 5**). This is consistent with our findings in Arabidopsis with BST4$_{WR/EE}$ (**Fig. 4C**),

demonstrating that localization of BST4 is driven via RBM-Rubisco interaction. Thus,

the C-terminus is necessary for BST4 localization to the pyrenoid tubules in the

presence of the matrix. Collectively these data indicate that BST4 may require a pre-

existing pyrenoid tubule network to be localized in the pyrenoid rather than driving the

inclusion of thylakoid membranes into the Rubisco matrix or is redundant as a tether

protein.

When BST4$_{\Delta C\text{-term}}$-Venus was expressed in WT Chlamydomonas (i.e., that still

produced non-truncated BST4), we observed fluorescence throughout the thylakoids,

but with the majority of the signal still localized to the pyrenoid tubules (**Fig. S5**). We

conclude that BST4_{ΔC-term}-Venus is recruited to the pyrenoid through an interaction
with the native full-length BST4, which is further evidence that BST4 oligomerizes.

We next investigated whether BST4 localizes to the pyrenoid tubules, or whether
BST4 localizes to the tubules through an interaction with Rubisco. To do so, we utilized
a Chlamydomonas mutant generated by Genkov et al. (2010) that expresses *At*RBCS
but lacks both isoforms of *Cr*RBCS (*Crrbcs::AtRBCS*), and thus lacks a Rubisco matrix
as EPYC1 does not interact with *At*RBCS (Atkinson et al., 2019). *Crrbcs::AtRBCS*
retains reticulated thylakoid membranes at the canonical pyrenoid site, which are likely
the nascent pyrenoid tubule network (Caspari et al., 2017) (**Fig. S6A**). Caspari et al.
(2017) show large starch granules accumulating at the canonical pyrenoid site, which
we confirmed by expressing STA2-Venus as a starch marker (Mackinder et al., 2017)
in *Crrbcs::AtRBCS* (**Fig. S6B**). We expressed BST4-Venus in *Crrbcs::AtRBCS* and
found that BST4 localized to a punctum adjacent to the canonical pyrenoid site, which
we attribute to the nascent pyrenoid tubules (**Fig. S6C**). To confirm this, we also
expressed a known pyrenoid tubule marker protein, PsaF (Emrich-Mills et al., 2021)
in *Crrbcs::AtRBCS*, which showed a similar localization pattern to that of BST4 (Fig.
S6D). This suggests there may be an additional Rubisco-binding independent
mechanism for localization of BST4 as well as other proteins to the pyrenoid tubules.
Further investigation of the role of the whole C-terminal domain will be required to
understand the mode of BST4 pyrenoid tubule localization.

Chlamydomonas *bst4* mutant is not defective in growth at air level CO_2 but increased H_2O_2 production

To test whether BST4 has a role in the operation of the CCM, we measured the growth of *bst4* compared the WT control strain under various CO_2 conditions (**Figs. 6, S7**). Spot assays did not reveal any reduction in growth under CO_2-limiting conditions (**Fig. S7**). When grown in liquid medium, *bst4* even seemed to grow slightly better than WT when sparged with 0.04% CO_2 (**Fig. 6A**). However, when comparing the calculated specific growth rates (μ h^{-1}) for both the exponential growth phase (days 0-3, *bst4* 0.0402 ± 0.0003 μ h^{-1} and WT 0.0389 ± 0.0007 μ h^{-1}) or the full growth assay (days 0-5, *bst4* 0.0257 ± 0.0001 μ h^{-1} and WT 0.0241 ± 0.001 μ h^{-1}), there was no statistically significant increase in specific growth rates between *bst4* and WT (two-tailed t-test, p = 0.17 and 0.20, respectively, n=3, full results **Table S1**). We conclude from these experiments that BST4 is not essential for growth at air levels of CO_2 and might not be necessary for the functioning of the CCM. We also included the complemented *bst4::BST4* and *bst4::BST4$_{\Delta C\text{-}term}$-mScarlet-I* (hereafter *bst4::BST4$_{\Delta C\text{-}term}$*) lines in the spot and liquid growth assays. While all lines grew well in the spot assay, in the liquid growth *bst4::BST4$_{\Delta C\text{-}term}$* grew comparably to WT and *bst4* whereas *bst4::BST4* exhibited a slightly reduced growth than the other lines in both CO_2 conditions.

Figure 6. The *bst4* mutant does not have an impaired growth phenotype under CCM induced conditions but has increased H₂O₂ production. A. *Chlamydomonas* strains were subjected to a liquid growth assay using pH 7.4 TP media that was bubbled with 0.04% CO_2 or 3% CO_2 (+/- 2 ppm). Error bars are ± SEM (n=3). **B.** Dot assay of WT, *bst4,* complemented *bst4::BST4* and *bst4* complemented with C-terminal truncation of BST4 (*bst4::BST4*ₐC-term) on minimal pH 7.4 TP agar at indicated light intensities and CO_2 concentrations. **C.** H₂O₂ assay. Chlamydomonas cells were grown in pH 7.4 TP liquid media and exposed to 150 µmol photons m⁻²s⁻¹ for 24 h with or without the photosynthetic inhibitor DCMU (10 µM). The concentration of H₂O₂ was subsequently quantified using Amplex Red (n=4), and is presented both proportionately to cell density and chlorophyll content. Chlorophyll content was quantified for all cell lines (n=3). Different letters indicate significance ($p<0.05$) as determined by a one-way Anova and Tukey's post-hoc test.

One noticeable difference between *bst4* and WT lines from the growth assays
conducted on solid media (**Fig. 6B and S7**) was that *bst4* cells had a distinct halo of
diffuse cells on the periphery of the colony. We used a range of CO_2 and light
conditions to investigate the diffuse colony phenotype (**Fig. 6B**) and found it was most
apparent under high light (300 µmol photons m^{-2} s^{-1}) and low or very low CO_2
conditions (0.04% CO_2, 0.01% CO_2, respectively) (**Fig. 6B**). WT, *bst4::BST4* and
bst4::BST4-mScarlet-I complemented lines had little or no diffusivity, although WT did
display a slightly diffuse colony phenotype at 300 µmol photons m^{-2} s^{-1}. Interestingly,
*bst4::*BST4$_{\Delta C-term}$ was unable to rescue the diffuse colony phenotype suggesting that
either the presence of the C-terminus or localization to the tubules is essential for the
function of BST4.

The diffuse colony phenotype was most apparent under high light (300 µmol photons
m^{-2} s^{-1}), to a slightly lesser extent at medium light (150 µmol photons m^{-2} s^{-1}), and
exacerbated by low CO_2 (0.01% and 0.04%). These are conditions where carbon
fixation can be limiting and therefore, the energy production by photosynthesis could
exceed the energetic demand required to fix CO_2 by the CCM and the Calvin cycle.
This imbalance can result in the release of Reactive Oxygen Species (**ROS**) (Erickson
et al., 2015). Because cells exposed to ROS have altered phototactic responses to
light (Wakabayashi et al., 2011), we thus thought to assess the phototactic capacity of
bst4 and its control WT strain by exposing them to directional light in liquid culture
(**Fig. S8A**). In this assay, *bst4* cells displayed strong positive phototaxis, whereas WT
and complemented *bst4::BST4* lines displayed negative phototaxis. To test whether
the phototactic response of *bst4* was due to an increase in ROS production, we
recorded the direction of phototaxis in cells exposed to either ROS or a ROS quencher.
In the presence of the ROS quencher N,N'-dimethylthiourea (**DMTU**), *bst4* positive
phototaxis was disrupted, resulting in a negative response to directional light (**Fig.
S8A**). When the ROS H_2O_2 (75 µM) was added, WT, *bst4::BST4* and *bst4* displayed
positive phototaxis (**Fig. S8A**). To directly quantify the difference in ROS generation,
we analysed the H_2O_2 produced by cells exposed to 150 µmol photons m^{-2} s^{-1} (**Fig.
6C**). The *bst4* and *bst4::*BST4$_{\Delta C-term}$ lines had significantly higher H_2O_2 production than
WT and *bst4::BST4* when normalized to both chlorophyll content and cell density. We
validated the assay by using the same ROS quencher DTMU from **Fig. S8A**, and saw
a consistent reduction in H_2O_2 detected in all lines **Fig. S8B**. To test the involvement

of photosynthetic activity in the increased ROS production, we also treated cells with the PS II plastoquinone binding site inhibitor 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) whereby the concentration of H_2O_2 produced was reduced in all lines (**Fig. 6C**). The cellular chlorophyll content was also assessed, we found that *bst4* and *bst4::BST4$_{\Delta C-term}$* had significantly lower cellular chlorophyll content than WT and *bst4::BST4.* The reduction in cellular chlorophyll and the increase in ROS scavenging pigments is a known physiological response to high light exposure and subsequent elevated ROS production (Bonente et al., 2012) and is consistent with other studies (Ma et al 2020; Erikson et al 2015).

ROS generation occurs when light energy exceeds the capacity of electron transport, which can be caused by either high light conditions or limited CO_2 fixation due to an impaired CCM or low CO_2 availability (Santhanagopalan et al., 2021; Choi et al., 2022). As *bst4* can maintain similar growth to WT under low CO_2 conditions, we conclude that the absence of BST4 might affect photosynthesis more broadly and not specifically influence the sustained delivery of CO_2 to Rubisco.

BST4 regulates the lumenal pH in Chlamydomonas during dark to light transition

To assess the impact of BST4 on photosynthesis, we used a pulsed amplitude modulation fluorimeter (**PAM**) to measure chlorophyll fluorescence upon a dark to light transition in all the strains (**Fig. 7**). We used the PAM to assess the quantum yield of PSII (**Y(II)**) and the amount of non-photochemical quenching of chlorophyll fluorescence (**NPQ**) (**Fig. S9**). NPQ is mediated by multiple mechanisms and harbors multiple components (Erickson et al., 2015), one of them, termed energy-dependent quenching, (**qE**) is quickly induced and relaxed and is mostly mediated by the activity of the proton-sensing Light Harvesting Complex Stress related 3 (**LCHSR3**) protein (Bonente et al., 2011; Peers et al., 2009; Steen et al., 2022; Tian et al., 2019). The magnitude of qE has recently been shown to be an indicator of the lumenal pH, the lower the pH, the higher the qE (Burlacot et al., 2022; Tian et al., 2019). While no differences in Y(II) were observed between the strains upon a dark to light transition (**Fig. S9C, D**), the NPQ of *bst4* mutant was transiently higher than WT (**Fig. 7A**), the NPQ becoming undistinguishable from WT after three minutes of illumination (**Fig.**

7A). The complementing *bst4::BST4* strain was undistinguishable from the WT control during

Figure 7. Chlamydomonas *bst4* mutant has an enhanced initial NPQ response under limiting Ci conditions.
Wild type (WT) and mutants were grown in HS medium at 80 μmol photons m^{-2} s^{-1} and measured at 10 μg Chl ml^{-1} (prepared by dilution). The cells were dark adapted for 5 min before the measurements. **A.** Dynamics of Non-photochemical quenching (NPQ) on transition from dark to high light. Kinetics for induction of chlorophyll fluorescence were recorded during 5 min of illumination at 896 μmol photons m^{-2} s^{-1} followed by 5 min in darkness. **B.** Dynamics of NPQ during dark to light transition with one minute of light exposure. Shown are individual data points (dots) and their average (lines) **C.** Calculated fast relaxing NPQ after one minute of light exposure as determined by NPQ at the light to dark transition minus the minimum NPQ in the dark. **D.** Immunoblot of NPQ protein LHCSR3 in each strain plus PetB and Histone as loading controls.* **E.** Quantification of LHCSR3 band

the first three minutes of illumination and *bst4::BST4*$_{\Delta C\text{-}term}$ had a NPQ similar to *bst4*

(**Fig. 7A**). To establish the nature of the transiently increased NPQ in *bst4*, we used a

shorter illumination time (**Fig. 7B**). The entirety of the 45% increase in NPQ for *bst4*

mutant was quickly dissipated (**Fig. 7 B, C**) suggesting that it could be attributed to

qE. Similar trends of NPQ kinetics were observed when cells were supplemented with

HCO_3^- before the measurement, although NPQ relaxation in the dark was different

between lines (**Fig. S10**). Since *bst4* did not accumulate more LHCSR3 as compared

to WT (**Fig. 7D, E**), we conclude that BST4 is involved in the lumenal pH regulation

upon a dark to light transition.

The build-up of lumenal H^+ concentration is usually accompanied with a build-up in the

proton motive force (**PMF**) across the thylakoid membrane, which is used by the

ATPase to generate ATP. We used Electrochromic Shift (**ECS**) measurements

(Bailleul et al., 2010) to measure the total PMF size, as well as initial PMF dissipation

rate (g_H^+) (**Fig. 7F, G & S11**). Interestingly, after one minute of illumination, which are

conditions where the lumenal pH was higher in *bst4* mutant, neither the PMF, g_H^+ nor

proton flux (v_H^+) differed between *bst4* and its WT strain. We conclude that the

changes induced by BST4 on the lumenal pH are either a small contribution to the pmf

formation or are compensated for in the *bst4* mutant.

To assess a longer timescale acclimation of the *bst4* mutant, we performed similar

measurements but on cells pre-treated with three hours of high light (150 µmol photons

m^{-2} s^{-1}) and for a longer time-period (**Fig. S12**). We found that *bst4* was able to

maintain a higher F_v/F_m than WT **(Fig. S12A),** had higher sustained NPQ compared to

WT, which was concurrent with a higher LHCSR3 expression as compared to the WT

strain **(Fig. S12B, G, H).** No significant differences in PMF or g_H^+ were observed

although v_H^+ was higher in *bst4* compared to WT (**Fig. S12D, E, I, J**). The higher v_H^+

together with the higher expression of LHCSR3 suggest a build-up of protons in the

tubule lumen. Since ROS induce photoprotective mechanisms (Roach & Na, 2017),

we propose that in the absence of BST4, enhanced production of ROS **(Fig. 6)** might

lead to enhanced NPQ and photoprotective mechanisms.

As a result of the NPQ difference seen between *bst4* and WT lines in response to

illumination, we proposed that BST4 might be an anion channel involved in regulating

the pH of the thylakoid lumen. Bestrophins are typically permeable to Cl^- and HCO_3^-.

A plant thylakoid bestrophin, *At*VCCN1, is permeable to Cl^- and is also active in the

first minutes of illumination to modulate the lumenal pH, although *vccn1* mutants have

lower NPQ (Herdean et al., 2016). Alternatively, like BST1-3, BST4 may be permeable

to HCO_3^- (Mukherjee et al., 2019). To determine what BST4 may be permeable to, we

expressed BST4 in Xenopus oocytes and measured currents in the presence of

different anions (**Fig. S13A**). No currents for BST4 were detected in the presence of

100 mM KCl or 100 mM $NaHCO_3$. As some bestrophins are autoinhibited by their C-

terminus (Qu et al., 2006), we also tested two C-terminal truncations of BST4 (0-386

and 0-591), but no currents were detected. Another possibility is that BST4 is

permeable to organic ions, similarly to *Hs*Best1, which has been shown to be

permeable to γ-aminobutyric acid (GABA) (Lee et al., 2010) and glutamate (Woo et

al., 2012), as well as Cl^- and HCO_3^- anions. Fei et al. (2022) proposed a model

whereby RuBP acts as proton carrier to increase H^+ concentration in the pyrenoid

tubules. It is possible that BST4 is the channel that facilitates RuBP translocation in

the tubules in this model. To test this hypothesis, we used small molecule analogues

K-PEP and K-Gluconate but no currents were detected for these either (**Fig. S13A**).

Therefore, we were unable to draw conclusions as to what BST4 is permeable to.

BST4 may require certain conditions to be open that are not met in the oocyte system,

such as post-translational modifications, a specific pH, a specific voltage, or an

interaction partner. BST4 was found to be phosphorylated and had an oxidized

methionine residue in its C-terminus (Bergner et al., 2015). Methionine oxidation can

serve as channel-regulating post translational modification (Ciorba et al., 1999), which

would fit with the role that BST4 appears to have in preventing oxidative stress.

BST4 has no impact on growth and photosynthesis in Arabidopsis S2$_{Cr}$ line

Although the permeability is undefined, if BST4 is a functional ion channel, it is possible

that its presence in plant thylakoid membranes might have an effect. We generated

three independent BST4 no tag lines in the S2$_{Cr}$ background and used them to assess

the impact of BST4 on Arabidopsis physiology (**Fig. S14A**). We found there was no
difference in growth between plant expressing BST4 and their azygous segregants,
as determined by the rosette area (**Fig. S14B**). We also found that BST4 expressing
plants tended to have slightly lower F_v/F_m as compared to azygous segregants and
the parent line, although this was not significant (**Fig. S15A**). Further measurements
were made for lines 2 and 3 on the kinetics of NPQ, Y(II) and for PMF size and
partitioning but no consistent differences were observed (**Fig. S15**).

Proposed model for the role of BST4 in the Chlamydomonas pyrenoid

Based on these findings, we propose that BST4 moderates the thylakoid lumen
through its function as a bestrophin channel and is influential in this role during the
transition from dark to light. Specifically, we propose that BST4 is a pentameric HCO_3^-
transmembrane channel found within the pyrenoid tubules but is not crucial for
Rubisco matrix tethering. Rather, BST4 is targeted to the tubules by the C-terminal
RBMs to facilitate its role in the pyrenoid. In its proposed primary role as a HCO_3^-
channel, BST4 may form part of a recovery system to take up HCO_3^- from within the
pyrenoid, allowing HCO_3^- to be transported back into the tubules where it could be
dehydrated to CO_2 and thus available to be fixed by Rubisco **(Fig. 8).** The close
proximity of BST4 to the luminal carbonic anhydrase CAH3 further supports this
hypothesis. As indicated by our NPQ data, this role might be helpful during the dark to
light transition where HCO_3^- is needed immediately to both moderate lumenal H^+
concentration and begin Ci uptake when Ci has not yet been transported into the
thylakoid tubules. In this scenario, the absence of BST4 would lead to an initial build-
up of protons as they are not being consumed by HCO_3^- dehydration, thus the initial
high NPQ. After a minute, the standard BST1-3 HCO_3^- uptake system would take over,
protons being used for HCO_3^- dehydration and resulting in pH stabilization and NPQ
relaxation. Over the longer term, a lack of BST4 may lead to a slight reduction in CCM
operation efficiency, perhaps not sufficient to cause a growth penalty but enough to
alter the homeostasis of photosynthesis and thus increase ROS production. Future
work is needed to determine what species BST4 is permeable to and how it is
regulated. In summary, BST4 might play a role in modulating the intricate dynamics
between the CCM and photosynthetic ion management, two physiological processes
that are key to algal primary productivity.

Figure 8. The role of BST4 in light and CO₂ limiting conditions. The data presented here indicate that BST4
moderates the thylakoid lumen through its function as a bestrophin channel and is influential in this role during the
transition from dark to light. 1) When the light is turned on, protons are generated by the electron transport chain
(ETC) resulting in a luminal pH decrease and NPQ is initiation. 2) BST4 allows the passage of HCO_3^- from the
pyrenoid Rubisco-matrix into the lumen. The HCO_3^- is then dehydrated by the carbonic anhydrase CAH3 to
generate CO_2, which can subsequently diffuse out of the lumen into the pyrenoid Rubisco-matrix for fixation. The
conversion of HCO_3^- to CO_2 also acts as a sink for the protons generated by the ETC, increasing the luminal pH
and reducing NPQ. This mechanism (shown in 1, 2) is most impactful in the first minute of illumination, after which
(3) HCO_3^- also begins to enter the thylakoid lumen through BST1-3 and the impact of BST4 uptake is reduced.

MATERIALS AND METHODS

Phylogenetic Analysis

Amino acid sequences for phylogenetic analysis were compiled by blasting BST4 (Cre06.g261750) in the NCBI database (Sayer et al 2022) and the manual addition of other well characterized bestrophin proteins including BEST1 *Homo sapiens* (XP_011543531.1), KpBEST *Klebsiella aerogenes* (WP_049046555.1), VCCN1 from *Arabidopsis thaliana* (Q9M2D2) and BST1-3 *Chlamydomonas reinhardtii* (Cre16.g662600, Cre16.g663400 and Cre16.g663450). Sequences were aligned using MAFFT (Katoh and Standley, 2013) visually inspected and manually trimmed (specified). Finalized alignments were run through IQTREE webserver (Minh et al., 2020) to identify the most appropriate substitution model. Maximum likelihood trees were then generated in Geneious v11 using the PhyML 3.0 (Guindon et al., 2010) plugin with a LG substitution model (Le and Gascuel, 2008) with Gamma distribution (4 categories) and 500 bootstrap iterations. Full alignments are found in **Fig. S16**.

Alphafold structure prediction

Five chains of BST4 were submitted to Alphafold Multimer v2 with default settings. using the ColabFold server (Mirdita et al., 2022). All protein structure figures were generated using UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases (Pettersen et al., 2021). The top ranked model of five was used for figure generation.

Generation of plasmids

The plasmids for BST4 mutant complementation in *Chlamydomonas* were prepared using a recombineering method described previously (Emrich-Mills et al., 2021). BST4 is expressed under native promotor with either without a tag or with an mScarlet-I C-terminal tag, and a hygromycin AphVII selection marker. The same approach was used to generate STA2-Venus lines in the *Crrbcs::AtRBCS* background (Genkov et al., 2010).

Plasmids for plant expression were generated using the MoClo system (Engler et al., 2014). For visualisation, BST4 was combined with 35S promoter (pICH51277),

mNeonGreen C-terminal tag (pICSL50015), HSP terminator and acceptor plasmid (pICH47732) and a pFAST-R section cassette used for selection. For all other experiments a no-tag BST4 construct was generated with the 35S protomer and HSP terminator parts and a Kanamycin resistance cassette was used for selection.

Plasmids for Xenopus expression were generated by using the Gateway system by cloning the coding sequence for BST4 into pGT vector (Grefen et al., 2010). *BST4* was amplified with Gateway adaptor sequences from a synthesized as g-block (IDT) with the PredAlgo (Tardif et al., 2012) predicted transit peptide removed from the N-terminal (sequence begins R35), any subsequent mutations were made via PCR.

Arabidopsis transformation

Arabidopsis was transformed by floral dip as previously described in (Atkinson et al., 2016). BST4-mNeon primary transformants were screened for transgene insertion by seed fluorescence from pFAST-R and BST4 expression was confirmed by checking for mNeon fluorescence and by immunoblot. BST4 no tag primary transformants were screened using kanamycin resistance and immunoblot. Zygosity was checked via seed fluorescence from pFAST-R or kanamycin resistance.

Chlamydomonas cell culture conditions and strain details

Chlamydomonas reinhardtii cultures were maintained as previously described (Ma et al., 2011). Tris-Acetate-Phosphate (TAP) and minimal (TP) media (acetate free) were prepared according to Sueoka (1960). TAP and TP agar plates for growth were made by adding 1.5% (w/v) agar. CMJ030 (CC-4533; *cw15, mt⁻*) and *bst4* (BST4 knock-out LMJ.RY0402.159478) were obtained from the CLiP collection at the *Chlamydomonas* culture collection (Zhang et al., 2014; Li et al., 2019). This CLiP mutant has two other mapped CIB1 cassette insertions at loci Cre04.g230046 and Cre08.g367750. The insertion of the CIB1 cassette in *BST4* locus was confirmed by PCR amplifying the insertion locus from genomic DNA (Fig. S3A) using loci specific primers (forward GAGCTTCGTGGATGGATGTT and reverse GTATGAAGGTCACCGCCTGT) in parallel with a control locus (forward ATGCTTCTCTGCATCCGTCT and reverse ATGTTTTACGTCCAGTCCGC). The additional two insertions were also confirmed using loci specific primers for Cre04.g230046 (forward TGTGCCTCTGTCAGTCTTGG and reverse TGCGTGGATGGGTAACAGTA), Cre08.g367750 (forward AATCAAGAAGCTTCCCAGCA and reverse CCTACCGCTATCTCAGCCAG) and

657 STT7 locus as a control (forward GCACGAACCAAGACACACATAG and reverse
658 GTAGACGATGTCACCGCACTT). Therefore, the *bst4* knock-out was complemented
659 with *BST4* constructs described herein. All complemented lines were validated by
660 western blotting of BST4 and specified epitope tags, described below (Fig. S3B-E).

Chlamydomonas transformation

For each Chlamydomonas transformation, 28 ng kbp^{-1} of plasmid was linearized by
restriction digest. Cells were grown to 2 - 4 x 10^6 cell mL^{-1}, harvested by centrifugation
at 1000 xg for 10 min and resuspended in TAP with 40 mM sucrose at a concentration
of 2 × 10^8 cells mL^{-1}. Linearized plasmid was mixed with 250 µL of cells at 15°C in a
0.4 cm gap electroporation cuvette and transformed immediately by electroporation
using a Gene Pulser II (Bio-Rad) set to 800V and 25 µF. Cells were recovered
overnight in TAP sucrose while shaking gently (140 rpm) in the dark. Transformed cell
were subsequently subjected to selection by growth on TAP agar plates with
paromomycin (20 µg mL^{-1}) or hygromycin (25 µg mL^{-1}) which were kept in low light
(5–10 µmol photons m^{-2} s^{-1}) until screening positive transformants.

Chlamydomonas Growth Assays

Spot Tests: Cells were grown heterotrophically in TAP media. Once cultures reached
2 - 4 x 10^6 cell mL^{-1}, 1 x 10^6 cells were harvested by centrifugation at 1000 xg for 10
min. Cell were washed and resuspended at a concentration of 1 x 10^6 cell ml^{-1} in TP
media. Liquid cultures were spotted onto TP agar (1.5%) in 1000, 100 and 10 cell
spots at a range of pHs (specified). The plates were incubated in 3, 0.04 and 0.01%
CO$_2$ and illuminated under constant light at 400 µmol photons m^{-2} s^{-1}. Growth was
monitored for up to 10 days.

Liquid Growth: Cells were grown heterotrophically in TAP media. Once cultures
reached 2-4 x 10^6 cell mL^{-1}, cells were harvested by centrifugation at 1000 xg and
resuspended at a starting concentration of 1 x 10^5 cell ml^{-1} in TP media pH 7.4.
Cultures were incubated in a CellDEG HDC 22.10 culture platform (CellDeg GMBH,
Berlin) bubbled with 0.04 and 3% CO$_2$, illuminated at 150 µmol photons m^{-2} s^{-1} and
consistently stirred at 180 rpm. Cell density and optical destiny (750 nm)
measurements were taken daily for up to 10 days. Specific growth rates (SGR) per h
were calculated using the following formula: $\mu = ln(N_2/N_1)/t$ whereby N = cell density

Dot tests: Cultures were grown in a 96 format on agar plates and replicated by a Rotor+
(Singer Instruments) high throughput replication robot. The cultures were stamped
onto pH 7.8 TP agar plates, incubated in 3, 0.04 and 0.01% CO_2 and illuminated under
constant light at a range of intensities (specified). Growth was monitored for up to 10
days.

Phototaxis Assays

Chlamydomonas cells were grown heterotrophically in TAP media until they reached
2 - 4 x 10^6 cell mL^{-1} and harvested by centrifugation at 1000 xg for 10 min. Pelleted
cells were either re-suspended in TP media or, for ROS manipulation assays, a
phototaxis buffer described previously by Ueki et al. (2016) (5 mM Hepes pH 7.4, 0.2
mM EGTA, 1 mM KCl, and 0.3 mM $CaCl_2$). The assays took place in 12-well dishes
with a thin layer of TP agar (0.8%) on the well bottom and approximately 1.5 x 10^7
cells in 400 μL of homogenous suspension laid on top. The dishes were illuminated
from one direction with 150 μmol photons m^{-2} s^{-1} illumination for up to 3 h. Plates were
imaged using a Flatbed Scanner at specified intervals.

Quantification of H_2O_2

Cells were grown heterotrophically in TAP media. Once cultures reached 2 - 4 x 10^6
cell mL^{-1}, cells were harvested by centrifugation at 1000 xg and resuspended at a
concentration of 2 x 10^6 cell ml-1 in TP media pH 7.4, illuminated at 150 μmol photons
m^{-2} s^{-1} for 24 h and shaken at 140 rpm. For H_2O_2 quantification, 1 mL of culture was
diluted at a 1:1 ratio with fresh TP media, containing 1U of horseradish peroxidase
and 5 μM of Amplex Red (ThermoFisher) and incubated for 1 h (illuminated at 150
μmol photons m^{-2} s^{-1}, shaking 140 rpm). Cells were removed by centrifugation. The
H_2O_2 of the media was immediately quantified using a ClarioStar Plate Reader
Excitation/Emission 520/570-600 and compared against a linear H_2O_2 standard curve
up to 5 μM. Additional controls were included; some cells were treated with the ROS
quencher N,N' dimethylthiourea (DMTU) at a final concentration of 150 μM; and some
with the PSII plastoquinone binding site blocker 3-(3,4-dichlorophenyl)-1,1-
dimethylurea (DCMU) was dissolved in methanol at a final concertation of 10 μM prior
to H_2O_2 quantification (specified). All measurements were conducted with a minimum
of four technical replicates. The data shown represents one of a multiple of
experimental repeats conducted on different days with fresh cultures. All H_2O_2

concentrations were normalized to cell density, calculated as described previously,
and chlorophyll content, described below.

Total chlorophyll was calculated by resuspending 1 mL of harvested cells in 1 mL of
methanol. All samples were protected from the light after menthol addition. After
vortexing for 1 min to resuspend the pellet and incubating for 10 min, the cells were
removed by centrifugation. The absorbance of the supernatant was analysed by
spectrophotometer at 652 and 665 nm. Total chlorophyll was calculated using the
formula below. All measurements are averaged from three technical replicates. Total
chlorophyll (µg/mL) = 22.12 x Abs652 + 2.71 x Abs665.

Growth of Arabidopsis
Arabidopsis seeds were sown on moist F2+S soil and stratified in the dark at 4℃ for 2
days. For growth experiments seeds were grown in a Percival SE-41AR3cLED
chamber (CLF PlantClimatics GmbH, Wertingen, Germany) equipped with cool white
LED lights under 12 h light (175-180 µmol photons m^{-2} s^{-1})/12 h dark cycles at 21°C,
respectively and 70% relative humidity.

Chlamydomonas Confocal Microscopy
Transgenic fluorescent strains were initially grown heterotrophically in TAP media until
reaching 2 – 4 ×10^6 cells mL^{-1} and resuspended in TP media overnight prior to
imaging. Cells were mounted on 8-well chamber slides and overlayed with 1.5% low
melting point agarose made with TP-medium. Images were collected on a LSM880
(Zeiss) equipped with an Airyscan module using a 63× objective. Laser excitation and
Emission setting for each channel used are set as below: Venus (Excitation: 514 nm;
Emission 525 – 500 nm); mScarlet-I (Excitation: 561 nm; Emission 570 – 620 nm);
Chlorophyll (Excitation: 633 nm; Emission 670 – 700 nm).

Yeast-2-Hybrid
Yeast two-hybrid to detect interactions between BST4 C-terminus and RbcS1 was
carried out as described in He et al., (2020). BST4 C-terminus (amino acids 387-end)
was cloned into the two-hybrid vector pGBKT7 to create a fusion with the GAL4 DNA-
binding domain. Point mutations were introduced by PCR into BST4 RBMs, which was
then cloned into the same vector. Mature CrRBCS1 was cloned into the vector
pGADT7 to create a fusion with the GAL4 activation domain. Yeast cells were then co-

transformed with binding and activation domain vectors. Successful transformants

were cultured, diluted to an optical density at 600 nm (OD_{600}) of 0.5 or 0.1 and plated

onto SD-L-W (double drop out, DDO) and SD-L-W-H (triple drop out, TDO) media. The

plates were imaged after three days and **Fig. S2** shows yeast spots from cultures

diluted to an OD_{600} of 0.5.

Slimfield microscopy

Chlamydomonas lines *bst4::BST4-mScarlet-I* and the unlabelled control line *bst4* were

prepared overnight in TP media. Each was harvested and spotted onto a slide-

mounted agar pad (GeneFrames, ThermoFisher), consisting of TP media with 1.5%

low melting point agarose. Fluorescence imaging with single-molecule sensitivity was

performed using a custom Slimfield microscope (Syeda et al., 2019). The setup used

a high-magnification objective (NA 1.49 Apo TIRF 100× oil immersion, Nikon) and the

detector was a Prime95B sCMOS camera (Teledyne Photometrics) operating in 12-

bit 'sensitivity' gain at a high total magnification of 53 nm/pixel. The samples were

illuminated either in brightfield, or for Slimfield fluorescence in camera-triggered

frames by a collimated 561 nm wavelength, Gaussian mode OPSL laser (Coherent,

Obis LS) at a peak intensity of 5 kW/cm^2 at the sample plane. This beam was tilted

into a HILO configuration (Payne-Dwyer and Leake, 2022) to reduce out-of-focus

excitation while retaining quantitative molecular sensitivity. The fluorescence image

was split into two parallel channels comprising emission bandpass filters (Semrock

BrightLine®): one with a 585/15 emission filter (central wavelength/spectral bandwidth

in nm) optimized to isolate the mScarlet-I signal, and a second with a 525/25 emission

filter, used only to indicate autofluorescence background. The total length of each

acquisition sequence was ~5 s; sufficient to observe the full course of mScarlet-I

photobleaching, from the initial unbleached state to single-molecule blinking, while

also rapid enough (10 ms exposure/frame at 180 frames/s) to capture the motion of

individual molecular assemblies.

Single particle tracking and molecular counting

779 Slimfield image sequences were segmented manually in ImageJ to include only the

780 pyrenoids in downstream analysis. The centroid positions of fluorescent tracks were

781 identified from local intensity maxima in each frame using ADEMScode software in

782 MATLAB (Wollman et al., 2022). The summed intensity of each candidate track was

calculated in each frame by adding all pixel values within 5 pixels of the centroid, then subtracting the local background averaged between 5 - 8 pixels from the centroid. Candidates with a summed intensity <0.4× the standard deviation in the background region were discarded.

Fluorescent proteins are known to exhibit a characteristic integrated intensity per molecule under stable Slimfield imaging conditions and within the quasi-uniformly illuminated area within half the beam waist (Shepherd et al., 2021). After sufficient photobleaching of mScarlet-I in the Slimfield image sequences, only step-like blinking was observed at the end of each track. The modal integrated intensity of these steps was used to estimate this characteristic single molecule brightness, equivalent to 56 ± 9 photoelectrons per frame per molecule.

At the start of each track, we obtained an initial integrated intensity (independent of photobleaching) by linearly extrapolating the summed intensity backwards over the first 4 frames of the exposure. This initial intensity was then divided by the characteristic brightness of a single mScarlet-I to estimate the number of molecules, or stoichiometry, in that track. This estimate was precise enough to detect stoichiometry steps of up to 12 tagged molecules without ambiguity.

Stoichiometry distributions may exhibit peaks which are separated by a characteristic interval. The smallest consistent interval between peaks can be used to infer the size of a physical repeat unit or 'periodicity' within assemblies (Hunter et al., 2022; Payne-Dwyer et al., 2022). A kernel width of 0.7 molecules was chosen to generate the stoichiometry distribution (Fig. 2F), reflecting the background standard deviation. Peaks were pinpointed using MATLAB's *findpeaks*.

The intervals between all peaks for each acquisition were aggregated across the pyrenoid population, weighted by inverse square-root distance (thereby accounting for shot-noise in broader intervals). A second distribution **(Fig. 2G)** was then generated from this weighted population of intervals. The kernel width in this estimate was 0.7 molecules multiplied by the square root of the mean stoichiometry divided by the root number of intervals (thereby accounting for shot-noise in intervals between peaks of higher stoichiometry). The periodicity was then reported as the mode of this distribution and its 95% confidence interval.

Arabidopsis Confocal microscopy

Small sections of 3 – 4 week-old leaf tissue (~5 - 10 mm^2) were adhered to slides using double-sided tape with basal side up. A x40 water immersion objective lens was used. Samples were excited by 488 nm at 1% laser power, chlorophyll autofluorescence was collected at 680 – 750 nm and mNeonGreen fluorescence at 503–532 nm. For dual tagged lines, we used sequential acquisition to minimize bleed-through. mCherry was excited using the 542 nm laser and emission collected at 601-620 nm and mNeon as before. Images were acquired using the SP8 Confocal system and Leica LAS AF software (http://www.leica-microsystems.com/). Figures were prepared using ImageJ (http://fiji.sc/Fiji).

Immunoblot detection

Two leaf disks (6 mm diameter) were harvested and immediately frozen in liquid nitrogen. Two steel balls (3 mm) were added and tissue was homogenized using a tissue-lyser twice for 30 Hz for 30 s. A four times volume of cold extraction buffer (20 mM Tris-HCl pH = 7.5, 5 mM MgCl$_2$, 300 mM NaCl, 5 mM DTT, 1% (v/v) Triton X-100, 1 x Protease inhibitor (Roche)) was added and samples vortexed for 30 s. Samples were solubilized on ice for 5 mins and then centrifuged at 5000 xg for 5 min at 4℃. 17.5 µL of supernatant was used to make up 1x LDS and 100 uM DTT. 20 µL was loaded on a Novex™ 4 - 12% Bis-Tris Mini Gel, (Thermo Fisher, Catalogue number: NP0322BOX). The gel was run at 150 V for 60 minutes. Proteins were transferred to a nitrocellulose membrane using an iBlot 2, programme 0. The membrane was probed with primary antibody in 5% milk in 1 x TBST at the following dilutions: BST4 (1:1000; generated for this study, peptide from C-terminus: SDTELSEANRPRTRPDWRN) (YenZym, Antibodies LLC, USA), AtpB (1:2000; Agrisera:AS05085), RbcL (1:1000; kind gift from Griffiths lab), CP43 (1:3000; Agrisera: AS111787), PsaB (1:1000; Agrisera: AS10695), PsbO (1:2000; Agrisera:AS06142-33). Secondary antibody (goat α-rabbit IR-800; Li-COR: 925-32211). Membrane was imaged using the Li-COR Odyssey CLx scanner.

In order to quantify BST4 protein in Chlamydomonas lines, cells were grown in TP media at ambient CO$_2$ until reaching 2 - 4 x 10^6 cells mL^{-1}. Cells were harvested by centrifugation at 1000 x g for 10 mins, normalized to Chl content and resuspended in the extraction buffer described above. Samples were freeze-thawed three times and

spun at 20, 000 x g for 20 min at 4℃. Protein extractions containing 5 µg of Chl with 1 x SDS loading buffer were boiled at 100℃ for 5 min and loaded onto a 4 - 20% polyacrylamide gel (Mini Protean TGX, Biorad Laboratories). Proteins were transferred to a PVDF-FL membrane on a Biorad semidry blotting system. BST4 primary antibody was used as described above alongside alpha-tubulin primary antibody raised in mouse (Agrisera), as a loading control. Anti-rabbit and anti-mouse fluorescent secondary antibodies, Invitrogen AlexaFluor 488 and 555 respectively, were used at a 1:20 000 dilution. Immunoblots were imaged using an Amersham typhoon 5 scanner with 488 and 535 excitation lasers and Cy2 and Cy3 emission filters. BST4 band fluorescent intensity was quantified using FIJI (Image J) (Schindelin et al., 2012) and normalized to alpha-tubulin loading control. All Chlamydomonas lines for quantification were extracted and analysed in triplicate. For LHCSR3 protein quantification, cells were seeded at 0.1 OD_{750} in TAP media for 4 days and then switched to TP media at a concentration of 30 µg Chl ml^{-1} and exposed to 150 µmol photons m^{-2} s^{-1} for 3 h. Protein was extracted according to Burlacot et al. 2022 and separated by SDS-PAGE as described for BST4 immunoblots.

Oocytes for expression of BST4 were collected after recording and prepared for western blot as described in Lefoulon et al., 2018. The BST4 primary antibody was used as described above and the secondary antibody used was horseradish peroxidase-coupled goat, anti-rabbit (dilution 1:10000 Abcam). Proteins were detected with ECL Advance kit (GE Healthcare,Poole UK).

Blue-Native PAGE

A crude thylakoid enrichment was performed according to (Aro et al., 2004). Thylakoid membranes were solubilized in 0.5% DDM, 1x NativePAGE buffer (Thermo: BN2003), 1x cOmplete protease inhibitor tab (Roche;10x stock made by dissolving 1 tablet in 1 mL dH_2O) for at a concentration of 0.8 µg Chl/µl for 15 min on ice. Unsolubilized material was removed by two rounds of centrifugation at 17,000 xg at 4°C for 15 min. 19.5 µL of supernatant was combined with 0.5 µL of Coomassie additive and loaded immediately onto a 4-16% Bis-tris gel (Thermo: BN1002BOX). Electrophoresis was performed at room temperature at 150 V for 90 min. Cathode buffer was swapped from dark to light when the dye front was a third way through the gel. Separated

proteins were transferred to a nitrocellulose membrane by electrophoresis at 100 V for
90 min at 4 ℃.

Proteins were visualized using chemiluminescence. Secondary antibody (goat α-
rabbit, HRP;1:10,000; Abcam: ab6721). Chemiluminescence substrate SuperSignal
West Pico PLUS (ThermoScientific, ref number: 34579) according to manufacturer's
instructions. Chemiluminescence was detected using clear blue X-Ray Film CL-
Xposure^{TM} Film (ThermoScientific, ref number: 34090).

Chlorophyll fluorescence measurements in Chlamydomonas for Fig. 7
To measure PSII activity in Chlamydomonas, cells were grown in HS media under 80
µmol photons m^{-2} s^{-1} for 3 days at 120 rpm (in a Multitron, Infors-ht) to reach
logarithmic phase and then maintained at ~10 µg Chl mL^{-1}. Two millilitres of cells were
added to a cuvette and bubbled with air (10 cc/min) and continuous stirring. Cells were
incubated in the dark for five minutes before recording Chl fluorescence using DUAL-
PAM-100 (Walz, Effeltrich, Germany). A saturating pulse of 8,000 µmol photons m^{-2}
s^{-1} of 300 ms was applied to the samples for determination of the maximal
fluorescence yield in the dark state (F_m) and maximal fluorescence yield during the
period with actinic light (F_m'). The maximal quantum efficiency of PSII was calculated
as (F_m'-F)/F_m where F is the stationary fluorescence. NPQ was calculated as (F_m–
F_m')/F_m' . Far-red light (4 µmol photons m^{-2} s^{-1}) was used throughout the entire
experiment to limit state transition from contributing to the NPQ. NPQ and Y(II) were
calculated based on changes in Chl fluorescence as (F_m–F_m')/F_m' and (F_m'–F)/ F_m',
respectively, according to Genty et al., (1989). When indicated, cells were
supplemented with a final concentration of 500 µM HCO_3^- at the beginning of the dark
adaptation.

Electrochromic shift (ECS) in Chlamydomonas for Fig. 7
Electrochromic shift (ECS) in Chlamydomonas was assessed by measuring the
absorbance changes of cells at 520 and 545 nm using a JTS-100 spectrophotometer
(BioLogic). Cells were grown and prepared as for Chl fluorescence experiments
described previously except cells were resuspended to 150 µg Chl mL^{-1} before being
loaded into a custom vertical light path cuvette. Cells were dark adapted for 1 min and
then exposed to 890 µmol photons m^{-2} s^{-1} red light (630 nm) for 1 min. The light was
switched off and decay kinetics were measured. ECS signal was calculated as the

difference between absorbance changes measured at 520 and 545 nm. For each
biological replicate, 3 technical replicates were taken and averaged. PMF size was
calculated as the difference between the ECS signal in light and the minimum value
of the ECS signal immediately after the light was turned off. The g_H^+ parameter was
calculated as 1/τ (time constant for decay during the first 100 ms (Cruz et al., 2005).

Chlorophyll fluorescence measurements in Chlamydomonas for Fig. S12

To measure PSII activity in Chlamydomonas, cells were grown in TAP in low light (20
µmol photons m^{-2} s^{-1}) for 4 days at 50 rpm to reach logarithmic phase. Cells were
then washed, resuspended to 30 µg Chl mL^{-1} in TP media and exposed to 150 µmol
photons m^{-2} s^{-1} light for 3 h, followed by 1 h incubation in darkness at 50 rpm before
recording Chl fluorescence using DUAL-PAM-100 (Walz, Effeltrich, Germany). A
saturating pulse of 3,000 µmol photons m^{-2} s^{-1} of 800 ms was applied to the samples
in a cuvette under continuous stirring for determination of the maximal fluorescence
yield in the dark state (F_m) and maximal fluorescence yield during the period with
actinic light (F_m'). The maximal photochemical efficiency of PSII (F_v/F_m) was
calculated. NPQ was determined from slow kinetics during actinic illumination at 1,500
µmol m^{-2} s^{-1} for 17 min followed by 5 min of dark relaxation. NPQ and Y(II) were
calculated based on changes in Chl fluorescence as (F_m–F_m')/F_m' and (F_m'–F)/F_m',
respectively, according to Genty et al., (1989).

Electrochromic shift (ECS) in Chlamydomonas for Fig. S12

ECS measurements in Chlamydomonas were carried out using a the Dual-PAM-100
equipped with a P515/535 module (Walz). Cells grown and prepared as for Chl
fluorescence experiments described previously were layered on a glass slide and
exposed to actinic red light for the given period. The light was switched off and decay
kinetics were measured. PMF size was calculated as the difference between the ECS
signal in light and the minimum value of the ECS signal immediately after the light was
turned off. Calculation of ΔpH and ΔΨ was performed using the steady-state time point
of the ECS signal in darkness (Cruz et al., 2001). Before each ECS measurement, a
3 saturating 50-µs actinic red flashes of 200,000 µmol photons m^{-2} s^{-1} was applied to
determine the ECS$_{ST}$; subsequently, the ECS$_{ST}$ amplitude was used to normalize the
ECS signal before the calculation of PMF size and partitioning values. To determine
H$^+$ conductivity (g_H^+), the light was switched off at specific time points to record the

ECS signal decay during 620 ms dark intervals. The g_H^+ parameter was calculated as
$1/\tau$ (time constant for decay during the first 100 ms (Cruz et al., 2005). The total proton
flux across the membrane was calculated as $v_H^+ = PMF \times g_H^+$ (Cruz et al, 2001).

Chlorophyll fluorescence measurements for Arabidopsis

Plants were grown for 8 weeks on S-Hasselfors soil in a Percival AR-82L chamber
(CLF Plant Climatics, Wertingen, Germany) using 12 h light (180 µmol photons m^{-2}
s^{-1})/ 12 h dark cycles at 21°C/19°C, respectively, and 70% relative humidity. Slow
kinetics of Chl a fluorescence induction were recorded with a pulse-amplitude
modulated fluorometer DUAL-PAM 100 equipped with DUAL-DB and DUAL-E emitter-
detector module (Walz) on attached leaves of 30 min dark-adapted plants using actinic
red light of 830 µmol photons m^{-2} s^{-1} for 10 min, followed by a 5 min dark period. The
saturating pulse applied was 5,000 µmol photons m^{-2} s^{-1} and of 800 ms duration. NPQ
and (Y(II)) were calculated based on changes in Chl fluorescence as $(F_m–F_m')/F_m'$ and
$(F_m'–F)/ F_m'$, respectively (Genty et al., 1989).

Electrochromic shift (ECS) for Arabidopsis

ECS was recorded with a DUAL-PAM 100 system equipped with a P515/535
emitter/detector module (Walz). First, plants were dark adapted for 30 min, then
illuminated with actinic red light at 830 µmol photons m^{-2} s^{-1} for 3 min followed by a
60 s dark period in which the ECS decay kinetics were recorded. Before each
measurement, three pulses of 5 µs and 200,000 µmol photons m^{-2} s^{-1} were applied to
determine ECS$_{ST}$, which was used to normalize the ECS$_T$ values of each
measurement.

Sedimentation of proto-pyrenoid

Cross-linked samples were prepared by vacuum infiltrating intact leaves 1%
formaldehyde prior to sedimentation 200 mg of leaf tissue was flash frozen in liquid
nitrogen and ground by bead beating twice at 30 Hz for 30 s. Four times volume of
extraction buffer (50 mM HEPES-KOH pH 7.5, 17.4% (v/v) glycerol, 2% (v/v) Triton X-
100, cOmplete protease inhibitor tab) was added and sample mixed by bead beating
again. Extract was filtered through one layer of miracloth. A small aliquot of filtered
extract was saved as the input. Extract was then centrifuged at 500 x g for 3 min at 4
℃ and the pellet discarded. The supernatant was centrifuged at 500 x g for 12min, 4
℃. The pellet was washed once with extraction buffer and then centrifuged again. The

pellet was resuspended in 100 μl extraction buffer then centrifuged for a further 5 min.

Pellet was finally resuspended in 25 μl extraction buffer. Fractions were made up in

1x LDS loading buffer and 200 mM DTT. Ten microlitres was subjected to SDS-PAGE

(NuPAGE™ 4-12% Bis-Tris Mini Gel, Thermo Fisher, Catalogue number:

NP0322BOX) at 150 V for 60 min.

Chloroplast fractionation

979 In order to biochemically localize transgenically expressed BST4, chloroplasts were

980 fractionated as described in Herdean et al. (2016) using 100 g of leaf tissue from four-

981 five week-old BST4 transgenic plants. The stromal fraction was concentrated using a

982 10,000 MWCO centrifugal concentrator (Sartorius Stedim Biotech GmbH, product

983 number: VS1502).

Electron microscopy of Chlamydomonas

High (3%) and low (0.04%) CO_2 acclimated cells were harvested by centrifugation

(×1000g, 4 minutes, 20°C). Primary fixation was performed in 1.25% glutaraldehyde

in 50 mM sodium cacodylate (pH 7.15) in TP medium for 15 minutes followed by 2.5%

Glutaraldehyde in 50 mM cacodylate for 2 hours. Fixed samples were washed three

times with 50 mM sodium cacodylate by centrifugation. Samples were then osmicated

with 1% OsO4 in 50 mM sodium 25 cacodylate for 1 hour on ice and washed with de-

ionised water. Samples were block stained in 1% uranyl acetate in the dark for 1 hour.

Samples were washed twice with dH_2O and twice with 50 mM sodium cacodylate.

Fixed samples were dehydrated in an acetone series (25%, 50%, 75%, 90% and

100%) ~20 minutes each step. Dehydrated samples were infiltrated with Spurr's resin

by incubating in 25% then 50% Spurr resin in acetone for 30 minutes and transferred

to 75% for 45 minutes at room temperature. They were then incubated in 100% Spurr

resin overnight before polymerising at 70 °C for 24 hours. Sections ~70 nm thick were

collected on Copper grids and stained with saturated uranyl acetate and lead citrate.

Images were collected with a FEI Tecnai 12 BT at 120kV using a Ceta camera.

Electron microscopy of Arabidopsis

1001 Leaves were cut into small 5 mm strips and fixed in 4% (v/v) Paraformaldehyde/0.5%

1002 glutaraldehyde in 0.05M sodium cacodylate (pH = 7.2) by vacuum infiltration three

1003 times for 15 min and incubation at 4 °C overnight with gentle agitation followed by

1004 dehydration in increasing amounts of ethanol 50/70/80/90/100% 1 hr each then

overnight rotation. 100% ethanol was repeated 3 times, 1 hr each and a final over night at 4 °C. Samples were then fixed in LR resin by infiltrating with increasing concentration (50/70/100%) with a repeat of 100% and then polymerized overnight at 50 °C. Ultrathin sections were cut and mounted onto plastic-coated copper grids. Grids were stained with 2% uranyl acetate and visualized by transmission electron microscopy (TEM).

Protease protection assay

Investigation of the orientation of BST4 in isolated thylakoid membranes was conducted as described in Stepien and Johnson (2018). Briefly, trypsin made up in 50 mM acetic acid according to the manufacturer's instructions (Thermo Scientific, ref number: 90057). To disrupt the thylakoid membrane and allow degradation to lumen-facing peptides, 1% (v/v) Triton X-100 was added, and tubes gently agitated prior to addition of trypsin.

Xenopus oocyte electrophysiology

Destination clones containing BST4 for Xenopus expression were linearized with EcoRI, before proceeding to the *in vitro* transcription (mMessage mMachine T7 transcription kit, Thermofisher Scientific, ref number AM1344). Stage IV oocytes were injected with 20 ng of RNA per oocyte. Measurements of ion transport was done by voltage clamp using an Axoclamp 2B amplifier (Axon instruments, Foster City, CA) (Lefoulon et al., 2014; Leyman et al., 1999). They were performed under perfusion of either 100 mM KCl, K-HCO$_3$, Na-HCO$_3$, K-PEP, or K-Gluconate, with 1 mM CaCl$_2$, 1.5 mM MgCl$_2$, and 10 mM HEPES-NaOH, pH 7.3. Recordings were obtained and analysed using Henry IV software (Y-Science, Glasgow, UK). BST4 expression in oocytes was validated by western blotting (Fig S10B), as described above.

ACKNOWLEDGEMENTS

The authors thank the University of York Horticulture team who helped with setting up the Chlamydomonas phototaxis assays. TEM for Arabidopsis was carried out with the support of Stephen Mitchell and the Wellcome Trust Multi User Equipment Grant (WT104915MA)

AUTHOR CONTRIBUTIONS

LA, CW, CSL, AB, CS, AM and LM conceived and designed the study with input from all authors. LA performed all Arabidopsis experiments, BN-PAGE and immunoblots. CW performed Chlamydomonas physiology experiments and immunoblots. CSL performed Chlamydomonas microscopy imaging and immunoblots. LA, AB, KMS, KAvM, and ED performed Chlamydomonas and Arabidopsis Chl fluorescence and ECS experiments and generated figures. APD and CW performed molecular tracking experiments with input from ML. CL and MB performed xenopus oocyte work. PG and JB did *in silico* structure predictions. TEM contributed to the complementation of Chlamydomonas *bst4* mutant. NA contributed to Arabidopsis transgenic line generation. GP assisted with experimental design and data interpretation. LA and CW generated figures. LA led the writing of the manuscript with input from CW. All other authors provided manuscript feedback.

REFERENCES

Adler L, Díaz-Ramos A, Mao Y, Pukacz KR, Fei C, Mccormick AJ (2022) New horizons for building pyrenoid-based CO_2-concentrating mechanisms in plants to improve yields. Plant Physiol 190: 1609-1627

Aro E-M, Rokka A, Vener A V (2004) Determination of Phosphoproteins in Higher Plant Thylakoids BT - Photosynthesis Research Protocols. *In* R Carpentier, ed,Humana Press, Totowa, NJ, pp 271–285

Atkinson N, Feike D, Mackinder LCM, Meyer MT, Griffiths H, Jonikas MC, Smith AM, Mccormick AJ (2016) Introducing an algal carbon-concentrating mechanism into higher plants: Location and incorporation of key components. Plant Biotechnol J 14: 1302–1315

Atkinson N, Leitão N, Orr DJ, Meyer MT, Carmo-Silva E, Griffiths H, Smith AM, McCormick AJ (2017) Rubisco small subunits from the unicellular green alga Chlamydomonas complement Rubisco-deficient mutants of Arabidopsis. New Phytol 214: 655–667

Atkinson N, Mao Y, Chan KX, McCormick AJ (2020) Condensation of Rubisco into a proto-pyrenoid in higher plant chloroplasts. Nat Commun 11: 6303

Bailleul B, Cardol P, Breyton C, Finazzi G. (2010) Electrochromism: a useful probe to study algal photosynthesis. Photosynthesis research 106:179-89

Bergner SV, Scholz M, Trompelt K, Barth J, Gäbelein P, Steinbeck J, Xue H, Clowez S, Fucile G, Fufezan C, et al (2015) STATE TRANSITION7-Dependent Phosphorylation Is Modulated by Changing Environmental Conditions , and Its Absence Triggers Remodeling of Photosynthetic Protein Complexes 1. Plant Physiol *168*: 615-634

Bharill S, Fu Z, Palty R, Isacoff EY (2014) Stoichiometry and specific assembly of Best ion channels. Proc Natl Acad Sci U S A 111: 6491–6496

Bonente G, Ballottari M,Truong TB, Morosinotto T, Ahn TK, Fleming GR, Niyogi KK, Bassi R (2011) Analysis of LhcSR3, a Protein Essential for Feedback De-

Excitation in the Green Alga *Chlamydomonas reinhardtii* PLoS biology. 2011 Jan 18;9(1):e1000577

Bonente G, Pippa S, Castellano S, Bassi R, Ballottari M. (2012) Acclimation of Chlamydomonas reinhardtii to different growth irradiances. J Biol Chem 287:5833-47

Burlacot A, Dao O, Auroy P, Cuiné S, Li-Beisson Y, Peltier G (2022) Alternative photosynthesis pathways drive the algal CO_2-concentrating mechanism. Nature 605: 366–371

Caspari OD, Meyer MT, Tolleter D, Wittkopp TM, Cunniffe NJ, Lawson T, Grossman AR, Griffiths H (2017) Pyrenoid loss in *Chlamydomonas reinhardtii* causes limitations in CO_2 supply, but not thylakoid operating efficiency. J Exp Bot 68: 3903–3913

Choi BY, Kim H, Shim D, Jang S, Yamaoka Y, Shin S, Yamano T, Kajikawa M, Jin ES, Fukuzawa H, et al (2022) The Chlamydomonas bZIP transcription factor BLZ8 confers oxidative stress tolerance by inducing the carbon-concentrating mechanism. Plant Cell 34: 910–926

Ciorba MA, Heinemann SH, Weissbach H, Brot N, Hoshi T (1999) Regulation of voltage-dependent K^+ channels by methionine oxidation: effect of nitric oxide and vitamin C. FEBS Lett 442: 48–52

Cruz JA, Avenson TJ, Kanazawa A, Takizawa K, Edwards GE, Kramer DM (2005) Plasticity in light reactions of photosynthesis for energy production and photoprotection. J Exp Bot 56: 395–406

Cruz JA, Sacksteder CA, Kanazawa A, Kramer DM (2001) Contribution of electric field ($\Delta\psi$) to steady-state transthylakoid proton motive force (pmf) in vitro and in vivo. Control of pmf parsing into $\Delta\psi$ and ΔpH by ionic strength. Biochemistry 40: 1226–1237

Emrich-Mills TZ, Yates G, Barrett J, Girr P, Grouneva I, Lau CS, Walker CE, Kwok TK, Davey JW, Johnson MP, et al (2021) A recombineering pipeline to clone large and complex genes in Chlamydomonas. Plant Cell 33: 1161–1181

Engel BD, Schaffer M, Kuhn Cuellar L, Villa E, Plitzko JM, Baumeister W (2015) Native architecture of the Chlamydomonas chloroplast revealed by in situ cryo-electron tomography. Elife 4: e11383

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ, Marillonnet S (2014) A Golden Gate Modular Cloning Toolbox for Plants. ACS Synth Biol 3: 839–843

Erickson E, Wakao S, Niyogi KK (2015) Light stress and photoprotection in *Chlamydomonas reinhardtii*. Plant J 82: 449–465

Fei C, Wilson AT, Mangan NM, Wingreen NS, Jonikas MC (2022) Modelling the pyrenoid-based CO_2-concentrating mechanism provides insights into its operating principles and a roadmap for its engineering into crops. Nat Plants 8: 583–595

Genkov T, Meyer M, Griffiths H, Spreitzer RJ (2010) Functional hybrid rubisco enzymes with plant small subunits and algal large subunits: Engineered rbcS cDNA for expression in chlamydomonas. J Biol Chem 285: 19833–19841

Genty B, Briantais JM, Baker NR (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. Biochim Biophys Acta - Gen Subj 990: 87–92

Grefen C, Chen Z, Honsbein A, Donald N, Hills A, Blatt MR (2010) A Novel Motif Essential for SNARE Interaction with the K^+ Channel KC1 and Channel Gating in Arabidopsis . Plant Cell 22: 3076–3092

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies:
Assessing the Performance of PhyML 3.0. Syst Biol 59: 307–321
Hagino T, Kato T, Kasuya G, Kobayashi K, Kusakizako T, Hamamoto S, Sobajima T,
Fujiwara Y, Yamashita K, Kawasaki H, et al (2022) Cryo-EM structures of
thylakoid-located voltage-dependent chloride channel VCCN1. Nat Commun 13:
2505
Hasegawa M, Shiina T, Terazima M, Kumazaki S (2010) Selective excitation of
photosystems in chloroplasts inside plant leaves observed by near-infrared
laser-based fluorescence spectral microscopy. Plant Cell Physiol 51: 225–238
He S, Chou H, Matthies D, Wunder T, Meyer MT, Martinez-sanchez A, Jeffrey PD,
Port SA, He G, Chen VK, et al (2020) The Structural Basis of Rubisco Phase
Separation in the Pyrenoid. 6: 1480–1490
Hennacy JH, Jonikas MC (2020) Prospects for Engineering Biophysical CO_2
Concentrating Mechanisms into Land Plants to Enhance Yields. Annu Rev Plant
Biol 71: 461-485
Herdean A, Teardo E, Nilsson AK, Pfeil BE, Johansson ON, Ünnep R, Nagy G,
Zsiros O, Dana S, Solymosi K, et al (2016) A voltage-dependent chloride
channel fine-tunes photosynthesis in plants. Nat Commun 7: 11654
Horton P, Long SP, Smith P, Banwart SA, Beerling DJ (2021) Technologies to
deliver food and climate security through agriculture. Nat Plants 7: 250–255
Hunter P, Payne-Dwyer AL, Shaw M, Signoret N, Leake MC (2022) Single-molecule
and super-resolved imaging deciphers membrane behavior of onco-
immunogenic CCR5. iScience 25: 105675
Izumi M, Tsunoda H, Suzuki Y, Makino A, Ishida H (2012) RBCS1A and RBCS3B,
two major members within the Arabidopsis RBCS multigene family, function to
yield sufficient Rubisco content for leaf photosynthetic capacity. J Exp Bot 63:
2159–2170
Karlsson J, Ciarke AK, Chen ZY, Hugghins SY, Park Y II, Husic HD, Moroney J V.,
Samuelsson G (1998) A novel α-type carbonic anhydrase associated with the
thylakoid membrane in *Chlamydomonas reinhardtii* is required for growth at
ambient CO_2. EMBO J 17: 1208–1216
Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. Mol Biol Evol 30: 772–
780
Le SQ, Gascuel O (2008) An Improved General Amino Acid Replacement Matrix.
Mol Biol Evol 25: 1307–1320
Lee S, Yoon B-E, Berglund K, Oh S-J, Park H, Shin H-S, Augustine GJ, Lee CJ
(2010) Channel-mediated tonic GABA release from glia. Science 330: 790–796
Lefoulon C, Waghmare S, Karnik R, Blatt MR (2018) Gating control and K^+ uptake by
the KAT1 K^+ channel leaveraged through membrane anchoring of the trafficking
protein SYP121. Plant Cell Environ 41: 2668–2677
Li X, Patena W, Fauser F, Jinkerson RE, Saroussi S, Meyer MT, Ivanova N,
Robertson JM, Yue R, Zhang R, Vilarrasa-Blasi J, Wittkopp TM, Ramundo S,
Blum SR, Goh A, Laudon M, Srikumar T, Lefebvre PA, Grossman AR, Jonikas
MC (2019)*Nature Genetics* 51: 627-35
Long S, Burgess S, Causton I (2019) Redesigning Crop Photosynthesis. Sustain.
Glob. Food Secur. Nexus Sci. Policy. pp 131–141
Ma X, Zhang B, Miao R, Deng X, Duan Y, Cheng Y, Zhang W, Shi M, Huang K, Xia
XQ. (2020) Transcriptomic and Physiological Responses to Oxidative Stress in

a *Chlamydomonas reinhardtii* Glutathione Peroxidase Mutant. Genes (Basel) 11: 463.

Ma Y, Pollock S V., Xiao Y, Cunnusamy K, Moroney J V. (2011) Identification of a novel gene, *CIA6*, required for normal pyrenoid formation in *Chlamydomonas reinhardtii*. Plant Physiol 156: 884–896

Mackinder LCM, Chen C, Leib RD, Patena W, Blum SR, Rodman M, Ramundo S, Adams CM, Jonikas MC (2017) A Spatial Interactome Reveals the Protein Organization of the Algal CO_2-Concentrating Mechanism. Cell 171: 133–147

McGrath JM, Long SP (2014) Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis. Plant Physiol 164: 2247–2261

Meyer MT, Itakura AK, Patena W, Wang L, He S, Emrich-Mills T, Lau CS, Yates G, MacKinder LCM, Jonikas MC (2020) Assembly of the algal CO_2-fixing organelle, the pyrenoid, is guided by a Rubisco-binding motif. Sci Adv 6: eabd2408

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol 37: 1530–1534

Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M (2022) ColabFold: making protein folding accessible to all. Nat Methods 19: 679–682

Mitra M, Mason CB, Xiao Y, Ynalvez RA, Lato SM, and Moroney JV. 2005. The carbonic anhydrase gene families of *Chlamydomonas reinhardtii*. *Canadian Journal of Botany*. 83: 780-795

Mukherjee A, Lau CS, Walker CE, Rai AK, Prejean CI, Yates G, Emrich-Mills T, Lemoine SG, Vinyard DJ, Mackinder LCM, et al (2019) Thylakoid localized bestrophin-like proteins are essential for the CO_2 concentrating mechanism of Chlamydomonas reinhardtii. Proc Natl Acad Sci U S A 116: 16915–16920

Payne-Dwyer AL, Leake MC (2022) Single-Molecular Quantification of Flowering Control Proteins Within Nuclear Condensates in Live Whole Arabidopsis Root BT - Chromosome Architecture: Methods and Protocols. *In* MC Leake, ed,Springer US, New York, NY, pp 311–328

Payne-Dwyer AL, Syeda AH, Shepherd JW, Frame L, Leake MC (2022) RecA and RecB: probing complexes of DNA repair proteins with mitomycin C in live Escherichia coli with single-molecule sensitivity. J R Soc Interface 19: 20220437

Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M, Niyogi KK (2009) An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. Nature 462: 518–521

Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE (2021) UCSF ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci 30: 70–82

Plank M, Wadhams GH, Leake MC (2009) Millisecond timescale slimfield imaging and automated quantification of single fluorescent protein molecules for use in probing complex biological processes. Integr Biol (Camb) 1: 602–612

Price GD, Pengelly JJL, Forster B, Du J, Whitney SM, von Caemmerer S, Badger MR, Howitt SM, Evans JR (2013) The cyanobacterial CCM as a source of genes for improving methylation and chromatin patterning photosynthetic CO_2 fixation in crop species. J Exp Bot 64: 753–768

Qu Z, Cui Y, Hartzell C (2006) A short motif in the C-terminus of mouse bestrophin 4 inhibits its activation as a Cl^- channel. FEBS Lett 580: 2141–2146

Qu Z, Hartzell HC (2008) Bestrophin Cl^- channels are highly permeable to HCO_3^-. Am J Physiol Physiol 294: C1371–C1377

RAVEN J A (1997), CO_2-concentrating mechanisms: a direct role for thylakoid lumen
acidification?. Plant, Cell & Environment, 20: 147-154
Roach T, Na CS (2017) LHCSR3 affects de-coupling and re-coupling of LHCII to
PSII during state transitions in Chlamydomonas reinhardtii. Scientific Reports
7:43145
Roberts SK, Milnes J, Caddick M (2011) Characterisation of *An*BEST1 , a functional
anion channel in the plasma membrane of the filamentous fungus , *Aspergillus*
nidulans. Fungal Genet Biol 48: 928–938
Santhanagopalan I, Wong R, Mathur T, Griffiths H (2021) Orchestral manoeuvres in
the light: Crosstalk needed for regulation of the Chlamydomonas carbon
concentration mechanism. J Exp Bot 72: 4604–4624
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,
Preibisch S, Rueden C, Saalfeld S, Schmid B, et al (2012) Fiji: an open-source
platform for biological-image analysis. Nat Methods 9: 676–682
Shepherd JW, Payne-Dwyer AL, Lee JE, Syeda A, Leake MC (2021) Combining
single-molecule super-resolved localization microscopy with fluorescence
polarization imaging to study cellular processes. JPhys Photonics. 3: 034010
Steen C J, Burlacot A, Short A H, Niyogi K K & Fleming, G R (2022) Interplay
between LHCSR proteins and state transitions governs the NPQ response
in *Chlamydomonas* during light fluctuations. Plant, Cell &
Environment. 45: 2428–2445
Stepien P, Johnson GN (2018) Plastid terminal oxidase requires translocation to the
grana stacks to act as a sink for electron transport. Proc Natl Acad Sci U S A
115: 9634–9639
Sueoka N (1960) Mitotic Replication of Deoxyribonucleic Acid in Chlamydomonas
Reinhardi. Proc Natl Acad Sci 46: 83–91
Syeda AH, Wollman AJM, Hargreaves AL, Howard JAL, Brüning J-G, McGlynn P,
Leake MC (2019) Single-molecule live cell imaging of Rep reveals the dynamic
interplay between an accessory replicative helicase and the replisome. Nucleic
Acids Res 47: 6287–6298
Tardif M, Atteia A, Specht M, Cogne G, Rolland N, Brugière S, Hippler M, Ferro M,
Bruley C, Peltier G, et al (2012) PredAlgo: a new subcellular localization
prediction tool dedicated to green algae. Mol Biol Evol 29: 3625–3639
Tian L, Nawrocki W J, Liu X, Polukhina I, Van Stokkum I H, Croce R (2019) pH
dependence, kinetics and light-harvesting regulation of nonphotochemical
quenching in *Chlamydomonas*. Proc. Natl. Acad. Sci. U. S. A. 116: 8320– 8325
Ueki N, Ide T, Mochiji S, Kobayashi Y, Tokutsu R, Ohnishi N, Yamaguchi K,
Shigenobu S, Tanaka K, Minagawa J, et al (2016) Eyespot-dependent
determination of the phototactic sign in *Chlamydomonas reinhardtii*. Proc Natl
Acad Sci U S A 113: 5299–5304
Van Dijk M, Morley T, Rau ML, Saghai Y. (2021) A meta-analysis of projected global
food demand and population at risk of hunger for the period 2010–2050. Nature
Food 7:494-501
Wollman AJM, Fournier C, Llorente-Garcia I, Harriman O, Payne-Dwyer AL,
Shashkova S, Zhou P, Liu T-C, Ouaret D, Wilding J, et al (2022) Critical roles for
EGFR and EGFR–HER2 clusters in EGF binding of SW620 human carcinoma
cells. J R Soc Interface 19: 20220088
Woo DH, Han K-S, Shim JW, Yoon B-E, Kim E, Bae JY, Oh S-J, Hwang EM,
Marmorstein AD, Bae YC, et al (2012) TREK-1 and Best1 Channels Mediate
Fast and Slow Glutamate Release in Astrocytes upon GPCR Activation. Cell

151: 25–40

Wu A, Brider J, Busch FA, Chen M, Chenu K, Clarke VC, Collins B, Ermakova M, Evans JR, Farquhar GD, et al (2023) A cross-scale analysis to understand and quantify the effects of photosynthetic enhancement on crop growth and yield across environments. Plant Cell Environ 46: 23–44

Yang T, Liu Q, Kloss B, Bruni R, Kalathur RC, Guo Y, Kloppmann E, Rost B, Colecraft HM, Hendrickson WA (2014) Structure and selectivity in bestrophin ion channels. Science 346: 355–359

Zhang R, Patena W, Armbruster U, Gang SS, Blum SR, Jonikas MC (2014) High-throughput genotyping of green algal mutants reveals random distribution of mutagenic insertion sites and endonucleolytic cleavage of transforming DNA. Plant Cell 26: 1398–1409

Supplemental Figures

0.04% CO_2

	Strain	Days 0-3			Days 0 to 5		
		Mean	SE	*P(T<=t)* two-tail	Mean	SE	*P(T<=t)* two-tail
SGR (μ h^{-1})	WT	0.0389	0.0007		0.0241	0.0010	
	bst4	0.0402	0.0003	0.1652	0.0257	0.0001	0.2036
Doubling time (h)	WT	17.8462	0.3380		28.8436	1.2467	
	bst4	17.2379	0.1255	0.1669	26.9798	0.1491	0.2119

3% CO_2

	Strain	Days 0-2			Days 0-3		
		Mean	SE	*P(T<=t)* two-tail	Mean	SE	*P(T<=t)* two-tail
SGR (μ h^{-1})	WT	0.0679	0.0013		0.0456	0.0011	
	bst4	0.0682	0.0010	0.8626	0.0455	0.0003	0.9436
Doubling time (h)	WT	10.2145	0.1930		15.2239	0.3725	
	bst4	10.1669	0.1510	0.8555	15.2353	0.0878	0.9777

Supplemental Table 1 compares specific growth rates (μ h^{-1}) and cell doubling times of WT and *bst4* strains (n=3) during liquid growth assays at 0.04 and 3 % CO_2 (+/- 2 ppm).

Supplemental Figure 1. BST4 structure and sequence compared to other bestrophins. Phylogenetic analysis of the full length Chlamydomonas BST4 amino acid sequence (bold). The evolutionary history of BST4 was inferred by using the maximum likelihood method based on the Le and Gascuel substitution model with discrete Gamma distribution (5 categories) and 500 bootstrap replicates. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

Supplemental Figure 2. Interaction between BST4 and RBCS. A. Yeast-2-hybrid experiment. The disordered region of BST4 C-terminus (amino acid 387-end) and with residues WR from RBMs are changed to EE. Abbreviations: AD, activation domain; BD, binding domain; DDO, double drop out media; TDO, triple drop out media. Growth on TDO indicates an interaction. **B**. FRET efficiency between BST4-mScarlet-I and CrRBCS-Venus. Sensitized-FRET measurement on dual-tagged cells (BST4-Venus and RBCS1-mCherry) showed a moderate FRET efficiency of 35% (n=10).

Supplemental Figure 3. Validation of Chlamydomonas BST4 lines. **A.** PCR amplification of the mapped CIB1 insertion sites and control loci of *bst4* (LMJ.RY0402.159478) and WT control strain (CMJ030 (CC-4533; cw15, mt -) gDNA to confirm the insertion of the CIB1 cassette in *bst4*. **B.** Immunoblots confirming the production of BST4 in WT, *bst4, bst4::BST4* and *bst4::BST-truncated* Chlamydomonas lines. **C.** Immunoblot showing the absence of BST4 in the *bst4* knock-out line and presence in the WT and *bst4::BST4* complemented lines. **D.** Western blot used to quantify the amount of BST4 protein detected in WT and *bst4::BST4.* **E.** Immunoblot was used for the quantification of fluorescent intensity as a measure of BST4 protein abundance. All fluorescent intensity measurements were normalized to respective alpha-tubulin loading controls and conducted in triplicate. No statistical difference was observed between WT and *bst4::BST4* BST4 protein abundance (Paired two-tail t-test *p* = 0.22, n=3)

Supplemental Figure 4. Pyrenoid morphology in bst4 vs WT. A. The pyrenoid area to cell area ratio of WT control strain (black) and *bst4* (white) strains at high (3%) and low (0.04%) CO_2 concentrations. Each bar represents the mean value ± SEM (n=40–50). The data for wild type and *bst4* strains were compared using a two sample T-test. The p-values (showing no significant difference) are displayed above the bars. **B.** Pyrenoid area vs cell area of WT and *bst4* at 0.04% [CO_2].

Supplemental Figure 5. Localization of C-terminally truncated BST4 in WT background. BST4$_{\Delta Cterm}$-mScarlet-I fluorescence shown in green and chlorophyll autofluorescence in magenta. Scale is bar 1 μm.

Supplemental Figure 6. Localization of proteins in a pyrenoid matrix-less background. A. Diagram showing a section of the *Crrbcs::AtRBCS* matrix-less mutant (*-matrix*). The nascent pyrenoid tubules (pt) are shown in dark green, starch granules (sg) in white. **B.** Confocal image of starch binding protein STA2 fused to Venus in *Crrbcs::AtRBCS.* Scale bar is 2 μm. **C.** BST4-Venus in a WT Chlamydomonas cell. Scale bar is 1 μm. **D.** PsaF-mNeon in *Crrbcs::AtRBCS*. Venus and mNeonGreen fluorescence are shown in green and chlorophyll autofluorescence in magenta. Scale bar is 1 μm.

Supplemental Figure 7. Spot test of WT, *bst4* and BST4 complemented lines under CCM induced conditions. *Chlamydomonas* strains were grown in serial dilution on agar plates in saturating light (400 µmol m^{-2} s^{-1}) under a range of CO_2 (+/- 2 ppm) and pH conditions (specified) to induce the CCM. All lines grew comparably to WT across the conditions used and on the low light TAP control plate.

Supplemental Figure 8. Phototaxis and ROS assay. A. TP liquid cultures of Chlamydomonas cells were uniformly distributed on 0.8% (w/v) agar and subjected to directional light (150 μmol m^{-2} s^{-1}). Cell phototaxis was monitored at 0, 1, and 3 h. The assay was performed in the presence of 75 μM ROS hydrogen peroxide (H_2O_2) or 0.3 M ROS quencher N,N'-dimethylthiourea (DMTU). **B.** H_2O_2 assay. Chlamydomonas cells were grown in minimal TP liquid media and exposed to 150 μmol photons m^{-2}s^{-1}. A subset of cells were treated with the quencher DMTU. The concentration of H_2O_2 was subsequently quantified using Amplex Red (n=4), and is presented both proportionately to cell density and chlorophyll content. Different letters indicate significance (p<0.05) as determined by a one-way ANOVA and Tukey's post-hoc test.

Supplemental Figure 9. Chlorophyll fluorescence measurements. A. Y(II) during 5 min illumination and **B.** One minute illumination. **C.** and **D.** Raw fluorescence curves for A. and B., respectively. Curves are normalized to F_o. Genotypes are artificially spaced by 5 s for clarity. **E.** F_v/F_m measurements **F.** Light curve for WT in the presence of bicarbonate. Points represent the mean of three technical replicates ±SEM.

Supplemental Figure 10. NPQ with supplemented bicarbonate. Wild type (WT) and mutants were grown in HS medium at 80 µmol photons m^{-2} s^{-1} and measured at 10 µg Chl ml^{-1}. Cells were supplemented with 500 µM HCO_3^- and then dark adapted for 5 min before the measurements. Dynamics of Non-photochemical photochemical quenching (NPQ) on transition from dark to high light. Kinetics for induction of chlorophyll fluorescence were recorded during 1 min of illumination at 896 µmol photons m^{-2} s^{-1} followed by 5 min in darkness.

Supplemental Figure 11. Electrochromic shift (**ECS) traces across four biological repeats. A.** Cells resuspended in HS at 150 µg Chl ml⁻¹ were dark-adapted for 1 min and then illuminated for 1 min with 890 µmol photons m⁻² s⁻¹ after which the light was switched off to record ECS in darkness. We were unable to calculate the partitioning of the PMF due to non-canonical ECS slow kinetics. **B.** Normalized ECS decay curves. ECS decay of the first 100 ms was fitted to calculate g_H^+ (s⁻¹) =1/time constant for decay. Data are the means of n=3 technical replicates.

Supplemental Figure 12. Chlamydomonas bst4 mutant has an enhanced NPQ and proton conductance under high light and limiting Ci conditions. Wild type (WT) and mutants were grown on TAP medium at 20 µmol photons m^{-2} s^{-1}, resuspended in TP at 30 µg Chl ml^{-1} and exposed for 3 h to light at 150 µmol photons m^{-2} s^{-1}. The cells were dark adapted for 1 h before the measurements. **A.** Maximum quantum yield of photosystem II. **B.** Dynamics of photosynthesis on transition from dark to high light. Kinetics for induction of chlorophyll fluorescence were recorded during 17 min of illumination at 1500 µmol photons m^{-2} s^{-1} followed by 5 min in darkness. Non-photochemical quenching (NPQ) and **C.** Photosystem II quantum yield (Y(II)). **D to F.** ECS decay kinetics were performed on cells pre-exposed for 10 min to high light and the **D.** total PMF, **E.** g_{H^+}, and **F.** total H^+ flux (v_{H^+}) were determined as described in Methods. Data are the means ± SEM (n=3 replicates). **G.** Immunoblot of NPQ protein LHCSR3 in each genotype compared to tubulin after exposure of cells to 3 h 150 µmol photons m^{-2} s^{-1} in TP. **H.** Fluorescence of LHCSR3 protein band normalized to α-tubulin. Data are the means ± SEM (n=3 replicates). Different letters indicate statistically significant difference among the genotypes (one-way ANOVA test, followed by Tukey's post hoc test, P < 0.05). **I.** Representative Electrochromic shift (ECS) curves used to determine PMF values in D. **J.** Representative ECS decay curves. To determine the gH+ parameter in E., ECS kinetics were recorded during 600 ms dark intervals. The ECS decay of the first 100 ms was fitted to calculate gH+ (s^{-1}) =1/time

A

B

α-BST4

Supplemental Figure 13. No currents were detected for BST4 with any anions tested in Xenopus oocytes
A. Steady state currents analysis of oocytes injected with BST4, full length or truncated, compared to non-injected oocytes. The voltage steps start from 60 mV to -100 mV (Cl⁻, PEP⁻, and Gluconate conditions) 40 mV to -120 mV (HCO3⁻ conditions, right panel), holding voltage is -20 mV. Recordings were performed on n>4 oocytes. There is no differences between the currents recorded in non-injected oocytes and expressing the protein. **B.** Western Blot analysis of oocytes. Lane 1 is the marker, lane 2 are non-injected oocytes, lane 3 is the full-length protein (about 66 kDa), lanes 3 and 4 are the truncated proteins. We can only detect full length BST4 as the antibody is directed against the C-terminal part of the protein that is removed in the two truncated versions of the channel.

Supplemental Figure 14. Phenotype of BST4 transgenic Arabidopsis line. A. Immunoblot against BST4 of proteins extracted from BST4 no tag lines and Azygous segregants (Az). S2$_{Cr}$ is the parent line. **B.** Fresh weight of 28-day old rosettes, bars represent mean weight for each genotype n=18-21. Different letters indicate statistically significant difference among the genotypes (one-way ANOVA test, followed by Turkey's post hoc test, P < 0.05).

Supplemental Figure 15. Photosynthetic measurements of BST4 transgenic Arabidopsis lines 2 and 3. A. F_v/F_m values measured on attached, 30 min dark adapted leaves of 8-week-old plants (n=15-21). The letters in indicate nonsignificant differences between plants expressing BST4 and their azygous segregants using Tukey post hoc test (P > 0.05). **B.** Non-photochemical quenching (NPQ) as a measure of photoprotection and **C.** photosystem II quantum yield (Y(II)) were recorded during 10 min of illumination at 830 µmol photons m^{-2} s^{-1} followed by a 5-minute dark period. Data are presented as means ± SEM (n=4-6). Asterisks indicate statistical difference between plants expressing BST4 and their Azygous (Az) segregants according to unpaired t-test (P ≤ 0.05). **D.** Proton motive force (PMF) size and **E.** partitioning to pH gradient (ΔpH) after 3 min illumination at 830 µmol photons m^{-2} s^{-1}. Data are means ± SEM (n=5-6). The letters in E and F indicate non-significant differences between plants expressing BST4 and their azygous segregants according to unpaired t-test (P > 0.05).

A. Full length BST4 and homologous sequences

```
>Brassica_rapa_(PAC:30641593)
--------------------------------------------------------------------------------
------------------------------------------------------------------------------
RTLYTHEKWVEHRSSLRHVHHLFSSFSSRVILSLIPPVFFFTSVAIFIASYNSAVAL-------------
DWLPSVFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--DDLARQVICSVDGSG--
---------------DELVIK-----DLLLRYV----AAFPVALKCHVTYGSD-----VARDLRN-----
LIEGDDLSLI--IESKHR------------PRCVIEFISQSLQL----LKLDDT--------------
KRDLLESKMLHLHEGIGVCEQLMGIPIPLAYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSES---------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
-----------------------------------
>Q9M2D2_-_VCCN1_Athaliana
MY-----------------QSMNLSVSSNFTHRSLLES-----------RFPI-----FSTGFR-----
KSVNLKPPRVSSGPE---SNDSGH--------------ETLTDKLIHLLRAVP--------
DWADEIKERGMQQKRSLYTHEKWVEHRSSLRHVRHLLSSFSSRVILSLIPPVFFFTSVAVVIASYNSAVAL----
---------DWLPGIFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
NDLARQVICSVDSSG----------------DELIIK-----DLLLRYI----AAFPVALKCHVIYGSD-----
IARDLRN-----LIEADDLSLI--LQAKHR------------PRCVIEFISQSIQL----LKLDDA--------
------KRDLLESKMLHLHEGIGVCEQLMGIPIPLSYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSEKVIRNRIIAKIKLHEFK-
-----HSS-----------------------NGRHRS-----------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
-----------------------------------
>Nicotiana_tomentosiformis_(XP_009629643.1)
M------------------------TNSRTLFSIQSPTNASFSSHF-TLKTPSK--LQQQSFPSKL-
NFKKLRFSTFKVRCCPQ----QTPQN--------------QNPTSALISILRIIP--------
DWADRIQEEGMKKKRSLYTHESWMQHRSSLRHVRHLFSSLNSRVILSLVPPVIAFTSVAVVIASYNSAVSM----
---------HWLPELFPVLR---ASPLPYQLTAPALALLLVFRTEASYSRFETGKKAWTKVIAGT--
NDFARQVIACVDKS-----------------DAVLK-----AALLQYI----MAFPVALKCHITYGSD-----
IASDLKN-----LLEADDLAVV--LSSKHR------------PRCIIGFISQCLQS----LHLEGT--------
------KLTQLESKISCFHEGIGVCEQLAGIPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCQL-VHDNIQESMANEKKIQERLSAKRKRRFSE-
-----HSQ-----------------------NGWPTS-----------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
-----------------------------------
>XP_028961536_Malus_domestica
MLLPSSLSLQTLAPPNADTIQKPHQTLPQNLTLFQLVHP------QHFPTLQFPN-----LPSGPKTL-
KFKLLCSQSPNPNPSP-------PSS--------------SSPVQTLISILRIIP-------
DWSDRTQERGMRQHRTLYDHEKWMHHRSSYRHLRHLLSSLSSRVILSLIPPVIAFTLVAVVIASYNTAVAL----
---------DLLPGIFPLLR---SSSLPYQLTAPALALLLVFRTEASYSRFEEGRKAWTEVIAGA--
NDFARQIISSVETSG----------------DAQLK-----KALLQYI----VAFPVALKCHVIYGSD-----
IARDLQN-----LLEVDDLLVV--LNSKHR------------PGCIIQFISRSLQL----LKLEES--------
------RRIMLQSKISCFHEGIGICEQLIGTPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEQVGVLIEEPFPM----LALDDLCNS-VRNNVQEALASEKLIRARLAAKGRIQSEQ-
-----QFQ-----------------------NGQPRP-----------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
--------------------------------------------------------------------------------
-----------------------------------
>Chlamydomonas_eustigma_(GAX83184.1)
```

```
M--------------------------------------------------------------
LVHRVHTRSLGNRNQCGRK--------------------LHRVSTFVVKTPTEKPVVA----
DYVLPRSEEARRYFRTVYDFPQWQKHRSPTRLIDRLLQIPRSHVLQNILPSIAWCSSVAGLLTLYMQAYDA----
---------HILPDGFPSFATNNACTSFVNTTTVALSLLLVFRTNVSYGRWDEARKMKGLLVNRS--
RDLMRQVCAMVPEE-----------------DVATK-----AMMAKWT----AAFCRVLRIHFQPEVS----
LEDEMKG-----LLSPEELEWL--IESKHR------------PCSVIHMLSQIIYD----SQISAI--------
------CQAQMCNNLTAFEDVLGGCERLLRAPIPVSYTRH-TARFLFTWLTLL-------PFALYNSC-----
GVWTLPVVAGVSAVLCGIEEIGVQIEEPFGI----LPLEAICGR-IQADVMATLKEDAKTRNLRNKVLLIRGPE-
-----HWAMV----------------------NGHSNK--------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
------------------------------------
>Cre16.g662600.t1.2_(BST1)
M----------------------------QMQA-----------
NRSSLRASPVRGLGARPLLRALPAGRVARLNVSAQA--KDPNAPIQ---------------
SNPLGTLSSQSGQVA----------
TLPRSEEARKYFRTVYDFPQWQKHRSSYRFAERLFQLSQSHILQNALPAISWVTLVATLVASYGYSYDQ------
-------HMLPDVFPSISPNASCTAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-----------------QVEAK-----KALARWT----VAFSRALRIHFQPEVT----
IESELQN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VPMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYPTT-----
GWGVVPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-
IQADVMATLKDDADTKTILAEAGLISLIPS----ATSATPVASA----EPVLVSARPSAAPAPNNGLQVRVAM--
------------------------------------------------------------------GG-
-------------------ERK*---------------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
------------------------------------------------------------
>Cre16.g663400.t2.1_(BST2)
M-------------------------------QC-------------LSSRPVAMGRAGSSALPRL-
PLRAGRVCHLGVRCQAANKDPNAPIQ---------------SNPLGSFSSQLQNQP----------
TLPRSEEARKYFRTVYDFPQWQTHRNQYRLMKRLFSIPQSHVIQNALPSIMWVAFTSTCVAAYMYGYDQ------
-------HMLPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-----------------QVEAK-----KALARWV----VAFSRALRIHFQPEVT----
IESELKN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VPMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCSGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDDADTKTILAEAGLISLRA-
-----NSAMAVENALPDLDSINAAAPNGNGSHNGNGAAV-------------------------------
------------------------------------------------------------PVS-----
-------------------VSAGA--------------------------------------
--------------------------------------------------------------
---------------------------------------------------------SGNGMNV---
----------------------RISPR*-----
>Cre16.g663450.t1.2_(BST3)
M-------------------------------QVSK-----------VPSS------ASARCLPRL-
PVRTSRVCQLSVRCQAANKDPNAPIQ---------------SNPLGSFSSQNSSGAVV--------
TAPRNEDARKYFRTVYDFPQWQKHRSQSRLVRRLFTIPQSHVIQNALPSIMWVTFTSTCVAAYMYGYDL------
-------HILPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-----------------QVEAK-----KALARWT----VAFARALRIHFQPEVT----
IESELQN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VRMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDDADTKTVLAEAGLISLIP-
-----SMSLPPTEHASPSDPVTAAAAAALAAANGNGAASHSNG-------------------------
------------------------------------NG--------------------
SKPVSTQVPP----------------PVLAPVTVS-------------------------------
--------------------------------------------------------------
--------------------------------------------------------------
SSGSMNV------------------------RISPR*-----
>Chlorella_sorokiniana_(PRW33726.1)
M---------------------STAMLAGSRIQLQQPAG--------LGGSRLQ--RAAAPVAAAA-
RLGSVRPAGLQARSTAARRADRSALR---------------VSATASPEAAPVKLS----------
```

```
GDDLKEANRKHMRSVFDFDLWKKHRSSSRYLRHIVGLGESRIVSGLMAPLTYVMTLSLAVACYNAAAEA------
-------GYLP-VFPELKL--ATNAPFGLTSFALSLLLVFRTNSSYGRWDEARKMWGLIVNRS--
RDFIRQGLGYIPPE------------------QEELQ-----KMLVRWT----VAYSRSLMCHLRPGED----
LRVELKD-----TLKPEELEAL--LASTHR------------PNYVVQVLTAIIKT----
AQLPAAVTNNRDSTGCVPAGAAYRMDENLTVFADVTGGCERILRTPIPLSYTRH-TSRFMMIWLTLL------
PFTLWDSC-----HWAMLPIAGIVSFLLLGIEEIGVQIEEPFTI----LPLEVISRT-IEGNVWELYRMHSGEA-
--LEKEQAELAN------GQDVQVLNA----QDLVALMAPSAVGNTANGTSRKSLV-----------------
VNYGL----------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
--------------------------------------------------
>Cre06.g261750.t1.2_(BST4)
M----------------------QC-------QLKH---------------------------GA-
RPQSQRPNWLPARAATLRPAVQHGVR--------------RGLTLGVKAAAAPLE--------
DKKMPADMTTRQYRRVVYDFALWAKHRDVNRYLYNLRTIPGSRIIRQLSQPMGVVLAWAALFGFYETCLEA----
---------GVLPSYLPKMTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-----------------DQAGR-----EAVGKWT----VGFTRALQAHLQEDID-----
LRKELEKA--TPRWSKEEIDML--VNAQHR------------PIKAISVLSELTRQ----LSITQF--------
------QALQMQENCTFFYDALGGCERLLRTPIPVSYTRH-TARFLTIWLAML-------PLGLWERY-----
HWSMLPVIALIGFLLLGIDEIGISIEEPFGI----LPLDAICGR-AQTDVNSLLKEDPAVMKYVDDVRSGRVKS-
-----PPPLPPAPA----APAAA-------AAAAAAAARSVSP----
QPDVAKTLGSLFTNVRAGVGAVAPGAPLMPQAPVRSPSPTRSVSP--------SFPRASAGTGMPPPVGMNG---
-------ATPRVAAAPPTPPPVSRPAA-----------------
PAAAPAAGSGFTMPNFSASLSGLTGGAAAAAKSAADAASSKLTKMADSMSSGAAAPAPPAAPAR-----------
----------PSTSPRPSASSPISSSADADRSDSSRR---PVNWRDELQSLKATRE---------------
PNGNGNGSGVAPAA--GRADADEEALRRFGNLAGRSR-------------SGNG---------------
GGGSSDTELSEANRPRTRPDWRNQL*
>XP_002945810.1_Volvox_carteri_f._nagariensis
M----------------------QS-------QLQP---------------------------RL-
QLQGTRLNWLPQRSCVQRRSLRVDAT--------------SG------AAPPPPA--------
GKELSNDMVTRQYRRTVYDFSLWAKHRDVNRYLYNLKTIPGSRIIRTLGQPMGIVLAWAAMFGFYETCLES----
---------GVLPSYFPKLTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-----------------DVAGR-----EAVGKWA----VGFCRALQAHLQEDAN-----
LREELQKA--QPRWSREEIDML--CSAQHSWQQLQSCVNAFW-PIKAISMLSELTRQ----LPISQF--------
------QALQMQENVTFFYDALGGCERLLRTPIPVSYTRI--------------------------------
----------------------------LPLDAICTR-AQTDVVSLLKDDPAVVKYISDVRQGRIAP---
---PTEPPVAGA----
APVAAAPPPPPPASAGGGISRSGSPTAQQQPDVMKTVTSMLHNVKAGIGAVAPAPPRPPSPQPRARSP-
RAASPGGPS----PFPRASAGTG-------G----------
AAAAVPSPPPIKPLTSSSSSSSGAVSKDSNNSTATAKKPASAPAASSA----GFSMGFSGLADGAAAAAKSAS-
AAAAKFSKIADSVVAG--TPAAPASEAKRETAA--AAAMQAQPRN---
TPSSSSSTPSAAPANGSSDDDRSSSGRRTAAAVNWREELAALRAGREDAEEPASASASYDREFPSSSWSFSSASS
AAVVQSGDAEDEARRRFGGLAGRGARSDTTTSAAAVMRGNGNGLSENGYGNGYGNDN-
GNGNGNTVEARGARPRTRPDWRNQL-
>Chrysochromulina_sp._(KOO32217.1)
M---------------------REHPLSYEEYMRQRS----------AGRDPLA--EAVQGQSASM-
ERGVVPPPPVKATPERTEAAFTPPAEFDFFQDVFKPTVESVVKAVVSPGAQQDSDESYMRVPWWEQGSTYSEDQR
KDRRTVFMHDDWKRHRSSERFFRNIKTWPSSGINQALRKELTFVTSVSVFVVLANMLLYQYQDFGGVVHPGPLSF
LDGPIKSLS---LPALPFSMASPVLSLLLVFRTNTAYFRWNEARTLWGGLINNC--RNIVRQTTTMFPNDA----
------------YHNALK-----KRLATET----ATFIKSLRNFLRGPEDDAT--
LRKELYAYVNQGLMTSAQAEAT--LAAKNR------------PMFALAAMSATLRK----ANIDEM-------
------YISRMDSTISVLVDLTGANERIFKSPIPLVYTRL-TARFLSVFLTLL-------
PLAMWAALGESWNHWATIPATFILSVFLFGVEEVGIQIEEPFSI----LPLEAMCNGAIEAVQLEMLAAE-----
-----------------QSQVFEAAG----DAVAVA--------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
--------------------------------------------------
>Emiliania_huxleyi_(XP_005770556.1)
----------------------------------------------------------------------------
--------------------------------------------------------------
```

```
EFKEAGREFRQDVYSYNDWRWHRESGHIASAISSVFTSGVGKAMWRETFFVIATAAAVYLYNIGVPVLAAKTAAS
L----PIVAALLGRLPLLHLSLLPLTLSSPALFLLLVFRTNNSYDRWWEARKVWGGVINAS--
RDLARQALALVR-------------------DAELK-----KLMVSQI----
ASYARVLKYHLGPPTPEARDLLRNELVD----NRLPADQVRVI--MEAKHK------------
PMALLGLVSASLHDSGR-TGLDTV--------------QASKLDQTLSLLTDYLGKCERIVKTPLPLVYTRH-
TARFLSWWLLFL-------PVCLYNQLRA---NWMIVPVSGLIGFFLVGIEDLGNQIEEPFSI----
LPLTAMGTG-IQQSIFEAL---------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
>Phaeodactylum_tricornutum_(XP_002180738.1)
MM----------------------------------------------------RNFASVLLLL-
SSGAAAFAPVQHNGVRTIATPSTPLY--------------GNTKQPPALP--PIK---------
DISYGEESRKYRRTVYSHDDWVKHRSSDRFLRNLLAIGSSGVYKSLAKEVLATTGVATFIVLYNCLVGGYTDLEG
IKHS---ALIESVWAPLMA--LPLAPFTLSSPSLGLLLVFRTNTSYQRWDEARKNWGMNINHT--
RDLVRMGTSFYDNAA----------------VSSEQRAKDL-KALSLAT----WSFVRAMKRHLSPESEDEQD-
FRRELFE-----RLPAPQAQAI--IDAAHR------------PNRALFDLSVAIEN----LPMHFL--------
------RKNQVHQAVTIFEDNLGSSERLLTSPVPLFYSRH-TARFLSFWLLLL-------
PFALWDPFAGTWNHVGMIPATAVISIFLFGIEELATQMEEPFTI----LPMQAFCDK-
IGNWCNEIVSWQAGDNGMAVNMPSMISPEGLPELKEPAPVPAMA----VASVAAAMPVMANGDINGDTTGITM--
-DQP-------------------------------------------------HNAIP----------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
---------------------------------------------------------------
>Thalassiosira_pseudonana_(XP_002289965.1)
M---------------------------------------------------------------------------
----------------------------------------GPPIDPSVPVT-----------
DQVGEGSRKYRRTVYTHDDWVRHRSPDRFGNNLSTLFNSGIYKQVANEVFATTAVATFVFLWNMIAGGYTDLAGV
QHG---PIIDSPLAQMVG--LPMTAFTILTPSLGLLLVFRTNTSYGRWDEARKMWGLNINHT--
RDLNRMATAWYGNEGNMDSVAFMGGDIPYSQPIDPVQRAYDL-GQVSLFT----WAFVRSMKRHLSPPEEDEED-
FKAELRA-----RLTPEQAENI--INAAHR------------PNRALFDLSVAIEN----LPMHFL--------
------RKNAINTNLSIFEDTLGGCERLLSSPVPLFYSRH-TARFLSTWLLLL-------
PFGLYEQFKDSWNHIAMIPATAFISVCLFGIEELATQLEEPFTI----LPMQGFCDK-IGGWCDEIVSW-
AGQGQQEYTEENAMSNE------QEMTY---------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
---WR-----------------------------------------------------------------------
-----------------------------------------------------------
>WP_049046555.1_Kiebsiella_aerogenes
----------------------------------------------------------------------------
----------------------------------------------------------------------------
------MIIRPEQHWFFRLFDWHGSVLSKIVFRLLLNVLMSVIAIISYQWYEQL-------------GI------
-----HLTVAPFSLLGIAIAIFLGFRNSASYSRFVEARNLWGTVLIAE--RTLVRQLKNILPD-----------
--------DEETH-----KTLVSYL----VAFSWSLKHQLRK-TD-----PAVDLYR-----LLPKEKVAEI--
LASSMP------------TNRILLLIGNELGRLREQGKLSDI--------------
TYGLMDNKLDELAHVLGGCERLASTPVPFAYTLI-LQRTVYLFCTLL-------PFALVGDL-----
HYMTPFVSVFISYTFLSWDSLAEELEDPFGTSANDLPLNAMCNT-IERNLMDMTGQHPLPEKMQPDRYYNLT---
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------------------------------------------------
----------------------------------
>XP_011543531.1_Homo_sapiens
M---------------------------------------------------------------------------
----------------------------------------------------------------------------
TITYTSQVANARLGSFSRLLLCWRGSIYKLLYGEFLIFLLCYYIIRFIYRLALTEEQQL-------------
MFEKLTL--YCDSYIQL--IPISFVLGFYVTLVVTRW------WNQYENLPWPDRLMSLVSGFVEGK--------
-----------DEQGRL--LRRTLIRYANLGNVLILRSVSTAVYKRFP-----SAQHLVQ---
```

```
AGFMTPAEHKQLEKLSLPHNM---------FWVPWVWFANLSMKAWLG---GRIRDPI------------
LLQSLLNEMNTLRTQCGHLYAYDWISIPLVYTQVVTVAVYSFFLTCLVGRQFLNPAKAYPGHEL---
DLVVPVFTFLQFFFYVGWLKVAEQLINPFGEDDDDFETNWIVDRNLQVSLLAVDEMHQDLPRMEPDMYWNKPEP-
-----QPPYTAASA----QFRRASFMGSTFNISLNKEEMEFQPNQEDEED-----------
AHAGIIGRFLGLQSHDHHPPRANSRTKLLWPKRESLLHEGLPKNHKAAKQ----
NVRGQEDNKAWKLKAVDAFKSAPLYQRPGYYSAPQ-----------TPLSPTPMFFPLEPSA---
PSKLHSVTGI-DTKDKSLKTVSSGAKKSFELLSES--
DGALMEHPEVSQVRRKTVEFNLTDMPEIPENHLKEPLEQSPTNIHTTLKDHMDPYWALENR----------
SVLHLNQGHCIALCPTPASLALSLPFLHNFLGFHHCQSTLDLRPALAWGIYLATFTGILGKC------------
--SGPFLTSPWY---HPEDFLGPGEGR------------------
```

B. BST4 and homologous sequences trimmed after the bestrophin domain

>Brassica_rapa_(PAC:30641593)
```
----------------------------------------------------------------------
----------------------------------------------------------------------
RTLYTHEKWVEHRSSLRHVHHLFSSFSSRVILSLIPPVFFFTSVAIFIASYNSAVAL------------
DWLPSVFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--DDLARQVICSVDGSG--
--------------DELVIK-----DLLLRYV----AAFPVALKCHVTYGSD-----VARDLRN-----
LIEGDDLSLI--IESKHR------------PRCVIEFISQSLQL----LKLDDT-------------
KRDLLESKMLHLHEGIGVCEQLMGIPIPLAYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSE
```
>Q9M2D2_-_VCCN1_Athaliana
```
MY----------------QSMNLSVSSNFTHRSLLES------------RFPI-----FSTGFR-----
KSVNLKPPRVSSGPE---SNDSGH--------------ETLTDKLIHLLRAVP--------
DWADEIKERGMQQKRSLYTHEKWVEHRSSLRHVRHLLSSFSSRVILSLIPPVFFFTSVAVVIASYNSAVAL----
---------DWLPGIFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
NDLARQVICSVDSSG----------------DELIIK-----DLLLRYI----AAFPVALKCHVIYGSD-----
IARDLRN-----LIEADDLSLI--LQAKHR------------PRCVIEFISQSIQL----LKLDDA--------
------KRDLLESKMLHLHEGIGVCEQLMGIPIPLSYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSE
```
>Nicotiana_tomentosiformis_(XP_009629643.1)
```
M-----------------------TNSRTLFSIQSPTNASFSSHF-TLKTPSK--LQQQSFPSKL-
NFKKLRFSTFKVRCCPQ----QTPQN--------------QNPTSALISILRIIP--------
DWADRIQEEGMKKKRSLYTHESWMQHRSSLRHVRHLFSSLNSRVILSLVPPVIAFTSVAVVIASYNSAVSM----
---------HWLPELFPVLR---ASPLPYQLTAPALALLLVFRTEASYSRFETGKKAWTKVIAGT--
NDFARQVIACVDKS------------------DAVLK-----AALLQYI----MAFPVALKCHITYGSD-----
IASDLKN-----LLEADDLAVV--LSSKHR------------PRCIIGFISQCLQS----LHLEGT--------
------KLTQLESKISCFHEGIGVCEQLAGIPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCQL-VHDNIQESMANE
```
>XP_028961536_Malus_domestica
```
MLLPSSLSLQTLAPPNADTIQKPHQTLPQNLTLFQLVHP------QHFPTLQFPN-----LPSGPKTL-
KFKLLCSQSPNPNPSP-------PSS--------------SSPVQTLISILRIIP--------
DWSDRTQERGMRQHRTLYDHEKWMHHRSSYRHLRHLLSSLSSRVILSLIPPVIAFTLVAVVIASYNTAVAL----
---------DLLPGIFPLLR---SSSLPYQLTAPALALLLVFRTEASYSRFEEGRKAWTEVIAGA--
NDFARQIISSVETSG----------------DAQLK-----KALLQYI----VAFPVALKCHVIYGSD-----
IARDLQN-----LLEVDDLLVV--LNSKHR------------PGCIIQFISRSLQL----LKLEES--------
------RRIMLQSKISCFHEGIGICEQLIGTPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEQVGVLIEEPFPM----LALDDLCNS-VRNNVQEALASE
```
>Chlamydomonas_eustigma_(GAX83184.1)
```
M---------------------------------------------------------------
LVHRVHTRSLGNRNQCGRK--------------------LHRVSTFVVKTPTEKPVVA----
DYVLPRSEEARRYFRTVYDFPQWQKHRSPTRLIDRLLQIPRSHVLQNILPSIAWCSSVAGLLTLYMQAYDA----
---------HILPDGFPSFATNNACTSFVNTTTVALSLLLVFRTNVSYGRWDEARKMKGLLVNRS--
RDLMRQVCAMVPEE------------------DVATK-----AMMAKWT----AAFCRVLRIHFQPEVS-----
LEDEMKG-----LLSPEELEWL--IESKHR------------PCSVIHMLSQIIYD----SQISAI--------
------CQAQMCNNLTAFEDVLGGCERLLRAPIPVSYTRH-TARFLFTWLTLL-------PFALYNSC-----
GVWTLPVVAGVSAVLCGIEEIGVQIEEPFGI----LPLEAICGR-IQADVMATLKED
```
>Cre16.g662600.t1.2_(BST1)
```
M-----------------------------QMQA-----------
NRSSLRASPVRGLGARPLLRALPAGRVARLNVSAQA--KDPNAPIQ--------------
```
```
                                  64
```

```
SNPLGTLSSQSGQVA----------
TLPRSEEARKYFRTVYDFPQWQKHRSSYRFAERLFQLSQSHILQNALPAISWVTLVATLVASYGYSYDQ------
-------HMLPDVFPSISPNASCTAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD------------------QVEAK-----KALARWT----VAFSRALRIHFQPEVT----
IESELQN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VPMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYPTT-----
GWGVVPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Cre16.g663400.t2.1_(BST2)
M-----------------------------QC-------------LSSRPVAMGRAGSSALPRL-
PLRAGRVCHLGVRCQAANKDPNAPIQ--------------SNPLGSFSSQLQNQP----------
TLPRSEEARKYFRTVYDFPQWQTHRNQYRLMKRLFSIPQSHVIQNALPSIMWVAFTSTCVAAYMYGYDQ------
-------HMLPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD------------------QVEAK-----KALARWV----VAFSRALRIHFQPEVT----
IESELKN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VPMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCSGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Cre16.g663450.t1.2_(BST3)
M-----------------------------QVSK-----------VPSS------ASARCLPRL-
PVRTSRVCQLSVRCQAANKDPNAPIQ--------------SNPLGSFSSQNSSGAVV--------
TAPRNEDARKYFRTVYDFPQWQKHRSQSRLVRRLFTIPQSHVIQNALPSIMWVTFTSTCVAAYMYGYDL------
-------HILPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD------------------QVEAK-----KALARWT----VAFARALRIHFQPEVT----
IESELQN-----ILTPAELQML--AKSQHR------------PVRAIHAISQIIQS----VRMSSI--------
------HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Chlorella_sorokiniana_(PRW33726.1)
M----------------------STAMLAGSRIQLQQPAG--------LGGSRLQ--RAAAPVAAAA-
RLGSVRPAGLQARSTAARRADRSALR--------------VSATASPEAAPVKLS----------
GDDLKEANRKHMRSVFDFDLWKKHRSSSRYLRHIVGLGESRIVSGLMAPLTYVMTLSLAVACYNAAAEA------
--------GYLP-VFPELKL--ATNAPFGLTSFALSLLLVFRTNSSYGRWDEARKMWGLIVNRS--
RDFIRQGLGYIPPE------------------QEELQ-----KMLVRWT----VAYSRSLMCHLRPGED-----
LRVELKD-----TLKPEELEAL--LASTHR------------PNYVVQVLTAIIKT----
AQLPAAVTNNRDSTGCVPAGAAYRMDENLTVFADVTGGCERILRTPIPLSYTRH-TSRFMMIWLTLL------
PFTLWDSC-----HWAMLPIAGIVSFLLLGIEEIGVQIEEPFTI----LPLEVISRT-IEGNVWELYRMH
>Cre06.g261750.t1.2_(BST4)
M--------------------QC--------QLKH----------------------------GA-
RPQSQRPNWLPARAATLRPAVQHGVR--------------RGLTLGVKAAAAPLE--------
DKKMPADMTTRQYRRVVYDFALWAKHRDVNRYLYNLRTIPGSRIIRQLSQPMGVVLAWAALFGFYETCLEA----
----------GVLPSYLPKMTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE------------------DQAGR-----EAVGKWT----VGFTRALQAHLQEDID-----
LRKELEKA--TPRWSKEEIDML--VNAQHR------------PIKAISVLSELTRQ----LSITQF--------
------QALQMQENCTFFYDALGGCERLLRTPIPVSYTRH-TARFLTIWLAML-------PLGLWERY-----
HWSMLPVIALIGFLLLGIDEIGISIEEPFGI----LPLDAICGR-AQTDVNSLLKED
>XP_002945810.1_Volvox_carteri_f._nagariensis
M--------------------QS--------QLQP--------------------------RL-
QLQGTRLNWLPQRSCVQRRSLRVDAT--------------SG------AAPPPPA--------
GKELSNDMVTRQYRRTVYDFSLWAKHRDVNRYLYNLKTIPGSRIIRTLGQPMGIVLAWAAMFGFYETCLES----
----------GVLPSYFPKLTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE------------------DVAGR-----EAVGKWA----VGFCRALQAHLQEDAN-----
LREELQKA--QPRWSREEIDML--CSAQHSWQQLQSCVNAFW-PIKAISMLSELTRQ----LPISQF--------
------QALQMQENVTFFYDALGGCERLLRTPIPVSYTRI--------------------------------
-----------------------------LPLDAICTR-AQTDVVSLLKDD
>Chrysochromulina_sp._(KOO32217.1)
M--------------------REHPLSYEEYMRQRS----------AGRDPLA--EAVQGQSASM-
ERGVVPPPPVKATPERTEAAFTPPAEFDFFQDVFKPTVESVVKAVVSPGAQQDSDESYMRVPWWEQGSTYSEDQR
KDRRTVFMHDDWKRHRSSERFFRNIKTWPSSGINQALRKELTFVTSVSVFVVLANMLLYQYQDFGGVVHPGPLSF
LDGPIKSLS---LPALPFSMASPVLSLLLVFRTNTAYFRWNEARTLWGGLINNC--RNIVRQTTTMFPNDA----
-------------YHNALK-----KRLATET----ATFIKSLRNFLRGPEDDAT--
LRKELYAYVNQGLMTSAQAEAT--LAAKNR------------PMFALAAMSATLRK----ANIDEM--------
------YISRMDSTISVLVDLTGANERIFKSPIPLVYTRL-TARFLSVFLTLL-------
PLAMWAALGESWNHWATIPATFILSVFLFGVEEVGIQIEEPFSI----LPLEAMCNGAIEAVQLEMLAAE
>Emiliania_huxleyi_(XP_005770556.1)
--------------------------------------------------------------------------
```

```
----------------------------------------------------------------
EFKEAGREFRQDVYSYNDWRWHRESGHIASAISSVFTSGVGKAMWRETFFVIATAAAVYLYNIGVPVLAAKTAAS
L----PIVAALLGRLPLLHLSLLPLTLSSPALFLLLVFRTNNSYDRWWEARKVWGGVINAS--
RDLARQALALVR-------------------DAELK-----KLMVSQI----
ASYARVLKYHLGPPTPEARDLLRNELVD----NRLPADQVRVI--MEAKHK-------------
PMALLGLVSASLHDSGR-TGLDTV-------------QASKLDQTLSLLTDYLGKCERIVKTPLPLVYTRH-
TARFLSWWLLFL-------PVCLYNQLRA---NWMIVPVSGLIGFFLVGIEDLGNQIEEPFSI----
LPLTAMGTG-IQQSIFEAL---
>Phaeodactylum_tricornutum_(XP_002180738.1)
MM--------------------------------------------------------RNFASVLLLL-
SSGAAAFAPVQHNGVRTIATPSTPLY--------------GNTKQPPALP--PIK---------
DISYGEESRKYRRTVYSHDDWVKHRSSDRFLRNLLAIGSSGVYKSLAKEVLATTGVATFIVLYNCLVGGYTDLEG
IKHS---ALIESVWAPLMA--LPLAPFTLSSPSLGLLLVFRTNTSYQRWDEARKNWGMNINHT--
RDLVRMGTSFYDNAA----------------VSSEQRAKDL-KALSLAT----WSFVRAMKRHLSPESEDEQD-
FRRELFE-----RLPAPQAQAI--IDAAHR------------PNRALFDLSVAIEN----LPMHFL--------
------RKNQVHQAVTIFEDNLGSSERLLTSPVPLFYSRH-TARFLSFWLLLL-------
PFALWDPFAGTWNHVGMIPATAVISIFLFGIEELATQMEEPFTI----LPMQAFCDK-IGNWCNEIVSWQ
>Thalassiosira_pseudonana_(XP_002289965.1)
M---------------------------------------------------------------
------------------------------------GPPIDPSVPVT-----------
DQVGEGSRKYRRTVYTHDDWVRHRSPDRFGNNLSTLFNSGIYKQVANEVFATTAVATFVFLWNMIAGGYTDLAGV
QHG---PIIDSPLAQMVG--LPMTAFTILTPSLGLLLVFRTNTSYGRWDEARKMWGLNINHT--
RDLNRMATAWYGNEGNMDSVAFMGGDIPYSQPIDPVQRAYDL-GQVSLFT----WAFVRSMKRHLSPPEEDEED-
FKAELRA-----RLTPEQAENI--INAAHR------------PNRALFDLSVAIEN----LPMHFL--------
------RKNAINTNLSIFEDTLGGCERLLSSPVPLFYSRH-TARFLSTWLLLL-------
PFGLYEQFKDSWNHIAMIPATAFISVCLFGIEELATQLEEPFTI----LPMQGFCDK-IGGWCDEIVSW-
>WP_049046555.1_Kiebsiella_aerogenes
----------------------------------------------------------------
----------------------------------------------------------------
------MIIRPEQHWFFRLFDWHGSVLSKIVFRLLLNVLMSVIAIISYQWYEQL-------------GI------
-----HLTVAPFSLLGIAIAIFLGFRNSASYSRFVEARNLWGTVLIAE--RTLVRQLKNILPD-----------
--------DEETH-----KTLVSYL----VAFSWSLKHQLRK-TD-----PAVDLYR-----LLPKEKVAEI--
LASSMP-------------TNRILLLIGNELGRLREQGKLSDI--------------
TYGLMDNKLDELAHVLGGCERLASTPVPFAYTLI-LQRTVYLFCTLL-------PFALVGDL-----
HYMTPFVSVFISYTFLSWDSLAEELEDPFGTSANDLPLNAMCNT-IERNLMDMTGQH
>XP_011543531.1_Homo_sapiens
M---------------------------------------------------------------
-----------------------------------------------------------------
TITYTSQVANARLGSFSRLLLCWRGSIYKLLYGEFLIFLLCYYIIRFIYRLALTEEQQL--------------
MFEKLTL--YCDSYIQL--IPISFVLGFYVTLVVTRW------WNQYENLPWPDRLMSLVSGFVEGK--------
----------DEQGRL--LRRTLIRYANLGNVLILRSVSTAVYKRFP-----SAQHLVQ---
AGFMTPAEHKQLEKLSLPHNM---------FWVPWVWFANLSMKAWLG---GRIRDPI------------
LLQSLLNEMNTLRTQCGHLYAYDWISIPLVYTQVVTVAVYSFFLTCLVGRQFLNPAKAYPGHEL---
DLVVPVFTFLQFFFYVGWLKVAEQLINPFGEDDDDFETNWIVDRNLQVSLLAVDEMH
```

Supplemental Figure 16. Sequences used to generate MAFFT alignments used to generate the phylogenetic trees included in this manuscript. FASTA files of MAFFT alignments of **A.** Full length and **B.** trimmed after the bestrophin domain BST4 and homologous amino acid sequences.