Short Title: Role of BST4 in Chlamydomonas

Title: The role of BST4 in the pyrenoid of *Chlamydomonas reinhardtii*

Authors: Liat Adler[1,2,], Chun Sing Lau[3], Kashif M. Shaikh[4], Kim A. van Maldegem[4],

Alex L. Payne-Dwyer[3,5], Cecile Lefoulon[6], Philipp Girr[3], Nicky Atkinson[1,2], James

Barrett[3], Tom Z. Emrich-Mills[3], Emilija Dukic[4], Michael R. Blatt[6], Mark C. Leake[3,5],

Gilles Peltier[7], Cornelia Spetea[4], Alistair J. McCormick[†,1,2], Luke C. M. Mackinder[†,3],

Charlotte E. Walker[†,3]

[1]Institute of Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, EH9 3BF, United Kingdom.

[2]Centre for Engineering Biology, University of Edinburgh, EH9 3BF, United Kingdom.

[3]Centre for Novel Agricultural Products (CNAP), Department of Biology, University of

York, Heslington, York YO10 5DD, United Kingdom.

[4]Department of Biological and Environmental Sciences, University of Gothenburg,

Gothenburg 40530, Sweden.

[5]School of Physics, Engineering and Technology, University of York, Heslington, York

YO10 5DD, United Kingdom.

[6]Laboratory of Plant Physiology and Biophysics, Bower Building, University of

Glasgow, Glasgow, United Kingdom.

[7]Aix-Marseille Université, CEA, CNRS, Institut de Biosciences et Biotechnologies Aix-
Marseille, CEA Cadarache, 13108 Saint-Paul-lez-Durance, France

[†]The author(s) responsible for distribution of materials integral to the findings

presented in this article in accordance with the policy described in the Instructions for

Authors (https://academic.oup.com/plcell/pages/General-Instructions) are:

• Alistair McCormick, School of Biological Sciences, University of Edinburgh, +44

1316 505316, alistair.mccormick@ed.ac.uk

• Luke Mackinder, Department of Biology, University of York, +44 1904 328984,

luke.mackinder@york.ac.uk

• Charlotte Walker, Department of Biology, University of York, +44 1904 328500,

charlotte.walker@york.ac.uk

One-sentence summary: In Chlamydomonas, the pyrenoid-localized bestrophin-like protein BST4 is a putative ion channel involved in pH regulation of the thylakoid lumen.

Funding information: This work was funded by the Biotechnology and Biological Sciences Research Council (BBSRC) (grant number (BB/S015531/1) and Leverhulme Trust (RPG-2017-402). LA acknowledges funding support by EASTBIO DTP (BB/J01446X/1). CEW acknowledges funding support from BBSRC Discovery Fellowship (BB/W009587/1). CSL was supported by Bill & Melinda Gates Agricultural Innovations (Investment ID 53197). CS acknowledges grants from the Swedish Research Council (VR 2016-03836 and 2021-03790). KMS was recipient of a postdoctoral fellowship from the Carl Tryggers Foundation (CTS 20:406). A United Kingdom Research and Innovation Future Leaders Fellowship to LCMM (MR/T020679/1), Biotechnology and Biological Sciences Research Council grants (BB/T017589/1, BB/S015337/1, BB/R001014/1) and Engineering and Physical Research Council Grant (EP/W024063/1) with ML and APD. The authors would like to thank the University of York Biosciences Technology Facility for confocal microscopy access and support. CL was funded by BBSRC grants BB/P011586/1 and BB/T013508/1 to MRB. PG was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – project number 456013262.

ABSTRACT

The eukaryotic algal CO_2-concentrating mechanism (CCM) is based on a Rubisco-rich organelle called the pyrenoid, which is typically traversed by a network of thylakoid membranes. BST4 is a bestrophin-like transmembrane protein that has previously been identified in the model alga *Chlamydomonas reinhardtii* as a putative tether that could link the traversing thylakoid membrane network to the Rubisco matrix. In the present study, we show that BST4 forms a higher order complex assembly that localizes to the thylakoid network within the pyrenoid. However, investigation of a *bst4* knock-out mutant in Chlamydomonas showed that the absence of BST4 did not result in a CCM-deficient phenotype and that BST4 is not necessary for the formation of the trans-pyrenoid thylakoids. Furthermore, heterologous expression of BST4 was not sufficient to facilitate the incorporation of thylakoids into a reconstituted Rubisco condensate in the land plant Arabidopsis. Subsequent analyses revealed that *bst4* was under oxidative stress and showed enhanced non-photochemical quenching

67 associated with CO_2 limitation and over acidification of the thylakoid lumen. We

68 conclude that the primary role of BST4 is not as a tethering protein, but rather as an

69 ion channel involved in pH regulation in pyrenoid-based CCMs.

INTRODUCTION

Maintaining improvement in crop yields to keep pace with the rising demands for food

is becoming increasingly challenging (Horton et al., 2021). A possible solution to

overcome this challenge is engineering a biophysical CO_2 concentrating mechanism

(CCM) into C3 crop plants, which has the potential to improve crop yields by up to

60% as well as water-use and nitrogen-use efficiency (Price et al., 2013; McGrath and

Long, 2014; Long et al., 2019; Fei et al., 2022; Wu et al., 2023). The biophysical CCMs

in algae typically function by concentrating CO_2 into a Rubisco-rich, liquid-liquid phase

separated microcompartment called a pyrenoid. This raises the $[CO_2]:[O_2]$ ratio around

the primary CO_2-fixing enzyme Rubisco, which brings Rubisco closer to its maximal

carboxylation rate and minimizes the competing oxygenation reaction.

Chlamydomonas reinhardtii (hereafter Chlamydomonas) has the most well

understood pyrenoid-based CCM and has become the blueprint for engineering such

a CCM into C3 plants (Hennacy and Jonikas, 2020; Adler et al., 2022).

An important yet little understood aspect of the Chlamydomonas CCM is the function

and biogenesis of the thylakoid tubule network that traverses the pyrenoid (Engel et

al., 2015). The pyrenoid tubules are continuous with the thylakoid membrane and are

thought to function as a delivery system for inorganic carbon (Ci) to the Rubisco matrix.

In the current model, bicarbonate (HCO_3^-) is channelled into the thylakoid lumen by

bestrophin-like proteins 1-3 (BST1-3) and diffuses to the pyrenoid tubules where it is

converted to CO_2 by carbonic anhydrase 3 (CAH3) (Karlsson et al., 1998; Mukherjee

et al., 2019; Burlacot et al., 2022). Association of the condensed Rubisco to the

thylakoid membrane is predicted to be essential for an efficient CCM. Therefore,

understanding the mechanisms of pyrenoid tubule formation will be crucial (Fei et al.,

2022).

The protein BST4 (bestrophin-like protein 4, also known as Rubisco binding

membrane protein 1, RBMP1; Cre06.g261750) localizes exclusively to the pyrenoid

tubules and has been proposed to function as a tether protein, linking the Rubisco

matrix to the tubules (Meyer et al., 2020). BST4 is a predicted transmembrane protein

and has two Rubisco binding motifs (RBMs) on its long, disordered C-terminus (Fig.
1A) (He et al., 2020; Meyer et al., 2020). RBMs facilitate the targeting of proteins to
the pyrenoid and are also hypothesized to underpin the assembly of the pyrenoid.
Meyer et al. (2020) proposed that BST4, together with other tether proteins, may
recruit Rubisco to the tubule network.

BST4 also has a well-conserved bestrophin domain similar to those present in the
thylakoid-localized BST1-3 proteins (Mukherjee et al. 2019). Bestrophins primarily act
as anion channels and are found in a wide diversity of organisms, including animals,
plants and fungi. They are best known to have permeability to chloride and HCO_3^- (Qu
and Hartzell, 2008), although some are reportedly permeable to cations (Yang et al.,
2014) and larger organic anions (Roberts et al., 2011). It is possible that BST4
functions as an HCO_3^- channel, like that proposed for BST1-3 (Mukherjee et al. 2019),
but there are currently no data to support this hypothesis.

In this study we aimed to elucidate the role of BST4 in the Chlamydomonas CCM. We
tested BST4 as a thylakoid-Rubisco tethering protein as well as its suitability in
promoting a thylakoid-Rubisco matrix interface in the model land plant Arabidopsis.
We also explored an alternative hypothesis that BST4 functions as an ion channel in
the pyrenoid tubules and discussed the implications this has for our current
understanding of CO_2 fixation by the pyrenoid.

RESULTS AND DISCUSSION
BST4 is a bestrophin-like protein that is localized in the intra-pyrenoid thylakoid
tubules in Chlamydomonas
The amino acid sequence of BST4 has two key unique features compared to BST1-3.
(Fig. 1), the latter of which play a key role in Ci uptake in the CCM (Mukherjee et al.
2019). First, BST4 has an extended disordered C-terminus that contains two RBMs
(Fig. 1A) (He et al., 2020; Meyer et al., 2020). Second, BST4 has a phenylalanine
residue in the first position of the putative selection pore, as opposed to valine which
is conserved throughout BST1-3 (Fig.1B, C). In corroboration, we found the
evolutionary history of BST4 to be distinct from BST1-3 when investigated using
maximum likelihood phylogenetic analysis. We analysed the full-length sequence (Fig.
S1) and a truncated version without the disordered C-terminal, leaving the bestrophin

domain and N-terminal (Fig. 1D), both of which found BST4 to resolve in a distinct

clade from BST1-3 but within the wider green algae group.

Figure 1. BST4 is a bestrophin protein that is distinct from BST1-3. A. Schematic of the topology of BST4.
BST4 is predicted to have four transmembrane domains and a disordered C-terminus containing two Rubisco
Binding Motifs (RBMs) **B.** AlphaFold models of two BST4 bestrophin domains (amino acid residues 53-386 shown
for clarity) to show a typical bestrophin channel pore. Pore lining residues are highlighted in red. **C.** AlphaFold
structures of the BST4 bestrophin domains alongside structures of bestrophins from *Chlamydomonas reinhardtii*
(*Cr*BST1; Alphafold), *Malus domestica* (*Md*VCCN1;7EK1) and *Homo sapiens* (*Hs*BEST1; 8D1I). Residues known
to line the channel pore are highlighted. **D.** Phylogenetic analysis of the BST4 bestrophin domain (bold) with the
disordered C-terminal removed. The alignment used was trimmed at residue 369. The evolutionary history of BST4
was inferred by using the maximum likelihood method based on the Le and Gascuel substitution model with discrete
Gamma distribution (5 categories) and 500 bootstrap replicates. The tree is drawn to scale, with branch lengths
measured in the number of substitutions per site.

As well as being distinct at a sequence level, BST4 also localizes differently from

BST1-3. While BST1-3 localize throughout the thylakoid membrane and are enriched

at the pyrenoid periphery (Mukherjee et al., 2019), BST4 localizes to the centre of the

pyrenoid in a pattern that resembles the pyrenoid thylakoid tubule system (Fig. 2A, B;

Meyer et al., 2020). To confirm tubule localisation, we generated a dual-tagged line

that expressed BST4-mScarlet-I and the Chlamydomonas Rubisco small subunit 1

(*Cr*RBCS1) fused to Venus. BST4-mScarlet-I was enriched where *Cr*RBCS1-Venus was depleted (Fig. 2C, D), indicating that BST4 is located in the tubules and not the Rubisco-enriched pyrenoid matrix. Nevertheless, previous work has shown that the C-terminal RBMs of BST4 enable the protein to interact with *Cr*RBCS1 (Meyer et al. 2020). We confirmed using a yeast-2-hybrid approach that the C-terminus of BST4 interacts with *Cr*RBCS1 (Fig. S2A). We also measured the efficiency of Förster resonance energy transfer (FRET) from Venus to mScarlet-I, and found that the FRET efficiency was ~35%, supporting the proximity of *Cr*BST4 to *Cr*RBCS1 (Fig. S2B).

Bestrophins typically form pentameric assemblies (Bharill et al., 2014; Hagino et al., 2022). To investigate if this is the case for BST4, we initially used an AlphaFold-multimer modelling approach. When five chains of BST4 were inputted to AlphaFold a pentameric structure was predicted (Fig. 2E). To test the complex assembly of BST4 *in vivo*, we utilized a Slimfield microscopy molecular tracking method (Plank et al., 2009). We first developed a Chlamydomonas line that only expressed fluorescently tagged BST4 by expressing BST4-mScarlet-I in a *bst4* mutant background (Fig. S3). We then tracked individual fluorescent molecules of BST4-mScarlet-I and quantified the number of BST4 monomers per complex. The resulting probability distribution revealed that the most common BST4 complex is made up of five molecules (Fig. 2F). Other peaks showed complexes with numbers of molecules divisible by five, which may be multiple pentameric channels grouping together. The interval between the probability peaks was also five (Fig. 2G). To further support higher order complex assembly of BST4 we ran purified Chlamydomonas thylakoid membranes on a Blue Native-PAGE gel (BN-PAGE) and immunoblotted for BST4 (Fig. 2H). BST4 formed a smear at ~1000 kDa, which is considerably larger than a pentamer (~330kDa). This could be due to higher order assemblies of BST4, pentameric BST4 in a complex with other proteins and/or aberrant migration during BN-PAGE due to influences by complex shape. Collectively, our data support *in vivo* higher order assembly of BST4 potentially as a pentamer.

Figure 2. BST4 forms higher order assemblies in the pyrenoid tubules in Chlamydomonas. A. Diagram of a
Chlamydomonas cell with the pyrenoid Rubisco matrix and pyrenoid tubules indicated. **B.** Confocal image of a
Chlamydomonas cell expressing BST4-Venus. Scale bar is 2 μm. **C.** Pyrenoid in dual-tagged Chlamydomonas.
BST4-mScarlet-I and RBCS-Venus are shown in green and magenta, respectively. Overlap appears white. The
yellow line shows the 1D cross-section used for generating the line plot in (D). Scale bar is 1 μm. **D**. Plot of
normalized fluorescence intensity values from a 1D cross-section from (C). mScarlet-I and RBCS-Venus are shown
in green and magenta, respectively. **E**. AlphaFold prediction of pentameric BST4 structure. Top structure includes
the disordered C-terminus, bottom structure displays amino acid residues 53-386 only for clarity. **F**. BST4-mScarlet-
I stoichiometry probability distribution based on single particle tracking and molecular counting in
live Chlamydomonas (n=347 tracks) using Slimfield microscopy. **G**. Averaging the intervals between peaks in
BST4-mScarlet-I stoichiometry (n=22 intervals) indicates a consistent pentameric unit. **H**. Immunoblot of proteins
from *bst4* mutant and wild type (WT) Chlamydomonas thylakoids separated by Blue Native-PAGE.

BST4 localizes to the stroma lamellae thylakoid membrane in Arabidopsis

We used Arabidopsis as a heterologous system to examine if BST4 was able to tether
the Rubisco matrix to thylakoid membranes. In previous work, an Arabidopsis Rubisco
small subunit (*At*RBCS) double mutant (*1a3b*) was complemented with *Cr*RBCS2,

resulting in a line with hybrid Rubisco representing ~50% of the Rubisco pool (S2$_{Cr}$)

(Izumi et al., 2012; Atkinson et al., 2017). Subsequent expression of the pyrenoid linker

protein Essential PYrenoid Component 1 (EPYC1), in S2$_{Cr}$ resulted in the formation of

an EPYC1-hybrid Rubisco condensate or 'proto-pyrenoid' (Atkinson et al., 2020). The

S2$_{Cr}$ line was therefore used as a platform to test whether BST4 acts as a tether

protein.

We initially localized a BST4-mNeon fusion protein following stable expression in S2$_{Cr}$

(Fig. 3). BST4-mNeon was observed in chloroplasts, demonstrating that the native

chloroplast signal peptide was compatible with the chloroplast targeting mechanism in

land plants (Fig. 3A), as seen previously for other Chlamydomonas proteins (Atkinson

et al., 2016). The fluorescence signal from BST4-mNeon had a sponge-like pattern

that was inversely correlated with the punctate chlorophyll autofluorescence signal that

represents the grana stacks (Fig. 3B, C). The sponge-like pattern was similar to

previous observations of autofluorescence originating from Photosystem I (Hasegawa

et al., 2010), which is enriched in the stroma lamellae of thylakoids. We subsequently

generated a stable Arabidopsis transgenic line expressing untagged BST4 to confirm

its location by biochemical fractionation. BST4 was detected in the thylakoid fraction

and not in the stromal fraction (Fig. 3D). The thylakoids were then further fractionated

into grana stacks and stroma lamellae sub-fractions. BST4 was found in the stroma

lamellae fraction (Fig. 3E), which is consistent with the observed sponge-like

fluorescence pattern.

Figure 3. BST4 assembles as a complex in the stroma lamellae of thylakoids in Arabidopsis. A. Confocal image of WT protoplasts expressing BST4-mNeon. Scale bar is 5 μm. **B.** Mesophyll chloroplast from S2$_{Cr}$ Arabidopsis expressing BST4-mNeon. BST4mNeon and chlorophyll autofluorescence are shown in green and magenta, respectively. Overlap appears white. Scale bar is 2 μm. Yellow lines indicate selections for profile plot in (C). **C.** Plot of normalized fluorescence intensity values from a 1D cross-section (yellow line) through two grana stacks from (B). mNeon and chlorophyll autofluorescence are shown in green and magenta, respectively. **D.** Immunoblots of sub-chloroplast fractions isolated from Arabidopsis line S2$_{Cr}$ expressing BST4. RbcL and CP43 were probed for as stromal and thylakoid controls, respectively. **E.** Immunoblots of fractionation thylakoids from Arabidopsis line S2$_{Cr}$ expressing BST4. CP43 and PsaB were used for grana stack and stroma lamellae controls, respectively **F.** Trypsin protease protection assay. Intact thylakoids containing BST4 subjected to 0 or 100 μg/ml trypsin with or without the addition of 1% (v/v) Triton. AtpB and PsbO used as controls for stromal facing (exposed) and lumen facing (protected), respectively. **G.** Immunoblot of proteins from thylakoids separated by Blue Native-PAGE from either BST4 stable line or S2$_{Cr}$ background. Abbreviations: Clp, whole chloroplast; CP43, CP43-like chlorophyll binding protein; RbcL, Rubisco large subunit; PsbO, photosystem II manganese-stabilizing polypeptide;AtpB, ATP synthase subunit beta; PsaB, photosystem I P700 chlorophyll a apoprotein A2; Str, stromal fraction; Thy, thylakoid fraction.

As BST4 has two RBMs on its C-terminus, we hypothesized that BST4 should be orientated with the C-terminus facing the stroma so that the RBMs are available to interact with *Cr*RBCS. To determine the orientation of BST4, we performed a protease protection assay on thylakoids isolated from the untagged BST4 transgenic plants. Our antibody was raised against the C-terminal end of BST4, so it could be used to assess whether the C-terminus was exposed for degradation in the stroma or if it was protected in the lumen. We found that the BST4 C-terminus was fully degraded after a 60 min treatment of trypsin, indicating that it faced the stroma (Fig. 3F). There was some degradation of the lumenal control, PsbO, which we attribute to a portion of the thylakoid membrane preparation not being fully intact. However, PsbO was fully degraded when the membranes were solubilized, indicating that they were sufficiently intact to differentiate between lumenal- and stromal-facing peptides. Therefore, BST4 was observed in the expected location and orientation in plant thylakoid membranes.

Finally, we also tested whether BST4 forms a complex in the Arabidopsis thylakoid membrane. We subjected thylakoids from the untagged BST4 transgenic plants to BN-PAGE and detected a single band of ~850 kDa (Fig. 3G). Thus, BST4 forms a similar order complex in Arabidopsis to that in Chlamydomonas but may be lacking additional interaction partners present in Chlamydomonas.

BST4 is not sufficient for integration of thylakoids into the Arabidopsis proto-pyrenoid

To test whether BST4 could facilitate incorporation of thylakoids into hybrid Rubisco condensates (i.e., proto-pyrenoids) in plants, BST4-mCherry and EPYC1-tGFP were co-expressed in the S2$_{Cr}$ background (Fig. 4). When we expressed EPYC1 alone,

Rubisco condensates formed in chloroplasts as expected (Atkinson et al., 2020), which were visible as a ~2 μm wide puncta in the tGFP channel (Fig. 4A). When BST4-mCherry and EPYC1-tGFP were co-expressed, approximately 60% of the BST4-mCherry fluorescence signal was observed in the condensate region (Fig. 4B, D), with the remaining signal exhibiting the same sponge-like pattern as seen in Fig. 3A.

To confirm that the observed co-localisation was due to an interaction between BST4 and CrRBCS2, we mutated the first two residues of each core RBM motif of BST4 to glutamic acid (WR to EE). Previously, these substitutions have been reported to disrupt the binding interface between EPYC1 and CrRBCS2 (He et al., 2020). Using yeast-2-hybrid we confirmed that the interaction between the C-terminus of BST4 and CrRBCS2 was disrupted by these mutations (Fig. S3). When BST4$_{WR/EE}$-mCherry was expressed with EPYC1-tGFP in S2$_{Cr}$, no enrichment of the mCherry signal was observed in the Rubisco condensate (Fig. 4C, D). Thus, the RBMs of BST4 were responsible for the enrichment of BST4 in the vicinity of the Rubisco condensate in Arabidopsis.

However, when condensates were sedimented and analysed by immunoblotting, BST4-mCherry was not detected in the condensate fraction (Fig. 4E). When leaf samples were subjected to formaldehyde cross-linking prior to sedimentation of the condensate, a small amount of BST4 was present in the condensate fraction. We concluded that BST4 was not present in the condensate itself but preferentially occupied the thylakoid membranes surrounding the CrRBCS2-enriched condensate (Atkinson et al., 2020), likely due to interactions with CrRBCS2.

Figure 4. BST4 is not sufficient to enable the inclusion of thylakoid membranes in a Rubisco condensate in Arabidopsis and is not required for pyrenoid tubule inclusion in the Chlamydomonas pyrenoid. **A**. Confocal images showing EPYC1-tGFP in S2$_{Cr}$ Arabidopsis background. Scale bar is 10 µm. **B**. Confocal images showing BST4-mCherry co-expressed with EPYC1-tGFP in S2$_{Cr}$ Arabidopsis background. Scale bar is 10 µm. **C**. Confocal images showing BST4 with mutated RBMs (BST4$_{(WR/EE)}$) fused to mCherry co-expressed with EPYC1-tGFP in the S2$_{Cr}$ Arabidopsis background. **D.** Ratio of mean mCherry fluorescence associated with the Rubisco condensate compares do the rest of chloroplast when mCherry was fused to either BST4 or BST4$_{(WR/EE)}$ (n=22-23). **E**. Immunoblot of sedimented Rubisco condensates from S2$_{Cr}$ Arabidopsis expressing EPYC1-tGFP and BST4-mCherry. Abbreviations: I, input; S, supernatant; P, pellet. Rubisco condensates are enriched in the pelleted fraction. Cross-linked samples were prepared by vacuum infiltrating intact leaves with 1% (v/v) formaldehyde prior to sedimentation. **F**. Transmission electron micrograph of a chloroplast from S2$_{Cr}$ Arabidopsis expressing either EPYC1 alone or EPYC1-tGFP and BST4-mCherry. Rc is the Rubisco condensate. Scale bar is 1 µm. **G**. Transmission electron micrograph of pyrenoid from WT Chlamydomonas or *bst4* mutant. Abbreviations: tt, thylakoids tubules; p, pyrenoid; s, starch sheath. Scale bar is 1 µm.

Although BST4 partially co-localized with the Rubisco condensate, we found no evidence to suggest that that BST4 could facilitate the inclusion of thylakoid membranes. Confocal microscopy was not sufficient to determine if chlorophyll autofluorescence was in the proto-pyrenoids of S2$_{Cr}$ lines expressing BST4-mCherry and EPYC1-tGFP. However, transmission electron microscopy (TEM) revealed no visible indication of thylakoid membranes in the condensates, which were structurally similar to condensates in S2$_{Cr}$-EPYC1 lines lacking BST4 (Fig. 4F). Thus, BST4

appeared insufficient to enable thylakoid inclusion in the Rubisco condensate in
Arabidopsis.

BST4 is not necessary for tubule formation in Chlamydomonas
To investigate whether BST4 was necessary for normal formation of the thylakoid
tubule-structure in Chlamydomonas we compared the structure of pyrenoids of WT
and a BST4 knock-out line (*bst4*) (Fig. S3). TEM images showed that pyrenoids from
bst4 were structurally comparable to the WT control, including the presence of
pyrenoid tubules (Fig. 4G). There were also no differences in pyrenoid size or shape
between the two lines when comparing pyrenoids from 40-50 cells from each genotype
(Fig. S4). As a result, we conclude that BST4 is not necessary for the pyrenoid tubule-
Rubisco matrix interface in Chlamydomonas. We speculate that BST4 may require a
pre-existing pyrenoid tubule network to be localized in the pyrenoid rather than driving
the inclusion of thylakoid membranes into the Rubisco matrix, or is redundant as a
tether protein.

The C-terminus of BST4 is required for localisation to the pyrenoid tubules
Meyer et al., 2020 previously demonstrated multiple copies of the BM are sufficient t
target proteins to the pyrenoid. In Fig. S2A we show that the BST4 RBMs are required
for RBCS interaction via yeast-2-hybrid and in Fig. 4B-D we show they are required
for enrichment with the proto-pyrenoid in Arabidopsis. To investigate the role of the
RBMs in BST4 localization in Chlamydomonas, we generated a truncated version of
BST4 (residues 1 to 386) that lacked the C-terminus containing the two RBMs
(BST4$_{\Delta C\text{-term}}$) and compared localisation with full-length BST4 expressed in WT and
the *bst4* mutant (Fig. 5 and Fig. S5). BST4$_{\Delta C\text{-term}}$-mScarlet expressed in the *bst4*
mutant line did not localize to the pyrenoid tubules but was found throughout the
thylakoid membrane (Fig. 5). This is consistent with our findings in Arabidopsis with
BST4$_{WR/EE}$ (Fig. 4C), demonstrating that localization of BST4 is driven via RBM-
Rubisco interaction. Thus, the C-terminus is necessary for BST4 localisation to the
pyrenoid tubules in the presence of the matrix.

When BST4$_{\Delta C\text{-term}}$-Venus was expressed in WT Chlamydomonas (i.e., that still
produced non-truncated BST4), we observed fluorescence throughout the thylakoids,
but with the majority of the signal still localized to the pyrenoid tubules (Fig. S5). We

conclude that BST4$_{\Delta C\text{-term}}$-Venus is recruited to the pyrenoid through an interaction
with the native full-length BST4, which is further evidence that BST4 oligomerizes.

We next investigated whether BST4 localizes to the pyrenoid tubules, or whether
BST4 localizes to the tubules through an interaction with Rubisco. To do so, we utilized
a Chlamydomonas mutant generated by Genkov et al. (2010) that expresses *At*RBCS
but lacks both isoforms of *Cr*RBCS (*Crrbcs::AtRBCS*), and thus lacks a Rubisco matrix
as EPYC1 does not interact with *At*RBCS (Atkinson et al., 2019). *Crrbcs::AtRBCS*
retains reticulated thylakoid membranes at the canonical pyrenoid site, which are likely
the nascent pyrenoid tubule network (Caspari et al., 2017) (Fig. S6A). Caspari et al.
(2017) show large starch granules accumulating at the canonical pyrenoid site, which
we confirmed by expressing STA2-Venus as a starch marker (Mackinder et al., 2017)
in *Crrbcs::AtRBCS* (Fig. S6B). We expressed BST4-Venus in *Crrbcs::AtRBCS* and
found that BST4 localized to a punctum adjacent to the canonical pyrenoid site, which
we attribute to the nascent pyrenoid tubules (Fig. S6C). To confirm this, we also
expressed a known pyrenoid tubule marker protein, PsaF (Emrich-Mills et al., 2021)
in *Crrbcs::AtRBCS*, which showed a similar localisation pattern to that of BST4 (Fig.
S6D). This suggests there may be an additional Rubisco-binding independent
mechanism for localisation of BST4 as well as other proteins to the pyrenoid tubules.
Further investigation of the role of the whole C-terminal domain will be required to
understand the mode of BST4 pyrenoid tubule localisation.

Chlamydomonas KO mutant *bst4* does not have a defective CCM but is under oxidative stress

To test whether BST4 has a role in the operation of the CCM, we measured the growth of *bst4* compared to WT under various CO_2 conditions (Fig. 6). Semi-quantitative spot assays indicated no reduction in growth under CO_2-limiting conditions (Fig. S7). The liquid growth assay also demonstrated that *bst4* grew well in both CO_2 replete and limiting conditions, 3% and 0.04%, respectively (Fig. 6A). The *bst4* mutant had a slightly increased cell density compared to WT (Fig. 6A) in the 0.04% CO_2 treatment. However, when comparing the calculated specific growth rates (μ h^{-1}) for both the exponential growth phase (days 0-3, *bst4* 0.0402 ± 0.0003 μ h^{-1} and WT 0.0389 ± 0.0007 μ h^{-1}) or the full growth assay (days 0-5, *bst4* 0.0257 ± 0.0001 μ h^{-1} and WT 0.0241 ± 0.001 μ h^{-1}), there was no statistically significant increase in specific growth rates between *bst4* and WT (two-tailed t-test, p = 0.17 and 0.20, respectively, n = 3, full results Table S1). In combination, we concluded that BST4 is not an essential component of the CCM and may therefore have a redundant or regulatory role. We also included the complemented *bst4::BST4* and *bst4::BST4$_{\Delta C-term}$-mScarlet-I* (hereafter *bst4::BST4$_{\Delta C-term}$*) lines in the spot and liquid growth assays. While all lines grew well in the spot assay, in the liquid growth *bst4::BST4$_{\Delta C-term}$* grew comparably to WT and *bst4* whereas *bst4::BST4* exhibited a slightly reduced growth than the other lines in both CO_2 conditions.

Figure 6. The *bst4* mutant does not have an impaired growth phenotype under CCM induced conditions but is associated with oxidative stress. A. *Chlamydomonas* strains were subjected to a liquid growth assay using pH 7.4 TP media that was bubbled with 0.04% CO_2 or 3% CO_2. Error bars are ± SEM, n=3. **B.** Dot assay of WT, *bst4,* complemented *bst4::BST4* and *bst4* complemented with C-terminal truncation of BST4 (*bst4::BST4*$_{\Delta C\text{-term}}$) on minimal TP agar at indicated light intensities and CO_2 concentrations. **C**. Phototaxis assay. Liquid cultures of Chlamydomonas cells were uniformly distributed on 0.8% (w/v) agar and directional light applied. Cell phototaxis was monitored by aerial photographs of wells at 0, 1, and 3 h. The assay was performed in the presence of 75 µM ROS hydrogen peroxide (H_2O_2) or 0.3 M ROS quencher N,N'-dimethylthiourea (DMTU).

One noticeable difference between *bst4* and WT lines from the growth assays conducted on solid media (Fig. 6B and S8) was that *bst4* cells had a distinct halo of

diffuse cells on the periphery of the colony. We used a range of CO_2 and light

conditions to investigate the diffuse colony phenotype (Fig. 6B) and found it was most

apparent under high light (300 µmol photons m^{-2} s^{-1}) and low or very low CO_2

conditions (0.04% CO_2, 0.01% CO_2, respectively). WT, *bst4::BST4* and *bst4::BST4-*

mScarlet-I complemented lines had little or no diffusivity, although WT did display a

slightly diffuse colony phenotype at 300 µmol photons m^{-2} s^{-1}. Interestingly, BST4$_{\Delta C\text{-}}$

$_{term}$-mScarlet-I was unable to rescue the diffuse colony phenotype suggesting that

either the presence of the C-terminus or localization to the tubules is essential for the

function of BST4.

The diffuse colony phenotype was most apparent under high light (300 µmol photons

m^{-2} s^{-1}) and exacerbated by low CO_2 (0.01 and 0.04%), at medium light (150 µmol

photons m^{-2} s^{-1}) which are conditions where carbon fixation can be limiting and

therefore, the light dependent and light independent reactions of photosynthesis are

unbalanced. This imbalance can result in the release of Reactive Oxygen Species

(ROS) (Erickson et al., 2015). It has previously been shown that cells exposed to ROS

have altered phototactic responses to light (Wakabayashi et al., 2011), which may

explain the diffuse colony phenotype we observed. To test this hypothesis, we

exposed Chlamydomonas cells in liquid culture to directional light (150 µmol photons

m^{-2} s^{-1}) (Fig. 6C). In this assay, *bst4* cells displayed strong positive phototaxis,

whereas WT and complemented *bst4::BST4* lines displayed negative phototaxis,

indicating that *bst4* cells may be experiencing oxidative stress.

To confirm that the phototactic response of *bst4* was due to oxidative stress, we

recorded the direction of phototaxis in cells exposed to either ROS or a ROS quencher.

In the presence of the ROS quencher N,N'-dimethylthiourea (0.3 M DMTU), *bst4*

positive phototaxis was disrupted, resulting in a negative response to directional light

(Fig. 6C). When the ROS H_2O_2 (75 µM) was added, WT, *bst4::BST4* and *bst4*

displayed positive phototaxis (Fig. 6C). Together, our data indicates that the direction

of phototaxis in *bst4* is due to the degree of oxidative stress.

ROS generation occurs under excess light which can be caused by either high light

conditions or limited CO_2 fixation due to an impaired CCM or low CO_2 availability

(Santhanagopalan et al., 2021; Choi et al., 2022). As *bst4* can maintain similar growth

to WT under low CO_2 conditions, we hypothesized that CO_2 fixation is not limiting in

this case. Rather, the absence of BST4 may disrupt regulation of the light reactions.

BST4 is associated with regulation of the lumenal pH in Chlamydomonas.

To assess the impact of BST4 on the light reactions, we measured photosynthetic

parameters of WT, *bst4* and *bst4* complemented with either full length *BST4* or

BST4$_{\Delta C\text{-term}}$. Chlorophyll and carotenoid levels were slightly lower in *bst4*, however this

was not significantly different. Complementation with BST4 recovered chlorophyll and

carotenoid levels to a higher level than WT whilst in *bst4::BST4$_{\Delta C\text{-term}}$* lines levels were

comparable to the *bst4* mutant (Fig. S9A).

Next, we measured the slow kinetics of chlorophyll fluorescence upon transition from

dark to high light to determine the impact of BST4 on the distribution of absorbed light

between non-photochemical quenching (NPQ) and electron transport through

photosystem II (Y(II)) (Fig. 7). F_v/F_m was slightly higher in *bst4, bst4::BST4* and

bst4::BST4$_{\Delta C\text{-term}}$ compared to WT (Fig. 7A). NPQ was induced faster and reached a

significantly higher steady state in *bst4* than in WT (Fig. 7B). In contrast, *bst4::BST4*

had WT-like NPQ. *bst4::BST4$_{\Delta C\text{-term}}$* had an intermediate phenotype indicating

incomplete rescue of the high NPQ phenotype via BST4$_{\Delta C\text{-term}}$. No differences in Y(II)

were observed during the light phase (Fig. 7C). However, in the subsequent dark

phase, Y(II) in *bst4* recovered faster in line with the higher F_v/F_m (Fig. 7C).

The increase in NPQ suggested that BST4 influences the reactions occurring in the

thylakoid membrane. A higher steady-state NPQ indicates that the *bst4* cells

experience high light stress and respond by accelerating photoprotective excess light

dissipation to maintain optimal electron transport. This observation aligns with

increased ROS production observed in *bst4* (Fig. 6C), which would result in elevated

NPQ to compensate for the increase in oxidative stress. In the case of *bst4::BST4$_{\Delta C\text{-}}$*

$_{term}$, it is possible that the truncation directly affects BST4 activity. Alternatively, the

mis-localization of BST4 activity throughout the thylakoid membranes, instead of solely

localized to the thylakoid tubules, results in the disruption of optimal electron transport

and the need for increased photoprotection.

NPQ is induced in response to a drop in lumenal pH, a process that is mediated by

the quencher protein LHCSR3. Thylakoid ion channels can influence NPQ through

altering the partitioning of the proton motive force (PMF) into pH gradient (ΔpH) and
electrical gradient (ΔΨ). To investigate whether BST4 was altering lumenal pH we
measured the slow decay kinetics of electrochromic shift (ECS), which allows the
determination of PMF partitioning. WT and *bst4::BST4* displayed similar standard
kinetics. However, both *bst4* and *bst4::BST4*$_{\Delta C\text{-}term}$ had abnormal kinetics and did not
reach steady state (Fig. S8A, B). Therefore, we could not derive PMF partitioning for
all the genotypes. The abnormal ECS kinetics in *bst4* and *bst4::BST4*$_{\Delta C\text{-}term}$ may
indicate altered proton-related processes, such as over acidification of the thylakoid
lumen. We also measured the accumulation of protein LHCSR3 after exposing cells
to 150 µmol m^{-2} s^{-1} for 3 h and found that *bst4* accumulated more LHCSR3 as
compared to WT (Fig. 7G, H). In the complemented lines, *bst4::BST4* accumulated
similar levels to WT and *bst4::BST4*$_{\Delta C\text{-}term}$ accumulated less. The increased NPQ
observed in *bst4* may therefore be due to a combination of both upregulation of
LHCSR3 expression and altered lumenal pH.

Figure 7. Chlamydomonas *bst4* mutant has an enhanced NPQ and ATP synthase activity under limiting Ci conditions. Wild type (WT) and mutants were grown on TAP medium at 20 µmol photons m^{-2} s^{-1}, resuspended in TP at 30 µg Chl ml^{-1} and exposed for 3 h to light at 150 µmol photons m^{-2} s^{-1}. The cells were dark adapted for 1 h before the measurements. **A.** Maximum quantum yield of photosystem II. **B.** Dynamics of photosynthesis on transition from dark to high light. Kinetics for induction of chlorophyll fluorescence were recorded during 17 min of illumination at 1500 µmol photons m^{-2} s^{-1} followed by 5 min in darkness. Non-photochemical quenching (NPQ) and **C.** Photosystem II quantum yield (Y(II)). **D to F.** ECS decay kinetics were performed on cells pre-exposed for 10 min to high light and the **D.** total PMF, **E.** g_H^+, and **F.** total H^+ flux (v_H^+) were determined as described in Methods. Data are the means ± SEM (n=3 replicates). Different letters indicate statistically significant difference among the genotypes (one-way ANOVA test, followed by Turkey's post hoc test, P < 0.05). Statistical analysis was performed separately for measurements made with and without HCO_3^- (black and green letters, respectively). **G.** Immunoblot of NPQ protein LHCSR3 in each genotype compared to tubulin after exposure of cells to 3 h 150 µmol photons m^{-2} s^{-1} in TP. **H.** Fluorescence of LHCSR3 protein band normalized to α-tubulin. Data are the means ± SEM (n=3 replicates). Different letters indicate statistically significant difference among the genotypes (one-way ANOVA test, followed by Turkey's post hoc test, P < 0.05).

To further explore the proton related processes, we recorded fast decay ECS kinetics which allows for the determination of the total PMF size, as well as H^+ conductivity of ATP synthase (g_H^+) (Fig. S9E). The PMF after 10 min of illumination did not significantly differ among any of the genotypes (Fig. 7D). At the same timepoint, the g_H^+ was significantly higher in *bst4* and *bst4::BST4$_{\Delta C\text{-term}}$* than in WT and *bst4::BST4* (Fig. 7E). We also calculated the total proton flux (v_H^+), which showed a similar trend although v_H^+ was also significantly lower in *bst4::BST4,* compared to WT (Fig. 7F). Based on the proton flux data, we postulate that more ATP is produced in *bst4* and *bst4::BST4$_{\Delta C\text{-term}}$* and less in *bst4::BST4* and WT. In the *bst4* mutant, H^+ in the lumen are likely in excess and are used in H^+-dependent processes such as NPQ and ATP synthesis.

We next proposed that BST4 might be an anion channel involved in regulating the pH of the thylakoid lumen. To determine what BST4 may be permeable to, we expressed BST4 in Xenopus oocytes and measured currents in the presence of different anions (Fig. S9A). Bestrophins are typically permeable to Cl^- and HCO_3^- but no currents for BST4 were detected in the presence of 100 mM KCl or 100 mM $NaHCO_3$. As some bestrophins are autoinhibited by their C-terminus (Qu et al., 2006), we also tested two C-terminal truncations of BST4 (0-386 and 0-591), but no currents were detected. Another possibility is that BST4 is permeable to organic ions, similarly to *Hs*Best1, which has been shown to be permeable to **γ**-aminobutyric acid (GABA) (Lee et al., 2010) and glutamate (Woo et al., 2012), as well as Cl^- and HCO_3^- anions. Fei et al. (2022) proposed a model whereby RuBP acts as proton carrier to increase H^+ concentration in the pyrenoid tubules. It is possible that BST4 is the channel that facilitates RuBP translocation in the tubules in this model. To test this hypothesis, we used small molecule analogues K-PEP and K-Gluconate but no currents were detected for these either (Fig. S9A). Therefore, we were unable to draw conclusions as to what BST4 is permeable to. BST4 may require certain conditions to be open that are not met in the oocyte system, such as post-translational modifications, a specific pH or an interaction partner. BST4 was found to be phosphorylated and had an oxidized methionine residue in its C-terminus (Bergner et al., 2015). Methionine oxidation can serve as channel-regulating post translational modification (Ciorba et al., 1999), which would fit with the role that BST4 appears to have in preventing oxidative stress.

BST4 has no impact on growth in Arabidopsis S2$_{Cr}$ line

Although the permeability is undefined, if BST4 is a functional anion channel, it is possible that its presence in plant thylakoid membranes might have an effect. We generated three independent BST4 no tag lines in the S2$_{Cr}$ background and used them to assess the impact of BST4 on Arabidopsis physiology (Fig. S10A). We found there was no difference in growth between plant expressing BST4 and their azygous segregants, as determined by the rosette area (Fig. S10B). We also found that BST4 expressing plants tended to have slightly lower F_v/F_m as compared to azygous segregants and the parent line, although this was not significant (Fig. S11C). Further measurements were made for lines 2 and 3 on the kinetics of NPQ, Y(II) and for PMF size and partitioning but no consistent differences were observed (Fig. S11).

Proposed model for the role of BST4 in the Chlamydomonas pyrenoid

Based on these findings, we speculate that BST4 may have a regulatory role in the CCM through its function as a bestrophin channel. We propose two possible models to explain our data. The first model is that BST4 is functioning as an anion channel to import anions from the pyrenoid into the thylakoid lumen. This would counterbalance the charge of protons in the tubules, possibly contributing to pH regulation or maintaining a high flux of protons towards the pyrenoid to power the CCM (Figure 8). The absence of BST4 would cause disruption of lumen pH or cause a drop in CCM efficiency that causes stress under high light and low CO_2. The second model is that BST4 is part of a HCO_3^- recovery system to takes up HCO_3^- from within the pyrenoid, allowing HCO_3^- to be transported back into the tubules where it can be dehydrated to CO_2 and thus available to be fixed by Rubisco. In this scenario, the absence of BST4 would lead to a slight reduction in CCM operation efficiency, perhaps not sufficient to cause a growth penalty but enough to cause a build-up of protons in the tubule lumen and alter the homeostasis of the light reactions. Further work is required to elucidate the function of BST4, primarily, determining what species BST4 is permeable to, any modes of channel regulation and possible interaction partners.

Model 1. BST4 as an anion channel   **Model 2. BST4 as a HCO₃⁻ recovery channel**

Figure 8. The role of BST4 in light and CO₂ limiting conditions. The data presented here show that in the *bst4*
knock-out line there is increased ROS production. There is also likely an over acidified thylakoid lumen, which
causes elevated NPQ and increased ATP generated through ATP Synthase activity. We suggest that BST4
functions as a thylakoid membrane channel, which plays a role in balancing the light reactions with carbon fixation
while the CCM is active. In this role, we propose BST4 is a channel that facilitates the transport of anions from the
pyrenoid matrix into the thylakoid lumen whereby they act as a sink for available protons (Model 1). It is possible
that BST4 is specifically a HCO₃⁻ channel (Model 2) that facilitates the recovery of HCO₃⁻ from the pyrenoid matrix.
This would subsequently enable CAH3 to catalyse the dehydration of HCO₃⁻ into CO₂, which is then able to diffuse
back into the pyrenoid and be fixed by Rubisco.

MATERIALS AND METHODS

Phylogenetic Analysis

Sequences for phylogenetic analysis were compiled by blasting of BST4
(Cre06.g261750) in the NCBI database (Sayer et al 2022) and the manual addition of
other well characterized bestrophin proteins including BEST1 *Homo sapiens*
(XP_011543531.1), KpBEST *Klebsiella aerogenes* (WP_049046555.1), VCCN1 from
Arabidopsis thaliana (Q9M2D2) and BST1-3 *Chlamydomonas reinhardtii*
(Cre16.g662600, Cre16.g663400 and Cre16.g663450). Sequences were aligned
using MAFFT (Katoh and Standley, 2013) visually inspected and manually trimmed
(specified). Finalized alignments were run through IQTREE webserver (Minh et al.,

2020) to identify the most appropriate substitution model. Maximum likelihood trees
were then generated in Geneious v11 using the PhyML 3.0 (Guindon et al., 2010)
plugin with a LG substitution model (Le and Gascuel, 2008) with Gamma distribution
(4 categories) and 500 bootstrap iterations. Full alignments are found in Fig. S12.

Alphafold structure prediction

565 Five chains of BST4 were submitted to Alphafold Multimer v2 with default settings.
566 The top ranked model of five was used for figure generation.

Generation of plasmids

The plasmids for BST4 mutant complementation in *Chlamydomonas* were prepared
using a recombineering method described previously (Emrich-Mills et al., 2021). BST4
is expressed under native promotor with either without a tag or with an mScarlet-I C-
terminal tag, and a hygromycin AphVII selection marker.

Plasmids for plant expression were generated using the MoClo system (Engler et al.,
2014). For visualisation, BST4 was combined with 35S promoter (pICH51277),
mNeonGreen C-terminal tag (pICSL50015), HSP terminator and acceptor plasmid
(pICH47732) and a pFAST-R section cassette used for selection. For all other
experiments a no-tag BST4 construct was generated with the 35S protomer and HSP
terminator parts and a Kanamycin resistance cassette was used for selection.

Plasmids for Xenopus expression were generated by using the Gateway system by
cloning the coding sequence for BST4 into pGT vector (Grefen et al., 2010) was
amplified with Gateway adaptor sequences from a synthesized as g-block (IDT) with
the PredAlgo (Tardif et al., 2012) predicted transit peptide removed from the N-
terminal (sequence begins R35), any subsequent mutations were made via PCR.

Arabidopsis transformation

Arabidopsis was transformed by floral dip as previously described in (Atkinson et al.,
2016). BST4-mNeon primary transformants were screened for transgene insertion by
seed fluorescence from pFAST-R and BST4 expression was confirmed by checking
for mNeon fluorescence and by immunoblot. BST4 no tag primary transformants were
screened using kanamycin resistance and immunoblot. Zygosity was checked via
seed fluorescence from pFAST-R or kanamycin resistance.

Chlamydomonas cell culture conditions and strain details

Chlamydomonas reinhardtii cultures were maintained as previously described (Ma et al., 2011). Tris-Acetate-Phosphate (TAP) and minimal (TP) media (acetate free) were prepared according to Sueoka (1960). TAP and TP agar plates for growth were made by adding 1.5% (w/v) agar. CMJ030 (CC-4533; *cw15, mt⁻*) and *bst4* (BST4 knock-out LMJ.RY0402.159478) were obtained from the CLiP collection at the *Chlamydomonas* culture collection (Zhang et al., 2014; Li et al., 2015). The insertion of the CIB1 cassette in *BST4* was confirmed by PCR amplifying the insertion locus from genomic DNA (Fig. S3A) using loci specific primers (forward GAGCTTCGTGGATGGATGTT and reverse GTATGAAGGTCACCGCCTGT) in parallel with a control locus (forward ATGCTTCTCTGCATCCGTCT and reverse ATGTTTTACGTCCAGTCCGC). The *bst4* knock-out was complemented with *BST4* constructs described herein. All complemented lines were validated by western blotting of BST4 and specified epitope tags, described below (Fig. S3B-E).

Chlamydomonas transformation

For each Chlamydomonas transformation, 28 ng kbp^{-1} of plasmid was linearized by restriction digest. Cells were grown to 2-4 x 10^6 cell mL^{-1}, harvested by centrifugation at 1000 xg for 10 min and resuspended in TAP with 40 mM Sucrose at a concentration of 2×10^8 cells mL^{-1}. Linearized plasmid was mixed with 250 μL of cells at 15°C in a 0.4 cm gap electroporation cuvette and transformed immediately by electroporation using a Gene Pulser II (Bio-Rad) set to 800V and 25 μF. Cells were recovered overnight in TAP Sucrose while shaking gently (140 rpm) in the dark. Transformed cell were subsequently subjected to selection by growth on TAP agar plates with paromomycin (20 μg mL^{-1}) or hygromycin (25 μg mL^{-1}) which were kept in low light (5–10 μmol photons m^{-2} s^{-1}) until screening positive transformants.

Chlamydomonas Growth Assays

Spot Tests: Cells were grown heterotrophically in TAP media. Once cultures reached 2-4 x 10^6 cell mL^{-1}, 1 x 10^6 cells were harvested by centrifugation at 1000 xg for 10 min. Cell were washed and resuspended at a concentration of 1 x 10^6 cell ml^{-1} in TP media. Liquid cultures were spotted onto TP agar (1.5%) in 1000, 100 and 10 cell spots at a range of pHs (specified). The plates were incubated in 3, 0.04 and 0.01% CO_2 and illuminated under constant light at 400 μmol photons m^{-2} s^{-1}. Growth was monitored for up to 10 days.

Liquid Growth: Cells were grown heterotrophically in TAP media. Once cultures reached $2-4 \times 10^6$ cell mL^{-1}, cells were harvested by centrifugation at 1000 xg and resuspended at a starting concentration of 1×10^5 cell ml^{-1} in TP media pH 7.4. Cultures were incubated in a CellDEG HDC 22.10 culture platform (CellDeg GMBH, Berlin) bubbled with 0.04 and 3% CO_2, illuminated at 150 µmol photons m^{-2} s^{-1} and consistently stirred at 180 rpm. Cell density and optical destiny (750 nm) measurements were taken daily for up to 10 days. Specific growth rates (SGR) per h were calculated using the following formula, *ln(N$_2$/N$_1$)/t* whereby N = cell density.

Dot tests: Cultures were grown in a 96 format on agar plates and replicated by a Rotor+ (Singer Instruments) high throughput replication robot. The cultures were stamped onto pH 7.8 TP agar plates, incubated in 3, 0.04 and 0.01% CO_2 and illuminated under constant light at a range of intensities (specified). Growth was monitored for up to 10 days.

Phototaxis Assays

Chlamydomonas cells were grown heterotrophically in TAP media until they reached $2-4 \times 10^6$ cell mL^{-1} and harvested by centrifugation at 1000 xg for 10 min. Pelleted cells were either re-suspended in TP media or, for ROS manipulation assays, a phototaxis buffer described previously by Ueki et al. (2016) (5 mM Hepes pH 7.4, 0.2 mM EGTA, 1 mM KCl, and 0.3 mM $CaCl_2$). The assays took place in 12-well dishes with a thin layer of TP agar (0.8%) on the well bottom and approximately 1.5×10^7 cells in 400 µL of homogenous suspension laid on top. The dishes were illuminated from one direction with 150 µmol photons m^{-2} s^{-1} illumination for up to 3 h. Plates were imaged using a Flatbed Scanner at specified intervals.

Growth of Arabidopsis

Arabidopsis seeds were sown on moist F2+S soil and stratified in the dark at 4°C for 2 days. For growth experiments seeds were grown in a Percival SE-41AR3cLED chamber (CLF PlantClimatics GmbH, Wertingen, Germany) equipped with cool white LED lights under 12 h light (175-180 µmol photons m^{-2} s^{-1})/12 h dark cycles at 21°C, respectively and 70% relative humidity.

Chlamydomonas Confocal Microscopy

Transgenic fluorescent strains were initially grown heterotrophically in TAP media until
reaching $2 - 4 \times 10^6$ cells mL^{-1} and resuspended in TP media overnight prior to
imaging. Cells were mounted on 8-well chamber slides and overlayed with 1.5% low
melting point agarose made with TP-medium. Images were collected on a LSM880
(Zeiss) equipped with an Airyscan module using a 63× objective. Laser excitation and
Emission setting for each channel used are set as below: Venus (Excitation: 514 nm;
Emission 525 – 500 nm); mScarlet-I (Excitation: 561 nm; Emission 570 – 620 nm);
Chlorophyll (Excitation: 633 nm; Emission 670 – 700 nm).

Slimfield microscopy

Chlamydomonas lines *bst4::BST4-mScarlet-I* and the unlabelled control line *bst4* were
prepared overnight in TP media. Each was harvested and spotted onto a slide-
mounted agar pad (GeneFrames, ThermoFisher), consisting of TP media with 1.5%
low melting point agarose. Fluorescence imaging with single-molecule sensitivity was
performed using a custom Slimfield microscope (Syeda et al., 2019). The setup used
a high-magnification objective (NA 1.49 Apo TIRF 100× oil immersion, Nikon) and the
detector was a Prime95B sCMOS camera (Teledyne Photometrics) operating in 12-
bit 'sensitivity' gain at a high total magnification of 53 nm/pixel. The samples were
illuminated either in brightfield, or for Slimfield fluorescence in camera-triggered
frames by a collimated 561 nm wavelength, Gaussian mode OPSL laser (Coherent,
Obis LS) at a peak intensity of 5 kW/cm^2 at the sample plane. This beam was tilted
into a HILO configuration (Payne-Dwyer and Leake, 2022) to reduce out-of-focus
excitation while retaining quantitative molecular sensitivity. The fluorescence image
was split into two parallel channels comprising emission bandpass filters (Semrock
BrightLine®): one with a 585/15 emission filter (central wavelength/spectral bandwidth
in nm) optimized to isolate the mScarlet-I signal, and a second with a 525/25 emission
filter, used only to indicate autofluorescence background. The total length of each
acquisition sequence was ~5 s; sufficient to observe the full course of mScarlet-I
photobleaching, from the initial unbleached state to single-molecule blinking, while
also rapid enough (10 ms exposure/frame at 180 frames/s) to capture the motion of
individual molecular assemblies.

Single particle tracking and molecular counting

Slimfield image sequences were segmented manually in ImageJ to include only the
pyrenoids in downstream analysis. The centroid positions of fluorescent tracks were
identified from local intensity maxima in each frame using ADEMScode software in
MATLAB (Wollman et al., 2022). The summed intensity of each candidate track was
calculated in each frame by adding all pixel values within 5 pixels of the centroid, then
subtracting the local background averaged between 5-8 pixels from the centroid.
Candidates with a summed intensity <0.4× the standard deviation in the background
region were discarded.

Fluorescent proteins are known to exhibit a characteristic integrated intensity per
molecule under stable Slimfield imaging conditions and within the quasi-uniformly
illuminated area within half the beam waist (Shepherd et al., 2021). After sufficient
photobleaching of mScarlet-I in the Slimfield image sequences, only step-like blinking
was observed at the end of each track. The modal integrated intensity of these steps
was used to estimate this characteristic single molecule brightness, equivalent to 56
± 9 photoelectrons per frame per molecule.

At the start of each track, we obtained an initial integrated intensity (independent of
photobleaching) by linearly extrapolating the summed intensity backwards over the
first 4 frames of the exposure. This initial intensity was then divided by the
characteristic brightness of a single mScarlet-I to estimate the number of molecules,
or stoichiometry, in that track. This estimate was precise enough to detect
stoichiometry steps of up to 12 tagged molecules without ambiguity.

Stoichiometry distributions may exhibit peaks which are separated by a characteristic
interval. The smallest consistent interval between peaks can be used to infer the size
of a physical repeat unit or 'periodicity' within assemblies (Hunter et al., 2022; Payne-
Dwyer et al., 2022). A kernel width of 0.7 molecules was chosen to generate the
stoichiometry distribution (Fig. 2F), reflecting the background standard deviation.
Peaks were pinpointed using MATLAB's *findpeaks*.

The intervals between all peaks for each acquisition were aggregated across the
pyrenoid population, weighted by inverse square-root distance (thereby accounting for
shot-noise in broader intervals). A second distribution (Fig. 2G) was then generated
from this weighted population of intervals. The kernel width in this estimate was 0.7

molecules multiplied by the square root of the mean stoichiometry divided by the root number of intervals (thereby accounting for shot-noise in intervals between peaks of higher stoichiometry). The periodicity was then reported as the mode of this distribution and its 95% confidence interval.

Arabidopsis Confocal microscopy

Small sections of 3–4-week-old leaf tissue (~5-10 mm^2) were adhered to slides using double-sided tape with basal side up. A x40 water immersion objective lens was used. Samples were excited by 488 nm at 1% laser power, chlorophyll autofluorescence was collected at 680 – 750 nm and mNeonGreen fluorescence at 503–532 nm. For dual tagged lines, we used sequential acquisition to minimise bleed-through. mCherry was excited using the 542 nm laser and emission collected at 601-620 nm and mNeon as before. Images were acquired using the SP8 Confocal system and Leica LAS AF software (http://www.leica-microsystems.com/). Figures were prepared using ImageJ (http://fiji.sc/Fiji).

Immunoblot detection

Two leaf disks (6 mm diameter) were harvested and immediately frozen in liquid nitrogen. Two steel balls (3 mm) were added and tissue was homogenized using a tissue-lyser twice for 30 Hz for 30 s. A four times volume of cold extraction buffer (20 mM Tris-HCl pH = 7.5, 5 mM MgCl$_2$, 300 mM NaCl, 5 mM DTT, 1% (v/v) Triton X-100, 1 x Protease inhibitor (Roche)) was added and samples vortexed for 30 s. Samples were solubilized on ice for 5 mins and then centrifuged at 5000 x g for 5 min at 4℃. 17.5 µL of supernatant was used to make up 1x LDS and 100 uM DTT. 20 µL was loaded on a Novex™ 4-12% Bis-Tris Mini Gel, (Thermo Fisher, Catalogue number: NP0322BOX). The gel was run at 150 V for 60 minutes. Proteins were transferred to a nitrocellulose membrane using an iBlot 2, programme 0. The membrane was probed with primary antibody in 5% milk in 1 x TBST at the following dilutions: BST4 (1:1000; generated for this study, peptide from C-terminus: SDTELSEANRPRTRPDWRN) (YenZym, Antibodies LLC, USA), AtpB (1:2000; Agrisera:AS05085), RbcL (1:1000; kind gift from Griffiths lab), CP43 (1:3000; Agrisera: AS111787), PsaB (1:1000; Agrisera: AS10695), PsbO (1:2000; Agrisera:AS06142-33). Secondary antibody (goat α-rabbit IR-800; Li-COR: 925-32211). Membrane was imaged using the Li-COR Odyssey CLx scanner.

In order to quantify BST4 protein in Chlamydomonas lines, cells were grown in TP media at ambient CO_2 until reaching 2-4 x 10^6 cells mL^{-1}. Cells were harvested by centrifugation at 1000 x g for 10 mins, normalized to Chl content and resuspended in the extraction buffer described above. Samples were freeze-thawed three times and spun at 20, 000 x g for 20 min at 4°C. Protein extractions containing 5 μg of Chl with 1 x SDS loading buffer were boiled at 100°C for 5 min and loaded onto a 4-20% polyacrylamide gel (Mini Protean TGX, Biorad Laboratories). Proteins were transferred to a PVDF-FL membrane on a Bio-rad semidry blotting system. BST4 primary antibody was used as described above alongside alpha-tubulin primary antibody raised in mouse (Agrisera), as a loading control. Anti-rabbit and anti-mouse fluorescent secondary antibodies, Invitrogen AlexaFluor 488 and 555 respectively, were used at a 1:20 000 dilution. Immunoblots were imaged using an Amersham typhoon 5 scanner with 488 and 535 excitation lasers and Cy2 and Cy3 emission filters. BST4 band fluorescent intensity was quantified using FIJI (Image J) (Schindelin et al., 2012) and normalized to alpha-tubulin loading control. All Chlamydomonas lines for quantification were extracted and analysed in triplicate. For LHCSR3 protein quantification, cells were seeded at 0.1 OD_{750} in TAP media for 4 days and then switched to TP media at a concentration of 30 μg Chl ml^{-1} and exposed to 150 μmol photons m^{-2} s^{-1} for 3 h. Protein was extracted according to Burlacot et al. 2022 and separated by SDS-PAGE as described for BST4 immunoblots.

Oocytes for expression of BST4 were collected after recording and prepared for western blot as described in Lefoulon et al., 2018. The BST4 primary antibody was used as described above and the secondary antibody used was horseradish peroxidase-coupled goat, anti-rabbit (dilution 1:10000 Abcam). Proteins were detected with ECL Advance kit (GE Healthcare,Poole UK).

Blue-Native PAGE
A crude thylakoid enrichment was performed according to (Aro et al., 2004). Thylakoid membranes were solubilized in 0.5% DDM, 1x NativePAGE buffer (Thermo: BN2003), 1x cOmplete protease inhibitor tab (Roche;10x stock made by dissolving 1 tablet in 1 mL dH_2O) for at a concentration of 0.8 μg Chl/μl for 15 min on ice. Unsolubilized material was removed by two rounds of centrifugation at 17,000 xg at 4°C for 15 min. 19.5 μL of supernatant was combined with 0.5 μL of Coomassie additive and loaded

immediately onto a 4-16% Bis-tris gel (Thermo: BN1002BOX). Electrophoresis was

performed at room temperature at 150 V for 90 min. Cathode buffer was swapped

from dark to light when the dye front was a third way through the gel. Separated

proteins were transferred to a nitrocellulose membrane by electrophoresis at 100 V for

90 min at 4 ℃.

Proteins were visualized using chemiluminescence. Secondary antibody (goat α-

rabbit, HRP;1:10,000; Abcam: ab6721). Chemiluminescence substrate SuperSignal

West Pico PLUS (ThermoScientific, ref number: 34579) according to manufacturer's

instructions. Chemiluminescence was detected using clear blue X-Ray Film CL-

Xposure™ Film (ThermoScientific, ref number: 34090).

Chlorophyll fluorescence measurements in Chlamydomonas

To measure PSII activity in Chlamydomonas, cells were grown in TAP in low light (20

μmol photons m^{-2} s^{-1}) for 4 days at 50 rpm to reach logarithmic phase. Cells were

then washed, resuspended to 30 μg Chl mL^{-1} in TP media and exposed to 150 μmol

photons m^{-2} s^{-1} light for 3 h, followed by 1 h incubation in darkness at 50 rpm before

recording Chl fluorescence using DUAL-PAM-100 (Walz, Effeltrich, Germany). A

saturating pulse of 3,000 μmol photons m^{-2} s^{-1} of 800 ms was applied to the samples

in a cuvette under continuous stirring for determination of the maximal fluorescence

yield in the dark state (F_m) and maximal fluorescence yield during the period with

actinic light (F_m'). The maximal photochemical efficiency of PSII (F_v/F_m) was

calculated. NPQ was determined from slow kinetics during actinic illumination at 1,500

μmol m^{-2} s^{-1} for 17 min followed by 5 min of dark relaxation. NPQ and Y(II) were

calculated based on changes in Chl fluorescence as (F_m-F_m')/F_m' and ($F_m'-F$)/ F_m',

respectively, according to Genty et al., (1989).

Electrochromic shift (ECS) in Chlamydomonas

ECS measurements in Chlamydomonas were carried out using a the Dual-PAM-100

equipped with a P515/535 module (Walz). Cells grown and prepared as for Chl

fluorescence experiments described previously were layered on a glass slide and

exposed to actinic red light for the given period. The light was switched off and decay

kinetics were measured. PMF size was calculated as the difference between the ECS

signal in light and the minimum value of the ECS signal immediately after the light was

turned off. Calculation of ΔpH and ΔΨ was performed using the steady-state time point

of the ECS signal in darkness (Cruz et al., 2001). Before each ECS measurement, a

3 saturating 50-µs actinic red flashes of 200,000 µmol photons m^{-2} s^{-1} was applied to

determine the ECS_{ST}; subsequently, the ECS_{ST} amplitude was used to normalize the

ECS signal before the calculation of PMF size and partitioning values. To determine

H^+ conductivity of the ATP synthase (g_{H^+}), the light was switched off at specific time

points to record the ECS signal decay during 620 ms dark intervals. The g_{H^+} parameter

was calculated as $1/\tau$ (time constant for decay during the first 100 ms (Cruz et al.,

2005). The total proton flux across the membrane was calculated as $v_{H^+} = PMF \times g_{H^+}$

(Cruz et al, 2001).

Chlorophyll fluorescence measurements for Arabidopsis

Plants were grown for 8 weeks on S-Hasselfors soil in a Percival AR-82L chamber

(CLF Plant Climatics, Wertingen, Germany) using 12 h light (180 µmol photons m^{-2}

s^{-1})/ 12h dark cycles at 21°C /19°C, respectively, and 70% relative humidity. Slow

kinetics of Chl *a* fluorescence induction were recorded with a pulse-amplitude

modulated fluorometer DUAL-PAM 100 equipped with DUAL-DB and DUAL-E emitter-

detector module (Walz) on attached leaves of 30 min dark-adapted plants using actinic

red light of 830 µmol photons m^{-2} s^{-1} for 10 min, followed by a 5 min dark period. The

saturating pulse applied was 5,000 µmol photons m^{-2} s^{-1} and of 800 ms duration. NPQ

and (Y(II)) were calculated based on changes in Chl fluorescence as $(F_m-F_m')/F_m'$ and

$(F_m'-F)/F_m'$, respectively (Genty et al., 1989).

Electrochromic shift (ECS) for Arabidopsis

ECS was recorded with a DUAL-PAM 100 system equipped with a P515/535

emitter/detector module (Walz). First, plants were dark adapted for 30 min, then

illuminated with actinic red light at 830 µmol photons m^{-2} s^{-1} for 3 min followed by a

60 s dark period in which the ECS decay kinetics were recorded. Before each

measurement, three pulses of 5 µs and 200,000 µmol photons m^{-2} s^{-1} were applied to

determine ECS_{ST}, which was used to normalize the ECS_T values of each

measurement.

Sedimentation of proto-pyrenoid

Cross-linked samples were prepared by vacuum infiltrating intact leaves 1%

formaldehyde prior to sedimentation 200 mg of leaf tissue was flash frozen in liquid

nitrogen and ground by bead beating twice at 30 Hz for 30 s. Four times volume of

extraction buffer (50 mM HEPES-KOH pH 7.5, 17.4% (v/v) glycerol, 2% (v/v) Triton X-

100, cOmplete protease inhibitor tab) was added and sample mixed by bead beating

again. Extract was filtered through one layer of miracloth. A small aliquot of filtered

extract was saved as the input. Extract was then centrifuged at 500 x g for 3 min at 4

℃ and the pellet discarded. The supernatant was centrifuged at 500 x g for 12min, 4

℃. The pellet was washed once with extraction buffer and then centrifuged again. The

pellet was resuspended in 100 µl extraction buffer then centrifuged for a further 5 min.

Pellet was finally resuspended in 25 µl extraction buffer. Fractions were made up in

1x LDS loading buffer and 200 mM DTT. Ten microlitres was subjected to SDS-PAGE

(NuPAGE™ 4-12% Bis-Tris Mini Gel, Thermo Fisher, Catalogue number:

NP0322BOX) at 150 V for 60 min.

Chloroplast fractionation

In order to biochemically localize transgenically expressed BST4, chloroplasts were

fractionated as described in Herdean et al. (2016) using 100 g of leaf tissue from four-

five week-old BST4 transgenic plants. The stromal fraction was concentrated using a

10,000 MWCO centrifugal concentrator (Sartorius Stedim Biotech GmbH, product

number: VS1502).

Electron microscopy of Chlamydomonas

High (3%) and low (0.04%) CO_2 acclimated cells were harvested by centrifugation

(×1000g, 4 minutes, 20°C). Primary fixation was performed in 1.25% glutaraldehyde

in 50 mM sodium cacodylate (pH 7.15) in TP medium for 15 minutes followed by 2.5%

Glutaraldehyde in 50 mM cacodylate for 2 hours. Fixed samples were washed three

times with 50 mM sodium cacodylate by centrifugation. Samples were then osmicated

with 1% OsO4 in 50 mM sodium 25 cacodylate for 1 hour on ice and washed with de-

ionised water. Samples were block stained in 1% uranyl acetate in the dark for 1 hour.

Samples were washed twice with dH_2O and twice with 50 mM sodium cacodylate.

Fixed samples were dehydrated in an acetone series (25%, 50%, 75%, 90% and

100%) ~20 minutes each step. Dehydrated samples were infiltrated with Spurr's resin

by incubating in 25% then 50% Spurr resin in acetone for 30 minutes and transferred

to 75% for 45 minutes at room temperature. They were then incubated in 100% Spurr

resin overnight before polymerising at 70 °C for 24 hours. Sections ~70 nm thick were

collected on Copper grids and stained with saturated uranyl acetate and lead citrate.

Images were collected with a FEI Tecnai 12 BT at 120kV using a Ceta camera.

Electron microscopy of Arabidopsis

Leaves were cut into small 5 mm strips and fixed in 4% (v/v) Paraformaldehyde/0.5%

glutaraldehyde in 0.05M sodium cacodylate (pH = 7.2) by vacuum infiltration three

times for 15 min and incubation at 4 °C overnight with gentle agitation followed by

dehydration in increasing amounts of ethanol 50/70/80/90/100% 1 hr each then

overnight rotation. 100% ethanol was repeated 3 times, 1 hr each and a final over

night at 4 °C. Samples were then fixed in LR resin by infiltrating with increasing

concentration (50/70/100%) with a repeat of 100% and then polymerised overnight at

50 °C. Ultrathin sections were cut and mounted onto plastic-coated copper grids. Grids

were stained with 2% uranyl acetate and visualized by transmission electron

microscopy (TEM).

Protease protection assay

Investigation of the orientation of BST4 in isolated thylakoid membranes was

conducted as described in Stepien and Johnson (2018). Briefly, trypsin made up in 50

mM acetic acid according to the manufacturer's instructions (Thermo Scientific, ref

number: 90057). To disrupt the thylakoid membrane and allow degradation to lumen-

facing peptides, 1% (v/v) Triton X-100 was added, and tubes gently agitated prior to

addition of trypsin.

Xenopus oocyte electrophysiology

Destination clones containing BST4 for Xenopus expression were linearized with

EcoRI, before proceeding to the *in vitro* transcription (mMessage mMachine T7

transcription kit, Thermofisher Scientific, ref number AM1344). Stage IV oocytes were

injected with 20 ng of RNA per oocyte. Measurements of ion transport was done by

voltage clamp using an Axoclamp 2B amplifier (Axon instruments, Foster City, CA)

(Lefoulon et al., 2014; Leyman et al., 1999). They were performed under perfusion of

either 100 mM KCl, K-HCO$_3$, Na-HCO$_3$, K-PEP, or K-Gluconate, with 1 mM CaCl$_2$, 1.5

mM MgCl$_2$, and 10 mM HEPES-NaOH, pH 7.3. Recordings were obtained and

analysed using Henry IV software (Y-Science, Glasgow, UK). BST4 expression in

oocytes was validated by western blotting (Fig S10B), as described above.

ACKNOWLEDGEMENTS

The authors thank the University of York Horticulture team who helped with setting up the Chlamydomonas phototaxis assays. TEM for Arabidopsis was carried out with the support of Stephen Mitchell and the Wellcome Trust Multi User Equipment Grant (WT104915MA)

For the purpose of open access, a Creative Commons Attribution (CC BY) licence is applied to any Author Accepted Manuscript version arising from this submission.

AUTHOR CONTRIBUTIONS

LA, CW, CSL, CS, AM and LM conceived and designed the study with input from all authors. LA performed all Arabidopsis experiments, BN-PAGE and immunoblots. CW performed Chlamydomonas physiology experiments and immunoblots. CSL performed Chlamydomonas microscopy imaging and immunoblots. KMS, KAvM, and ED performed Chlamydomonas and Arabidopsis Chl fluorescence and ECS experiments and generated figures. APD and CW performed molecular tracking experiments with input from ML. CL and MB performed xenopus oocyte work. PG and JB did *in silico* structure predictions. TEM contributed to the complementation of Chlamydomonas *bst4* mutant.   NA contributed to Arabidopsis transgenic line generation. GP assisted with experimental design and data interpretation. LA and CW generated figures. LA led the writing of the manuscript with input from CW. All other authors provided manuscript feedback.

REFERENCES

Adler L, Díaz-Ramos A, Mao Y, Pukacz KR, Fei C, Mccormick AJ (2022) New
    horizons for building pyrenoid-based CO_2 -concentrating mechanisms in plants
    to improve yields. Plant Physiol 190: 1609-1627

Aro E-M, Rokka A, Vener A V (2004) Determination of Phosphoproteins in Higher
    Plant Thylakoids BT - Photosynthesis Research Protocols. *In* R Carpentier,
    ed,Humana Press, Totowa, NJ, pp 271–285

Atkinson N, Feike D, Mackinder LCM, Meyer MT, Griffiths H, Jonikas MC, Smith AM,
    Mccormick AJ (2016) Introducing an algal carbon-concentrating mechanism into
    higher plants: Location and incorporation of key components. Plant Biotechnol J
    14: 1302–1315

Atkinson N, Leitão N, Orr DJ, Meyer MT, Carmo-Silva E, Griffiths H, Smith AM,
    McCormick AJ (2017) Rubisco small subunits from the unicellular green alga
    Chlamydomonas complement Rubisco-deficient mutants of Arabidopsis. New

Phytol 214: 655–667

Atkinson N, Mao Y, Chan KX, McCormick AJ (2020) Condensation of Rubisco into a proto-pyrenoid in higher plant chloroplasts. Nat Commun 11: 6303

Bergner SV, Scholz M, Trompelt K, Barth J, Gäbelein P, Steinbeck J, Xue H, Clowez S, Fucile G, Fufezan C, et al (2015) STATE TRANSITION7-Dependent Phosphorylation Is Modulated by Changing Environmental Conditions , and Its Absence Triggers Remodeling of Photosynthetic Protein Complexes 1. Plant Physiol *168*: 615-634

Bharill S, Fu Z, Palty R, Isacoff EY (2014) Stoichiometry and specific assembly of Best ion channels. Proc Natl Acad Sci U S A 111: 6491–6496

Burlacot A, Dao O, Auroy P, Cuiné S, Li-Beisson Y, Peltier G (2022) Alternative photosynthesis pathways drive the algal CO_2-concentrating mechanism. Nature 605: 366–371

Caspari OD, Meyer MT, Tolleter D, Wittkopp TM, Cunniffe NJ, Lawson T, Grossman AR, Griffiths H (2017) Pyrenoid loss in *Chlamydomonas reinhardtii* causes limitations in CO_2 supply, but not thylakoid operating efficiency. J Exp Bot 68: 3903–3913

Choi BY, Kim H, Shim D, Jang S, Yamaoka Y, Shin S, Yamano T, Kajikawa M, Jin ES, Fukuzawa H, et al (2022) The Chlamydomonas bZIP transcription factor BLZ8 confers oxidative stress tolerance by inducing the carbon-concentrating mechanism. Plant Cell 34: 910–926

Ciorba MA, Heinemann SH, Weissbach H, Brot N, Hoshi T (1999) Regulation of voltage-dependent K^+ channels by methionine oxidation: effect of nitric oxide and vitamin C. FEBS Lett 442: 48–52

Cruz JA, Avenson TJ, Kanazawa A, Takizawa K, Edwards GE, Kramer DM (2005) Plasticity in light reactions of photosynthesis for energy production and photoprotection. J Exp Bot 56: 395–406

Cruz JA, Sacksteder CA, Kanazawa A, Kramer DM (2001) Contribution of electric field ($\Delta\psi$) to steady-state transthylakoid proton motive force (pmf) in vitro and in vivo. Control of pmf parsing into $\Delta\psi$ and ΔpH by ionic strength. Biochemistry 40: 1226–1237

Emrich-Mills TZ, Yates G, Barrett J, Girr P, Grouneva I, Lau CS, Walker CE, Kwok TK, Davey JW, Johnson MP, et al (2021) A recombineering pipeline to clone large and complex genes in Chlamydomonas. Plant Cell 33: 1161–1181

Engel BD, Schaffer M, Kuhn Cuellar L, Villa E, Plitzko JM, Baumeister W (2015) Native architecture of the Chlamydomonas chloroplast revealed by in situ cryo-electron tomography. Elife 4: e11383

Engler C, Youles M, Gruetzner R, Ehnert T-M, Werner S, Jones JDG, Patron NJ, Marillonnet S (2014) A Golden Gate Modular Cloning Toolbox for Plants. ACS Synth Biol 3: 839–843

Erickson E, Wakao S, Niyogi KK (2015) Light stress and photoprotection in *Chlamydomonas reinhardtii*. Plant J 82: 449–465

Fei C, Wilson AT, Mangan NM, Wingreen NS, Jonikas MC (2022) Modelling the pyrenoid-based CO_2-concentrating mechanism provides insights into its operating principles and a roadmap for its engineering into crops. Nat Plants 8: 583–595

Genkov T, Meyer M, Griffiths H, Spreitzer RJ (2010) Functional hybrid rubisco enzymes with plant small subunits and algal large subunits: Engineered rbcS cDNA for expression in chlamydomonas. J Biol Chem 285: 19833–19841

Genty B, Briantais JM, Baker NR (1989) The relationship between the quantum yield

of photosynthetic electron transport and quenching of chlorophyll fluorescence. Biochim Biophys Acta - Gen Subj 990: 87–92

Grefen C, Chen Z, Honsbein A, Donald N, Hills A, Blatt MR (2010) A Novel Motif Essential for SNARE Interaction with the K^+ Channel KC1 and Channel Gating in Arabidopsis . Plant Cell 22: 3076–3092

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. Syst Biol 59: 307–321

Hagino T, Kato T, Kasuya G, Kobayashi K, Kusakizako T, Hamamoto S, Sobajima T, Fujiwara Y, Yamashita K, Kawasaki H, et al (2022) Cryo-EM structures of thylakoid-located voltage-dependent chloride channel VCCN1. Nat Commun 13: 2505

Hasegawa M, Shiina T, Terazima M, Kumazaki S (2010) Selective excitation of photosystems in chloroplasts inside plant leaves observed by near-infrared laser-based fluorescence spectral microscopy. Plant Cell Physiol 51: 225–238

He S, Chou H, Matthies D, Wunder T, Meyer MT, Martinez-sanchez A, Jeffrey PD, Port SA, He G, Chen VK, et al (2020) The Structural Basis of Rubisco Phase Separation in the Pyrenoid. 6: 1480–1490

Hennacy JH, Jonikas MC (2020) Prospects for Engineering Biophysical CO_2 Concentrating Mechanisms into Land Plants to Enhance Yields. Annu Rev Plant Biol 71: 461-485

Herdean A, Teardo E, Nilsson AK, Pfeil BE, Johansson ON, Ünnep R, Nagy G, Zsiros O, Dana S, Solymosi K, et al (2016) A voltage-dependent chloride channel fine-tunes photosynthesis in plants. Nat Commun 7: 11654

Horton P, Long SP, Smith P, Banwart SA, Beerling DJ (2021) Technologies to deliver food and climate security through agriculture. Nat Plants 7: 250–255

Hunter P, Payne-Dwyer AL, Shaw M, Signoret N, Leake MC (2022) Single-molecule and super-resolved imaging deciphers membrane behavior of onco-immunogenic CCR5. iScience 25: 105675

Izumi M, Tsunoda H, Suzuki Y, Makino A, Ishida H (2012) RBCS1A and RBCS3B, two major members within the Arabidopsis RBCS multigene family, function to yield sufficient Rubisco content for leaf photosynthetic capacity. J Exp Bot 63: 2159–2170

Karlsson J, Ciarke AK, Chen ZY, Hugghins SY, Park Y ll, Husic HD, Moroney J V., Samuelsson G (1998) A novel α-type carbonic anhydrase associated with the thylakoid membrane in *Chlamydomonas reinhardtii* is required for growth at ambient CO_2. EMBO J 17: 1208–1216

Katoh K, Standley DM (2013) MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Mol Biol Evol 30: 772–780

Le SQ, Gascuel O (2008) An Improved General Amino Acid Replacement Matrix. Mol Biol Evol 25: 1307–1320

Lee S, Yoon B-E, Berglund K, Oh S-J, Park H, Shin H-S, Augustine GJ, Lee CJ (2010) Channel-mediated tonic GABA release from glia. Science 330: 790–796

Lefoulon C, Waghmare S, Karnik R, Blatt MR (2018) Gating control and K^+ uptake by the KAT1 K^+ channel leaveraged through membrane anchoring of the trafficking protein SYP121. Plant Cell Environ 41: 2668–2677

Li X, Zhang R, Patena W, Gang SS, Blum SR, Ivanova N, Yue R, Robertson JM, Lefebvre PA, Fitz-Gibbon ST, et al (2015) An indexed, mapped mutant library enables reverse genetics studies of biological processes in chlamydomonas

reinhardtii. Plant Cell 28: 367–387

Long S, Burgess S, Causton I (2019) Redesigning Crop Photosynthesis. Sustain. Glob. Food Secur. Nexus Sci. Policy. pp 131–141

Ma Y, Pollock S V., Xiao Y, Cunnusamy K, Moroney J V. (2011) Identification of a novel gene, *CIA6*, required for normal pyrenoid formation in *Chlamydomonas reinhardtii*. Plant Physiol 156: 884–896

Mackinder LCM, Chen C, Leib RD, Patena W, Blum SR, Rodman M, Ramundo S, Adams CM, Jonikas MC (2017) A Spatial Interactome Reveals the Protein Organization of the Algal CO_2-Concentrating Mechanism. Cell 171: 133–147

McGrath JM, Long SP (2014) Can the cyanobacterial carbon-concentrating mechanism increase photosynthesis in crop species? A theoretical analysis. Plant Physiol 164: 2247–2261

Meyer MT, Itakura AK, Patena W, Wang L, He S, Emrich-Mills T, Lau CS, Yates G, MacKinder LCM, Jonikas MC (2020) Assembly of the algal CO_2-fixing organelle, the pyrenoid, is guided by a Rubisco-binding motif. Sci Adv 6: eabd2408

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R (2020) IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol Evol 37: 1530–1534

Mukherjee A, Lau CS, Walker CE, Rai AK, Prejean CI, Yates G, Emrich-Mills T, Lemoine SG, Vinyard DJ, Mackinder LCM, et al (2019) Thylakoid localized bestrophin-like proteins are essential for the CO_2 concentrating mechanism of Chlamydomonas reinhardtii. Proc Natl Acad Sci U S A 116: 16915–16920

Payne-Dwyer AL, Leake MC (2022) Single-Molecular Quantification of Flowering Control Proteins Within Nuclear Condensates in Live Whole Arabidopsis Root BT - Chromosome Architecture: Methods and Protocols. *In* MC Leake, ed,Springer US, New York, NY, pp 311–328

Payne-Dwyer AL, Syeda AH, Shepherd JW, Frame L, Leake MC (2022) RecA and RecB: probing complexes of DNA repair proteins with mitomycin C in live Escherichia coli with single-molecule sensitivity. J R Soc Interface 19: 20220437

Plank M, Wadhams GH, Leake MC (2009) Millisecond timescale slimfield imaging and automated quantification of single fluorescent protein molecules for use in probing complex biological processes. Integr Biol (Camb) 1: 602–612

Price GD, Pengelly JJL, Forster B, Du J, Whitney SM, von Caemmerer S, Badger MR, Howitt SM, Evans JR (2013) The cyanobacterial CCM as a source of genes for improving methylation and chromatin patterning photosynthetic CO_2 fixation in crop species. J Exp Bot 64: 753–768

Qu Z, Cui Y, Hartzell C (2006) A short motif in the C-terminus of mouse bestrophin 4 inhibits its activation as a Cl^- channel. FEBS Lett 580: 2141–2146

Qu Z, Hartzell HC (2008) Bestrophin Cl^- channels are highly permeable to HCO_3^-. Am J Physiol Physiol 294: C1371–C1377

Roberts SK, Milnes J, Caddick M (2011) Characterisation of *An*BEST1 , a functional anion channel in the plasma membrane of the filamentous fungus , *Aspergillus nidulans*. Fungal Genet Biol 48: 928–938

Santhanagopalan I, Wong R, Mathur T, Griffiths H (2021) Orchestral manoeuvres in the light: Crosstalk needed for regulation of the Chlamydomonas carbon concentration mechanism. J Exp Bot 72: 4604–4624

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, et al (2012) Fiji: an open-source platform for biological-image analysis. Nat Methods 9: 676–682

Shepherd JW, Payne-Dwyer AL, Lee JE, Syeda A, Leake MC (2021) Combining

single-molecule super-resolved localization microscopy with fluorescence
polarization imaging to study cellular processes. JPhys Photonics. 3: 034010
Stepien P, Johnson GN (2018) Plastid terminal oxidase requires translocation to the
grana stacks to act as a sink for electron transport. Proc Natl Acad Sci U S A
115: 9634–9639
Sueoka N (1960) Mitotic Replication of Deoxyribonucleic Acid in Chlamydomonas
Reinhardi. Proc Natl Acad Sci 46: 83–91
Syeda AH, Wollman AJM, Hargreaves AL, Howard JAL, Brüning J-G, McGlynn P,
Leake MC (2019) Single-molecule live cell imaging of Rep reveals the dynamic
interplay between an accessory replicative helicase and the replisome. Nucleic
Acids Res 47: 6287–6298
Tardif M, Atteia A, Specht M, Cogne G, Rolland N, Brugière S, Hippler M, Ferro M,
Bruley C, Peltier G, et al (2012) PredAlgo: a new subcellular localization
prediction tool dedicated to green algae. Mol Biol Evol 29: 3625–3639
Ueki N, Ide T, Mochiji S, Kobayashi Y, Tokutsu R, Ohnishi N, Yamaguchi K,
Shigenobu S, Tanaka K, Minagawa J, et al (2016) Eyespot-dependent
determination of the phototactic sign in *Chlamydomonas reinhardtii*. Proc Natl
Acad Sci U S A 113: 5299–5304
Wollman AJM, Fournier C, Llorente-Garcia I, Harriman O, Payne-Dwyer AL,
Shashkova S, Zhou P, Liu T-C, Ouaret D, Wilding J, et al (2022) Critical roles for
EGFR and EGFR–HER2 clusters in EGF binding of SW620 human carcinoma
cells. J R Soc Interface 19: 20220088
Woo DH, Han K-S, Shim JW, Yoon B-E, Kim E, Bae JY, Oh S-J, Hwang EM,
Marmorstein AD, Bae YC, et al (2012) TREK-1 and Best1 Channels Mediate
Fast and Slow Glutamate Release in Astrocytes upon GPCR Activation. Cell
151: 25–40
Wu A, Brider J, Busch FA, Chen M, Chenu K, Clarke VC, Collins B, Ermakova M,
Evans JR, Farquhar GD, et al (2023) A cross-scale analysis to understand and
quantify the effects of photosynthetic enhancement on crop growth and yield
across environments. Plant Cell Environ 46: 23–44
Yang T, Liu Q, Kloss B, Bruni R, Kalathur RC, Guo Y, Kloppmann E, Rost B,
Colecraft HM, Hendrickson WA (2014) Structure and selectivity in bestrophin ion
channels. Science 346: 355–359
Zhang R, Patena W, Armbruster U, Gang SS, Blum SR, Jonikas MC (2014) High-
throughput genotyping of green algal mutants reveals random distribution of
mutagenic insertion sites and endonucleolytic cleavage of transforming DNA.
Plant Cell 26: 1398–1409

Supplemental Figures

0.04% CO₂

	Strain	Days 0-3		P(T<=t) two-tail	Days 0 to 5		P(T<=t) two-tail
		Mean	SE	two-tail	Mean	SE	two-tail
SGR (μ h⁻¹)	WT	0.0389	0.0007		0.0241	0.0010	
	bst4	0.0402	0.0003	0.1652	0.0257	0.0001	0.2036
Doubling time (h)	WT	17.8462	0.3380		28.8436	1.2467	
	bst4	17.2379	0.1255	0.1669	26.9798	0.1491	0.2119

3% CO₂

	Strain	Days 0-2		P(T<=t) two-tail	Days 0-3		P(T<=t) two-tail
		Mean	SE	two-tail	Mean	SE	two-tail
SGR (μ h⁻¹)	WT	0.0679	0.0013		0.0456	0.0011	
	bst4	0.0682	0.0010	0.8626	0.0455	0.0003	0.9436
Doubling time (h)	WT	10.2145	0.1930		15.2239	0.3725	
	bst4	10.1669	0.1510	0.8555	15.2353	0.0878	0.9777

Supplemental Table 1 compares specific growth rates (μ h⁻¹) and cell doubling times of WT and *bst4* strains (n=3) during liquid growth assays at 0.04 and 3 % CO_2.

Supplemental Figure 1. BST4 structure and sequence compared to other bestrophins. Phylogenetic analysis of the full length Chlamydomonas BST4 amino acid sequence (bold). The evolutionary history of BST4 was inferred by using the maximum likelihood method based on the Le and Gascuel substitution model with discrete Gamma distribution (5 categories) and 500 bootstrap replicates. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site.

Supplemental Figure 2. Interaction between BST4 and RBCS. A. Yeast-2-hybrid experiment. The disordered region of BST4 C-terminus (a.a 387-end) and with residues WR from RBMs are changed to EE. Abbreviations: AD, activation domain; BD, binding domain; DDO, double drop out media; TDO, triple drop out media **B**. FRET efficiency between BST4-mScarlet-I and CrRBCS-Venus. Sensitized-FRET measurement on dual-tagged cells (BST4-Venus and RBCS1-mCherry) showed a moderate FRET efficiency of 35% (n=10).

Supplemental Figure 3. Validation of Chlamydomonas BST4 lines. **A.** PCR amplification of the *BST4* and control loci of WT (CMJ030 (CC-4533; cw15, mt -) and *bst4* (LMJ.RY0402.159478) gDNA to confirm the insertion of the CIB1 cassette in *bst4*. **B.** Immunoblots confirming the production of BST4 in WT, *bst4, bst4::BST4* and *bst4::BST-truncated* Chlamydomonas lines. **C.** Immunoblot showing the absence of BST4 in the *bst4* knock-out line and presence in the WT and *bst4::BST4* complemented lines. **D.** Western blot used to quantify the amount of BST4 protein detected in WT and *bst4::BST4*. **E.** Immunoblot was used for the quantification of fluorescent intensity as a measure of BST4 protein abundance. All fluorescent intensity measurements were normalized to respective alpha-tubulin loading controls and conducted in triplicate. No statistical difference was observed between WT and *bst4::BST4* BST4 protein abundance (Paired two-tail t-test p = 0.22, n=3)

Supplemental Figure 4. Pyrenoid morphology in *bst4* vs WT. A. The pyrenoid area to cell area ratio of wild type (black) and *bst4* (white) strains at high (3%) and low (0.04%) CO_2 concentrations. Each bar represents the mean value ± SEM (n=40–50). The data for wild type and *bst4* strains were compared using a two sample T-test. The p-values (showing no significant difference) are displayed above the bars. **B.** Pyrenoid area vs cell size of WT and bst4 at 0.04% [CO_2].

Supplemental Figure 5. Localisation of C-terminally truncated BST4 in WT background. BST4$_{\Delta Cterm}$-mScarlet-I fluorescence shown in green and chlorophyll autofluorescence in magenta. Scale is bar 1 µm.

Supplemental Figure 6. Localisation of proteins in a pyrenoid matrix-less background. A. Diagram showing a section of the *Crrbcs::AtRBCS* matrix-less mutant (*-matrix*). The nascent pyrenoid tubules (pt) are shown in dark green, starch granules (sg) in white. **B.** Confocal image of starch binding protein STA2 fused to Venus in *Crrbcs::AtRBCS.* Scale bar is 2 µm. **C.** BST4-Venus in a WT Chlamydomonas cell. Scale bar is 1 µm. **D.** PsaF-mNeon in *Crrbcs::AtRBCS*. Venus and mNeonGreen fluorescence are shown in green and chlorophyll autofluorescence in magenta. Scale bar is 1 µm.

Supplemental Figure 7. **Spot test of WT, *bst4* and BST4 complemented lines under CCM induced conditions**. *Chlamydomonas* strains were grown in serial dilution on agar plates in saturating light (400 µmol m^{-2} s^{-1}) under a range of CO_2 and pH conditions (specified) to induce the CCM. All lines grew comparably to WT across the conditions used and on the low light TAP control plate.

Supplemental Figure 8. **Photosynthetic parameters of WT Chlamydomonas and mutants.** Wild type (WT) and mutants were grown on TAP medium in the light at 20 µmol photons m^{-2} s^{-1}. **A.** Chlorophyll and carotenoid contents of cells were determined in methanol extracts. **B.** In a separate experiment, cells resuspended in TP at 30 µg Chl ml^{-1} were dark-adapted for 1 h and then illuminated for 10 min with 530 µmol photons m^{-2} s^{-1} after which the light was switched off to record ECS in darkness. **C.** Representative Electrochromic shift (ECS) decay curves. The cells were prepared as in B. To determine the g_H^+ parameter, ECS kinetics were recorded during 600 ms dark intervals. The ECS decay of the first 100 ms was fitted to calculate g_H^+ (s^{-1}) =1/time constant for decay. Data are the means ±SEM (n=3 replicates).

Supplemental Figure 9. No currents were detected for BST4 with any anions tested in Xenopus oocytes A. Steady state currents analysis of oocytes injected with BST4, full length or truncated, compared to non-injected oocytes. The voltage steps start from 60 mV to -100 mV (Cl⁻, PEP⁻, and Gluconate conditions) 40 mV to -120 mV (HCO3⁻ conditions, right panel), holding voltage is -20 mV. Recordings were performed on n>4 oocytes. There is no differences between the currents recorded in non-injected oocytes and expressing the protein. **B.** Western Blot analysis of oocytes. Lane 1 is the marker, lane 2 are non-injected oocytes, lane 3 is the full-length protein (about 66 kDa), lanes 3 and 4 are the truncated proteins. We can only detect full length BST4 as the antibody is directed against the C-terminal part of the protein that is removed in the two truncated versions of the channel.

Supplemental Figure 10. Phenotype of BST4 transgenic Arabidopsis line. A. Immunoblot against BST4 of proteins extracted from BST4 no tag lines and Azygous segregants (Az). S2$_{Cr}$ is the parent line. **B.** Fresh weight of 28-day old rosettes, bars represent mean weight for each genotype n=18-21. Different letters indicate statistically significant difference among the genotypes (one-way ANOVA test, followed by Turkey's post hoc test, P < 0.05).

Supplemental Figure 11. Photosynthetic measurements of BST4 transgenic Arabidopsis lines 2 and 3. A. F_v/F_m values measured on attached, 30 min dark adapted leaves of 8-week-old plants (n=15-21). The letters in indicate nonsignificant differences between plants expressing BST4 and their azygous segregants using Tukey post hoc test (P > 0.05). **B.** Non-photochemical quenching (NPQ) as a measure of photoprotection and **C.** photosystem II quantum yield (Y(II)) were recorded during 10 min of illumination at 830 µmol photons m^{-2} s^{-1} followed by a 5-minute dark period. Data are presented as means ± SEM (n=4-6). Asterisks indicate statistical difference between plants expressing BST4 and their Azygous (Az) segregants according to unpaired t-test (P ≤ 0.05). **D.** Proton motive force (PMF) size and **E.** partitioning to pH gradient (ΔpH) after 3 min illumination at 830 µmol photons m^{-2} s^{-1}. Data are means ± SEM (n=5-6). The letters in E and F indicate non-significant differences between plants expressing BST4 and their azygous segregants according to unpaired t-test (P > 0.05).

A. Full length BST4 and homologous sequences

>Brassica_rapa_(PAC:30641593)
```
--------------------------------------------------------------------------------------------
------------RTLYTHEKWVEHRSSLRHVHHLFSSFSSRVILSLIPPVFFFTSVAIFIASYNSAVAL-------------
DWLPSVFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
DDLARQVICSVDGSG-----------------DELVIK-----DLLLRYV----AAFPVALKCHVTYGSD-----VARDLRN---
--LIEGDDLSLI--IESKHR-------------PRCVIEFISQSLQL----LKLDDT--------------
KRDLLESKMLHLHEGIGVCEQLMGIPIPLAYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSES---------------------------
--------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------------------
---------------------------------------------------------------------------
```

>Q9M2D2_-_VCCN1_Athaliana
```
MY-------------------QSMNLSVSSNFTHRSLLES------------RFPI-----FSTGFR-----KSVNLKPPRVSSGPE---
SNDSGH---------------ETLTDKLIHLLRAVP--------
DWADEIKERGMQQKRSLYTHEKWVEHRSSLRHVRHLLSSFSSRVILSLIPPVFFFTSVAVVIASYNSA
VAL-------------DWLPGIFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
NDLARQVICSVDSSG-----------------DELIIK-----DLLLRYI----AAFPVALKCHVIYGSD-----IARDLRN-----
LIEADDLSLI--LQAKHR-------------PRCVIEFISQSIQL----LKLDDA--------------
KRDLLESKMLHLHEGIGVCEQLMGIPIPLSYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSEKVIRNRIIAKIKLHEFK--
-----HSS-------------------------NGRHRS---------------------------------------------------------
--------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------------------
--------------
```

>Nicotiana_tomentosiformis_(XP_009629643.1)
```
M--------------------------TNSRTLFSIQSPTNASFSSHF-TLKTPSK--LQQQSFPSKL-
NFKKLRFSTFKVRCCPQ----QTPQN---------------QNPTSALISILRIIP--------
DWADRIQEEGMKKKRSLYTHESWMQHRSSLRHVRHLFSSLNSRVILSLVPPVIAFTSVAVVIASYNSA
VSM-------------HWLPELFPVLR---ASPLPYQLTAPALALLLVFRTEASYSRFETGKKAWTKVIAGT--
NDFARQVIACVDKS-------------------DAVLK-----AALLQYI----MAFPVALKCHITYGSD-----IASDLKN-----
LLEADDLAVV--LSSKHR-------------PRCIIGFISQCLQS----LHLEGT--------------
KLTQLESKISCFHEGIGVCEQLAGIPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCQL-
VHDNIQESMANEKKIQERLSAKRKRRFSE------HSQ------------------------NGWPTS------------------------
--------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------------------
-----------------------------------------------------------------------
```

>XP_028961536_Malus_domestica
```
MLLPSSLSLQTLAPPNADTIQKPHQTLPQNLTLFQLVHP------QHFPTLQFPN-----LPSGPKTL-
KFKLLCSQSPNPNPSP-------PSS---------------SSPVQTLISILRIIP--------
DWSDRTQERGMRQHRTLYDHEKWMHHRSSYRHLRHLLSSLSSRVILSLIPPVIAFTLVAVVIASYNTA
VAL-------------DLLPGIFPLLR---SSSLPYQLTAPALALLLVFRTEASYSRFEEGRKAWTEVIAGA--
NDFARQIISSVETSG-----------------DAQLK-----KALLQYI----VAFPVALKCHVIYGSD-----IARDLQN-----
LLEVDDLLVV--LNSKHR-------------PGCIIQFISRSLQL----LKLEES--------------
RRIMLQSKISCFHEGIGICEQLIGTPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEQVGVLIEEPFPM----LALDDLCNS-
VRNNVQEALASEKLIRARLAAKGRIQSEQ------QFQ------------------------NGQPRP------------------------
--------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------------------
-----------------------------------------------------------------------
```

>Chlamydomonas_eustigma_(GAX83184.1)
```
M------------------------------------------------------------LVHRVHTRSLGNRNQCGRK---------------------
LHRVSTFVVKTPTEKPVVA----
DYVLPRSEEARRYFRTVYDFPQWQKHRSPTRLIDRLLQIPRSHVLQNILPSIAWCSSVAGLLTLYMQA
YDA-------------HILPDGFPSFATNNACTSFVNTTTVALSLLLVFRTNVSYGRWDEARKMKGLLVNRS--
RDLMRQVCAMVPEE-------------------DVATK-----AMMAKWT----AAFCRVLRIHFQPEVS-----LEDEMKG-
----LLSPEELEWL--IESKHR-------------PCSVIHMLSQIIYD----SQISAI--------------
CQAQMCNNLTAFEDVLGGCERLLRAPIPVSYTRH-TARFLFTWLTLL-------PFALYNSC-----
```

```
GVWTLPVVAGVSAVLCGIEEIGVQIEEPFGI----LPLEAICGR-
IQADVMATLKEDAKTRNLRNKVLLIRGPE------HWAMV----------------------NGHSNK----------------------
----------------------------------------------------------------------------------------------------------------------
----------------------------------------------------------------------------------------------------------------------
------------------------------------------------------------------
```

>Cre16.g662600.t1.2_(BST1)
```
M--------------------------------QMQA-----------NRSSLRASPVRGLGARPLLRALPAGRVARLNVSAQA--
KDPNAPIQ--------------SNPLGTLSSQSGQVA----------
TLPRSEEARKYFRTVYDFPQWQKHRSSYRFAERLFQLSQSHILQNALPAISWVTLVATLVASYGYSY
DQ-------------HMLPDVFPSISPNASCTAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWT----VAFSRALRIHFQPEVT-----IESELQN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VPMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYPTT-----
GWGVVPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-
IQADVMATLKDDADTKTILAEAGLISLIPS----ATSATPVASA----EPVLVSARPSAAPAPNNGLQVRVAM-
----------------------------------------------------------------GG-------------------ERK*----------------------------
----------------------------------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------
```

>Cre16.g663400.t2.1_(BST2)
```
M--------------------------------QC-------------LSSRPVAMGRAGSSALPRL-
PLRAGRVCHLGVRCQAANKDPNAPIQ----------------SNPLGSFSSQLQNQP----------
TLPRSEEARKYFRTVYDFPQWQTHRNQYRLMKRLFSIPQSHVIQNALPSIMWVAFTSTCVAAYMYGY
DQ-------------HMLPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWV----VAFSRALRIHFQPEVT-----IESELKN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VPMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCSGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-
IQADVMATLKDDADTKTILAEAGLISLRA------NSAMAVENALPDLDSINAAAPNGNGSHNGNGAAV-----
----------------------------------------------------------------------------------PVS-------------------------
VSAGA----------------------------------------------------------------------------------------------------------------
----------------------------------------------------------------SGNGMNV-------------------------RISPR*-----
```

>Cre16.g663450.t1.2_(BST3)
```
M--------------------------------QVSK-----------VPSS------ASARCLPRL-
PVRTSRVCQLSVRCQAANKDPNAPIQ----------------SNPLGSFSSQNSSGAVV--------
TAPRNEDARKYFRTVYDFPQWQKHRSQSRLVRRLFTIPQSHVIQNALPSIMWVTFTSTCVAAYMYGY
DL-------------HILPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWT----VAFARALRIHFQPEVT-----IESELQN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VRMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDDADTKTVLAEAGLISLIP-
-----SMSLPPTEHASPSDPVTAAAAAALAAANGNGAASHSNG---------------------------------------------------
----------------------NG--------------------SKPVSTQVPP-----------------PVLAPVTVTS----------------------------
----------------------------------------------------------------------------------------------------------------------
----------------------SSGSMNV-------------------------RISPR*-----
```
>Chlorella_sorokiniana_(PRW33726.1)
```
M-----------------------STAMLAGSRIQLQQPAG--------LGGSRLQ--RAAAPVAAAA-
RLGSVRPAGLQARSTAARRADRSALR---------------VSATASPEAAPVKLS----------
GDDLKEANRKHMRSVFDFDLWKKHRSSSRYLRHIVGLGESRIVSGLMAPLTYVMTLSLAVACYNAAA
EA-------------GYLP-VFPELKL--ATNAPFGLTSFALSLLLVFRTNSSYGRWDEARKMWGLIVNRS--
RDFIRQGLGYIPPE-------------------QEELQ-----KMLVRWT----VAYSRSLMCHLRPGED-----LRVELKD---
--TLKPEELEAL--LASTHR-------------PNYVVQVLTAIIKT----
AQLPAAVTNNRDSTGCVPAGAAYRMDENLTVFADVTGGCERILRTPIPLSYTRH-TSRFMMIWLTLL---
-----PFTLWDSC-----HWAMLPIAGIVSFLLLGIEEIGVQIEEPFTI----LPLEVISRT-
IEGNVWELYRMHSGEA---LEKEQAELAN------GQDVQVLNA----
QDLVALMAPSAVGNTANGTSRKSLV------------------VNYGL----------------------------------------------------
----------------------------------------------------------------------------------------------------------------------
----------------------------------------------------------------------------------------------------------------------
-----------------
```
>Cre06.g261750.t1.2_(BST4)
```
M----------------------QC--------QLKH---------------------------GA-RPQSQRPNWLPARAATLRPAVQHGVR-
```

```
--------------RGLTLGVKAAAAPLE--------
DKKMPADMTTRQYRRVVYDFALWAKHRDVNRYLYNLRTIPGSRIIRQLSQPMGVVLAWAALFGFYET
CLEA-------------GVLPSYLPKMTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-------------------DQAGR-----EAVGKWT----VGFTRALQAHLQEDID-----LRKELEKA--
TPRWSKEEIDML--VNAQHR-------------PIKAISVLSELTRQ----LSITQF--------------
QALQMQENCTFFYDALGGCERLLRTPIPVSYTRH-TARFLTIWLAML-------PLGLWERY-----
HWSMLPVIALIGFLLLGIDEIGISIEEPFGI----LPLDAICGR-
AQTDVNSLLKEDPAVMKYVDDVRSGRVKS------PPPLPPAPA----APAAA-------AAAAAAAARSVSP----
QPDVAKTLGSLFTNVRAGVGAVAPGAPLMPQAPVRSPSPTRSVSP--------
SFPRASAGTGMPPPVGMNG----------ATPRVAAAPPTPPPVSRPAA-----------------
PAAAPAAGSGFTMPNFSASLSGLTGGAAAAAKSAADAASSKLTKMADSMSSGAAAPAPPAAPAR-----
-----------------PSTSPRPSASSPISSSADADRSDSSRR---PVNWRDELQSLKATRE----------------
PNGNGNGSGVAPAA--GRADADEEALRRFGNLAGRSR-------------SGNG---------------
GGGSSDTELSEANRPRTRPDWRNQL*
>XP_002945810.1_Volvox_carteri_f._nagariensis
M----------------------QS--------QLQP---------------------------RL-QLQGTRLNWLPQRSCVQRRSLRVDAT--
-------------SG------AAPPPPA--------
GKELSNDMVTRQYRRTVYDFSLWAKHRDVNRYLYNLKTIPGSRIIRTLGQPMGIVLAWAAMFGFYET
CLES-------------GVLPSYFPKLTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-------------------DVAGR-----EAVGKWA----VGFCRALQAHLQEDAN-----LREELQKA-
-QPRWSREEIDML--CSAQHSWQQLQSCVNAFW-PIKAISMLSELTRQ----LPISQF--------------
QALQMQENVTFFYDALGGCERLLRTPIPVSYTRI-------------------------------------------------------------------
LPLDAICTR-AQTDVVSLLKDDPAVVKYISDVRQGRIAP------PTEPPVAGA----
APVAAAPPPPPPASAGGGISRSGSPTAQQQPDVMKTVTSMLHNVKAGIGAVAPAPPRPPSPQPRAR
SP-RAASPGGPS----PFPRASAGTG--------G----------
AAAAVPSPPPIKPLTSSSSSSSGAVSKDSNNSTATAKKPASAPAASSA----
GFSMGFSGLADGAAAAAKSAS-AAAAKFSKIADSVVAG--TPAAPASEAKRETAA--AAAMQAQPRN---
TPSSSSSTPSAAPANGSSDDDRSSSGRRTAAAVNWREELAALRAGREDAEEPASASASYDREFPSS
SWSFSSASSAAVVQSGDAEDEARRRFGGLAGRGARSDTTTSAAAVMRGNGNGLSENGYGNGYGN
DN-GNGNGNTVEARGARPRTRPDWRNQL-
>Chrysochromulina_sp._(KOO32217.1)
M----------------------REHPLSYEEYMRQRS-----------AGRDPLA--EAVQGQSASM-
ERGVVPPPPVKATPERTEAAFTPPAEFDFFQDVFKPTVESVVKAVVSPGAQQDSDESYMRVPWWE
QGSTYSEDQRKDRRTVFMHDDWKRHRSSERFFRNIKTWPSSGINQALRKELTFVTSVSVFVVLANM
LLYQYQDFGGVVHPGPLSFLDGPIKSLS---
LPALPFSMASPVLSLLLVFRTNTAYFRWNEARTLWGGLINNC--RNIVRQTTTMFPNDA-----------------
YHNALK-----KRLATET----ATFIKSLRNFLRGPEDDAT--LRKELYAYVNQGLMTSAQAEAT--LAAKNR---
----------PMFALAAMSATLRK----ANIDEM--------------YISRMDSTISVLVDLTGANERIFKSPIPLVYTRL-
TARFLSVFLTLL-------PLAMWAALGESWNHWATIPATFILSVFLFGVEEVGIQIEEPFSI----
LPLEAMCNGAIEAVQLEMLAAE----------------------QSQVFEAAG----DAVAVA--------------------------------
-------------------------------------------------------------------------------------------------------------------------
-------------------------------------------------------------------------------------------------------------------------
----------------------------------------------------------------------------
>Emiliania_huxleyi_(XP_005770556.1)
-------------------------------------------------------------------------------------------------------------------------
--
EFKEAGREFRQDVYSYNDWRWHRESGHIASAISSVFTSGVGKAMWRETFFVIATAAAVYLYNIGVPV
LAAKTAASL----PIVAALLGRLPLLHLSLLPLTLSSPALFLLLVFRTNNSYDRWWEARKVWGGVINAS--
RDLARQALALVR--------------------DAELK-----KLMVSQI----ASYARVLKYHLGPPTPEARDLLRNELVD--
--NRLPADQVRVI--MEAKHK-------------PMALLGLVSASLHDSGR-TGLDTV--------------
QASKLDQTLSLLTDYLGKCERIVKTPLPLVYTRH-TARFLSWWLLFL-------PVCLYNQLRA---
NWMIVPVSGLIGFFLVGIEDLGNQIEEPFSI----LPLTAMGTG-IQQSIFEAL------------------------------------
-------------------------------------------------------------------------------------------------------------------------
-------------------------------------------------------------------------------------------------------------------------
--------------------------------------------------------------------------------------------------------------------
>Phaeodactylum_tricornutum_(XP_002180738.1)
MM------------------------------------------------------RNFASVLLLL-SSGAAAFAPVQHNGVRTIATPSTPLY---
-------------GNTKQPPPALP--PIK----------
DISYGEESRKYRRTVYSHDDWVKHRSSDRFLRNLLAIGSSGVYKSLAKEVLATTGVATFIVLYNCLVG
GYTDLEGIKHS---ALIESVWAPLMA--LPLAPFTLSSPSLGLLLVFRTNTSYQRWDEARKNWGMNINHT-
```

-RDLVRMGTSFYDNAA-----------------VSSEQRAKDL-KALSLAT----WSFVRAMKRHLSPESEDEQD-
FRRELFE-----RLPAPQAQAI--IDAAHR-------------PNRALFDLSVAIEN----LPMHFL--------------
RKNQVHQAVTIFEDNLGSSERLLTSPVPLFYSRH-TARFLSFWLLLL-------
PFALWDPFAGTWNHVGMIPATAVISIFLFGIEELATQMEEPFTI----LPMQAFCDK-
IGNWCNEIVSWQAGDNGMAVNMPSMISPEGLPELKEPAPVPAMA----
VASVAAAMPVMANGDINGDTTGITM---DQP---HNAIP-----------------

>Thalassiosira_pseudonana_(XP_002289965.1)
M--
GPPIDPSVPVT-----------
DQVGEGSRKYRRTVYTHDDWVRHRSPDRFGNNLSTLFNSGIYKQVANEVFATTAVATFVFLWNMIA
GGYTDLAGVQHG---PIIDSPLAQMVG--
LPMTAFTILTPSLGLLLVFRTNTSYGRWDEARKMWGLNINHT--
RDLNRMATAWYGNEGNMDSVAFMGGDIPYSQPIDPVQRAYDL-GQVSLFT----
WAFVRSMKRHLSPPEEDEED-FKAELRA-----RLTPEQAENI--INAAHR-------------PNRALFDLSVAIEN--
--LPMHFL--------------RKNAINTNLSIFEDTLGGCERLLSSPVPLFYSRH-TARFLSTWLLLL-------
PFGLYEQFKDSWNHIAMIPATAFISVCLFGIEELATQLEEPFTI----LPMQGFCDK-IGGWCDEIVSW-
AGQGGQQEYTEENAMSNE------QEMTY--

---WR--

>WP_049046555.1_Klebsiella_aerogenes

---------------------MIIRPEQHWFFRLFDWHGSVLSKIVFRLLLNVLMSVIAIISYQWYEQL-------------GI------
-----HLTVAPFSLLGIAIAIFLGFRNSASYSRFVEARNLWGTVLIAE--RTLVRQLKNILPD-------------------
DEETH-----KTLVSYL----VAFSWSLKHQLRK-TD-----PAVDLYR-----LLPKEKVAEI--LASSMP------------
TNRILLLIGNELGRLREQGKLSDI--------------TYGLMDNKLDELAHVLGGCERLASTPVPFAYTLI-
LQRTVYLFCTLL-------PFALVGDL-----
HYMTPFVSVFISYTFLSWDSLAEELEDPFGTSANDLPLNAMCNT-
IERNLMDMTGQHPLPEKMQPDRYYNLT--

>XP_011543531.1_Homo_sapiens
M--
----------------TITYTSQVANARLGSFSRLLLCWRGSIYKLLYGEFLIFLLCYYIIRFIYRLALTEEQQL----------
----MFEKLTL--YCDSYIQL--IPISFVLGFYVTLVVTRW------WNQYENLPWPDRLMSLVSGFVEGK-------
-----------DEQGRL--LRRTLIRYANLGNVLILRSVSTAVYKRFP-----SAQHLVQ---
AGFMTPAEHKQLEKLSLPHNM---------FWVPWVWFANLSMKAWLG---GRIRDPI-------------
LLQSLLNEMNTLRTQCGHLYAYDWISIPLVYTQVVTVAVYSFFLTCLVGRQFLNPAKAYPGHEL---
DLVVPVFTFLQFFFYVGWLKVAEQLINPFGEDDDDFETNWIVDRNLQVSLLAVDEMHQDLPRMEPDM
YWNKPEP------QPPYTAASA----QFRRASFMGSTFNISLNKEEMEFQPNQEDEED-----------
AHAGIIGRFLGLQSHDHHPPRANSRTKLLWPKRESLLHEGLPKNHKAAKQ----
NVRGQEDNKAWKLKAVDAFKSAPLYQRPGYYSAPQ-----------TPLSPTPMFFPLEPSA---
PSKLHSVTGI-DTKDKSLKTVSSGAKKSFELLSES--
DGALMEHPEVSQVRRKTVEFNLTDMPEIPENHLKEPLEQSPTNIHTTLKDHMDPYWALENR----------
SVLHLNQGHCIALCPTPASLALSLPFLHNFLGFHHCQSTLDLRPALAWGIYLATFTGILGKC--------------
SGPFLTSPWY---HPEDFLGPGEGR--------------------

B. BST4 and homologous sequences trimmed after the bestrophin domain

>Brassica_rapa_(PAC:30641593)

------------RTLYTHEKWVEHRSSLRHVHHLFSSFSSRVILSLIPPVFFFTSVAIFIASYNSAVAL-------------
DWLPSVFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
DDLARQVICSVDGSG-----------------DELVIK-----DLLLRYV----AAFPVALKCHVTYGSD-----VARDLRN---
--LIEGDDLSLI--IESKHR-------------PRCVIEFISQSLQL----LKLDDT--------------

KRDLLESKMLHLHEGIGVCEQLMGIPIPLAYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSE
>Q9M2D2_-_VCCN1_Athaliana
MY------------------QSMNLSVSSNFTHRSLLES------------RFPI-----FSTGFR-----KSVNLKPPRVSSGPE---
SNDSGH---------------ETLTDKLIHLLRAVP--------
DWADEIKERGMQQKRSLYTHEKWVEHRSSLRHVRHLLSFSSRVILSLIPPVFFFTSVAVVIASYNSA
VAL-------------DWLPGIFPILR---SSSLPYQLTAPALALLLVFRTEASYSRYEEGRKAWVGIIAGT--
NDLARQVICSVDSSG-----------------DELIIK-----DLLLRYI----AAFPVALKCHVIYGSD-----IARDLRN-----
LIEADDLSLI--LQAKHR-------------PRCVIEFISQSIQL----LKLDDA--------------
KRDLLESKMLHLHEGIGVCEQLMGIPIPLSYTRL-TSRFLVFWHLTL-------PIILWDEC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCDL-VHSNIQEAVKSE
>Nicotiana_tomentosiformis_(XP_009629643.1)
M-------------------------TNSRTLFSIQSPTNASFSSHF-TLKTPSK--LQQQSFPSKL-
NFKKLRFSTFKVRCCPQ----QTPQN---------------QNPTSALISILRIIP--------
DWADRIQEEGMKKKRSLYTHESWMQHRSSLRHVRHLFSSLNSRVILSLVPPVIAFTSVAVVIASYNSA
VSM-------------HWLPELFPVLR---ASPLPYQLTAPALALLLVFRTEASYSRFETGKKAWTKVIAGT--
NDFARQVIACVDKS------------------DAVLK-----AALLQYI----MAFPVALKCHITYGSD-----IASDLKN-----
LLEADDLAVV--LSSKHR-------------PRCIIGFISQCLQS----LHLEGT--------------
KLTQLESKISCFHEGIGVCEQLAGIPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEEVGVLIEEPFPM----LALDELCQL-VHDNIQESMANE
>XP_028961536_Malus_domestica
MLLPSSLSLQTLAPPNADTIQKPHQTLPQNLTLFQLVHP------QHFPTLQFPN-----LPSGPKTL-
KFKLLCSQSPNPNPSP-------PSS---------------SSPVQTLISILRIIP--------
DWSDRTQERGMRQHRTLYDHEKWMHHRSSYRHLRHLLSSLSSRVILSLIPPVIAFTLVAVVIASYNTA
VAL-------------DLLPGIFPLLR---SSSLPYQLTAPALALLLVFRTEASYSRFEEGRKAWTEVIAGA--
NDFARQIISSVETSG------------------DAQLK-----KALLQYI----VAFPVALKCHVIYGSD-----IARDLQN-----
LLEVDDLLVV--LNSKHR-------------PGCIIQFISRSLQL----LKLEES--------------
RRIMLQSKISCFHEGIGICEQLIGTPIPLSYTRL-TSRFLVLWHLTL-------PIILWDDC-----
HWIVVPATFISAASLFCIEQVGVLIEEPFPM----LALDDLCNS-VRNNVQEALASE
>Chlamydomonas_eustigma_(GAX83184.1)
M---LVHRVHTRSLGNRNQCGRK---------------------
LHRVSTFVVKTPTEKPVVA----
DYVLPRSEEARRYFRTVYDFPQWQKHRSPTRLIDRLLQIPRSHVLQNILPSIAWCSSVAGLLTLYMQA
YDA-------------HILPDGFPSFATNNACTSFVNTTTVALSLLLVFRTNVSYGRWDEARKMKGLLVNRS--
RDLMRQVCAMVPEE-------------------DVATK-----AMMAKWT----AAFCRVLRIHFQPEVS-----LEDEMKG-
----LLSPEELEWL--IESKHR-------------PCSVIHMLSQIIYD----SQISAI--------------
CQAQMCNNLTAFEDVLGGCERLLRAPIPVSYTRH-TARFLFTWLTLL-------PFALYNSC-----
GVWTLPVVAGVSAVLCGIEEIGVQIEEPFGI----LPLEAICGR-IQADVMATLKED
>Cre16.g662600.t1.2_(BST1)
M--------------------------------QMQA-----------NRSSLRASPVRGLGARPLLRALPAGRVARLNVSAQA--
KDPNAPIQ---------------SNPLGTLSSQSGQVA----------
TLPRSEEARKYFRTVYDFPQWQKHRSSYRFAERLFQLSQSHILQNALPAISWVTLVATLVASYGYSY
DQ-------------HMLPDVFPSISPNASCTAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWT----VAFSRALRIHFQPEVT-----IESELQN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VPMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYPTT-----
GWGVVPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Cre16.g663400.t2.1_(BST2)
M--------------------------------QC-------------LSSRPVAMGRAGSSALPRL-
PLRAGRVCHLGVRCQAANKDPNAPIQ---------------SNPLGSFSSQLQNQP----------
TLPRSEEARKYFRTVYDFPQWQTHRNQYRLMKRLFSIPQSHVIQNALPSIMWVAFTSTCVAAYMYGY
DQ-------------HMLPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWV----VAFSRALRIHFQPEVT-----IESELKN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VPMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCSGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Cre16.g663450.t1.2_(BST3)
M--------------------------------QVSK-----------VPSS------ASARCLPRL-
PVRTSRVCQLSVRCQAANKDPNAPIQ---------------SNPLGSFSSQNSSGAVV--------
TAPRNEDARKYFRTVYDFPQWQKHRSQSRLVRRLFTIPQSHVIQNALPSIMWVTFTSTCVAAYMYGY

DL-------------HILPEGFPTLAPNAACSAFISNTSVALSLLLVFRTNSSYGRWDEARKMWGGLLNRS--
RDIMRQGATCFPDD-------------------QVEAK-----KALARWT----VAFARALRIHFQPEVT-----IESELQN-----
ILTPAELQML--AKSQHR-------------PVRAIHAISQIIQS----VRMSSI--------------
HQQQMSNNLTFFHDVLGGCERLLRAPIPVSYTRH-TARFLFAWLTLL-------PFALYGSC-----
GVSVIPVCTGIAAVLCGIEEIGVQCEEPFGI----LPLDVICNR-IQADVMATLKDD
>Chlorella_sorokiniana_(PRW33726.1)
M----------------------STAMLAGSRIQLQQPAG--------LGGSRLQ--RAAAPVAAAA-
RLGSVRPAGLQARSTAARRADRSALR--------------VSATASPEAAPVKLS----------
GDDLKEANRKHMRSVFDFDLWKKHRSSSRYLRHIVGLGESRIVSGLMAPLTYVMTLSLAVACYNAAA
EA-------------GYLP-VFPELKL--ATNAPFGLTSFALSLLLVFRTNSSYGRWDEARKMWGLIVNRS--
RDFIRQGLGYIPPE-------------------QEELQ-----KMLVRWT----VAYSRSLMCHLRPGED-----LRVELKD---
--TLKPEELEAL--LASTHR-------------PNYVVQVLTAIIKT----
AQLPAAVTNNRDSTGCVPAGAAYRMDENLTVFADTGGCERILRTPIPLSYTRH-TSRFMMIWLTLL---
-----PFTLWDSC-----HWAMLPIAGIVSFLLLGIEEIGVQIEEPFTI----LPLEVISRT-IEGNVWELYRMH
>Cre06.g261750.t1.2_(BST4)
M----------------------QC--------QLKH---------------------------GA-RPQSQRPNWLPARAATLRPAVQHGVR-
--------------RGLTLGVKAAAAPLE--------
DKKMPADMTTRQYRRVVYDFALWAKHRDVNRYLYNLRTIPGSRIIRQLSQPMGVVLAWAALFGFYET
CLEA-------------GVLPSYLPKMTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-------------------DQAGR-----EAVGKWT----VGFTRALQAHLQEDID-----LRKELEKA--
TPRWSKEEIDML--VNAQHR-------------PIKAISVLSELTRQ----LSITQF--------------
QALQMQENCTFFYDALGGCERLLRTPIPVSYTRH-TARFLTIWLAML-------PLGLWERY-----
HWSMLPVIALIGFLLLGIDEIGISIEEPFGI----LPLDAICGR-AQTDVNSLLKED
>XP_002945810.1_Volvox_carteri_f._nagariensis
M----------------------QS--------QLQP---------------------------RL-QLQGTRLNWLPQRSCVQRRSLRVDAT--
--------------SG------AAPPPPA--------
GKELSNDMVTRQYRRTVYDFSLWAKHRDVNRYLYNLKTIPGSRIIRTLGQPMGIVLAWAAMFGFYET
CLES-------------GVLPSYFPKLTL--MSAEPQGLTSFALSLLLVFRTNSSYGRFDEARKIWGGILNRA--
RNIANQAVTFIPAE-------------------DVAGR-----EAVGKWA----VGFCRALQAHLQEDAN-----LREELQKA-
-QPRWSREEIDML--CSAQHSWQQLQSCVNAFW-PIKAISMLSELTRQ----LPISQF--------------
QALQMQENVTFFYDALGGCERLLRTPIPVSYTRI--
LPLDAICTR-AQTDVVSLLKDD
>Chrysochromulina_sp._(KOO32217.1)
M----------------------REHPLSYEEYMRQRS----------AGRDPLA--EAVQGQSASM-
ERGVVPPPPVKATPERTEAAFTPPAEFDFFQDVFKPTVESVVKAVVSPGAQQDSDESYMRVPWWE
QGSTYSEDQRKDRRTVFMHDDWKRHRSSERFFRNIKTWPSSGINQALRKELTFVTSVSVFVVLANM
LLYQYQDFGGVVHPGPLSFLDGPIKSLS---
LPALPFSMASPVLSLLLVFRTNTAYFRWNEARTLWGGLINNC--RNIVRQTTTMFPNDA----------------
YHNALK-----KRLATET----ATFIKSLRNFLRGPEDDAT--LRKELYAYVNQGLMTSAQAEAT--LAAKNR---
----------PMFALAAMSATLRK----ANIDEM--------------YISRMDSTISVLVDLTGANERIFKSPIPLVYTRL-
TARFLSVFLTLL-------PLAMWAALGESWNHWATIPATFILSVFLFGVEEVGIQIEEPFSI----
LPLEAMCNGAIEAVQLEMLAAE
>Emiliania_huxleyi_(XP_005770556.1)

--
EFKEAGREFRQDVYSYNDWRWHRESGHIASAISSVFTSGVGKAMWRETFFVIATAAAVYLYNIGVPV
LAAKTAASL----PIVAALLGRLPLLHLSLLPLTLSSPALFLLLVFRTNNSYDRWWEARKVWGGVINAS--
RDLARQALALVR--------------------DAELK-----KLMVSQI----ASYARVLKYHLGPPTPEARDLLRNELVD--
--NRLPADQVRVI--MEAKHK-------------PMALLGLVSASLHDSGR-TGLDTV--------------
QASKLDQTLSLLTDYLGKCERIVKTPLPLVYTRH-TARFLSWWLLFL-------PVCLYNQLRA---
NWMIVPVSGLIGFFLVGIEDLGNQIEEPFSI----LPLTAMGTG-IQQSIFEAL---
>Phaeodactylum_tricornutum_(XP_002180738.1)
MM--RNFASVLLLL-SSGAAAFAPVQHNGVRTIATPSTPLY---
-------------GNTKQPPALP--PIK----------
DISYGEESRKYRRTVYSHDDWVKHRSSDRFLRNLLAIGSSGVYKSLAKEVLATTGVATFIVLYNCLVG
GYTDLEGIKHS---ALIESVWAPLMA--LPLAPFTLSSPSLGLLLVFRTNTSYQRWDEARKNWGMNINHT-
-RDLVRMGTSFYDNAA------------------VSSEQRAKDL-KALSLAT----WSFVRAMKRHLSPESEDEQD-
FRRELFE-----RLPAPQAQAI--IDAAHR-------------PNRALFDLSVAIEN----LPMHFL--------------
RKNQVHQAVTIFEDNLGSSERLLTSPVPLFYSRH-TARFLSFWLLLL-------
PFALWDPFAGTWNHVGMIPATAVISIFLFGIEELATQMEEPFTI----LPMQAFCDK-IGNWCNEIVSWQ

>Thalassiosira_pseudonana_(XP_002289965.1)
M--
GPPIDPSVPVT-----------
DQVGEGSRKYRRTVYTHDDWVRHRSPDRFGNNLSTLFNSGIYKQVANEVFATTAVATFVFLWNMIA
GGYTDLAGVQHG---PIIDSPLAQMVG--
LPMTAFTILTPSLGLLLVFRTNTSYGRWDEARKMWGLNINHT--
RDLNRMATAWYGNEGNMDSVAFMGGDIPYSQPIDPVQRAYDL-GQVSLFT----
WAFVRSMKRHLSPPEEDEED-FKAELRA-----RLTPEQAENI--INAAHR-------------PNRALFDLSVAIEN--
--LPMHFL--------------RKNAINTNLSIFEDTLGGCERLLSSPVPLFYSRH-TARFLSTWLLLL-------
PFGLYEQFKDSWNHIAMIPATAFISVCLFGIEELATQLEEPFTI----LPMQGFCDK-IGGWCDEIVSW-
>WP_049046555.1_Kiebsiella_aerogenes

---------------------MIIRPEQHWFFRLFDWHGSVLSKIVFRLLLNVLMSVIAIISYQWYEQL-------------GI------
-----HLTVAPFSLLGIAIAIFLGFRNSASYSRFVEARNLWGTVLIAE--RTLVRQLKNILPD--------------------
DEETH-----KTLVSYL----VAFSWSLKHQLRK-TD-----PAVDLYR-----LLPKEKVAEI--LASSMP-------------
TNRILLLIGNELGRLREQGKLSDI--------------TYGLMDNKLDELAHVLGGCERLASTPVPFAYTLI-
LQRTVYLFCTLL-------PFALVGDL-----
HYMTPFVSVFISYTFLSWDSLAEELEDPFGTSANDLPLNAMCNT-IERNLMDMTGQH
>XP_011543531.1_Homo_sapiens
M---
----------------TITYTSQVANARLGSFSRLLLCWRGSIYKLLYGEFLIFLLCYYIIRFIYRLALTEEQQL----------
----MFEKLTL--YCDSYIQL--IPISFVLGFYVTLVVTRW------WNQYENLPWPDRLMSLVSGFVEGK-------
-----------DEQGRL--LRRTLIRYANLGNVLILRSVSTAVYKRFP-----SAQHLVQ---
AGFMTPAEHKQLEKLSLPHNM---------FWVPWVVFANLSMKAWLG---GRIRDPI-------------
LLQSLLNEMNTLRTQCGHLYAYDWISIPLVYTQVVTVAVYSFFLTCLVGRQFLNPAKAYPGHEL---
DLVVPVFTFLQFFFYVGWLKVAEQLINPFGEDDDDFETNWIVDRNLQVSLLAVDEMH

Supplemental Figure 12. Sequences used to generate MAFFT alignments used to generate the phylogenetic trees included in this manuscript. FASTA files of MAFFT alignments of **A.** Full length and **B.** trimmed after the bestrophin domain BST4 and homologous amino acid sequences.