Surface temperatures are influenced by handling stress independently of

glucocorticoid levels in wild king penguins

Agnès Lewden[1,2†], Chelsea Ward[3], Aude Noiret[4], Sandra Avril[5,6], Lucie Abolivier[5,6], Caroline Gérard[5,6],

Tracey L. Hammer[5], Émilie Raymond[4,5], Jean-Patrice Robin[5], Vincent A. Viblanc[5*], Pierre Bize[3,7*]and

Antoine Stier[4,5,8*†]

[1] Faculty of Biological Sciences, University of Leeds, Leeds, UK

[2] Université de Brest - UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des sciences de

l'environnement marin - IUEM - Rue Dumont D'Urville - 29280 - Plouzané

[3] School of Biological Sciences, University of Aberdeen, Aberdeen, UK

[4] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne,

[5] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[6] Institut Polaire Français Paul-Émile Victor, Plouzané, France

[7] Swiss Ornithological Institute, CH-6204 Sempach, Switzerland

[8] Department of Biology, University of Turku, Turku, Finland

[†]: corresponding authors: agnes.lewden@univ-brest.fr & antoine.stier@iphc.cnrs.fr

[*]: share co-senior authorship

Highlights:

• Experimental increase in CORT does not affect king penguin's surface temperature

• Acute handling stress increases eye but decreases beak surface temperature in adults

• Acute handling decreases both eye and beak surface temperatures in young chicks

• Parental CORT treatment does not affect egg surface temperature during incubation

• Chicks brooded by non-implanted partners of CORT parents are warmer

Abstract

Assessing the physiological stress responses of wild animals opens a window for understanding how organisms cope with environmental challenges. Since stress response is associated with changes in body temperature, the use of body surface temperature through thermal imaging could help to measure acute and chronic stress responses non-invasively. We used thermal imaging, acute handling-stress protocol and an experimental manipulation of glucocorticoid hormone levels in breeding king penguins (*Aptenodytes patagonicus*), to assess: 1. the potential contribution of the Hypothalamo-Pituitary-Adrenal (HPA) axis in mediating chronic and acute stress-induced changes in adult surface temperature, 2. the influence of HPA axis manipulation on parental investment through thermal imaging of eggs and brooded chicks, and 3. the impact of parental treatment on offspring thermal's response to acute handling.

Eye temperature (T_{eye}) increased and beak temperature (T_{beak}) decreased in response to handling stress in adults, but neither basal nor stress-induced surface temperatures were significantly affected by glucocorticoid implants. While egg temperature was not significantly influenced by parental treatment, we found a surprising pattern for chicks: chicks brooded by the (non-implanted) partner of glucocorticoid-implanted individuals exhibited higher surface temperature (both T_{eye} and T_{beak}) than those brooded by glucocorticoid-implanted or control parents. Chick's response to handling in terms of surface temperature was characterized by a drop in both T_{eye} and T_{beak} independently of parental treatment.

We conclude that the HPA seems unlikely to play a major role in determining chronic or acute changes in surface temperature in king penguins. Changes in surface temperature may primarily be mediated by the Sympathetic-Adrenal-Medullary axis (SAM) in response to stressful situations. Our experiment did not reveal a direct impact of parental HPA manipulation on parental investment (egg or chick temperature), but a potential influence on the partner's brooding behaviour (*i.e. family-transmitted stress* hypothesis).

Key words: Thermal imaging, Stress, Corticosterone, Heterothermy, Bird, Beak, Eye

Abbreviations:

Ambient air temperature: T_a

Glucocorticoid hormones (corticosterone or cortisol): CORT

Egg surface temperature: T_{egg}

Hypothalamic-pituitary-adrenal axis: HPA

Maximum eye region surface temperature: T_{eye}

Minimum beak surface temperature: T_{beak}

Sympathetic-adrenal-medullary axis: SAM

1. Introduction

Measuring physiological stress is of central interest in wild animals to understand how they cope with environmental change (Ellenberg et al., 2007; Romero, 2004). Traditionally, ecologists have evaluated individual stress by assessing the activation of the Sympathetic-Adrenal-Medullary (SAM) or Hypothalamic-Pituitary-Adrenal (HPA) axes, either at baseline levels or in response to acute disturbances in the environment, by metrics such as increased heart rate (Cabanac and Guillemette, 2001; Viblanc et al., 2012), ventilation rate (Carere and Oers, 2004) or glucocorticoid hormone (GC) secretion (Sapolsky et al., 2000). Stress exposure is also known to trigger core body and surface temperature changes (Oka, 2018). As the latter can be measured using thermal imaging (McCafferty, 2013), it led to a recent burgeoning of thermal imaging studies to measure the stress response of captive and wild endotherms (McCafferty et al., 2021). Both the SAM and HPA can potentially influence changes in body surface temperature, but the importance of these two pathways in mediating the stress response(s) measured through thermal imaging remains largely unknown. This information is however essential for the proper interpretation of the physiological stress response measured by thermal imaging.

SAM responds within seconds to a stressor by releasing catecholamines hormones (adrenaline and noradrenaline) into the blood stream, which induces an immediate increase in heart rate (tachycardia) and vasoconstriction of peripheral blood vessels (Wingfield and Romero, 2015). This redistribution of peripheral blood flow to essential internal organs generally leads to a response called stress-induced hyperthermia along with a decrease of surface temperature (Cabanac and Guillemette, 2001; Herborn et al., 2015; Oka, 2018). The SAM response is followed by the slower (within a few minutes) activation of HPA that releases

GC into the blood stream (Wingfield and Romero, 2015). GC are metabolic hormones that

helps responding to a stressor by mobilizing energy resources and by triggering an increase in

behavioural activity and metabolic rate (Wingfield and Romero, 2015). Circulating GC levels

are usually considered at baseline (*i.e.* without an inducing stressor) or at stress-induced

levels, which are often interpreted as mirroring chronic stress/metabolic demand and acute

stress response respectively. Yet, the actual interpretation of circulating GC levels is likely

more complex (MacDougall-Shackleton et al., 2019; Lemmonier et al. 2022). GC-mediated

changes in metabolism may cause an increase in internal heat production that, in turn, may

lead to changes in body surface temperature (Oka, 2018). Additionally, previous evidence

suggest that glucocorticoids might be necessary for catecholamine-induced thermogenic

effects such as shivering, free fatty acid mobilization or vasoconstriction (Deavers and

Musacchia, 1979).

Therefore, both the SAM and HPA axes could trigger, independently from each other,

or together, a change in body core and/or surface temperatures (Ouyang et al., 2021). The

immediate activation of SAM could lead to a decrease of peripheral temperature

independently of CORT release. On the other hand, the short to long-term elevation of

baseline CORT might lead to increased metabolic rate, heat production and, ultimately, higher

peripheral temperature (to facilitate heat dissipation and maintain homeothermy)

independently of SAM. To date, the use of thermal imaging as a non-invasive tool to measure

stress in unmanipulated animals, as well as the roles of SAM and HPA axes in mediating

changes in surface temperature, remains debated. For instance, data in captive and wild birds

show that acute stress exposure is often leading to a decrease in body surface temperature

(reviewed in Table 1, but note some discrepancies between species and/or body parts), as

predicted if these changes are driven by the SAM axis.

Table 1: **Summary table of avian studies testing the impact of acute stress exposure on**
changes in body surface temperatures measured with thermal imaging (studies related to
heat and food stress are not included). *Area* corresponds to the body region being measured,
response to the direction of the temperature response (=: no significant change, ↘: significant
decrease and ↗: significant increase, *Δ T°C* to the amplitude of the temperature response, *stressor* the
nature and duration of the experimental stressor. Some studies measured temperature responses at
different time points, and when the effects of stress exposure change through time, the responses are
shown with consecutive symbols (*e.g.* =, ↗: initial absence of change followed later on by an increase
in surface temperature).

Species	Condition	State	Stressor	Area	Response	Δ T°C	Ref.
Blue tit *Cyaniste caerulescens*	Wild	Adult	Trap closure + handling (3min)	T_{eye}	↘, ↗,=	-1.0, +1.0°C, =	Jerem et al. 2019
Budgerigars *Melopsittacus undulatus*	Captive	Adult	Handling (30 min)	T_{eye}	↗, =	+0.7°C, =	Ikkatai and Watanabe 2015
				T_{legs}	=	=	
Chicken *Gallus gallus domesticus*	Captive	Adult	Air puff (10 min)	T_{eye}	↘	-1.0°C	Edgar et al. 2011
Chicken *Gallus gallus domesticus*	Captive	Adult	Handling (20 min)	T_{eye}	↘, =	-0.5°C, =	Edgar et al. 2013
				T_{head}	=, ↗	=, +1.0°C	
				T_{comb}	↘, =	-1.5°C, =	
Chicken *Gallus gallus domesticus*	Captive	Chick (9 weeks old)	Air puff (10 min)	T_{eye}	↘	-1.0°C	Edgar et al. 2018
Chicken *Gallus gallus domesticus*	Captive	Adult	Handling: mild/severe (20 min)	T_{eye}	↘	-0.4°C	Herborn et al. 2015
				T_{face}	=	=	
				T_{wattle}	↘	-0.7 / -1.3°C	
				T_{comb}	↘	-0.5 / -2.2°C	
Chicken *Gallus gallus domesticus*	Captive	Adult	Enrichment removal (within 2h)	T_{eye}	↘	-0.8°C	Herborn et al. 2018
				T_{face}	↘	-0.7°C	
				T_{comb}	↘	-3.5°C	
Chicken *Gallus gallus domesticus*	Captive	Chick (30 days old)	Handling (10 min)	T_{head}	↗	+0.76°C	Moe et al. 2017
				T_{feet}	↘	-0.45°C	
Chicken *Gallus gallus domesticus*	Captive	Chick (9 days old)	Visual (45 min)	T_{eye}	=	=	Pijpers et al. 2022
				T_{beak}	=, ↗	+0.5°C	
Domestic pigeon *Columbia livia domestica*	Captive	Adult	Handling (3.5 min)	T_{eye}	=	-0.4°C	Tabh et al. 2021
				T_{beak}	↘	-2.1°C	
King penguin *Aptenodytes patagonicus*	Wild	Adult	Handling (5-10 min)	T_{eye}	↗	+0.8°C	This study
				T_{beak}	↘	-1.5°C	
	Wild	Chick (20 days old*)	Handling (5-10 min)	T_{eye}	↘	-0.9°C	
				T_{beak}	↘	-3.3°C	
Little Auk *Alle alle*	Wild	Adult	Handling (30 min)	T_{eye}	↗	+1,7°C	Jakubas et al. 2022
Pheasant *Phasianus colchicus*	Captive	Juvenile (42 to 49 days old)	Agression (20 s)	T_{head}	↗	+0.6°C	Knoch et al. 2022
Song sparrow *Melospiza melodia*	Captive	Adult	Handling (13 min)	T_{eye}	↘, =	-1.5°C, =	Zuluaga et al. 2023
				T_{beak}	↘, ↗,=	-2.6°C, +2.6°C, =	
* king penguins chicks are not thermally emancipated at this age							

Evidence for an impact of chronic stressors on body surface temperature (Herborn et

al., 2018; Robertson et al., 2020a, 2020b; Winder et al., 2020) or for a correlation between

circulating GC levels and body surface temperature (Giloh et al., 2012; Jerem et al., 2018;

Weimer et al., 2020), however, remains mixed. Therefore, the question of whether acute or

chronic elevation of GC causally induces changes in surface temperature remains largely

untested (but see Ouyang et al., 2021). Since both HPA and SAM axes provide strong and very

reactive responses to stress, teasing apart their contribution to changes in peripheral

temperature is challenging.

In this study, we took up the challenge of distinguishing the role of HPA and SAM on

changes in body surface temperatures using an experimental approach in adult king penguins

(*Aptenodytes patagonicus*). The king penguin is a heterothermic bird whose body temperature

largely varies between core and peripheral tissues (Lewden et al., 2017a, 2017b) . Its thick

body plumage provides a very efficient insulation, while counter-current vascular heat

exchangers in the appendages allow to decrease peripheral temperature loss (Thomas et al.,

2011). These adaptations allow birds to conserve heat when foraging in cold Antarctic waters

(Handrich et al., 1997), and to rapidly recover to normothermia when exiting the water

(Lewden et al., 2020). Thermal imaging has been frequently used in the last years to

investigate acute (see an overview in Table 1) and chronic (Herborn et al., 2018; Robertson et

al., 2020a, 2020b; Tabh et al., 2022; Winder et al., 2020) stress-response in birds. Because of

their dense and watertight plumage, the use of thermal imaging in penguins is restricted to

measuring bare parts of the body, such as the eye (periorbital region) and beak (Lewden et

al., 2020). These regions are highly vascularised and provide valuable information on body

surface temperatures.

Breeding king penguins were subcutaneously implanted with either a corticosterone

(CORT) implant or a placebo implant. The use of CORT implants in king penguin (see methods

and Fig. 1), as in many other bird species (see Torres-Medina et al., 2018), leads to higher

baseline GC (or CORT) levels, while inhibiting the acute CORT responses in the days following

implantation. Indeed, CORT-implanted birds no longer exhibit high blood CORT release in

response to acute stress induced by a handling stress protocol. Hence, the use of CORT

implants in conjunction with handling stress allows for the separation of the contributions of

the HPA and SAM axes to acute changes in body surface temperatures. Here, we therefore

measured the effects of our CORT treatment, together with a handling stress protocol, on

surface temperatures in the king penguin. If changes in surface temperatures are mainly

driven by activating the HPA axis, we expect to find higher initial body surface temperatures

in CORT-implanted individuals (with elevated baseline CORT) prior to capture and handling.

However, because CORT implants shut-down the release of CORT in response to handling, we

expect to find no change in body surface temperature in response to handling stress in CORT

individuals if such changes are mediated by the HPA only. On the contrary, if changes in

surface temperatures are mainly driven by the activity of the SAM axis, we expect our CORT

treatment to have no additional effect on body surface temperatures (both before or after

capture and handling). We also measured egg surface temperature on the same day as the

body surface temperatures of implanted parents and the surface temperatures of chicks at

day 20 after hatching, before they were fully thermally emancipated. Based on experimental

data in Adélie penguin showing that elevated baseline CORT led to lower parental care (Thierry

et al., 2013a), we hypothesized that increasing CORT levels may lower the surface

temperatures of eggs and/or chicks during incubation and brooding respectively, as parents

may pay less attention in keeping the egg/chick covered and warm under their brood pouch.

2. Material and Methods

2.1. Study site and species

The study was conducted in 2018-2019 in a king penguin colony of ca 22,000 breeding birds at 'La Baie du Marin' on Possession Island (Barbraud et al. 2020), Crozet Archipelago, in the Southern Indian Ocean (46°26' S, 51°52' E). We followed 49 breeding pairs from courtship (early November) until the onset of the Austral winter (April). All adults were identified by a hair dye mark on the breast feathers. During the breeding season, male and female alternate between periods on-land caring for their single-egg or chick and periods foraging at sea (Weimerskirch et al., 1992). The male takes care of the first incubation shift while the female forages at sea. The female returns ~15 days later to relieve her partner, and the parents continue to alternate shifts throughout the incubation (~53 days) and early chick brooding. Chicks become thermally emancipated from the parents around one month of age, allowing both parents to go back at sea simultaneously to forage (Weimerskirch et al., 1992).

2.2. Use of sub-cutaneous CORT and SHAM implants

We randomly divided the 49 breeding pairs into two groups of either female-treated pairs (N = 23) or male-treated pairs (N = 26). In the female-treated pairs, we randomly implanted females with either a CORT (NG-111, 50mg corticosterone, 90-days release; N = 12 females) or a Placebo (NC-111, same vehicle but no CORT; N = 11 females) implant. Sub-cutaneous implants were purchased from Innovative Research of America (Sarasota, FL, USA). Females were implanted during their first incubation shift, 3 days after returning from sea. Similarly, in the male-treated pairs, we randomly implanted 13 males with a CORT and 13 males with a Placebo implant, 3 days after returning from sea. In males, the first incubation

shift is immediately after egg laying, and males have already been fasting for ~15 days during

courtship. Therefore, we chose to implant the males during their second incubation shift so

that they would have comparable fasting times to the females, and therefore both sexes

would be in a comparable physiological state during measurements. Male and female partners

in, respectively, the female-treated and male-treated pairs were not implanted. Hereafter,

they are referred to as "Partner-CORT" or "Partner-Placebo" regardless of their sex. Partner

penguins were only considered in the chick analysis (see below).

On the day of the implantation, adults were captured while incubating their egg and

restrained by one experimenter with a hood covering the head to keep them calm. Sub-

cutaneous CORT or placebo implants were inserted in the upper part of the back by a

veterinary surgeon (SA) under local anaesthesia (*ca.* 0.5mg/kg xylocaine and 0.0001mg/kg

adrenalin, Aspen Pharma). Immediately after implantation, the small incision was closed using

3 sterile surgical staples, and a prophylactic dose of antibiotics (cephalexin *ca.* 50mg/kg,

Rilexine®, Virbac) was injected to prevent any risk of infection. The wound was checked 3 days

later, when taking the thermal images without any sign of infection on individuals, nor implant

rejection, and staples were removed 6 days after implantation.

To verify the efficacy of our CORT treatment, blood samples were taken 6 days after

implantation in less than 4 min (baseline CORT) and after 30 min of standardized handling

(stress-induced, see Stier et al., 2019; Viblanc et al., 2018). Plasma total CORT levels were

determined by immunoassay according to guidelines provided by the manufacturer

(Corticosterone EIA Kit, Arbor Assay, USA). Intra-plate coefficient of variation based on

duplicates was 8.17 ± 0.90%, and inter-plate coefficient of variation based on one repeated

sample was 8.08%. Implants were successful in raising circulating baseline CORT levels and in

inhibiting stress-induced CORT release (Fig. 1).

Fig. 1: **Baseline and stress-induced plasma corticosterone levels of breeding king penguins** (N = 20
individuals per group) implanted either with a Placebo, or a CORT subcutaneous implant (50 mg, 90
days release) during incubation. Birds were blood sampled 6 days after implantation. CORT levels were
influenced by the interaction between handling stress (baseline: < 4 min vs. stress-induced: 30 min of
handling stress) and CORT treatment (LMM on log-transformed plasma CORT: $F_{1,37.2}$ = 62.4, $p < 0.001$).
Letters indicate significant differences according to post-hoc tests with Tukey adjustments ($p < 0.030$).

2.3. Thermal image collection and analysis

Thermal pictures of implanted adults (Fig. 2A) and their egg (Fig. 2B) were taken 3 days

after implantation and thermal pictures of chicks (Fig. 2C) were taken at day 20 after hatching

(40 to 60 days after CORT or Placebo implant of the focal parent, *i.e.* within the 90-days release

given by the CORT pellet manufacturer). Thermal pictures were taken using FLIR E8 thermal

imaging camera, Lewden et al., 2020). When approaching the adults and/or chicks, we started

a stopwatch when the focal individual showed the first signs of alarm behaviour to measure

the duration of our disturbance at the time the different pictures were taken. In adults, we

took a first set of pictures (1 to 2 pictures per individual) before handling, on average (mean ±

SE) 38 ± 4 sec after the first sign of alarm behaviour at a distance of *ca*. 2 meters from the

focal individuals. Adults were then captured-restrained, wound healing was checked, and we

took a second set of pictures (1 to 2 pictures per individual) just after terminating handling

stress (mean ± SE handling stress duration was 8.3 ± 0.3 min; a timeframe within which CORT

is known to increase by *ca.* 100% above baseline levels in king penguins (Viblanc et al., 2018).

Just after capturing the adult, we removed the single-egg from the brood pouch and

immediately took a thermal picture at a distance of *ca.* 0.5 m. We then measured egg length

and breadth using callipers and estimated egg volume (cm^3) following Narushin (2005). When

chicks were 20 days old, we took a first set of pictures (1 to 2 pictures per individual) at a

distance of *ca.* 0.5 m immediately after they were removed from the brood pouch (*i.e.* before

handling: 44 ± 5 sec after the first sign of alarm behavior from the parent) and a second set of

pictures (1-2 pictures per individual) was taken after handling, 5.4 ± 0.3 min later. Thermal

images of the chicks were only taken in non-rainy days, which reduced the sample size from

38 chicks that survived until day 20, to 16 chicks with thermal images available. Chick sex was

unknown and body mass was not significantly influenced by the treatment (CORT: 1.14 ± 0.11

vs. Placebo 1.32 ± 0.09, N = 8/group, t = -1.28, p = 0.22). Chicks were kept at ambient

temperature (14.9 ± 0.9°C) between the first and second set of pictures, a temperature at

which they are known to be able to maintain their internal body temperature constant

(Duchamp et al., 2002). In adults and chicks, we defined the first set of measures taken < 1

minute as 'before handling' surface temperatures and the second set of measures taken after

> 5 minutes of handling stress as 'after handling' surface temperatures.

274

Fig. 2: **Infrared images of (A) breeding adult, (B) incubated egg and (C) 20 days-old chick**

Thermal images were analysed using the ThermaCAM TM Researcher Pro 2.10

software (Flir systems, Wilsonville, Oregon, USA). Only profile pictures were analysed to avoid

surface temperature errors related to head's orientation (Playà-Montmany and Tattersall,

2021; Tabh et al., 2021). For each image we set the emissivity at 0.98 and controlled in the

analyses for daily variation in air temperature (T_a) and relative humidity using daily climatic

measures from a permanent weather station located 2 km inland from the colony

(https://rp5.ru/Archives_m%C3%A9t%C3%A9o_sur_la_base_Alfred-Faure). We extracted

measures of surface temperatures on maximum eye temperature (T_{eye}) and minimum beak

temperature (T_{beak}) in adults and chicks. We used maximum T_{eye} as recommended by (Jerem

et al., 2015, 2019) and the minimum T_{beak} to gain insight on the maximum state of

vasoconstriction for this body surface. Finally, we estimated egg surface temperature (T_{egg}) as

the mean of average length and width surface temperature.

2.4. Ethical note

All the procedures were approved by the French Ethical Committee (APAFIS#16465-

2018080111195526 v4) and the Terres Australes et Antarctiques Françaises (Arrêté TAAF A-

2018-118).

2.5. Statistical analysis

We ran separate analyses for adults, eggs and chicks. In adults, we investigated variations in T_{eye} and T_{beak} using two separated linear mixed models (LMMs) where treatment (CORT *vs.* Placebo), handling stress (before *vs.* after), Sex and T_a were specified as fixed effects. Relative humidity was initially tested but removed from final models since it was never significant. Two-ways interactions were also initially included but removed from final model if $p > 0.10$ using a backward stepwise procedure. The p-values (just before removal in the backward stepwise procedure) for the focal interaction between treatment and handling stress are reported in tables. Bird identity was included as a random intercept to control for repeated measures. We investigated variation in T_{egg} by entering parental CORT treatment (CORT *vs.* Placebo), the sex of the parent, T_a and egg volume as fixed effects. We investigated variation chick T_{eye} and T_{beak} by entering parental treatment (CORT *vs.* Placebo), the category of the brooding parent at the time of measurement (implanted individual *vs.* non-implanted partner), handling stress (before *vs.* after), and T_a as fixed effects, and chick identity as a random intercept. Interactions were treated as described above for adults. All statistical analyses were conducted using R Studio (version 3.6.2) and *lme4*, *emmeans* and *pbkrtest* packages. Results are reported as least-square means ± SE.

3. Results

3.1. Adult surface temperatures

Fig. 3: **Adult king penguin surface temperature responses to an experimental manipulation of glucocorticoid levels (CORT *vs.* Placebo subcutaneous implants) and handling stress: (A) Maximum eye temperature response to handling stress, (B) Minimum beak temperature response to handling stress, and (C) Sex-specific response of eye temperature to handling stress.** In panels A and B, CORT (blue) *vs.* Placebo (green) individual responses (raw data) are presented. Since CORT treatment had no significant effect either alone or in interaction with handling stress (see Table 2 for details), the overall mean ± SE is shown in black. Least-square means ± SE from final statistical models (Table 2) are presented. Different letters indicate significant differences in panels A and B, and ** in panel C represent the significant increase of T_{eye} observed only in females.

Treatment (CORT *vs.* Placebo), either alone or in interaction with handling stress, had

no significant effect on eye (T_{eye}) or beak (T_{beak}) surface temperature in incubating adults

(Table 2, Fig. 3A and 3B). Handling stress significantly affected T_{eye} and T_{beak}, in opposite

directions (Table 2): over an 8-min handling stress, T_{eye} increased by 0.8 ± 0.3°C, on average

(Fig. 3A), while T_{beak} decreased by -1.5 ± 0.4°C, on average (Fig. 3B). The increase in T_{eye}

appeared to be sex-specific (marginally significant interaction Handling * Sex: $p = 0.060$), with

only females showing a significant increase in T_{eye} in response to handling stress (Fig. 3C;

males: $t = 0.63$, $p = 0.53$; females: $t = 3.08$, $p = 0.003$). As expected, T_{eye} and T_{beak} were

significantly and positively associated with T_a (Table 2).

Table 2: Summary of linear mixed models (LMMs) investigating the determinants of (A) maximum

eye temperature (T_{eye}) and (B) minimum beak temperature (T_{beak}) in adult king penguins.

A. T_{eye} (n = 138 observations; N = 49 individuals)

Random effects:		Variance			
Bird ID	Intercept	5.04			
Residual		3.21			
Fixed effects:	**Estimate**	**Std. Error**	**t**	**df**	**p**
Intercept	24.94	1.71	14.54	47.9	<0.001
Treatment (CORT)	-0.10	0.72	-0.15	45.8	0.89
Handling (after)	**1.45**	**0.47**	**3.04**	**89.3**	**0.003**
Sex (Male)	0.76	0.78	0.97	63.7	0.34
T_a	**0.29**	**0.13**	**2.33**	**45.0**	**0.024**
*Handling*Sex*	*-1.19*	*0.63*	*-1.90*	*89.7*	*0.060*
(Treatment x Handling)					(0.44)

B. T_{beak} (n = 138 observations; N = 49 individuals)

Random effects:		Variance			
Bird ID	Intercept	15.91			
Residual		6.40			
Fixed effects:	**Estimate**	**Std. Error**	**t**	**df**	**p**
Intercept	13.84	2.92	4.74	45.0	<0.001
Treatment (CORT)	-1.13	1.24	-0.91	44.5	0.37
Handling (after)	**-1.47**	**0.45**	**-3.29**	**84.6**	**0.001**
Sex (Male)	0.90	1.23	0.73	44.6	0.47
T_a	**0.54**	**0.22**	**2.45**	**43.8**	**0.018**
(Treatment x Handling)					(0.61)

 3.2. Egg surface temperature

 Neither parental treatment ($F_{1,45}$ = 1.38, p = 0.25; CORT: 34.5 ± 0.2 *vs.* Placebo: 34.9 ±

 0.3°C; N = 46), sex ($F_{1,45}$ = 0.47, p = 0.49), ambient temperature ($F_{1,45}$ = 0.45, p = 0.50), nor egg

 volume ($F_{1,45}$ = 0.01, p = 0.93) significantly affected T_{egg}.

 3.3. Chick surface temperatures

 Fig. 4: **King penguin chick surface temperature in response to parental glucocorticoid manipulation**
 and handling: (A) Maximum eye temperature according to parental treatment and category of the
 parent brooding the chick (*i.e.* partners are not implanted), (B) Minimum beak temperature
 according to parental treatment and category of the parent brooding the chick, (C) Maximum eye
 temperature response to handling, and (D) Minimum eye temperature response to handling. In
 panels C and D, CORT (blue) vs. Placebo (green) chick's individual responses (raw data) are presented,
 but only the global mean ± SE are presented since there was no significant interaction between
 parental CORT treatment and handling (see Table 3 for details). Least-square means ± SE from final
 statistical models (Table 3) are presented. Different letters indicate significant differences.

The interaction between parental treatment (CORT *vs.* Placebo) and the category of

the parent brooding the chick (implanted *vs.* non-implanted partner) at the time of

measurement significantly affected chick T_{eye} ($F_{1,11}$ = 6.37, p = 0.028, Table 3) and chick T_{beak}

($F_{1,11}$ = 5.49, p = 0.039, Table 3). Chicks brooded by the partner of CORT-implanted individuals

had higher overall T_{eye} and T_{beak} than chicks brooded by CORT parents, or by Placebo and

associated partners (Fig 4A and 4B). While we found no significant interaction between

parental treatment and Handling (Table 3), both chick T_{eye} (Fig 4C) and chick T_{beak} (Fig 4D)

significantly decreased after handling (by -0.9 ± 0.3°C and -3.3 ± 0.6°C respectively, Table 3).

As expected, chick T_{eye} and T_{beak} were significantly positively associated with T_a (Table 3).

Table 3: Summary of linear mixed models (LMMs) investigating the determinants of (A) maximum
eye temperature (T_{eye}) and (B) minimum beak temperature (T_{beak}) in king penguin chicks. Only one
parent was implanted with CORT or Placebo implant, thus we considered both parental treatment
(CORT *vs.* Placebo), the category of the parent brooding the chick at the time of measurement
(implanted parent *vs.* non-implanted partner) and their interaction in our analyses.

A. T_{eye} (n = 63 observations; N = 16 chicks)

Random effects:		Variance			
Bird ID	Intercept	0.81			
Residual		1.18			
Fixed effects:	**Estimate**	**Std. Error**	**t**	**df**	**p**
Intercept	29.47	1.39	21.26	11.1	<0.001
Treatment (CORT)	-0.88	0.68	-1.29	11.3	0.22
Parent (implanted)	**-2.84**	**0.82**	**3.48**	**11.1**	**0.005**
Handling (after)	**-0.85**	**0.28**	**-3.08**	**47.2**	**0.003**
T_a	**0.21**	**0.09**	**2.41**	**10.9**	**0.035**
Treatment*Parent	**2.78**	**1.10**	**2.53**	**11.0**	**0.028**
(Treatment x Handling)					(0.79)

B. T_{beak} (n = 63 observations; N = 16 chicks)

Random effects:		Variance			
Bird ID	Intercept	5.53			
Residual		4.89			
Fixed effects:	**Estimate**	**Std. Error**	**t**	**df**	**p**
Intercept	13.55	3.42	3.96	11.1	0.002
Treatment (CORT)	1.15	1.68	0.69	11.3	0.51
Parent (implanted)	**-6.82**	**2.02**	**-3.38**	**11.1**	**0.006**
Handling (after)	**-3.26**	**0.56**	**-5.77**	**46.9**	**<0.001**
T_a	**0.57**	**0.21**	**2.65**	**11.0**	**0.023**
Treatment*Parent	**6.37**	**2.72**	**2.34**	**11.0**	**0.039**
(Treatment x Handling)					(0.94)

4. Discussion

By experimentally manipulating CORT levels in adult breeding king penguins, we show that glucocorticoids are unlikely to play a major role in determining changes in surface temperatures (both before or in response to handling stress) in this large seabird species. Indeed, we found no significant difference in baseline surface temperature between CORT and Placebo implanted individuals. However, whatever the parental CORT treatment, our handling stress protocol led to an increase in T_{eye} and a decrease in T_{beak} in response to capture and handling. The contrasting results between parental treatment and handling stress suggest that changes in surface temperature are probably driven primarily by the activity of the SAM axis in king penguins (but see Ouyang et al. (2021), for a potential causal involvement of CORT in house sparrows *Passer domesticus*). Contrary to our prediction, parental treatment had no significant effect on egg temperature. Yet, we found an unexpected pattern on chick surface temperatures: chicks brooded by the non-implanted partner of CORT individuals had higher surface temperatures (both T_{eye} and T_{beak}) than chicks brooded by Placebo (or associated non-implanted partner) or CORT individuals. Finally, in the thermally non-emancipated chicks, both T_{eye} and T_{beak} decreased with handling, but irrespectively of the parental treatment.

4.1. CORT and adult surface temperatures

Previous correlative studies have considered baseline and stress-induced CORT levels to investigate the relationship between GC signalling and changes in body surface temperature in birds (blue tit, *Cyanistes caerulescens*, Jerem et al. (2018); domestic chicken, *Gallus gallus domesticus,* Giloh et al. (2012); Weimer et al. (2020)). To go deeper in the involved mechanisms, we experimentally manipulated CORT in king penguin to help to causally distinguish the contribution of the HPA and SAM axes to changes in body surface temperature. Since we did not detect any significant effect of CORT manipulation on baseline or stress-induced surface temperatures, our results provide little support for a causal involvement of the HPA in mediating stress-related changes in surface temperature.

Our results contrast with a recent study showing that an acute stimulation of the HPA axis with ACTH induced a decrease in T_{eye} within 5 min in captive-held house sparrows (Ouyang et al., 2021). While we used an experimental approach and the lack of significant effect of baseline CORT on body surface temperatures appears to be robust, we cannot fully exclude

that our handling stress protocol for thermal imaging was too short to detect a potential role

of stress-induced CORT in influencing changes in surface temperatures, since in 8 minutes of

handling, CORT levels are expected to double in king penguin, while the maximum CORT levels

are only attained after *ca.* 80 minutes of handling in our study species (Viblanc et al., 2018).

Overall, our results suggest that changes in surface temperature are probably more likely

mediated by the SAM rather than the HPA in adult king penguins. Future studies testing for a

direct acute stimulation of the HPA axis (using ACTH) or SAM axis (using catecholamines) in

the king penguin and other bird species could be useful to gain greater insight on the

contribution of the HPA and SAM axes in mediating changes in body surface temperature

(Ouyang et al., 2021).

4.2. Stress-induced changes in surface temperatures

In breeding adults, we observed a mild (sex-specific) significant increase of T_{eye} and a

significant decrease of T_{beak} in response to acute capture-handling stress. Differences in

temperature change between body regions in response to acute stress are not rare (see Table

1). For instance, Moe et al. (2017) measured a decrease of T_{eye} contrasting with an increase of

T_{feet} in chicken, whereas in captive budgerigars (*Melopsittacus undulatus*) T_{eye} increased

transiently after exposure to a stressor while T_{feet} did not change (Ikkatai and Watanabe,

2015); Table 1). Anatomical and functional reasons can likely explain the difference of stress-

induced changes in surface temperature between body regions in penguins, with for instance

a potential interest in maintaining or increasing eye blood flow to maintain or enhance visual

acuity when acutely threatened by an environmental challenge.

T_{eye} has also been shown to positively correlate with internal (cloacal) temperature in

various avian species (budgerigars: Ikkatai and Watanabe (2015); chicken: Cândido et al.

(2020)). Consequently, it is possible that the increase in T_{eye} we observe in adults could reflect

an increase in internal body temperature (Oka, 2018). Consistently, in closely related Emperor

penguins (*Aptenodytes forsteri*), acute stress leads to an approximate 1.5°C increase in

internal (stomach) temperature (Regel and Pütz, 1997). Such an increase in T_{eye} in response to

handling stress has been reported two times in avian species to the best of our knowledge

(Ikkatai and Watanabe, 2015; Jakubas et al., 2022), while most previous studies have shown

either a significant decrease (6 studies) or no significant change (2 studies; Table 1). Those

contrasted findings might be explained by at least three factors: 1. variation in body size, since

smaller individuals/species are expected to favour internal heat conservation by reducing

more markedly peripheral blood flow and surface temperature compared to large ones

(*thermoprotective hypothesis*; Robertson et al., 2020a); 2. differences in the thermal

environment, since it has been shown that during acute stress response, heat conservation is

favoured below the thermoneutral zone while heat dissipation is favoured above its upper

limit (Robertson et al., 2020a); 3. the various delays at which T_{eye} was measured in response

to acute stress, since for instance the increase in T_{eye} found by Ikkatai and Watanabe (2015)

was short-lived (visible 5 min after the stressor, but not later on). Although the interaction

between handling stress and sex was only marginally significant (p = 0.060), it appeared that

only females exhibited a noticeable increase in T_{eye} in response to handling stress in our study.

There are known sex-differences in stress physiology (Handa and McGivern, 2017), but sex

differences in surface temperature changes induced by acute stress have been rarely

investigated (but see Robertson et al. (2020a), for a result opposite to ours between sexes).

We previously observed no sex effect in HPA responsiveness between males and females king

penguins (Viblanc et al., 2016), but as mentioned above SAM is the likely driver of acute

changes in surface temperatures. Consequently, this suggests that sexes might differ in the

stress-sensitivity of their SAM, although this would need to be confirmed by direct

measurements of the SAM activity.

The decrease we observe in T_{beak} in response to acute stress in adults appears like the

typical peripheral vasoconstriction response previously reported for instance by Tabh et al.

(2021). Such peripheral vasoconstriction enables the redistribution of the peripheral blood

circulation to internal organs and tissues, such as the brain or muscles, favouring their

oxygenation and nutrition to sustain the fight-or-flight response. King penguins breed in a

highly dense and aggressive colonial environment (up to 100 aggressive interactions per hour;

Côté, 2000). Aggressive interactions, frequently leading to injuries, are known to result in

increased heart rates (Viblanc et al., 2012). A decrease of T_{beak} during acute stress response

likely reflects peripheral vasoconstriction, that is likely widespread to other peripheral body

parts, which according to the *haemoprotective hypothesis* could help to reduce blood loss in

case of injury (Robertson et al., 2020a).

The changes observed in surface temperatures in chicks differ from those found in

adults. Indeed, we observed a significant decrease in chick's T_{eye} whereas T_{eye} increased in

adult females and stayed stable in adult males. While T_{beak} significantly decreased as observed

in adults, the decrease was even more pronounced in chicks (-3.3 ± 0.6°C *vs.* -1.5 ± 0.4°C, in

chicks and adults, respectively). This is likely explained by greater peripheral thermal losses in

chicks suddenly exposed to an ambient temperature drop (*ca.* 35°C under the brood pouch of

the adults *vs.* 15°C when taken out for measurement and handling), despite their ability to

maintain their internal temperature at this stage and ambient temperature (Duchamp et al.,

2002). The more pronounced drop in surface temperature we observed in chicks during acute

handling could reflect their greater need to conserve heat at a relatively mild T_a compared to

adults.

4.3. Impact of parental CORT on egg and chick surface temperature

Contrary to our prediction, we observed no significant impact of parental CORT

treatment on egg temperature, suggesting little alteration of incubation behaviour by

increased CORT levels, contrary to a previous study in Adelie penguin reporting a reduction of

1.3 ± 0.2°C in T_{egg} (Thierry et al., 2013a). It is possible that incubating two eggs in the colder

environment of Adélie penguins is more challenging (and thus more likely to be impacted by

high CORT levels) than incubating a single egg under the milder climate experienced by king

penguins. Alternatively, it is also possible that king penguins are more resilient to increased

CORT levels than Adelie penguins, which is supported by the good reproductive success of

king penguins implanted with CORT (Stier et *al.* unpublished), contrary to what has been

observed in Adelie penguins (Thierry et al., 2013b). Measurements of incubation quality using

a dummy egg with temperature and rotation sensors (Thierry et al., 2013a) may provide more

accurate data on this question, but it is important to note that the T_{egg} measured in this study

(Placebo: 34.9 ± 0.3°C) was close to the T_{egg} measured in the same penguin colony using

internal sensors in dummy eggs (35.7 ± 0.4°C; Groscolas et al., 2000). Our results therefore

suggest that further studies may benefit from the use of minimally invasive thermal imaging

to measure incubation quality, for instance in the context of parental behaviour and climate

change (Cook et al., 2020).

The measure of chick surface temperatures revealed some surprising patterns. While

we expected CORT chicks to have lower surface temperatures due to poorer parental care

(less efficient brooding), we found the opposite result, but only when chicks were brooded by

the non-implanted partner of the CORT-parent. This result should be considered with caution

considering the limited sample size available for chick surface temperatures (N = 16). One

possibility is that partners from CORT-implanted individuals somehow perceived the 'stress'
levels of their partner (as shown between siblings; Noguera et al., 2017) and somehow
compensated for parental care through more efficient brooding. This hypothesis requires
further study to determine whether the concept of 'family-transmitted stress' (Noguera et al.,
2017) applies to the king penguin. In this species, both parents must rely on each other over
more than one year to successfully fledge their single chick, which itself is entirely dependent
on its parents for its food supply throughout its growth. Hence, the king penguin could be
promising system to investigate the 'family-transmitted stress' hypothesis.

4.4. Conclusion

Evaluating stress levels and responses of free-living animals in a non-invasive manner
using thermal imaging is a burgeoning field of research (Jerem et al., 2015; Tabh et al., 2022).
However, little is still known about the underlying pathways of stress physiology that influence
the changes in surface temperature. Our experimental study in king penguin points toward a
likely preponderant role of the SAM. Additionally, by summarizing the current evidence for
acute changes in surface temperatures during stress exposure (Table 1), we highlight the
relative complexity and inconsistency between the effects observed by different studies.
Further experimental studies related to SAM and HPA involvement are thus required to clarify
what stress component(s) we are measuring through non-invasive thermal imaging.

5. Declarations

5.1. Author contribution

Study design: AS, VAV, PB and AL. Funding acquisition: JPR, PB, VAV, AS. Data collection in the
field: AS, SA, TH, LA, CG, and JPR. Data collection in the lab: ER and AS. Data analysis: CW, AL,
PB, AS. Writing original draft: AL, PB and AS. Writing review and editing: VAV, JPR, AN, SA, TH,
LA, ER and CG.

5.2 Data availability

The datasets used in this manuscript are available on *FigShare* through a private link for review

purposes (https://figshare.com/s/73b39dff978b0dfd2545), and public access with the

following doi (10.6084/m9.figshare.20134181) will be enabled pending acceptance of the

manuscript.

5.3. Acknowledgments

We are grateful to the French Polar Institute (IPEV) and the Terres Australes et Antarctiques

Françaises for providing financial and logistical support for this study through the polar

program #119 (ECONERGY). We wish to thank the Zone Atelier Antarctique et Terres Australes

(ZATA) from the CNRS for financial support, and the members of the Alfred Faure field station

for their help and support in the field. AL was supported by ISblue project, Interdisciplinary

graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the

French government under the program "Investissements d'Avenir" embedded in France 2030.

AS was supported by a 'Turku Collegium for Science and Medicine' Fellowship and a Marie

Sklodowska-Curie Postdoctoral Fellowships (#894963).

5.4. Competing interests

Antoine Stier is an Associate Editor of *J Thermal Biol* but took no part in the peer review and

decision-making processes for this research article.

6. References

Barbraud, C., Delord, K., Bost, C.A., Chaigne, A., Marteau, C., Weimerskirch, H., 2020.
Population trends of penguins in the French Southern Territories. Polar Biology 1–18.
https://doi.org/10.1007/s00300-020-02691-6

Cabanac, A.J., Guillemette, M., 2001. Temperature and heart rate as stress indicators of
handled common eider. Physiol Behav 74, 475–479. https://doi.org/10.1016/s0031-
9384(01)00586-8

Cândido, M.G.L., Tinôco, I.F.F., Albino, L.F.T., Freitas, L.C.S.R., Santos, T.C., Cecon, P.R., Gates,
R.S., 2020. Effects of heat stress on pullet cloacal and body temperature. Poultry Sci 99,
2469–2477. https://doi.org/10.1016/j.psj.2019.11.062

Carere, C., Oers, K. van, 2004. Shy and bold great tits (Parus major): body temperature and
breath rate in response to handling stress. Physiol Behav 82, 905–912.
https://doi.org/10.1016/j.physbeh.2004.07.009

Cook, T.R., Martin, R., Roberts, J., Häkkinen, H., Botha, P., Meyer, C., Sparks, E., Underhill,
L.G., Ryan, P.G., Sherley, R.B., 2020. Parenting in a warming world: thermoregulatory
responses to heat stress in an endangered seabird. Conservation Physiology 8, 2283–13.
https://doi.org/10.1093/conphys/coz109

CôTé, S.D., 2000. Aggressiveness in king penguins in relation to reproductive status and
territory location. Anim Behav 59, 813–821. https://doi.org/10.1006/anbe.1999.1384

Deavers, D.R., Musacchia, X.J., 1979. The function of glucocorticoids in thermogenesis. Fed
Proc 38, 2177–81.

Duchamp, C., Rouanet, J.-L., Barre, H., 2002. Ontogeny of thermoregulatory mechanisms in
king penguin chicks (Aptenodytes patagonicus). Comparative Biochemistry and
Physiology, Part A 131, 765–773. https://doi.org/10.1016/s1095-6433(02)00014-4

Edgar, J.L., Lowe, J.C., Paul, E.S., Nicol, C.J., 2011. Avian maternal response to chick distress.
Proc Royal Soc B Biological Sci 278, 3129–3134. https://doi.org/10.1098/rspb.2010.2701

Edgar, J.L., Nicol, C.J., Pugh, C.A., Paul, E.S., 2013. Surface temperature changes in response
to handling in domestic chickens. Physiol Behav 119, 195–200.
https://doi.org/10.1016/j.physbeh.2013.06.020
571
Edgar, J.L., Nicol, C.J., 2018. Socially-mediated arousal and contagion within domestic chick
broods. Sci Rep 8, 10509. https://doi.org/10.1038/s41598-018-28923-8

Ellenberg, U., Setiawan, A.N., Cree, A., Houston, D.M., Seddon, P.J., 2007. Elevated hormonal
stress response and reduced reproductive output in Yellow-eyed penguins exposed to
unregulated tourism. Gen Comp Endocr 152, 54–63.
https://doi.org/10.1016/j.ygcen.2007.02.022

Giloh, M., Shinder, D., Yahav, S., 2012. Skin surface temperature of broiler chickens is
correlated to body core temperature and is indicative of their thermoregulatory status.
Poultry Sci 91, 175–188. https://doi.org/10.3382/ps.2011-01497

Groscolas, R., Decrock, F., Thil, M.-A., Fayolle, C., Boissery, C., Robin, J.-P., 2000. Refeeding
signal in fasting-incubating king penguins: changes in behavior and egg temperature. Am J
Physiology-regulatory Integr Comp Physiology 279, R2104–R2112.
https://doi.org/10.1152/ajpregu.2000.279.6.r2104

Handa, R.J., McGivern, R.F., 2017. Reference Module in Neuroscience and Biobehavioral
Psychology. https://doi.org/10.1016/b978-0-12-809324-5.02865-0

Handrich, Y., Bevan, R.M., Charrassin, J.-B., Butler, P.J., Ptz, K., Woakes, A.J., Lage, J., Maho,
Y.L., 1997. Hypothermia in foraging king penguins. Nature 388, 64–67.
https://doi.org/10.1038/40392

Herborn, K.A., Graves, J.L., Jerem, P., Evans, N.P., Nager, R., McCafferty, D.J., McKeegan,
D.E.F., 2015. Skin temperature reveals the intensity of acute stress. Physiol Behav 152,
225–230. https://doi.org/10.1016/j.physbeh.2015.09.032

Herborn, K.A., Jerem, P., Nager, R.G., McKeegan, D.E.F., McCafferty, D.J., 2018. Surface
temperature elevated by chronic and intermittent stress. Physiol Behav 191, 47–55.
https://doi.org/10.1016/j.physbeh.2018.04.004

Ikkatai, Y., Watanabe, S., 2015. Eye surface temperature detects stress response in
budgerigars (Melopsittacus undulatus). Neuroreport 26, 642–646.
https://doi.org/10.1097/wnr.0000000000000403

Jakubas, D., Wojczulanis-Jakubas, K., Grissot, A., Devogel, M., Cendrowska, M., Chastel, O.,
2022. Eye Region Surface Temperature and Corticosterone Response to Acute Stress in a
High-Arctic Seabird, the Little Auk. Animals 12, 499. https://doi.org/10.3390/ani12040499

Jerem, P., Herborn, K., McCafferty, D., McKeegan, D., Nager, R., 2015. Thermal Imaging to
Study Stress Non-invasively in Unrestrained Birds. J Vis Exp Jove 53184.
https://doi.org/10.3791/53184

Jerem, P., Jenni-Eiermann, S., Herborn, K., McKeegan, D., McCafferty, D.J., Nager, R.G., 2018.
Eye region surface temperature reflects both energy reserves and circulating
glucocorticoids in a wild bird. Sci Rep-uk 8, 1907. https://doi.org/10.1038/s41598-018-
20240-4

Jerem, P., Jenni-Eiermann, S., McKeegan, D., McCafferty, D.J., Nager, R.G., 2019. Eye region
surface temperature dynamics during acute stress relate to baseline glucocorticoids
independently of environmental conditions. Physiology & Behavior 112627.
https://doi.org/10.1016/j.physbeh.2019.112627

Knoch, S., Whiteside, M.A., Madden, J.R., Rose, P.E., Fawcett, T.W., 2022. Hot-headed
peckers: thermographic changes during aggression among juvenile pheasants (*Phasianus
colchicus*). Philosophical Transactions Royal Soc B 377, 20200442.
https://doi.org/10.1098/rstb.2020.0442

Lewden, A., Bonnet, B., Nord, A., 2020. The metabolic cost of subcutaneous and abdominal
rewarming in king penguins after long-term immersion in cold water. J Therm Biol 91,
102638. https://doi.org/10.1016/j.jtherbio.2020.102638

Lewden, A., Enstipp, M.R., Bonnet, B., Bost, C., Georges, J.-Y., Handrich, Y., 2017a. Thermal
strategies of king penguins during prolonged fasting in water. J Exp Biol 220, 4600–4611.
https://doi.org/10.1242/jeb.168807

Lewden, A., Enstipp, M.R., Picard, B., Walsum, T. van, Handrich, Y., 2017b. High peripheral
temperatures in king penguins while resting at sea: thermoregulation versus fat
deposition. Journal Of Experimental Biology 220, 3084–3094.
https://doi.org/10.1242/jeb.158980

Lewden, A., Nord, A., Bonnet, B., Chauvet, F., Ancel, A., McCafferty, D.J., 2020. Body surface
rewarming in fully and partially hypothermic king penguins. Journal Of Comparative
Physiology B-Biochemical Systemic And Environmental Physiology 190, 597–609.
https://doi.org/10.1007/s00360-020-01294-1

MacDougall-Shackleton, S.A., Bonier, F., Romero, L.M., Moore, I.T., 2019. Glucocorticoids
and "Stress" Are Not Synonymous. Integr Org Biology 1, obz017.
https://doi.org/10.1093/iob/obz017

McCafferty, D.J., 2013. Applications of thermal imaging in avian science. Ibis 155, 4–15.
https://doi.org/10.1111/ibi.12010

McCafferty, D.J., Koprowski, R., Herborn, K., Tattersall, G.J., Jerem, P., Nord, A., 2021.
Advances in Thermal Imaging. J Therm Biol 102, 103109.
https://doi.org/10.1016/j.jtherbio.2021.103109

Moe, R.O., Bohlin, J., Flø, A., Vasdal, G., Stubsjøen, S.M., 2017. Hot chicks, cold feet. Physiol
Behav 179, 42–48. https://doi.org/10.1016/j.physbeh.2017.05.025

Narushin, V.G., 2005. Egg geometry calculation using the measurements of length and
breadth. Poultry Sci 84, 482–484. https://doi.org/10.1093/ps/84.3.482

Noguera, J.C., Kim, S.-Y., Velando, A., 2017. Family-transmitted stress in a wild bird.
Proceedings of the National Academy of Sciences 27, 201706164.
https://doi.org/10.1086/680688

Oka, T., 2018. Chapter 35 Stress-induced hyperthermia and hypothermia. Handb Clin
Neurology 157, 599–621. https://doi.org/10.1016/b978-0-444-64074-1.00035-5

Ouyang, J.Q., Macaballug, P., Chen, H., Hodach, K., Tang, S., Francis, J.S., 2021. Infrared
thermography is an effective, noninvasive measure of HPA activation. Ann Ny Acad Sci 24,
1–6. https://doi.org/10.1080/10253890.2020.1868431

Palme, R., 2018. Non-invasive measurement of glucocorticoids: Advances and problems.
Physiol Behav 199, 229–243. https://doi.org/10.1016/j.physbeh.2018.11.021

Pijpers, N., Heuvel, H. van den, Duncan, I.H., Yorzinski, J., Neethirajan, S., 2022.
Understanding Chicks' Emotions: Are Eye Blinks & Facial Temperatures Reliable Indicators?
Biorxiv 2022.01.31.478468. https://doi.org/10.1101/2022.01.31.478468

Playà-Montmany, N., Tattersall, G.J., 2021. Spot size, distance and emissivity errors in field
applications of infrared thermography. Methods Ecol Evol 12, 828–840.
https://doi.org/10.1111/2041-210x.13563

Regel, J., Pütz, K., 1997. Effect of human disturbance on body temperature and energy
expenditure in penguins. Polar Biol 18, 246–253. https://doi.org/10.1007/s003000050185

Robertson, J.K., Mastromonaco, G., Burness, G., 2020a. Evidence that stress-induced
changes in surface temperature serve a thermoregulatory function. J Exp Biol 223,
jeb213421. https://doi.org/10.1242/jeb.213421

Robertson, J.K., Mastromonaco, G.F., Burness, G., 2020b. Social hierarchy reveals
thermoregulatory trade-offs in response to repeated stressors. J Exp Biol 223, jeb229047.
https://doi.org/10.1242/jeb.229047

Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research.
Trends Ecol Evol 19, 249–255. https://doi.org/10.1016/j.tree.2004.03.008

Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How Do Glucocorticoids Influence Stress
Responses? Integrating Permissive, Suppressive, Stimulatory, and Preparative Actions.
Endocr Rev 21, 55–89. https://doi.org/10.1210/edrv.21.1.0389

Stier, A., Schull, Q., Bize, P., Lefol, E., Haussmann, M., Roussel, D., Robin, J.-P., Viblanc, V.A.,
2019. Oxidative stress and mitochondrial responses to stress exposure suggest that king
penguins are naturally equipped to resist stress. Scientific Reports 9, 8545.
https://doi.org/10.1038/s41598-019-44990-x

Tabh, J.K.R., Burness, G., Wearing, O.H., Tattersall, G.J., Mastromonaco, G.F., 2021. Infrared
thermography as a technique to measure physiological stress in birds: Body region and
image angle matter. Physiological Reports 9, e14865.
https://doi.org/10.14814/phy2.14865

Tabh, J.K.R., Mastromonaco, G.F., Burness, G., 2022. Stress-induced changes in body surface
temperature are repeatable, but do not differ between urban and rural birds. Oecologia
1–15. https://doi.org/10.1007/s00442-022-05120-z

Thierry, A.-M., Massemin, S., Handrich, Y., Raclot, T., 2013a. Elevated corticosterone levels
and severe weather conditions decrease parental investment of incubating Adélie
penguins. Hormones And Behavior 63, 475–483.
https://doi.org/10.1016/j.yhbeh.2012.12.011

Thierry, A.-M., Ropert-Coudert, Y., Raclot, T., 2013b. Elevated corticosterone levels decrease
reproductive output of chick-rearing Adélie penguins but do not affect chick mass at
fledging. Conservation Physiology 1, cot007. https://doi.org/10.1093/conphys/cot007

Thomas, D.B., Ksepka, D.T., Fordyce, R.E., 2011. Penguin heat-retention structures evolved in
a greenhouse Earth. Biol Letters 7, 461–464. https://doi.org/10.1098/rsbl.2010.0993

Torres-Medina, F., Cabezas, S., Marchant, T.A., Wikelski, M., Romero, L.M., Hau, M., Carrete,
M., Tella, J.L., Blas, J., 2018. Corticosterone implants make stress hyporesponsive birds.
The Journal of Experimental Biology jeb.173864-43. https://doi.org/10.1242/jeb.173864

Viblanc, V.A., Dobson, F.S., Stier, A., Schull, Q., Saraux, C., Gineste, B., Pardonnet, S.,
Kauffmann, M., Robin, J.-P., Bize, P., 2016. Mutually honest? Physiological "qualities'
signalled by colour ornaments in monomorphic king penguins. Biological Journal of the
Linnean Society 118, 200–214. https://doi.org/10.1111/bij.12729

Viblanc, V.A., Schull, Q., Cornioley, T., Stier, A., Ménard, J.-J., Groscolas, R., Robin, J.-P., 2018.
An integrative appraisal of the hormonal and metabolic changes induced by acute stress
using king penguins as a model. Gen Comp Endocr 269, 1–10.
https://doi.org/10.1016/j.ygcen.2017.08.024

Viblanc, V.A., Valette, V., Kauffmann, M., Malosse, N., Groscolas, R., 2012. Coping with social
stress: heart rate responses to agonistic interactions in king penguins. Behavioral Ecology
23, 1178–1185. https://doi.org/10.1093/beheco/ars095

Weimer, S.L., Wideman, R.F., Scanes, C.G., Mauromoustakos, A., Christensen, K.D., Vizzier-
Thaxton, Y., 2020. Broiler stress responses to light intensity, flooring type, and leg
weakness as assessed by heterophil-to-lymphocyte ratios, serum corticosterone, infrared
thermography, and latency to lie. Poultry Sci 99, 3301–3311.
https://doi.org/10.1016/j.psj.2020.03.028

Weimerskirch, H., Stahl, J.C., Jouventin, P., 1992. The breeding biology and population
dynamics of king penguins Aptenodytes patagonica on the Crozet Islands. Ibis 134, 107–
117.

Winder, L.A., White, S.A., Nord, A., Helm, B., McCafferty, D.J., 2020. Body surface
temperature responses to food restriction in wild and captive great tits. Journal Of
Experimental Biology 223, jeb220046-8. https://doi.org/10.1242/jeb.220046

Wingfield, J.C., Romero, L.M., 2015. Tempests, Poxes, Predators, and People 149–196.
https://doi.org/10.1093/acprof:oso/9780195366693.003.0005

Zuluaga, J.D., Danner, R.M., 2023. Acute stress and restricted diet reduce bill-mediated heat
dissipation in the song sparrow (*Melospiza melodia*): implications for optimal
thermoregulation. J Exp Biol 226. https://doi.org/10.1242/jeb.245316

722