Surface temperatures are influenced by handling stress

independently of corticosterone levels in wild king penguins

(Aptenodytes patagonicus)

Agnès Lewden[1,2†], Chelsea Ward[3], Aude Noiret[4], Sandra Avril[5], Lucie Abolivier[5], Caroline

Gérard[5], Tracey L. Hammer[5], Émilie Raymond[4,5], Jean-Patrice Robin[5], Vincent A. Viblanc[5*],

Pierre Bize[3,6*] and Antoine Stier[4,5,7*†]

[1] Faculty of Biological Sciences, University of Leeds, Leeds, UK

[2] Université de Brest - UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des sciences de

l'environnement marin - IUEM - Rue Dumont D'Urville - 29280 - Plouzané

[3] School of Biological Sciences, University of Aberdeen, Aberdeen, UK

[4] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622,

Villeurbanne,

[5] Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

[6] Swiss Ornithological Institute, CH-6204 Sempach, Switzerland

[7] Department of Biology, University of Turku, Turku, Finland

[†]: corresponding authors: agnes.lewden@univ-brest.fr & antoine.stier@iphc.cnrs.fr

[*]: share co-senior authorship

Highlights:

- Experimental increase in corticosterone does not affect king penguin's surface temperature
- Acute handling stress increases eye but decreases beak surface temperature in adults
- Acute handling decreases both eye and beak surface temperatures in young chicks
- Parental corticosterone treatment does not affect egg surface temperature during incubation
- Chicks brooded by non-implanted partners of corticosterone parents are warmer than chicks brooded by others adults.

Abstract

Assessing the physiological stress responses of wild animals opens a window for understanding how organisms cope with environmental challenges. Since stress response is associated with changes in body temperature, the use of body surface temperature through thermal imaging could help to measure acute and chronic stress responses non-invasively. We used thermal imaging, acute handling-stress protocol and an experimental manipulation of corticosterone (the main glucocorticoid hormone in birds) levels in breeding king penguins (*Aptenodytes patagonicus*), to assess: 1. the potential contribution of the Hypothalamo-Pituitary-Adrenal (HPA) axis in mediating chronic and acute stress-induced changes in adult surface temperature, 2. the influence of HPA axis manipulation on parental investment through thermal imaging of eggs and brooded chicks, and 3. the impact of parental treatment on offspring thermal's response to acute handling.

Maximum eye temperature (T_{eye}) increased and minimum beak temperature (T_{beak}) decreased in response to handling stress in adults, but neither basal nor stress-induced surface temperatures were significantly affected by corticosterone implant. While egg temperature was not significantly influenced by parental treatment, we found a surprising pattern for chicks: chicks brooded by the (non-implanted) partner of corticosterone-implanted individuals exhibited higher surface temperature (both T_{eye} and T_{beak}) than those brooded by glucocorticoid-implanted or control parents. Chick's response to handling in terms of surface temperature was characterized by a drop in both T_{eye} and T_{beak} independently of parental treatment.

We conclude that the HPA axis seems unlikely to play a major role in determining chronic or acute changes in surface temperature in king penguins. Changes in surface temperature may primarily be mediated by the Sympathetic-Adrenal-Medullary (SAM) axis in response to stressful situations. Our experiment did not reveal a direct impact of parental HPA axis manipulation on parental investment (egg or chick temperature), but a potential influence on the partner's brooding behaviour.

Key words: Thermal imaging, Stress, Corticosterone, Heterothermy, Bird, Beak, Eye

Abstract:

Abbreviations:

Adrenocorticotropic hormone: ACTH

Ambient air temperature: T_a

Corticosterone: CORT

Glucocorticoid hormones: GC

Egg surface temperature: T_{egg}

Hypothalamic-pituitary-adrenal: HPA

Linear mixed models: LMMs

Maximum eye region surface temperature: T_{eye}

Minimum beak surface temperature: T_{beak}

Sympathetic-adrenal-medullary: SAM

1. Introduction

Measuring stress response is of central interest in wild animals to understand how they cope with environmental change (Ellenberg et al., 2007; Romero, 2004). Traditionally, ecologists have evaluated acute stress by assessing the activation of the Sympathetic-Adrenal-Medullary (SAM) or Hypothalamic-Pituitary-Adrenal (HPA) axes, either at baseline levels or in response to acute disturbances in the environment, by metrics such as increased heart rate (Cabanac and Guillemette, 2001; Viblanc et al., 2012), ventilation rate (Carere and Oers, 2004) or glucocorticoid hormone (GC) secretion (Sapolsky et al., 2000). Stress exposure is also known to trigger core body and surface temperature changes (Oka, 2018). As the latter can be measured using thermal imaging (McCafferty, 2013), it led to a recent burgeoning of thermal imaging studies to measure the stress response of captive and wild endotherms (McCafferty et al., 2021). Both the SAM and HPA axes can potentially influence changes in body surface temperature, but the importance of these two pathways in mediating the stress response(s) measured through thermal imaging remains largely unknown (but see Jerem and Romero (2023)). Considering that both HPA and SAM axes provide strong and very reactive responses to stress, teasing apart their contribution to changes in peripheral temperature is challenging.

SAM axis responds within seconds to a stressor by releasing catecholamines hormones (adrenaline and noradrenaline) into the blood stream, which induces an immediate increase in heart rate (tachycardia) and vasoconstriction of peripheral blood vessels (Wingfield and Romero, 2015). This redistribution of peripheral blood flow to essential internal organs generally leads to a response called stress-induced hyperthermia along with a decrease of surface temperature (Cabanac and Guillemette, 2001; Herborn et al., 2015; Oka, 2018). The SAM axis response is followed by the slower activation of HPA axis that releases GC into the blood stream within a few minutes (Wingfield and Romero, 2015). GC are metabolic hormones

that helps responding to a stressor by mobilizing energy resources and by triggering an

increase in behavioural activity and metabolic rate (Wingfield and Romero, 2015). Circulating

GC levels are usually considered at baseline (*i.e.* without an inducing stressor) or at stress-

induced levels, which are often interpreted as mirroring chronic stress/metabolic demand and

acute stress response respectively. Yet, the actual interpretation of circulating GC levels is

likely more complex (MacDougall-Shackleton et al., 2019). GC-mediated changes in

metabolism may cause an increase in internal heat production that, in turn, may lead to

changes in body surface temperature (Oka, 2018). Additionally, previous evidence suggests

that GC might be necessary for enabling the SAM axis to induce shivering, free fatty acid

mobilization or vasoconstriction (Deavers and Musacchia, 1979).

Therefore, both the SAM and HPA axes could trigger, independently from each other, or

together, a change in body core and/or surface temperatures (Ouyang et al., 2021). The

immediate activation of SAM axis could lead to a decrease of peripheral temperature

independently of GC release by the HPA axis. On the other hand, the short to long-term

elevation of baseline GC might lead to increased metabolic rate, heat production and,

ultimately, higher peripheral temperature (to facilitate heat dissipation and maintain

homeothermy) independently of the SAM axis. The use of thermal imaging as a non-invasive

tool to measure stress in unmanipulated animals is actively developing, but the implication of

the SAM and HPA axes in mediating changes in surface temperature remains uncertain (Jerem

and Romero, 2023; Ouyang et al., 2021). For instance, data in captive and wild birds show that

acute stress exposure often leads to a decrease in body surface temperature (reviewed in

Table 1, but note some discrepancies between species and/or body parts), as predicted if

these changes are driven by the SAM axis. However, evidence for an impact of chronic

stressors on body surface temperature (Herborn et al., 2018; Robertson et al., 2020a, 2020b;

Winder et al., 2020) or for a correlation between circulating GC levels and body surface

temperature (Giloh et al., 2012; Jerem et al., 2018; Weimer et al., 2020) remains unclear.

In this study, we took up the challenge of distinguishing the role of HPA and SAM axes

on changes in body surface temperatures using an experimental approach in adult king

penguins (*Aptenodytes patagonicus*). Breeding king penguins were subcutaneously implanted

with either a corticosterone (CORT, the main GC hormone in avian species) implant or a

placebo implant. The use of CORT implants in king penguin (see methods and Fig. 1), as in

many other bird species (see Torres-Medina et al., 2018), leads to higher baseline CORT levels,

while inhibiting the acute CORT response in the days following implantation. Indeed, CORT-

implanted birds no longer exhibit high blood CORT release in response to acute stress induced

by a handling stress protocol because of the HPA negative feedback loop (Torres-Medina et

al., 2018). Specifically, high circulating CORT levels are inhibiting the activity of the

paraventricular nucleus in the hypothalamus through binding with glucocorticoid receptors.

This inhibits the release of corticotrophin-releasing hormone and the afferent HPA cascade

through the pituitary (secreting the adrenocorticotropic hormone, ACTH) and adrenal

(secreting CORT) glands (Smulders, 2021). Hence, the use of CORT implants in conjunction

with handling stress allows for the separation of the contributions of the HPA and SAM axes

to acute changes in body surface temperatures. Here, we therefore measured the effects of

our CORT treatment, together with a handling stress protocol, on surface temperatures in the

king penguin. If changes in surface temperatures are mainly driven by activating the HPA axis,

we expect to find higher initial body surface temperatures in CORT-implanted individuals (with

elevated baseline CORT) prior to capture and handling. However, because CORT implants shut-

down the release of CORT in response to handling, we expect to find no change in body surface

temperature in response to handling stress in CORT individuals if such changes are mediated

by the HPA axis only. On the contrary, if changes in surface temperatures are mainly driven by

the activity of the SAM axis, we expect our CORT treatment to have no effect on body surface

temperatures (both before or after capture and handling).

In addition to direct effects on adults, parental stress exposure or increased CORT

levels may also affect the following generation, for instance by affecting parental care (Thierry

et al., 2013a) or by conveying information about a stressful environment (Brandl et al., 2022).

The latter idea has been framed as the concept of family-transmitted stress by Noguera et al.

(2017). Therefore, we also aimed to investigate whether thermal imaging could be used to

quantify indirect effects of stress on king penguin's progeny. To this end, we measured egg

surface temperature on the same day as the body surface temperatures of implanted parents,

and the surface temperatures of their chicks. Based on experimental data in Adélie penguin

(*Pygoscelis adeliae*) showing that elevated baseline CORT levels in adults led to lower parental

care (Thierry et al., 2013a), we hypothesized that increasing CORT levels may lower the surface

temperatures of eggs and/or chicks during incubation and brooding respectively, as parents

may pay less attention in keeping the egg/chick covered and warm under their brood pouch.

2. Material and Methods

2.1. Study site and species

The king penguin is a heterothermic bird whose body temperature largely varies between core and peripheral tissues (Lewden et al., 2017a; Lewden et al., 2017b). Its thick body plumage provides a very efficient insulation, while counter-current vascular heat exchangers in the appendages allows for a decrease in peripheral temperature loss (Thomas et al., 2011). These adaptations allow birds to conserve heat when foraging in cold Antarctic waters (Handrich et al., 1997), and to rapidly recover to normothermia when exiting the water (Lewden et al., 2020).

The study was conducted in 2018-2019 in a king penguin colony of *ca.* 22,000 breeding birds at 'La Baie du Marin' on Possession Island (Barbraud et al. 2020), Crozet Archipelago, in the Southern Indian Ocean (46°26' S, 51°52' E). We followed 49 breeding pairs from courtship (early November) until the onset of the Austral winter (April). All adults were identified by a hair dye mark on the breast feathers. During the early breeding season, male and female alternate between periods on-land caring for their single-egg or chick, and periods foraging at sea (Weimerskirch et al., 1992). The male takes the first incubation shift while the female forages at sea. The female returns ~15 days later to relieve her partner, and the parents continue to alternate shifts throughout incubation (~53 days) and early chick brooding. Chicks become thermally emancipated from the parents around one month of age, allowing both parents to go back at sea simultaneously to forage (Weimerskirch et al., 1992). Twenty days after hatching, king penguin chicks are able to maintain their core body temperature under mild environmental conditions (> 10°C) but still rely on their parent's brooding behaviour (Duchamp et al., 2002), and are able to express a HPA-dependent stress response (X et al. unpublished).

2.2. Use of sub-cutaneous CORT and SHAM implants

We divided the 49 breeding pairs into two groups of either female-treated pairs (N = 23) or male-treated pairs (N = 26). In the female-treated pairs, we implanted females with either a CORT (NG-111, 50 mg corticosterone, 90-days release, 8 mm diameter and 3mm thickness; N = 12 females) or a Placebo (NC-111, same vehicle but no CORT; N = 11 females) implant. Sub-cutaneous implants were purchased from Innovative Research of America (Sarasota, FL, USA). Females were implanted during their first incubation shift, 3 days after returning from sea. Similarly, in the male-treated pairs, we implanted 13 males with a CORT and 13 males with a Placebo implant, 3 days after returning from sea. In males, the first incubation shift is immediately after egg laying, and males have already been fasting for ~15 days during courtship. Therefore, we chose to implant the males during their second incubation shift so that they would have comparable fasting times compared to females, and therefore both sexes would be in a comparable physiological state during measurements. Allocation of birds to specific groups was designed to avoid spatial or temporal biases within and between groups. Male and female partners in, respectively, the female-treated and male-treated pairs were not implanted. Hereafter, they are referred to as "Partner-CORT" or "Partner-Placebo" regardless of their sex. Partner penguins were only considered in the chick analysis (see below).

On the day of the implantation, adults were captured by hand while incubating their egg and restrained by one experimenter with a hood covering the head to keep them calm. While maintaining individuals on their territory within the colony to avoid risking reproductive abandonment, sub-cutaneous CORT or placebo implants were inserted in the upper part of the back by a veterinary surgeon under local anaesthesia (*ca.* 0.5 mg/kg xylocaine combined

with 0.0001 mg/kg adrenalin, Aspen Pharma). Immediately after implantation, the small

incision (*ca.* 15 mm) was closed using 3 sterile surgical staples (SurgiClose^{TM} Skin Stapler), and

a prophylactic dose of antibiotics (cephalexin *ca.* 50 mg/kg, Rilexine®, Virbac) was injected to

prevent any risk of infection. The incision was checked 3 days later for implant rejection; no

sign of infection or implant rejection were noticed and staples were removed 6 days after

implantation.

To verify the efficacy of our CORT treatment, 6 days after implantation, 1 mL of blood

was sampled from the flipper vein using a heparinized syringe in less than 4 min after being

detected by the focal individual (baseline CORT) and after 30 min of standardized handling

(stress-induced, see Stier et al. (2019) and Viblanc et al. (2018)). Blood samples were taken

between 09:00 and 13:00 to limit variation linked to circadian variation in plasma CORT levels.

Blood samples were centrifuged within one hour of sampling at 3000 g for 10 minutes. The

plasma fraction was then removed and plasma aliquots were stored at −20°C until the end of

the day, before being transferred to −80°C until laboratory analyses in 2023. Plasma total

CORT levels were measured from 25 μL of plasma by immunoassay according to guidelines

provided by the manufacturer (Corticosterone Enzyme Immunoassay Kit, Arbor Assay, USA).

Intra-plate coefficient of variation based on duplicates was 8.17 ± 0.90% (mean ± SE), and

inter-plate coefficient of variation based on one repeated sample was 8.08%. We tested the

effect of implant on log-transformed plasma CORT using a linear mixed model (LMM) including

as fixed factors: Handling stress (Baseline *vs.* Handling stress), Treatment (CORT *vs.* Placebo)

and their interaction, as well as penguin ID as random effect. Implants were successful in

raising circulating baseline CORT levels and in inhibiting stress-induced CORT release (Fig. 1).

2.3. Thermal image collection and analysis

Because of their dense and watertight plumage, the use of thermal imaging in seabirds

is usually restricted to measuring the eye (periorbital region) and beak regions (Gauchet and

Grémillet, 2022). These regions are highly vascularised and provide valuable information on

body surface temperatures. Thermal pictures of implanted adults (Fig. 2A) and their egg (Fig.

2B) were taken 3 days after implantation and thermal pictures of chicks (Fig. 2C) were taken

at day 20 after hatching (mean ± SE: 49.8 ± 1.4 days after CORT or Placebo implant of the focal

parent, *i.e.* within the 90-days release given by the CORT pellet manufacturer). Thermal

pictures were taken at a distance between *ca.* 0.8 and 2 m using FLIR E8 thermal imaging

camera (320 x 240 pixels) resulting in a spotsize between 1.95 and 4.88 mm (Playa-Montmany

and Tatersall 2021). King penguins' beaks being approximately 120 mm long, and the eye

region being approximately 18 mm of diameter, our target areas exceeded the 3 times larger

spotsize recommended by Playa-Montmany and Tatersall (2021). When approaching the

adults and/or chicks, we started a stopwatch when the focal individual showed the first signs

of alarm behaviour (*e.g.* stopping activity, looking towards the experimenter) to measure the

duration of our disturbance at the time the different pictures were taken. In adults, we took

a first set of pictures (1 to 2 pictures per individual) before handling, on average (mean ± SE)

0.63 ± 0.07 min after the first sign of alarm behaviour at a distance of *ca*. 1.5 to 2 m from the

focal individuals. Adults were then captured-restrained, incision healing was checked, and we

took a second set of pictures (1 to 2 pictures per individual) after terminating handling stress

(mean ± SE handling stress duration was 8.3 ± 0.3 min; a timeframe within which CORT is

known to increase by *ca.* 100% above baseline levels in king penguins (Viblanc et al., 2018).

Just after capturing the adult, we removed the single-egg from the brood pouch and

immediately (*ca.* within 10 s) took a thermal picture at a distance of *ca.* 0.8 m. We then

measured egg length and breadth using a Vernier calliper (Mitutoyo; accuracy ± 0.02 mm) and

estimated egg volume (cm^3) following (Narushin, 2005). When chicks were 20 days old, we

took a first set of pictures (1 to 2 pictures per individual) at a distance of *ca.* 0.8 m immediately

(*ca.* within 10 s) after they were removed from the brood pouch (*i.e.* before handling: 0.73 ±

0.08 min after the first sign of alarm behavior from the parent) and a second set of pictures

(1-2 pictures per individual) was taken after handling, 5.40 ± 0.30 min later. Thermal images

of the chicks were only taken on non-rainy days, which reduced the sample size from 38 chicks

that survived until day 20, to 16 chicks with thermal images available. Chick sex was unknown

and body mass was not significantly influenced by the parental treatment (CORT: 1.14 ± 0.11

kg *vs.* Placebo 1.32 ± 0.09 kg, N = 8/group, t-test: t = -1.28, p = 0.22). Chicks were kept within

the colony at ambient temperature (14.9 ± 0.9°C) between the first and second set of pictures,

a temperature at which they are known to be able to maintain their internal body temperature

(Duchamp et al., 2002). In adults and chicks, we defined 'before handling' surface

temperatures, the temperatures extracted from the first set of pictures taken in < 1 min and

'after handling' surface temperatures, the temperatures extracted from the second set of

pictures taken after > 5 minutes of handling stress.

We analysed thermal images using the ThermaCAM TM Researcher Pro 2.10 software

(Flir systems, Wilsonville, Oregon, USA). Only profile pictures were analysed to avoid surface

temperature errors related to head orientation (Playà-Montmany and Tattersall, 2021; Tabh

et al., 2021). For each image we set the emissivity at 0.98, reflected temperature at 20°C and

controlled in the analyses for daily variation in air temperature (T_a) and relative humidity using

daily climatic measures from a permanent Météo France weather station located 2 km inland

from the colony (https://rp5.ru/Archives_m%C3%A9t%C3%A9o_sur_la_base_Alfred-Faure).

We extracted measures of surface temperatures on maximum periorbital temperature (T_{eye})

and minimum beak temperature (T_{beak}) in adults and chicks. We used maximum T_{eye} as

recommended by Jerem et al. (2015 & 2019) as well as recorded in recent studies (*e.g.*

Robertson et al. 2020a; Zuluaga & Danner 2023, and the minimum T_{beak} to gain insight

on the maximum state of vasoconstriction for this body surface (Tattersall et al., 2009). Finally,

we estimated egg surface temperature (T_{egg}) as the mean of average length and width surface

temperature.

2.4. Ethical note

All the procedures were approved by the French Ethical Committee (APAFIS#16465-

2018080111195526 v4) and the Terres Australes et Antarctiques Françaises (Arrêté TAAF A-

2018-118).

2.5. Statistical analysis

We ran separate analyses for adults, eggs and chicks. In adults, we investigated

variations in T_{eye} and T_{beak} using two separated linear mixed models (LMMs) where Treatment

(CORT *vs.* Placebo), Handling stress (before *vs.* after), Sex and T_a were specified as fixed effects.

Relative humidity was initially tested but removed from final models since it was never

significant. All two-ways interactions were also initially included but removed from the final

model if $p > 0.10$ using a backward stepwise procedure. The p-values (just before removal in

the backward stepwise procedure) for the focal interaction between Treatment and Handling

stress are reported in tables. Bird identity was included as a random intercept to control for

repeated measures. We investigated variation in T_{egg} by entering parental CORT Treatment

(CORT *vs.* Placebo), the Sex of the parent, T_a and egg volume as fixed effects. Eggs were always

measured when incubated by the implanted parents (*i.e.* 3 days after implantation of CORT or

Placebo implants), while chicks at day 20 could be brooded by the implanted parent or its non-

implanted partner. Thus, we investigated variation chick T_{eye} and T_{beak} by entering parental

Treatment (CORT *vs.* Placebo), the category of the brooding parent at the time of

measurement (implanted individual *vs.* non-implanted partner), Handling stress (before *vs.*

after), and T_a as fixed effects, and chick identity as a random intercept. Parental sex was not

included in the models for chick T_{eye} and T_{beak} due to limited sample size available for chicks.

Interactions were treated as described above for adults. All statistical analyses were

conducted using *R* (version 3.5.3) and the packages *lme4* (Bates et al., 2015), *emmeans* (Lenth

et al., 2018) and *pbkrtest* (Halekoh and Højsgaard, 2014). Results are reported as least-square

means ± SE.

3. Results

3.1. Adult surface temperatures

Treatment (CORT *vs.* Placebo), either alone or in interaction with Handling stress, had no significant effect on eye (T_{eye}) or beak (T_{beak}) surface temperature in incubating adults (Table 2, Fig. 3A and 3B). Handling stress significantly affected T_{eye} and T_{beak}, in opposite directions (Table 2). Over an 8-min handling stress, T_{eye} increased by 0.8 ± 0.3°C, on average (Fig. 3A), while T_{beak} decreased by -1.5 ± 0.4°C, on average (Fig. 3B). The increase in T_{eye} appeared to be sex-specific (marginally significant interaction Handling * Sex: $p = 0.060$), with only females showing a significant increase in T_{eye} in response to Handling stress (Fig. 3C; males: $t = 0.63$, $p = 0.53$; females: $t = 3.08$, $p = 0.003$). As expected, T_{eye} and T_{beak} were significantly and positively associated with T_a (Table 2).

3.2. Egg surface temperature

Neither parental Treatment ($F_{1,45} = 1.38$, $p = 0.25$; CORT: 34.5 ± 0.2°C *vs.* Placebo: 34.9 ± 0.3°C; N = 46), Sex ($F_{1,45} = 0.47$, $p = 0.49$), ambient temperature ($F_{1,45} = 0.45$, $p = 0.50$), nor egg volume ($F_{1,45} = 0.01$, $p = 0.93$) significantly affected T_{egg}.

3.3. Chick surface temperatures

The interaction between parental Treatment (CORT *vs.* Placebo) and the category of the parent brooding the chick (implanted *vs.* non-implanted partner) at the time of measurement significantly affected chick T_{eye} ($F_{1,11} = 6.37$, $p = 0.028$, Table 3) and chick T_{beak} ($F_{1,11} = 5.49$, $p = 0.039$, Table 3). Chicks brooded by the partner of CORT-implanted individuals had higher overall T_{eye} and T_{beak} than chicks brooded by CORT parents, or by Placebo and associated partners (Fig 4A and 4B). While we found no significant interaction between

343   parental Treatment and Handling stress (Table 3), both chick T_{eye} (Fig 4C) and chick T_{beak} (Fig

344   4D) significantly decreased after handling (by -0.9 ± 0.3°C and -3.3 ± 0.6°C respectively, Table

345   3). As expected, chick T_{eye} and T_{beak} were significantly positively associated with T_a (Table 3).

4. Discussion

By experimentally manipulating CORT levels in adult breeding king penguins, we show that corticosterone is unlikely to play a major role in determining changes in surface temperatures (both before or in response to handling stress) in king penguins. Indeed, we found no significant difference in baseline surface temperature between CORT and Placebo implanted individuals. However, whatever the parental CORT treatment, our handling stress protocol led to an increase in T_{eye} and a decrease in T_{beak} in response to capture and handling. The absence of effect of the CORT treatment on baseline surface temperature and its dynamics during acute handling stress suggest that changes in surface temperature are probably driven primarily by the activity of the SAM axis in king penguins (but see below for a potential causal involvement of CORT in house sparrow *Passer domesticus* (Ouyang et al., 2021)). Contrary to our prediction, parental treatment had no significant effect on egg temperature. Yet, we found an unexpected pattern for chick surface temperatures: chicks brooded by the non-implanted partner of CORT individuals had higher surface temperatures (both T_{eye} and T_{beak}) than chicks brooded by Placebo (or associated non-implanted partner) or CORT individuals. Finally, in the thermally non-emancipated chicks, both T_{eye} and T_{beak} decreased with handling, but irrespectively of the parental treatment.

4.1. CORT and adult surface temperatures

Previous correlative studies have considered baseline and stress-induced CORT levels to investigate the relationship between GC signalling and changes in body surface temperature in birds (blue tit, *Cyanistes caerulescens*, (Jerem et al., 2018); domestic chicken, *Gallus gallus domesticus,* (Giloh et al., 2012; Weimer et al., 2020)). To go deeper in the involved mechanisms, we experimentally manipulated CORT in the king penguin to help to causally distinguish the contribution of the HPA and SAM axes to changes in body surface temperature. Since we did not detect any significant effect of CORT manipulation on baseline or stress-induced surface temperatures, our results provide little support for a causal involvement of the HPA axis in mediating stress-related changes in surface temperature.

Our results contrast with a recent study showing that an acute stimulation of the HPA axis with adrenocorticotropic hormone (ACTH) induced a decrease in T_{eye} within 5 min in captive-held house sparrows (Ouyang et al., 2021), which suggests a causal involvement of

the HPA axis in mediating stress-related changes in surface temperature. Yet, a recent study

in the same species found a significant correlation between the acute change in surface

temperature and heart rate variability, a measure of SAM axis activation (Jerem and Romero,

2023). This suggests that both the HPA and SAM axes could be involved in mediating stress-

related changes in surface temperature in house sparrow. While we used an experimental

approach and the lack of significant effect of baseline CORT on body surface temperatures

appears to be robust, we cannot fully exclude that our handling stress protocol for thermal

imaging was too short to detect a potential role of stress-induced CORT in influencing changes

in surface temperatures. Indeed, in 8 minutes of handling, CORT levels are expected to double

in king penguin, while maximum CORT levels are only attained after *ca.* 80 minutes of handling

in our study species (Viblanc et al., 2018). Additionally, we may have missed some early

changes in surface temperatures occurring in response to the stress of our approach prior to

bird capture and handling. Indeed, changes in T_{eye} and T_{beak} within 25 seconds before capture

have been observed in house sparrow (*Passer domesticus*) (Jerem and Romero, 2023) and our

pre-handling thermal pictures were taken on average 38 seconds after the first sign of alarm

behaviour. Yet, our results suggest that changes in surface temperature are probably more

likely mediated by the SAM axis rather than the HPA axis in adult king penguins. Future studies

testing for a direct acute stimulation of the HPA axis (using ACTH) or SAM axis (using

catecholamines) in the king penguin and other bird species could be useful to gain greater

insight on the contribution of the HPA and SAM axes in mediating changes in body surface

temperature (Jerem and Romero, 2023; Ouyang et al., 2021).

4.2. Stress-induced changes in surface temperatures

In breeding adults, we observed a mild sex-specific significant increase of T_{eye} and a

significant decrease of T_{beak} in response to acute capture-handling stress. Differences in

temperature change between body regions in response to acute stress are not rare (see Table

1). For instance, (Moe et al., 2017) measured a decrease of T_{eye} contrasting with an increase

of T_{feet} in chicken, whereas in captive budgerigars (*Melopsittacus undulatus*) T_{eye} increased

transiently after exposure to a stressor while T_{feet} did not change (Ikkatai and Watanabe, 2015;

Table 1). Anatomical and functional reasons can likely explain the difference of stress-induced

changes in surface temperature between body regions in penguins, with for instance a

potential interest in maintaining or increasing eye blood flow to maintain or enhance visual
acuity when acutely threatened by an environmental challenge.

T_{eye} has also been shown to positively correlate with internal (cloacal) temperature in
various avian species (budgerigars: Ikkatai and Watanabe, 2015; chicken: Cândido et al.,
2020). Consequently, it is possible that the increase in T_{eye} we observe in adults could reflect
an increase in internal body temperature (Oka, 2018). Consistently, in closely related Emperor
penguins (*Aptenodytes forsteri*), acute stress leads to an approximate 1.5°C increase in
internal (stomach) temperature (Regel and Pütz, 1997). Such an increase in T_{eye} in response to
handling stress has been reported two times in avian species to the best of our knowledge
(Ikkatai and Watanabe, 2015; Jakubas et al., 2022), while most previous studies have shown
either a significant decrease (4 studies) or no significant change (4 studies; Table 1). Those
contrasted findings might be explained by at least three factors: 1. variation in body size, since
smaller individuals/species are expected to favour internal heat conservation by reducing
more markedly peripheral blood flow and surface temperature compared to large ones
(*thermoprotective hypothesis*; Robertson et al., 2020a); 2. differences in the thermal
environment, since it has been shown that during acute stress response, heat conservation is
favoured below the thermoneutral zone while heat dissipation is favoured above its upper
limit (Robertson et al., 2020a); 3. the various delays at which T_{eye} was measured in response
to acute stress, since for instance the increase in T_{eye} found by Ikkatai and Watanabe (2015)
was short-lived (visible 5 min after the stressor, but not later on during the 30 min
measurement period). It is important to note that T_{eye} and T_{beak} can show very rapid and non-
linear changes in response to acute stress, that are likely to be missed by most studies not
using continuous thermal image recording (Jerem et al., 2019; Jerem and Romero, 2023;
Zuluaga and Danner, 2023). For instance, Zuluaga et al. (2023) found an initial drop in T_{beak}
following exposure to stressor, with a recovery to baseline levels after only 2 minutes, while
Jerem et al. 2019 found an initial drop in T_{eye} lasting less than 15s, followed by an increase
above baseline from 15 to 75 seconds.

Although the interaction between handling stress and sex was only marginally
significant (p = 0.060), it appeared that only females exhibited a noticeable increase in T_{eye} in
response to handling stress in our study. Differences in body size or mass between sexes are
very modest (*ca.* 5%) in king penguin (Kriesell et al., 2018), which thus seems unlikely to fully
explain the sex-difference observed here. There are known sex-differences in stress

physiology (Handa and McGivern, 2017), but sex differences in surface temperature changes

induced by acute stress have been rarely investigated (but see Robertson et al. (2020a) for a

result opposite to ours between sexes). We previously observed no sex effect in HPA axis

responsiveness between males and females king penguins (Viblanc et al., 2016), but as

mentioned above, the SAM axis is the likely driver of acute changes in surface temperatures.

Consequently, this suggests that sexes might differ in the stress-sensitivity of their SAM axis,

although this would need to be confirmed by direct measurements of SAM axis activity.

The decrease we observe in T_{beak} in response to acute stress in adults appears like the

typical peripheral vasoconstriction response previously reported for instance by Tabh et al.

(2021). Such peripheral vasoconstriction enables the redistribution of the peripheral blood

circulation to internal organs and tissues, such as the brain or muscles, favouring their

oxygenation and nutrition to sustain the fight-or-flight response. King penguins breed in a

highly dense and aggressive colonial environment (up to 500 aggressive interactions per hour;

(CôTé, 2000). Aggressive interactions, frequently leading to injuries, are known to result in

increased heart rates (Viblanc et al., 2012). A decrease of T_{beak} during acute stress response

likely reflects peripheral vasoconstriction, that is likely widespread to other peripheral body

parts, which according to the *haemoprotective hypothesis* could help to reduce blood loss in

case of injury (Robertson et al., 2020a).

The changes observed in surface temperatures in chicks differ from those found in

adults. Indeed, we observed a significant decrease in chick's T_{eye}, whereas T_{eye} increased in

adult females and stayed stable in adult males. While T_{beak} significantly decreased as observed

in adults, the decrease was more pronounced in chicks (-3.3 ± 0.6°C *vs.* -1.5 ± 0.4°C, in chicks

and adults, respectively). This is likely explained by greater peripheral thermal losses in chicks

suddenly exposed to an ambient temperature drop (*ca.* 35°C under the brood pouch of the

adults *vs.* 15°C when taken out for measurement and handling), despite their ability to

maintain their internal temperature at this stage at ambient temperature (Duchamp et al.,

2002). The more pronounced drop in surface temperature we observed in chicks during acute

handling could reflect their greater need to conserve heat at a relatively mild T_a compared to

adults.

4.3. Impact of parental CORT on egg and chick surface temperature

Contrary to our prediction, we observed no significant impact of parental CORT

treatment on egg temperature, suggesting little alteration of incubation behaviour by

increased CORT levels, contrary to a previous study in Adelie penguin reporting a reduction of

$1.3 \pm 0.2°C$ in T_{egg} (Thierry et al., 2013a). It is possible that incubating two eggs in the colder

environment of Adélie penguins is more challenging (and thus more likely to be impacted by

high CORT levels) than incubating a single egg under the milder climate experienced by king

penguins. Alternatively, it is also possible that king penguins are more resilient to increased

CORT levels than Adelie penguins, which is supported by the good reproductive success of

king penguins implanted with CORT (X et *al.* unpublished), contrary to what has been observed

in Adelie penguins (Thierry et al., 2013b). Measurements of incubation quality using a dummy

egg with temperature and rotation sensors (Thierry et al., 2013a) may provide more accurate

data on this question, but it is important to note that the T_{egg} measured in this study (Placebo:

34.9 ± 0.3 °C) was close to the T_{egg} measured in the same penguin colony using internal sensors

in dummy eggs ($35.7 \pm 0.4°C$; (Groscolas et al., 2000)). Our results therefore suggest that

further studies may benefit from the use of minimally invasive thermal imaging to measure

incubation quality, for instance in the context of parental behaviour and climate change (Cook

et al., 2020).

The measure of chick surface temperatures revealed some surprising patterns. While

we expected CORT chicks to have lower surface temperatures due to poorer parental care

(less efficient brooding), we found some opposite result: chicks had higher surface

temperatures when brooded by the non-implanted partner of the CORT parent. This result

should be considered with caution considering the limited sample size available for chick

surface temperatures (N = 16). One possibility is that partners from CORT-implanted

individuals somehow perceived the 'stress' levels of their partner (as shown between siblings

in yellow legged gulls (*Larus michahellis*); Noguera et al., 2017) and somehow compensated

for parental care through more efficient brooding. This hypothesis requires further study to

determine whether the concept of 'family-transmitted stress' (Noguera et al., 2017) applies

to the king penguin. In this species, both parents must rely on each other over more than one

year to successfully fledge their single chick, which itself is entirely dependent on its parents

for its food supply throughout its growth. Hence, the king penguin could be promising system

to investigate the 'family-transmitted stress' hypothesis.

4.4. Conclusion

Evaluating stress levels and responses of free-living animals in a non-invasive manner

using thermal imaging is a burgeoning field of research (Jerem et al., 2015; Tabh et al., 2022).

However, little is still known about the underlying pathways of stress physiology that influence

the changes in surface temperature. Our experimental study in king penguin points toward a

likely preponderant role of the SAM axis. Additionally, by summarizing the current evidence

for acute changes in surface temperatures during stress exposure (Table 1), we highlight the

relative complexity and inconsistency between the effects observed by different studies.

Further experimental studies related to SAM and HPA axes involvement are thus required to

clarify what stress component(s) are measured through non-invasive thermal imaging.

5. Declarations

5.1. Author contribution

Study design: AS, VAV, PB and AL. Funding acquisition: JPR, PB, VAV, AS. Data collection in the

field: AS, SA, TH, LA, CG, and JPR. Data collection in the lab: ER and AS. Data analysis: CW, AL,

PB, AS. Writing original draft: AL, PB and AS. Writing review and editing: VAV, JPR, AN, SA, TH,

LA, ER and CG.

5.2 Data availability

The   datasets   used   in   this   manuscript   are   available   on   *FigShare*

(10.6084/m9.figshare.20134181).

5.3. Acknowledgments

We are grateful to the French Polar Institute (IPEV) and the Terres Australes et Antarctiques

Françaises for providing financial and logistical support for this study through the polar

program #119 (ECONERGY). This study is part of the long-term Studies in Ecology and

Evolution (SEE-Life) program of the CNRS. We wish to thank the Zone Atelier Antarctique et

Terres Australes (ZATA) from the CNRS for financial support, and the members of the Alfred

Faure field station for their help and support in the field. AL was supported by ISblue project,

Interdisciplinary graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a

grant from the French government under the program "Investissements d'Avenir" embedded

in France 2030. AS was supported by a 'Turku Collegium for Science and Medicine' Fellowship,

a Marie Sklodowska-Curie Postdoctoral Fellowships (#894963) and the IdEx Université de

Strasbourg (*HotPenguin*). SA and CG were funded by the French Polar Institute.

5.4. Competing interests

None

6. References

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67. https://doi.org/10.18637/jss.v067.i01

Brandl, H.B., Pruessner, J.C., Farine, D.R., 2022. The social transmission of stress in animal collectives. Proc Royal Soc B 289, 20212158. https://doi.org/10.1098/rspb.2021.2158

Cabanac, A.J., Guillemette, M., 2001. Temperature and heart rate as stress indicators of handled common eider. Physiol Behav 74, 475–479. https://doi.org/10.1016/s0031-9384(01)00586-8

Cândido, M.G.L., Tinôco, I.F.F., Albino, L.F.T., Freitas, L.C.S.R., Santos, T.C., Cecon, P.R., Gates, R.S., 2020. Effects of heat stress on pullet cloacal and body temperature. Poultry Sci 99, 2469–2477. https://doi.org/10.1016/j.psj.2019.11.062

Carere, C., Oers, K. van, 2004. Shy and bold great tits (Parus major): body temperature and breath rate in response to handling stress. Physiol Behav 82, 905–912. https://doi.org/10.1016/j.physbeh.2004.07.009

Cook, T.R., Martin, R., Roberts, J., Häkkinen, H., Botha, P., Meyer, C., Sparks, E., Underhill, L.G., Ryan, P.G., Sherley, R.B., 2020. Parenting in a warming world: thermoregulatory responses to heat stress in an endangered seabird. Conservation Physiology 8, 2283–13. https://doi.org/10.1093/conphys/coz109

CôTé, S.D., 2000. Aggressiveness in king penguins in relation to reproductive status and territory location. Anim Behav 59, 813–821. https://doi.org/10.1006/anbe.1999.1384

Deavers, D.R., Musacchia, X.J., 1979. The function of glucocorticoids in thermogenesis. Fed Proc 38, 2177–81.

Duchamp, C., Rouanet, J.-L., Barre, H., 2002. Ontogeny of thermoregulatory mechanisms in king penguin chicks (Aptenodytes patagonicus). Comparative Biochemistry and Physiology, Part A 131, 765–773. https://doi.org/10.1016/s1095-6433(02)00014-4

Edgar, J.L., Lowe, J.C., Paul, E.S., Nicol, C.J., 2011. Avian maternal response to chick distress. Proc Royal Soc B Biological Sci 278, 3129–3134. https://doi.org/10.1098/rspb.2010.2701

Edgar, J.L., Nicol, C.J., 2018. Socially-mediated arousal and contagion within domestic chick broods. Sci Rep-uk 8, 10509. https://doi.org/10.1038/s41598-018-28923-8

Edgar, J.L., Nicol, C.J., Pugh, C.A., Paul, E.S., 2013. Surface temperature changes in response to handling in domestic chickens. Physiol Behav 119, 195–200. https://doi.org/10.1016/j.physbeh.2013.06.020

Ellenberg, U., Setiawan, A.N., Cree, A., Houston, D.M., Seddon, P.J., 2007. Elevated hormonal stress response and reduced reproductive output in Yellow-eyed penguins exposed to unregulated tourism. Gen Comp Endocr 152, 54–63. https://doi.org/10.1016/j.ygcen.2007.02.022

Gauchet, L., Grémillet, D., 2022. Using facial infrared thermography to infer avian body temperatures in the wild. Marine Biology.

Giloh, M., Shinder, D., Yahav, S., 2012. Skin surface temperature of broiler chickens is correlated to
body core temperature and is indicative of their thermoregulatory status. Poultry Sci 91, 175–
188. https://doi.org/10.3382/ps.2011-01497

Groscolas, R., Decrock, F., Thil, M.-A., Fayolle, C., Boissery, C., Robin, J.-P., 2000. Refeeding signal in
fasting-incubating king penguins: changes in behavior and egg temperature. Am J Physiology-
regulatory Integr Comp Physiology 279, R2104–R2112.
https://doi.org/10.1152/ajpregu.2000.279.6.r2104

Halekoh, U., Højsgaard, S., 2014. A Kenward-Roger Approximation and Parametric Bootstrap
Methods for Tests in Linear Mixed Models - The R Package pbkrtest. J Stat Softw 59.
https://doi.org/10.18637/jss.v059.i09

Handa, R.J., McGivern, R.F., 2017. Reference Module in Neuroscience and Biobehavioral Psychology.
https://doi.org/10.1016/b978-0-12-809324-5.02865-0

Handrich, Y., Bevan, R.M., Charrassin, J.-B., Butler, P.J., Ptz, K., Woakes, A.J., Lage, J., Maho, Y.L.,
1997. Hypothermia in foraging king penguins. Nature 388, 64–67. https://doi.org/10.1038/40392

Herborn, K.A., Graves, J.L., Jerem, P., Evans, N.P., Nager, R., McCafferty, D.J., McKeegan, D.E.F., 2015.
Skin temperature reveals the intensity of acute stress. Physiol Behav 152, 225–230.
https://doi.org/10.1016/j.physbeh.2015.09.032

Herborn, K.A., Jerem, P., Nager, R.G., McKeegan, D.E.F., McCafferty, D.J., 2018. Surface temperature
elevated by chronic and intermittent stress. Physiol Behav 191, 47–55.
https://doi.org/10.1016/j.physbeh.2018.04.004

Ikkatai, Y., Watanabe, S., 2015. Eye surface temperature detects stress response in budgerigars
(Melopsittacus undulatus). Neuroreport 26, 642–646.
https://doi.org/10.1097/wnr.0000000000000403

Jakubas, D., Wojczulanis-Jakubas, K., Grissot, A., Devogel, M., Cendrowska, M., Chastel, O., 2022. Eye
Region Surface Temperature and Corticosterone Response to Acute Stress in a High-Arctic
Seabird, the Little Auk. Animals Open Access J Mdpi 12, 499.
https://doi.org/10.3390/ani12040499

Jerem, P., Herborn, K., McCafferty, D., McKeegan, D., Nager, R., 2015. Thermal Imaging to Study
Stress Non-invasively in Unrestrained Birds. J Vis Exp Jove 53184. https://doi.org/10.3791/53184

Jerem, P., Jenni-Eiermann, S., Herborn, K., McKeegan, D., McCafferty, D.J., Nager, R.G., 2018. Eye
region surface temperature reflects both energy reserves and circulating glucocorticoids in a wild
bird. Sci Rep-uk 8, 1907. https://doi.org/10.1038/s41598-018-20240-4

Jerem, P., Jenni-Eiermann, S., McKeegan, D., McCafferty, D.J., Nager, R.G., 2019. Eye region surface
temperature dynamics during acute stress relate to baseline glucocorticoids independently of
environmental conditions. Physiology & Behavior 112627.
https://doi.org/10.1016/j.physbeh.2019.112627

Jerem, P., Romero, L.M., 2023. It's cool to be stressed: Body surface temperatures track sympathetic
nervous system activation during acute stress. J. Exp. Biol. 226, jeb246552.
https://doi.org/10.1242/jeb.246552

Knoch, S., Whiteside, M.A., Madden, J.R., Rose, P.E., Fawcett, T.W., 2022. Hot-headed peckers:
thermographic changes during aggression among juvenile pheasants (Phasianus colchicus).
Philosophical Transactions Royal Soc B 377, 20200442. https://doi.org/10.1098/rstb.2020.0442

Kriesell, H.J., Aubin, T., Planas-Bielsa, V., Benoiste, M., Bonadonna, F., Gachot-Neveu, H., Maho, Y.L.,
Schull, Q., Vallas, B., Zahn, S., Bohec, C.L., 2018. Sex identification in King Penguins Aptenodytes
patagonicus through morphological and acoustic cues. Ibis 160, 755–768.
https://doi.org/10.1111/ibi.12577

Lenth, R., Singmann, H., Love, J., Buerkner, P., Herve, M., 2018. Emmeans: Estimated marginal
means, aka least-squares means. R package version ...
https://doi.org/10.1080/00031305.1980.10483031>;

Lewden, A., Bonnet, B., Nord, A., 2020. The metabolic cost of subcutaneous and abdominal
rewarming in king penguins after long-term immersion in cold water. J Therm Biol 91, 102638.
https://doi.org/10.1016/j.jtherbio.2020.102638

Lewden, Agnès, Enstipp, M.R., Bonnet, B., Bost, C., Georges, J.-Y., Handrich, Y., 2017a. Thermal
strategies of king penguins during prolonged fasting in water. J Exp Biol 220, 4600–4611.
https://doi.org/10.1242/jeb.168807

Lewden, Agnes, Enstipp, M.R., Picard, B., Walsum, T. van, Handrich, Y., 2017b. High peripheral
temperatures in king penguins while resting at sea: thermoregulation versus fat deposition.
Journal Of Experimental Biology 220, 3084–3094. https://doi.org/10.1242/jeb.158980

MacDougall-Shackleton, S.A., Bonier, F., Romero, L.M., Moore, I.T., 2019. Glucocorticoids and
"Stress" Are Not Synonymous. Integr Org Biology 1, obz017. https://doi.org/10.1093/iob/obz017

Moe, R.O., Bohlin, J., Flø, A., Vasdal, G., Stubsjøen, S.M., 2017. Hot chicks, cold feet. Physiol Behav
179, 42–48. https://doi.org/10.1016/j.physbeh.2017.05.025

Narushin, V.G., 2005. Egg geometry calculation using the measurements of length and breadth.
Poultry Sci 84, 482–484. https://doi.org/10.1093/ps/84.3.482

Noguera, J.C., Kim, S.-Y., Velando, A., 2017. Family-transmitted stress in a wild bird. Proceedings of
the National Academy of Sciences 27, 201706164. https://doi.org/10.1086/680688

Oka, T., 2018. Chapter 35 Stress-induced hyperthermia and hypothermia. Handb Clin Neurology 157,
599–621. https://doi.org/10.1016/b978-0-444-64074-1.00035-5

Ouyang, J.Q., Macaballug, P., Chen, H., Hodach, K., Tang, S., Francis, J.S., 2021. Infrared
thermography is an effective, noninvasive measure of HPA activation. Ann Ny Acad Sci 24, 1–6.
https://doi.org/10.1080/10253890.2020.1868431

Pijpers, N., Heuvel, H. van den, Duncan, I.H., Yorzinski, J., Neethirajan, S., 2022. Understanding
Chicks' Emotions: Are Eye Blinks & Facial Temperatures Reliable Indicators? Biorxiv
2022.01.31.478468. https://doi.org/10.1101/2022.01.31.478468

Playà-Montmany, N., Tattersall, G.J., 2021. Spot size, distance and emissivity errors in field
applications of infrared thermography. Methods Ecol Evol 12, 828–840.
https://doi.org/10.1111/2041-210x.13563

Regel, J., Pütz, K., 1997. Effect of human disturbance on body temperature and energy expenditure in
penguins. Polar Biol 18, 246–253. https://doi.org/10.1007/s003000050185

Robertson, J.K., Mastromonaco, G., Burness, G., 2020a. Evidence that stress-induced changes in
surface temperature serve a thermoregulatory function. J Exp Biol 223, jeb213421.
https://doi.org/10.1242/jeb.213421

Robertson, J.K., Mastromonaco, G.F., Burness, G., 2020b. Social hierarchy reveals thermoregulatory
trade-offs in response to repeated stressors. J Exp Biol 223, jeb229047.
https://doi.org/10.1242/jeb.229047

Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research. Trends Ecol
Evol 19, 249–255. https://doi.org/10.1016/j.tree.2004.03.008

Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How Do Glucocorticoids Influence Stress
Responses? Integrating Permissive, Suppressive, Stimulatory, and Preparative Actions. Endocr Rev
21, 55–89. https://doi.org/10.1210/edrv.21.1.0389

Smulders, T.V., 2021. Telencephalic regulation of the HPA axis in birds. Neurobiol. Stress 15, 100351.
https://doi.org/10.1016/j.ynstr.2021.100351

Stier, A., Schull, Q., Bize, P., Lefol, E., Haussmann, M., Roussel, D., Robin, J.-P., Viblanc, V.A., 2019.
Oxidative stress and mitochondrial responses to stress exposure suggest that king penguins are
naturally equipped to resist stress. Scientific Reports 9, 8545. https://doi.org/10.1038/s41598-
019-44990-x

Tabh, J.K.R., Burness, G., Wearing, O.H., Tattersall, G.J., Mastromonaco, G.F., 2021. Infrared
thermography as a technique to measure physiological stress in birds: Body region and image
angle matter. Physiological Reports 9, e14865. https://doi.org/10.14814/phy2.14865

Tabh, J.K.R., Mastromonaco, G.F., Burness, G., 2022. Stress-induced changes in body surface
temperature are repeatable, but do not differ between urban and rural birds. Oecologia 1–15.
https://doi.org/10.1007/s00442-022-05120-z

Tattersall, G.J., Andrade, D.V., Abe, A.S., 2009. Heat Exchange from the Toucan Bill Reveals a
Controllable Vascular Thermal Radiator. Science 325, 468–470.
https://doi.org/10.1126/science.1175553

Thierry, A.-M., Massemin, S., Handrich, Y., Raclot, T., 2013a. Elevated corticosterone levels and
severe weather conditions decrease parental investment of incubating Adélie penguins.
Hormones And Behavior 63, 475–483. https://doi.org/10.1016/j.yhbeh.2012.12.011

Thierry, A.-M., Ropert-Coudert, Y., Raclot, T., 2013b. Elevated corticosterone levels decrease
reproductive output of chick-rearing Adélie penguins but do not affect chick mass at fledging.
Conservation Physiology 1, cot007. https://doi.org/10.1093/conphys/cot007

Thomas, D.B., Ksepka, D.T., Fordyce, R.E., 2011. Penguin heat-retention structures evolved in a
greenhouse Earth. Biol Letters 7, 461–464. https://doi.org/10.1098/rsbl.2010.0993

Torres-Medina, F., Cabezas, S., Marchant, T.A., Wikelski, M., Romero, L.M., Hau, M., Carrete, M.,
Tella, J.L., Blas, J., 2018. Corticosterone implants make stress hyporesponsive birds. The Journal of
Experimental Biology jeb.173864-43. https://doi.org/10.1242/jeb.173864

Viblanc, V.A., Dobson, F.S., Stier, A., Schull, Q., Saraux, C., Gineste, B., Pardonnet, S., Kauffmann, M.,
Robin, J.-P., Bize, P., 2016. Mutually honest? Physiological "qualities' signalled by colour
ornaments in monomorphic king penguins. Biological Journal of the Linnean Society 118, 200–
214. https://doi.org/10.1111/bij.12729

Viblanc, V.A., Schull, Q., Cornioley, T., Stier, A., Ménard, J.-J., Groscolas, R., Robin, J.-P., 2018. An
integrative appraisal of the hormonal and metabolic changes induced by acute stress using king
penguins as a model. Gen Comp Endocr 269, 1–10. https://doi.org/10.1016/j.ygcen.2017.08.024

Viblanc, V.A., Valette, V., Kauffmann, M., Malosse, N., Groscolas, R., 2012. Coping with social stress:
heart rate responses to agonistic interactions in king penguins. Behavioral Ecology 23, 1178–
1185. https://doi.org/10.1093/beheco/ars095

Weimer, S.L., Wideman, R.F., Scanes, C.G., Mauromoustakos, A., Christensen, K.D., Vizzier-Thaxton,
Y., 2020. Broiler stress responses to light intensity, flooring type, and leg weakness as assessed by
heterophil-to-lymphocyte ratios, serum corticosterone, infrared thermography, and latency to lie.
Poultry Sci 99, 3301–3311. https://doi.org/10.1016/j.psj.2020.03.028

Weimerskirch, H., Stahl, J.C., Jouventin, P., 1992. The breeding biology and population dynamics of
king penguins Aptenodytes patagonica on the Crozet Islands. Ibis 134, 107–117.

Winder, L.A., White, S.A., Nord, A., Helm, B., McCafferty, D.J., 2020. Body surface temperature
responses to food restriction in wild and captive great tits. Journal Of Experimental Biology 223,
jeb220046-8. https://doi.org/10.1242/jeb.220046

Wingfield, J.C., Romero, L.M., 2015. Tempests, Poxes, Predators, and People 149–196.
https://doi.org/10.1093/acprof:oso/9780195366693.003.0005

Zuluaga, J.D., Danner, R.M., 2023. Acute stress and restricted diet reduce bill-mediated heat
dissipation in the song sparrow (Melospiza melodia): implications for optimal thermoregulation. J
Exp Biol 226. https://doi.org/10.1242/jeb.245316

Figures

Fig. 1: **Baseline and stress-induced plasma corticosterone levels of breeding king penguins** implanted either with a Placebo, or a CORT subcutaneous implant (50 mg, 90 days release; n = 80 observations, N = 40 individuals) during incubation. Birds were blood sampled 6 days after implantation. CORT (red) *vs.* Placebo (blue) individual responses (raw data) are presented along with group and time specific mean ± SE. CORT levels were influenced by the interaction between handling stress (baseline: < 4 min vs. stress-induced: 30 min of handling stress) and CORT treatment (LMM on log-transformed plasma CORT: $F_{1,37.2}$ = 62.4, p < 0.001). Letters indicate significant differences according to post-hoc tests with Tukey adjustments (p < 0.030).

Fig. 2: **Infrared images of (A) breeding adult, (B) incubated egg and (C) 20 days-old chick**

752
753
Fig. 3: **Adult king penguin surface temperature responses to an experimental manipulation**
of glucocorticoid levels (CORT *vs.* Placebo subcutaneous implants) and handling stress: (A)
Maximum eye temperature response to handling stress, (B) Minimum beak temperature
response to handling stress, and (C) Sex-specific response of eye temperature to handling
stress. In panels A and B, CORT (red) *vs.* Placebo (blue) individual responses (raw data) are
presented. Since CORT Treatment had no significant effect either alone or in interaction with
Handling stress (see Table 2 for details), the overall mean ± SE is shown in black. Least-square
means ± SE from final statistical models (Table 2; n = 138 observations; N = 49 individuals) are
presented. Different letters indicate significant differences in panels A and B, and ** in panel
C represent the significant increase of T_{eye} observed only in females.
764

Fig. 4: **King penguin chick surface temperature in response to parental glucocorticoid manipulation and handling: (A) Maximum eye temperature according to parental treatment and category of the parent brooding the chick (*i.e.* partners are not implanted), (B) Minimum beak temperature according to parental treatment and category of the parent brooding the chick, (C) Maximum eye temperature response to handling, and (D) Minimum eye temperature response to handling.** In panels C and D, CORT (red) vs. Placebo (blue) chick's individual responses (raw data) are presented, but only the global mean ± SE are presented since there was no significant interaction between parental CORT Treatment and Handling (see Table 3 for details). Least-square means ± SE from final statistical models (Table 3) are presented (n = 63 observations, N = 16 chicks). Different letters indicate significant differences.

Tables

Table 1: Summary table of avian studies testing the impact of acute stress exposure on

changes in body surface temperatures measured with thermal imaging (studies related to

heat and food stress are not included). *Area* corresponds to the body region being measured,

response to the direction of the temperature response (=: no significant change, ↓: significant

decrease and ↑: significant increase, *Δ T°C* to the amplitude of the temperature response,

stressor the nature and duration of the experimental stressor. Some studies measured

temperature responses at different time points, and when the effects of stress exposure

change through time, the responses are shown with consecutive symbols (*e.g.* =, ↑: initial

absence of change followed later on by an increase in surface temperature). Please note that

the area of interest for T_{eye} differs between studies (periorbital region *vs.* eyeball region) and

that the surface temperature being measured can differ in nature between studies (*i.e.*

maximum, average or minimum temperature of the area of interest).

795
796

Species	Condition	Age	Stressor	Area	Response	Δ T°C	Ref.
Blue tit *Cyaniste caerulescens*	Wild	Adult	Trap closure + handling (3min)	Eye	↓, ↑,=	-1.0, +1.0°C, =	(Jerem et al., 2019)
Budgerigars *Melopsittacus undulatus*	Captive	Adult	Handling (30 min)	Eye / Legs	↑, = / =	+0.7°C, = / =	(Ikkatai and Watanabe, 2015)
Chicken *Gallus gallus domesticus*	Captive	Adult	Air puff (10 min)	Eye	↓	-1.0°C	(Edgar et al., 2011)
Chicken *Gallus gallus domesticus*	Captive	Adult	Handling (20 min)	Eye / Head / Comb	↓, = / =, ↑ / ↓, =	-0.5°C, = / =, +1.0°C / -1.5°C, =	(Edgar et al., 2013)
Chicken *Gallus gallus domesticus*	Captive	Chick (9 weeks)	Air puff (10 min)	Eye	↓	-1.0°C	(Edgar and Nicol, 2018)
Chicken *Gallus gallus domesticus*	Captive	Adult	Handling: mild/severe (20 min)	Eye / Face / Wattle / Comb	↓ / = / ↓ / ↓	-0.4°C / = / -0.7 / -1.3°C / -0.5 / -2.2°C	(Herborn et al., 2015)
Chicken *Gallus gallus domesticus*	Captive	Adult	Enrichment removal (within 2 h)	Eye / Face / Comb	↓ / ↓ / ↓	-0.8°C / -0.7°C / -3.5°C	(Herborn et al., 2018)
Chicken *Gallus gallus domesticus*	Captive	Chick (30 days)	Handling (10 min)	Head / Feet	↑ / ↓	+0.76°C / -0.45°C	(Moe et al., 2017)
Chicken *Gallus gallus domesticus*	Captive	Chick (9 days)	Visual (45 min)	Eye / Beak	= / =, ↑	= / +0.5°C	(Pijpers et al., 2022)
Domestic pigeon *Columbia livia domestica*	Captive	Adult	Handling (3.5 min)	Eye / Beak	= / ↓	-0.4°C / -2.1°C	(Tabh et al., 2021)
House sparrow *Passer domesticus*	Captive	Adult	Handling and injection (~1min)	Eye / Beak	↓ / ↓	-1.0°C / -4.0°C	(Jerem and Romero, 2023)
King penguin *Aptenodytes patagonicus*	Wild	Adult	Handling (5-10 min)	Eye / Beak	↑ / ↓	+0.8°C / -1.5°C	This study
	Wild	Chick (20 days*)	Handling (5-10 min)	Eye / Beak	↓ / ↓	-0.9°C / -3.3°C	
Little Auk *Alle alle*	Wild	Adult	Handling (30 min)	Eye	↑	+1,7°C	(Jakubas et al., 2022)
Pheasant *Phasianus colchicus*	Captive	Juvenile (42 to 49 days)	Agression (20 s)	Head	↑	+0.6°C	(Knoch et al., 2022)
Song sparrow *Melospiza melodia*	Captive	Adult	Handling (13 min)	Eye / Beak	↓, = / ↓, ↑, =	-1.5°C, = / -2.6°C, +2.6°C, =	(Zuluaga and Danner, 2023)

* king penguins chicks are not thermally emancipated at this age

 Table 2: Summary of linear mixed models (LMMs) investigating the determinants of (A)

 maximum eye temperature (T_{eye}) and (B) minimum beak temperature (T_{beak}) in adult king

 penguins.

A. T_{eye} (n = 138 observations; N = 49 individuals)

Random effects:		Variance			
Bird ID	Intercept	5.04			
Residual		3.21			
Fixed effects:	**Estimate**	**SE**	**t**	**df**	**p**
Intercept	24.94	1.71	14.54	47.9	<0.001
Treatment (CORT)	-0.10	0.72	-0.15	45.8	0.89
Handling (after)	**1.45**	**0.47**	**3.04**	**89.3**	**0.003**
Sex (Male)	0.76	0.78	0.97	63.7	0.34
T_a	**0.29**	**0.13**	**2.33**	**45.0**	**0.024**
*Handling*Sex*	*-1.19*	*0.63*	*-1.90*	*89.7*	*0.060*
(Treatment x Handling)					(0.44)

B. T_{beak} (n = 138 observations; N = 49 individuals)

Random effects:		Variance			
Bird ID	Intercept	15.91			
Residual		6.40			
Fixed effects:	**Estimate**	**SE**	**t**	**df**	**p**
Intercept	13.84	2.92	4.74	45.0	<0.001
Treatment (CORT)	-1.13	1.24	-0.91	44.5	0.37
Handling (after)	**-1.47**	**0.45**	**-3.29**	**84.6**	**0.001**
Sex (Male)	0.90	1.23	0.73	44.6	0.47
T_a	**0.54**	**0.22**	**2.45**	**43.8**	**0.018**
(Treatment x Handling)					(0.61)

Table 3: Summary of linear mixed models (LMMs) investigating the determinants of (A) maximum eye temperature (T_{eye}) and (B) minimum beak temperature (T_{beak}) in king penguin chicks. Only one parent was implanted with CORT or Placebo implant, thus we considered both parental treatment (CORT *vs.* Placebo), the category of the parent brooding the chick at the time of measurement (implanted parent *vs.* non-implanted partner) and their interaction in our analyses.

A. T_{eye} (n = 63 observations; N = 16 chicks)

Random effects:		Variance			
Bird ID	Intercept	0.81			
Residual		1.18			
Fixed effects:	**Estimate**	**SE**	**t**	**df**	**p**
Intercept	29.47	1.39	21.26	11.1	<0.001
Treatment (CORT)	-0.88	0.68	-1.29	11.3	0.22
Parent (implanted)	**-2.84**	**0.82**	**3.48**	**11.1**	**0.005**
Handling (after)	**-0.85**	**0.28**	**-3.08**	**47.2**	**0.003**
T_a	**0.21**	**0.09**	**2.41**	**10.9**	**0.035**
Treatment*Parent	**2.78**	**1.10**	**2.53**	**11.0**	**0.028**
(Treatment x Handling)					(0.79)

B. T_{beak} (n = 63 observations; N = 16 chicks)

Random effects:		Variance			
Bird ID	Intercept	5.53			
Residual		4.89			
Fixed effects:	**Estimate**	**SE**	**t**	**df**	**p**
Intercept	13.55	3.42	3.96	11.1	0.002
Treatment (CORT)	1.15	1.68	0.69	11.3	0.51
Parent (implanted)	**-6.82**	**2.02**	**-3.38**	**11.1**	**0.006**
Handling (after)	**-3.26**	**0.56**	**-5.77**	**46.9**	**<0.001**
T_a	**0.57**	**0.21**	**2.65**	**11.0**	**0.023**
Treatment*Parent	**6.37**	**2.72**	**2.34**	**11.0**	**0.039**
(Treatment x Handling)					(0.94)