A novel Na$_v$1.5-dependent feedback mechanism driving glycolytic acidification in breast cancer metastasis

Theresa K Leslie[1,2], Aurelien Tripp[3], Andrew D James[1,2], Scott P Fraser[4], Michaela Nelson[1,2], Michael Toss[5], Wakkas Fadhil[5], Samantha C Salvage[8], Mar Arias Garcia[3], Melina Beykou[3,10], Emad Rakha[5], Valerie Speirs[6], Chris Bakal[3], George Poulogiannis[3], Mustafa B A Djamgoz[4,7], Antony P Jackson[8], Hugh R Matthews[8], Christopher L-H Huang[8,9], Andrew N Holding[1,2], Sangeeta Chawla[1,2], and William J Brackenbury[1,2*]

[1] Department of Biology, University of York, York, UK

[2] York Biomedical Research Institute, University of York, York, UK

[3] Division of Cancer Biology, Institute of Cancer Research, London, UK

[4] Department of Life Sciences, Imperial College London, London SW7 2AZ, UK

[5] Department of Pathology, School of Medicine, University of Nottingham, NG5 1PB, UK

[6] Institute of Medical Sciences, University of Aberdeen, Aberdeen, AB24 2ZD, UK

[7] Biotechnology Research Centre, Cyprus International University, Haspolat, TRNC, Mersin 10, Türkiye

[8] Department of Biochemistry, University of Cambridge, Cambridge, UK

[9] Physiological Laboratory, University of Cambridge, Cambridge, UK

[10] Department of Electrical and Electronic Engineering, Imperial College London, London, SW7 2AZ, UK

*Corresponding author: Dr William J. Brackenbury, Department of Biology and York Biomedical Research Institute, University of York, Wentworth Way, Heslington, York YO10 5DD, UK. Email: william.brackenbury@york.ac.uk Tel: +44 1904 328284.

Abstract

Solid tumours have abnormally high intracellular $[Na^+]$. The activity of various Na^+ transporting proteins including channels may underlie this Na^+ accumulation. Here, we show that voltage-gated Na^+ channels (VGSCs) are functionally active in a subset of breast cancer cell lines, cancer-associated fibroblasts, xenograft tumours and metastases. Downregulation of the $Na_v1.5$ VGSC in xenograft breast tumours suppresses expression of invasion-regulating genes, consistent with previous studies showing that $Na_v1.5$ promotes invasion in cancer cells. We also show that $Na_v1.5$ activity increases glycolysis, promoting extracellular acidification that would facilitate this invasion. In a reciprocal interaction, acidic extracellular pH elevates persistent Na^+ influx through $Na_v1.5$ in breast cancer cells. Using a mathematical model, we show that $Na_v1.5$ activity can sustain production of extracellular H^+. We show that likely VGSC currents are detectable in patient-derived breast tumour cells and tissues. Furthermore, protein expression of $Na_v1.5$ strongly correlates with increased metastasis and shortened cancer-specific survival in breast cancer patients. Together, these findings show positive feedback between extracellular acidification and movement of Na^+ into cancer cells which can facilitate invasion. They also highlight the clinical significance of $Na_v1.5$ as a potentiator of breast cancer metastasis and provide further evidence supporting the use of VGSC inhibitors in cancer treatment.

Keywords: Breast cancer / Electrophysiology / Glycolysis / Invasion / Metastasis / $Na_v1.5$ / Sodium / Na^+ channel / voltage-gated sodium channel

Introduction

Breast cancer is the leading cause of cancer-related deaths in women worldwide (Sung *et al.*, 2021) and most deaths are due to metastatic disease resulting from poor treatment options and therapy resistance (Anderson *et al.*, 2019). Around 20-30% of patients with primary breast cancer will go on to develop distant metastasis and once this has been diagnosed, the average survival time is 2-3 years despite treatment (Cardoso *et al.*, 2012). Thus, there is a great need for improved treatments to prevent or reduce breast cancer metastasis.

Increasing evidence points to ion channels as key regulators of cancer progression (Fraser & Pardo, 2008; Elingaard-Larsen *et al.*, 2022; Rahrmann *et al.*, 2022; Folcher *et al.*, 2023). Members of the voltage-gated Na^+ channel (VGSC) family are upregulated in multiple cancer types (Leslie & Brackenbury, 2022). In solid cancers, including breast, prostate, lung, and colon cancer, VGSC activity promotes cellular invasion (Djamgoz *et al.*, 2019; Lopez-Charcas *et al.*, 2021). In breast cancer, the $Na_v1.5$ subtype is upregulated at the mRNA level compared to normal tissue and is associated with recurrence and metastasis (Yang *et al.*, 2012). $Na_v1.5$ is also upregulated in breast cancers at the protein level (Fraser *et al.*, 2005; Nelson *et al.*, 2015b), predominantly in its neonatal D1:S3 splice form (Yamaci *et al.*, 2017); however, the sample sizes of these studies were too small to reliably determine the relationship between $Na_v1.5$ expression and clinical outcome. $Na_v1.5$ activity, as measured by the presence of Na^+ currents, has not yet been investigated in breast cancer tissue or near-patient primary cells. Nonetheless, Na^+ currents carried by $Na_v1.5$ have been detected in a small number of breast cancer cell lines and in tissue slices from murine tumour xenografts (Roger *et al.*, 2003; Fraser *et al.*, 2005; Nelson *et al.*, 2015b; Fraser *et al.*, 2016). In these cells, the persistent Na^+ current (as distinct from the transient, inactivating Na^+ current), which passes through the

channels at the resting membrane potential (V_m), has been shown to potentiate cellular invasion *in vitro* and tumour growth and metastasis *in vivo* (Roger *et al.*, 2003; Fraser *et al.*, 2005; Brackenbury *et al.*, 2007; Yang *et al.*, 2012; Driffort *et al.*, 2014; Nelson *et al.*, 2015b). Importantly, the metastasis-promoting function of $Na_v1.5$ can be inhibited in preclinical models using VGSC blockers, including phenytoin and ranolazine, suggesting that $Na_v1.5$ may represent a novel anti-metastatic target for therapeutic intervention (Driffort *et al.*, 2014; Nelson *et al.*, 2015a). Furthermore, perioperative administration of the VGSC blocker lidocaine has recently been shown to significantly improve disease-free survival in women with early breast cancer (Badwe *et al.*, 2023).

The mechanism by which VGSCs increase invasion of cancer cells is not yet fully understood (Lopez-Charcas *et al.*, 2021). However, it has been shown that Na^+ influx via $Na_v1.5$ leads to extracellular acidification via the Na^+/H^+ exchanger, NHE1, activating pH-dependent cathepsins and promoting invasion (Gillet *et al.*, 2009; Brisson *et al.*, 2013; Driffort *et al.*, 2014). Because an increase in cytosolic $[Na^+]$ reduces the Na^+ electrochemical gradient powering H^+ extrusion via NHE1, an allosteric interaction between $Na_v1.5$ and NHE1 has been proposed to explain the $Na_v1.5$-dependent increase in H^+ extrusion by NHE1 (Brisson *et al.*, 2013). An additional possibility is that Na^+ influx through $Na_v1.5$, rather than the $Na_v1.5$ protein itself, is responsible indirectly for increasing H^+ extrusion through NHE1 and/or other pH regulators. Extracellular acidification may additionally positively regulate $Na_v1.5$ function, given that low pH increases the persistent Na^+ current carried by $Na_v1.5$ in cardiomyocytes (Khan *et al.*, 2006; Jones *et al.*, 2011). However, the link between $Na_v1.5$ function, Na^+ homeostasis and pH regulation in breast cancer cells has not been fully delineated.

A second consequence of $Na_v1.5$-mediated Na^+ influx is steady state depolarisation of the V_m, which activates the small GTPase Rac1 and increases migration of breast cancer cells (Yang *et al.*, 2020). Via V_m depolarisation, $Na_v1.5$ has also been shown to upregulate the PKA/ERK/c-JUN/ELK-1/ETS-1 transcriptional pathway increasing expression of a network of invasion-related genes in colorectal carcinoma cells (House *et al.*, 2010; House *et al.*, 2015). $Na_v1.5$ has also been shown to positively regulate epithelial-mesenchymal transition (EMT)-promoting genes in breast cancer cells (Gradek *et al.*, 2019). Together, these studies point towards a possible function of $Na_v1.5$ in transcriptional regulation of a range of invasion-promoting genes.

In the present study, we set out to delineate the relationship between $Na_v1.5$, Na^+ flux, pH regulation and breast cancer progression. We also explored VGSC activity in breast cancer tissue and primary cultures of breast cancer cells and we sought to evaluate protein expression of $Na_v1.5$ in a large cohort of patient breast cancer samples. We show that suppression of $Na_v1.5$ expression using shRNA leads to downregulation of multiple invasion-regulating genes in xenograft tumours. Furthermore, persistent Na^+ current is elevated in breast cancer cells upon exposure to low pH and $Na_v1.5$ activity increases glycolysis. We also show, for the first time, that inward Na^+ currents can be recorded from primary breast carcinoma cell samples from patients. In addition, high $Na_v1.5$ protein expression correlates with reduced cancer-specific survival and increased risk of metastasis. Together, these findings highlight the clinical significance of $Na_v1.5$ as a potentiator of breast cancer metastasis.

Materials and Methods

Ethical approvals

All animal procedures were carried out after approval by the University of York Animal Welfare and Ethical Review Body and under authority of a UK Home Office Project Licence and associated Personal Licences. Human tissue experiments were conducted in accordance with the ethical standards of the Declaration of Helsinki and according to national and international guidelines and were approved by the University of York Ethical Review Process.

Breast cancer cell lines

MDA-MD-231, MCF7, T47D, MDA-MB-453, CAL51, BT549 and Hs578T cells were cultured in Dulbecco's modified eagle medium (DMEM) supplemented with 5 % foetal bovine serum (FBS) and 4 mM L-glutamine (James *et al.*, 2023). Molecular identity was confirmed by short tandem repeat analysis (Masters *et al.*, 2001). MDA-MB-231 cells stably expressing shRNA targeting *SCN5A* were maintained in medium containing G418 (400 μg/ml) (Nelson *et al.*, 2015b). MDA-MD-231 cells stably expressing the VGSC β1 subunit with a C-terminal fusion to eGFP (Chioni *et al.*, 2009) (MDA-MB-231-β1-GFP) were maintained in medium containing hygromycin B (100 μg/ml). LS11-083 hTERT-immortalized primary breast cancer associated fibroblast cells were also cultured in DMEM with 5 % FBS and 4 mM L-glutamine (Roberts *et al.*, 2016). Cultures were confirmed to be *Mycoplasma*-free using the 4′,6-diamidino-2-phenylindole (DAPI) method (Uphoff *et al.*, 1992).

Orthotopic xenograft breast cancer model and tissue slice preparation

$Rag2^{-/-}$ $Il2rg^{-/-}$ mice were bred in-house and females over the age of 6 weeks were used for tumour implantation. A suspension of 1×10^{6} MDA-MB-231 cells in Matrigel (Corning; 50% v/v in phosphate-buffered saline (PBS)) was implanted into the left inguinal mammary fat pad of each animal whilst under isoflurane anaesthesia. Mice were weighed and their body condition and tumour size were checked at least every

2 days. Tumours were measured using callipers and the tumour volume was calculated using the modified ellsoidal formula, volume = ½(length × width^2). Mice were euthanized after approximately four weeks. Tumours were dissected immediately after euthanasia and sliced in ice-cold PBS using a Campden 5100MZ vibratome to a thickness of 200 μm for patch clamp recording or 500 μm for pH-sensitive microelectrode recording. To make lung tissue slices, following euthanasia, the trachea was cannulated, and 1% low melting point agarose (37 ˚C) was injected into the lungs. The lungs were then dissected and embedded in 1% low melting point agarose and sliced to a thickness of 250 μm for patch clamp recording.

Human breast cancer tissue and primary cells

Biopsy samples from four patient breast tumours that were excess to pathology requirements were acquired via the Breast Cancer Now Tissue Bank (BCNTB). The samples were transported in culture medium on ice and arrived within 24 h of surgical resection. Fresh tissue slices (250 μm thick) were cut in ice-cold PBS using a vibratome (Campden 5100MZ). For isolation of single cells from the tumour tissue, fragments were cut and then dissociated using a MACS Tumour Dissociation Kit (Miltenyi Biotec). Primary cultures of breast cancer cells, enriched for carcinoma cells, and primary cultures of purified normal breast epithelial cells were also acquired via the BCNTB. Human primary cells and tumour slices were cultured in DMEM:F12 with 1 ml/100 ml penicillin/streptomycin, 2.5 μg/ml Fungizone, 10 % FBS, 0.5 μg/ml hydrocortisone, 10 μg/ml apo-transferrin, 10 ng/ml human EGF and 5 μg/ml insulin. The cells were cultured on collagen-coated glass coverslips in plastic dishes.

Whole-cell patch clamp recording

The whole-cell patch clamp technique was used to record cell membrane currents from cells grown on glass coverslips (Leslie *et al.*, 2020). Filamented borosilicate capillary tubes were pulled and fire-polished to a resistance of ~5 MΩ for recording

from cell lines and ~10 MΩ for recording from primary cells. The extracellular physiological saline solution (PSS) contained (in mM) NaCl 144, KCl 5.4, $MgCl_2$ 1, $CaCl_2$ 2.5, HEPES 5, D-glucose 5.6, and was adjusted to pH 7.2 (unless otherwise stated) using NaOH. The intracellular recording solution for measuring Na^+ currents contained (in mM) NaCl 5, CsCl 145, $MgCl_2$ 2, $CaCl_2$ 1, HEPES 10, EGTA 11 and was adjusted to pH 7.4 (unless otherwise stated) using CsOH (Leslie *et al.*, 2020). The intracellular recording solution for measuring K^+ currents contained (in mM) NaCl 5, KCl 145, $MgCl_2$ 2, $CaCl_2$ 1, HEPES 10, EGTA 11 and was adjusted to pH 7.4 using KOH. Recordings were made using a MultiClamp 700B amplifier (Molecular Devices). Currents were digitized using a Digidata 1440A interface (Molecular Devices), low pass filtered at 10 kHz, sampled at 50 kHz, and analysed using pCLAMP 10.7 software (Molecular Devices). For detection of small currents in primary cells, patient tumour slices and a panel of cell lines, signals were post-filtered at 1 kHz. For examination of pH dependency in MDA-MB-231 cells, currents were noise-corrected by subtracting half the peak-to-peak noise measured during the 10 ms period before depolarisation (Leslie *et al.*, 2020). Series resistance was compensated by 40-60% and linear components of leak were subtracted using a P/6 protocol (Armstrong & Bezanilla, 1977). Cells were clamped at a holding potential of -120 mV for 250 ms. Two main voltage clamp protocols were used, as follows:

1. To assess the voltage-dependence of activation of VGSCs and K^+ channels, cells were held at -120 mV for 250 ms and then depolarised to test potentials in 5-10 mV steps between -120 mV and +30 mV for 50 ms.

2. To assess the voltage-dependence of steady-state inactivation, cells were held at -120 mV for 250 ms followed by prepulses for 250 ms in 5-10 mV steps between -120 mV and +30 mV and a test pulse to -10 mV for 50 ms.

Pharmacology

Tetrodotoxin citrate (TTX, HelloBio HB1035) was diluted in sterile-filtered water to a stock concentration of 1 mM TTX/8.44 mM citrate and stored at -30 °C. The working concentration was 30 µM TTX/253 µM citrate. Ouabain octahydrate (Sigma O3125) was diluted in DMSO to a stock concentration of 50 mM and stored at -30 °C. The working concentration was 300 nM. Cariporide (Santa Cruz Biotechnology SC337619) was diluted in DMSO to a stock concentration of 50 mM and stored at -30 °C. The working concentration was 20 µM. Sodium iodoacetate (Acros Organics 170970250) was diluted in water and made up immediately before each experiment. The working concentration was 2 µM. Oligomycin (Santa Cruz Biotechnology SC201551) was diluted to a stock concentration of 10 mM in DMSO and stored at -30 °C. The working concentration was 1 µM. The background concentration of DMSO in these solutions was ≤0.01%.

RNA sequencing

Mice were housed (up to 4 per cage) and were chosen at random for cell implantation ensuring that both cell types were represented within each cage/block. RNA was extracted from 12 xenograft tumours (6 control MDA-MB-231 tumours, and 6 $Na_v1.5$-knock-down MDA-MB-231 tumours) tissue using TRIzol (Invitrogen), according to the manufacturer's instructions. RNA quality assessment, library preparation, and 150 bp short-read, paired-end sequencing were conducted by Novogene Europe (Cambridge UK) with 1 µg of total RNA used for sequencing library construction with NEBNext Ultra RNA Library Prep Kit for Illumina (NEB, USA). Raw FastQ files were mapped using Bbsplit from the Bbtools 39.01 suite against Mm10 and hg38 (Bushnell, 2022) and ambiguous reads were excluded. The disambiguated hg38 reads were aligned and count matrix created using Rsubread v2.12.3 (Liao *et al.*, 2019). Differential gene expression was calculated using DESeq2 v2_1.38.1 (Love *et al.*, 2014). Gene set enrichment analysis (GSEA) was performed using the implementation in VULCAN v1.20.0 (Holding *et al.*, 2019). All

code for the analysis is available from https://github.com/andrewholding/RNASeq-SCN5A.

pH-selective microelectrodes

Unfilamented borosilicate capillary tubes were pulled to a resistance of ~5 MΩ (measured after silanization and when filled with PSS and in a recording bath). Silanization was performed at 200 °C for 15 min with N,N-dimethyltrimethylsilyamine. Microelectrodes were back filled with the following solution (in mM): NaCl 100, HEPES 20, NaOH 10, adjusted to pH 7.5. Microelectrodes were then front-filled with H^+ ionophore I – cocktail A (Sigma) (Voipio *et al.*, 1994). Recordings were made using a MultiClamp 900A amplifier (Molecular Devices) linked to a computer running MultiClamp 900A Commander software (Molecular Devices). The headstage amplifier was a high impedance 0.0001MU Axon HS-2 (Molecular Devices). Currents were digitized using a ITC018 A/D converter (HEKA Instruments), regular oscillatory noise was reduced with a HumBug noise eliminator (Quest Scientific) and the voltage signal was low-pass filtered at 10 Hz. Voltage was recorded using Axograph software (version 1.7.6). Electrodes were calibrated and offset from junction potentials empirically measured every 12 measurements to avoid drift with repeated electrode placement. A straight line was fitted to the offset-corrected voltage/pH calibration points, and the equation of this straight line was used to calculate the corresponding tissue slice pH. Tissue slices were maintained at 30 °C and 100 % humidity in the recording chamber at the interface between air and perfused PSS. Measurements were made on the top surface of tumour tissue slices within one hour of euthanasia. In total, 12 measurements were made from each region of the slice, alternating between regions, with calibrations and bath measurements taken before, half-way through and at the end of the series of measurements.

Immunohistochemistry

Tissue cryopreservation, sectioning and immunohistochemistry were performed as described previously (Nelson *et al.*, 2015b). The following primary antibodies were used: rabbit anti active caspase 3 (1:200; R&D Systems AF835), and rabbit anti-Ki67 (1:5000; Abcam AB15580). The secondary antibody was Alexa-568 conjugated goat anti-rabbit (1:500; Invitrogen A11036). Sections were mounted in Prolong Gold + DAPI (Thermofisher). Stained sections were imaged on a Zeiss AxioScan.Z1 slide scanner at 20X. Images were viewed using Zen 3.4 (blue edition) software (Zeiss) and the maximal intensity in the red and blue (DAPI) channels were changed to maximise the visibility of positively stained cells. The minimum intensity was not changed. Images were then converted from .czi format to 8-bit .tif format. In ImageJ, a whole section was viewed at a time and the shape was matched to the drawing of the tissue slice during pH-selective microelectrode recording. Regions of interest (ROIs; 1000 x 1000 pixels) were chosen in both "core" and "peripheral" regions identified during recording. Six ROIs were selected from each region. Image analysis was performed using an ImageJ macro. Briefly, a nuclear count was performed by a particle count in the DAPI channel. A minimum intensity threshold was applied to the Alexa-568 channel, and this value was kept consistent within all ROIs from each tissue section. Activated caspase 3 staining was assessed by a particle count. Nuclear Ki67 staining was quantified by a particle count where the DAPI signal colocalised with the Alexa-568 signal. Particle counts were expressed as a percentage of the DAPI particle count in the same ROI to give a percentage of positively stained cells in each ROI for each antibody and averaged to give a single value for each section (one section per tumour).

Measurement of intracellular pH

Cells were grown on glass coverslips for 48 h, then were incubated for 10 minutes at 21 °C in 1 µM 2′,7′-Bis(2-carboxyethyl)-5(6)-carboxyfluorescein acetoxymethyl ester (BCECF-AM, Biotium) in PSS, washed twice then left in PSS at pH 7.2 for 30

minutes before incubation for 10 minutes in PSS at pH 6.0 or 7.2. Coverslips were mounted in a Warner RC-20H recording chamber used in open configuration at room temperature with PSS perfusion at 1 ml/min. Two-point calibration was performed at the end of every experiment using K^+-based PSS (where Na^+ was replaced by K^+) at pH 7.0 with 13 µM nigericin (Sigma) for 7 minutes, followed by K^+-based PSS at pH 8.0 for a further 7 minutes. For each individual cell, a standard curve was derived using these buffers of known pH value, and the resting pH calibrated. Exposures of 0.15 s duration were taken every 15 s with a Nikon Eclipse TE200 epi-fluorescence microscope using SimplePCI 6.0 software to control the imaging system. Images were captured with a RoleraXR Fast1394 CCD camera (Q-imaging) with a 10X Plan Fluor objective. Images were saved as 16-bit .tif files and analysed in ImageJ 1.53c. Circular ROIs were placed over cells. The mean intensity at each wavelength was calculated for each ROI. Background fluorescence was calculated for each excitation wavelength by selecting a ROI containing no cells. Background fluorescence was subtracted from the mean intensity of each ROI before fluorescence ratio calculation. Each experimental repeat was the mean measurement from ~40 cells per coverslip.

Measurement of intracellular [Na^+]

Cells were seeded at 2×10^4 (MCF7) or 2.5×10^4 (MDA-MB-231) cells/well in a 96 well, black walled, µclear polymer-bottomed plate (Greiner 655097). Medium was exchanged and drug incubations started after 36 h. Before dye loading, wells were washed with PBS, and 60 µl DMEM containing SBFI-AM (10 µM) and Pluronic F-127 (0.1%) ± drug treatment was added to each well. Cells were incubated in SBFI-AM at 37 °C for 2 h. Wells were then washed twice in PSS ± drug and left in PSS ± drug for imaging on a BMG Clariostar plate reader with excitation at 340 and 380 nm and emission collected at 510 nm. Background fluorescence was subtracted from each wavelength before fluorescence ratio calculation. Each experimental repeat was the mean fluorescence ratio of five wells from a single plate.

Viability assay

Cells were cultured in 6 well plates. Culture medium from each well was removed from wells into 14 ml Falcon tubes then adherent cells were detached using trypsin-EDTA and added to the same tube. Cells were centrifuged at 800 x g for 5 minutes and resuspended in medium. A 10 µl sample was mixed with an equal volume of trypan blue and the number of viable and dead cells was counted using an Invitrogen Countess automated cell counter.

Metabolic profiling

Oxygen consumption rate (OCR) and extracellular acidification rate (ECAR) measurements were conducted using a Seahorse XFe 96 Extracellular Flux Analyzer (Agilent), based on methods described previously (Epstein *et al.*, 2014; van Weverwijk *et al.*, 2019). The day prior to the measurements, MCF7 (2.0 x 10^4/well) and MDA-MB-231 (3.0 x 10^4/well) were seeded in a Seahorse XF96 cell culture microplate (101085-004, Seahorse) in 100 µl DMEM supplemented with 5% FBS and left at room temperature for 1 hour. Cells were then transferred to a 5% CO_2 incubator set at 37 °C and incubated overnight. The next day, cells were washed twice with 200 µl of assay medium (DMEM powder, D5030, Sigma; resuspended in 1 L UF water, adjusted to pH 7.4, sterile-filtered and supplemented on the day of the experiment with 17.5 mM glucose, 2 mM glutamine and 0.5 mM sodium pyruvate) and a final volume of 180 µl was left per well. Then, the plate was incubated for equilibration in a 37°C non-CO_2 incubator for 1 hour. In the meantime, the sensor cartridge was loaded with 20 µl of a 10X concentrated TTX solution to give a final assay concentration in the well of 30 µM. The experimental protocol consisted of consecutive cycles of 6 minutes that included 2 minutes of mixing followed by 4 minutes of measuring OCR and ECAR. After calibration of the cartridge and equilibration of the cell plate in the Seahorse Analyzer, basal measurements were

acquired for 6 cycles, followed by the injection of the TTX or water vehicle control and measurement for 10 more cycles. There were 6 replicate wells per plate and the experiment was repeated independently 3 times.

Human breast cancer tissue microarray

A tissue microarray consisting of formalin-fixed, paraffin-embedded cores from 1740 primary breast tumours was obtained from the BCNTB. Immunohistochemistry was performed as described previously (Nelson *et al.*, 2014). Samples were incubated with rabbit anti-human $Na_V1.5$ antibody (1:100; Alomone ASC-013, which recognises residues 1978-2016 of human $Na_V1.5$) and stained with EnVision+ Dual Link System/HRP (DAB+; Dako), following the manufacturer's instructions and counterstained with Mayer's haematoxylin. Specificity of staining was evaluated using antibody pre-adsorbed with immunising peptide. Slides were imaged on a Zeiss AxioScan.Z1 slide scanner with a 20X objective and staining was visualised using Zen 3.4 (blue edition) software (Zeiss). Staining was quantified using a modification to the Allred scoring system, as described previously (Harvey *et al.*, 1999; Nelson *et al.*, 2015b). TKL did the scoring and 10% of the series was independently verified by WF. In both cases, scoring was performed without prior knowledge of the associated clinical data. Concordance between investigators was assessed using Cohen's weighted kappa (0.814; 95% CI 0.766-0.865; P < 0.001) and Intraclass Correlation Coefficient (0.954; 95% CI 0.938-0.966; P < 0.001), indicating excellent agreement between scorers.

Statistical analysis

Linear regression was used for calibration of ratiometric indicators and pH electrodes over the recording ranges used. Statistical analysis was performed on raw (non-normalized) data using GraphPad Prism 9 for most analyses. IBM SPSS Statistics 27 was used to compute Intraclass Correlation Coefficient, Cohen's weighted kappa and

perform Cox multivariate proportional hazard analysis. Pairwise statistical significance was determined with Student's paired or unpaired, or one-sample t tests for normally distributed data and Mann-Whitney tests for non-parametric data. Multiple comparisons were made using ANOVA and Tukey's or Dunnett's multiple comparisons tests for normally distributed data and using Kruskal-Wallis or Friedman's tests for non-parametric data. Seahorse data were analysed using 2-way ANOVA. Time to event data were analysed using Kaplan Meier plots and log-rank (Mantel-Cox) tests computed with hazard ratio (HR) and 95% confidence intervals (CI). Correlations between $Na_v1.5$ and other protein markers in the tissue microarray were evaluated using Spearman's test. Results were considered significant at P < 0.05 or Benjamini-Hochberg (BH) adjusted P < 0.05 for differential gene expression.

Results

Inward currents are present in several triple-negative breast cancer cell lines, cancer-associated fibroblasts and lung metastases

We first set out to evaluate the presence of voltage-gated inward currents across a panel of breast cancer cell lines and a breast tumour-derived cancer-associated fibroblast (CAF) cell line. Inward currents were present in MDA-MB-231 cells with activation at around -50 mV (Figure 1A, B), consistent with previous reports (Roger *et al.*, 2003; Fraser *et al.*, 2005). Similar inward currents were detected in two other cell lines: Hs578T and BT549 (Figure 1A). Notably, all three cell lines in which inward currents were detected are from TNBC (Supplementary Table 1). In addition, inward currents were present in the CAF line LS11-083, with activation at around -25 mV (Figure 1A, C, D). The whole cell capacitance was similar across all cell lines (Figure 1E). Thus, VGSCs are functionally active in a subset of TNBC and CAF cell lines.

We have shown previously that inward Na^+ currents are present in tissue slices taken from orthotopic xenograft MDA-MB-231 tumours (Nelson *et al.*, 2015b). Here, to evaluate whether Na^+ currents are retained following metastasis *in vivo*, we performed whole cell patch clamp recording on primary tumour tissue slices and lung tissue slices from the same animals. MDA-MB-231 cells stably expressing the VGSC β1 subunit tagged with GFP (β1-GFP) were used for these xenografts because they form lung metastases more readily than wild-type MDA-MB-231 cell tumours (Nelson *et al.*, 2014; Haworth *et al.*, 2022). The GFP tag also permitted identification of tumour cells within the lung parenchyma for recording (Figure 1F). This experiment revealed that Na^+ currents were of similar size in cells from primary tumours and lung metastases (P = 0.90; n = 24 cells in primary tumour and 8 cells in metastases; *t* test; Figure 1G). In summary, voltage-gated inward currents consistent with VGSCs are variably expressed across TNBC cell lines and CAFs and are retained in primary tumours and metastases in mice *in vivo*.

$Na_v1.5$ regulates expression of migration and invasion-promoting genes

Given that $Na_v1.5$ activity increases invasion and metastasis in preclinical models (Roger *et al.*, 2003; Fraser *et al.*, 2005; Brackenbury *et al.*, 2007; Gillet *et al.*, 2009; Driffort *et al.*, 2014; Nelson *et al.*, 2015a; Nelson *et al.*, 2015b), we hypothesised that knockdown of $Na_v1.5$ would suppress expression of invasion-related genes. To test this hypothesis, we next compared gene expression in six MDA-MB-231 xenograft tumours with six tumours in which *SCN5A* expression had been stably suppressed using shRNA (Nelson *et al.*, 2015b). We previously showed that $Na_v1.5$ protein expression and Na^+ current was ablated in these cells (Nelson *et al.*, 2015b). *SCN5A* mRNA expression was significantly reduced in the shRNA tumours compared to control tumours (adjusted P < 0.01; n = 6; Supplementary Figure 1A). Principal component analysis (PCA) showed that the *SCN5A* knockdown aligned with PC2, explaining 25% of the variance (Supplementary Figure 1B), and there was no

relationship to mouse cage/block (Supplementary Figure 1C). The *SCN5A* shRNA tumours displayed 136 differentially expressed genes, compared to the control tumours (BH-adjusted P < 0.05; Figure 2A-C).

We next performed GSEA to evaluate the relationship between *SCN5A* knockdown and invasion-promoting genes (Subramanian *et al.*, 2005). We used the MSigDB gene set SCHUETZ_BREAST_CANCER_DUCTAL_INVASIVE_UP which describes genes up-regulated in invasive ductal carcinoma vs ductal carcinoma in situ (Schuetz *et al.*, 2006). GSEA of the differential expression in *SCN5A* shRNA tumours showed a significant reduction in invasion transcriptional response (P < 0.001; Figure 2D). There was also a smaller reduction in expression of invasion-downregulated genes (normalised enrichment score -2.35 vs. -3.58), although this was only significant at a reduced stringency (P < 0.05; Supplementary Figure 1D).

Extracellular pH is low in xenograft tumours and correlates with cellularity and proliferation

The tumour microenvironment is reported to be acidic (White *et al.*, 2017), with a low intratumoural pH facilitating various metastatic hallmarks including ECM degradation and invasion. Given that $Na_v1.5$ activity has previously been shown to increase breast cancer cell invasion via acidification of extracellular fluid (Brisson *et al.*, 2011; Brisson *et al.*, 2013), we next assessed the extracellular pH (pH_e) of MDA-MB-231 xenograft tumours using pH-sensitive microelectrodes. Measurements were recorded in several locations on tissue slices directly prepared from xenograft tumours, alternately from the opaque core of the tumour, and from more translucent periphery (Figure 3A). The mean overall pH_e was 6.9 ± 0.1, which is significantly lower than pH 7.4 normally found in the extracellular fluid of healthy tissue (P = 0.001; n = 9; one sample *t* test) (White *et al.*, 2017). The mean pH_e in the core was 7.0 ± 0.1; in the periphery, it was significantly lower, at 6.8 ± 0.1 (P < 0.01; n = 9; paired *t* test; Figure

3B). The differences between core and periphery were investigated in more detail using immunohistochemistry (Figure 3C). Cellularity (mean nuclear count per ROI) was significantly higher in the periphery compared to the core (988 ± 29 vs. 838 ± 47; P < 0.01; n = 9 tumours; paired t test; Figure 3D). Similarly, proliferation, as measured by Ki67-positive nuclei, was significantly higher in the periphery than the core (20.1 ± 7.0% in the periphery vs. 6.9 ± 2.4% in core; P < 0.05; n = 9 tumours; paired t test; Figure 3E). Conversely, apoptosis, as measured by cleaved caspase 3 positivity, was significantly higher in the core compared to the periphery (10.9 ± 3.5% in core vs. 2.1 ± 0.7% in periphery; P < 0.05; n = 9 tumours; paired t test, Figure 3F). Taken together, these results suggest that in this model the pH_e is lower in peripheral regions with high cellularity which are proliferating rapidly, and higher towards the core of the tumours, where there is more apoptosis.

Persistent Na$_v$1.5 current is increased by low extracellular pH

Given that the tumour pH_e is acidic *in vivo*, we next assessed the effect of pH alteration on Na$_v$1.5 activity in MDA-MB-231 cells using patch clamp recording. A pH_e of 7.2 was compared with a pH_e of 6.2 since the pH_e of solid tumours has been reported to approach this level of acidity (Gatenby & Gillies, 2004). Lowering pH_e to 6.2 reduced the transient Na$^+$ current from -13.5 ± 2.3 pA/pF to -9.6 ± 1.5 pA/pF (P < 0.01; n = 11; Wilcoxon matched pairs test; Figure 4A, C). In contrast, the persistent Na$^+$ current, measured at 20-25 ms after depolarisation, was increased by acidification to pH_e 6.2, from -0.31 ± 0.04 pA/pF to -0.71 ± 0.11 pA/pF (P < 0.01; n = 10; paired t test; Figure 4B, D). Analysis of the voltage dependence of activation and steady-state inactivation (Figure 4E-G) revealed that the slope factor (k) and voltage at half activation (V½) did not significantly change when the pH_e was reduced from 7.2 to 6.2 (P = 0.077 and 0.087, respectively; n = 10; paired t tests, Table 1). However, the voltage at half inactivation (V½) was significantly depolarised at pH_e 6.2, from -80.4 ± 1.4 mV to -73.3 ± 2.8 mV (P < 0.01; n = 10; paired t test; Table 1)

and the slope factor for inactivation was also significantly changed, from -8.4 ± 0.8 mV to -11.9 ± 0.9 mV (P < 0.01; n = 10; paired t test; Table 1). This depolarising shift in steady-state inactivation thus increased the size of the window current (Figure 4G, H). Indeed, at the reported resting V_m of MDA-MB-231 cells, -18.9 mV (Fraser et $al.$, 2005), reducing the pH_e from 7.2 to 6.2 more than doubled channel availability from 1.9 ± 0.6% to 4.9 ± 0.7% of maximum (P < 0.05; n = 10; paired t test; Figure 4I). When pH_e was reduced further to 6.0, the effect on channel availability was even greater, increasing nearly five-fold from 2.1 ± 0.9% at pH_e 7.2 to 10.3 ± 2.2% of maximum at pH_e 6.0 (P < 0.001, n = 8, paired t test; Figure 4J & Supplementary Figure 2A, B). In addition, the time to peak Na^+ current was slower at pH_e 6.2 (3.02 ± 0.35 ms vs. 2.07 ± 0.24 ms at -10 mV; P < 0.001; n = 10; paired t test; Table 1; Figure 4K).

To explore whether the pH_e-dependent electrophysiological changes were mediated by a change in intracellular pH (pH_i), we assessed the effect of pH_e on pH_i using the ratiometric fluorescent pH indicator BCECF-AM to measure pH_i following incubation in different pH_e. Lowering pH_e from 7.2 to 6.2 led to intracellular acidification to pH_i 6.3 ± 0.1 (Supplementary Figure 3A, B). However, altering pH_i from 7.2 to 7.6 (empirically the maximum range of intracellular patch pipette solution pH which still allowed the formation of giga-Ohm seals onto MDA-MB-231 cells) had no effect on transient or persistent Na^+ current or voltage dependence of activation or steady-state inactivation (Supplementary Figure 3C-G). In summary, Na^+ entry into breast cancer cells through $Na_v1.5$ is increased in acidic pH_e but is not sensitive to changes in pH_i under the range of pH_i tested. These data suggest that the increased persistent Na^+ current generated in an acidic tumour microenvironment would potentiate a constant leak of Na^+ into breast cancer cells at steady state, which in turn may impact on the intracellular milieu.

Na$^+$ influx via Na$_v$1.5 promotes glycolytic H$^+$ production

Na$_v$1.5 activity has been shown to increase H$^+$ extrusion from breast cancer cells via NHE1, thus providing an optimal pH$_e$ for cysteine cathepsin-dependent extracellular matrix degradation and increased invasion (Gillet *et al.*, 2009; Brisson *et al.*, 2011; Brisson *et al.*, 2013). The mechanistic link between VGSC (Na$_v$1.5) activity and NHE1-mediated H$^+$ extrusion is however unclear, given that increased Na$^+$ influx would be expected to reduce the driving force for H$^+$ export via NHE1. One possibility is that Na$^+$ influx via Na$_v$1.5 increases Na$^+$/K$^+$ ATPase (NKA) activity to remove additional intracellular Na$^+$ (Pellerin & Magistretti, 1994; Skou, 1998). NKA preferentially uses ATP derived from glycolysis in many tissues including breast cancer cells (James *et al.*, 1996; Dutka & Lamb, 2007; Epstein *et al.*, 2014). Accordingly, Na$^+$ entry through Na$_v$1.5 may lead to an increase in glycolytic rate and thus H$^+$ production. To test this hypothesis, we firstly examined the source of ATP used by NKA in breast cancer cells. As expected, the NKA inhibitor ouabain (300 nM; 6 h incubation) increased [Na$^+$]$_i$, measured using the ratiometric Na$^+$ indicator SBFI-AM, by 2.2-fold in MDA-MB-231 cells and by 2.0-fold in MCF7 cells (P < 0.001 for both; n = 6; one sample *t* tests; Figure 5A, B). Interestingly, inhibition of mitochondrial respiration using the ATP synthase inhibitor oligomycin (1 μM; 2 h) did not alter [Na$^+$]$_i$ in MDA-MB-231 or MCF7 cells (P = 0.62 and P = 0.14, respectively; n = 6; one sample *t* test; Figure 5A, B). However, inhibition of glycolysis with the GAPDH inhibitor sodium iodoacetate (2 mM; 2 h) significantly increased [Na$^+$]$_i$ in both cell lines (P < 0.01 for both cell lines; n = 6; one sample *t* test; Figure 5A, B). These data suggest that NKA was able to maintain a steady [Na$^+$]$_i$ in the absence of mitochondrial respiration but not in the absence of glycolysis. Oligomycin had no effect on the viability of either cell line (P = 0.74 for MDA-MB-231 and P = 0.84 for MCF7; n = 3; one sample *t* tests), whereas sodium iodoacetate significantly reduced viability in both cell lines (P < 0.01 for MDA-MB-231 and P < 0.05 for MCF7; n = 3; one sample *t* tests; Supplementary Figure 4). Ouabain reduced viability of MCF7

cells by 16% over 2 hours (P < 0.05; n = 3; one sample t test) but had no effect on MDA-MB-231 cells (P = 0.18; n = 3; one sample t test; Supplementary Figure 4). Thus, these data suggest that in both MDA-MB-231 and MCF7 cells, NKA activity requires ATP derived from glycolysis, rather than mitochondrial respiration, to export Na^+. Furthermore, both NKA activity and glycolysis are necessary to maintain cell survival in these experimental conditions.

Next, we used a Seahorse XFe96 analyser to test the hypothesis that Na^+ influx through $Na_v1.5$ increases the rate of glycolysis via increasing ATP demand from NKA. The effects of TTX (30 µM) on the extracellular acidification rate (ECAR; a measure of glycolysis) and oxygen consumption rate (OCR; a measure of mitochondrial respiration) were compared between MDA-MB-231 and MCF7 cells. In control experiments, addition of TTX to wells containing medium without cells transiently reduced the measured ECAR, which then returned to baseline levels within 15 minutes (Supplementary Figure 5). Therefore, measurements on cells ± TTX were compared after this period. Addition of TTX (30 µM) to MDA-MB-231 cells caused a rapid and sustained reduction in ECAR (P < 0.0001; n = 3 experimental repeats containing 6 wells each; 2-way ANOVA; Figure 5C, D). However, TTX had no significant effect on the ECAR of MCF7 cells, which do not express functional $Na_v1.5$ channels (Chioni *et al.*, 2005) (P = 0.07; n = 3 experimental repeats containing 6 wells each; 2-way ANOVA; Figure 5E, F). In contrast, OCR was unaffected by TTX in both cell lines (P = 0.99 and 0.11 for MDA-MB-231 and MCF7, respectively; n = 3 experimental repeats containing 6 wells each; 2-way ANOVA; Figure 5G-J). Together, these data are consistent with Na^+ influx via $Na_v1.5$ increasing the rate of glycolysis. In turn, the latter increases H^+ production and export, thereby acidifying the extracellular space.

Prediction of $Na_v1.5$-dependent extracellular acidification rate

The expected ECAR due to VGSC activity can be calculated if it is assumed that the persistent Na^+ current into cells through VGSCs is counteracted by activity of NKA to maintain a stable $[Na^+]_i$. The other assumptions used in this calculation are that NKA pumps 3 Na^+ ions out of the cell per cycle in which it hydrolyses one molecule of ATP (Post & Jolly, 1957). Glycolytic production of lactic acid produces 2 molecules of ATP and 2 molecules of lactate per glucose. The production of H^+ by this reaction coupled to the ATPase hydrolysis of ATP to ADP generates 2 H^+ per glucose (Hochachka & Mommsen, 1983). There is therefore net production of one H^+ per ATP molecule generated by glycolytic fermentation to produce lactate. Using these assumptions, we calculated the ECAR due to $Na_v1.5$ activity to be 1.3 mpH/min (full calculations delineated in Supplementary Materials). This predicted ECAR is within an order of magnitude of the measured change in ECAR due to TTX inhibition of $Na_v1.5$ (9.8 ± 1.7 mpH/min; Figure 5D). Thus, our model can explain how $Na_v1.5$ activity can increase the bulk of H^+ extrusion through NHE1, considering experimental variability in determination of ECAR and persistent Na^+ current, and estimation of pH_e during the Seahorse assay.

Protein-protein interactions of NHE1

NHE1 is the pH regulator which has been implicated as most important in $Na_v1.5$-dependent extracellular acidification and protein-protein interactions have been suggested to play a significant role in regulating NHE1 activity, including by $Na_v1.5$ (Brisson *et al.*, 2013; Sjogaard-Frich *et al.*, 2021; Lopez-Charcas *et al.*, 2022). We therefore looked for other protein interactions of NHE1. Using the STRING database (v11.5) we searched for the top 50 likely protein interactions of NHE1, only considering the proteins sharing a physical complex. This identified several subunits of NKA as likely binding partners of NHE1 (Supplementary Figure 6). An interaction between NKA and NHE1 further supports a model in which NKA is an intermediate step by which Na^+ flux through channels such as $Na_v1.5$ can then alter NHE1 activity.

In summary, as well as showing how $Na_v1.5$ can increase tumour acidification via enhanced rate of glycolysis, we have demonstrated that the acidic tumour microenvironment can increase Na^+ influx via $Na_v1.5$. Together, these findings imply that there is a positive feedback loop in breast cancer cells, whereby Na^+ influx and H^+ release into the tumour microenvironment synergise to promote invasion and metastasis.

Primary breast epithelial and carcinoma cells exhibit voltage-sensitive inward and outward membrane currents

Our findings here build upon several studies which have independently shown that $Na_v1.5$ channels are functional in a subset of breast cancer cell lines (Roger *et al.*, 2003; Fraser *et al.*, 2005; Brackenbury *et al.*, 2007; Gillet *et al.*, 2009; Driffort *et al.*, 2014). We have also shown that $Na_v1.5$ is functionally active in tumour tissue slices from mice. However, there is a lack of evidence of functional VGSC currents from near patient models such as breast cancer patient-derived tumour tissue or primary cells. To address this critical knowledge gap, we next undertook electrophysiological characterization of membrane currents from primary human mammary epithelial and breast cancer cells in culture. We recorded membrane currents from a total of four normal human mammary epithelial cell samples and thirteen breast cancer samples enriched for carcinoma cells (Supplementary Table 3). Voltage-sensitive inward currents were present in two out of 13 breast cancer samples and none of four normal human mammary epithelial cell samples (Figure 6A, B, Table 2). Notably, inward currents were only present in cells from tumours which had metastasised to the lymph nodes, although this correlation was not statistically significant, likely due to the small number of inward currents detected (P = 0.16; Fisher's exact test; Table 2). The mean current-voltage relationship of the inward currents displayed activation

at approximately -50 mV, again consistent with VGSC currents recorded from breast cancer cell lines (Figure 6C) (Roger *et al.*, 2003; Yang *et al.*, 2020).

Voltage-sensitive, non-inactivating outward currents were present in all normal human mammary epithelial cell samples tested and 10 out of 13 breast cancer samples (Figure 6D, E; Table 2). Interestingly, outward currents were significantly more prevalent in cells from tumours which had metastasised to the lymph nodes (P < 0.01; Fisher's exact test; Table 2). However, the outward current density varied considerably between samples (Figure 6F). There was no difference in the size of the normal vs. cancer cells as estimated by measuring whole cell capacitance, although the range was larger in cancer cells than for normal cells (P = 0.18; n = 28 normal and 96 cancer cells; Mann-Whitney test; Figure 6G). In addition, the outward current density was not significantly different between breast cancer cells and normal cells (P = 0.08; n = 17 normal and 68 cancer cells; Mann-Whitney test; Figure 6H). However, the outward current density was significantly larger in ER+/HER2+ than TNBC samples (P < 0.001; n = 42 ER+/HER2+ and 26 TNBC cells; Figure 6I). In summary, (i) non-inactivating voltage-gated outward currents were common in normal human mammary epithelial cells and primary breast cancer cells whilst (ii) voltage-gated inward currents were detectable in malignant cells, albeit in a small subset.

Cells in patient breast cancer tissue exhibit voltage-sensitive inward and outward membrane currents

We next performed electrophysiological experiments to record membrane currents in a small cohort of three available breast tumour fresh tissue specimens. In tissue slices freshly prepared from patient tumour biopsies, there were few cellular areas and most of the slices were composed of connective tissue or fat. Thus, we took patch clamp recordings from pockets of cancer cells within the connective tissue at the top surface of the slice. Adipocytes were identified based on their large size and

avoided. Portions of each tumour were also dissociated and seeded onto coverslips to enable patch clamp recording from isolated cells.

Voltage clamp recording revealed that cells in tumour slices expressed both voltage-sensitive inward and outward currents (Figure 7A; Supplementary Table 4). The mean outward current-voltage relationship is typical of voltage-gated K^+ channels, with activation at approximately -50 mV (Figure 7B). Voltage-sensitive outward currents were present in all three specimens and 9/17 recordings made (Figure 7C). Small voltage-sensitive inward currents were found in two out of three patient specimens and in 4/17 recordings made (Figure 7D). The mean inward current-voltage relationship was noisy due to the small current density; however, it displayed activation at approximately -50 mV (Figure 7E), consistent with VGSC currents recorded from breast cancer cell lines (Roger *et al.*, 2003; Yang *et al.*, 2020).

Comparisons were then made between the cells in tumour slices, and cells dissociated from the same tumours plated onto glass coverslips (Supplementary Table 4). The outward current density was significantly larger in dissociated cells vs. those in slices (P < 0.05; t test; Figure 7F). The size of the cells, estimated by whole cell capacitance, also differed between cells in slices and on coverslips, with cells in slices being generally larger than dissociated cells (P < 0.05; *t* test; Figure 7G). However, the mean outward current-voltage relationship was broadly similar between cells in slices and on coverslips (Figure 7H). It was not possible to compare inward currents due to the small number of cells exhibiting these currents. In summary, both outward and inward currents were detected in cells in breast cancer tissue slices taken direct from patients, although making such recordings was technically challenging due to limited availability and the fibrous composition of the tissue. We therefore next moved to immunohistochemical characterization of $Na_v1.5$ protein expression in a breast cancer tissue microarray.

Na$_v$1.5 protein expression associates with poor clinical outcome in breast cancer patients

Expression of Na$_v$1.5 protein in breast cancer has previously been demonstrated in a small, qualitative study of 6 patients (Fraser *et al.*, 2005) and later in a study of 36 patients (Nelson *et al.*, 2015b). To test the prognostic value of Na$_v$1.5 in a larger cohort of patients, we evaluated Na$_v$1.5 expression in a breast cancer tissue microarray containing 1740 cases. In agreement with previous studies (Fraser *et al.*, 2005; House *et al.*, 2010; Nelson *et al.*, 2015b), Na$_v$1.5 was expressed mostly intracellularly, with some staining observed at the plasma membrane. Specificity of staining was confirmed by pre-incubation with the immunising peptide (Figure 8A). To explore correlation between Na$_v$1.5 expression and histoclinical characteristics of the patient population, staining scores between 0-3 were classed as 'low' and scores between 4-8 were classed as 'high' (Figure 8A).

High Na$_v$1.5 protein expression was correlated with larger tumour size ($P < 0.001$; Mann-Whitney U test; Figure 8B), lymph node positivity ($P < 0.05$; Mann-Whitney U test; Figure 8C), higher Nottingham Prognostic Index ($P < 0.001$; Mann-Whitney U test; Figure 8D), and higher tumour grade ($P < 0.001$; χ^2 test; Table 3). In addition, Na$_v$1.5 expression was significantly higher in patients who developed a distant metastasis ($P < 0.001$; Mann-Whitney U test; Figure 8E). Na$_v$1.5 was negatively associated with estrogen receptor (ER; $P < 0.05$; Fisher's exact test; Table 3) and progesterone receptor (PgR; $P < 0.001$; Fisher's exact test; Table 3) expression, but positively associated with human epidermal growth factor receptor 2 (HER2; $P < 0.01$; Fisher's exact test; Table 3). There was no association between Na$_v$1.5 expression and triple negative breast cancer (TNBC; $P = 0.12$; Fisher's exact test; Table 3), age ($P = 0.52$; Fisher's exact test; Table 3), menopause ($P = 0.46$; Fisher's exact test; Table 3) or endocrine therapy ($P = 0.51$; Fisher's exact test; Table 3).

However, high Na$_v$1.5 expression was correlated with recorded chemotherapy use (P < 0.05; Fisher's exact test; Table 3). These relationships are also displayed as violin plots in Supplementary Figure 7A-L.

High Na$_v$1.5 protein expression was associated with a significant reduction in metastasis-free survival (HR 2.18; 95% CI 1.63-2.92; P < 0.001; log-rank test; Figure 8F). This result was reflected in a significant reduction in overall survival (Supplementary Figure 8A), cancer-specific survival (Supplementary Figure 8B), disease-free survival (Supplementary Figure 8C), and local recurrence-free survival (Supplementary Figure 8D). When subdivided by receptor status, high Na$_v$1.5 expression was associated with significantly reduced overall survival in ER+ (P < 0.05), but not HER2+ (P = 0.14) or TNBC patients (P = 0.53), although the sample sizes for HER2+ and TNBC patients were considerably smaller than for ER+ patients (Supplementary Figure 8E-G).

The prognostic value of Na$_v$1.5 protein expression was considered in a Cox proportional hazards model including tumour size, grade, and lymph node status as categorical variables. Na$_v$1.5 expression was an independent predictor of survival alongside the other variables in this model (HR = 1.58; 95% CI 1.05-2.37, P < 0.05; Table 4). Finally, the correlation between Na$_v$1.5 expression and other protein markers previously scored in the same breast cancer tissue microarray was explored (Althobiti *et al.*, 2020). This analysis revealed that Na$_v$1.5 expression was significantly positively correlated with several other invasion-related protein markers (Table 5). In summary, high Na$_v$1.5 protein expression associated with worse prognosis in combined subtypes of breast cancer patients across a range of clinical measures, highlighting the proposed role of this ion channel in promoting invasion and metastasis.

Discussion

In this study on breast cancer, we show that widespread upregulation of $Na_v1.5$ expression associates positively with metastasis. We also identify a novel activity-dependent positive feedback role for this channel in pH regulation. Thus, we link Na^+ transport through $Na_v1.5$ to altered gene expression and metabolic function, providing an integrated mechanism by which $Na_v1.5$ can promote metastatic dissemination. We show specifically that acidity in the invasive perimeter of the tumour microenvironment enhances persistent Na^+ current through $Na_v1.5$ into breast cancer cells, thus promoting glycolytic metabolic activity. We therefore propose that $Na_v1.5$ is a key regulator of the ionic tumour microenvironment, facilitating local invasion during the early stage of metastasis.

Clinical significance of $Na_v1.5$ expression in breast cancer

Our findings further support, for the first time, a pro-metastatic role for $Na_v1.5$ in the clinical setting. We show that $Na_v1.5$ protein expression correlates with lymph node positivity, increased metastasis, higher tumour grade and consequently reduced survival. In addition, the prognostic potential of $Na_v1.5$ expression is comparable to lymph node status, tumour grade and tumour size. Although there have been several previous reports indicating that VGSCs are likely to be important prognostic indicators of breast cancer (Fraser *et al.*, 2005; Yang *et al.*, 2012; Nelson *et al.*, 2015b), no study has previously examined $Na_v1.5$ protein expression in a large cohort of breast cancer patients. Previously, VGSC protein expression in breast cancer has mostly been extrapolated from experiments on cell lines with differing metastatic potential (Roger *et al.*, 2003; Fraser *et al.*, 2005) and small cohorts of patients (Fraser *et al.*, 2005; Nelson *et al.*, 2015b; Yamaci *et al.*, 2017). Interestingly, we found that $Na_v1.5$ expression was negatively associated with ER status, and positively correlated with HER2 status, but not with TNBC status. Together, these

findings suggest possible functional linkage between ER, HER2 and $Na_v1.5$ expression. In agreement with this notion, ER+ MCF-7 cells have low VGSC expression (Fraser *et al.*, 2005), and silencing ER in MCF-7 cells increases Na^+ current and VGSC-dependent invasion (Mohammed *et al.*, 2016). In addition, various growth factor receptors, including EGFR, have been shown to regulate VGSC expression in carcinoma cells (Fraser *et al.*, 2014). Further work is required to determine how these receptors interact with $Na_v1.5$ in breast cancer cells.

$Na_v1.5$ expression appeared largely intracellular in most tumour samples. Therefore, we attempted to investigate functional channel activity at the plasma membrane using whole cell patch clamp recording. Although non-inactivating outward currents were widespread, small inward currents, indicative of $Na_v1.5$ activity, were rarer and harder to detect. This result is surprising given the high proportion of $Na_v1.5$ positive cells in the tissue microarray; however, given that a large proportion of the channels may be present on intracellular membranes, these would be unavailable to carry Na^+ current at the plasma membrane. Another possibility is that the tissue slice, dissociation, and cell culture conditions resulted in transport of channels away from the plasma membrane, as has been shown for $K_{Ca}3.1$ (Catacuzzeno *et al.*, 2012). A few studies have reported the presence of Na^+ currents carried by VGSCs in primary cells dissociated from mesothelioma and cervical tumour tissue, although the latter had been maintained in long-term culture (Fulgenzi *et al.*, 2006; Diaz *et al.*, 2007; Hernandez-Plata *et al.*, 2012). A recent study also demonstrated Na^+ currents in primary colorectal carcinoma cells (Lopez-Charcas *et al.*, 2022). To our knowledge, ours is the first report of putative VGSC currents in breast cancer tissue and primary cells from patients. Thus, we provide the first evidence in support of $Na_v1.5$ channel activity in clinical breast cancer specimens. Nevertheless, more work is required to refine the procedures for recording VGSC currents from such clinical material.

Functional consequences of $Na_v1.5$ expression

Numerous studies have shown that $Na_v1.5$ promotes invasion and migration of breast cancer cell lines *in vitro* (Roger *et al.*, 2003; Fraser *et al.*, 2005; Brackenbury *et al.*, 2007; Gillet *et al.*, 2009; Chioni *et al.*, 2010; Brisson *et al.*, 2011; Brisson *et al.*, 2013; Yang *et al.*, 2020). $Na_v1.5$ also potentiates tumour growth and metastasis of xenograft tumours *in vivo* (Driffort *et al.*, 2014; Nelson *et al.*, 2015a; Nelson *et al.*, 2015b). Consistent with these, VGSC activity in cancer cells enables the production of an elongated, mesenchymal phenotype (Gradek *et al.*, 2019; Yang *et al.*, 2020). Understanding the mechanisms behind these functions may help development of new cancer treatments targeting VGSC signalling pathways. Here, we found that $Na_v1.5$ expression in breast cancer xenografts regulates expression of a set of genes controlling invasion (Schuetz *et al.*, 2006). These results are consistent with the observation that VGSC activity in cancer cells aids degradation of the extracellular matrix (Gillet *et al.*, 2009) and agree with data showing that $Na_v1.5$ regulates invasion genes in colon cancer cells (House *et al.*, 2010; House *et al.*, 2015).

Our finding that the proliferating and invasive tumour periphery was more acidic than the hypoxic core agrees with a number of other studies (Helmlinger *et al.*, 1997; Grillon *et al.*, 2011; Grillon *et al.*, 2015; White *et al.*, 2017; Jardim-Perassi *et al.*, 2019). Low extracellular pH in tumours promotes invasion by increasing activity of low pH-dependent enzymes that degrade the ECM, such as cysteine cathepsins (Gillet *et al.*, 2009; Brisson *et al.*, 2011). We found that low pH_e, as found in tumours, increases the persistent Na^+ current through $Na_v1.5$ in breast cancer cells, thereby promoting Na^+ influx. Greater Na^+ influx into cancer cells in tumour regions of lower pH_e may be partially responsible for the heterogeneity of apparent tumour $[Na^+]$ as measured by [23]Na-MRI (Ouwerkerk *et al.*, 2007; James *et al.*, 2022).

Increased $Na_v1.5$-mediated Na^+ influx into cancer cells would be expected to promote activity of NKA (Nyblom *et al.*, 2013). There is substantial evidence that NKA utilises glycolysis as its main ATP source (Paul *et al.*, 1979; McBride & Early, 1989; Epstein *et al.*, 2014; Sepp *et al.*, 2014). NKA activity, and consequent glycolytic metabolism, would therefore increase the rate of H^+ production. In agreement with this paradigm, we found that $Na_v1.5$ activity in breast cancer cells increased glycolysis, as measured by extracellular H^+ accumulation, without affecting oxidative phosphorylation. Increasing $[Na^+]_i$, via the ionophore gramicidin, has previously been shown to potentiate the rate of glycolysis in breast cancer cells, whereas the NKA inhibitor ouabain decreased H^+ production (Epstein *et al.*, 2014). These findings echo those where glycolysis was found to be the ATP source for another plasma membrane ion pump, the plasma membrane Ca^{2+} ATPase (PMCA) in pancreatic cancer cells (James *et al.*, 2015; Bruce *et al.*, 2021). We found that inhibition of glycolysis induced a large increase in $[Na^+]_i$ similar to that caused by ouabain, and also rapidly led to cell death. From these results it is not possible to judge definitively whether NKA is more reliant on glycolysis than other ATP-consuming processes. However, in our *in vitro* experimental conditions, NKA activity is mainly powered by glycolysis, similar to other plasma membrane ATPases (James *et al.*, 2015), and the rate of glycolysis increases due to $Na_v1.5$-dependent Na^+ influx.

$Na_v1.5$-induced extracellular acidification – a positive feedback mechanism promoting invasion

Elevated steady-state Na^+ entry via persistent current through $Na_v1.5$, leading to increased glycolysis to power NKA, would in turn, be expected to increase H^+ extrusion through pH regulators such as CAIX and NHE1. Reciprocally, the reduction in pH_e serves to increase persistent Na^+ entry into breast cancer cells via $Na_v1.5$. Together, these mechanisms would lead to a positive feedback loop linking $Na_v1.5$, Na^+ entry, increased H^+ production via glycolysis and extracellular acidification

(Figure 9). This model fits with previous studies showing that $Na_v1.5$ activity increases NHE1-mediated H^+ extrusion in breast cancer cells, leading to ECM degradation and increased invasion (Brisson *et al.*, 2011; Brisson *et al.*, 2013). It would also explain how Na^+ influx via $Na_v1.5$ increases H^+ efflux via NHE1, despite a collapse of the inward Na^+ gradient (Leslie & Brackenbury, 2022). Given our evidence that extracellular acidification in breast tumours occurs particularly in the highly proliferative peripheral region, and that the persistent Na^+ current through $Na_v1.5$ is larger in acidic conditions, more Na^+ would be likely to enter breast cancer cells at the invading edges of the tumour.

Conclusion

Here, we have shown that $Na_v1.5$ is associated with poor prognosis and increased metastasis in breast cancer. Since $Na_v1.5$ is a negative prognostic indicator, and its expression increases tumour growth and metastasis in preclinical models (Nelson *et al.*, 2015b), it is a promising target for drug repurposing and discovery (Yang *et al.*, 2012; Djamgoz & Onkal, 2013; Capatina *et al.*, 2020). In agreement with this notion, we recently showed that exposure to certain persistent Na^+ current-inhibiting Class 1c and 1d antiarrhythmic drugs is associated with significantly improved cancer-specific survival (Fairhurst *et al.*, 2023). In addition, VGSC-inhibiting drugs have been shown to decrease tumour growth and metastasis in murine breast cancer models (Driffort *et al.*, 2014; Nelson *et al.*, 2015a). Furthermore, a recent clinical trial has shown that presurgical peritumoral treatment with lidocaine significantly improves disease-free and overall survival in women with early breast cancer (Badwe *et al.*, 2023). In conclusion, our results reveal a positive feedback mechanism by which Na^+ influx through $Na_v1.5$ promotes glycolytic H^+ production to increase invasive capacity and drive breast cancer metastasis. This novel mechanism, together with the emerging clinical data, underscores the value of $Na_v1.5$ as a prognostic marker and potential anti-metastatic therapeutic target.

Acknowledgements

The authors wish to acknowledge the roles of the Breast Cancer Now Tissue Bank in collecting and making available the samples and data, and the patients who have generously donated their tissues and shared their data to be used in the generation of this publication. The authors also thank Prof Miles Whittington (Hull-York Medical School, UK), Dr John Davey and Dr Katherine Newling (Technology Facility, University of York, UK) and Prof Lýdia Vargová (Charles University, Czechia) for providing invaluable advice. For the purpose of open access, a Creative Commons Attribution (CC BY) licence is applied to any Author Accepted Manuscript version arising from this submission.

Author contributions

Study conception and design: TKL, SC, WJB. Data collection: TKL, AT, ADJ, SPF, MAG, MB, MN, MT, WF, WJB. Analysis and interpretation of results: TKL, AT, ADJ, SPF, MT, WF, SCS, ER, GP, MBAD, APJ, HRM, CLHH, ANH, SC, WJB. Manuscript preparation: TKL, AT, ADJ, SPF, MT, WF, SCS, ER, VS, CB, GP, MBAD, APJ, HRM, CLHH, ANH, SC, WJB. All authors reviewed the results and approved the final version of the manuscript.

Data and code availability

The RNA-seq data are deposited in the GEO database, accession number GSE228621. The code used to analyse the data are available from https://github.com/andrewholding/RNASeq-SCN5A.

Funding

WJB received funding from Breast Cancer Now (2015NovPhD572) and Cancer Research UK (A25922). ANH received funding from the BBSRC (BB/V000071/1). SPF received funding from the Pro Cancer Research Fund. SCS, CLHH and APJ received funding from the British Heart Foundation (PG/14/79/31102 and PG/19/59/34582).

Competing interests

VS is one of the founders of the Breast Cancer Now Tissue Bank. MBAD is CEO and Director of CELEX Oncology Innovations Ltd.

References

Althobiti M, El Ansari R, Aleskandarany M, Joseph C, Toss MS, Green AR & Rakha EA. (2020). The prognostic significance of ALDH1A1 expression in early invasive breast cancer. *Histopathology* **77,** 437-448.

Anderson RL, Balasas T, Callaghan J, Coombes RC, Evans J, Hall JA, Kinrade S, Jones D, Jones PS, Jones R, Marshall JF, Panico MB, Shaw JA, Steeg PS, Sullivan M, Tong W, Westwell AD, Ritchie JWA, Cancer Research UK & Cancer Therapeutics CRCAMWG. (2019). A framework for the development of effective anti-metastatic agents. *Nat Rev Clin Oncol* **16,** 185-204.

Armstrong CM & Bezanilla F. (1977). Inactivation of the sodium channel. II. Gating current experiments. *J Gen Physiol* **70,** 567-590.

Badwe RA, Parmar V, Nair N, Joshi S, Hawaldar R, Pawar S, Kadayaprath G, Borthakur BB, Rao Thammineedi S, Pandya S, Balasubramanian S, Chitale PV, Neve R, Harris C, Srivastava A, Siddique S, Vanmali VJ, Dewade A, Gaikwad V & Gupta S. (2023). Effect of Peritumoral Infiltration of Local Anesthetic Before Surgery on Survival in Early Breast Cancer. *J Clin Oncol* **41,** 3318-3328.

Brackenbury WJ, Chioni AM, Diss JK & Djamgoz MB. (2007). The neonatal splice variant of Nav1.5 potentiates in vitro invasive behaviour of MDA-MB-231 human breast cancer cells. *Breast Cancer Res Treat* **101,** 149-160.

Brisson L, Driffort V, Benoist L, Poet M, Counillon L, Antelmi E, Rubino R, Besson P, Labbal F, Chevalier S, Reshkin SJ, Gore J & Roger S. (2013). NaV1.5 Na(+) channels allosterically regulate the NHE-1 exchanger and promote the activity of breast cancer cell invadopodia. *J Cell Sci* **126,** 4835-4842.

Brisson L, Gillet L, Calaghan S, Besson P, Le Guennec JY, Roger S & Gore J. (2011). Na(V)1.5 enhances breast cancer cell invasiveness by increasing NHE1-dependent H(+) efflux in caveolae. *Oncogene* **30,** 2070-2076.

Bruce JIE, Sanchez-Alvarez R, Sans MD, Sugden SA, Qi N, James AD & Williams JA. (2021). Insulin protects acinar cells during pancreatitis by preserving glycolytic ATP supply to calcium pumps. *Nat Commun* **12,** 4386.

Bushnell B. (2022). BBTools. sourceforge.net/projects/bbmap/.

Capatina AL, Lagos D & Brackenbury WJ. (2020). Targeting Ion Channels for Cancer Treatment: Current Progress and Future Challenges. *Rev Physiol Biochem Pharmacol*.

Cardoso F, Costa A, Norton L, Cameron D, Cufer T, Fallowfield L, Francis P, Gligorov J, Kyriakides S, Lin N, Pagani O, Senkus E, Thomssen C, Aapro M, Bergh J, Di Leo A, El Saghir N, Ganz PA, Gelmon K, Goldhirsch A, Harbeck N, Houssami N, Hudis C, Kaufman B, Leadbeater M, Mayer M, Rodger A, Rugo H, Sacchini V, Sledge G, van't Veer L, Viale G, Krop I & Winer E. (2012). 1st International consensus guidelines for advanced breast cancer (ABC 1). *Breast* **21,** 242-252.

Catacuzzeno L, Fioretti B & Franciolini F. (2012). Expression and Role of the Intermediate-Conductance Calcium-Activated Potassium Channel KCa3.1 in Glioblastoma. *J Signal Transduct* **2012,** 421564.

Chioni AM, Brackenbury WJ, Calhoun JD, Isom LL & Djamgoz MB. (2009). A novel adhesion molecule in human breast cancer cells: voltage-gated Na+ channel beta1 subunit. *Int J Biochem Cell Biol* **41,** 1216-1227.

Chioni AM, Fraser SP, Pani F, Foran P, Wilkin GP, Diss JK & Djamgoz MB. (2005). A novel polyclonal antibody specific for the $Na_v1.5$ voltage-gated Na^+ channel 'neonatal' splice form. *J Neurosci Methods* **147,** 88-98.

Chioni AM, Shao D, Grose R & Djamgoz MB. (2010). Protein kinase A and regulation of neonatal Nav1.5 expression in human breast cancer cells: activity-dependent positive feedback and cellular migration. *Int J Biochem Cell Biol* **42,** 346-358.

Diaz D, Delgadillo DM, Hernandez-Gallegos E, Ramirez-Dominguez ME, Hinojosa LM, Ortiz CS, Berumen J, Camacho J & Gomora JC. (2007). Functional expression of voltage-gated sodium channels in primary cultures of human cervical cancer. *J Cell Physiol* **210,** 469-478.

Djamgoz MB & Onkal R. (2013). Persistent current blockers of voltage-gated sodium channels: a clinical opportunity for controlling metastatic disease. *Recent Pat Anticancer Drug Discov* **8,** 66-84.

Djamgoz MBA, Fraser SP & Brackenbury WJ. (2019). In Vivo Evidence for Voltage-Gated Sodium Channel Expression in Carcinomas and Potentiation of Metastasis. *Cancers (Basel)* **11**.

Driffort V, Gillet L, Bon E, Marionneau-Lambot S, Oullier T, Joulin V, Collin C, Pages JC, Jourdan ML, Chevalier S, Bougnoux P, Le Guennec JY, Besson P & Roger S. (2014). Ranolazine inhibits NaV1.5-mediated breast cancer cell invasiveness and lung colonization. *Mol Cancer* **13,** 264.

Dutka TL & Lamb GD. (2007). Na+-K+ pumps in the transverse tubular system of skeletal muscle fibers preferentially use ATP from glycolysis. *Am J Physiol Cell Physiol* **293,** C967-977.

Elingaard-Larsen LO, Rolver MG, Sørensen EE & Pedersen SF. (2022). How Reciprocal Interactions Between the Tumor Microenvironment and Ion Transport Proteins Drive Cancer Progression. In *From Malignant Transformation to Metastasis: Ion Transport in Tumor Biology*, ed. Stock C & Pardo LA, pp. 1-38. Springer International Publishing, Cham.

Epstein T, Xu L, Gillies RJ & Gatenby RA. (2014). Separation of metabolic supply and demand: aerobic glycolysis as a normal physiological response to fluctuating energetic demands in the membrane. *Cancer Metab* **2,** 7.

Fairhurst C, Martin F, Watt I, Bland M, Doran T & Brackenbury WJ. (2023). Sodium channel-inhibiting drugs and cancer-specific survival: a population-based study of electronic primary care data. *BMJ Open* **13,** e064376.

Folcher A, Gordienko D, Iamshanova O, Bokhobza A, Shapovalov G, Kannancheri-Puthooru D, Mariot P, Allart L, Desruelles E, Spriet C, Diez R, Oullier T, Marionneau-Lambot S, Brisson L, Geraci S, Impheng H, Lehen'kyi V, Haustrate A, Mihalache A, Gosset P, Chadet S, Retif S, Laube M, Sobilo J, Lerondel S, Villari G, Serini G, Pla AF, Roger S, Fromont-Hankard G, Djamgoz M, Clezardin P, Monteil A & Prevarskaya N. (2023). NALCN-mediated sodium influx confers metastatic prostate cancer cell invasiveness. *EMBO J,* e112198.

Fraser SP, Diss JK, Chioni AM, Mycielska ME, Pan H, Yamaci RF, Pani F, Siwy Z, Krasowska M, Grzywna Z, Brackenbury WJ, Theodorou D, Koyuturk M, Kaya H, Battaloglu E, De Bella MT, Slade MJ, Tolhurst R, Palmieri C, Jiang J, Latchman DS, Coombes RC & Djamgoz MB. (2005). Voltage-gated sodium channel expression and potentiation of human breast cancer metastasis. *Clin Cancer Res* **11,** 5381-5389.

Fraser SP, Hemsley F & Djamgoz MBA. (2016). Caffeic acid phenethyl ester: Inhibition of metastatic cell behaviours via voltage-gated sodium channel in human breast cancer in vitro. *Int J Biochem Cell Biol* **71,** 111-118.

Fraser SP, Ozerlat-Gunduz I, Brackenbury WJ, Fitzgerald EM, Campbell TM, Coombes RC & Djamgoz MB. (2014). Regulation of voltage-gated sodium channel expression in cancer: hormones, growth factors and auto-regulation. *Philos Trans R Soc Lond B Biol Sci* **369,** 20130105.

Fraser SP & Pardo LA. (2008). Ion channels: functional expression and therapeutic potential in cancer. Colloquium on Ion Channels and Cancer. *EMBO Rep* **9,** 512-515.

Fulgenzi G, Graciotti L, Faronato M, Soldovieri MV, Miceli F, Amoroso S, Annunziato L, Procopio A & Taglialatela M. (2006). Human neoplastic mesothelial cells express voltage-gated sodium channels involved in cell motility. *Int J Biochem Cell Biol* **38,** 1146-1159.

Gatenby RA & Gillies RJ. (2004). Why do cancers have high aerobic glycolysis? *Nat Rev Cancer* **4,** 891-899.

Gillet L, Roger S, Besson P, Lecaille F, Gore J, Bougnoux P, Lalmanach G & Le Guennec JY. (2009). Voltage-gated Sodium Channel Activity Promotes Cysteine Cathepsin-dependent Invasiveness and Colony Growth of Human Cancer Cells. *J Biol Chem* **284,** 8680-8691.

Gradek F, Lopez-Charcas O, Chadet S, Poisson L, Ouldamer L, Goupille C, Jourdan ML, Chevalier S, Moussata D, Besson P & Roger S. (2019). Sodium Channel Nav1.5 Controls Epithelial-to-Mesenchymal Transition and Invasiveness in Breast Cancer Cells Through its Regulation by the Salt-Inducible Kinase-1. *Sci Rep* **9,** 18652.

Grillon E, Farion R, Fablet K, De Waard M, Tse CM, Donowitz M, Remy C & Coles JA. (2011). The spatial organization of proton and lactate transport in a rat brain tumor. *PLoS One* **6,** e17416.

Grillon E, Farion R, Reuveni M, Glidle A, Remy C & Coles JA. (2015). Spatial profiles of markers of glycolysis, mitochondria, and proton pumps in a rat glioma suggest coordinated programming for proliferation. *BMC Res Notes* **8,** 207.

Harvey JM, Clark GM, Osborne CK & Allred DC. (1999). Estrogen receptor status by immunohistochemistry is superior to the ligand-binding assay for predicting response to adjuvant endocrine therapy in breast cancer. *J Clin Oncol* **17,** 1474-1481.

Haworth AS, Hodges SL, Capatina AL, Isom LL, Baumann CG & Brackenbury WJ. (2022). Subcellular dynamics and functional activity of the cleaved intracellular domain of the Na(+) channel beta1 subunit. *J Biol Chem* **298,** 102174.

Helmlinger G, Yuan F, Dellian M & Jain RK. (1997). Interstitial pH and pO2 gradients in solid tumors in vivo: high-resolution measurements reveal a lack of correlation. *Nat Med* **3,** 177-182.

Hernandez-Plata E, Ortiz CS, Marquina-Castillo B, Medina-Martinez I, Alfaro A, Berumen J, Rivera M & Gomora JC. (2012). Overexpression of Na(V) 1.6 channels is associated with the invasion capacity of human cervical cancer. *Int J Cancer* **130,** 2013-2023.

Hochachka PW & Mommsen TP. (1983). Protons and anaerobiosis. *Science* **219,** 1391-1397.

Holding AN, Giorgi FM, Donnelly A, Cullen AE, Nagarajan S, Selth LA & Markowetz F. (2019). VULCAN integrates ChIP-seq with patient-derived co-expression networks to identify GRHL2 as a key co-regulator of ERa at enhancers in breast cancer. *Genome Biol* **20,** 91.

House CD, Vaske CJ, Schwartz A, Obias V, Frank B, Luu T, Sarvazyan N, Irby RB, Strausberg RL, Hales T, Stuart J & Lee NH. (2010). Voltage-gated Na+ channel SCN5A is a key regulator of a gene transcriptional network that controls colon cancer invasion. *Cancer Res* **70,** 6957-6967.

House CD, Wang BD, Ceniccola K, Williams R, Simaan M, Olender J, Patel V, Baptista-Hon DT, Annunziata CM, Gutkind JS, Hales TG & Lee NH. (2015). Voltage-gated Na+ Channel Activity Increases Colon Cancer Transcriptional Activity and Invasion Via Persistent MAPK Signaling. *Sci Rep* **5,** 11541.

James AD, Leslie TK, Kaggie JD, Wiggins L, Patten L, Murphy O'Duinn J, Langer S, Labarthe MC, Riemer F, Baxter G, McLean MA, Gilbert FJ, Kennerley AJ & Brackenbury WJ. (2022). Sodium accumulation in breast cancer predicts malignancy and treatment response. *Br J Cancer* **127,** 337-349.

James AD, Patel W, Butt Z, Adiamah M, Dakhel R, Latif A, Uggenti C, Swanton E, Imamura H, Siriwardena AK & Bruce JI. (2015). The Plasma Membrane Calcium Pump in Pancreatic Cancer Cells Exhibiting the Warburg Effect Relies on Glycolytic ATP. *J Biol Chem* **290,** 24760-24771.

James AD, Unthank KP, Jones I, Sajjaboontawee N, Sizer RE, Chawla S, Evans GJO & Brackenbury WJ. (2023). Sodium regulates PLC and IP(3) R-mediated calcium signaling in invasive breast cancer cells. *Physiol Rep* **11,** e15663.

James JH, Fang CH, Schrantz SJ, Hasselgren PO, Paul RJ & Fischer JE. (1996). Linkage of aerobic glycolysis to sodium-potassium transport in rat skeletal muscle. Implications for increased muscle lactate production in sepsis. *J Clin Invest* **98,** 2388-2397.

Jardim-Perassi BV, Huang S, Dominguez-Viqueira W, Poleszczuk J, Budzevich MM, Abdalah MA, Pillai SR, Ruiz E, Bui MM, Zuccari D, Gillies RJ & Martinez GV. (2019). Multiparametric MRI and Coregistered Histology Identify Tumor Habitats in Breast Cancer Mouse Models. *Cancer Res* **79,** 3952-3964.

Jones DK, Peters CH, Tolhurst SA, Claydon TW & Ruben PC. (2011). Extracellular proton modulation of the cardiac voltage-gated sodium channel, Nav1.5. *Biophys J* **101,** 2147-2156.

Khan A, Kyle JW, Hanck DA, Lipkind GM & Fozzard HA. (2006). Isoform-dependent interaction of voltage-gated sodium channels with protons. *J Physiol* **576,** 493-501.

Leslie TK & Brackenbury WJ. (2022). Sodium channels and the ionic microenvironment of breast tumours. *J Physiol*.

Leslie TK, Bruckner L, Chawla S & Brackenbury WJ. (2020). Inhibitory Effect of Eslicarbazepine Acetate and S-Licarbazepine on Nav1.5 Channels. *Front Pharmacol* **11,** 555047.

Liao Y, Smyth GK & Shi W. (2019). The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res* **47,** e47.

Lopez-Charcas O, Poisson L, Benouna O, Lemoine R, Chadet S, Petereau A, Lahlou W, Guyetant S, Ouaissi M, Pukkanasut P, Dutta S, Velu SE, Besson P,

Moussata D & Roger S. (2022). Voltage-Gated Sodium Channel Na(V)1.5 Controls NHE-1-Dependent Invasive Properties in Colon Cancer Cells. *Cancers (Basel)* **15**.

Lopez-Charcas O, Pukkanasut P, Velu SE, Brackenbury WJ, Hales TG, Besson P, Gomora JC & Roger S. (2021). Pharmacological and nutritional targeting of voltage-gated sodium channels in the treatment of cancers. *iScience* **24**, 102270.

Love MI, Huber W & Anders S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550.

Masters JR, Thomson JA, Daly-Burns B, Reid YA, Dirks WG, Packer P, Toji LH, Ohno T, Tanabe H, Arlett CF, Kelland LR, Harrison M, Virmani A, Ward TH, Ayres KL & Debenham PG. (2001). Short tandem repeat profiling provides an international reference standard for human cell lines. *Proc Natl Acad Sci U S A* **98**, 8012-8017.

McBride BW & Early RJ. (1989). Energy expenditure associated with sodium/potassium transport and protein synthesis in skeletal muscle and isolated hepatocytes from hyperthyroid sheep. *Br J Nutr* **62**, 673-682.

Mohammed FH, Khajah MA, Yang M, Brackenbury WJ & Luqmani YA. (2016). Blockade of voltage-gated sodium channels inhibits invasion of endocrine-resistant breast cancer cells. *Int J Oncol* **48**, 73-83.

Nelson M, Millican-Slater R, Forrest LC & Brackenbury WJ. (2014). The sodium channel beta1 subunit mediates outgrowth of neurite-like processes on breast cancer cells and promotes tumour growth and metastasis. *Int J Cancer* **135**, 2338-2351.

Nelson M, Yang M, Dowle AA, Thomas JR & Brackenbury WJ. (2015a). The sodium channel-blocking antiepileptic drug phenytoin inhibits breast tumour growth and metastasis. *Mol Cancer* **14**, 13.

Nelson M, Yang M, Millican-Slater R & Brackenbury WJ. (2015b). Nav1.5 regulates breast tumor growth and metastatic dissemination in vivo. *Oncotarget* **6**, 32914-32929.

Nyblom M, Poulsen H, Gourdon P, Reinhard L, Andersson M, Lindahl E, Fedosova N & Nissen P. (2013). Crystal structure of Na+, K(+)-ATPase in the Na(+)-bound state. *Science* **342**, 123-127.

Ouwerkerk R, Jacobs MA, Macura KJ, Wolff AC, Stearns V, Mezban SD, Khouri NF, Bluemke DA & Bottomley PA. (2007). Elevated tissue sodium concentration in malignant breast lesions detected with non-invasive 23Na MRI. *Breast Cancer Res Treat* **106**, 151-160.

Paul RJ, Bauer M & Pease W. (1979). Vascular smooth muscle: aerobic glycolysis linked to sodium and potassium transport processes. *Science* **206**, 1414-1416.

Pellerin L & Magistretti PJ. (1994). Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization. *Proc Natl Acad Sci U S A* **91**, 10625-10629.

Post RL & Jolly PC. (1957). The linkage of sodium, potassium, and ammonium active transport across the human erythrocyte membrane. *Biochim Biophys Acta* **25,** 118-128.

Rahrmann EP, Shorthouse D, Jassim A, Hu LP, Ortiz M, Mahler-Araujo B, Vogel P, Paez-Ribes M, Fatemi A, Hannon GJ, Iyer R, Blundon JA, Lourenco FC, Kay J, Nazarian RM, Hall BA, Zakharenko SS, Winton DJ, Zhu L & Gilbertson RJ. (2022). The NALCN channel regulates metastasis and nonmalignant cell dissemination. *Nat Genet* **54,** 1827-1838.

Roberts GC, Morris PG, Moss MA, Maltby SL, Palmer CA, Nash CE, Smart E, Holliday DL & Speirs V. (2016). An Evaluation of Matrix-Containing and Humanised Matrix-Free 3-Dimensional Cell Culture Systems for Studying Breast Cancer. *PLoS One* **11,** e0157004.

Roger S, Besson P & Le Guennec JY. (2003). Involvement of a novel fast inward sodium current in the invasion capacity of a breast cancer cell line. *Biochim Biophys Acta* **1616,** 107-111.

Schuetz CS, Bonin M, Clare SE, Nieselt K, Sotlar K, Walter M, Fehm T, Solomayer E, Riess O, Wallwiener D, Kurek R & Neubauer HJ. (2006). Progression-specific genes identified by expression profiling of matched ductal carcinomas in situ and invasive breast tumors, combining laser capture microdissection and oligonucleotide microarray analysis. *Cancer Res* **66,** 5278-5286.

Sepp M, Sokolova N, Jugai S, Mandel M, Peterson P & Vendelin M. (2014). Tight coupling of Na+/K+-ATPase with glycolysis demonstrated in permeabilized rat cardiomyocytes. *PLoS One* **9,** e99413.

Sjogaard-Frich LM, Prestel A, Pedersen ES, Severin M, Kristensen KK, Olsen JG, Kragelund BB & Pedersen SF. (2021). Dynamic Na(+)/H(+) exchanger 1 (NHE1) - calmodulin complexes of varying stoichiometry and structure regulate Ca(2+)-dependent NHE1 activation. *Elife* **10,** e60889.

Skou JC. (1998). The influence of some cations on an adenosine triphosphatase from peripheral nerves. *J Am Soc Nephrol* **9,** 2170-2177.

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES & Mesirov JP. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102,** 15545-15550.

Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A & Bray F. (2021). Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin* **71,** 209-249.

Uphoff CC, Gignac SM & Drexler HG. (1992). Mycoplasma contamination in human leukemia cell lines. I. Comparison of various detection methods. *J Immunol Methods* **149,** 43-53.

van Weverwijk A, Koundouros N, Iravani M, Ashenden M, Gao Q, Poulogiannis G, Jungwirth U & Isacke CM. (2019). Metabolic adaptability in metastatic breast

cancer by AKR1B10-dependent balancing of glycolysis and fatty acid oxidation. *Nat Commun* **10,** 2698.

Voipio J, Pasternack M & MacLeod K. (1994). Ion-sensitive microelectrodes. . In *Microelectrode Techniques; the Plymouth Workshop Handbook, 2nd edn*, ed. Ogden D, pp. 275-316. The Company of Biologists Ltd, Cambridge.

White KA, Grillo-Hill BK & Barber DL. (2017). Cancer cell behaviors mediated by dysregulated pH dynamics at a glance. *J Cell Sci* **130,** 663-669.

Yamaci RF, Fraser SP, Battaloglu E, Kaya H, Erguler K, Foster CS & Djamgoz MBA. (2017). Neonatal Nav1.5 protein expression in normal adult human tissues and breast cancer. *Pathol Res Pract* **213,** 900-907.

Yang M, James AD, Suman R, Kasprowicz R, Nelson M, O'Toole PJ & Brackenbury WJ. (2020). Voltage-dependent activation of Rac1 by Nav 1.5 channels promotes cell migration. *J Cell Physiol* **235,** 3950-3972.

Yang M, Kozminski DJ, Wold LA, Modak R, Calhoun JD, Isom LL & Brackenbury WJ. (2012). Therapeutic potential for phenytoin: targeting Na(v)1.5 sodium channels to reduce migration and invasion in metastatic breast cancer. *Breast Cancer Res Treat* **134,** 603-615.

Table 1. Effect of reduced pH_e on VGSC Na^+ current parameters.

Parameter	pH 7.2	pH 6.2	P	n
Peak current density (pA/pF)	-13.5 ± 2.3	-9.6 ± 1.5	0.006	11
Persistent current density (pA/pF)	-0.31 ± 0.04	-0.71 ± 0.11	0.006	10
Activation V½ (mV)	-15.2 ± 2.0	-12.9 ± 1.8	0.087	10
Activation k (mV)	10.7 ± 0.5	9.5 ± 0.4	0.077	10
Inactivation V½ (mV)	-80.4 ± 1.4	-73.3 ± 2.8	0.003	10
Inactivation k (mV)	-8.4 ± 0.8	-11.9 ± 0.9	0.005	10
T_p at -10 mV (ms)	2.07 ± 0.24	3.02 ± 0.35	< 0.001	10

V½: half (in)activation voltage; k: slope factor for (in)activation; T_p: time to peak current; The holding potential was -120 mV. Results are mean ± SEM. Statistical comparisons were made with paired *t* tests on non-normalised data.

Table 2. Clinical characteristics and membrane current expression in primary cells.

I Inward currents

Variable	No inward I, n (%)[1]	Inward I, n (%)	P
Cancer vs normal			
Normal	28/28 (100)	0/28 (0)	0.32
Cancer	64/68 (94.1)	4/68 (5.9)	
Subtype			
ER+/HER2+	38/42 (90.5)	4/42 (9.5)	0.29
TNBC	26/26 (100)	0/26 (0)	
Lymph node status			
Negative	27/27 (100)	0/27 (0)	0.16
Positive	37/41 (90.2)	4/41 (9.8)	

II Outward currents

Variable	No outward I, n (%)[1]	Outward I, n (%)	P
Cancer vs normal			
Normal	9/17 (52.9)	8/17 (47.1)	0.60
Cancer	31/68 (45.6)	37/68 (54.4)	
Subtype			
ER+/HER2+	13/42 (31.0)	29/42 (69.0)	<0.05
TNBC	18/26 (69.2)	8/26 (30.8)	
Lymph node status			
Negative	18 (66.7)	9 (33.3)	<0.01
Positive	13 (31.7)	28 (68.3)	

[1] n = cells recorded. Results are from Fisher's exact tests.

Table 3. Patient histoclinical characteristics and Na$_v$1.5 expression.

Variable	Na$_v$1.5 expression		P
	Low (%)	High (%)	
All cases			
ER			
-	24 (1.6)	348 (23.5)	< 0.05
+	113 (7.6)	997 (67.3)	
PR			
-	34 (2.4)	539 (38.2)	< 0.001
+	94 (6.7)	744 (52.7)	
HER2			
-	123 (8.8)	1106 (78.7)	< 0.01
+	7 (0.5)	170 (12.1)	
TNBC			
-	113 (7.9)	1065 (74.4)	0.12
+	16 (1.1)	238 (16.6)	
Age			
≤ 50	58 (3.9)	530 (35.9)	0.52
> 50	78 (5.3)	811 (54.9)	
Menopause status			
Pre-menopause	58 (4.0)	515 (35.2)	0.46
Post-menopause	79 (5.4)	813 (55.5)	
Lymph node status			
-	95 (6.5)	807 (54.9)	< 0.05
+	42 (2.9)	527 (35.8)	
Grade			
1	42 (2.9)	209 (14.3)	< 0.001
2	39 (2.7)	447 (30.5)	
3	54 (3.7)	674 (46.0)	
Chemotherapy			
-	115 (8.1)	1026 (72.4)	< 0.05
+	16 (1.1)	260 (18.4)	
Endocrine therapy			
-	83 (5.9)	765 (54.0)	0.51
+	49 (3.5)	520 (36.7)	

P values are from Fisher's exact tests except for grade, which is from χ^2 tests.

Table 4. Cox multivariate analysis of cancer-specific survival.

Variable	Hazard ratio (95% CI)	P value
Tumour size		
≤20 mm	1	
>20 mm	1.16 (1.10-1.22)	<0.001
Grade		
1	1	
2	1.46 (1.01-2.11)	<0.05
3	2.62 (1.86-3.70)	<0.001
Lymph node status		
Negative	1	
Positive	1.65 (1.36-2.00)	<0.001
$Na_v1.5$ expression		
Low	1	
High	1.58 (1.05-2.37)	<0.05

Table 5. Correlation between $Na_v1.5$ and other protein markers.

Protein marker	Spearman's correlation coefficient	P value	n
ALDH1A1	-0.05	0.265	579
AMPK	-0.15	< 0.001	1214
Carbonic anhydrase IX	-0.07	0.111	479
Cathepsin D	0.11	0.003	762
CD44	-0.13	0.001	701
CD56	-0.19	< 0.001	372
HIF-1	-0.05	0.576	130
IDH2	0.13	< 0.001	715
Kank1M	0.08	0.069	546
Lactate dehydrogenase 5	0.00	0.942	434
MMP11	-0.02	0.849	117
MMP13 (cytoplasmic)	-0.02	0.475	977
MMP13 (nuclear)	-0.20	< 0.001	977
MMP14 (cytoplasmic)	-0.07	0.033	868
MMP14 (membrane)	-0.09	0.009	867
MMP14 (nuclear)	0.01	0.754	867
MMP15 (cytoplasmic)	-0.05	0.189	842
MMP-15 (membrane)	0.19	< 0.001	842
MMP15 (nuclear)	0.19	< 0.001	842
MMP2 (cytoplasmic)	-0.20	< 0.001	847
MMP2 (nuclear)	-0.10	0.003	847
MMP7	-0.15	< 0.001	866
MMP9	0.11	0.007	599
MX1	0.05	0.157	741
NHERF1	0.07	0.214	321
SNAT2	-0.12	0.005	593

Figure 1. Characterisation of voltage-sensitive inward currents in breast cancer cell lines and cancer-associated fibroblasts. (A) Peak inward current density measured across a panel of breast cancer cell lines and cancer-associated fibroblasts (CAFs). (B) Inward current-voltage relationship of MDA-MB-231 cells, from a holding potential of -120 mV with 5 mV depolarising steps in the range -80 mV to +30 mV (n = 9). (C) Example voltage-sensitive inward currents in an LS11-083 CAF. (D) Inward current-voltage relationship of LS11-083 CAFs, from a holding potential of -120 mV with 5 mV depolarising steps in the range -80 mV to +30 mV (n = 5). (E) Whole cell capacitance across panel of breast cancer cell lines and CAFs. (F) MDA-MB-231-β1-GFP cells were implanted into the mammary fat pad. After 4 weeks, mice were killed, the lungs were filled with low melting point agarose and the primary tumour and lungs were sliced for whole cell patch clamp recording. Phase contrast and GFP fluorescence images of microelectrode recording from a cell in a primary tumour slice and a lung slice. (G) Peak inward Na$^+$ current density recorded in primary tumours (n

= 24) and lung metastases (n = 8). Data are mean ± SEM; ns, not significant (Student's *t* test).

Figure 2. Altered human gene expression in Na$_v$1.5 knockdown MDA-MB-231 xenograft tumours compared to control MDA-MB-231 tumours. (A) Volcano plot showing transcriptome changes in Na$_v$1.5 knockdown tumours, with genes involved in invasion, and *SCN5A*, highlighted. (B) The top 20 human genes which were significantly up-regulated in Na$_v$1.5 knock-down tumours. (C) The top 20 human genes which were significantly down-regulated in Na$_v$1.5 knock-down tumours. (D) Gene set enrichment analysis (GSEA) of invasion-upregulated genes defined in the MSigDB gene set SCHUETZ_BREAST_CANCER_DUCTAL_INVASIVE_UP analysed in Na$_v$1.5 knockdown vs. control tumours (n = 6 tumours in each condition).

Figure 3. Altered extracellular pH in tissues slices from MDA-MB-231 xenograft tumours. (A) Photograph of a tumour slice showing the difference in appearance between the translucent periphery and opaque core. (B) Tumour slice extracellular pH (pH$_e$) comparing core and peripheral regions in each slice; n = 9 tumours (one slice from each). (C) Examples of peripheral and core regions with H&E staining (left), Ki67 (red, middle), and cleaved caspase 3 (red, right) with DAPI (blue). (D) Cellularity in the peripheral and core regions of each section, based on DAPI nuclear count (n = 9 tumours). (E) Ki67-positive cells (%) in peripheral and core regions (n = 9 tumours). (F) Cleaved caspase 3-positive cells (%) in peripheral and core regions (n = 9 tumours). Data are mean ± SEM; *P < 0.05, **P < 0.01 (paired *t* tests).

Figure 4. Effect of low pH_e on Na^+ current in MDA-MB-231 cells. (A) Example Na^+ currents elicited by depolarisation to 0 mV from a holding voltage of -120 mV. (B) Data from (A) expanded between 20 and 25 ms after depolarisation. (C) Peak Na^+ current density (P < 0.01; n = 11; Wilcoxon matched pairs test). (D) Mean persistent Na^+ current density measured between 20 and 25 ms after depolarisation (P < 0.01; n = 10; paired *t* test). (E)(i) Example family of Na^+ currents generated by the activation voltage clamp protocol (inset). (ii) Example family of Na^+ currents generated by the steady-state inactivation voltage clamp protocol (inset). (F) Current density-voltage relationship (n = 17). (G) Overlay of activation and inactivation curves at pH 7.2 and 6.2 (n = 10 cells with the largest currents). (H) Expanded data from (G) showing the window current. (I) Fraction of channels available at the reported resting membrane potential of -18.9 mV at pH 7.2 and 6.2 (n = 10 cells with the largest currents; *t* test). (J) Fraction of channels available at the reported resting membrane potential of -18.9 mV, at pH 7.2 and 6.0 (P < 0.001; n = 8 cells; *t* test). (K) Time to peak current (n = 10 cells). Data are mean ± SEM.

Figure 5. Effect of voltage-gated Na$^+$ channel activity on glycolytic H$^+$ production. (A) SBFI-AM fluorescence ratios of MDA-MB-231 cells after 6 h treatment with ouabain (300 nM), or 2 h treatment with sodium iodoacetate (2 mM) or oligomycin (1 µM).

Data are normalised to the ratio in vehicle-treated cells (n = 6 experimental repeats; one sample t tests). (B) SBFI-AM fluorescence ratios of MCF7 cells after 6 h treatment with ouabain (300 nM), or 2 h treatment with sodium iodoacetate (2 mM) or oligomycin (1 µM). Data are normalised to the ratio in vehicle-treated cells (n = 6 experimental repeats; one sample t tests). (C) Representative measurements of the extracellular acidification rate (ECAR) of MDA-MB-231 cells. TTX was added to the treated cells at the indicated timepoint, to give a final concentration of 30 µM (n = 6 wells). (D) ECAR of MDA-MB-231 cells compared between control and TTX (30 µM) wells. (E) Representative measurements of the ECAR of MCF7 cells. TTX was added to the treated cells at the indicated timepoint, to give a final concentration of 30 µM (n = 6 wells). (F) ECAR of MCF7 cells compared between control and TTX (30 µM) wells. (G) Representative measurements of the oxygen consumption rate (OCR) of MDA-MB-231 cells. TTX was added to the treated cells at the indicated timepoint, to give a final concentration of 30 µM (n = 6 wells). (H) OCR of MDA-MB-231 cells compared between control and TTX (30 µM) wells. (I) Representative measurements of the OCR of MCF7 cells. TTX was added to the treated cells at the indicated timepoint, to give a final concentration of 30 µM (n = 6 wells). (J) OCR of MCF7 cells compared between control and TTX (30 µM) wells. Each data point in (D), (F), (H), (J) represents the mean of the last 6 timepoints for each well (n = 3 experimental repeats each containing 6 wells; experimental repeats colour-coded black, red, blue). Data are mean ± SEM. ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$, *$P < 0.05$, ns, not significant (2-way ANOVA for D, F, H, J).

Figure 6. Characterisation of membrane currents in primary breast epithelial and cancer cells. (A) Example inward currents recorded from a mammary carcinoma cell, elicited by depolarising steps between -40 mV and -5 mV from a holding potential of -120 mV. (B) Peak inward current density recorded from two mammary carcinoma cell samples. (C) Mean inward current-voltage relationship from both mammary carcinoma cell samples (n = 4 cells). (D) Example outward currents recorded from a mammary carcinoma cell, using a holding potential of -120 mV with 10 mV depolarising steps in the range -80 mV to +30 mV. (E) Mean outward current-voltage relationship from mammary carcinoma cells (n = 30 cells). (F) Outward current density at +30 mV of normal mammary epithelial and carcinoma cells, subdivided by patient sample. (G) Whole cell capacitance of normal mammary epithelial cells and carcinoma cells. Samples colour-coded according to (E). (H) Outward current density at +30 mV of normal mammary epithelial cells and carcinoma cells. Samples colour-coded according to (E). (I) Outward current density at +30 mV for ER+/HER2+ and TNBC carcinoma cells. Samples colour-coded according to (E). Data are mean ± SEM; ***P < 0.001, ns, not significant (Mann-Whitney test).

Figure 7. Characterisation of membrane currents recorded from cells in tumour tissue slices and following dissociation. (A) Example inward and outward membrane currents recorded from a cell in a patient tumour slice, from a holding potential of -120 mV with 10 mV depolarising steps in the range -40 mV to +30 mV. (B) Mean outward current-voltage relationship including cells from all patient tumour specimens (n = 9). (C) Outward current density elicited by depolarisation from -120 mV to +30 mV, subdivided by patient tumour specimen. (D) Peak inward current density, subdivided by patient tumour specimen. (E) Mean inward current-voltage relationship including cells from both patient tumour specimens (n = 4). (F) Outward current density at +30 mV recorded from cells in patient tumour slices compared to cells which had been dissociated and plated onto glass coverslips (data from 2 patients). Samples colour coded according to (C). (G) Whole cell capacitance recorded from cells in patient tumour slices compared to cells which had been dissociated and plated onto glass coverslips. Samples colour coded according to (C). (H) Mean outward current-voltage relationship from cells in patient tumour slices compared to cells which had been dissociated and plated onto glass coverslips (in slice: n = 10 cells from 3 patients; dissociated: n = 6 cells from 2 patients). Data are mean ± SEM; ns, not significant; *P < 0.05; (Student's t tests).

Figure 8. Na$_v$1.5 expression in a breast cancer tissue microarray. (A) Examples of low and high Na$_v$1.5 staining in carcinoma cells scored using a modified Allred system. Right: staining using anti-Na$_v$1.5 antibody which had been preincubated with the immunising peptide. (B) Tumour size compared to Na$_v$1.5 score. (C) Number of affected lymph nodes compared to Na$_v$1.5 score. (D) Nottingham Prognostic Index (NPI) compared to Na$_v$1.5 score. (E) Na$_v$1.5 score in patients with or without recorded distant metastasis. Data are median + quartiles. *P < 0.05, ***P < 0.001, ****P < 0.0001; Mann-Whitney U tests (n = 1740). (F) Metastasis-free survival compared to Na$_v$1.5 score. HR = 2.18 (95% CI 1.63-2.92); P < 0.001; log-rank test.

Figure 9. Mechanism for Na$_v$1.5-mediated cellular invasion. Elevated steady-state Na$^+$ entry through Na$_v$1.5, leads to increased NKA activity. The increased ATP demand is satisfied by glycolysis, increasing H$^+$ production and extrusion through pH regulators such as CAIX and NHE1. Extracellular acidification in turn increases persistent Na$^+$ entry via Na$_v$1.5, leading to a positive feedback loop linking Na$_v$1.5, Na$^+$ entry, increased H$^+$ production via glycolysis, and extracellular acidification. This results in ECM degradation and increased invasion (Brisson *et al.*, 2011; Brisson *et al.*, 2013; Leslie & Brackenbury, 2022).