miR-252 targeting temperature receptor *CcTRPM* to mediate the transition from
summer-form to winter-form of *Cacopsylla chinensis*

Songdou Zhang, Jianying Li, Dongyue Zhang, Zhixian Zhang, Shili Meng, Zhen Li, Xiaoxia Liu[*]

Department of Entomology and MOA Key Lab of Pest Monitoring and Green Management,

College of Plant Protection, China Agricultural University, 100193 Beijing, China

[*] Corresponding author. E-mail address: liuxiaoxia611@cau.edu.cn (Xiaoxia Liu).

Abstract:

Temperature determines the geographical distribution of organisms and affects the outbreak and

damage of pests. Insects seasonal polyphenism is a successful strategy adopted by some species to

adapt the changeable external environment. *Cacopsylla chinensis* (Yang & Li) showed two

seasonal morphotypes, summer-form and winter-form, with significant differences in

morphological characteristics. Low temperature is the key environmental factor to induce its

transition from summer-form to winter-form. However, the detailed molecular mechanism

remains unknown. Here, we firstly confirmed that low temperature of 10°C induced the transition

from summer-form to winter-form by affecting the cuticle thickness and chitin content.

Subsequently, we demonstrated that *CcTRPM* functions as a temperature receptor to regulate this

transition. In addition, miR-252 was identified to mediate the expression of *CcTRPM* to involve in

this behavioral transition. Finally, we found *CcTre1* and *CcCHS1*, two rate-limiting enzymes of

insect chitin biosyntheis, act as the critical down-stream signal of *CcTRPM* in mediating this

behavioral transition. Taken together, our results revealed that a novel signal transduction cascade

mediates the seasonal polyphenism in *C. chinensis*. These findings not only lay a solid foundation

for fully clarifying the ecological adaptation mechanism of *C. chinensis* outbreak, but also

broaden our understanding about insect polymorphism.

Keywords: *Cacopsylla chinensis*, miR-252, *CcTRPM*, polyphenism, low temperature

Introduction

Currently, global climate change is the most important environmental problem facing with mankind, and also one of the most complex challenges in the 21st century. Insects, as the largest animal population on the earth, adapt to the complex and changeable external environment by displaying polyphenism, thus achieving the purpose of survival and reproduction (Simpson et al., 2011). Insect polyphenism is the phenomenon allows them within a given genotype to produce two or more distinct phenotypes under the challenges posed by environmental variability via beneficial shifts in morphology, behavior, or physiology (Cridge et al., 2015). Polyphenism is commonly observed in many insect species, for examples: the solitary and gregarious phases of *Locusta migratoria* (Ma et al., 2011), the short-winged and long-winged morph of *Nilaparvata lugens* (Xu et al., 2015), the wingless and winged aphid of *Acyrthosiphon pisum* (Vellichirammal et al., 2017), and the catkin and twig morphs in caterpillars of the moth *Nemoria arizonaria* (Simpson et al., 2011). With the global climate problems are increasingly prominent, the study of insect polyphenism has become a new hotspot in insect ecology, epigenetics and developmental biology.

Pear psylla is a serious pest of pear trees worldwide, including *Cacopsylla chinensis* (Yang & Li), *Psylla Betulaefoliae*, *Psylla Pyri*, and so on (Butt and Stuart, 1986; Wei et al., 2020). *C. chinensis*, belongs to Hemiptera: Psyllidae, which exclusively feeds on pear trees, harms the young shoots and leaves with adult and nymph, and secretes a large amount of honey dew to breed coal pollution bacteria of *Gloeodes pomigena* (Wei et al., 2020). It is also an important vector of the destructive disease of pear trees, *Erwinia amovora*, and currently one of the key pests in the major pear production areas in many East Asian countries, including China and Japan (Hildebrand et al., 2010; Ge et al., 2019). In most pear production areas of China, the summer-form of *C. chinensis* are usually present from April to September, whereas winter-form of *C. chinensis* are present during the rest of the year (Ge et al., 2019; Wei et al., 2020). The summer-form has lighter body color and stronger activity ability, and causes more serious damage compare with winter-form. For better adapt to the harsh environmental conditions, the eggs of summer-form develop to the winter-form before the pear fallen leaves. The winter-form has brown to dark brown body color and larger body size, but has the weaker activity ability and stronger resistance ability compare with summer-form. With the gradual increase of temperature in early spring, the winter-form lays eggs at the tender buds of pear trees and then develop into summer-form (Ge et al., 2019; Tougeron et al., 2021). Hence, the transition between summer-form and winter-form is a significant ecological strategy for *C. chinensis* to avoid adverse environmental conditions and then burst into disaster, which has prominence biological and ecological significance. The seasonal polyphenism of summer-form and winter-form in *C. chinensis* has been recognized for many years and could be induced by low temperature and short day (Oldfield, 1970; Tougeron et al., 2021), but until recently no molecular mechanism had been advanced.

Temperature is a significant factor limiting the geographic ranges of species, which determines the distribution and diffusion area of species. As a type of small poikilothermic animal, insect is sensitive to temperature changes and temperature stress response is one of the most conservative mechanism to resist extreme environment (Teets et al., 2023). Different environmental conditions affect the polyphenism of many kinds of insects, for examples, wing dimorphism of Hemiptera *Schizaphis graminum* (An et al., 2012), diapause of Lepidoptera

Bombyx mori (Tsuchiya et al., 2021) and Coleoptera *Colaphellus bowringi* (Guo et al., 2021).
Under the low temperature conditions, insects can improve cold tolerance by regulating their
internal energy metabolism and external exoskeleton system. Such as: darken their cuticles with
melanin (Fedorka et al., 2013), thickened wax layer or chorion (Kreß et al., 2016), and store more
lipids to increase energy supply and increase the accumulation of antifreeze protective substances
(Terblanche et al., 2011). However, studies about how insect temperature receptor response to
temperature change and then activates the downstream signal molecules to regulate the
polyphenism have been very limited.

Transient receptor potential (TRP) channel family gene was firstly found in *Drosophila*
melanogaster (Minke et al., 1975). The TRP channel genes involved in temperature sensing are
called "thermo-sensitive TRP" or "temperature receptor" (Lee et al., 2005). In mammals, there are
six thermo-sensitive TRP receptor genes, of which TRPV1-4 mainly feel moderate high
temperature (Venkatachalam and Montell, 2007) and TRPM8 can be activated by low temperature
(8°C-28°C) or cooling substances, such as menthol and borneol (Peier et al., 2002). The activation
mode is mainly due to the depolarization of cell membrane caused by the influx of Ca^{2+} (McKemy
et al., 2002). Presently, researches on the function of temperature receptor gene TRP were mainly
focused on some model insects, but very few on the agricultural pests. For example, *D.*
melanogaster TRPM is highly expressed in dendritic sensory neurons and can sense cold
stimulation (Turner et al., 2016). In *B. mori*, high temperature receptor *BmTRPA1* regulates the
diapause of offspring eggs by acting on the neuroendocrine hormone signal of
GABAergic-GnRH/Crz-DH (Sato et al., 2014; Tsuchiya et al., 2021). However, studies on
agricultural insects, such as *Apolygus lucorum* (Fu et al., 2016), *N. lugens* (Wang et al., 2021a),
Bemisia tabaci (Dai et al., 2018), and *Tuta absoluta* (Wang et al., 2021b), also only focused on the
identification of the related temperature receptor gene TRPs and its correlation with temperature,
no detailed molecular mechanism were reported. In agricultural insects, members of TRPA
subfamily are mainly responsible for answering the high-temperature stimulation (Tracey et al.,
2003; Neely et al., 2011), while members of TRPM subfamily are mainly involved in
low-temperature stimulation as temperature receptors (Rosenzweig et al., 2008; Turner et al.,
2016). Temperature significantly affected the annual generations, adult survival rate, hibernation
period and egg laying period of *C. chinensis* (Butt and Stuart, 1986). Our preliminary research
results found that low temperature is the key factor to induce the transition from summer-form to
winter-form in *C. chinensis*. Therefore, we speculated that *TRPM*, functions as a temperature
receptor gene, may participate in the transition from summer-form to winter-form in response to
low temperature, but the specific molecular mechanism needs to be further explored.

MicroRNAs (miRNAs), a class of non-coding RNA molecules, were encoded by endogenous
genes in a length of about 18-25 nt. They mainly regulate many life processes, including cell
differentiation, organ development, cell apoptosis, biological rhythm and metamorphosis
development in animals and plants at the post-transcriptional level by inhibiting the translation
process of target genes or degrading the mRNA of target genes (Asgari, 2013). In recent years,
more and more studies reported that miRNAs are involved in regulating the insect polymorphism,
such as the caste differentiation of honeybees, the phenotypic plasticity of *L. migratoria* and the
wing pleomorphism of *N. lugens* (Liu et al., 2012; Yang et al., 2014; Ye et al., 2019). In addition,
miRNAs also play important roles in insect adaptation to temperature stress. For examples, miR-7
and miR-8 involved in the process of temperature adaptation of *D. melanogaster* (Kennell et al.,

2012). MiRNAs were also found in responding to low temperature stress in rice water weevil (*Lissorhoptrus oryzophilus*) and gall midge (*Epiblema Scudderiana*) (Yang et al., 2017a; Lyons et al., 2015). There were also reported that miR-204 targeted *TRPM3* to regulate mammalian eye development (Shiels, 2020), and miR-135a targeted *TRPC1* to participate in podocyte injury (Yang et al., 2017b). Here, we screened and validated miR-252, a miRNA whose expression was down-regulated under low temperature, negatively targeted *CcTRPM* to modulate the transition from summer-form to winter-form by acting on the cuticle thickness and cuticle chitin content. Chitin biosynthesis signal was also proved to involve in the transition from summer-form to winter-summer in response to *CcTRPM* and miR-252. Our work not only promotes the comprehensive understanding of the adaptation mechanism for *C. chinensis* outbreak at the physiological and ecological level as well as the insect polymorphism, but also helpful in identifying potential target genes in pest control through blocking temperature adaptation of pests.

Results

Low temperature induced the transition from summer-form to winter-form in *C. chinensis*

C. chinensis was displayed as seasonal polyphenism along with seasonal changes: the summer-form and the winter-form. The significant differences in morphological characteristics are as following: (1) Eggs are long oval shape with a thin tip and a blunt end. Eggs of summer-form are milky white at the initial stage and gradually become yellow white at the later stage (Figure 1A). Eggs of winter-form are mostly yellowish brown or orange (Figure 1A'). (2) The nymph has five instars and its body size is gradually increasing with age. The body of the newly hatched nymph is oval and the compound eyes are red. The 1st instar nymph of summer-form is light yellow and transparent (Figure 1B). The 3rd and 5th instar nymphs of summer-form are yellow-green or blue-yellow, and the wing buds are oblong, protruding and sticking to both sides of the body (Figure 1C and 1D). The 1st instar nymph of winter-form has many black stripes on the chest back plate, and the end of the abdomen and back are black (Figure 1B'). The 3rd and 5th instar nymphs of winter-form are overall black brown, with dark brown stripe on the left and right sides of each section on the chest and back of the abdomen, and the end of the abdomen and back becomes a large piece with black brown color (Figure 1C' and 1D'). (3) The antennae of adults are filamentous with 10 nodes, and the front wings are translucent and oblong. The body color of summer-form adult is green or yellow-green with yellow-brown stripes on the shield of the chest and back plate, and no stripes on the wings (Figure 1E). The body color of winter-form adult is brown or dark brown, with dark brown stripes on the whole body, and clearly brown stripes on the rear edges of the front wings (Figure 1E').

To clarify the effect of temperatures on the transition from summer-form to winter-form, the newly-hatched 1st instar summer-form nymphs were treated with four different temperatures (25°C, 20°C, 15°C, and 10°C). The results indicated that all the survived treated nymphs develop to the 3rd instar summer-form nymphs at 25°C, approximately 17% and 40% of survived treated nymphs develop to the 3rd instar winter-form nymphs at 20°C and 15°C, but all the survived treated nymphs develop to the 3rd instar winter-form nymphs at 10°C (Figure 1F). It meant that low temperature (especially 10°C) was the key environment factor to induce the transition from summer-form to winter-form and 10°C was selected for the following study. In addition, two cooling agents, menthol and borneol, were also used to mimic the low temperature effect of 10°C.

We observed that about 49% and 37% of the survived treated nymphs develop to the 3rd instar
winter-form nymphs after menthol and borneol treatment at 25°C, respectively (Figure 1G).
To further explore the differences between summer-form and winter-form nymphs, cuticle
thicknesses and cuticle chitin content were determined. The ultrastructure results of transmission
electron microscopy assays showed that cuticle thicknesses of 3rd instar winter-form (about 3.86
μm) were obviously higher than 3rd instar summer-form (about 1.82 μm) (Figure 1H and 1I). We
also found that winter-form nymphs exhibited higher chitin level in cuticle compared to the
summer-form nymphs by using the WGA-FITC staining and chitin content determination (Figure
1J and 1K). Taken together, these findings demonstrated that low temperature of 10°C induced the
transition of summer-form to winter-form by affecting the cuticle thicknesses and cuticle chitin
content.
167
CcTRPM* is a conventional cold-sensing TRP channel activated by 10°C in *C. chinensis
A cDNA fragment, which was putatively encoded a cold-sensing TRPM channel gene, was
identified as *CcTRPM* from *C. chinensis*. The full-length sequence of *CcTRPM* was amplified and
verified by RT-PCR with specific primers (Table S1) and sequencing. Sequence analysis
displayed that the ORF of *CcTRPM* was 4566 bp with a 455 bp 3'UTR and encoded a deduced
polypeptide of 1521 amino acids with six transmembrane domains. Multiple alignment analysis of
the amino acid sequence indicated that the transmembrane domains of *CcTRPM* shared very high
amino acid identity with TRPM sequences from other five selected insect species (Figure S1A).
As shown in Figure S1B, phylogenetic analysis revealed that *CcTRPM* was most closely related to
the *DcTRPM* homologue (*Diaphorina citri*, XP_017299512.2), which was consistent with the
evolutionary relationships predicted from the multiple alignment of amino acid sequences. The
potential tertiary protein structure of *CcTRPM* was constructed with the online server Phyre^2 and
modified with PyMOL-v1.3r1 software, and the conserved ankyrin repeats (ANK) and six
transmembrane domains were identified (Figure 2A).
The results of temperature treatment indicated that the mRNA expression of *CcTRPM* was
obviously induced by 10°C treatment at 3, 6, and 10 days compared with 25°C treatment by
qRT-PCR (Figure 2B). It is well known that the TRPM subfamily channels in vertebrates are
activated by a variety of chemical compounds, including menthol (Peier et al., 2002). We
therefore attempted to identify chemical compounds that either activate or inhibit *CcTRPM*.
qRT-PCR results revealed that menthol treatment with all three concentrations significantly
increased the mRNA expression of *CcTRPM* and antagonist (HY-112430, MedChemExpress,
Shanghai, China) treatment with all three concentrations markedly reduced *CcTRPM* transcription
(Figure S2F-S2G). To further confirm *CcTRPM* is a cold-sensing TRP channel gene, fluorescence
detection of Fluo-4 AM and Ca^{2+} imaging was performed to determine whether
CcTRPM-expressing cells increased intracellular Ca^{2+} concentration upon menthol treatment. The
results of fluorescence detection of Fluo-4 AM indicated that there was no change of Ca^{2+}
concentration in CcTRPM-expressing cells was treated with DMSO, but a clear increase of Ca^{2+}
concentration was observed when treated with menthol which was similar with the trend of
pcDNA-expressing cells was treated with the positive control of ionomycin (Figure 2C). The
effect of menthol on *CcTRPM* was further tested using Ca^{2+} imaging and showed that menthol
obviously elevated intracellular Ca^{2+} levels in CcTRPM-expressing cells (Figure 2D). We
therefore concluded that *CcTRPM* is a cold-sensitive ion channel activated by 10°C low

temperature and menthol.

CcTRPM modulated the transition from summer-form to winter-form of *C. chinensis* in response to low temperature

204   Temporal expression profiles by qRT-PCR revealed that *CcTRPM* mRNA expressed in all determined stages, with the highest expression level in the 1st instar nymph, and showed a decreased trend from 2nd instar nymph to adult 1 day (Figure 2E). The FISH results also indicated that *CcTRPM* mRNA has a greatest expression level in the 1st instar nymph and relative lower expression in the 3rd and 5th instar nymph (Figure 2F).

209   To study whether *CcTRPM* modulated the transition from summer-form to winter-form of *C. chinensis*, the newly-hatched 1st instar nymphs of summer-form were fed with dsCcTRPM or dsEGFP. For the RNAi efficiency under 10°C condition, feeding of dsCcTRPM reduced the transcription level of *CcTRPM* by about 61% and 62% compare to dsEGFP feeding at 3 d and 6 d by qRT-PCR (Figure S3A), and decreased the CcTRPM-Cy3 signal at 6 d by FISH (Figure S3B). Both results of cuticle ultrastructure and cuticle chitin content determination showed that cuticle thicknesses (about 1.97 μm) and cuticle chitin content of dsCcTRPM-treated nymphs were markedly lower than dsEGFP-treated nymphs (about 3.71 μm) under 10°C condition (Figure 2G and 2H). Additionally, dsCcTRPM feeding and *CcTRPM* antagonist treatment both significantly decreased the transition percent from summer-form to winter-form, but menthol feeding only partially rescued the transition percent and morphological phenotype under dsCcTRPM feeding (Figure 2I and 2J). To exclude the effect of *CcTRPM* itself on the transition percent and nymph development, we also performed the RNAi assays under 25°C condition. After successfully interfered the mRNA expression of *CcTRPM*, the nymph development and morphological phenotype were not affected compared to the dsEGFP treatment (Figure S3C-S3D). Together, these findings indicated that *CcTRPM* functions as a molecular switch to modulate the transition from summer-form to winter-form of *C. chinensis* in response to low temperature.

Confirming of miR-252 targeted *CcTRPM*

228   To explore the regulation mechanism of *CcTRPM* at the post-transcriptional level, two software programs of miRanda and Targetscan were used to predict the potential miRNAs. There are four miRNAs (miR-252, miR-5728, PC-3p-173924, and PC-3p-137568) were predicted to have the corresponding target-binding sites with the 3'UTR of *CcTRPM* (Figure 3A and S4A). miR-252 showed the opposite expression patterns with *CcTRPM* after 25°C and 10°C treatments (Figure 2B and 3B). To further determine whether miR-252 specifically targeted *CcTRPM*, the following four experimental approaches were performed: (1) The dual-luciferase reporter assays were carried out by cloning a 419 bp 3'UTR fragment of *CcTRPM* containing the predicted target binding sites of the miRNAs. The relative luciferase activities were obviously decreased when the agomir-252 and CcTRPM-3'UTR-pmirGLO were cotransfected, while the activity recovered to the control when using the mutated 3'UTR sequence without the binding sites of miR-252 (Figure 3C). However, the relative luciferase activities were no change when the construct of CcTRPM-3'UTR-pmirGLO was cotransfected with the agomir of other three miRNAs, respectively (Figure S4B). (2) An RNA immunoprecipitation assay indicated that the transcripts of *CcTRPM* and miR-252 increased 6.35-fold and 9.21-fold from the agomir-252 feeding nymphs of the Ago-1 antibody-mediated RNA complex compared to an IgG control, separately (Figure 3D

and S5). (3) FISH results showed that *CcTRPM* and miR-252 were co-expressed in the nymphs and displayed an opposite expression trend along with the increasing instar (Figure 3E). (4) qRT-PCR results indicated that agomir-252 feeding significantly inhibited *CcTRPM* mRNA expression and antagomir-252 feeding markedly increased *CcTRPM* mRNA expression compare to their controls, separately (Figure 3F). These data together confirmed that miR-252 directly targeted *CcTRPM*.

miR-252 mediated the transition from summer-form to winter-form in *C. chinensis* by suppressing *CcTRPM*

To test the efficiency of agomir-252, 1st instar nymphs of summer-form were fed with agomir-252 under 10°C condition. qRT-PCR results showed that miR-252 expression was both significantly increased after agomir-252 treatment at 3 d and 6 d (Figure 4A). To further clarify the relationship between *CcTRPM* and miR-252, we determined miR-252 expression after dsCcTRPM treatment under 10°C condition. Interestingly, miR-252 expression was dramatically inhibited after successful interference of *CcTRPM* at 3 d and 6 d (Figure 4B). This result implied that up-regulation of *CcTRPM* expression may have a positive feedback to miR-252 expression in response to 10°C temperature.

To deeply decipher the role of miR-252 in the transition from summer-form to winter-form, 1st instar nymphs of summer-form were fed with agomir-NC or agomir-252 under 10°C condition. Results of cuticle ultrastructure indicated that cuticle thickness of agomir-252 treated nymphs (about 1.85 μm) was markedly lower than agomir-NC treated nymphs (about 3.70 μm) (Figure 4C and 4D). Both WAG-FITC staining and chitin content determination revealed that cuticle chitin content of agomir-252 treated nymphs were obviously lower than agomir-NC treated nymphs (Figure 4E and 4F). Moreover, agomir-252 feeding significantly decreased the transition percent from summer-form to winter-form, but antagomir-252 feeding rescued the defect of transition percent and morphological phenotype under dsCcTRPM treatment (Figure 4G and 4H). To further evaluate the effect of agomir-252 under 25°C condition, we determined *CcTRPM* expression and observed the nymph development after agomir-252 feeding. As expected, *CcTRPM* expression was still significantly decreased after agomir-252 feeding (Figure S6C), but the nymph development and morphological phenotype were not affected by agomir-252 treatment compared to the agomir-NC treatment (Figure S6D). Above all, these results displayed that miR-252 mediated the transition from summer-form to winter-form in *C. chinensis* by suppressing *CcTRPM* via affecting the cuticle thickness and cuticle chitin content.

Chitin biosynthesis signaling involved in the transition from summer-form to winter-summer of *C. chinensis* in response to low temperature and *CcTRPM*

Since cuticle thickness and cuticle chitin content were obviously affected by dsCcTRPM treatment and agomir-252 treatment, we speculated that chitin biosyntheis signaling pathway participated in the transition from summer-form to winter-form. To demonstrate this hypothesis, a total of 14 transcripts corresponding to 8 genes involved in chitin biosyntheis were identified in *C. chinensis*, with several genes were present in two or more paralogous forms (Figure 5A). Next, we used qRT-PCR to determine the expression profiles of these 14 chitin biosynthesis transcripts after 25°C and 10°C treatments at 3 d, 6 d, and 10 d. Based on the relative gene expression in the heat map, we found that the expression of 13 transcripts exhibited overall up-regulation under 10°C

condition at 3 d, 6 d, and 10 d compared to those under 25°C condition, while levels for CcUAP2 were almost unchanged (Figure 5B).

Subsequently, we used RNAi to further uncover the impact of chitin biosyntheis genes in the transition from summer-form to winter-form. Compared with dsEGFP feeding control, only knockdown of *CcTre1* and *CcCHS1* obviously decreased the transition percent from summer-form to winter-form, but not by knockdown of the remaining transcripts (Figure 5C). To further dissect the roles of *CcTre1* and *CcCHS1*, cuticle thickness and cuticle chitin content were also determined. Compared with dsEGFP feeding control, knockdown of *CcTre1* and *CcCHS1* markedly reduced the cuticle thickness and cuticle chitin content (Figure S7C-S7D, 5D-5F), which were similar with dsCcTRPM feeding and agomir-252 treatment. To confirm whether *CcTre1* and *CcCHS1* function as the downstream of *CcTRPM* and miR-252, we next determine the expression of *CcTre1* and *CcCHS1* after *CcTRPM* knockdown and agomir-252 feeding. qRT-PCR results indicated that *CcTRPM* knockdown and agomir-252 feeding were both obviously reduced the mRNA expression of *CcTre1* and *CcCHS1* at 3 d and 6 d, respectively (Figure 5G-5J). These phenotypic and molecular data suggested that chitin biosynthesis signaling involved in the transition from summer-form to winter-summer of *C. chinensis* in response to low temperature and *CcTRPM*.

Discussion

Polyphenism offers the opportunity for insects to deploy different phenotypes generated from the same genome to best suit the extreme environmental changes (Simpson et al., 2011). In recent years, studies on insect polyphenism have mainly focused on the molecular regulation by endocrine hormones, neuropeptides, and neurotransmitters, while the regulation mechanism by external environmental factors remains poorly understood. For examples, ecdysone signaling and insulin/IGF signaling pathway regulated the wing polyphenism of pea aphid (Vellichirammal et al., 2017) or planthoppers (Zhang et al., 2022), neuropeptide F was essential for shaping locomotor plasticity in locust (Hou et al., 2017), and dopamine metabolic pathway modulated the behavioral phase changes of the migratory locust (Ma et al., 2011). Temperature is important in defining the distribution of insects, but how it activates the temperature receptor and then regulates polyphenism need further deeply study. Here, we firstly demonstrated that the transition from summer-form to winter-form in *C. chinensis* was "temperature-dependent pattern", and low temperature of 10°C completely induced this transition. Just like people wear more clothes in the winter, we also proved that 10°C induced the summer-form change to winter-form by increasing the cuticle thickness and cuticle chitin content. Several similar phenotype changes have been reported in other insects: the cricket of *Allonemobius socius* exhibited darker cuticles to improve thermo-regulation in colder environments (Fedorka et al., 2013), the mosquito of *Aedes albopictus* increased the thickness of the egg middle serosa cuticle and wax layer as an adaptation strategy for cold acclimation (Kreß et al., 2016).

TRPM8, a member of the TRP family, has been reported as the principal sensor of environmental cold in mammals (Peier et al., 2002). As a kind of poikilothermic animals, insects also have an evolutionarily conserved TRPM gene which is homologous to mammalian temperature receptor of TRPM8 (Venkatachalam and Montell, 2007). In this study, we identified a classical TRPM gene in *C. chinensis* which was designated as *CcTRPM* and functioned as a low temperature receptor by using qRT-PCR and calcium imaging. After knockdown of *CcTRPM* by

feeding dsCcTRPM, the treated nymphs displayed thinner cuticle thickness and less cuticle chitin content compared with the control of dsEGFP treatment under 10°C condition. In addition, *CcTRPM* knockdown by RNAi or *CcTRPM* antagonist feeding both caused obvious defect of the transition percent from summer-form to winter-form, but this defect could be partially rescued by a cooling agent menthol treatment. To exclude the developmental defect or genetic background, we also knocked down *CcTRPM* under 25°C condition, and not found any death defect and cuticle defect. These results were the first report about *CcTRPM* was essential for insect polyphenism in response to low temperature. Until now, the researches about TRPM on other agricultural insects only focused on the description of its correlation with temperature, but no deeply molecular function was reported (Fu et al., 2016; Wang et al., 2021a). Therefore, the molecular regulation mechanism of *CcTRPM* in response to low temperature needs further study.

miRNAs played significant roles in insect polyphenism and cold acclimation. For example, miR-34 and miR-9b mediated the wing dimorphism of *N. lugens* (Ye et al., 2019) and *A. citricidus* (Shang et al., 2020), miR-133 inhibited behavioral aggregation by controlling dopamine synthesis in *L. migratoria* (Yang et al., 2014), and miR-31-5p via asc-C9 to regulate cold acclimation in *Monochamus alternatus* (Zhang et al., 2020). Here, we demonstrated that a key regulator, miR-252, targets *CcTRPM* via a negative feedback loop to control the transition from summer-form to winter-form in response to low temperature. Interestingly, RNAi-mediated knockdown of *CcTRPM* also decreased the expression of miR-252. This result implied that up-regulation of *CcTRPM* in response to low temperature may be have a feedback to miR-252, then miR-252 inhibits the expression of *CcTRPM* under low temperature condition to avoid its excessive expression. Upon miR-252 agomir feeding, the treated nymphs showed thinner cuticle thickness and less cuticle chitin content compared with the control of agomir-NC feeding under 10°C condition. Besides, miR-252 antagomir feeding could rescue the transition percent defect and morphological phenotype defect caused by *CcTRPM* knockdown. These results were the first report about miR-252 functions as the up-stream regulator to control the transition from summer-form to winter-form in response to low temperature via acting on the insect cuticle thickness and cuticle chitin content. Similar result was reported in *Daphnia pulex*, miR-252 might increase cadmium tolerance by switching cellular growth and cuticle protein pathways to detoxification processes (Chen et al., 2016). In *Leptinotarsa decemlineata*, 42 differentially expressed miRNAs were identified following cold exposure including miR-9a-3p, miR-210-3p, miR-276-5p and miR-277-3p (Morin et al., 2017). Therefore, the regulatory network of miRNAs is need further investigating in the transition between two phases in *C. chinensis*, especially transboundary miRNAs from host plant.

Insect cuticle serves as a protective barrier and plays significant roles in supporting the body, resisting the adverse environmental conditions, preventing predators from predation, and preventing pathogens from invading (Evans and Sanson, 2005; Vargas et al., 2014). Chitin as an extremely important structural component of insects and represents up to 60% of the dry weight in some species (Richards, 1978). Many studies have reported that insect cuticles are essential for cold acclimation (Fedorka et al., 2013; Kreß et al., 2016). In this study, cuticle thickness and cuticle chitin content were the major differences between summer-form and winter-form. We have demonstrated that *CcTRPM* and miR-252 function as the up-stream signaling to mediate the transition from summer-form to winter-form in response to low temperature via affecting the cuticle thickness and cuticle chitin content. Therefore, it is necessary to study whether chitin

biosynthesis pathway acting as the down-stream signaling in this transition. Further RNAi results showed that two rate-limiting enzyme genes, *CcTre1* and *CcCHS1*, play essential roles in the transition from summer-form to winter-form in response to low temperature and affect the cuticle thickness and cuticle chitin content. In addition, *CcTRPM* knockdown and miR-252 agomir feeding both markedly decreased the mRNA expression of *CcTre1* and *CcCHS1*. These results proved our hypothesis and were the first report that chitin biosynthesis pathway functions as the down-stream signaling to control the transition from summer-form to winter-form in *C. chinensis*.

In summary, this study reported a novel signaling pathway of low temperature-miR-252-*CcTRPM* acting on cuticle chitin biosynthesis genes to mediate the transition from summer-form to winter-form in *C. chinensis*. We proposed a working model to describe this novel mechanism in Figure 6. Under 25°C condition, the expression of temperature receptor *CcTRPM* was depressed by miR-252, so the 1st instar nymphs of summer-form normally developed into 3rd instar nymphs of summer-form. However, under 10°C condition, low temperature greatly activated the expression of *CcTRPM*, but miR-252 functioned as a negative feedback regulator to mediate *CcTRPM* expression to avoid its excessive expression in response to low temperature. Then, *CcTRPM* obviously improved the expression of two rate-limiting enzyme genes from cuticle chitin biosynthesis pathway, *CcTre1* and *CcCHS1*, to increase cuticle thickness and raise cuticle chitin content. Lastly, the 1st instar nymphs of summer-form developed to 3rd instar nymphs of winter-form as an ecological adaptation strategy under low temperature stress. Future researches will focus on the following directions: (1) Whether and how transboundary miRNAs and hormones from host plant involve in the transition from summer-form to winter-form in *C. chinensis*. (2) How neuropeptides and endocrine hormones response to low temperature and *CcTRPM* to mediate the transition from summer-form to winter-form in *C. chinensis*.

Materials and Methods

Insects and host plants

The summer-form and winter-form colonies of *C. chinensis* (Yang & Li) were initiated collected in June and December 2018 from a pear orchard in Daxing district, Beijing, China, respectively. These two populations are now reared in the integrated pest management (IPM) laboratory of China Agricultural University. The summer-form population was raised in growth chambers at 25 ± 1°C under a photoperiod of 12L: 12D with 65 ± 5% RH (relative humidity). The feeding conditions of winter-form population were raised at 10 ± 1 ° C under a photoperiod of 12L: 12D with 25 ± 5% RH. Unless otherwise specified, the photoperiod of all the following different temperature treatments is 12L: 12D.

Host plants were 2-3 years old Korla fragrant pear seedlings with a height of 60-100 cm under conventional water and fertilizer management.

Different temperatures and cooling agents induced the transition from summer-form to winter-form

For different temperatures treatment, the newly-hatched 1st instar summer-form nymphs (within 12 h) were transferred to the young vegetative shoots of healthy Korla fragrant pear seedlings at four different temperatures (25°C, 20°C, 15°C, and 10°C) by using a fine brush. Then,

observed the morphological characteristics every two days, and counted the number of summer-form and winter-form individuals according to the body color and external characteristics until the 3rd instar. Each treatment was performed with three independent biological replications of at least 30 nymphs for each biological replication.

Menthol (HY-N1369) was purchased from MedChemExpress LLC (Shanghai, China) and borneol (420247) was bought from Sigma-Aldrich LLC (Saint Louis, USA) (Figure S2A-S2B) (Peier et al., 2002; Turner et al., 2016). For the treatment of different cooling agents, feeding setup through a pear leaf was used as previously described in *Aphis citricidus* with minor adjustment (Shang et al., 2020). Briefly, a stem with a fresh pear leaf was inserted into a sterile PCR tube of 250 μL containing 200 μL of menthol (1 mg/mL) or borneol (0.1 mg/mL). Then, the PCR tube with leaf was transferred to a 50 mL sterile plastic tube, termed as stem-leaf device. Finally, transferred the newly-hatched 1st instar summer-form nymphs (within 12 h) into the stem-leaf device, observed the morphological characteristics every two days, and counted the number of summer-form and winter-form individuals according to the body color and external characteristics until the 3rd instar under 25°C. Each treatment was performed with three independent biological replications of at least 30 nymphs for each biological replication.

Transmission electron microscopy assay of cuticle

To analyze the difference of nymph cuticle ultrastructure and thickness between different samples, transmission electron microscopy was used as previously described (Ekino et al., 2017; Ge et al., 2019). Briefly, samples were firstly fixed in 4% polyformaldehyde (PFA) for 48 h and followed by post-fixation in 1% osmium tetroxide for 1.5 h. Next, the samples were dehydrated in a standard ethanol/acetone series, infiltrated and embedded in spurr resin embedding medium. Then, the superthin sections (-70 nm) were cut and stained with 5% uranyl acetate followed by Reynolds' lead citrate solution. Finally, the sections were observed, photographed and measured under a Hitachi HT7800 transmission electron microscope operated at 120 kv. Five nymphs were used for each sample.

Cuticle chitin staining assay with WGA-FITC and determination of cuticle chitin content

Previous studies have demonstrated that cuticle thickness and hardness are important parameters for adapting to environment and natural enemies (Evans and Sanson, 2005). Therefore, cuticle chitin staining with WGA-FITC (Cat# F8520, Solarbio, Beijing, China) and cuticle chitin content were measured to analyze the difference between the summer-form and winter-form nymphs, or dsRNA treatment and miRNA agomir treatment. For the WGA-FITC staining, nymph samples were firstly fixed with 4% PFA for 48 h and dehydrated with a series of sucrose solution with varied concentrations (10% for 4 h, 20% for 4 h, 30% for 12 h), then rinsed three times for 15 min per rinse with sterile PBS buffer. The dehydrated samples were gradually embedded in Tissue-Tek O.C.T. compound (Cat# 4583, SAKURA, Ningbo, China) after the pre-embedding stages under -25°C. The ultra-thin sections (approximately 70 nm thickness) of embedded material were cut using Leica freezing ultra-cut microtome (CM1850, Leica, Weztlar, Germany). The sections were stained with WGA-FITC (50 μg/mL) and DAPI (10 μg/mL) for 15 min and then rinsed three times for 15 min per rinse with sterile PBS buffer as previously described (Farnesi et al., 2012; Xie et al., 2022). Fluorescence images were acquired using a Leica SP8 confocal microscopy (Weztlar, Germany).

For the determination of cuticle chitin content, chitin Elisa kit (Cat# YS80663B, Yaji
Biotechnology, Shanghai, China) was used according to the manufacturer's instruction. In short,
each 50 nymphs for one sample were homogenized in 0.2 mL distilled water using an electric
homogenizer. The testing samples and standard samples were hatched in enzyme-labeled plates at
37°C for 30 min, separately. The washing buffer was used to rinse the enzyme-labeled plates five
times for 30 s per time. Then, enzyme labeling reagent was added to the plates for hatching and
washing. Finally, chromogenic agent A solution and B solution were mixed in holes for coloration
at 37°C and protect from light. The absorbance (OD value) of each hole was measured at 450nm
wavelength and zeroed with a blank hole.

Sequence characterization of *CcTRPM*

The GenBank accession number of *CcTRPM* was OQ658558. The tertiary protein structures
of   *CcTRPM*   was   predicted   with   online   server   Phyre^2
(http://www.sbg.bio.ic.ac.uk/phyre2/html/page.cgi?id=index) and modified with PyMOL-v1.3r1
software. NCBI database was used to search the homologous protein sequences of *CcTRPM* from
various species. Multiple alignments of the amino acid sequences for *CcTRPM* with other
homologies was performed using DNAman software. Phylogenetic analysis was carried out based
on the neighbor-joining (NJ) method using MEGA10.1.8 software.

qRT-PCR for mRNAs and miRNAs

To examine the temporal expression profile of *CcTRPM*, summer-form samples were
collected at different developmental stages, including egg; nymphs of the 1st, 2nd, 3rd, 4th, and
5th instar; adults of 1, 5, and 10 days. For the effect of different temperatures treatment on the
mRNA expression of *CcTRPM*, miR-252, and chitin biosynthesis signaling genes, summer-form
1st instar nymphs were treated at 25°C and 10°C. Samples of nymphs were collected at 3, 6, and
10 days after different temperatures treatment and immediately stored at -80°C. Each sample was
performed in three replications, with about 100 individuals for each replication of egg samples and
at least 50 insects were included for each nymph or adult samples. All samples were immediately
stored at -80°C for total RNA extraction.
Total RNA was isolated from the above samples using TRNzol Universal (Cat# DP424,
TIANGEN, Beijing, China) and miRcute miRNA isolation kit (Cat# DP501, TIANGEN, Beijing,
China) for mRNA and miRNA according to the manufacturer' instruction, respectively. The
first-strand cDNAs of mRNA and mature miRNA were synthesized from 500 ng of total RNA
using PrimeScript™ RT reagent kit with gDNA Eraser (Cat# RR047A, Takara, Kyoto, Japan) and
miRcute Plus miRNA First-Strand cDNA Synthesis Kit (Cat# KR211, TIANGEN, Beijing, China)
following the instruction manual. The relative expression of mRNA and miRNA was respectively
quantified using TB Green® *Premix Ex Taq*™ II (Tli RNaseH Plus) (Cat# RR820A, Takara, Kyoto,
Japan) and miRcute Plus miRNA qPCR Detection Kit (Cat# FP411, TIANGEN, Beijing, China)
in a 20 μL reaction mixture on a Bio-Rad CFX Connect™ Real-Time PCR System (Bio-Rad,
Hercules, CA, USA). To reduce the error, two reference genes of *C. chinensis*, *Ccβ-actin*
(GenBank accession number: OQ658571) and *CcEF-1* (GenBank accession number: OQ658572),
were used as internal controls for real-time PCR. U6 snRNA of *C. chinensis* was used as reference
gene for miRNA expression. To check for the specificity, melting curves were analyzed for each
data point (Figure S2, S6-S7). The specific reverse primer for miRNA expression was

complementary to the poly (T) adapter that was provided in the miRcute plus miRNA qPCR
detection kit. qRT-PCR data were quantified using the $2^{-\Delta\Delta CT}$ method, where $\Delta\Delta CT$ is equal to
$\Delta CT_{treated\ sample}$ -$\Delta CT_{control}$ (Livak and Schmittgen, 2001).

Ca^{2+} imaging and fluorescence detection of Fluo-4 AM

The ORF sequence of *CcTRPM* was amplified by PCR with specific primers shown in Table
S1 and inserted into the pcDNA3.1(+)-mCherry vector to construct the recombinant vector of
CcTRPM-pcDNA3.1-mCherry using the *pEASY*®-Basic Seamless Cloning and Assembly Kit
(Cat# CU201, TransGen, Beijing, China). This recombinant vector was transfected into HEK293T
cells using TransIntro® EL Transfection Reagent (Cat# FT201, TransGen, Beijing, China) and the
cells were cultured for 20-40 h at 37°C. These transient-transfected cells were incubated in 500 μL
Fluo-4 AM (Cat# S1061, Beyotime, Shanghai, China) for 30 min at 37°C for staining and washed
for three times with Hank's Balanced Salt Solution. Then, these cells were treated with different
cooling agents and used for Ca^{2+} imaging by a Leica SP8 confocal microscopy (Weztlar, Germany)
or for fluorescence detection of Fluo-4 AM by a Molecular Devices i3x microplate reader (San
Jose, USA).

Fluorescence *in situ* hybridization (FISH)

FISH experiment was carried out as previously described with minor adjustment (Kliot et al.,
2014). Briefly, 20 nymphs for each sample were immersed in Carnoy's fixative (glacial acetic acid:
ethanol: chloroform, 1: 3: 6, vol/vol/vol) inside eppendorf tubes for at least 48 h at room
temperature. The fixed samples were washed with 50% ethanol for three times and decolorized in
6% H$_2$O$_2$ in ethanol for 48 h at room temperature. Prehybridization was performed with three
times for 10 min per time using the hybridization buffer (20 mM Tris-HCl, pH 8.0; 0.9 M NaCl;
0.01% sodium dodecyl sulfate; 30% formamide) without the probe in the dark. The probe
sequences of *CcTRPM*-Cy3 and miR-252-FAM are listed in Table S1 and synthesized in
GenePharma (Shanghai, China). Hybridize samples for 12 h in the dark in hybridization buffer
containing 1 μM fluorescent probe. Combine two probes with different fluorescent dyes for the
co-localization. After washing three times with PBST, cell nuclei were stained with DAPI (1
μg/mL) for 15 min and rinsed with PBST for 15 min. At last, the samples were transferred to the
microscope slide, mounted in mounting medium using a cover slip, and viewed under a Leica SP8
confocal microscopy (Weztlar, Germany). The dsCcTRPM treated samples or no-probe samples
were used as negative controls for the specificity of probe signal.

dsRNA synthesis and RNAi assay

Double-stranded RNAs (dsRNAs) of *EGFP*, *CcTRPM*, and 14 transcripts of chitin
biosynthesis signaling genes were prepared in vitro using primers ligated with T7 RNA
polymerase promoter sequences (Table S1) using MEGAscript™ RNAi kit (AM1626, Ambion,
California, USA). dsRNAs were further purified with MEGAclear columns (Ambion) and eluted
with diethyl pyrocarbonate (DEPC)-treated nuclease-free water. The purity and concentration of
dsRNA were then measured via ultraviolet spectrophotometry and gel electrophoresis. The above
stem-leaf device was used for dsRNA feeding and RNAi assay.
For the RNAi assays, summer-form 1st instar nymphs were fed with different dsRNAs (500
ng/μL) and divided into three groups. (1) The samples were collected after dsRNAs feeding at 2 d

and 4 d under 25°C condition, and at 3 d and 6 d under 10°C condition, then for the RNAi

efficiency analysis and miR-252 expression by qRT-PCR. (2) Observed the morphological

characteristics every two days and counted the number of summer-form and winter-form

individuals based on the above description until the 3rd instar under 10°C condition. (3) The

samples were collected after dsRNA feeding at 12 d for cuticle ultrastructure comparison, cuticle

chitin staining with WGA-FITC, and determination of cuticle chitin content under 10°C condition

according to the above methods.

For the rescue by menthol feeding under dsCcTRPM treatment, summer-form 1st instar

nymphs were fed with the mixture of dsCcTRPM (500 ng/μL) and menthol (1 mg/mL), then

counted the number of summer-form and winter-form individuals following the above method

until the 3rd instar under 10°C condition. All the experiments were performed in three replications

with at least 30 nymphs for each replication.

Prediction of miRNAs for *CcTRPM*

The 3'UTR sequence of *CcTRPM* gene was amplified by the 3'-Full RACE Core Set with

PrimeScript™ RTase kit (Cat# 6106, Takara, Kyoto, Japan). miRNAs for *CcTRPM* were

predicted by a service provider LC science with two software programs of miRanda

(http://www.microrna.org) and Targetscan (https://www.targetscan.org/vert_72/) with the default

parameters (Enright et al., 2003; Agarwal et al., 2015). For software miRanda, the alignment

scores of 145 and -15 kcal/mol were set to the threshold. The context score percentile of

Targetscan was set as larger than 50 to evaluate the target relationship between miRNA and

potential targets. The overlapped miRNAs predicted through both methods were selected.

In vitro luciferase activity assay

The 419 bp sequence of 3'UTR containing predicted target sites for miR-252 in *CcTRPM*

was cloned into the pmirGLO vector (Promega, Wisconsin, USA) downstream of the luciferase

gene to construct the recombinant plasmid of CcTRPM-3'UTR-pmirGLO using the *pEASY®*-Basic

Seamless Cloning and Assembly Kit. The mutated 3'UTR sequence was amplified and cloned into

the pmirGLO vector to generate the CcTRPM-3'UTR mutant-pmirGLO plasmid. The agomir and

antagomir of miRNAs were chemically synthesized and modified by GenePharma (Shanghai,

China) with chemically modified RNA oligos of the same sequence or anti-sense oligonucleotides

of the miRNA. The negative control for agomir and antagomir was provided by the manufacturer.

HEK293T cells were cultured in a 24-well plate at 37°C in 5% CO_2 overnight and transfected with

CcTRPM-3'UTR-pmirGLO plasmid (approximate 500 ng) or CcTRPM-3'UTR mutant-pmirGLO

plasmid (approximately 500 ng) and 275 nM agomir-miRNA or agomir-NC using the Calcium

Phosphate Cell Transfection Kit (Cat# C0508, Beyotime, Nanjing, China). The activity of the two

luciferase enzymes was measured 24 h after co-transfection following the manufacturer's

instruction of Dual-Luciferase® Reporter Assay System (Cat# E1910, Promega, Wisconsin, USA).

Nine biological replications were carried out for each transfection.

In vivo RNA-binding protein immunoprecipitation Assay (RIP)

589   A Magna RIP Kit (Cat# 17-704, Merck, Millipore, Germany) was used to perform the RIP

590 assay following the manufacturer's instruction. Specifically, approximately 50 summer-form 1st

591 instar nymphs were fed with agomir-252 or agomir-NC for 24 h. The treated nymphs were

collected and crushed with an auto homogenizer in ice-cold RIP lysis buffer, then stored at -80°C overnight. Next, 50 μL magnetic beads were incubated with 5 μg of Ago-1 antibody (Merck, Millipore, Germany) or IgG antibody (Merck, Millipore, Germany) to form bead-antibody complex. The frozen lysates were thawed and centrifuged, and the supernatants were divided into three equal parts. One part is considered as an "input" sample and stored at -80°C for qRT-PCR analysis. The other two parts were incubated with the magnetic bead-antibody complex (anti-Ago1 or IgG) at 4°C overnight. The immunoprecipitated RNAs were released by digestion with protease K and extracted by the above methods. Finally, the abundance of *CcTRPM* and miR-252 were quantified by qRT-PCR. The "input" samples and IgG controls were assayed to normalize the relative expression of target genes and ensure the specificity of the RNA-protein interactions. Six replications were carried out for each treatment.

Feeding treatment of agomir-252 and antagomir-252

To determine the effect of miR-252 on *CcTRPM* expression, summer-form 1st instar nymphs were fed with agomir-252 (1 μM) or antagomir-252 (1 μM) and collected at 6 d after treatment for qRT-PCR. Agomir/antagomir negative control was fed as the control. To test the effect of miR-252 on the transition of summer-form to winter-form, summer-form 1st instar nymphs were fed with agomir-252 (1 μM) or agomir-NC (1 μM) and divided into three groups. (1) The samples were collected after different agomir feeding at 3 d and 6 d under 10°C condition, then for miR-252 expression analysis using qRT-PCR. (2) The samples were collected after agomir feeding at 12 d for cuticle ultrastructure comparison, cuticle chitin staining with WGA-FITC, and determination of cuticle chitin content under 10°C condition according to the above methods. (3) Observed the morphological characteristics every two days and counted the number of summer-form and winter-form individuals based on the above description until the 3rd instar under 10°C condition.

For the rescue by antagomir-252 feeding under dsCcTRPM treatment, summer-form 1st instar nymphs were fed with the mixture of antagomir-252 (1 μM) and dsCcTRPM (500 ng/μL), then counted the number of summer-form and winter-form individuals based the above method until the 3rd instar under 10°C condition. All the experiments were performed in nine replications with at least 30 nymphs for each replication.

Statistical analysis

GraphPad Prism 8.0 software and IBM SPSS Statistics 26.0 were used for making figures and statistical analysis, respectively. All the data were shown as means ± standard deviation (SD) with different independent biological replications. Student's *t*-test was performed for pairwise comparisons to determine the statistically significant differences between treatments and controls (***$P < 0.001$). One-way ANOVA followed by the Tukey's HSD multiple comparison test was used for multiple comparisons (different letters denoted by $P < 0.05$).

Acknowledgements
Thanks for the funding support by the National Natural Science Foundation of China (32202291) and China Agriculture Research System (CARS-28).

Authors' contributions

636   S.Z., and X.L. designed research; S.Z., J.L., and D.Z. performed research; Z.Z., and S.M.
637   contributed new analytic tools; S.Z., J.L., D.Z., Z.Z., S.M., and Z.L. analyzed data; S.Z. wrote the
638   paper; and S.Z., and X.L. provided the fund.
639

640   **Data accessibility**
641   The published article includes all data generated or analyzed during this study. The full sequences
642   were submitted to NCBI, accession number: OQ658558 for *CcTRPM*, OQ734934 for *CcTre1*, and
643   OQ658570 for *CcCHS1*.
644

645   **Funding**
646   This work was funded by the National Natural Science Foundation of China (32202291) and
647   China Agriculture Research System (CARS-28).
648

649   **Competing interests**
650   The authors declared no competing financial interests.
651

References

Agarwal, V., Bell, G. W., Nam, J.W., & Bartel, D. P. (2015). Predicting effective microRNA
target sites in mammalian mRNAs. *ELife*, *4*, e05005. https://doi.org/10.7554/eLife.05005
An, C., Fei, X., Chen, W., & Zhao, Z. (2012). The integrative effects of population density,
photoperiod, temperature, and host plant on the induction of alate aphids in *Schizaphis*
graminum. *Archives of Insect Biochemistry and Physiology*, *79*(4–5), 198–206.
https://doi.org/10.1002/arch.21005
Asgari, S. (2013). MicroRNA functions in insects. *Insect Biochemistry and Molecular Biology*,
43(4), 388–397. https://doi.org/10.1016/j.ibmb.2012.10.005
Butt, B. A., & Stuart, C. (1986). Oviposition by summer and winter forms of *Pear Psylla*
(Homoptera: Psyllidae) on dormant pear budwood. *Environmental Entomology*, *5*,
1109–1110. https://doi.org/10.1093/ee/15.5.1109
Chen, S., Nichols, K. M., Poynton, H. C., & Sepúlveda, M. S. (2016). MicroRNAs are involved in
cadmium tolerance in *Daphnia pulex. Aquatic Toxicology (Amsterdam, Netherlands)*, *175*,
241–248. https://doi.org/10.1016/j.aquatox.2016.03.023
Cridge, A. G., Leask, M. P., Duncan, E. J., & Dearden, P. K. (2015). What do studies of insect
polyphenisms tell us about nutritionally-triggered epigenomic changes and their
consequences? *Nutrients*, *7*(3), 1787–1797. https://doi.org/10.3390/nu7031787
Dai, T.M., Wang, Y.S., Liu, W.X., Lü, Z.C., & Wan, F.H. (2018). Thermal discrimination and
transgenerational temperature response in *Bemisia tabaci* Mediterranean (Hemiptera:
Aleyrodidae): putative involvement of the thermo-sensitive receptor *BtTRPA*.
Environmental Entomology, *47*(1), 204–209. https://doi.org/10.1093/ee/nvx202
Ekino, T., Yoshiga, T., Takeuchi-Kaneko, Y., & Kanzaki, N. (2017). Transmission electron
microscopic observation of body cuticle structures of phoretic and parasitic stages of
Parasitaphelenchinae   nematodes.   *PloS   One*,   *12*(6),   e0179465.
https://doi.org/10.1371/journal.pone.0179465
Enright, A. J., John, B., Gaul, U., Tuschl, T., Sander, C., & Marks, D. S. (2003). MicroRNA
targets in *Drosophila. Genome Biology*, *5*(1), R1. https://doi.org/10.1186/gb-2003-5-1-r1
Evans, A. R., & Sanson, G. D. (2005). Biomechanical properties of insects in relation to
insectivory: Cuticle thickness as an indicator of insect "hardness" and "intractability."
Australian Journal of Zoology, *53*(1), 9–19. https://doi.org/10.1071/zo04018
Farnesi, L. C., Brito, J. M., Linss, J. G., Pelajo-Machado, M., Valle, D., & Rezende, G. L. (2012).
Physiological and morphological aspects of *Aedes aegypti* developing larvae: effects of
the   chitin   synthesis   inhibitor   novaluron.   *PLoS   ONE*,   *7*(1),   e30363.
https://doi.org/10.1371/journal.pone.0030363
Fedorka, K. M., Copeland, E. K., & Winterhalter, W. E. (2013). Seasonality influences cuticle
melanization and immune defense in a cricket: support for a temperature-dependent
immune investment hypothesis in insects. *The Journal of Experimental Biology*, *216*(Pt
21), 4005–4010. https://doi.org/10.1242/jeb.091538
Fu, T., Hull, J. J., Yang, T., & Wang, G. (2016). Identification and functional characterization of
four transient receptor potential ankyrin 1 variants in *Apolygus lucorum* (Meyer-Dür).
Insect Molecular Biology, *25*(4), 370–384. https://doi.org/10.1111/imb.12231

Ge, Y., Zhang, L., Qin, Z., Wang, Y., Liu, P., Tan, S., Fu, Z., Smith, O. M., & Shi, W. (2019).
Different predation capacities and mechanisms of *Harmonia axyridis* (Coleoptera:
Coccinellidae) on two morphotypes of pear psylla *Cacopsylla chinensis* (Hemiptera:
Psyllidae). *PLoS ONE*, *14*(4), e0215834. https://doi.org/10.1371/journal.pone.0215834

Guo, S., Tian, Z., Wu, Q.W., King-Jones, K., Liu, W., Zhu, F., & Wang, X.P. (2021). Steroid
hormone ecdysone deficiency stimulates preparation for photoperiodic reproductive
diapause. *PLoS Genetics*, *17*(2), e1009352. https://doi.org/10.1371/journal.pgen.1009352

Hildebrand, M., Dickler, E., & Geider, K. (2010). Occurrence of *Erwinia amylovora* on insects in
a fire blight orchard. *Journal of Phytopathology*, *148*(4), 251–256.
https://doi.org/10.1046/j.1439-0434.2000.00504.x

Hou, L., Yang, P., Jiang, F., Liu, Q., Wang, X., & Kang, L. (2017). The neuropeptide F/nitric
oxide pathway is essential for shaping locomotor plasticity underlying locust phase
transition. *ELife*, *6*, e22526. https://doi.org/10.7554/eLife.22526

Kennell, J. A., Cadigan, K. M., Shakhmantsir, I., & Waldron, E. J. (2012). The microRNA miR-8
is a positive regulator of pigmentation and eclosion in *Drosophila*. *Developmental
Dynamics: An Official Publication of the American Association of Anatomists*, *241*(1),
161–168. https://doi.org/10.1002/dvdy.23705

Kliot, A., Kontsedalov, S., Lebedev, G., Brumin, M., Cathrin, P. B., Marubayashi, J. M., Skaljac,
M., Belausov, E., Czosnek, H., & Ghanim, M. (2014). Fluorescence in situ hybridizations
(FISH) for the localization of viruses and endosymbiotic bacteria in plant and insect
tissues. *Journal of Visualized Experiments: JoVE*, *84*, 51030.
https://doi.org/10.3791/51030

Kreß, A., Kuch, U., Oehlmann, J., & Müller, R. (2016). Effects of diapause and cold acclimation
on egg ultrastructure: new insights into the cold hardiness mechanisms of the Asian tiger
mosquito *Aedes (Stegomyia) albopictus*. *Journal of Vector Ecology: Journal of the
Society for Vector Ecology*, *41*(1), 142–150. https://doi.org/10.1111/jvec.12206

Lee, Y., Lee, Y., Lee, J., Bang, S., Hyun, S., Kang, J., Hong, S.T., Bae, E., Kaang, B.K., & Kim, J.
(2005). Pyrexia is a new thermal transient receptor potential channel endowing tolerance
to high temperatures in *Drosophila melanogaster*. *Nature Genetics*, *37*(3), 305–310.
https://doi.org/10.1038/ng1513

Liu, F., Peng, W., Li, Z., Li, W., Li, L., Pan, J., Zhang, S., Miao, Y., Chen, S., & Su, S. (2012).
Next-generation small RNA sequencing for microRNAs profiling in *Apis mellifera*:
Comparison between nurses and foragers. *Insect Molecular Biology*, *21*(3), 297–303.
https://doi.org/10.1111/j.1365-2583.2012.01135.x

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time
quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods*, *25*(4), 402–408.
https://doi.org/10.1006/meth.2001.1262

Lyons, P. J., Crapoulet, N., Storey, K. B., & Morin, P. (2015). Identification and profiling of
miRNAs in the freeze-avoiding gall moth *Epiblema scudderiana* via next-generation
sequencing. *Molecular and Cellular Biochemistry*, *410*(1–2), 155–163.
https://doi.org/10.1007/s11010-015-2547-3

Ma, Z., Guo, W., Guo, X., Wang, X., & Kang, L. (2011). Modulation of behavioral phase changes
of the migratory locust by the catecholamine metabolic pathway. *Proceedings of the*

National Academy of Sciences of the United States of America, *108*(10), 3882–3887.

https://doi.org/10.1073/pnas.1015098108

McKemy, D. D., Neuhausser, W. M., & Julius, D. (2002). Identification of a cold receptor reveals

a general role for TRP channels in thermosensation. *Nature*, *416*(6876), 52–58.

https://doi.org/10.1038/nature719

Minke, B., Wu, C., & Pak, W. L. (1975). Induction of photoreceptor voltage noise in the dark in

Drosophila mutant. *Nature*, *258*(5530), 84–87. https://doi.org/10.1038/258084a0

Morin, M. D., Frigault, J. J., Lyons, P. J., Crapoulet, N., Boquel, S., Storey, K. B., & Morin, P. J.

(2017). Amplification and quantification of cold-associated microRNAs in the Colorado

potato beetle (*Leptinotarsa decemlineata*) agricultural pest. *Insect Molecular Biology*,

26(5), 574–583. https://doi.org/10.1111/imb.12320

Neely, G. G., Keene, A. C., Duchek, P., Chang, E. C., Wang, Q.P., Aksoy, Y. A., Rosenzweig, M.,

Costigan, M., Woolf, C. J., Garrity, P. A., & Penninger, J. M. (2011). TrpA1 regulates

thermal nociception in *Drosophila*. *PloS One*, *6*(8), e24343.

https://doi.org/10.1371/journal.pone.0024343

Oldfield, & G., N. (1970). Diapause and polymorphism in california populations of *Psylla*

pyricola (Homoptera: Psyllidae). *Annals of the Entomological Society of America*, *63*(1),

180–184. https://doi.org/10.1093/aesa/63.1.180

Peier, A. M., Moqrich, A., Hergarden, A. C., Reeve, A. J., Andersson, D. A., Story, G. M., Earley,

T. J., Dragoni, I., McIntyre, P., Bevan, S., & Patapoutian, A. (2002). A TRP channel that

senses cold stimuli and menthol. *Cell*, *108*(5), 705–715.

https://doi.org/10.1016/S0092-8674(02)00652-9

Richards, A. G. (1978). The chemistry of insect cuticle. *Biochemistry of Insects*, 205–232.

https://doi.org/10.1016/B978-0-12-591640-0.50010-8

Rosenzweig, M., Kang, K., & Garrity, P. A. (2008). Distinct TRP channels are required for warm

and cool avoidance in *Drosophila melanogaster*. *Proceedings of the National Academy of*

Sciences of the United States of America, *105*(38), 14668–14673.

https://doi.org/10.1073/pnas.0805041105

Sato, A., Sokabe, T., Kashio, M., Yasukochi, Y., Tominaga, M., & Shiomi, K. (2014). Embryonic

thermosensitive TRPA1 determines transgenerational diapause phenotype of the

silkworm, *Bombyx mori*. *Proceedings of the National Academy of Sciences of the United*

States of America, *111*(13), E1249–E1255. https://doi.org/10.1073/pnas.1322134111

Shang, F., Niu, J., Ding, B.Y., Zhang, W., Wei, D.D., Wei, D., Jiang, H.B., & Wang, J.J. (2020).

The miR-9b microRNA mediates dimorphism and development of wing in aphids.

Proceedings of the National Academy of Sciences, *117*(15), 8404–8409.

https://doi.org/10.1073/pnas.1919204117

Shiels, A. (2020). TRPM3_miR-204: A complex locus for eye development and disease. *Human*

Genomics, *14*(1), 7. https://doi.org/10.1186/s40246-020-00258-4

Simpson, S. J., Sword, G. A., & Lo, N. (2011). Polyphenism in insects. *Current Biology*, *21*(18),

R738–R749. https://doi.org/10.1016/j.cub.2011.06.006

Teets, N. M., Marshall, K. E., & Reynolds, J. A. (2023). Molecular mechanisms of winter survival.

Annual Review of Entomology, *68*, 319–339.

https://doi.org/10.1146/annurev-ento-120120-095233

Terblanche, J. S., Hoffmann, A. A., Mitchell, K. A., Rako, L., le Roux, P. C., & Chown, S. L. (2011). Ecologically relevant measures of tolerance to potentially lethal temperatures. *The Journal of Experimental Biology*, *214*(Pt 22), 3713–3725. https://doi.org/10.1242/jeb.061283

Tougeron, K., Iltis, C., Renoz, F., Albittar, L., Hance, T., Demeter, S., & Le Goff, G. J. (2021). Ecology and biology of the parasitoid *Trechnites insidiosus* and its potential for biological control of pear psyllids. *Pest Management Science*, *77*(11), 4836–4847. https://doi.org/10.1002/ps.6517

Tracey, W. D., Wilson, R. I., Laurent, G., & Benzer, S. (2003). Painless, a *Drosophila* gene essential for nociception. *Cell*, *113*(2), 261–273. https://doi.org/10.1016/s0092-8674(03)00272-1

Tsuchiya, R., Kaneshima, A., Kobayashi, M., Yamazaki, M., Takasu, Y., Sezutsu, H., Tanaka, Y., Mizoguchi, A., & Shiomi, K. (2021). Maternal GABAergic and GnRH/corazonin pathway modulates egg diapause phenotype of the silkworm Bombyx mori. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(1), e2020028118. https://doi.org/10.1073/pnas.2020028118

Turner, H. N., Armengol, K., Patel, A. A., Himmel, N. J., Sullivan, L., Iyer, S. C., Bhattacharya, S., Iyer, E. P. R., Landry, C., Galko, M. J., & Cox, D. N. (2016). The TRP channels Pkd2, NompC and Trpm act in cold-sensing neurons to mediate unique aversive behaviors to noxious cold in Drosophila. *Current Biology: CB*, *26*(23), 3116–3128. https://doi.org/10.1016/j.cub.2016.09.038

Vargas, H. C. M., Farnesi, L. C., Martins, A. J., Valle, D., & Rezende, G. L. (2014). Serosal cuticle formation and distinct degrees of desiccation resistance in embryos of the mosquito vectors Aedes aegypti, Anopheles aquasalis and Culex quinquefasciatus. *Journal of Insect Physiology*, *62*, 54–60. https://doi.org/10.1016/j.jinsphys.2014.02.001

Vellichirammal, N. N., Gupta, P., Hall, T. A., & Brisson, J. A. (2017). Ecdysone signaling underlies the pea aphid transgenerational wing polyphenism. *Proceedings of the National Academy of Sciences of the United States of America*, *114*(6), 1419–1423. https://doi.org/10.1073/pnas.1617640114

Venkatachalam, K., & Montell, C. (2007). TRP channels. *Annual Review of Biochemistry*, *76*, 387–417. https://doi.org/10.1146/annurev.biochem.75.103004.142819

Wang, L.-X., Niu, C.-D., Wu, S.-F., & Gao, C.-F. (2021). Molecular characterizations and expression profiles of transient receptor potential channels in the brown planthopper, Nilaparvata lugens. *Pesticide Biochemistry and Physiology*, *173*, 104780. https://doi.org/10.1016/j.pestbp.2021.104780

Wang, X.-D., Lin, Z.-K., Ji, S.-X., Bi, S.-Y., Liu, W.-X., Zhang, G.-F., Wan, F.-H., & Lü, Z.-C. (2021). Molecular characterization of TRPA subfamily genes and function in temperature preference in *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *International Journal of Molecular Sciences*, *22*(13), 7157. https://doi.org/10.3390/ijms22137157

Wei, M., Chi, H., Guo, Y., Li, X., Zhao, L., & Ma, R. (2020). Demography of *Cacopsylla chinensis* (Hemiptera: Psyllidae) reared on four cultivars of *Pyrus bretschneideri* (Rosales: Rosaceae) and *P. communis* Pears with estimations of confidence intervals of specific life table statistics. *Journal of Economic Entomology*, *113*(5), 2343–2353. https://doi.org/10.1093/jee/toaa149

Xie, J., Peng, G., Wang, M., Zhong, Q., Song, X., Bi, J., Tang, J., Feng, F., Gao, H., & Li, B.
(2022). RR-1 cuticular protein TcCPR69 is required for growth and metamorphosis in
Tribolium castaneum. *Insect Science*, *29*(6), 1612–1628.
https://doi.org/10.1111/1744-7917.13038

Xu, H.J., Xue, J., Lu, B., Zhang, X.C., Zhuo, J.C., He, S.F., Ma, X.F., Jiang, Y.Q., Fan, H.W., Xu,
J.Y., Ye, Y.X., Pan, P.L., Li, Q., Bao, Y.Y., Nijhout, H. F., & Zhang, C.X. (2015). Two
insulin receptors determine alternative wing morphs in planthoppers. *Nature*, *519*(7544),
464–467. https://doi.org/10.1038/nature14286

Yang, M., Wei, Y., Jiang, F., Wang, Y., Guo, X., He, J., & Kang, L. (2014). MicroRNA-133
inhibits behavioral aggregation by controlling dopamine synthesis in Locusts. *PLoS*
Genetics, *10*(2), e1004206. https://doi.org/10.1371/journal.pgen.1004206

Yang, S., Zhang, J., Wang, S., Zhang, X., Liu, Y., & Xi, J. (2017). Identification and profiling of
miRNAs in overwintering *Lissorhoptrus oryzophilus* via next-generation sequencing.
Cryobiology, *74*, 68–76. https://doi.org/10.1016/j.cryobiol.2016.11.013

Yang, X., Wu, D., Du, H., Nie, F., Pang, X., & Xu, Y. (2017). MicroRNA-135a is involved in
podocyte injury in a transient receptor potential channel 1-dependent manner.
International Journal of Molecular Medicine, *40*(5), 1511–1519.
https://doi.org/10.3892/ijmm.2017.3152

Ye, X., Xu, L., Li, X., He, K., Hua, H., Cao, Z., Xu, J., Ye, W., Zhang, J., Yuan, Z., & Li, F.
(2019). MiR-34 modulates wing polyphenism in planthopper. *PLoS Genetics*, *15*(6),
e1008235. https://doi.org/10.1371/journal.pgen.1008235

Zhang, B., Zhao, L., Ning, J., Wickham, J. D., Tian, H., Zhang, X., Yang, M., Wang, X., & Sun, J.
(2020). MiR-31-5p regulates cold acclimation of the wood-boring beetle *Monochamus*
alternatus via ascaroside signaling. *BMC Biology*, *18*(1), 184.
https://doi.org/10.1186/s12915-020-00926-w

Zhang, J.L., Chen, S.J., Liu, X.Y., Moczek, A. P., & Xu, H.J. (2022). The transcription factor
Zfh1 acts as a wing-morph switch in planthoppers. *Nature Communications*, *13*(1), 5670.
https://doi.org/10.1038/s41467-022-33422-6

Figure 1. Low temperature induced the transition from summer form (SF) to winter form
(WF) in *C. chinensis*.

A-E and A'-E': The morphology of egg, 1st, 3rd, and 5th instar nymph, 5 days adult for SF and
WF of *C. chinensis*, respectively. The scale bars were labeled in the lower left of each figure.

F: Percent of SF and WF under different temperatures (25°C, 20°C, 15°C, and 10°C) when
induction of newly-hatched (within 12 h) 1st instar nymph of *C. chinensis*.

G: Effect of two chemical cooling agents (menthol and borneol) on the percent of SF and WF
when induction of newly-hatched (within 12 h) 1st instar nymph of *C. chinensis* under 25°C.

H: Ultrastructure comparison of the nymph cuticle for SF 3rd instar and WF 3rd instar of *C.
chinensis* using transmission electron microscopy. Scale bar is 1 μm. The two-way arrow
indicated the cuticle thickness.

I: Comparison of the nymph cuticle thickness for SF 3rd instar and WF 3rd instar of *C. chinensis*.

J: Chitin staining of nymph cuticle with WGA-FITC in SF 3rd instar and WF 3rd instar of *C.*

chinensis. DAPI: the cell nuclei were stained with DAPI and visualized in blue. WAG-FITC: the
cuticle chitin was labeled with FITC and visualized in green. Merge: merged imaging of DAPI,
and WAG-FITC signals.
K: Determination of cuticle chitin content in 2nd and 3rd instar nymph of SF and WF.
The data of nymph percent were shown as the mean ± SD with three independent biological
replications of at least 30 nymphs for each biological replication. Different letters above the bars
indicated statistically significant differences ($p < 0.05$), as determined by ANOVA followed with
a Turkey's HSD multiple comparison test in SPSS 20.0 software. The results of cuticle thickness
and cuticle chitin content were presented as mean ± SD of three independent biological samples
with three technical replications for each biological replication. Statistically significant differences
were determined with the pair-wise Student's *t*-test, and significance levels were denoted by *** (*p*
< 0.001).
880

881
Figure 2. Temperature receptor *CcTRPM* modulated the transition from SF to WF of *C.*
***chinensis* in response to low temperature.**
A: The predicted protein tertiary structure of *CcTRPM*. The conserved ankyrin repeat (ANK)
domain was indicated in the N-terminal. The conserved six transmembrane domain of ion
channels structure were shown as TM1-TM6.
B: The mRNA expression of *CcTRPM* in response to different temperatures of 25°C and 10°C by
qRT-PCR.
C: Fluorescence detection of Fluo-4 AM after heterologous expression of *CcTRPM* in mammalian
HEK293T cells in response to menthol treatment. The recombinant plasmid was generated by

inserting the full ORF sequence of *CcTRPM* into pcDNA3.1(+)-mCherry plasmid. DMSO

treatment and ionomycin treatment were used as negative control and positive control, separately.

D: Representative images of Ca^{2+} imaging after heterologous expression of *CcTRPM* in

mammalian HEK293T cells in response to menthol treatment. CcTRPM (+) means the

recombinant plasmid of *CcTRPM* with pcDNA3.1(+)-mCherry. CcTRPM (-) denotes the

pcDNA3.1(+)-mCherry plasmid without *CcTRPM*. mCherry and Fluo-4 signal are shown in red

and green, respectively.

E: The developmental expression pattern of *CcTRPM* for SF at mRNA level using qRT-PCR. 1st,

2nd, 3rd, 4th and 5th are the nymphs at the first, second, third, fourth and fifth instar, respectively.

A-1d, A-5d, and A-10d are the adults at 1 day, 5 days, and 10 days, separately.

F: Representative confocal images of *CcTRPM* in different developmental stages of SF using

FISH. Scale bar is 0.5 mm. BF: the bright field. DAPI: the cell nuclei were stained with DAPI and

visualized in blue. CcTRPM-Cy3: CcTRPM signal was labeled with Cy3 and visualized in red.

Merge: merged imaging of BF, DAPI, and CcTRPM-Cy3 signals.

G-H: Comparison of the nymph cuticle ultrastructure, cuticle thickness, and cuticle chitin content

of SF 1st instar treated with dsCcTRPM and dsEGFP at 15 d.

I: The transition percent of SF 1st instar nymphs treated with dsEGFP, DMSO, dsCcTRPM,

CcTRPM antagonist and dsCcTRPM+menthol at 15 d under 10°C.

J: The phenotypes of SF 1st instar nymphs treated with dsEGFP, dsCcTRPM, CcTRPM antagonist

and dsCcTRPM+menthol at 15 d under 10°C.

The data in 2B and 2E are shown as the mean ± SD with three independent biological replications

of at least 30 nymphs for each biological replication. Data in 2G and 2H are presented as mean ±

SD with three biological replications of three technical replicates for each biological replication.

Data in 2I are presented as mean ± SD with nine biological replications. Statistically significant

differences were determined with the pair-wise Student's *t*-test, and significance levels were

denoted by *** ($p < 0.001$). Different letters above the bars indicated statistically significant

differences ($p < 0.05$), as determined by ANOVA followed with a Turkey's HSD multiple

comparison test in SPSS 20.0 software.

920
Figure 3. Confirming of miR-252 targeted *CcTRPM*.

A: The putative miR-252 binding sites in *CcTRPM* 3'UTR were predicted by miRanda and Targetscan.

B: The expression profiles of miR-252 in response to different temperatures of 25°C and 10°C by qRT-PCR.

C: In vitro validation of the target interactions between miR-252 and *CcTRPM* by dual luciferase reporter assays.

D: In vivo demonstration of miR-252 targeting *CcTRPM* by RNA-binding protein immunoprecipitation (RIP) assay.

E: Representative confocal images of miR-252 and *CcTRPM* in 1st, 3rd and 5th instar of SF. Scale bar is 0.5 mm. The signals of DAPI and *CcTRPM*-Cy3 are same as the above describing.

miR-252-FAM: the miR-252 signal was labeled with FAM and visualized in green. Merge: merged imaging of co-localization of cell nucleus, *CcTRPM* and miR-252.

F: Effect of miR-252 agomir and antagomir treatment 3 days on the expression of *CcTRPM* at mRNA level.

The data in 3B and 3F are shown as the mean ± SD with three independent biological replications of at least 30 nymphs for each biological replication. Data in 3C are presented as mean ± SD with nine biological replications. Data in 3D are presented as mean ± SD with two biological replications of three technical replicates for each biological replication. Statistically significant differences were determined with the pair-wise Student's *t*-test, and significance levels were denoted by *** ($p < 0.001$). Different letters above the bars indicated statistically significant differences ($p < 0.05$), as determined by ANOVA followed with a Turkey's HSD multiple comparison test in SPSS 20.0 software.

Figure 4. miR-252 mediated the transition from SF to WF in *C. chinensis* by suppressing

CcTRPM.

A: Effect of miR-252 agomir treatment on the mRNA expression of miR-252 after 3 d and 6 d
under 10°C. agomir-NC treatment was used as the control. 3 d: 3 days. 6 d: 6 days.

B: The mRNA expression of miR-252 after dsCcTRPM treatment 3 d and 6 d compare with the
dsEGFP treatment under 10°C.

C-F: Comparison of the nymph cuticle thickness, cuticle ultrastructure, cuticle chitin staining with
WGA-FITC, and chitin content of SF 1st instar after treatment with agomir-NC and agomir-252 at
15 d. Scale bar is 1 μm. The two-way arrow indicated the cuticle thickness. The DAPI and
WAG-FITC signals were same as the above describing.

G: The transition percent of SF 1st instar nymphs treated with agomir-NC, agomir-252, dsEGFP,
dsCcTRPM, and dsCcTRPM+antagomir-252 at 15 d under 10°C.

H: The phenotypes of SF 1st instar nymphs treated with agomir-NC, agomir-252, dsCcTRPM, and
dsCcTRPM+antagomir-252 at 15 d under 10°C.

The data in 4A and 4B are shown as the mean ± SD with three independent biological replications
of at least 30 nymphs for each biological replication. Data in 4C and 4F are presented as mean ±
SD with three biological replications of three technical replications for each biological replication.
Data in 4G are presented as mean ± SD with nine biological replications. Statistically significant
differences were determined with the pair-wise Student's *t*-test, and significance levels were
denoted by *** ($p < 0.001$). Different letters above the bars indicated statistically significant
differences ($p < 0.05$), as determined by ANOVA followed with a Turkey's HSD multiple
comparison test in SPSS 20.0 software.

Figure 5. Chitin biosynthesis signaling involved in the transition from SF to WF of *C. chinensis* in response to low temperature and *CcTRPM*.

A: Diagram of de novo biosynthesis of insect chitin. Tre: trehalase. HK: hexokinase. G6PI: glucose-6-phosphate isomerase. GFAT: fructose-6-phosphate aminotransferase. GNA: glucosamine-6-phosphate acetyltransferase. AGM: N-acetylglucosamine phosphate mutase. UAP: UDP- N-acetylglucosamine pyrophosphorylase. CHS: chitin synthase.

B: A heat map was constructed from the expression levels of chitin biosynthesis enzyme

transcripts in *C. chinensis* after 25°C and 10°C treatment at 3 d, 6 d, and 10 d.

C: Effect of RNAi knockdown of 14 transcripts covering all the enzymes in the chitin biosynthesis

pathway on the transition percent of SF 1st instar nymphs under 10°C.

D-F: Comparison of the nymph cuticle thickness, cuticle chitin content, and cuticle chitin staining

with WGA-FITC of SF 1st instar after treatment with dsEGFP, dsCcTre1, and dsCcCHS1 at 15 d.

The WAG-FITC signal was same as the above describing.

G-H: Effect of dsCcTRPM and agomir-252 treatments on the mRNA expression of *CcTre1* at 3 d

and 6 d under 10°C.

I-J: Effect of dsCcTRPM and agomir-252 treatments on the mRNA expression of *CcCHS1* at 3 d

and 6 d under 10°C.

Figure 6. A model showing the key roles of miR-252 targeting *CcTRPM* to regulate the

transition from SF to WF of *C. chinensis* in response to low temperature.

Under 25°C condition, the temperature receptor *CcTRPM* was inactive as inhibiting by miR-252

in 1st instar nymphs of SF. The 1st instar nymphs of SF normally developed into 3rd instar

nymphs of SF. However, under 10°C condition, low temperature profoundly activated *CcTRPM*

expression, miR-252 functioned as a negative regulator to mediate the expression of *CcTRPM* in

case of over-expression in response to low temperature. Then, *CcTRPM* significantly increased the

activity of chitin biosynthesis pathway, especially *CcTre1* and *CcCHS1*, and led to cuticle chitin

content was obvious raise and cuticle thickness became thicker. Finally, the 1st instar nymphs of

SF developed to 3rd instar nymphs of WF in order to better adapt to low temperature. Red arrows

and black T-bars indicated the activation and inhibition, respectively. Gray arrows and T-bars

represented the inactive states of the genes or physiological processes, separately.

   Supplemental information

   Figure S1. Sequence characterization of TRPM from different insect species.
   A: Multiple alignments of the amino acid sequences of *CcTRPM* transmembrane domain with
   homologs from other insect species. Black represents 100% identity, red represents 75% identity
   and blue represents <75% identity. The transmembrane domain of TM1 to TM6 are indicated by
   black horizontal line. *CcTRPM* (*C. chinensis*, OQ658558), *DcTRPM* (*Diaphorina citri*,
   XP_017299512.2), *BtTRPM* (*Bemisia tabaci*, XP_018904971.1), *NlTRPM* (*Nilaparvata lugens*,
   AOR81476.1), *ApTRPM* (*Acyrthosiphon pisum*, XP_003245193.1), *SpTRPM* (*Schistocerca*
   piceifrons, XP_047116036.1). The same as below.
   B: Phylogenetic tree analysis of *CcTRPM* and its homologs in other nine insect species. *HsTRPM*
   (*Homo sapiens*, NP_001384540.1), *MmTRPM* (*Mus musculus*, AAL79553.1), *DmTRPM*
   (*Drosophila melanogaster*, NP_001401007.1), *CqTRPM* (*Culex quinquefasciatus*,
   XP_038115236.1).

1019

Figure S2. Effect of menthol and *CcTRPM* antagonist treatment on the mRNA expression of
CcTRPM under 10 °C condition.

A-B: The chemical structure of menthol and TRPM antagonist.

C-E: Melting curve for qRT-PCR primers of *CcTRPM*, *Ccβ-actin* and *CcEF-1*.

F: Effect of menthol treatment at different concentrations on the mRNA expression of *CcTRPM*
under 10 °C condition.

G: Effect of TRPM antagonist treatment at different concentrations on the mRNA expression of
CcTRPM under 10 °C condition.

Data are shown as mean ± SD with three biological replications. Different letters above the bars
indicated statistically significant differences ($p < 0.05$), as determined by ANOVA followed with
a Turkey's HSD multiple comparison test in SPSS 20.0 software.

1031

1032

Figure S3. RNAi efficiency of *CcTRPM* and effect of *CcTRPM* knockdown on the phenotype at 25 °C.

A: RNAi efficiency of *CcTRPM* after dsCcTRPM treatment at 3 d and 6 d by qRT-PCR under 10 °C condition.

B: Representative confocal images of *CcTRPM* after SF 1st instar nymphs treated with dsCcTRPM for 6 d at 10 °C by FISH. Scale bar is 0.5 mm. BF: the bright field. DAPI: the cell nuclei were stained with DAPI and visualized in blue. CcTRPM-Cy3: CcTRPM signal was labeled with Cy3 and visualized in red. Merge: merged imaging of BF, DAPI, and CcTRPM-Cy3 signals.

C: RNAi efficiency of *CcTRPM* after dsCcTRPM treatment at 2 d and 4 d by qRT-PCR under 25 °C condition.

D: The phenotype of SF 1st instar nymphs treated with dsEGFP and dsCcTRPM at 10 d under 25 °C condition.

Data are shown as mean ± SD with three biological replications. Statistically significant differences were determined with the pair-wise Student's *t*-test, and significance levels were denoted by *** ($p < 0.001$).

1049

1050

Figure S4. Validation of the target interactions between *CcTRPM* and other three miRNAs.

A: The putative binding sites of miR-5728, PC-3p-173924, and PC-3p-137568 in *CcTRPM* 3'UTR were predicted by miRanda and Targetscan.

B: In vitro validation of the target interactions between miR-5728, PC-3p-173924, and PC-3p-137568 with *CcTRPM* by dual luciferase reporter assays. Data are presented as mean ± SD with nine biological replications. Different letters above the bars indicated statistically significant differences ($p < 0.05$), as determined by ANOVA followed with a Turkey's HSD multiple comparison test in SPSS 20.0 software.

1059

Figure S5. Transcript of miR-252 was significantly enriched by antibody against Ago1 in agomir-252 treated group compared with agomir-NC group.

1062 Data are shown as mean ± SD with two biological replications of three technical replicates for

1063 each biological replication. Statistically significant differences were determined with the pair-wise

1064 Student's *t*-test, and significance levels were denoted by *** ($p < 0.001$).

1065

1066

Figure S6. Effect of agomir-252 on the phenotype of SF 1st instar nymphs at 25 °C.

1068 A-B: Melting curve for qRT-PCR primers of U6 and miR-252.

1069 C: Effect of agomir-252 treatment 2 d and 4 d on the mRNA expression of *CcTRPM* under 25 °C

1070 condition.

1071 D: The phenotype of SF 1st instar nymphs treated with agomir-NC and agomir-252 at 10 d at

1072 25 °C.

1073
Figure S7. RNAi efficiency of *CcTre1* and *CcCHS1* at 10 °C by qRT-PCR.

A-B: Melting curve for qRT-PCR primers of *CcTre1* and *CcCHS1*.

C-D: RNAi efficiency of *CcTre1* and *CcCHS1* after dsCcTre1 and dsCcCHS1 treatment at 3 d and 6 d under 10 °C condition by qRT-PCR, respectively.

Data are shown as mean ± SD with three biological replications. Statistically significant differences were determined with the pair-wise Student's *t*-test, and significance levels were denoted by *** ($p < 0.001$).

1081

1082 **Table S1. The primers used in current study.**

1083

Gene name (Abbreviation)	Accession number	Sequences (5'-3')	Production length (bp)	Purpose	Amplification efficiency
CcTRPM	OQ658558	Full-F1: GGATAGAGGCAACTTTTCAA Full-R1: ATCTATTCTGTCGTGCTCCA	1307	RT-PCR for sequence validation	—
		Full-F2: TAGACGAAGCCATGATGC Full-R2: ACAACTGTTTCACGGAGTGT	1162	RT-PCR for sequence validation	—
		Full-F3: TAGCCAGACATTCTCAACAA Full-R3: GAAGCATTACTATGGGTTGC	1436	RT-PCR for sequence validation	—
		Full-F4: AGCAGACCCTTTCCACTC Full-R4: TAACTGTTAACACATTGTTTCTGC	1334	RT-PCR for sequence validation	—
		qF: ACAATGCATTCTTCTTGACC qR: GACTTTGTACACTGGGGGTA	141	qRT-PCR	102.57
		RNAi-F1: GATCACTAATACGACTCACTATAGGGAGA GGTACCTTCGTATCCTCAAC RNAi-R1: GATCACTAATACGACTCACTATAGGGAGA TGGCATTGATCTCCATGAAA	472	dsRNA synthesis	—

		Probe: Cy3-CATAGATCGCGTAGGAGGTC	20	FISH	—
		3'UTR-Full-F1: CTAGTTGTTTAAACGAGCTC CATTATGTAACCACAGCGTAA 3'UTR-Full-R1: TGCATGCCTGCAGGTCGACTCTAGA TTCCTAGAAACTCAAAAATCG	419	Amplification of 3'UTR full sequence	—
		3'UTR-Mut-F1: CTAGTTGTTTAAACGAGCTC CATTATGTAACCACAGCGTAA 3'UTR-Mut-R1: TGCATGCCTGCAGGTCGACTCTAGA AAGTACTATCCACGCTCGAC	137	Amplification of 3'UTR mutant sequence	—
		Orf-F: GGACGAGCTGTACAAGTAGGGATCC ATGGAGAGGTCCCAGAGGTC Orf-R: GTTTAAACGGGCCCTCTAGACTCGAGC TTAACACATTGTTTCTGCGC	4618	Amplification of orf sequence	—
miR-252	-	qF: GCTAAGTACTGGTGCCGCAGGA	—	qRT-PCR	—
U6	-	qF: AGGATGACACGCAAAATCGT	—	qRT-PCR	—
CcTre1	OQ734934	qF: GGAACTCCCTCCTCTATGTT qR: CCAATAGTCAGCCAATCTGT	147	qRT-PCR	97.73

		RNAi-F1: GATCACTAATACGACTCACTATAGGGAGA AGATGTCAAAACCAATCCAG RNAi-R1: GATCACTAATACGACTCACTATAGGGAGA GGCCAAGTTGTAGTATTTGC	455	dsRNA synthesis	—
CcTre2	OQ734935	qF: CCAATCCTTCTCTACACCTG qR: CTGTCATTGTTGTACGGTTG	123	qRT-PCR	93.68
CcTre-like	OQ734936	qF: GACCGAACCTATGATCTCAA qR: AGATGTACCAAGCTGCTGAG	135	qRT-PCR	106.13
CcCHS1	OQ658570	F: AGAAGAGAAGAAACAGCAGG R: TAGTGTCCAATCGTTTCTCC	233	qRT-PCR	101.47
		RNAi-F1: GATCACTAATACGACTCACTATAGGGAGA TTGGCTACCACTGAGAAACT RNAi-R1: GATCACTAATACGACTCACTATAGGGAGA ATGCGTTTCGTACTCGTAAT	367	dsRNA synthesis	—
CcCHS2	OQ658569	F: TATTAGGCTGATGTTGCTGA R: GAGGCAACACAAGAGGTAAG	124	qRT-PCR	95.49
CcHK1	OQ658562	F: GAGTCTGCCATCTACTGTCC	130	qRT-PCR	96.21

		R: CGGGGGTGATATTTGTATAG			
CcHK2	OQ658560	F: AAATAGTCCGACTGGCATTA R: TGTACTCTTGGGCTCACTTT	137	qRT-PCR	98.05
CcHK3	OQ658561	F: AGCATTTGAGAGAGAACGTG R: CCGCTGATATTGTCTTCTTC	120	qRT-PCR	103.51
CcG6PI	OQ658563	F: CCTGCTATCTCTCACAATCC R: CTTTCAGTTCAGCTTCTGCT	127	qRT-PCR	107.23
CcGFAT	OQ658564	F: ACTGAAACATGGACCTCTTG R: CTTCCATCTCTGGCAATAAC	123	qRT-PCR	97.06
CcGNA	OQ658565	F: GGCGAGTATTTTGTGAGTGT R: CTCGGTACGTATCATCGACT	145	qRT-PCR	108.46
CcAGM	OQ658566	F: TTTATTTTGAGGCAAATGGT R: AAGAGTCTCCGACAGTTTCA	138	qRT-PCR	102.84
CcUAP1	OQ658567	F: TGGGGAACTTGTGTATAACC R: TGGAATCTTCTTGAGTGCTT	124	qRT-PCR	98.03
CcUAP2	OQ658568	F: ACATTGACTCCAAAGGATTG R: GGCAAAATCATGTTCTCTGT	134	qRT-PCR	89.74

CcEF-1	OQ658572	qF: TGTGTCGAGTCATTCTCTGA qR: TTACTTTACCGGAGGACAGA	124	Reference gene for qRT-PCR	98.63
Ccβ-Actin	OQ658571	qF: CAGCACCATGAAGATCAAG qR: GACTCGTCATACTCCTGCTT	129	Reference gene for qRT-PCR	102.87
EGFP	ACY56286	F: GATCACTAATACGACTCACTATAGGGAGA CTCCAGCAGGACCATGTGATC R: GATCACTAATACGACTCACTATAGGGAGA CCTGAAGTTCATCTGCACCAC	596	dsRNA synthesis	—

1084

1085   Note: The black boxes indicated the T7 promoter sequences in the primers of dsRNA synthesis. The underlines showed the homologous arm sequences for the

1086   seamless cloning in homologous recombination.

1087