Task-evoked pulse wave amplitude tracks cognitive load

Yuri G. Pavlov [1], Anastasia S. Gashkova [2], Dauren Kasanov [2], Alexandra I. Kosachenko [2], Alexander I. Kotyusov [2], Boris Kotchoubey [1]

[1] Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, 72076, Germany
[2] Department of Psychology, Ural Federal University, Department of Psychology, Ekaterinburg, 620000, Russian Federation

Corresponding author:
Yuri G. Pavlov
Email: pavlovug@gmail.com
Address: Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, 72076, Germany

Keywords: cognitive load, pulse wave amplitude, photoplethysmography, ECG, heart rate

Abstract

Mental workload is a crucial factor in cognitively demanding activities and holds significance across various research fields. This study aimed to investigate the effectiveness of pulse wave amplitude (PWA) as a measure for tracking cognitive load and associated mental effort in comparison to heart rate (HR) during a digit span task. The data from 78 participants were included in the analyses. Participants performed a memory task in which they were asked to memorize sequences of 5, 9, or 13 digits, and a control task where they passively listened to the sequences. PWA and HR were quantified from photoplethysmography (PPG) and electrocardiography (ECG), respectively. Pupil dilation was also assessed as a measure of cognitive load. We found that PWA showed a strong suppression with increasing memory load, indicating sensitivity to cognitive load. In contrast, HR did not show significant changes with task difficulty. Moreover, when memory load exceeded the capacity of working memory, a reversal of the PWA pattern was observed, indicating cognitive overload. In this respect, changes in PWA in response to cognitive load correlated with the dynamics of pupil dilation, suggesting a potential shared underlying mechanism. Additionally, both HR and PWA demonstrated a relationship with behavioral performance, with higher task-evoked HR and lower PWA associated with better memory performance. Our findings suggest that PWA is a more sensitive measure than HR for tracking cognitive load and overload. PWA, measured through PPG, holds significant potential for practical applications in assessing cognitive load due to its ease of use and sensitivity to cognitive overload. The findings contribute to the understanding of psychophysiological indicators of cognitive load and offer insights into the use of PWA as a non-invasive measure in various contexts.

Introduction

Mental effort expended during cognitive tasks often results in changes in autonomic nervous system (ANS) activity. The allocation of resources during demanding task performance is evident through various indicators of cardiovascular activity regulated by the ANS (Gendolla & Wright, 2012). Notable among these indicators are heart rate (HR) (Al abdi et al., 2018; Schapkin et al., 2012; Vincent et al., 1996), pulse wave amplitude (PWA) (Xuan et al., 2020), heart rate variability (HRV) (Mandrick et al., 2016), pre-ejection period and systolic blood pressure (Richter et al., 2008). However, the effectiveness of these indicators in assessing cognitive load and cognitive overload varies based on several factors: the relative contribution of sympathetic and parasympathetic branches of the ANS, the complexity of obtaining the measure, and the effects of individual differences and situational factors.

Cognitively demanding tasks affect cardiovascular indicators mediated by sympathetic rather than parasympathetic influences (Gendolla & Wright, 2012). However, systolic blood pressure serves as an indicator reflecting both sympathetic and parasympathetic influence (Gendolla & Wright, 2012). Similarly, heart rate (HR) has regulation by sympathetic mechanisms and parasympathetic mechanisms, with parasympathetic dominance observed during minor and moderate efforts (Wright & Kirby, 2001). Thus, recording HR and systolic blood pressure over longer epochs can average sympathetic and parasympathetic effects, which may interact with, or even cancel out, each other.

On the other hand, peripheral vasoconstriction detected via decreased pulse wave amplitude extracted from photoplethysmography (PPG) has primarily sympathetic influence (alpha-adrenergic sympathetic impact) (Babchenko et al., 2001; Shelley, 2007). Other indicators of sympathetic influence are pre-ejection period, which is an index beta-adrenergic sympathetic impact, and some measures of HRV. However, measuring pre-ejection period requires simultaneous recording of systolic blood pressure, ECG, and impedance cardiogram or seismocardiogram, and HRV can only be assessed over a long period of stationary activity (at least one minute (Quintana et al., 2016)). To the contrary, PPG is easily accessible (e.g., utilizing a smartphone camera (Matsumura & Yamakoshi, 2013)) and can potentially detect changes in cognitive load over a period of seconds.

Despite its potential, the utilization of photoplethysmography (PPG) in psychological research has been limited. The low adoption of this method may be attributed to individual differences in the absolute values of pulse wave amplitude, which can reduce the effect sizes when comparing different task conditions. Previous studies have observed a decrease in pulse wave amplitude during tasks of increasing difficulty, such as the Sternberg working memory task (Iani et al., 2004), Stroop task (Tulen et al., 1989), and mental arithmetic (Goldstein & Edelberg, 1997). However, these studies typically averaged activity over long periods related to the task, compared with a resting state or tasks of varying difficulty. Consequently, the long-standing tradition in psychophysiological research that focuses on task-evoked changes in activity has not been tested in relation to pulse wave amplitude. To enhance the usefulness and widespread adoption of this measure, it is important to explore the application of baseline normalization technique. Applied to single-trials, baseline normalization can account for individual

differences in vascular physiology and situational factors, thereby improving the reliability and validity of PPG measurements in psychological research.

In the current study, we aimed to characterize cardiovascular responses with high precision in time during performance of the digit span task of varying difficulty. We quantified absolute and, to account for individual differences in peripheral physiology, event-related pulse wave amplitude. For comparison, we also report event-related HR extracted from both PPG and ECG sensors. Additionally, we made a comparison with a more established measure of ANS activity, where quantifying task-evoked changes are the norm – pupil dilation (van der Wel & van Steenbergen, 2018).

Methods

The methods partially overlap with our previously published articles (Kosachenko et al., 2023; Pavlov et al., 2022) where the sample, task, and procedure, as well as the pupillometry data acquisition and analysis are described in greater detail.

Participants

Eighty-six participants completed the task. The participants had normal or corrected-to-normal vision and reported no history of neurological or mental diseases. All the participants were Russian native speakers. Informed consent was obtained from each participant. The experimental protocol was approved by the Ural Federal University ethics committee.

Eight participants were excluded because of technical issues or insufficient data quality in one or more conditions (including resting state). Thus, data of 78 participants (74 females, age: 20.5 ± 3.9 years) entered the analyses.

Task and procedure

Before the WM task, we recorded 4 minutes of eyes-closed resting state ECG and PPG. After that, the participants were given instructions and proceeded with the WM task (see Figure 2; for a more detailed description of the task see Kosachenko et al. (2023)). In this task, participants were either required to memorize sequences of 5, 9, or 13 digits (memory condition) or to listen to them (control condition) without trying to memorize them. Afterward, they were asked to recite each digit in the order in which it was presented (i.e., serial recall). There were 54 control trials and 108 memory trials in 9 blocks separated by self-paced breaks. Each block consisted of 3 control (one of each sequence length) followed by 12 memory (4 trials on each sequence length in random order) followed again by 3 control trials.

Figure 1 – Schematic representation of the English equivalent of the digit span task (stimuli are not to scale). The digits were presented auditorily every 2 s. After the last digit in the sequence, there was 3 s retention interval.

Cardiovascular measures

Cardiovascular measures (ECG and PPG) were continuously acquired using an actiCHamp amplifier (Brain Products) from the auxiliary (AUX) inputs. The ECG was recorded using one channel with the active electrode on the right wrist, the reference electrode on the left wrist, and the ground electrode on the left inner forearm 3 cm distal from the elbow. Finger photoplethysmography was recorded from the left index finger.

Preprocessing of PPG and ECG included the following steps. First, both PPG and ECG data were filtered between 0.5 and 100 Hz, and a notch filter 50 Hz was applied using *eegfiltnew* function in EEGLAB. Then, the data were downsampled to 250 Hz. Next, HRVTool v.1.07 (Vollmer, 2019) was used to detect R peaks in ECG and identify troughs and peaks in PPG. Each trough and peak was visually inspected and incorrectly extracted values were rejected from subsequent analysis. IBI < 0.3 and IBI > 2.4 s and HR and PWA values above and below 3 SDs were also rejected. Pulse wave amplitude was defined as the amplitude difference between consecutive trough and systolic peaks, starting from the trough. After that, the recording was segmented into trials starting from 5.5 s before first digit presentation and ending 12 s after last digit presentation. In the epochs, the instantaneous HR (60/inter-bit intervals) and PWA series were linearly interpolated in 0.5 s bins to obtain HR and PWA series with equally spaced values. Trials with no valid values were removed from the analysis. Out of 12619 trials, 14 were excluded based on this criterion from the HR (ECG) data analysis, 274 from the HR (PPG), and 324 from the PWA analysis.

For the time-locked analyses, after averaging in trials, 3s time interval before first digit onset was used for baseline normalization (averaged in tasks and sequence lengths separately for each participant). Absolute baseline normalization (subtraction) was used for HR values and relative (percent change) normalization was used for PWA values.

Statistics

We used a repeated-measures ANOVA for the analysis of the effect of Task (control, memory) and Load (13 levels corresponding to processing of individual digits irrespectively of the sequence length) on HR (extracted from ECG) and PWA. Another ANOVA comparing baseline

normalized and raw data included factors Task and Sequence (3 levels: 5, 9, and 13 digits). Greenhouse-Geisser adjusted degrees of freedom are reported where applicable.

Pearson correlations were calculated to investigate the relationships among behavioral accuracy, NASA-TLX and the physiological variables. cocor package for R (Diedenhofen & Musch, 2015) was used to compare the correlation coefficients between different measures. We used Hittner, May, and Silver's modification of Dunn and Clark's z using a backtransformed average Fisher's Z procedure to make the comparison (see Diedenhofen & Musch (2015) for references).

The statistical analyses were conducted in R (v. 4.2.2).

Results

Behavior

The average number of correctly recalled digits decreased as the sequence length increased. Specifically, the mean (±SD) for 5-digit sequences was 4.62 ± 0.31, for 9-digit sequences it was 4.18 ± 1.18, and for 13-digit sequences it was 3.57 ± 1.13. Overall, the average number of correctly recalled digits was 4.12 ± 0.81. Full results of the behavioral data analyses are reported elsewhere (Kosachenko et al., 2023).

Task and load effects

The HR extracted from PPG and ECG had similar temporal trajectories (Figure 2c,d) and strongly correlated (r(71) = 0.951, p < 0.001). Therefore, we analyzed statistically only HR from ECG to reduce the number of redundant analyses.

HR was generally higher during the memory task compared to the control task, as indicated by a significant main effect of Task (see Table 1). The effect of Load, however, was only observed in the control task, showing a significant relationship (F(4.02, 309.47) = 61.52, p < .001, η_p^2 = .44). During the memory task, HR exhibited an initial deceleration followed by acceleration, eventually reaching a plateau (Figure 2e). However, Load did not significantly affect HR in the memory task (F(2.21, 170.20) = 1.58, p = .206, η_p^2 = .02). The pairwise comparisons between levels of Load and Tasks are presented in Table S2.

PWA showed different patterns in the control and memory tasks (Table 1). In the memory task, PWA mostly decreased with increasing load (F(1.75, 134.86) = 35.64, p < .001, η_p^2 = .32), while in the control task, it mostly increased (F(2.65, 203.99) = 25.57, p < .001, η_p^2 = .25; see Table S1 for post hoc tests in both Tasks). As can be seen in Figure 2e, in the memory task, PWA initially dropped after the first digit and then remained relatively stable until the fifth digit, after which there was a significant decrease. The pattern reversed after the 12th and 13th digits in the sequence.

Table 1 – The Task (control vs memory) and Load (13 levels) effects on PWA and HR

	Pulse wave amplitude			
	df	*F*	*p*	η^2
Task	1, 77	97.95	<.001	.56
Load	2.45, 188.48	11.76	<.001	.13
Task x Load	1.80, 138.31	53.18	<.001	.41
	Heart rate (ECG)			
Task	1, 77	41.37	<.001	.35
Load	2.67, 205.87	21.66	<.001	.22
Task x Load	3.14, 241.59	31.44	<.001	.29

Figure 2. Temporal dynamics of (a) pupil size; (b) pulse wave amplitude (PWA) registered by PPG; (c) heart rate registered by ECG; and (d) heart rate registered by PPG. Each dashed line represents the onset of a digit presentation. Time 0 indicates presentation of the first digit in the sequence. (e) Pupil size, heart rate (from ECG), and PWA in the memory task averaged in the two-second time intervals corresponding to processing of individual digits in the sequence. Red lines mark a significant decrease and blue lines a significant increase in the parameter compared with the adjacent levels of load. * p<0.05, ** p<0.01, *** p<0.001.

The effect of baseline normalization

To assess the effect of baseline normalization on the cardiovascular measures, we separately averaged raw and baseline normalized PWA and HR in the whole interval corresponding to memorizing or listening to the entire sequence.

The most notable difference was observed in the estimation of PWA (Table 2). Interestingly, due to a cross-over Task x Sequence interaction effect in raw data, main effects were not significant (Figure 3, bottom left panel). Specifically, in the case of 5-digit sequences, raw PWA was larger in the memory task compared to the control task (p = 0.039). In the 9-digit sequences it was, in contrast, slightly (not significantly; p = 0.093) larger in the control task, and the same effect was highly significant in the 13-digit sequences (p = 0.002). After baseline normalization, the effect was highly consistent in all conditions, PWA being significantly larger in the control task than in the memory task. Furthermore, normalization visibly reduced the variability of the data, which may have led to the fact that, within the memory task, the difference between the 9 and 13-digit sequences became significant (p = 0.006) while it was not in raw data (p = .231).

Table 2 – The comparison of baseline normalized and raw PWA values between sequences of 5, 9, and 13 digits in the control and memory tasks

	df	F	p	η^2
	Baseline normalized PWA			
Task	1, 77	97.20	<.001	.56
Sequence	2, 154	3.87	.023	.05
Task x Sequence	2, 154	11.37	<.001	.13
	Raw PWA			
Task	1, 77	1.54	.219	.02
Sequence	1.68, 129.58	0.01	.987	< .01
Task x Sequence	2, 154	19.14	<.001	.20

HR showed a significant decrease in the control task compared to the memory task (Table 3). HR decreased as the sequence length increased in the control task: there was a significant difference in HR between 5-digit sequences and both 9- and 13-digit sequences, as observed in both baseline normalized and raw data (highest p = 0.002). However, in the memory task, neither the baseline normalized nor raw data showed significant pairwise differences in HR between different sequence lengths. Nevertheless, the difference between the tasks remained significant for every sequence length, considering both baseline normalized and raw data.

Table 3 – The comparison of baseline normalized and raw HR (extracted from ECG) values between sequences of 5, 9, and 13 digits in control and memory tasks

	df	F	p	η^2
Baseline normalized HR				
Task	1, 77	27.72	<.001	.26
Sequence	2, 154	2.68	.072	.03
Task x Sequence	2, 154	9.73	<.001	.11
Raw HR				
Task	1, 77	100.22	<.001	.57
Sequence	1.77, 136.53	25.54	<.001	.25
Task x Sequence	1.80, 138.30	26.90	<.001	.26

Figure 3. A comparison of baseline normalized and raw HR and PWA in the sequences of 5, 9, or 13 digits.

To investigate the changes in raw PWA and HR compared to the resting state, we conducted an ANOVA with three levels of Task factor: memory, control, and rest. As expected, the effects of Task were significant on both PWA ($F(1.05, 80.82) = 15.83$, $p < .001$, $\eta_p^2 = .17$) and HR ($F(1.19, 91.62) = 11.32$, $p < .001$, $\eta_p^2 = .13$).

PWA during rest was significantly larger compared to both memory (t(77) = 4.152, p < 0.001, d = 0.473) and control tasks (t(77) = 3.844, p < 0.001, d = 0.438), while the difference between memory and control tasks was marginally significant (t(77) = 2.007, p = 0.048, d = 0.229). The corresponding means (± standard deviations) were 5.14±3.76, 3.93±2.16, 4.03±2.16, for rest, memory, and control, respectively.

Regarding HR, there was a significant difference between the rest and memory tasks (mean+SD; rest: 77.8±11.5, memory: 79.5±10.9 b/min; t(77) = 2.627, p = 0.01, d = 0.29). However, no significant difference was observed between the rest and control tasks (control: 77±10.5 b/min; t(77) = 1.217, p = 0.227, d = 0.139). Additionally, the control task had significantly lower HR compared to the memory task (t(77) = 10.64, p < 0.001, d = 1.212).

The relationship between physiological responses and memory accuracy

Next, to understand the relationship between cardiovascular reactivity measures and behavioral performance, we conducted a correlational analysis. In this analysis, we used PWA and HR values averaged across listening and memorizing all 13 digits, along with the average number of correctly recalled digits in the correct order. Additionally, to compare with a well-established index of peripheral nervous system activity known to be associated with cognitive load and effort, we incorporated pupillometry data. Detailed pupillometry data analysis can be found elsewhere (Kosachenko et al., 2022). Here, a figure displaying pupil size dynamics is presented alongside the PPG and ECG data (Figure 2a).

The results revealed significant associations between pupil size and both HR and PWA (Figure 4). In the memory task, the correlation between pupil size and PWA displayed a numerically stronger relationship compared to the pupil size-HR correlation, although the difference did not reach statistical significance (z = -1.627, p = 0.052). Moreover, all physiological variables demonstrated correlations with accuracy in the task. Among the physiological variables examined, only PWA significantly correlated with subjective assessment of cognitive load (r(76) = -0.23, p = 0.038) as measured by the NASA-TLX scale (see the extended correlation matrix in Figure S1).

Figure 4. Correlations between the variables, density distributions, and scatterplots for each of the variables. The physiological parameters are averaged over all conditions and times (from presentation of the first digit till end of the retention interval after the last digit in the sequence). · p < 0.1 * p < 0.05, ** p < 0.01, *** p < 0.001

Discussion

In this study, we aimed to assess the effectiveness of PWA as a measure for tracking cognitive load in comparison to the commonly used cardiovascular reactivity measure, HR. Unlike HR, which showed no significant changes with difficulty of the task, we observed a strong suppression of PWA with increasing memory load. Moreover, when memory load exceeded the capacity of working memory, this pattern reversed, indicating memory overload. Additionally, only PWA during the memory task correlated with self-reported cognitive load. Furthermore, both HR and PWA demonstrated a relationship with behavioral performance, as a higher task-evoked HR and lower PWA were associated with a greater number of correctly recalled memory items. These findings strongly support the notion that PWA is sensitive to cognitive overload and holds significant potential for practical application due to the ease of obtaining PPG data.

We have demonstrated the comparable effectiveness of task-evoked HR measurements using PPG and ECG, suggesting the interchangeability of these measures in mental load research. However, heart rate was not sensitive to small changes in cognitive load and associated mental

effort. In studies examining cognitive load and its impact on physiological responses, it has been observed that HR is capable of distinguishing between control and memory tasks but lacks sensitivity in detecting differences between levels of cognitive load. This finding aligns with a working memory study conducted by Iani et al. (2004), which also failed to identify a general effect of memory load on HR. This lack of sensitivity may be due to the influence of different branches of ANS: the simultaneous activity of both branches of the ANS may have canceled out the weaker effects of memory load (as compared with the effect of task) on HR.

Peripheral vessel tone as expressed in PWA was changing as a function of cognitive task and load. Vasoconstriction is regulated solely by sympathetic activations (Babchenko et al., 2001; Shelley, 2007). As stated by motivation intensity theory, effort entails the mobilization of resources for instrumental behavior, and effort is more directly related to activation of the sympathetic nervous system (Gendolla & Wright, 2012). Among all studied physiological parameters, only PWA was related to self-assessed expression of cognitive load during the tasks as measured by NASA-TLX scales. The task-evoked PWA not only decreased in response to cognitive load but also displayed a pattern of overload that was reflected in a reverse of the trajectory of PWA after about 11-digit load. This uptrend in PWA can be interpreted as a release of effort due to cognitive overload. Although the exact timing of this event may not be precisely detected by PWA, the fact that the overload was detected makes the measure an important non-invasive marker of cognitive load, largely overlooked by previous studies.

HR strongly differed between control and memory tasks independently of whether baseline normalization was applied or not. Baseline normalization appeared to be more crucial for accurate PWA measurements compared to HR. Large individual variability of indexes derived from the pulse wave has been reported previously (Lyu et al., 2015). The dramatic improvement in the assessment of cognitive load in PWA after baseline normalization can shed light on the mixed results observed in previous studies that aimed to differentiate between levels of task difficulty (Iani et al., 2004; Xuan et al., 2020). Therefore, we suggest that future investigations consider trial-level baseline PWA values for accurate comparisons across cognitive load levels and to establish consistent associations with cardiovascular responses.

In our previous study, we showed that a well-established correlate of effort and cognitive load – pupil dilation – peaked at 8 items load before a period of constriction returning pupil size to baseline level. Here, PWA peaked after the presentation of the 10th digit. At the 11[th] and 12[th] digit, both pupil response and PWA reversed their patterns observed with lower load levels. HR remained roughly on the same level after presentation of the fifth digit in the sequence. It is known that pupil size is influenced by both the sympathetic and parasympathetic nervous systems. Steinhauer et al. (2004) have come to the conclusion that emotional arousal primarily affects pupil dilation through sympathetic activation, while the similar dilation in response to cognitive load is primarily affected by the inhibition of the parasympathetic system. However, the current study revealed a strong correlation between task-evoked pupil size and PWA, which included, first, their simultaneous increase with cognitive load, and second, a similar drop in the amplitude of both responses in the most challenging conditions, thus indicating a shared underlying mechanism. The similarity between pupil size and PWA is additionally stressed by the fact that no parallel drop of response was observed in HR, controlled by both autonomous branches. We propose, therefore, that alpha adrenergic sympathetic activation is involved in

both task-evoked pupil dilation and peripheral vasoconstriction during cognitive tasks. This activation grows with increasing load and decreases with overload.

In our study, PWA and pupil size have emerged as valuable metrics for investigating the intensity of effort, both reflecting the response of the sympathetic nervous system to the increased cognitive load. Compared with the pupil dilation, PWA offers advantages such as simplicity, independence from luminosity and lighting conditions, and versatility. The alternative means for monitoring sympathetic activation such as heart rate variability and skin conductance operate on much larger time scales, while measuring pre-ejection period is distinctively more technologically challenging. In contrast, the PWA response is immediate, and the recording of PPG is possible with modern smartphones and, thanks to recent developments, via a web camera remotely (van Es et al., 2023), which opens up new avenues for research and practical utilization of these measures. These developments have the potential to expand the application of PPG and PWA to enhance our understanding and assessment of cognitive load in various contexts.

Author contributions statement

Yuri G. Pavlov: Conceptualization, Funding acquisition, Data curation, Formal analysis, Project administration, Supervision, Visualization, Methodology, Writing - original draft, Writing - review & editing.

Anastasia S. Gashkova: Investigation, Formal analysis, Writing - original draft, Writing - review & editing.

Alexandra I. Kosachenko: Investigation, Methodology, Data curation, Writing - review & editing.

Dauren Kasanov: Investigation, Methodology, Data curation, Writing - review & editing.

Alexander I. Kotyusov: Investigation, Methodology, Data curation, Writing - review & editing.

Boris Kotchoubey: Writing - review & editing.

Data availability statement

The data are publicly available on OpenNeuro (https://openneuro.org/datasets/ds003838).

Potential Conflicts of Interest

Nothing to report.

Acknowledgements

The research funding from the Ministry of Science and Higher Education of the Russian Federation (Ural Federal University Program of Development within the Priority-2030 Program) is gratefully acknowledged.

References

Al abdi, R. M., Alhitary, A. E., Abdul Hay, E. W., & Al-bashir, A. K. (2018). Objective detection of chronic stress using physiological parameters. *Medical & Biological Engineering & Computing*, *56*(12), 2273–2286. https://doi.org/10.1007/s11517-018-1854-8

Babchenko, A., Davidson, E., Ginosar, Y., Kurz, V., Faib, I., Adler, D., & Nitzan, M. (2001). Photoplethysmographic measurement of changes in total and pulsatile tissue blood volume, following sympathetic blockade. *Physiological Measurement*, *22*(2), 389. https://doi.org/10.1088/0967-3334/22/2/310

Diedenhofen, B., & Musch, J. (2015). cocor: A comprehensive solution for the statistical comparison of correlations. *PloS One*, *10*(4), e0121945. https://doi.org/10.1371/journal.pone.0121945

Gendolla, G., & Wright, R. (2012). Effort Intensity: Some Insights From the Cardiovascular System. In *The Oxford Handbook of Human Motivation* (pp. 420–438). https://doi.org/10.1093/oxfordhb/9780195399820.013.0024

Goldstein, H. S., & Edelberg, R. (1997). A plethysmographic method for demonstrating the response specificity of the oral vascular bed. *Psychophysiology*, *34*(1), 124–128. https://doi.org/10.1111/j.1469-8986.1997.tb02423.x

Iani, C., Gopher, D., & Lavie, P. (2004). Effects of task difficulty and invested mental effort on peripheral vasoconstriction. *Psychophysiology*, *41*(5), 789–798. https://doi.org/10.1111/j.1469-8986.2004.00200.x

Kosachenko, A. I., Kasanov, D., Kotyusov, A. I., & Pavlov, Y. G. (2023). EEG and pupillometric signatures of working memory overload. *Psychophysiology*, *n/a*(n/a), e14275. https://doi.org/10.1111/psyp.14275

Lyu, Y., Luo, X., Zhou, J., Yu, C., Miao, C., Wang, T., … Kameyama, K. (2015). Measuring Photoplethysmogram-Based Stress-Induced Vascular Response Index to Assess Cognitive Load and Stress. *Proceedings of the 33rd Annual ACM Conference on Human Factors in Computing Systems*, 857–866. https://doi.org/10.1145/2702123.2702399

Mandrick, K., Peysakhovich, V., Rémy, F., Lepron, E., & Causse, M. (2016). Neural and psychophysiological correlates of human performance under stress and high mental workload. *Biological Psychology*, *121*, 62–73. https://doi.org/10.1016/j.biopsycho.2016.10.002

Matsumura, K., & Yamakoshi, T. (2013). iPhysioMeter: A new approach for measuring heart rate and normalized pulse volume using only a smartphone. *Behavior Research Methods*, *45*(4), 1272–1278. https://doi.org/10.3758/s13428-012-0312-z

Pavlov, Y. G., Kasanov, D., Kosachenko, A. I., Kotyusov, A. I., & Busch, N. A. (2022). Pupillometry and electroencephalography in the digit span task. *Scientific Data*, *9*(1), 325. https://doi.org/10.1038/s41597-022-01414-2

Quintana, D. S., Alvares, G. A., & Heathers, J. a. J. (2016). Guidelines for Reporting Articles on Psychiatry and Heart rate variability (GRAPH): Recommendations to advance research communication. *Translational Psychiatry*, *6*(5), e803–e803. https://doi.org/10.1038/tp.2016.73

Richter, M., Friedrich, A., & Gendolla, G. H. E. (2008). Task difficulty effects on cardiac activity. *Psychophysiology*, *45*(5), 869–875. https://doi.org/10.1111/j.1469-8986.2008.00688.x

Schapkin, S. A., Freude, G., Gajewski, P. D., Wild-Wall, N., & Falkenstein, M. (2012). Effects of Working Memory Load on Performance and Cardiovascular Activity in Younger and Older Workers. *International Journal of Behavioral Medicine*, *19*(3), 359–371. https://doi.org/10.1007/s12529-011-9181-6

Shelley, K. H. (2007). Photoplethysmography: Beyond the Calculation of Arterial Oxygen Saturation and Heart Rate. *Anesthesia & Analgesia*, *105*(6), S31. https://doi.org/10.1213/01.ane.0000269512.82836.c9

Steinhauer, S. R., Siegle, G. J., Condray, R., & Pless, M. (2004). Sympathetic and parasympathetic innervation of pupillary dilation during sustained processing. *International Journal of Psychophysiology*, *52*(1), 77–86. https://doi.org/10.1016/j.ijpsycho.2003.12.005

Tulen, J. H. M., Moleman, P., van Steenis, H. G., & Boomsma, F. (1989). Characterization of stress reactions to the Stroop Color Word Test. *Pharmacology Biochemistry and Behavior*, *32*(1), 9–15. https://doi.org/10.1016/0091-3057(89)90204-9

van der Wel, P., & van Steenbergen, H. (2018). Pupil dilation as an index of effort in cognitive control tasks: A review. *Psychonomic Bulletin & Review*, *25*(6), 2005–2015. https://doi.org/10.3758/s13423-018-1432-y

van Es, V. A. A., Lopata, R. G. P., Scilingo, E. P., & Nardelli, M. (2023). Contactless Cardiovascular Assessment by Imaging Photoplethysmography: A Comparison with Wearable Monitoring. *Sensors*, *23*(3), 1505. https://doi.org/10.3390/s23031505

Vincent, A., Craik, F. I. M., & Furedy, J. J. (1996). Relations among memory performance, mental workload and cardiovascular responses. *International Journal of Psychophysiology*, *23*(3), 181–198. https://doi.org/10.1016/S0167-8760(96)00058-X

Vollmer, M. (2019). HRVTool–an open-source matlab toolbox for analyzing heart rate variability. *2019 Computing in Cardiology (CinC)*, Page-1. https://doi.org/10.22489/CinC.2019.032

Wright, R. A., & Kirby, L. D. (2001). Effort determination of cardiovascular response: An integrative analysis with applications in social psychology. In *Advances in Experimental Social Psychology* (Vol. 33, pp. 255–307). https://doi.org/10.1016/S0065-2601(01)80007-1

Xuan, Q., Wu, J., Shen, J., Ji, X., Lyu, Y., & Zhang, Y. (2020). Assessing cognitive load in adolescent and adult students using photoplethysmogram morphometrics. *Cognitive Neurodynamics*, *14*(5), 709–721. https://doi.org/10.1007/s11571-020-09617-2