Histone binding of ASF1 is required for fruiting body development, but not for genome

stability in the filamentous fungus *Sordaria macrospora*

Jan Breuer[1], Tobias Busche[2,3], Jörn Kalinowski[2], Minou Nowrousian[1]

[1]Department of Molecular and Cellular Botany, Ruhr University Bochum, 44801 Bochum, Germany

[2]Center for Biotechnology, Bielefeld University, 33594 Bielefeld, Germany

[3]Medical School OWL, Bielefeld University, 33615 Bielefeld, Germany

running title:

ASF1 in fungal development

author for correspondence:

Minou Nowrousian

Department of Molecular and Cellular Botany

Ruhr University Bochum

Universitaetsstr. 150

44801 Bochum

minou.nowrousian@rub.de

Abstract

The highly conserved eukaryotic histone chaperone ASF1 is involved in the assembly and disassembly of nucleosomes during transcription, DNA replication and repair. It was the first chaperone discovered to be involved in all three of these processes. The filamentous fungus *Sordaria macrospora* is one of only two multicellular organisms where *asf1* deletions are viable, which makes it useful for *in vivo* analysis of this central part of eukaryotic chromatin structure. Deletion of *asf1* in *S. macrospora* leads to sterility, a reduction of DNA methylation, and upregulation of genes that are usually weakly expressed in the wild type. Here, we focused on the functions of the highly conserved core and the divergent C-terminal tail of ASF1, studied the effects of ASF1 on histone modifications and tested its relevance for genomic stability. By Co-IP and complementation analysis we showed that substitutions of amino acid V94 or truncations of the C-terminal tail abolish histone binding and do not complement the sterile mutant phenotype. Δasf1 is sensitive to the DNA damaging agent MMS, while complementation strains, even those with non-histone-binding variants, regain wild type-like resistance. To aid in subsequent ChIP-seq and Hi-C analyses, we generated a chromosome-resolved genome assembly of *S. macrospora*. ChIP-Seq analysis revealed a global increase of H3K27me3 in Δasf1, accompanied by a global decrease of H3K56ac. By using Hi-C we detected a tandem duplication of around 600 kb on chromosome 2 in the mutant. Crossing experiments indicated linkage between the viability of Δasf1 strains and the presence of the duplication.

Importance

Histone chaperones are proteins that are involved in nucleosome assembly and disassembly and can therefore influence all DNA-dependent processes including transcription, DNA replication and repair. ASF1 is a histone chaperone that is conserved throughout eukaryotes. In contrast to most other multicellular organisms, a deletion mutant of *asf1* in the fungus *Sordaria macrospora* is viable; however, the mutant is sterile. In this study, we could show that the histone binding ability of ASF1 is required for fertility in *S. macrospora*, whereas the function of ASF1 in maintenance of genome stability does not require histone binding. We also showed that the histone modifications H3K27me3 and H3K56ac are misregulated in the Δasf1 mutant. Furthermore, we identified a large duplication on chromosome 2 of the mutant strain that is genetically linked to the Δasf1 allele present on chromosome 6, suggesting that viability of the mutant might depend on the presence of the duplicated region.

Introduction

Multicellular development is a complex process that evolved independently in multiple branches of eukaryotes (1). In fungi, multicellular development evolved at least twice and possibly more times, the most prominent examples being the fruiting bodies of filamentous ascomycetes and basidiomycetes (2). To study the genetic background of multicellular development, simple, easy to handle model organisms are of particular interest. Several species of fungi have proven to be usable models for the analysis of developmental processes and the generation of complex, multicellular fruiting bodies of filamentous ascomycetes is an excellent example (3). *Sordaria macrospora*, a homothallic ascomycete that can form its perithecia in 7 days under laboratory conditions, has been used as a model to study fungal sexual development and multicellular development in general. During the formation of fruiting bodies in *S. macrospora*, at least 13 different cell types are generated, many of which cannot be found in vegetative mycelium (4). Sexual development and perithecia formation are accompanied by transcriptional changes, and transcriptomics can be a helpful tool to identify important developmental genes by reverse genetic approaches. A number of transcription factors and chromatin modifiers have been identified this way and were proven to be fundamental for multicellular development (5). Among them is the conserved histone chaperone ASF1 that was shown to be upregulated during sexual development and is essential for this process in *S. macrospora* (6). ASF1 is involved in the assembly and disassembly of nucleosomes during transcription, replication and DNA-repair and was the first chaperone found to be involved in all three of these processes. It mediates nucleosome assembly and disassembly in promoter regions during transcriptional activation and elongation in *Saccharomyces cerevisiae* (7). ASF1 is also important for histone modification by interacting with enzymes that mediate the corresponding modifications. For example, it has been shown to be involved in the acetylation of histone H3 at lysines 9 and 56 and also to play a role in the genome-wide deacetylation of histones (8, 9). The methylation of histones also appears to proceed in conjunction with ASF1. In *Mus musculus*, an altered distribution of H3K9me3 and H3K27me3 was shown under ASF1 deficiency. ASF1 was also proven to be important for histone recycling (10). Additionally, ASF1-depleted cells were shown to exhibit higher tendencies for double strand breaks, showcasing a role of ASF1 during the maintenance of genomic stability (11). The function and structure of ASF1 is highly conserved in all eukaryotes. The 155 amino acids of the N-terminus of ASF1 form a globular core with conserved acidic patches that mediate interaction with the C-terminus of histone H3. ASF1 has been shown to interact with the histone H3-H4 heterodimer *in vitro* and *in vivo*. In addition to binding of H3, binding of the H4 C-terminus to ASF1 also occurs. ASF1 and the H3-H4 dimer can form a heterotrimeric complex, although both histones can be bound individually by ASF1 (7). The central role of ASF1 for eukaryotic chromatin structure is underscored by findings in human cell lines that showed all non-DNA bound histones H3 and H4 are bound to ASF1 (12, 13). Therefore, researching ASF1 functions during multicellular

development might be fundamental for understanding the contribution of the chromatin landscape to

development in fungi and multicellular eukaryotes in general. Gaining insight into the functions of ASF1

is hindered by the fact that deletions of the respective gene are lethal in almost all higher eukaryotes.

Thus, research on ASF1 was mostly conducted with ASF1-depleted cell lines, *in vitro* experiments and

studies of the unicellular *S. cerevisiae* where an *asf1* deletion is viable (14-16)*. S. macrospora* and

Arabidopsis thaliana were shown to be exceptions in this regard and deletions are viable in these

organisms (6, 17). Therefore, the analysis of *S. macrospora* Δasf1 strains offers the unique opportunity

to perform ASF1 research *in vivo* in a multicellular fungus. These strains were shown to be sterile,

highlighting a central role of ASF1 during sexual development. Unexpectedly, the positioning of

nucleosomes was not affected in *asf1* deletion strains. A significant reduction of DNA methylation was

observed in *S. macrospora* Δasf1, and a large number of genes that are usually weakly expressed in

wild type strains are upregulated in Δasf1, indicating a role of ASF1 during gene regulation and DNA

methylation (18). The *S. macrospora* ASF1 has been shown to bind histones H3 and H4 (6), however, it

has not yet been analyzed if the histone-binding activity is actually required for fruiting body formation.

Therefore, in this study, we analyzed the histone binding function of ASF1 and its connection to sexual

development. We performed interaction studies and complementation analysis to verify the

importance of amino acid V94 (valine at position 94), a highly conserved histone interaction site known

from *in vitro* studies (7, 19). The C-terminal region of ASF1 is much less conserved than the first 155

amino acids and parts of it exist only in fungi (Figure 1). We generated truncated versions of this region

of *asf1* and used these to perform interaction studies and complementation analysis to study the

relevance of the divergent and functionally less well studied C-terminus of ASF1. Additionally, we

performed genomic stability assays based on growth under the influence of the genotoxic substance

methyl methansulfonate (MMS). We further focused on the role of ASF1 in histone modification and

chromatin structure by performing ChIP (chromatin immunoprecipitation)-seq and Hi-C (chromosome

conformation capture with high-throughput sequencing) experiments with the wild type and the Δasf1

mutant. For improved analysis of the ChIP-seq and Hi-C data, we first generated a chromosome-level

genome sequence of *S. macrospora*. Participation of ASF1 in histone acetylation and deacetylation has

been documented in *S. cerevisiae*, and a function during histone methylation is regarded as highly

probable and connections to the effect on DNA-methylation and gene regulation might be possible

(10). Therefore, ChIP-seq was performed for H3K27me3 and H3K56ac. Furthermore, we performed Hi-

C experiments to compare overall chromatin structure of young, vegetative mycelium to older, sexual

mycelium and *asf1* deletion mutants.

Materials and Methods

Strains, crosses and growth conditions

All strains used in this study have been summarized in Supplemental Table 1 and were cultivated on either solid or liquid cornmeal medium (BMM) or complete medium (CM) at 25 °C as previously described (20, 21). To perform genetic crosses, the spore color mutant fus was used as a partner to allow the identification of recombinant asci (22). Transformation of *S. macrospora* was performed as previously described (21).

Cloning procedures, oligonucleotides and plasmids

Plasmids encoding ASF1 variants with amino acid substitutions for complementation and interaction analysis were generated by PCR mutagenesis and yeast recombinant cloning (23). Golden Gate cloning was used to assemble plasmids encoding truncated ASF1 variants (24). Oligonucleotides for the generation of plasmids and integration tests have been listed in Supplemental Table 2. All plasmids used in this study are listed in Supplemental Table 3.

Histone interaction assay by Co-Immunoprecipitation (Co-Ip)

Plasmids coding for eGFP-tagged ASF1 variants were co-transformed together with plasmids encoding FLAG-tagged histones H3 or H4 into the *S. macrospora* wild type strain SN1693 by protoplast transformation (25). The resulting strains were grown for 4 d at 27 °C in 20 ml liquid BMM medium with 50 µg / ml Nourseothricin. The mycelium was harvested by filtration and frozen in liquid nitrogen before being ground into powder and mixed with protein extraction buffer (26). Co-Immunoprecipitation was performed according to the manufacturers' instructions using GFP-Trap® Beads (Chromotek) and ANTI-FLAG® M2 Beads (Sigma Aldrich). Raw and elution fractions were used for SDS-PAGE (27) before being transferred to a PVDF membrane by western blotting. Co-Ip results were generated with two biological replicates each for eGFP trapping and FLAG trapping. Signals for eGFP and FLAG were detected by using Mouse Anti-GFP JL-8 (Clontech), Mouse Anti-Flag (Sigma Aldrich) and Goat Anti-Mouse, HRP-linked (Cell Signaling) antibodies according to the manufacturers' instructions.

Complementation of *S. macrospora* with ASF1 variants and MMS tests

Plasmids for the expression of ASF1 variants were generated from wild type genomic DNA with specific primers amplifying *asf1* with specific substitutions or a shortened C-terminus and cloned by Golden Gate Cloning (24) or via yeast homologous recombination (28). The *asf1* deletion strain SN1983 was transformed with the respective plasmids by protoplast transformation (25). Complementation strains

that did not gain fertility on their own were crossed against the wild type, and ascospores were isolated

to obtain homokaryotic strains, while fertile transformants were used to isolate ascospores directly

(22). Phenotypical characterization was performed by documenting the growth and fertility on BMM

plates with a Stemi 2000-C stereomicroscope (Zeiss). To evaluate the resistance against methyl

methanesulfonate (MMS) all strains were inoculated on BMM plates with 0,007 % MMS and growth

observed over a span of 7 d.

Microscopic analysis

For analysis of the subcellular localization of ASF1-EGFP fusion proteins, strains were grown on glass

slides together with strains expressing H3-MRFP fusion proteins as described (29). Light and

fluorescence microscopy was performed with an AxioImager microscope (Zeiss, Jena, Germany) with

a Photometrix Cool SnapHQ camera (Roper Scientific). EGFP fluorescence was detected with Chroma

(Bellows Falls, VT, USA) filter set 41017 (HQ470/40, HQ525/50, Q495lp) and mRFP fluorescence was

detected with set 49008 (EG560/40x, ET630/75 m, T585lp). Images were further handled by using

MetaMorph (Molecular Devices).

Extraction and sequencing of genomic DNA from *S. macrospora*

For DNA preparation for nanopore sequencing, the wild type strain SN1693 was grown for 2 d at 27 °C

in 20 ml liquid CM medium (25) in petri dishes, harvested by filtration and ground in liquid nitrogen.

500 µg of ground mycelium was used for extraction of high molecular weight DNA with the NucleoBond

HMW-Kit (Macherey-Nagel, Düren, Germany) according to the manufacturer's instructions. A

sequencing library with wild type genomic DNA was prepared using the Nanopore DNA Ligation

Sequencing Kit (SQK-LSK109, Oxford Nanopore Technologies, Oxford, UK) according to the

manufacturer's instructions. Sequencing was performed on an Oxford Nanopore GridION Mk1

sequencer using an R9.4.1 flow cell, which was prepared according to the manufacturer's instructions.

Basecalling was performed using guppy v5.0.11 with the super-accurate basecalling model. A total of

290927 reads were obtained with an average read length of 15874 bp and an N50 of 23018 bp.

For DNA preparation for Illumina sequencing of the Δasf1 mutant, the Δasf1 strain SN1983 was grown

for 3 d at 27 °C in 20 ml liquid CM medium in petri dishes, harvested by filtration and ground in liquid

nitrogen. DNA preparation was performed as described previously for *Pyronema confluens* (30). Library

preparation and Illumina sequencing of a 300 bp insert library (150 bp paired-end sequencing) were

performed at Novogene (Cambridge, UK).

Genome assembly and annotation

Nanopore reads were assembled with SMARTdenovo (available at
https://github.com/ruanjue/smartdenovo, accessed on 2021/06/29) resulting in nine contigs with a
total size of 39.4 Mb. The initial assembly was subjected to four rounds of correction with Racon (31)
using the nanopore reads for correction, and subsequently three rounds of correction with pilon
(v1.24) (32) based on previously generated *S. macrospora* genomic Illumina reads (33) (NCBI SRA
database, accession number SRR5749461). BLAST comparisons (34) with a previously generated *S.*
macrospora genome assembly (35) showed that the smallest contig (161 kb) most likely contained
mitochondrial DNA. This contig was removed, and the remaining eight contigs were analyzed for
putative telomeric repeats (sequence TTAGGG) using a custom-made perl program. Telomeric repeats
were found at both ends of five contigs, and at one end each in the remaining three contigs. Of the
latter, one contig contained the rDNA repeats at the end without telomeric repeats. The other two
non-telomeric contig ends were compared to each other and found to have an overlap of 2.2 kb with
100 % sequence identity, making it likely that these contigs were part of the same chromosome. The
contigs were fused manually, and the junction was checked for continuity by mapping the nanopore
reads to the assembly with graphmap (36). The resulting assembly consists of 7 contigs with a total
size of 39.4 Mb and an N50 of 5.7 Mb. Comparison to the genome assembly of the closely related
ascomycete *Neurospora crassa* (FungiDB version 52) (37, 38) was performed with nucmer from the
MUMmer package (v4.0.0rc1) (39).
Genome annotation of protein-coding genes was performed with MAKER (v3.01.03) (40) based on
transcripts and proteins from a previous annotation of the *S. macrospora* genome (33) as well as on
the *N. crassa* proteins (38) from FungiDB version 52 (37). tRNAs were annotated with tRNAscan-SE
(v2.0.8) (41). rDNA repeats were annotated manually based on BLAST comparisons. *S. macrospora*
locus tags from a previous annotation (33) were mapped onto the predicted gene models based on
BLAST results. Approximately 1200 gene models were manually corrected in the genome browser
Artemis (42) based on RNA-Seq data generated previously (18, 43) (NCBI GEO accession numbers
GSE33668 and GSE92337) that were mapped to the assembly with Hisat2 (v2.2.1) (44).
223
Western blot analysis of histone modifications and chromatin immunoprecipitation
Potential targets for ChIP-seq experiments were selected based on semiquantitative western blot
screenings with antibodies against histone modifications previously suspected to be influenced by
ASF1. The wild type strain SN1693 and the Δasf1 strain SN1983 were cultivated in liquid CM medium
(25) for 3d and 4d, respectively, at 27 °C, harvested by filtration and washed with PPP (21). Samples
were frozen in liquid nitrogen and ground into a fine powder. Proteins were extracted by mixing the
powder with extraction buffer (50 mM Tris/HCl pH 7.5, 250 mM NaCl, 0.05 % NP-40, 0.05 % β-
mercaptoethanol, 0,3 % (v / v) Protease Inhibitor Cocktail Set IV (Calbiochem)) and 20 min

centrifugation. Protein concentrations were determined by Bradford assays (45) and equal amounts

were loaded and separated by SDS gelelectrophoresis and transferred to a PVDF membrane by western

blotting. Antibodies against target histone modifications were used to detect respective bands (Anti

H3K27Me3 #9733 Cell Signaling, Anti H3K9Me #ABE101 Merck Millipore, Anti H3K56Ac #61061 Active

Motif, Anti H3K9Ac #39038 Active Motif), detection protocols have been summarized in Supplemental

Table 4. Band strengths were compared by using a ChemiDoc XRS+ system (BioRad) with Image Lab 4.0

software (BioRad). Normalization for the total amounts of protein in the samples was based on

Coomassie-stained gels run in parallel. Modifications that exhibited significant differences between

wild type and deletion mutant were selected for ChIP-seq experiments. ChIP-samples were cultivated

similar to the samples for western blotting and crosslinking was performed in PPP buffer with 1%

formaldehyde for 15 min at room temperature, followed by a quenching time of 15 min by an excess

amount of glycin. After three PPP washing steps, lysis was conducted in extraction buffer (50 mM

Tris/HCl pH 7.5, 250 mM NaCl, 0.05 % NP-40, 0.05 % β-mercaptoethanol, 0,3 % (v/v) Protease Inhibitor

Cocktail Set IV (Calbiochem)) for 15 min on ice by frequent vortexing. After centrifugation, chromatin

solution was obtained and mixed with MNase Buffer (18) and proteinase inhibitors. Chromatin

digestion was conducted by incubation with 200 U/ml MNase (Thermo Fisher Scientific) for 105 min at

37 °C under constant shaking and stopped by adding EDTA to a concentration of 20 mM. Aliquots were

removed as input samples, and immunoprecipitation performed over night at 4 °C on a spinning wheel

by adding 10 µl of the respective antibody (Anti H3K27Me3 #9733 Cell Signaling, Anti H3 #2650 Cell

Signaling, Anti H3K56Ac #61061 Active Motif). Afterwards, 4h of incubation on Protein-A agarose

beads (Santa Cruz) were followed by 2 washing steps in low salt and 2 washing steps in high salt buffer.

Each washing step was performed for 5 min at 4 °C on a spinning wheel and followed by 2 min of

centrifugation at 2500 rpm to separate the beads from the buffer (46). Chromatin was washed off the

beads by 2 h of incubation in TE buffer with 1 % SDS at 65 °C and de-crosslinked over night at 65 °C

after adding 2 mg/ml Proteinase K (Genaxxon bioscience). Standard phenol/chloroform extraction was

performed to isolate DNA from the ChIP and input samples. Two biological replicates each were

created for H3, H3K27me3 and H3K56ac. Library generation and Illumina sequencing (150 bp paired-

end) of ChIP and input samples were performed by Novogene (Cambridge, UK). The detailed ChIP-seq

protocol can be found in Supplemental Text 1.

ChIP-seq peak detection

20 million paired-end reads per sample were obtained, trimmed with fastp 0.23.2 (47) and mapped to

the reference genome of *S. macrospora* SN1693 by using BWA-MEM 0.7.17 with default parameters

(48). Input corrected peak detection was performed for the duplicated samples of H3, which were

subsequently used together with the native input samples as control for H3K27me3 and H3K56ac by

using MACS2 callpeak 2.2.7.1 in paired end mode with a 0.01 minimum FDR (false discovery rate) cutoff
for broad regions and an effective genome size of 39 Mb (49, 50). Differential peak analysis was
conducted by discarding overlapping peaks in wildtype and Δasf1 using BEDTools 2.30.0 with a
minimum overlap of 10 % (51). Peaks were annotated to the reference genome using HOMER
annotatePeaks 4.9.1 (52).

Hi-C sample preparation
Young wild type samples of SN1693 where cultivated for 2 d at 27 °C in liquid CM medium (25). Samples
for the older wild type where grown for 5 d and Δasf1 SN1983 for 3 d under the same conditions.
Mycelium was harvested by filtration, washed with PPP (21) and frozen in liquid nitrogen. The Omni-C
library was prepared using the Dovetail® Omni-C® Kit according to the manufacturer's protocol with
additional changes to enable the use on *S. macrospora*. Briefly, the chromatin was fixed with 1 %
formaldehyde for 10 min at room temperature on a spinning wheel and quenched by bringing the
solution up to 200 mM glycine. The cross-linked chromatin was then digested in situ with DNase I
according to the manufacturer's instructions. Following digestion, the cells were lysed in 1 % SDS to
extract the chromatin fragments and the chromatin fragments were bound to Chromatin Capture
Beads. Next, the chromatin ends were repaired and ligated to a biotinylated bridge adapter followed
by proximity ligation of adapter-containing ends by using the reagents supplied by the Dovetail® Omni-
C® Kit. After proximity ligation, the crosslinks were reversed at 68 °C in Dovetail® Crosslink Reversal
Buffer and 1 µg / µl Proteinase K overnight. The DNA was purified with SPRIselect™ Beads (Beckmann
Coulter) and 150 ng were converted into a sequencing library using Illumina-compatible adaptors by
using the Dovetail® Library Module for Illumina®. Biotin-containing fragments were isolated using
streptavidin beads prior to PCR amplification. Illumina sequencing (150 bp paired-end) was performed
by Novogene (Cambridge, UK).

Generation of Hi-C maps
Hi-C maps were created by following the Dovetail® Omni-C® Pipeline with adjustments for the genome
of *S. macrospora*. 50 million paired end reads were obtained, trimmed with fastp 0.23.2 (47) and
mapped to the reference genome of *S. macrospora* SN1693 by using BWA-MEM 0.7.17 with default
parameters (48). Valid ligation events were recorded by using pairtools 1.02 (53) parse with a min-map
quality of 40 and a max-inter-align-gap of 30 before being sorted by pairtools sort. PCR duplicates were
removed by using pairtools dedup and the pairs file was generated by pairtools split. The hic matrix
was generated by using juicertools 1.6.2 (54) with standard parameters.

Data availability

Nanopore reads and the chromosome-resolved genome sequence for the *S. macrospora* wild type

strain SN1693 were deposited in the NCBI databases under BioProject ID PRJNA744349

(http://www.ncbi.nlm.nih.gov/bioproject/744349, SRA accession number SRR15901728 for the

nanopore reads, DDBJ/ENA/GenBank accession numbers CP083903-CP083909 for the seven

chromosomes). ChIP-seq and Hi-C data generated in this study were submitted to the NCBI GEO

database   as   a   superseries   under   accession   number   GSE216634

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE216634, accession numbers GSE216629

and GSE228529 for ChIP-seq data and GSE216632 for Hi-C data). Illumina reads from the Δasf1 strain

were   deposited   in   the   NCBI   SRA   database   under   BioProject   ID   PRJNA922040

(https://www.ncbi.nlm.nih.gov/sra/PRJNA922040).

Results

A chromosome-level genome assembly of *S. macrospora*

The nuclear genome of *S. macrospora* was previously shown by pulsed-field gel electrophoresis to

consist of seven chromosomes (21). Prior to our study, three genome assemblies of *S. macrospora*

have been generated with increasing continuity (33, 35, 55); however, even the latest genome

assembly still consists of 584 scaffolds (33) and would therefore be unsuited for analyses at

chromosome level. Therefore, to aid in the planned Hi-C and ChIP-seq analyses, we generated a

chromosome-level genome assembly of *S. macrospora* based on nanopore sequence reads. The new

assembly (*S. macrospora* genome version 04) consists of seven gapless contigs with a total size of 39.4

Mb (Figure S1) which is similar in size to previous assemblies and to estimates by pulsed-field gel

electrophoresis of 39.5 Mb (21, 33, 35, 55). Six of the seven contigs have telomeric repeats at both

ends, whereas one contig has telomeric repeats at one end and 10 copies of the ribosomal RNA (rRNA)

genes at the other end with the repetitive nature of the rRNA genes most likely preventing assembly

of telomeric repeats at that end. This indicates that the seven contigs represent the seven

chromosomes of *S. macrospora*. A comparison with the genome assembly of the closely related

ascomycete *Neurospora crassa* showed a high degree of synteny for the seven chromosomes of each

species, consistent with previous analyses of smaller genomic regions of the two fungi (35, 56) (Figure

S1). Thus, a chromosome-level assembly for *S. macrospora* is now available and can be used for

analyses at chromosome level.

***S. macrospora* ASF1 depends on V94 and large parts of the C-terminus to accomplish histone binding**

The function and structure of ASF1 is highly conserved in all eukaryotes and it has been described as a
central histone chaperone that binds the histones H3 and H4 as a dimer and individually. The first 155
amino acids form a globular core in which acidic patches allow for the interaction with both histones
(7). Especially the area surrounding position V94 has been shown to be essential for ASF1-histone
binding in yeast (19). However, little is known about the functions of the C-terminal tail. This region is
much more divergent than the globular core and parts of it exist only in fungi (Figure 1). To analyze if
V94 is required for histone binding of *S. macrospora* ASF1, we performed co-immunoprecipitation
experiments with an ASF1 variant that includes a V94R substitution and a variant with a D37A
substitution. The latter is a mutation of an interaction site with the histone chaperone HIRA that was
identified in *S. pombe* (57). In addition, ASF1 variants with truncations after positions 152, 185 and 210
were generated to gain insight into possible involvement of the C-terminal region in the process of
histone binding. In GFP- and FLAG-trap experiments, the V94R variant proved to be incapable of pulling
down the histones H3 and H4, while the D37A variant performed like the wild type ASF1 (Figure 2).
Thus, the conserved V94 is required for histone interactions in *S. macrospora*, whereas the putative
HIRA interaction site D37 is not. Truncated ASF1 variants were also unable to bind H3 and H4 when
the truncations occurred after positions 152 or 185. Truncation after position 210 had no detectable
effect and did not impair the capability to pull down the histones (Figure 3).

The histone binding function of ASF1 is essential for sexual development
Deletions of *asf1* are usually lethal in higher eukaryotes, making research into its involvement in
multicellular development very challenging. *S. macrospora* and *A. thaliana* are the only known
exceptions that have been discovered so far (17, 18). The qualities of *S. macrospora* as an easy to
handle and fast propagating model organism with the ability to form complex, multicellular structures
during sexual development make it an exceptional tool for *asf1 in vivo* research. We used the *asf1*
variants described above carrying substitutions in the histone binding site V94, the putative HIRA
interaction site D37 and variants with truncated C-terminal regions for complementation analysis in a
previously constructed *asf1* deletion strain (6). By correlating the ability of these variants for rescuing
the phenotype of the deletion mutant with information about their ability to bind histones, more
insights into the role of ASF1-histone interaction during multicellular development can be gained. The
asf1 deletion mutant was transformed with plasmids coding for the respective variants, and
homokaryotic strains were obtained by the isolation of ascospores from self-fertile perithecia or from
genetic crosses. PCR screenings confirmed the integration of *asf1* variants in the resulting
transformants (Figure S4). The phenotype of the resulting strains was documented after growth for a
week on BMM medium. *S. macrospora* Δasf1 is sterile and mostly consists only of vegetative mycelium,
but is sometimes able to generate small protoperithecia. Introducing the wild type *asf1* gene back into

the deletion mutant led to a complete reversal of the deletion phenotype and yielded strains indistinguishable from the wild type, gaining full fertility. Strains expressing the D37A variant also regained fertility and were able to form perithecia, but showed slight aberrations. A tendency to grow giant protoperithecia and a distinct brown color of the mycelium and ascospores was observed. A reintroduction of an ASF1 variant with a V94R substitution did not complement the deletion mutant and the resulting strains exhibited the same phenotype as Δasf1 strains (Figure 4, Table 1). Since ASF1 V94R does not complement the sterile phenotype of deletion mutants and is unable to bind histones, a connection between histone binding and sexual development seems likely. ASF1 D37A proved to have no disruption in histone binding and regained fertility, but the substitution in a putative HIRA-binding site still had impact on the life cycle of *S. macrospora*. However, histone binding capability was enough to reach fertility. Truncations of ASF1 at positions 152 and 185 led to strains that resembled the deletion mutant, but a truncation at position 210 still allowed for full complementation (Figure 4, Table 1). Thus, although the C-terminal tail of ASF1 is less conserved, parts of it are essential for histone binding and sexual development. Leaving only the core intact, or shortening the protein to 185 amino acids seems to have the same effect on both processes as a full deletion. A truncation after position 210 removes the part of ASF1 that is only present in fungi and does not affect its overall functionality. To rule out the possibility of mislocalization as a cause for the inability to complement the deletion mutant, the eGFP-tagged variants were tracked by fluorescence microscopy. All of them co-localized with an mRFP-tagged H3 in the nucleus (Figure S5), similar to what was previously shown for wild type ASF1 (6).

ASF1 can contribute to DNA damage protection even without histone binding capabilities

Even after confirming the correct localization of tested ASF1 variants, a possible complete loss of function of the protein, i.e. not specifically related to histone binding, could not be completely ruled out. Thus, we looked for a way to confirm functionality of the variants aside from histone binding *in vivo*. The complex network of ASF1 regarding chromatin modification is still poorly understood and direct functions beyond nucleosome assembly are not widely known. It was previously show in *S. cerevisiae* that ASF1 is involved in DNA damage protection, since Δasf1 mutants are unable to grow on the DNA damage-inducing agent MMS (58). However, in *S. cerevisiae* C-terminal truncations had no impact on DNA damage protection abilities of ASF1 (59, 60). Therefore, we tested *S. macrospora* Δasf1 and the strains expressing the variants for their reaction to the DNA damaging agent MMS, which causes DNA single- and double strand breaks. At a concentration of 0.007 % MMS (v/v), the wild type was able to grow normally on BMM medium, while the mutant did not grow at all (Figure 5). Complementation strains expressing ASF1 variants D37A and 210T showed wild type-like resistance. Although the V94R, 152T and 185T variants of ASF1 are indistinguishable from the deletion strain

under normal conditions, an increased resistance against MMS compared to the Δasf1 strain was clearly visible (Figure 5). The difference between *S. macrospora* strains expressing these ASF1 variants and the deletion mutant suggests that an uncharacterized function of ASF1 in DNA damage protection exists and these variants are functional in this regard even though they are not able to bind histones. Thus, the data suggest that the role of ASF1 in DNA damage protection does not depend on histone interaction.

Deletion of *asf1* changes global histone modification levels in *S. macrospora*

The functions of ASF1 as a central histone chaperone and its multiple interaction partners during chromatin modification suggest a participation in processes like histone modification. Indeed, ASF1 has been shown to be involved in the acetylation of H3 on the lysines 9 and 56 in *S. cerevisiae* (9) and changes in the disposition of H3K27me3 and H3K9me3 have been observed in *M. musculus* cells during ASF1 depletion conditions (10). To gain insight into the role of ASF1 in histone modification, we performed semi-quantitative western blot screens with protein extracts from *S. macrospora* wild type and Δasf1 for histone modifications H3K27me3, H3K9me, H3K56ac, and H3K9ac. The comparison of the band strength after incubation with antibodies specific for these modifications indicate significant changes in global histone modification levels for H3K27me3 and H3K56ac in the Δasf1 mutant (Figure 6 A-B). H3K27me3 levels were upregulated in the deletion mutant, while a depletion of H3K56Ac could be observed. Thus, we chose these histone modifications as candidates for ChIP-Seq experiments to verify the results of the screening and to gain information about the local distribution of both histone marks. Establishing this method for older mycelium of *S. macrospora* presented challenges regarding the crosslinking and shearing process, as crosslinking agents appeared to create an agglomeration of inaccessible cells. A combination of high input amounts, soft lysis conditions and DNase based shearing, allowed us to conduct reproducible ChIP-seq experiments on older *S. macrospora* samples for the first time (Supplemental Text 1). Matching the results of the Western blot analysis, ChIP-seq revealed that the amount of H3K27me3 was significantly increased on a global level in the deletion mutant (Figure 6 C). Local distribution regarding functional regions like transcription start sides and gene bodies did not undergo significant changes, albeit looking at the peaks which only exist in wild type or Δasf1, a slight shift from intergenic regions into exon regions occurred in the *asf1* deletion mutant (Figure 6 E). H3K56ac levels appeared to be reduced in the deletion mutant, although this euchromatin mark was still present. Focusing on the unique H3K56ac positions in either Δasf1 or the wild type revealed that almost no nucleosomes were uniquely marked in *S. macrospora* Δasf1 whereas the wild type exhibits a far higher abundance of specific H3K56ac positions (Figure 6 D). The specific H3K56ac positions in the mutant seemed to be more prevalent in exons, compared to the wild type (Figure 6 F).

Hi-C reveals a 600 kb duplication in *asf1* deletion mutants that might be linked to their survival

To study the effect of *asf1* and sexual development on overall chromatin structure, we performed Hi-C experiments to map the abundance of DNA-DNA interactions in mycelium at two different developmental stages of the wild type and in mycelium of Δasf1. Samples for the young wild type were grown for 2 d, thus most of the mycelium consisted of vegetative tissue and sexual development was in its early stages. Samples for older mycelium were grown for 5 d, at this stage fully melanized protoperithecia and some early perithecia can be expected. The deletion mutant was cultivated for 3 d (since it is growing somewhat slower than the wild type), but is blocked in its sexual development at the stage of very young protoperithecia. The chromatin of the samples was fixed with formaldehyde and treated with DNase to obtain crosslinked pairs of DNA that are in close proximity at that time. Ends were filled with biotin and re-ligation was performed under high dilution conditions, so ligation events of crosslinked strands are more likely. A streptavidin pulldown was used to generate samples containing interacting DNA fragments of a sample at the given time. By using Dovetail® Dual index primers, Illumina® libraries of around 50 ng were obtained and used for high throughput paired end sequencing. 50 million paired end reads per sample were used for the generation of Hi-C maps. All recorded interaction events were plotted against their position on the *S. macrospora* genome, each one represented by a red dot. The intensity of red dots at a specific area indicates more recorded DNA-DNA interactions in the sample at the tested point in time. The characteristic diagonal line is an indicator of the high interaction frequency of genomic regions that are always in close proximity to each other, because of their vicinity on the DNA strand itself (Figure 7). While the resolution of the generated HI-C maps was too low to make out specific compartments of the young and old wild type, a significant reduction of the general resolution in the sexual mycelium might hint at the development of multiple cell types with different chromatin conformations. During sexual development *S. macrospora* generates multiple different cell types (4), therefore it seems likely that different cell types are characterized by different chromatin conformations. The presence of more than one chromatin conformation in a sample could lead to a loss of signal for mappable DNA-DNA interactions at a specific point. The Hi-C map of the deletion mutant appeared to show a strong increase of DNA-DNA interactions of a specific region of 600 kb on chromosome 2 with all other genomic regions (enhanced horizontal and vertical bars in Figure 7). However, such an explanation is biologically unlikely, and a more likely explanation would be that this genomic region in the Δasf1 mutant differs from the wild type genome that was used as a reference for mapping. In this case, this strong increase hinted at a duplication of this region in Δasf1, which would result in twice the number of reads, which then all mapped to the single copy of the wild type genome that was used as a reference for the generation of the Hi-C map. This would also lead to twice the number of (spurious) interactions with other regions

of the genome, which would appear as the observed enhanced horizontal and vertical bars in the Hi-C

contact map. We detected read patterns in the Hi-C data that match the internal border of a putative

tandem duplication of the area of interest. To test this, we sequenced the genome of the Δasf1 mutant

by Illumina sequencing. Sequencing reads were mapped to the wild type reference genome, which

proved the existence of the duplication in the deletion mutant (Figure 8 A). The detected duplication

consists of two compartments, with the middle region of the area of interest not duplicated, hinting

at a loss event or two independent duplication events. PCR tests with primers amplifying the newly

created internal junctions of the duplicated area (Figure 8 B) revealed the existence of the duplication

in all used Δasf1 strains even those that resulted from backcrosses against the wild type. Since the *asf1*

gene is located on chromosome 6, whereas the duplication is present on chromosome 2, they should

segregate independently resulting in approximately 50 % of Δasf1 strains with and 50 % without the

duplication in a cross with a wild type strain. Therefore, we suspected that the viability of the *asf1*

deletion might depend on the presence of the duplication. Crossing experiments were performed to

analyze the correlation between viable Δasf1 strains and the duplication. The crosses with the spore

color mutant fus, which carries the wild type *asf1* allele, resulted 69 strains with an *asf1* wild type

allele, of which 21 contained the duplication, matching expectations for free recombination. The 37

Δasf1 mutant strains all possessed the duplication, suggesting that the viability of Δasf1 strains is

indeed linked to the occurrence of the duplication on chromosome 2 (Table 2). During screenings for

duplication-free Δasf1 strains, we also screened strains carrying the *asf1* variants we used in the

complementation analysis and detected the absence of the duplication in an ASF1-V94R strain

(Supplemental Table 5). This variant is unable to bind histones and does not complement the

developmental phenotype of *asf1* deletions, but is able to contribute to the resistance against

genotoxic stress. This suggests that the presence of an ASF1 protein even without histone binding

ability is sufficient to overcome the potential lethality of the complete absence of ASF1, whereas the

complementation studies described above show that ASF1 without histone binding ability is not

sufficient for fruiting body development.

Discussion

***S. macrospora* ASF1 needs an intact C-terminal region to perform histone binding, an essential**

function for sexual development

Next to *A. thaliana* (17), *S. macropspora* is one of only two known multicellular organisms that survive

a full deletion of *asf1* (even though this might be contingent on the presence of the duplication on

chromosome 2 as described above), making it a valuable model for ASF1 *in vivo* research, especially

when its fast vegetative growth and sexual cycle are taken in account. Therefore, we have the unique opportunity to study the importance of *asf1* for multicellular development. *asf1* deletion strains have been shown to exhibit a slowed vegetative growth rate, and sexual development is blocked at the stage of young protoperithecia (6).

The ability to interact with histones can be regarded as the central requirement of a histone chaperone to perform its functions. ASF1 is known to interact with the histones H3 and H4 and deposits them during nucleosome assembly during transcription, replication and DNA-repair. The general interaction mode of ASF1 with H3 and H4 has been analyzed by *in vitro* experiments with *S. cerevisiae* proteins. The central acidic patch in the core of ASF1 and especially the amino acid V94 are essential in this regard (7). Since the first 155 amino acids of ASF1 are highly conserved in all eukaryotes, similar functions can be assumed. In fact, our co-immunoprecipitation experiments showed a clear dependency of *S. macrospora* ASF1 on V94 to establish interactions with H3 and H4. However, the C-terminal region of ASF1 is quite divergent, with some parts existing only in fungi, and information about the functional relevance of this region is sparse. Although experiments in *S. cerevisiae* showed a contribution of the C-terminal region of ASF1 to histone interaction and gene silencing, the specific interaction mode still remains elusive (59). Since the unicellular nature of yeasts does not allow for the analysis of multicellular developmental processes and the specific region of ASF1 is quite divergent, experiments in multicellular eukaryotes like *S. macrospora* might shed more light on the functions of this part of ASF1. Co-immunoprecipitation showed that despite of this region's divergent nature it is essential for establishing a connection between ASF1 and H3-H4 up until amino acid 210. Thus, a function of the C-terminal tail during this interaction is highly likely. The acidic nature of the C-terminal tail might contribute to facilitating the connection between ASF1 and the histones. A physical interaction between the C-terminal region and H3-H4 has not yet been detected, but might be difficult to prove with structural analysis, because of the usually unfolded C-terminus (58). ASF1 is known to bind histones H3 and H4, but prior to this study, it was not clear if histone binding is required for sexual development. We therefore analyzed different ASF1 variants for their ability to bind histones as well as their ability to support fruiting body formation. Since all ASF1 variants that did not co-immunoprecipitate the histones H3 and H4 were unable to restore the deletion phenotype, the histone binding function of ASF1 appears to be fundamental for sexual and therefore complex multicellular development. Complementation experiments with ASF1 variants that were shown to be able to bind histones led to fully fertile strains, underscoring the importance of this function for normal development. With the exception of ASF1 D37A, these variants generated *S. macrospora* strains that resembled the wild type. D37A strains exhibited a light brown coloration of the mycelium and had a tendency to generate giant protoperithecia. Since D37 is the putative interaction site of ASF1 with the

545 histone chaperone complex HIRA (19), one might speculate about an involvement of HIRA during
546 melanin biosynthesis and sexual development.
547
Histone binding is not essential for the function of ASF1 during DNA-damage repair
A role of ASF1 during DNA-damage repair without usage of its histone chaperone activity has often
been suspected and indeed, in recent studies with human ASF1A, a connection to double strand repair
mechanisms independent of histone binding has been discovered (61). In *S. macrospora*, strains
expressing ASF1 variants defective in histone binding were able to survive genotoxic stress, whereas
Δasf1 proved to be highly sensitive. It can therefore be concluded that the constructed ASF1 variants
are all functional in DNA damage repair and that the observed defects in fruiting body development
are specifically related to the ability to perform canonical histone chaperone functions. One hypothesis
might be that ASF1 is not only a histone chaperone, but also functions as a chromatin modifier hub
that regulates and supports the activity of other chromatin-modifying proteins. Regarding the
involvement in DNA damage repair, multiple interaction partner candidates are conceivable. A histone-
independent interaction with RIF1 has already been demonstrated in human cell lines and contributes
to determining the choice of the appropriate double strand break repair mechanism (61). Another
candidate involved in DNA damage protection and possibly activated by ASF1 is the histone
acetyltransferase RTT109. RTT109 is responsible for H3K9 and K56 acetylation, modifications that are
known to be important for DNA damage resistance (62). Interaction between ASF1 and RTT109 has
been characterized for *S. cerevisiae* (9) and an activating effect on the histone acetyltransferase was
shown, although the exact mechanism remains elusive (63). A defect in acetylation of H3 in *asf1*
deletion mutants has been also demonstrated in our ChIP-Seq experiments, therefore a connection
between the increased sensitivity of *S. macrospora* Δasf1 to DNA damage and lowered activity of
RTT109 seems possible. Another ASF1-mediated factor for DNA stability is known in *S. cerevisiae* as
the checkpoint kinase RAD53 and is involved in DNA-double strand break repair mechanisms. RAD53
needs to be inactivated and bound to ASF1 after repair has been completed, and lack of ASF1 could
lead to an overabundance of active RAD53 and subsequent disturbances in the cell cycle and further
damage repairs (64). All in all, ASF1 seems to play a crucial role during multiple DNA-damage protection
and repair pathways, not all of them relying on ASF1-H3/H4 interactions.
574
ASF1 is involved in the histone modification landscape of *S. macrospora*
The histones are targets of various posttranslational modifications. The best known are acetylation,
methylation, phosphorylation, sumoylation, and ubiquitination. These modifications vary throughout
the genome and shape the histone code, which in turn influences access to the DNA for transcription
and the recruiting of transcription factors. The histone landscape of a cell is a significant determinant

of the transcriptome (65). ASF1 as a central eukaryotic histone chaperone has been reported to affect

multiple histone modifications, such as acetylation of H3K56 (9) and methylation of H3K27 (10). These

modifications undergo significant changes after deletion of *asf1* in *S. macrospora*. For H3K27me3, a

well-known heterochromatin modification (66), *asf1* deletion mutants showed almost three times as

many unique H3K27 trimethylations as the wild type. The increase in unique H3K27me3 positions did

not seem to be very site specific, as the relative distribution was comparable in the mutant and the

wild type. In general, we did not observe a significant change of H3K27me3 in specific functional

regions, like transcriptional start sites or gene bodies. The increase of H3K27me3 in *asf1* deletion

strains could either be a direct consequence of lack of activation of a corresponding methyltransferase,

or a compensatory reaction to the absence of ASF1. ASF1 is known to be involved in the recycling of

histones during DNA replication (7). H3K27me3 is an inheritable histone modification that can persist

during replication and the *de novo* integration of histones by ASF1 and CAF1 in parental cells was

shown to heavily dilute the inheritance of this mark in nascent chromatin in *A. thaliana* (67). Therefore,

a reduction in *de novo* nucleosome assembly through the deletion of a major histone chaperone like

ASF1 could reduce general recycling of histones and cause the already established H3K27me3 to persist

and accumulate. Especially genomic regions that do not carry H3K27me3 all the time under wild type

conditions would show a visible increase under such circumstances, matching our observations of

significantly more uniquely marked positions in the deletion mutant. The blocking of nucleosome

recycling and the resulting more permanent marking of histones with H3K27me3 might be an

explanation for the severe developmental phenotype of *S. macrospora* Δasf1. Genome wide

redistribution of H3K27me3 has been linked to defects in sexual development in *N. crassa*, a close

relative of *S. macrospora* (68), so the persistence of this mark in regions where it would exist only

temporarily in the wild type might cause similar effects. However, the changes in H3K27me3

distribution could also be a reaction to the reduced DNA methylation observed in Δasf1 (18). In

addition to a global reduction of DNA methylation, the deletion of the *S. macrospora asf1* leads to an

increase in transcription of many usually weakly expressed genes (18). DNA methylation and

H3K27me3 are mostly considered to be marks of transcriptionally silenced regions, but are often

regarded as antagonistic to each other (69). Thus, the observed global increase in H3K27me3 in a

deletion mutant that also shows a decrease in DNA methylation could be the result of a compensation

mechanism for the loss of a transcriptional silencing mechanism. The rising transcription levels

resulting from the DNA methylation loss might be kept at least in an acceptable range by using other

transcriptional silencers, such as H3K27me3 to keep the deletion mutant alive. This effect might

therefore be one of the reasons why *S. macrospora* survives full *asf1* deletion and other multicellular

organisms do not.

Histone acetylations on the other hand are known as euchromatin markers, their negative charge can lower the overall positive charge of histones and weaken their binding to the DNA. This can facilitate a more relaxed and open chromatin state that can be accessed by RNA polymerases (70). The higher transcription rates of usually weakly expressed genes after the deletion of *asf1* in *S. macrospora* might suggest that euchromatin marks could be in higher abundance in such a strain. For H3K56ac, our ChIP-Seq results showed that at least for this mark, the opposite is the case and the mutant exhibits less H3K56 acetylation on a global scale. As with the heterochromatin mark H3K27me3, we were not able to detect specific patterns of acetylation loss, the overall abundance of the mark appeared to be reduced. This coincides with the higher sensitivity of Δasf1 strains to genotoxic stress. H3K56ac has been described to be of high importance for DNA damage repair (71). Strikingly, we still detected H3K56ac during western blot screenings, as well as ChIP-Seq experiments, while it has been reported that the presence of ASF1 is required to generate this mark in *S. cerevisiae* (72). A significant reduction of H3K56ac levels in *S. macrospora* Δasf1, however, might be an explanation for the severe sensitivity to genotoxic stress of those strains. H3K56ac is known as an important factor for genomic stability and resistance against replication fork damage, as studies in yeast have shown (73). Since this mark is generated by the histone acetyltransferase RTT109, which is activated by ASF1 (63), it can be speculated that the lower abundance of H3K56ac and the genomic instability are resulting from a lack of ASF1-RTT109 interaction. Additionally, the presence of H3K56ac in *S. macrospora* Δasf1, even though ASF1 was expected to be required for the generation of this mark, hints at either an unknown second activation pathway for RTT109, or an undiscovered way to establish H3K56ac in multicellular ascomycetes. Taken together with the detected increased levels of the heterochromatin mark H3K27me3, lower euchromatic H3K56ac levels would suggest an all-around lower general transcriptional activity of *S. macrospora asf1* deletion mutants. Since the opposite has been documented (18), it can be hypothesized that the contribution of ASF1 to regulating the transcriptome is either not mainly mediated by histone modifications, or is even more complex than previously assumed.

Hi-C analysis reveals a 600 kb duplication in *S. macrospora* Δasf1 which might be linked to its viability
Methods such as Hi-C allow the analysis of the three-dimensional organization of the chromatin and its effects on the transcriptome and gene regulation. The relationship between chromatin structure and gene regulation seems to be quite complex and possibly both processes can regulate each other (74). In this study, we created DNA-DNA interaction maps of the young, mostly vegetative state of *S. macrospora* and compared it to older mycelium, which has progressed in the cycle of sexual development and an *asf1* deletion mutant with the aim of identifying differences in the chromatin state of those samples. The major challenge was the application of a method that has been fine tuned

for single cell assays and fungal protoplasts to older, more mature fungal mycelium. The resulting map

of the young wild type showed some areas of the chromatin that suggested higher condensation than

other areas, but in general the resolution was not high enough to confidently identify specific

interaction compartments. An even lower resolution was visible for the Hi-C map of the older, sexual

sample. But the lower DNA-DNA interaction in this case could be a sign of multiple different chromatin

states. Entering sexual development, *S. macrospora* begins generating more specialized cell types, at

least 13 different cell types have been identified during fruiting body formation of this fungus (4).

Different functions of cells require different transcriptomic processes, which can depend on specific

changes of the three-dimensional chromatin state. Higher differentiation of cells in a sample and

therefore more divergent chromatin states would reduce the signal-to-noise ratio and would lead to

lower detection rates of DNA-DNA interactions. Significant changes in the transcriptome during

development are well documented in *S. macrospora* (3). Other studies have demonstrated a strong

connection between detectable changes of chromatin architecture and gene expression (75), so it can

be assumed that the transcriptional changes of *S. macrospora* during sexual development might be

accompanied by significant changes of DNA-DNA interaction frequency. A lower resolution of regularly

detectable interactions in the older wild type that is generating multiple cell types supports these

assumptions and might be another sign of chromatin state changes that accompany multicellular

development.

The Hi-C analysis of the Δasf1 mutant revealed a large duplicated region on chromosome 2. We

hypothesized that the presence of this duplication might not be just a coincidence, but could be linked

to the surprising viability of the Δasf1 mutant in *S. macrospora*. To the best of our knowledge, this

ascomycete is one of only two multicellular organisms that are known to survive loss of *asf1* (6, 17).

Genetic crosses confirmed co-segregation of the Δasf1 allele on chromosome 6 with the duplication

on chromsome 2, whereas the wild type *asf1* allele and the duplication were able to recombine freely.

These findings support the hypothesis that the viability of the Δasf1 mutant is coupled to the presence

of the duplication. Which specific role this region on chromosome 2 plays in the network of ASF1 is still

unclear. So far, no documented interaction partners or parts of the ASF1 network seem to be encoded

in this region. We assume that candidate genes in the duplication contribute to DNA damage repair or

protection, since during the crossing experiments we included *asf1* strains which express ASF1 variants

with specific amino acid substitutions, and one ASF1-V94R variant did not contain the duplication of

the area of interest. This variant is unable to perform the histone chaperone function of ASF1, due to

its inability to bind histones and is also unable to complement the development phenotype of *S.*

macrospora Δasf1. However, it is able to complement the sensitivity to genotoxic stress of Δasf1. Thus,

a strain that is able to perform the DNA-damage protection function of ASF1 is not dependent on the

presence of the duplication on chromosome 2, therefore a role of the duplicated region in DNA-
damage protection might be conceivable.

Summarizing our work, we discovered that the ability of ASF1 to interact with histones is essential for
sexual development of *S. macrospora*, a function that also depends on the presence of large part of
the less conserved C-terminus of ASF1. Histone binding however, proved not to be necessary to allow
ASF1 to contribute to the genotoxic stress response. ChIP-seq experiments showed that ASF1 is
involved in the establishment of H3K56ac and the repression of H3K27me3. Furthermore, we
discovered a 600 kb duplication in the deletion mutant that appears to be linked to the surprising
viability of *S. macrospora* Δasf1. Future work might be able to shed light on how ASF1 regulates gene
expression by histone modification and which genes of the duplicated area are necessary to stabilize
asf1 deletion mutants.

Acknowledgements

The authors would like to thank Silke Nimtz, Susanne Schlewinski and Tanja Rollnik for excellent
technical assistance and Christopher Grefen for support at the Department of Molecular and Cellular
Botany. The authors acknowledge the NGS team of the Omics CF NGS Unit (in development) and
CeBiTec. This work was funded by the German Research Foundation (DFG, grant NO407/7-2 to MN).

References

1.   **Niklas KJ, Newman SA.** 2020. The many roads to and from multicellularity. Journal of
Experimental Botany **71:**3247-3253.
2.   **Nagy LG, Kovács GM, Krizsán K.** 2018. Complex multicellularity in fungi: evolutionary
convergence, single origin, or both? Biol Rev **93:**1778-1794.
3.   **Teichert I, Pöggeler S, Nowrousian M.** 2020. *Sordaria macrospora*: 25 years as a model
organism for studying the molecular mechanisms of fruiting body development. Appl
Microbiol Biotechnol **104:**3691-3704.
4.   **Lord KM, Read ND.** 2011. Perithecium morphogenesis in *Sordaria macrospora*. Fungal Genet
Biol **48:**388-399.
5.   **Nowrousian M.** 2018. Genomics and transcriptomics to study fruiting body development: An
update. Fungal Biology Reviews **32:**231-235.
6.   **Gesing S, Schindler D, Fränzel B, Wolters D, Nowrousian M.** 2012. The histone chaperone
ASF1 is essential for sexual development in the filamentous fungus *Sordaria macrospora*. Mol
Microbiol **84:**748-765.
7.   **English CM, Adkins MW, Carson JJ, Churchill ME, Tyler JK.** 2006. Structural basis for the
histone chaperone activity of ASF1. Cell **127:**495-508.
8.   **Yamane K, Mizuguchi T, Cui B, Zofall M, Noma K, Grewal SI.** 2011. ASF1/HIRA facilitate global
histone deacetylation and associate with HP1 to promote nucleosome occupancy at
heterochromatic loci. Mol Cell **41:**56-66.
9.   **Lercher L, Danilenko N, Kirkpatrick J, Carlomagno T.** 2018. Structural characterization of the
ASF1-RTT109 interaction and its role in histone acetylation. Nucleic Acids Res **46:**2279-2289.

10. **Clément C, Orsi GA, Gatto A, Boyarchuk E, Forest A, Hajj B, Miné-Hattab J, Garnier M, Gurard-**
Levin ZA, Quivy J-P, Almouzni G. 2018. High-resolution visualization of H3 variants during
replication reveals their controlled recycling. Nature Communications **9:**3181.
11. **Prado F, Cortés-Ledesma F, Aguilera A.** 2004. The absence of the yeast chromatin assembly
factor ASF1 increases genomic instability and sister chromatid exchange. EMBO Rep **5:**497-
502.
12. **Groth A, Ray-Gallet D, Quivy J-P, Lukas J, Bartek J, Almouzni G.** 2005. Human ASF1 Regulates
the Flow of S Phase Histones during Replicational Stress. Molecular Cell **17:**301-311.
13. **Tagami H, Ray-Gallet D, Almouzni G, Nakatani Y.** 2004. Histone H3.1 and H3.3 complexes
mediate nucleosome assembly pathways dependent or independent of DNA synthesis. Cell
116:51-61.
14. **Le S, Davis C, Konopka JB, Sternglanz R.** 1997. Two new S-phase-specific genes from
Saccharomyces cerevisiae. Yeast **13:**1029-1042.
15. **Singer MS, Kahana A, Wolf AJ, Meisinger LL, Peterson SE, Goggin C, Mahowald M, Gottschling**
DE. 1998. Identification of high-copy disruptors of telomeric silencing in *Saccharomyces*
cerevisiae. Genetics **150:**613-632.
16. **Tyler JK, Adams CR, Chen SR, Kobayashi R, Kamakaka RT, Kadonaga JT.** 1999. The RCAF
complex mediates chromatin assembly during DNA replication and repair. Nature **402:**555-
560.
17. **Zhu Y, Weng M, Yang Y, Zhang C, Li Z, Shen WH, Dong A.** 2011. *Arabidopsis* homologues of
the histone chaperone ASF1 are crucial for chromatin replication and cell proliferation in plant
development. Plant J **66:**443-455.
18. **Schumacher DI, Lütkenhaus R, Altegoer F, Teichert I, Kück U, Nowrousian M.** 2018. The
transcription factor PRO44 and the histone chaperone ASF1 regulate distinct aspects of
multicellular development in the filamentous fungus *Sordaria macrospora*. BMC Genetics
19:112.
19. **Mousson F, Lautrette A, Thuret JY, Agez M, Courbeyrette R, Amigues B, Becker E, Neumann**
JM, Guerois R, Mann C, Ochsenbein F. 2005. Structural basis for the interaction of ASF1 with
histone H3 and its functional implications. Proc Natl Acad Sci U S A **102:**5975-5980.
20. **Esser K.** Cryptogams : cyanobacteria, algae, fungi, lichens : textbook and practical guide, p. *In*
(ed),
21. **Walz M, Kück U.** 1995. Transformation of *Sordaria macrospora* to hygromycin B resistance:
characterization of transformants by electrophoretic karyotyping and tetrad analysis. Curr
Genet **29:**88-95.
22. **Kück U, Pöggeler S, Nowrousian M, Nolting N, Teichert I.** 2009. *Sordaria macrospora*, a Model
System for Fungal Development, p 17-39 doi:10.1007/978-3-642-00286-1_2.
23. **Oldenburg KR, Vo KT, Michaelis S, Paddon C.** 1997. Recombination-mediated PCR-directed
plasmid construction *in vivo* in yeast. Nucleic Acids Res **25:**451-452.
24. **Terfrüchte M, Joehnk B, Fajardo-Somera R, Braus GH, Riquelme M, Schipper K, Feldbrügge**
M. 2014. Establishing a versatile Golden Gate cloning system for genetic engineering in fungi.
Fungal Genet Biol **62:**1-10.
25. **Nowrousian M, Masloff S, Pöggeler S, Kück U.** 1999. Cell differentiation during sexual
development of the fungus *Sordaria macrospora* requires ATP citrate lyase activity. Mol Cell
Biol **19:**450-460.
26. **Bayram O, Krappmann S, Ni M, Bok JW, Helmstaedt K, Valerius O, Braus-Stromeyer S, Kwon**
NJ, Keller NP, Yu JH, Braus GH. 2008. VelB/VeA/LaeA complex coordinates light signal with
fungal development and secondary metabolism. Science **320:**1504-1506.
27. **Laemmli UK.** 1970. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. Nature **227:**680-685.
28. **Colot HV, Park G, Turner GE, Ringelberg C, Crew CM, Litvinkova L, Weiss RL, Borkovich KA,**
Dunlap JC. 2006. A high-throughput gene knockout procedure for Neurospora reveals
functions for multiple transcription factors. Proc Nat Acad Sci USA **103:**10352-10357.

29.   **Engh I, Würtz C, Witzel-Schlömp K, Zhang Hai Y, Hoff B, Nowrousian M, Rottensteiner H, Kück**
778   **U.** 2007. The WW Domain Protein PRO40 Is Required for Fungal Fertility and Associates with
779   Woronin Bodies. Eukaryotic Cell **6:**831-843.

780   30.   **Traeger S, Altegoer F, Freitag M, Gabaldon T, Kempken F, Kumar A, Marcet-Houben M,**
781   **Pöggeler S, Stajich JE, Nowrousian M.** 2013. The genome and development-dependent
782   transcriptomes of *Pyronema confluens*: a window into fungal evolution. PLoS Genet
783   **9:**e1003820.

784   31.   **Vaser R, Sović I, Nagarajan N, Šikić M.** 2017. Fast and accurate de novo genome assembly from
785   long uncorrected reads. Genome Res **27:**737-746.

786   32.   **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,**
787   **Wortman J, Young SK, Earl AM.** 2014. Pilon: an integrated tool for comprehensive microbial
788   variant detection and genome assembly improvement. PLOS ONE **9:**e112963.

789   33.   **Blank-Landeshammer B, Teichert I, Märker R, Nowrousian M, Kück U, Sickmann A.** 2019.
790   Combination of proteogenomics with peptide *de novo* sequencing identifies new genes and
791   hidden posttranscriptional modifications. mBio **10:**e02367-02319.

792   34.   **Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ.** 1997. Gapped
793   BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids
794   Res **25:**3389-3402.

795   35.   **Nowrousian M, Stajich JE, Chu M, Engh I, Espagne E, Halliday K, Kamerewerd J, Kempken F,**
796   **Knab B, Kuo HC, Osiewacz HD, Pöggeler S, Read ND, Seiler S, Smith KM, Zickler D, Kück U,**
797   **Freitag M.** 2010. *De novo* assembly of a 40 Mb eukaryotic genome from short sequence reads:
798   *Sordaria macrospora*, a model organism for fungal morphogenesis. PLoS Genet **6:**e1000891.

799   36.   **Sović I, Šikić M, Wilm A, Fenlon SN, Chen S, Nagarajan N.** 2016. Fast and sensitive mapping of
800   nanopore sequencing reads with GraphMap. Nat Commun **7:**11307.

801   37.   **Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D, Warrenfeltz S,**
802   **Aurrecoechea C, Stoeckert CJJ, Kissinger J, Roos DS, Hertz-Fowler C.** 2018. FungiDB: An
803   integrated bioinformatic resource for fungi and oomycetes. J Fungi **4:**39.

804   38.   **Galagan JE, Calvo SE, Borkovich KA, Selker EU, Read ND, Jaffe D, FitzHugh W, Ma LJ, Smirnov**
805   **S, Purcell S, Rehman B, Elkins T, Engels R, Wang S, Nielsen CB, Butler J, Endrizzi M, Qui D,**
806   **Ianakiev P, Bell-Pedersen D, Nelson MA, Werner-Washburne M, Selitrennikoff CP, Kinsey JA,**
807   **Braun EL, Zelter A, Schulte U, Kothe GO, Jedd G, Mewes W, Staben C, Marcotte E, Greenberg**
808   **D, Roy A, Foley K, Naylor J, Stange-Thomann N, Barrett R, Gnerre S, Kamal M, Kamvysselis**
809   **M, Mauceli E, Bielke C, Rudd S, Frishman D, Krystofova S, Rasmussen C, Metzenberg RL,**
810   **Perkins DD, Kroken S, et al.** 2003. The genome sequence of the filamentous fungus
811   *Neurospora crassa*. Nature **422:**859-868.

812   39.   **Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL.** 2004.
813   Versatile and open software for comparing large genomes. Genome Biol **5:**R12.

814   40.   **Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Sánchez Alvarado A, Yandell**
815   **M.** 2008. MAKER: An easy-to-use annotation pipeline designed for emerging model organism
816   genomes. Genome Res **18:**188-196.

817   41.   **Lowe TM, Eddy SR.** 1997. tRNAscan-SE: a program for improved detection of transfer RNA
818   genes in genomic sequence. Nucl Acids Res **25:**955-964.

819   42.   **Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA.** 2012. Artemis: an integrated
820   platform for visualization and analysis of high-throughput sequence-based experimental data.
821   Bioinf **28:**464-469.

822   43.   **Teichert I, Wolff G, Kück U, Nowrousian M.** 2012. Combining laser microdissection and RNA-
823   seq to chart the transcriptional landscape of fungal development. BMC Genomics **13:**511.

824   44.   **Kim D, Paggi JM, Park C, Bennett C, Salzberg SL.** 2019. Graph-based genome alignment and
825   genotyping with HISAT2 and HISAT-genotype. Nat Biotech **37:**907-915.

826   45.   **Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram
827   quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry
828   **72:**248-254.

46. **Huang X, Pan Q, Lin Y, Gu T, Li Y.** 2020. A native chromatin immunoprecipitation (ChIP)
protocol for studying histone modifications in strawberry fruits. Plant Methods **16:**10.
47. **Chen S, Zhou Y, Chen Y, Gu J.** 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34:**i884-i890.
48. **Li H, Durbin R.** 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform.
Bioinformatics **26:**589-595.
49. **Schulten A, Pietzenuk B, Quintana J, Scholle M, Feil R, Krause M, Romera-Branchat M, Wahl**
V, Severing E, Coupland G, Krämer U. 2022. Energy status-promoted growth and development
of Arabidopsis require copper deficiency response transcriptional regulator SPL7. Plant Cell
34:3873-3898.
50. **Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,**
Brown M, Li W, Liu XS. 2008. Model-based Analysis of ChIP-Seq (MACS). Genome Biology
9:R137.
51. **Quinlan AR, Hall IM.** 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics **26:**841-842.
52. **Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK.**
2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory
Elements Required for Macrophage and B Cell Identities. Molecular Cell **38:**576-589.
53. **Open2C, Nezar A, Geoffrey F, Ilya MF, Aleksandra AG, Anton G, Maxim I, Sergey VV.** 2023.
Pairtools: from   sequencing   data   to   chromosome   contacts.   bioRxiv
doi:10.1101/2023.02.13.528389:2023.2002.2013.528389.
54. **Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL.** 2016. Juicer
Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst **3:**95-
98.
55. **Nowrousian M, Teichert I, Masloff S, Kück U.** 2012. Whole-genome sequencing of *Sordaria*
macrospora mutants identifies developmental genes. G3 (Bethesda) **2:**261-270.
56. **Nowrousian M, Würtz C, Pöggeler S, Kück U.** 2004. Comparative sequence analysis of *Sordaria*
macrospora and *Neurospora crassa* as a means to improve genome annotation. Fungal Genet
Biol **41:**285-292.
57. **Malay AD, Umehara T, Matsubara-Malay K, Padmanabhan B, Yokoyama S.** 2008. Crystal
structures of fission yeast histone chaperone Asf1 complexed with the Hip1 B-domain or the
Cac2 C terminus. Journal of Biological Chemistry **283:**14022-14031.
58. **Daganzo SM, Erzberger JP, Lam WM, Skordalakes E, Zhang R, Franco AA, Brill SJ, Adams PD,**
Berger JM, Kaufman PD. 2003. Structure and function of the conserved core of histone
deposition protein ASF1. Curr Biol **13:**2148-2158.
59. **Dennehey BK, Noone S, Liu WH, Smith L, Churchill ME, Tyler JK.** 2013. The C terminus of the
histone chaperone ASF1 cross-links to histone H3 in yeast and promotes interaction with
histones H3 and H4. Mol Cell Biol **33:**605-621.
60. **Tamburini BA, Carson JJ, Linger JG, Tyler JK.** 2006. Dominant mutants of the *Saccharomyces*
cerevisiae ASF1 histone chaperone bypass the need for CAF-1 in transcriptional silencing by
altering histone and Sir protein recruitment. Genetics **173:**599-610.
61. **Tang M, Chen Z, Wang C, Feng X, Lee N, Huang M, Zhang H, Li S, Xiong Y, Chen J.** 2022. Histone
chaperone ASF1 acts with RIF1 to promote DNA end joining in BRCA1-deficient cells. Journal
of Biological Chemistry **298:**101979.
62. **Mao P, Wyrick JJ.** 2016. Emerging roles for histone modifications in DNA excision repair. FEMS
Yeast Research **16:**fow090.
63. **Tsubota T, Berndsen CE, Erkmann JA, Smith CL, Yang L, Freitas MA, Denu JM, Kaufman PD.**
2007. Histone H3-K56 acetylation is catalyzed by histone chaperone-dependent complexes.
Mol Cell **25:**703-712.
64. **Tsabar M, Waterman DP, Aguilar F, Katsnelson L, Eapen VV, Memisoglu G, Haber JE.** 2016.
ASF1 facilitates dephosphorylation of RAD53 after DNA double-strand break repair. Genes Dev
30:1211-1224.

65.   **Bannister AJ, Kouzarides T.** 2011. Regulation of chromatin by histone modifications. Cell Res
882   **21:**381-395.
883   66.   **Boros J, Arnoult N, Stroobant V, Collet JF, Decottignies A.** 2014. Polycomb repressive complex
884   2 and H3K27me3 cooperate with H3K9 methylation to maintain heterochromatin protein 1α
885   at chromatin. Mol Cell Biol **34:**3662-3674.
886   67.   **Hugues A, Jacobs CS, Roudier F.** 2020. Mitotic Inheritance of PRC2-Mediated Silencing:
887   Mechanistic Insights and Developmental Perspectives. Frontiers in Plant Science **11**.
888   68.   **Basenko EY, Sasaki T, Ji L, Prybol CJ, Burckhardt RM, Schmitz RJ, Lewis ZA.** 2015. Genome-
889   wide redistribution of H3K27me3 is linked to genotoxic stress and defective growth. Proc Natl
890   Acad Sci U S A **112:**E6339-6348.
891   69.   **Hagarman JA, Motley MP, Kristjansdottir K, Soloway PD.** 2013. Coordinate regulation of DNA
892   methylation and H3K27me3 in mouse embryonic stem cells. PLoS One **8:**e53880.
893   70.   **Lee CY, Grant PA.** 2019. Chapter 1-1 - Role of Histone Acetylation and Acetyltransferases in
894   Gene Regulation, p 3-30. *In* McCullough SD, Dolinoy DC (ed), Toxicoepigenetics
895   doi:https://doi.org/10.1016/B978-0-12-812433-8.00001-0. Academic Press.
896   71.   **Driscoll R, Hudson A, Jackson SP.** 2007. Yeast RTT109 promotes genome stability by
897   acetylating histone H3 on lysine 56. Science **315:**649-652.
898   72.   **Adkins MW, Carson JJ, English CM, Ramey CJ, Tyler JK.** 2007. The Histone Chaperone Anti-
899   silencing Function 1 Stimulates the Acetylation of Newly Synthesized Histone H3 in S-phase*.
900   Journal of Biological Chemistry **282:**1334-1340.
901   73.   **Wurtele H, Kaiser GS, Bacal J, St-Hilaire E, Lee EH, Tsao S, Dorn J, Maddox P, Lisby M, Pasero
902   P, Verreault A.** 2012. Histone H3 lysine 56 acetylation and the response to DNA replication
903   fork damage. Mol Cell Biol **32:**154-172.
904   74.   **Banigan EJ, Tang W, van den Berg AA, Stocsits RR, Wutz G, Brandão HB, Busslinger GA, Peters
905   J-M, Mirny LA.** 2023. Transcription shapes 3D chromatin organization by interacting with loop
906   extrusion. Proceedings of the National Academy of Sciences **120:**e2210480120.
907   75.   **Almassalha LM, Tiwari A, Ruhoff PT, Stypula-Cyrus Y, Cherkezyan L, Matsuda H, Dela Cruz
908   MA, Chandler JE, White C, Maneval C, Subramanian H, Szleifer I, Roy HK, Backman V.** 2017.
909   The Global Relationship between Chromatin Physical Topology, Fractal Structure, and Gene
910   Expression. Sci Rep **7:**41061.

920 **Tables**

924 **Table 1.** Transformants and fertility rates for complementation experiments of Δasf1 with ASF1

925 variants.

Encoded ASF1 variant	No. of primary transformants	Fertile transformants	Sterile transformants	Percantage of fertile transformants
Wild type	53	47	6	88.7
D37A	61	41	20	67.2
V94R	41	0	41	0
152T	19	0	19	0
185T	22	0	22	0
210T	18	15	3	83.3

929 **Table 2.** Presence of the duplication on chromosome 2 in wild type and *asf1* deletion strains. Strains

930 were obtained from crosses of Δasf1 with the spore color mutant fus, which carries the wild type *asf1*

931 gene. The full list of tested strains has been added as Supplemental table 5.

Genotype	No. of tested strains	No. of strains with duplication	Percentage of strains with duplication
Wild type	69	21	30.4
Δasf1	37	37	100

935   **Figures**

Figure 1. Alignment of ASF1 amino acid sequences of several animal and fungal model organisms. The
first 152 amino acids are highly conserved. The C-terminal region is more divergent and parts of it exist
only in fungi. Positions D37 and V94 were substituted for functional analysis in this study. D37 is known
as a HIRA interaction site in *Schizosaccharomyces pombe* (57), while V94 has been established as the
H3-H4 interaction site (7, 19). Positions 152, 185 and 210 were targeted to create truncated ASF1
variants. Abbreviations and accession numbers: *Mm*, *Mus musculus* (AAH27628.1); *Dm*, *Drosophila*
melanogaster (NP_524163.1); *Sm*, *Sordaria macrospora* (KAA8632619.1); *Nc*, *Neurospora crassa*
(XP_958588.1); *Sc*, *Saccharomyces cerevisiae* (NP_012420.1).

GFP-Trap

Figure 2. Co-immunoprecipitation results for ASF1 variants with amino acid substitutions. GFP-tagged ASF1 wild type and variants D37A and V94R were used as bait proteins for Flag-tagged histones H3 and H4 in a GFP trap. Results were checked by western blot analysis with antibodies against GFP and Flag tags. ASF1 wild type and the D37A variant showed signals for bait and prey proteins in the raw and eluate sample, indicating interaction, whereas the V94R variant showed the signal for the prey protein only in the raw sample. Strains expressing non-fused eGFP with the corresponding Flag-tagged histones were used as a negative control. The corresponding results with histones H3 and H4 as baits are shown in Supplemental Figure 2.

GFP-Trap

Detection of eGFP-Tags

ASF1-wild type-eGFP + H3-Flag		ASF1-152T-eGFP + H3-Flag		ASF1-185T-eGFP + H3-Flag		ASF1-210T-eGFP + H3-Flag		eGFP + H3-Flag	
Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate
57 kDa		45 kDa		48 kDa		51 kDa		27 kDa	

Detection of Flag-Tags

ASF1-wild type-eGFP + H3-Flag		ASF1-152T-eGFP + H3-Flag		ASF1-185T-eGFP + H3-Flag		ASF1-210T-eGFP + H3-Flag		eGFP + H3-Flag	
Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate
18 kDa		18 kDa		18 kDa		18 kDa		18 kDa	

Detection of eGFP-Tags

ASF1-wild type-eGFP + H4-Flag		ASF1-152T-eGFP + H4-Flag		ASF1-185T-eGFP + H4-Flag		ASF1-210T-eGFP + H4-Flag		eGFP + H4-Flag	
Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate
57 kDa		45 kDa		48 kDa		51 kDa		27 kDa	

Detection of Flag-Tags

ASF1-wild type-eGFP + H4-Flag		ASF1-152T-eGFP + H4-Flag		ASF1-185T-eGFP + H4-Flag		ASF1-210T-eGFP + H4-Flag		eGFP + H4-Flag	
Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate	Raw	Eluate
14 kDa		14 kDa		14 kDa		14 kDa		14 kDa	

Figure 3. Co-immunoprecipitation results for truncated ASF1 variants. GFP-tagged ASF1 wild type and variants 152T, 185T and 210T were used as bait proteins for Flag-tagged histones H3 and H4 in a GFP trap. Results were checked by western blot analysis with antibodies against GFP and Flag tags. ASF1 wild type and the 210T variant showed signals for bait and prey proteins in the raw and eluate sample, indicating interaction, whereas the 152T and 185T variants showed the signal for the prey protein only in the raw sample. Strains expressing non-fused eGFP with the corresponding Flag-tagged histones were used as a negative control. The corresponding results with histones H3 and H4 as baits are shown in Supplemental Figure 3.

Figure 4. Phenotypes of strains from Δasf1 complementation experiments. The wild type generates mature perithecia in 7 days, while the development of the deletion mutant is blocked at the stage of early protoperithecia and mostly generates only vegetative mycelium. Reintroducing the *asf1* wild type gene into the deletion mutant fully complements the phenotype and lead to strains that are indistinguishable from the wild type. Transforming Δasf1 with the *asf1* V94R variant did not complement the mutant strain resulting in strains with the sterile phenotype of Δasf1. A D37A variant partially complements the mutant, such strains are fertile, but show distinct developmental aberrations like a tendency to generate mostly large protoperithecia and a brown coloration of the mycelium. Transformation with truncated *asf1* variants did not lead to complementation with variants 152T and 185T, but the longer 210T variant led to strains that show a wild type phenotype again.

987
988

Figure 5. Genotoxic stress resistance of *S. macrospora* strains expressing ASF1 variants. At the given MMS concentration, the wild type was able to grow normally, while the *asf1* deletion mutant exhibited severe sensitivity (no growth). Reintroducing the wild type *asf1* into the mutant restored the resistance. All strains that expressed any kind of ASF1 variant showed at least partial resistance to genotoxic stress, even variants that were shown to be unable to interact with histones and do not complement the developmental defects of *S. macrospora* Δasf1.

Figure 6. Analysis of histone modifications in the wild type and Δasf1.

A. Semiquantitative screening for histone modifications affected by ASF1. Western blots with antibodies against the indicated histone modifications were used to compare the band strength of equal amounts of protein from wild type and Δasf1 extracts.

B. The band intensity in Western Blots was measured densiometrically in 5 independent biological replicates and the ratio of Δasf1 versus wild type was calculated. The deletion mutant showed a significant increase in H3K27me3 and significant decreases in H3K9me and H3K56ac, while H3K9ac was not affected by an *asf1* deletion. Student's t-test for significance: * = p-value < 0.05; ** = p-value < 0.01.

C and D. Uniquely H3K27 trimethylated positions (C) and uniquely H3K56 acetylated positions (D) across the *S. macrospora* genome in wild type and Δasf1. ChiP-Seq revealed a general increase in H3K27me3 and decrease in H3K56ac across the genome in Δasf1 compared to the wild type. 401 unique positions carried H3K27me3 in the wild type, while this mark was present at 1142 unique positions in the mutant. For H3K56ac, 4736 unique positions were found for H3K56ac in the wild type and 664 unique positions in the mutant (Supplemental table 6).

E and F. Relative distribution of H3K27me3 (E) and H3K56ac (F) in annotated regions of *S. macrospora* in wild type and Δasf1. Although the general amount of H3K27me3 was increased in the deletion mutant, we did not observe a significant change in the relative positioning of this mark across the shown genomic regions aside from a slight shift from intergenic to exonic regions. An *asf1* deletion leads to a significant decrease of unique H3K56ac positions compared to the wild type, but it appears that a relative increase can be observed in exonic regions.

Figure 7. Hi-C analysis of *S. macrospora* wild type and Δasf1. DNA-DNA interactions recorded in the samples at the given time were plotted against both interacting positions in the genome. Each red dot represents an interaction event. The characteristic red diagonal indicates highly increased interaction frequencies of adjacent positions that are always in close proximity to each other. Even in the younger, mostly vegetative state, the resolution was not high enough to make out specific compartments of the genome, even though some areas of heightened interaction frequencies can be observed. The resolution was much lower in the older sample that has progressed further in the developmental cycle. This might be an indicator of different chromatin states that could be present in cells that are only generated during sexual development. Strong horizontal and vertical lines were detectable in the deletion mutant that appeared to indicate higher interaction frequencies originating from a 600 kb are on chromosome 2 and were subsequently shown to be caused by a duplication of a region on chromosome 2.

Figure 8. Confirmation of a duplication on chromosome 2 in the *asf1* deletion mutant by genome sequencing.

A. Whole genome sequencing of *S. macrospora* Δasf1 confirmed the presence of a duplication in a region of around 600 kb on chromosome 2. The alignment of the reads from sequencing of the mutant genome to the wild type reference genome shows that the duplication consists of two parts, indicating either two separate duplication events or the loss of the internal area of the duplicated region.

B. Schematics for a putative duplication on chromosome 2 of *S. macrospora* Δasf1. Whole genome sequencing proved the presence of a duplication. To test our hypothesis of a tandem duplication with an internal loss event, PCR tests with primers amplifying the genomic borders in the wild type or the newly created, duplication-specific junctions were designed. While the wild type should only yield PCR products for the external borders of the area of interest (asf1_dupl1 x asf1_dupl2, asf1_dupl3 x asf1_dupl4), strains possessing a tandem duplication should also generate products for the new junctions of the duplicated area (asf1_dupl3 x asf1_dupl2) and of the internal region lost in the duplication (asf1_dupl5 x asf1_dupl6).