The G-alpha Gpa1 directs septin localization in the mating projection of *Saccharomyces*

cerevisiae through its Ubiquitination Domain and Endocytic Machinery

Cory P. Johnson[1], Andrew Hart[2], Katherine F. Jarvis[2,4], Sarah G. Latario[2], Sudati

Shrestha[2], Nicholas Leclerc[2], André Khalil[1,3,4], Joshua B. Kelley[1,2*]

[1]Graduate School of Biomedical Science and Engineering, University of Maine, Orono,

ME

[2]Department of Molecular and Biomedical Sciences, University of Maine, Orono, ME

[3]Department of Chemical and Biomedical Engineering, University of Maine, Orono, ME

[4]CompuMAINE Laboratory University of Maine, Orono, ME

*Corresponding Author

joshua.b.kelley@maine.edu

5735 Hitchner Hall, University of Maine, Orono, ME

31 **Keywords:** Saccharomyces cerevisiae, yeast, pheromone, septins, polarity, g-

32 alpha, g-protein, endocytosis, epsin, GPCR, Cdc42, GAP

Abstract

The yeast mating response uses a G-protein coupled receptor (GPCR), Ste2, to detect mating pheromone and initiate mating projection morphogenesis. The septin cytoskeleton plays a key role in the formation of the mating projection, forming structures at the base of the projection. Desensitization of the Gα, Gpa1, by the Regulator of G-protein Signaling (RGS), Sst2, is required for proper septin organization and morphogenesis. In cells where the Gα is hyperactive, septins are mislocalized to the site of polarity, and the cells are unable to track a pheromone gradient. We set out to identify the proteins that mediate Gα control of septins during the *Saccharomyces cerevisiae* mating response by making mutations to rescue septin localization in cells expressing the hyperactive Gα mutant *gpa1^{G302S}*. We found that single deletions of the septin chaperone Gic1, the Cdc42 GAP Bem3, and the epsins Ent1 and Ent2 rescued the polar cap accumulation of septins in the hyperactive Gα. We created an agent-based model of vesicle trafficking that predicts how changes in endocytic cargo licensing alters localization of endocytosis that mirrors the septin localization we see experimentally. We hypothesized that hyperactive Gα may increase the rate of endocytosis of a pheromone responsive cargo, thereby altering where septins are localized. Both the GPCR and the Gα are known to be internalized by clathrin-mediated endocytosis during the pheromone response. Deletion of the GPCR C-terminus to block internalization partially rescued septin organization. However, deletion of the Gpa1 ubiquitination domain required for its endocytosis completely abrogated septin accumulation at the polarity site. Our data support a model where the location of endocytosis serves as a spatial mark for septin structure assembly and that

58   desensitization of the Gα delays its endocytosis sufficiently that septins are placed

59   peripheral to the site of Cdc42 polarity.

63 **Abbreviations**

Introduction

The *Saccharomyces cerevisiae* mating response is driven by a G-protein coupled

receptor (GPCR) signaling pathway that initiates transcription of mating genes,

polarization of the actin cytoskeleton and growth of a mating projection. Mating

projection morphogenesis is dependent on septins, which are cytoskeletal filaments

most famously known for their role in formation of the bud neck (Bridges and Gladfelter,

2015; McMurray et al., 2011). At a high dose of pheromone, as when close to a mate,

yeast form a tight mating projection called a shmoo that is bounded by septins at the

base (Giot and Konopka, 1997; Longtine et al., 1998). At an intermediate dose of

pheromone, as when farther from a mate, the yeast undergo elongated growth towards

the source of pheromone in a gradient tracking process that requires septins to control

direction of growth (Kelley et al., 2015; Segall, 1993).

The α-Factor pheromone is detected by the GPCR Ste2, which initiates signaling

through the heterotrimeric G-protein comprised of the α (Gpa1), β (Ste4), and γ (Ste18)

subunits(Alvaro and Thorner, 2016). While Gβγ signaling has traditionally taken center

stage for its regulation of both Cdc42-mediated polarity and MAPK signaling, Gα also

controls downstream signaling. Gpa1 serves as a scaffold for the MAPK, Fus3, and

through this role promotes the phosphorylation and activation of the formin Bni1

(Matheos et al., 2004; Metodiev et al., 2002). Active Gpa1 traffics to the endosome

through its unique Ubiquitination Domain, a 109 amino acid insertion not found outside

of fungi, and once at the endosome, Gpa1 interacts with the PI3 Kinase Vps34,

promoting PI(3)P production (Dixit et al., 2014a; Slessareva and Dohlman, 2006). We

previously found that Gpa1 activity influences the spatial distribution of septins during

the pheromone response and that septins are required for gradient tracking (Kelley et

al., 2015).

Septins are evolutionarily conserved dynamic cytoskeletal filaments that are

known for their roles as diffusion barriers, cortical actin regulators, and scaffolds that

fortify membrane shape (Spiliotis and McMurray, 2020). During the yeast pheromone

response, septin filaments form at the base of the mating projection, where they serve

as scaffolds for signaling and promote morphogenesis and gradient tracking (Bharucha

et al., 2008; Changwei et al., 2007; Gladfelter et al., 2005; Kelley et al., 2015). Septin

deposition is controlled by the interplay of several Cdc42 effectors including the Rho-

family member Cdc42, the Cdc42-binding septin chaperones called GTPase interactive

components (Gic1 and Gic2), and the Cdc42 GTPase Activating Proteins (GAPs, Rga1,

Rga2, and Bem3) (Khan et al., 2015). The Cdc42 GAPs also interact with the epsins

Ent1 and Ent2, endocytic adaptor proteins that have also been shown to impact septin

function (Aguilar et al., 2006).

The mechanism that controls the spatial distribution of septins during the

pheromone response has not yet been elucidated, but we have previously found that

Gpa1 desensitization is required for the normal localization of septins to the base of the

mating projection (Kelley et al 2015). Gpa1 is desensitized through the actions of the

regulator of G-protein signaling (RGS), Sst2, which serves as a GAP for Gpa1

(Apanovitch et al., 1998; Dohlman et al., 1996). Either deletion of the RGS or mutation

of the Gα to disrupt RGS binding ($gpa1^{G302S}$) leads to septin mislocalization to the site of

polarity (DiBello et al., 1998; Kelley et al., 2015). These septin changes result in a loss

of gradient tracking ability and defects in mating projection morphogenesis (Kelley et al.,

2015).

Here, we have implemented a plan to uncover the pathway that Gpa1 uses to

influence septin regulation. Using microfluidics and live-cell fluorescence microscopy,

we looked for gene deletions that would restore peripheral septin distribution in cells

expressing the hyperactive Gα mutant, *gpa1^{G302S}*. We found that Gpa1 control of septin

distribution during the pheromone response is mediated by an Epsin-Bem3-Gic1

signaling axis. We developed a mathematical model that shows that endocytic events

could generate the patterning that we see with septins. Following this, we disrupted

endocytosis of both the receptor Ste2 and the Gα Gpa1, and found that, while both

contribute to septin organization, Gpa1 endocytosis plays a larger role. Thus, the

ubiquitination domain of Gpa1, which is required for its endocytosis, drives pheromone-

induced septin organization.

Methods

Yeast Strains

Strains used in this study are listed in Table S1. All strains were constructed in

the MATa haploid Saccharomyces cerevisiae parent strain, BY4741. All plasmids used

in this study are included in Table S2. Polar cap (Bem1) and septin (Cdc3) proteins

were tagged with GFP (Huh et al., 2003) and mCherry, respectively, at the native

chromosomal locus through oligonucleotide-directed homologous recombination using

primers listed in Table S3.

Cells were grown in rich medium (YPD) or synthetic medium (SC) at 30°C unless

otherwise indicated. PCR products were transformed into parent yeast strains using

standard lithium acetate transformation (Burke et al., 2000). Individual colonies were

isolated by growth on standard selective media (SC leu-, SC ura-, SC his-, or YPD

G418+). Transformants were verified using fluorescence microscopy, sequencing,

and/or PCR.

Cloning

gpa1$^{\Delta UD\text{-}G302S}$ Plasmid Cloning The gpa1$^{\Delta UD\text{-}G302S}$ plasmid was constructed using the

previously published pRS406-GPA1_truncΔUD (dixit et al 2014), which we received as

a gift from the Dohlman Lab. We mutagenized the pRS406-GPA1_truncΔUD using the

site-directed mutagenesis kit (NEB) in combination with the primers AHM-14 and AHM-

15 listed in table 3. The plasmid was sequenced via PlasmidSaurus

(https://www.plasmidsaurus.com/) to confirm successful mutagenesis of the glycine at

position 302 to serine. Following sequence confirmation, we linearized the plasmid for

genomic integration using AflII (NEB), a silent mutation inserted previously.

Gal-inducible Ste5-CTM Cloning. A galactose-inducible STE5 with the C-terminal

membrane domain (CTM) of SNC2 was excised from the parent plasmid pGS5-CTM

using *SacI-HF* and *ApaI* endonucleases. A pRSII416 plasmid (Chee and Haase, 2012)

was linearized using the same enzymes, followed by a treatment of Antarctic

phosphatase (New England Biolabs) to produce a new parent vector for the Gal-

inducible Ste5-CTM fragment. Verification of the fragment and vector was performed

using gel electrophoresis with a 0.8% TBE-agarose gel to confirm the size of the two

products. The fragment and vector were then ligated together overnight at 16 °C using

T4 DNA ligase. The ligation mixture was transformed into DH5α competent *E. coli* cells

(NEB #C2987), and colonies were isolated by growth on LB agar plates supplemented

with 0.1 mg/mL carbenicillin. Plasmids were extracted from these transformants and

were verified by restriction digest with *SacI-HF* and *ApaI*, followed by gel

electrophoresis.

Microfluidics Experiments

Microfluidic devices were made by mixing SYLGARD 184 silicone polymer at

10(part A):1(part B) and poured directly onto a microfluidics device mold (Suzuki et al.,

2021) that was fabricated by the University of Maine Frontier Institute for Research in

Sensor Technologies (FIRST, https://umaine.edu/first/). Microfluidic chambers were

made as previously described (Suzuki et al., 2021) Molds and coverslips were exposed

to oxygen plasma for 50 seconds in a Harrick Plasma Cleaner PDC32G (Harrick

Plasma) followed by fusion of the device mold to the coverslip. After fusing, the device

was then baked in an oven at 80°C for at least 20 minutes to facilitate fusion of the mold

to the coverslip.

Cultures were grown in 0.45μm-filtered Synthetic Complete + Dextrose to an

OD600 between 0.1-0.8 at 30°C. Live-cell microfluidics experiments were performed

using an IX83 (Olympus, Waltham MA) microscope and a Prime 95B CMOS Camera

(Photometrics) controlled by Cell Sens 1.17 (Olympus). Fluorescence and Differential

Interference Contrast (DIC) images were acquired using an Olympus-APON-60X-TIRF

objective. Z-stacks of GFP and mCherry images were acquired using an X-Cite 120

LEDBoost (Excelitas). Cells were imaged in a "Dial-a-wave" based microfluidic device

that allowed for rapid switching of media while yeast remained in place (Bennet et al.,

2011; Dixit et al., 2014b; Suzuki et al., 2021). Pheromone-containing media was verified

using AlexaFluor 647 dye at 1:8000 dilution (Life Technologies) and imaged in a single

plane. Cells were imaged at 20-minute intervals for 12 hours in Synthetic Complete

media containing Dextrose (SCD) with 300nM pheromone present in the media.

Imaging settings were determined based on experimental needs and were replicated for

repeat experiments.

Ste5-CTM Experiments

Yeast cultures containing the Ste5-CTM (pRSII416) plasmid were diluted and

grown overnight at 30 °C shaking in SC ura- media such that they did not exceed a

saturating concentration (OD600 = 0.8). These cultures were then diluted to OD600 =

0.2, treated with either 0.5% galactose or 0.5% galactose and 10µM pheromone, and

allowed to incubate for 4 hours shaking at 30 °C prior to imaging. Fluorescence

microscopy was used to image Bem1-GFP and Cdc3-mCherry.

Image Analysis

Images were deconvolved using Huygens Software (Scientific Volume Imaging,

Hilversum, Netherlands). For all fluorescent images, deconvolution was carried out with

a theoretical point-spread function, 5 signal-to-noise ratio (GFP images) or 7 signal-to-

noise ratio (mCherry images), maximum 500 iterations, and a quality change threshold

of 0.01 using the classical algorithm. All other settings were set to default. Images were

saved as 16-bit TIFF with linked-scale. Masks of fluorescent signal were created using a

merged 8-bit RGB version of GFP and mCherry channels in FIJI and final segmentation

was done prior to septin localization analysis in MATLAB. All septin localization analysis

was conducted using *whole_cell_cap_v4.m* for cell tracking and *combokeeper_v3.m* for

line scan analysis Septin data was aligned to the polar cap profile using the

regalignraw.m script (GitHub) and subsequently normalized to the fraction of total

signal, as previously described (Kelley et al., 2015; Simke et al., 2022). Data was

normalized to sum to 1 and 95% confidence intervals of the mean intensity were

determined by bootstrapping. The referenced MATLAB scripts for these analyses are

available on Github (https://github.com/Kelley-Lab-Computational-Biology/Spatial-

Normalization).

Agent-Based Model Description

The agent-based model of cargo and vesicle transport was made using MATLAB

(Mathworks). The code is available on Github at https://github.com/Kelley-Lab-

Computational-Biology/Trafficking-Model. The membrane consists of 100 bins, each 0.1

μm wide. The model has a time step of 0.01 seconds, and we ran the model for 200,000

steps per simulation.

-*Agents:* Individual cargo is tracked and can become phosphorylated with a

probability of 5e-5 per time step (termed 1x phosphorylation). Phosphorylated cargo can

be ubiquitinated, which happens with a probability of 0.01 per time step. Cargo which is

phosphorylated and ubiquitinated is now competent for endocytosis. If it diffuses into an

endocytic pit, it will be trapped and will no longer diffuse.

-*Diffusion:* The system is scaled such that diffusion can happen one bin at a time

and recapitulate realistic receptor diffusion rates (Valdez-Taubas and Pelham, 2003).

Movement probability incorporates cargo concentration in adjacent bins to weight

diffusion probability to be more likely to move down the concentration gradient, as

described previously (Azimi et al., 2011). The base probability to move one bin is

0.0024 per time step.

- *Vesicle Trafficking:* Exocytosis frequency has been measured by membrane

capacitance at 2.4 events per minute in yeast that have been spheroplasted (Carrillo et

al., 2015), and estimated to be 24 events per minute from microscopy data (Layton et

al., 2011; Marco et al., 2007). Based on these two values, we chose an intermediate

value of 8 events per minute, as yeast protoplasts may be experiencing cell wall stress

that could decrease exocytosis. Endocytosis is set to happen at twice the rate of

exocytosis with a single 2 bin exocytic event and 2 independent single bin endocytic

events. These rates and sizes are balanced to roughly maintain membrane size.

Exocytic events push the surrounding bins away from the site of exocytosis, while

endocytic events remove the bin affected, bringing the surrounding bins next to each

other. Location of exocytic events is determined by a probability distribution based on

the Bem1 and Exo84 distributions calculated during the pheromone response previously

(Brennwald, 2013; Kelley et al., 2015). We used the probability distribution equal to

$[Bem1]^2$ x [Exo84] to account for the role of Cdc42-GTP in both directing actin

organization and promoting vesicle fusion. Location of endocytic events is determined

by selecting the most mature endocytic pit. Whenever an endocytic pit is removed, a

new one is formed at the site of the highest ubiquitinated cargo, consistent with the

finding that the presence of cargo regulates endocytic pit location (Pedersen et al.,

2020).

Results

A scheme to identify proteins responsible for Gpa1 control of septins

We previously found that septin organization at the periphery of the polarity site

in the yeast mating projection is dependent on the ability of the RGS, Sst2, to stimulate

Gpa1 hydrolysis of GTP (Kelley et al., 2015). Here we set out to discover the signaling

process that leads from active Gpa1 to control of septins. Our strategy was to delete

genes involved in septin regulation (Figure 1A) from cells expressing either WT Gpa1 or

the hyperactive gpa1^{G302S} mutant that is defective for binding to Sst2. Each strain has

the Cdc42 scaffolding protein Bem1 tagged with GFP as a polar cap marker and the

septin Cdc3 tagged with mCherry as a marker for septins. We then assessed these

deletion strains for the ability to restore septin localization to the periphery of the polar

cap. In cells responding to pheromone, septins form structures at the base of the mating

projection, peripheral to the collection of polarity machinery that contains Cdc42 and

Bem1 and is collectively known as the polar cap (Figure 1B). To map our baseline

septin profiles we used live-cell fluorescence microscopy and computational image

analysis of Bem1-GFP and Cdc3-mCherry. We collected high-quality images at single

time-points to visualize discrete septin structures (Figure 1B). However, because septin

structures change through the pheromone response, we used live-cell microscopy in a

microfluidic device to monitor the polar cap and septins through 12 hours of pheromone

response using a flat dose of 300nM pheromone, which is sufficient to stimulate shmoo

formation and prevent sustained elongation (Figure 1C and D) (Kelley et al., 2015).

To quantify the average distribution of septins along the periphery of the cell, we

spatially normalized our septin profiles to the center of the polar cap – indicated by peak

fluorescence of Bem1-GFP, a Cdc42-GTP binding protein, as we have done previously

(Kelley et al., 2015; Simke et al., 2022). We averaged the spatially normalized septin

profiles across time points and cells to yield an average septin distribution for each

strain. We then bootstrapped 95% confidence intervals (indicated by shaded regions):

non-overlapping confidence intervals indicate statistically significant differences in

septin distribution for $p < 0.05$. Our metric for septin rescue is shifting the septin

distributions in the hyperactive $gpa1^{G302S}$ cells back toward the septin shape in WT cells

(Figure 1E).

Gpa1 regulation of septins occurs in addition to a default septin regulation

pathway in response to pheromone.

Septins are organized into a ring structure during mitosis in haploids, where

Gpa1 levels are high, and in diploids, where Gpa1 levels are much lower (de Godoy et

al., 2008). We would anticipate that mechanisms for septin organization during cell

division are independent of Gpa1, however, we do not know if Gpa1 is required for

septin organization during the pheromone response. We tested whether Gpa1 activity is

required for septin organization during the pheromone response by initializing MAPK

signaling and Cdc42-mediated polarity by using a mutant MAPK scaffold, Ste5, modified

on its C-terminus to contain a transmembrane domain (Ste5-CTM) (Pryciak and

Huntress, 1998). Expression is controlled by a GAL promoter such that the pheromone

pathway is inducible by galactose, without activation of the receptor Ste2, or activation

of the hetero-trimeric G-protein, Gpa1, Ste4, and Ste18. We grew cells in raffinose and

then treated with either 0.5% galactose for 4 hours, or 0.5% galactose and 10 μM

pheromone for 4 hours, and imaged Bem1-GFP and Cdc3-mCherry. Cells were able to

place septin structures at the base of the mating projection in the presence of galactose

alone, or with both galactose and pheromone (Figure 1F). When we quantified the

average location of septins in this experiment (Figure 1G), we find that in both

treatments, the average distribution is moderately flat, suggesting that the spatial

placement of septins is more random than in cells that are not expressing Ste5-CTM.

Thus, Gpa1 activation is not required for septin structures to be formed during the

pheromone response, indicating that Gpa1 control of septins is layered on top of normal

septin regulatory machinery.

Gic1, but not Gic2, contributes to Gpa1-driven septin organization

The Cdc42-interacting paralogs, Gic1 and Gic2, are Cdc42-GTP binding proteins

involved in the recruitment of septins to the polarity site during mitotic budding and in

septin bundling activity *in vitro* (Figure 2A) (Iwase et al., 2006; Sadian et al., 2013). We

constructed single-gene-deletions of Gic1 and Gic2 from yeast expressing hyperactive

gpa1^{G302S} and examined the polar cap (Bem1-GFP) and septin (Cdc3-mCherry)

distribution (Figure 2B, C, D). High resolution, single time point images reveal that

hyperactive *gpa1*^{G302S} cells lacking Gic1 placed septin structures further from the polar

cap compared to cells lacking Gic2 (Figure 2B). We found through live-cell imaging (as

conducted above) that cell morphology and septin distribution in hyperactive *gpa1*^{G302S}

cells lacking Gic1 resembled WT-like phenotypes (Figure 2C). This observation was

supported by septin distribution quantification performed as above (Figure 2E). Thus,

Gic1 has a role in septin organization during the pheromone response distinct from that

of Gic2 and is required for Gpa1 control of septins.

The Cdc42GAP, Bem3, mediates Gpa1 control of septin structures

Cdc42 GAP activity promotes septin structure assembly on the plasma

membrane (Figure 3A) (Gladfelter et al., 2002; Khan et al., 2015). Yeast express three

Cdc42 GAPs: Rga1, Rga2, and Bem3 (Smith et al., 2002). At high resolution we found

that all three single-gene deletions in hyperactive $gpa1^{G302S}$ mutants had at least some

overlap between septins and the polar cap, and at the same time all three mutants had

septin structures set up far from the polar cap as well (Figure 3B). However, live-cell

microscopy at lower pheromone dose revealed that cells expressing hyperactive

$gpa1^{G302S}$ and lacking Bem3 displayed more definitive separation between septins and

the polar cap compared to mutants lacking Rga1 or Rga2 (Figure 3C-E). When we

quantitatively examined the septin organization over time, we confirmed that

hyperactive $gpa1^{G302S}$ mutants lacking Bem3 robustly rescued septin distribution

(Figure3F). We also found that hyperactive $gpa1^{G302S}$ mutants lacking either Rga1 or

Rga2 mildly rescued aberrant septin distribution (Figure 3F). Thus, while all three Cdc42

GAPS influence Gpa1 control of septins, Bem3 contributes more significantly than Rga1

or Rga2.

The endocytosis adaptors Ent1 and Ent2 direct Gpa1 dependent pheromone-

induced septin organization

Cdc42 GAPs bind directly to the epsin endocytic-adaptor proteins, Ent1 and

Ent2, in an interaction thought to couple Cdc42 regulation to endocytosis (Aguilar et al.,

2006). Additionally, epsins are known to differentially contribute to septin organization

during cell division (Mukherjee et al., 2009). We hypothesized that epsins may also

contribute to septin organization during the pheromone response (Figure 4A). As

previously described, we investigated septin organization in hyperactive $gpa1^{G302S}$

mutants lacking either Ent1 or Ent2 at high resolution (Figure 4B), via time course

imaging (Figure 4C-D), and through time-averaged quantitation of peak septin

distribution in relation to the polar cap (Figure 4E). At high resolution we found cells

lacking Ent1 clearly rescued aberrant septin structures. Additionally, cells lacking Ent2

rescued the septin phenotype but with a pronounced peak of septins at the center of the

polar cap (Figure 4B). Timelapse imaging also revealed the same phenotype found at

high resolution (Figure 4C-D) and quantitation of peak septin distribution confirmed that

single-gene deletion of either epsin robustly rescues aberrant septin distributions in

hyperactive $gpa1^{G302S}$ mutants (Figure 4E). Thus, both Ent1 and Ent2 are mediators of

Gpa1 control of septin distribution during the pheromone response.

An agent-based model of vesicle trafficking shows that location of endocytosis

could explain septin organization.

We have found that the endocytic adaptor protein Ent1 and Ent2 are both

involved in the control of septin distribution by Gpa1. Furthermore, rescue of septin

organization in each experiment has been through a loss of function (deletion) mutation,

which should slow cellular processes in which these proteins are involved. Epsins are

endocytic adapter proteins, which led us to hypothesize that endocytosis itself may be

able to serve as a spatial mark for septin organization. In this scenario, changes in the

rate that cargo becomes competent for endocytosis could give rise to the septin

distribution changes that we see. To test this, we developed an agent-based stochastic

model of protein trafficking. This model is described in Figure 5. Cargo is delivered to

the membrane, diffuses, undergoes endocytosis, and is tracked spatially over time. Like

the receptor, Ste2, it has a probability to become phosphorylated, and phosphorylated

cargo has a probability to become ubiquitinated and therefore competent for

endocytosis (Figure 5A) (Lu et al., 2016). These reactions occur in a discretized 1

dimensional membrane that can grow and shrink with endo- and exocytosis (Figure 5B).

The location of exocytosis is based upon the distributions of Bem1 and Exo84 that we

have previously measured (Figure 5C) (Kelley et al., 2015). The number of endocytic

pits is kept constant, such that when an endocytic pit is endocytosed, a new pit forms at

a new site based upon the amount of ubiquitinated cargo, consistent with polarized

endocytic cargo driving the polarization of endocytosis (Figure 6 D) (Pedersen et al.,

2020). Diffusion within the membrane is carried out as described by Azimi et al, where

the direction of diffusion is biased by the occupancy of the adjacent bins such that

diffusion into bins with fewer molecules is more likely (Figure 5E) (Azimi et al., 2011).

Ubiquitinated cargo that enters an endocytic pit becomes trapped there, and eventually

will be internalized (Figure 6F). We performed simulations where we varied the rate of

phosphorylation and examined the distribution of ubiquitinated cargo and the location of

endocytic events.

To test the effects of increasing the rate of cargo endocytosis, we varied the rate

of phosphorylation in our model, and ran multiple simulations with each condition. We

simulated cargo location and modification on the surface of the cell over 30+ minutes in-

model time (Figure 6A). Increasing the rate of phosphorylation from baseline (1x) 5-fold

or 25-fold increases the fraction of endocytic events that occur at the center of the polar

cap (Figure 6B, C). The increased phosphorylation rates mean that cargo cannot travel

along the membrane as far before being internalized.

Altering the rate that cargo becomes competent for endocytosis allows us to

qualitatively recreate the septin distributions we have measured in WT and *gpa1*^{G302S}

strains at 300 nM. When we compare our simulation of endocytosis locations to our

experimentally measured septin distributions, we see that the 1X phosphorylation

simulation of endocytic events is similar to septin distribution in our WT cells treated

with 300 nM pheromone. The simulations of endocytosis at a high rate of endocytic

competency, the 25X phosphorylation rate, generate endocytic events with a distribution

similar to septins in the hyperactive *gpa1*^{G302S} mutant strain (Figure 7A). Our model

simulations suggest that a 5x higher rate of endocytosis competency would lead to

endocytosis events closer to the polar cap than we had measured (Figure 7B). If our

model is accurately recapitulating the system, then we would expect that the endocytic

distribution of the 5X phosphorylation rate would predict the septin distribution in the

presence of a high dose of pheromone. In this case, the pathway activity would be

high, but the Gα would still be able to turn off. We treated WT cells with 10 μM

pheromone for 2 hours and measured Cdc3-mCherry distribution relative to the polar

cap (Figure 7B). The measured septin distribution matches the simulation remarkably

well, even down to the same asymmetry in both simulated endocytic events and septin

distribution, which we attribute to happy coincidence, as we have no reason to believe

there is an underlying asymmetry in the biology and our model should not have any

intrinsic asymmetry. These data do not necessarily mean that *gpa1*^{G302S} increases

phosphorylation of a cargo *per se*, but it does suggest that increasing the rate that a

cargo becomes competent for endocytosis and coupling that endocytic event to

assembly of macromolecular complexes of septin would be capable of creating the

septin distributions we see in WT and *gpa1^{G302S}* mutants.

Endocytosis of Gα and the pheromone receptor directs pheromone-induced septin

distribution

We have found that the endocytic adaptor proteins Ent1 and Ent2 are required

for Gα control of septins and that our mathematical model can reconstruct the

distribution of septin peaks seen in our experimental data by altering the rate that cargo

become endocytically competent. These data raise the question: Can endocytic

cargo(s) drive septin deposition during the pheromone response?

The pheromone receptor, Ste2, is both phosphorylated and ubiquitinated at its C-

terminus to control its endocytosis (Dunn and Hicke, 2001; Hicke et al., 1998). We

constructed cells lacking the receptor C-terminus from codon 326 (*ste2$^{\Delta C}$*), a critical

region for control of receptor endocytosis (Konopka et al., 1988). We then investigated

the effect of the *ste2$^{\Delta C}$* on septin localization in cells expressing *gpa1^{G302S}*. We found

that *gpa1^{G302S}* mutants lacking the C-terminus of Ste2 partially rescued the septin defect

(Figure 8A,B). Morphologically, the *gpa1^{G302S} ste2$^{\Delta C}$* double mutants are very round,

with poor mating projection formation, consistent with septin defects. In contrast to this

partial rescue, epsin deletions completely rescued the septin organization and restored

elongated (albeit still abnormal) growth, suggesting there could be another cargo driving

septin distribution.

Gpa1 contains a ubiquitination domain (UD) that promotes endocytosis and

subsequent trafficking through the endomembrane system (Dixit et al., 2014a; Wang et

al., 2005a). We hypothesized that endocytosis of Gpa1 (via the UD) directs septin

deposition. We constructed an integrating plasmid to replace the wildtype Gpa1 locus

with a $gpa1^{G302S}$ lacking the UD ($gpa1^{G302S-\Delta UD}$) and monitored both the polar cap and

septins as we did in Figures 1-4 (Dixit et al., 2014a). Our single time point imaging of

hyperactive mutants lacking the UD of Gpa1 showed rescue of the $gpa1^{G302S}$-induced

septin defects, similar to the rescue seen in the $ste2^{\Delta C}$ (Figure 8A). The degree of

rescue was more apparent in live-cell microscopy carried out under a more

physiological pheromone dose of 300 nM in microfluidic devices (Figure 8C). Under

these conditions, there is clear elongation of the cells, with septin structures well

separated from the polar cap.

While the $gpa1^{G302S}$ mutant leads to increased pathway sensitivity, the gpa1$^{\Delta UD}$

mutant is known to lead to decreased pathway sensitivity (DiBello et al., 1998; Wang et

al., 2005b). It is noteworthy that cells expressing $gpa1^{G302S-\Delta UD}$ arrested in G1 for less

time than WT cells. When these cells were exposed to 300nM pheromone – as all other

strains were exposed to – we found that cells reverted to mitotic behavior after only 8

hours of pheromone exposure (Movie S1). Both WT and $gpa1^{G302S}$ cells under these

conditions will respond to pheromone for the entire 12 hours, indicating that the deletion

of the UD is dominant to the gpa1^{G302S} reduction of RGS interaction, at least for arrest of

the cell cycle. For the purposes of our septin experiments, we found that increasing the

concentration of pheromone to 1μM lead to the $gpa1^{G302S-\Delta UD}$ cells continuing to

develop mating projections for the full 12-hour time course (Movie S1). Additionally, we

quantified septin distribution in the $gpa1^{G302S-\Delta UD}$ mutants at 300nM pheromone and

1µM pheromone and found that pheromone concentration did not affect septin

distribution profiles during the pheromone response (Figure S1).

Finally, we quantified the septin distributions of cells expressing the hyperactive

$gpa1^{G302S}$ and lacking either the C-terminus of Ste2 ($gpa1^{G302S}$ $ste2^{\Delta C}$) or the UD of

Gpa1 ($gpa1^{G302S-\Delta UD}$) relative to the center of the polar cap as described above. We

found that hyperactive cells lacking the C-terminus of Ste2 partially rescued the

$gpa1^{G302S}$-induced septin defects, while cells lacking the Gpa1 UD fully rescued the

septin defect in relation to the polar cap (Figure 8E). Disruption of endocytosis of Gpa1,

and to a lesser degree, Ste2, must therefore slow septin structure formation in the

$gpa1^{G302S}$ mutant, allowing septins sufficient time to move to the periphery of the polarity

site prior to forming higher-order structures. Thus, Gpa1 influences septin organization

via protein domains required for endocytosis: the Gpa1 UD and the C-terminus of Ste2.

Discussion

Here we describe the mechanism for Gpa1 control of septins in the yeast

pheromone response. We tested for rescue of septin organization by deletion of known

components of the septin regulatory system, including the Cdc42 effectors Gic1 and

Gic2, and the Cdc42-GAPs Rga1, Rga2, and Bem3. We found that pheromone-induced

septin organization is dependent upon Gic1, but not its paralog Gic2 and preferentially

uses the Cdc42 GAP Bem3. Because Cdc42 GAPs bind to the Ent1 and Ent2 endocytic

adaptors, we examined the ability of epsin deletions to rescue Gpa1-induced septin

defects. We found that Ent1 and Ent2 contribute equally to pheromone-induced septin

organization. Based on Ent1/2 roles in endocytosis, we used an agent-based

mathematical model to test if the location of endocytic events could explain septin

distribution. Our model recapitulates the wild type and mutant Gpa1-induced septin

distributions when the following two assumptions are made: 1) that *gpa1^{G302S}* alters

when a cargo is marked for clathrin-mediated endocytosis and 2) that endocytosis is

spatially coupled to septin structure organization. Based on these results, we

hypothesized that a pheromone-dependent endocytic cargo was responsible for Gpa1

control of septins. A likely candidate was the pheromone responsive GPCR, Ste2, which

is known to undergo endocytosis in response to pheromone (Dores et al., 2010; Hicke

and Riezman, 1996; Lu et al., 2016). However, when we disrupted Ste2 endocytosis,

we only saw a moderate effect on septin organization. Gpa1 internalization is less well

characterized but is known to be dependent upon its Ubiquitin Domain (UD) (Dixit et al.,

2014a). Remarkably, blocking Gpa1 endocytosis by removal of the UD led to complete

recovery of septins to the periphery of the polar cap in the presence of the hyperactive

gpa1^{G302S} mutation.

Endocytosis of Cargo as a Marker for Septin Organization

Septin organization occurs during mitosis, with no Gα activity necessary, and we

have shown that Gpa1 activity is dispensable for septin organization during the

pheromone response, as initiating signaling with the MAPK scaffold, Ste5, leads to

septin structure formation at the base of the mating projection. Thus, Gpa1 adds an

additional layer of septin regulation on top of the normal process. Our data lead us to

the following model (Figure 9): In the presence of pheromone, when Gpa1 and Ste2 are

phosphorylated and then ubiquitinated, they recruit epsins and Cdc42 GAPs to the site

of endocytosis. The recruitment of the Cdc42GAP promotes Cdc42 hydrolysis of GTP,

and subsequent organization of higher order septin structures in a Gic dependent

manner. When Gpa1 is hyperactive, it becomes competent for endocytosis sooner,

recruiting epsins and Cdc42 GAPs earlier, and driving septin organization closer to the

polar cap.

Previous models of septin organization incorporated exocytosis as a driving force

for clearance of septins from the bud site, and Cdc42 GAP function as the "timer"

dictating how far septins were pushed before assembly into higher order structures

(Okada et al., 2013). Our data add a new wrinkle to this, suggesting that epsin-mediated

endocytosis may control *where* septins organize into higher order structures. Thus,

vesicle trafficking drives the movement of septins away from the polar cap on the

"conveyor belt" of the plasma membrane (moving away from exocytic events and

towards endocytic events) (Figure 9B). Coupled to this, endocytosis provides the spatial

cue that septins should transition from "passengers" on this conveyor belt to

establishing higher-order structures at the site of clathrin mediated endocytosis.

Disruption of receptor endocytosis showed a partial rescue of septin, suggesting

that endocytic cargoes, broadly, promote septin organization. But why does Gpa1 have

a larger effect? Both receptor and Gα undergo clathrin-mediated endocytosis in

response to pheromone, but Ste2 endocytosis is likely not linked to Gpa1 activation. So

Ste2 recruitment of epsins and GAPs may be contributing to a fraction of septin

organization as normal. As such, removal of the Ste2-contribution reduces some of the

septin organization signaling happening close to the polar cap, but hyperactive Gpa1

continues to drive significant amounts of septin organization close to the polar cap.

Our model for Gα control of septins during the pheromone response predicts that

the distance between the polar cap and septin structures would be smaller for high

doses and get larger at lower doses. We found that septins accumulated closer to the

polarity site at 10 uM than at 300nM (Figure 7B) and indeed, this is what we saw in our

previous work (Kelley et al., 2015), where cells responding to an intermediate of

pheromone dose (~75 nM, center of a 0-150 nM gradient) established septin structures

farther from the center of the polar cap than cells exposed to a high dose of pheromone

(300 nM pheromone). Our model couples pheromone dose to septin spatial regulation,

which is necessary for a mechanistic understanding of gradient tracking.

Implications for gradient tracking

We previously found that cells without functional septins grow in a straight line,

unable to correct their orientation by turning (Kelley et al., 2015). WT cells that turn have

septins on the inside of the turn. Thus, asymmetric distribution of septins on the inside

of the turn (up-gradient) appears to be required for error correction during gradient

tracking (Kelley et al., 2015). Our findings that Gpa1 activity level influences septin

organization through the endocytosis machinery provides a potential mechanism for

asymmetric septin organization (Figure 9B). Simply, higher active Gpa1 on the

upgradient side of the cell results in increased Gpa1 endocytosis. The endocytic

machinery recruited to the site of endocytosis promotes septin organization on the up-

gradient side of the cell. The accumulation of septins could then facilitate the turning of

the cell. Future studies will be needed to address how activation of Gpa1 influences

internalization through its UD.

Epsins and Cdc42-GAPs

These results raise the question of whether organization of the septin ring during

mitosis is also controlled by endocytic events? The Aguilar lab has found that

overexpression of the Epsin N-terminal Homology Domain from Ent2 (ENTH2) leads to

defects in the organization of septins during mitosis (Mukherjee et al., 2009).

Furthermore, they found that ENTH2 overexpression affects Bem3 localization. It may

be that Epsin recruitment of Bem3 is broadly used to control septin organization.

The Bi and Goryachev labs showed that exocytosis serves to drive septin

formation into a ring structure during mitosis, with the assumption that GAP activity

would serve as the limit on expansion of the ring, resulting in septin organization at the

edge of the polar cap (Okada et al., 2013). It may be that the initial cue to form septin

structures during mitosis arises from endocytic cargoes recruiting Epsin/Bem3

complexes, and that further action by Rga1 and Rga2 bound to these new septin

structures serves to reinforce structure formation. By contrast, Rga1 and Rga2 show

little localization during the pheromone response (Kelley et al., 2015) so the bulk of the

responsibility to control septins may fall to Bem3.

Gα endocytosis as a signaling event

Here, we show that Gpa1 influences septin organization through its UD, the

yeast-specific domain required for Gpa1 endocytosis (Figure 8) (Dixit et al., 2014a;

Wang et al., 2005b). While the Gpa1 UD is unique to yeast Gα, mammalian cells also

endocytically regulate Gα via β-arrestins (Smith et al., 2021). Mammalian endocytosis of

Gα occurs indirectly via receptor C-terminal interaction and not directly with the Gα

subunit (Shenoy and Lefkowitz, 2003). Indeed, many components of clathrin mediated

endocytosis are conserved between yeast and mammalian cells: 1) mammalian epsins

bind AP2 (Chen et al., 1998), 2) mammalian epsins bind Cdc42-GAP (Coon et al.,

2010), and mammalian epsins maintain the ubiquitin interacting motif necessary for

endocytic cargo binding (Wendland, 2002). Therefore, future research will need to

examine whether mammalian cells may also utilize Gα endocytosis as a spatial

landmark for septin organization during morphogenesis.

Acknowledgements

The authors wish to thank Holly Goodson and Melissa Maginnis for helpful discussions.

The authors would also like to thank Niklas Hase, the Shmoo Crew and other former

members of the Kelley Lab for experimental and logistical input including. This work was

supported by the National Institutes of Health Grant R15-GM140409 to JBK.

Author Contributions

CPJ wrote the manuscript -original draft, review, and edits, visualization; designed,

conducted, analyzed, and oversaw the experiments. AH wrote the manuscript -review

and edits, collected, and analyzed data. KJF wrote the manuscript -original draft,

review, and edits, designed and conducted experiments (mathematical modeling), and

created software. SGL wrote the manuscript -review and edits, collected, and analyzed

data. SS wrote the manuscript -review and edits, collected and analyzed data. NL wrote

the manuscript -review and edits, collected and analyzed data. AK wrote the

manuscript-review and edits, and contributed supervision. JBK wrote the manuscript -

original draft, review and edits, visualization, designed experiments, acquired funding,

contributed supervision, conceptualized and oversaw the project.

 Figure Legends

Figure 1. Pheromone-induced septin accumulation occurs at the center of the

polar cap in Hyper Gα mutants.

A) Schematic of key players in septin deposition in yeast. Cdc42-GTP binds to GICs,

which are required for septin recruitment. GAP activity promotes Cdc42 hydrolysis of

GTP to GDP and promotes septin structure formation. B) Representative images of

pheromone treated yeast (2hrs with saturating pheromone) expressing endogenous

Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) in WT (top panels) and in the

hyperactive Gα (*gpa1*^{G302S}) mutant (bottom panels). Scale bar represents 3μm and

white arrows indicate sites of peripheral septin localization, while black arrow indicates

septins localized close to the polar cap. C) Representative images of WT cells

expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to

300 nM pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top

row), 360min (middle-row), and 720min (bottom row) are displayed with split-channel

(left and middle columns) and merged images (right column). Arrows indicate septins

organizes at the base of projections, away from the polar cap. Scale bar represents

5μm. D) Representative images of *gpa1*^{G302S} yeast expressing endogenous Bem1-GFP

(Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating pheromone in a

microfluidic chamber for 12hrs. Images of cells at 0min (top row), 360min (middle-row),

and 720min (bottom row) are displayed with split-channel (left and middle columns) and

merged images (right column). Arrows indicate septins accumulating close to the polar

cap. Scale bar represents 5μm. E) Time-averaged quantification of the fraction (y-axis)

of septin and its distribution relative to the polar cap (x-axis). Lines are representations

of time-averaged data (37 time points; 20min intervals for 12hrs) and shaded regions

surrounding lines are 95% confidence intervals. Lines are representative of two

combined experiments (WT n = 51 cells and $gpa1^{G302S}$ n = 73 cells). F) Representative

images of cells expressing the GAL-inducible Ste5-CTM that activates the mating

pathway without activation of Ste2, Gpa1, or Ste4/Ste18. Cells were treated with 0.5%

galactose or 0.5% galactose & 10 µM pheromone for 4 hours. Arrows indicate sites of

septin organization at the base of the mating projection. G) Quantitation of septin

organization in cells expressing the GAL-inducible Ste5-CTM. Shaded area represents

the 95% confidence intervals. n = 128 for +Gal and n = 139 for +Gal +Pheromone,

across two separate experiments.

Figure 2. Gα control of septins is preferentially mediated by Gic1 rather than Gic2.

A) Schematic shows that we predict decreased septin recruitment in individual genetic

deletions of Gic1 or Gic2 in yeast with hyperactive Gα, *gpa1^G302S^*. B) Representative

images of pheromone treated yeast (2hrs with saturating pheromone) expressing

endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) in *gpa1^G302S^* cells

lacking Gic1 or Gic2. Scale bar represents 3μm and white arrows indicate sites of

peripheral septin localization. C) Representative images of *gpa1^G302S^-gic1Δ* cells

expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to

300 nM pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top

row), 360min (middle row), and 720min (bottom row) are displayed with split-channel

(left and middle columns) and merged images (right column). Scale bar represents 5μm.

D) Representative images of *gpa1^G302S^gic2Δ* yeast expressing endogenous Bem1-GFP

(Polar Cap) and Cdc3-mCherry (Septin) exposed to 300 nM pheromone in a microfluidic

chamber for 12hrs. Images of cells at 0min (top row), 360min (middle-row), and 720min

(bottom row) are displayed with split-channel (left and middle columns) and merged

images (right column). Scale bar represents 5μm. E) Time-averaged quantification of

the fraction (y-axis) of septin distribution relative to the polar cap (x-axis). Lines are

smoothed representations of time-averaged data (37 time points; 20min intervals for

12hrs) and shaded regions surrounding lines are 95% confidence intervals. Lines are

representative of two combined experiments (*gpa1^G302S^-gic1Δ* n = 63 cells and

gpa1^G302S^-gic2Δ n = 68 cells).

Figure 3. Gα control of septins is preferentially mediated by the Cdc42GAP Bem3,

rather than Rga1 or Rga2.

A) Schematic representation of our predicted outcome for individual genetic deletion of

the GAPs Rga1, Rga2, or Bem3 in *gpa1^{G302S}* yeast. We predicted the lack of individual

Cdc42-GAPs would reduce septin recruitment to sites of polarity during the pheromone

response and ultimately rescue the *gpa1^{G302S}* phenotype of early septin deposition (top-

panel). B) Representative images of pheromone treated yeast (2hrs with saturating

pheromone) at high magnification (200X prior to image cropping) expressing

endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) in *gpa1^{G302S}* cells

lacking Rga1 (top row), Rga2 (middle row), or Bem3 (bottom row). Scale bar represents

3µm. Back arrows indicate sites of septin structure close to the polar cap and white

arrows indicate sites of peripheral septin localization. C) Representative images of

gpa1^{G302S}-rga1Δ cells expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-

mCherry (Septin) exposed to saturating pheromone in a microfluidic chamber for 12hrs.

Images of cells at 0min (top row), 360min (middle-row), and 720min (bottom row) are

displayed with split-channel (left and middle columns) and merged images (right

column). Scale bar represents 5µm. D) Representative images of *gpa1^{G302S}-rga2Δ*

yeast expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin)

exposed to saturating pheromone in a microfluidic chamber for 12hrs. Images of cells at

0min (top row), 360min (middle-row), and 720min (bottom row) are displayed with split-

channel (left and middle columns) and merged images (right column). Scale bar

represents 5µm. E) Representative images of *gpa1^{G302S}-bem3Δ* yeast expressing

endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating

pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top row),

360min (middle-row), and 720min (bottom row) are displayed with split-channel (left and

middle columns) and merged images (right column). Scale bar represents 5μm. F)

Time-averaged quantification of the fraction (y-axis) of septin distribution relative to the

polar cap (x-axis). Lines are smoothed representations of time-averaged data (37 time

points; 20min intervals for 12hrs) and shaded regions surrounding lines are 95%

confidence intervals. Lines are representative of two combined experiments ($gpa1^{G302S}$-

$rga1\Delta$ n = 79 cells, $gpa1^{G302S}$-$rga2\Delta$ n = 80 cells, and $gpa1^{G302S}$-$bem3\Delta$ n = 87 cells).

Figure 4. Gα control of septins can be mediated by either of the endocytic adaptor proteins, Ent1 or Ent2.

A) Schematic representation of our predicted outcome for individual genetic deletion of Ent1 or Ent2 in $gpa1^{G302S}$ yeast. We predicted the lack of either Ent1 or Ent2 would reduce septin recruitment to sites of polarity during the pheromone response and ultimately rescue the $gpa1^{G302S}$ phenotype of early septin deposition (top-panel). B) Representative images of pheromone treated yeast (2hrs with saturating pheromone) expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) in $gpa1^{G302S}$ cells lacking Ent1 (middle-panels) and in $gpa1^{G302S}$ cells lacking Ent2 (bottom-panels). Scale bar represents 3μm and white arrows indicate sites of peripheral septin localization. C) Representative images of $gpa1^{G302S}$-$ent1\Delta$ cells expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top row), 360min (middle-row), and 720min (bottom row) are displayed with split-channel (left and middle columns) and merged images (right column). Scale bar represents 5μm. D) Representative images of $gpa1^{G302S}$-$ent2\Delta$ yeast expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top row), 360min (middle-row), and 720min (bottom row) are displayed with split-channel (left and middle columns) and merged images (right column). Scale bar represents 5μm. E) Time-averaged quantification of the fraction (y-axis) of septin distribution relative to the polar cap (x-axis). Lines are smoothed representations of time-averaged data (37 time points; 20min intervals for 12hrs) and shaded regions surrounding lines are 95% confidence intervals.

Lines are representative of two combined experiments ($gpa1^{G302S}$-$ent1\Delta$ n = 92 cells

and $gpa1^{G302S}$-$ent2\Delta$ n = 55 cells).

Figure 5. A stochastic agent-based model of protein trafficking.

Schematic representation of the mathematical model of protein trafficking in which, (A)

naïve cargo (unmodified) is first phosphorylated, then ubiquitinated. Upon ubiquitination,

the cargo is considered "endocytically competent". (B) The plasma membrane is

represented by a 1-dimensional line segmented using 100 initial bins. Bins are removed

via endocytosis and new bins are added via exocytosis. (C) The location of exocytosis

is restricted to the center of the membrane with the given probability distribution. (D) In

contrast, the location of endocytosis depends on two factors: location of endocytic pit

formation and endocytic pit maturity. Endocytic pits form at sites with a high

concentration of endocytosis competent cargo. Endocytic pits cannot diffuse, but their

relative position on the membrane changes as bins are added and removed by vesicle

trafficking. (E) To determine the directionality of cargo diffusion we employed an agent-

based probability model. Briefly, the probability a cargo will diffuse to a bin is inversely

correlated with how much cargo is currently in that bin. (F) Finally, once a ubiquitinated

cargo enters an endocytic pit, it can no longer diffuse, leading to accumulation of

endocytic cargo.

Figure 6. Simulations of cargo becoming competent for endocytosis at higher

rates move endocytic events closer to the polar cap.

To change endocytic competency rate, we modified the phosphorylation probability 1-,

5-, and 25-fold. (A) Kymographs from individual simulations showing all cargo on the

membrane over time. (B) Kymographs from individual simulations showing ubiquitinated

cargo on the membrane over time. (C) Histograms of the location of endocytic events

for 11 independent simulations per phosphorylation probability (1-, 5-, and 25-fold). 1X

phosphorylation rate yields a distribution of endocytic events that mirrors WT septin

localization. While increased phosphorylation probabilities (5-fold and 25-fold higher)

result in endocytic events occurring closer to the site of polarity.

Figure 7. Computational modelling predicts that endocytic cargo dynamics are capable of generating septin distribution patterns.

Comparisons of model-generated simulations of the distribution of endocytic events with experimentally determined septin distributions. All plots have been normalized to have a maximum value of 1. A) Comparison of the low (1x phosphorylation) and high (25x phosphorylation) rates of endocytosis competency from Figure 6 with the WT and $gpa1^{G302S}$ septin distributions in 300 nM pheromone from Figure 1. B) The model suggests that a higher rate of endocytosis competency would push endocytic events closer to the center of the polar cap (on the left). Since we hypothesize that rate of endocytosis competency is influenced by Gpa1 activity levels, we added a high level of pheromone (10 µM) and measured septin distribution (right). Shaded area represents 95% confidence intervals, n = 293 cells across two independent experiments.

Figure 8. Gα control of septin deposition is driven by its ubiquitination domain

(A) Schematic representation of our predicted outcome for genetic deletion of the

receptor C-terminus or the ubiquitin domain (UD) of Gα – both domains are required for

the endocytosis of either protein – in $gpa1^{G302S}$ mutant yeast. We predicted that cells

lacking the capacity to drive endocytosis of either the receptor, Ste2, or the Gα, Gpa1,

would reduce septin recruitment to sites of polarity during the pheromone response and

ultimately rescue the $gpa1^{G302S}$ phenotype of early septin deposition (top-panel). B)

Representative images of pheromone treated yeast (2hrs with saturating pheromone)

expressing endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) in

$gpa1^{G302S}$ cells lacking the receptor C-terminus ($gpa1^{G302S}$-$ste2^{\Delta C}$) (middle-panels) and

in $gpa1^{G302S}$ cells lacking the UD of Gα ($gpa1^{G302S-\Delta UD}$) (bottom-panels). Scale bar

represents 3µm. (C) Representative images of $gpa1^{G302S}$-$ste2^{\Delta C}$ cells expressing

endogenous Bem1-GFP (Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating

pheromone in a microfluidic chamber for 12hrs. Images of cells at 0min (top row),

360min (middle-row), and 720min (bottom row) are displayed with split-channel (left and

middle columns) and merged images (right column). Arrows show sites of septin

organization coincident with the polar cap. Scale bar represents 5µm. (D)

Representative images of $gpa1^{G302S-UD\Delta}$ yeast expressing endogenous Bem1-GFP

(Polar Cap) and Cdc3-mCherry (Septin) exposed to saturating pheromone in a

microfluidic chamber for 12hrs. Images of cells at 0min (top row), 360min (middle row),

and 720min (bottom row) are displayed with split-channel (left and middle columns) and

merged images (right column). Scale bar represents 5µm. (E) Time-averaged

quantification of the fraction (y-axis) of septin distribution relative to the polar cap (x-

axis). Lines are smoothed representations of time-averaged data (37 time points; 20min

intervals for 12hrs) and shaded regions surrounding lines are 95% confidence intervals.

Lines are representative of two combined experiments ($gpa1^{G302S}$-$ste2^{\Delta C}$ n = 58 cells

and $gpa1^{G302S\text{-}\Delta UD}$ n = 93 cells).

Figure 9. Diagram of proposed mechanism. A) Proposed molecular mechanism for

three scenarios: 1) The Gpa1 independent pathway, which would use Gics and Cdc42-

GAPs to promote septin organization, similar to mitotic processes. 2) The standard

Gpa1-regulated pathway, where we propose Gpa1 recruits epsins and Cdc42-GAPs to

sites of high Gpa1 activity, leading to septin organization. and 3) The Hyperactive Gpa1

scenario, where Gpa1 activity is high close to the center of the polar cap, leading to

faster recruitment of endocytic machinery, and therefore septin organization closer to

the center of the polar cap than normal. Consistent with our mathematical modeling,

proteins that stay on the membrane longer also travel further from the polar cap, thus

faster recruitment of endocytic adaptors leads to less distance traveled. The diagram

fades to the left to indicate that this is a symmetric process that we are showing only

one side of. B) Proposed mechanism for the asymmetric distribution of septins during

gradient tracking. Note that delivery of vesicles to the center of the polarity site and

removal distal to the polarity site creates a "treadmill" effect where plasma membrane

moves away from the center and towards the sites of endocytosis. On the left, when the

polar cap (where exocytosis is occurring) is oriented towards the source of pheromone,

the amount of signal to either side of the polar cap is equal, Gpa1 activation levels are

similar, leading to similar amounts of endocytosis, and symmetric septin organization.

On the right, the polarity site (where exocytosis is occurring) is not oriented towards the

source of pheromone, instead high pheromone to the right of the polarity site is driving

increased endocytosis relative to the opposite side of the polarity site, leading to

asymmetric distribution of septins.

Reference List

Aguilar, R.C., S.A. Longhi, J.D. Shaw, L.-Y. Yeh, S. Kim, A. Schön, E. Freire, A. Hsu,
W.K. McCormick, and H.A. Watson. 2006. Epsin N-terminal homology domains
perform an essential function regulating Cdc42 through binding Cdc42 GTPase-
activating proteins. *Proceedings of the National Academy of Sciences*. 103:4116-
4121.

Alvaro, C.G., and J. Thorner. 2016. Heterotrimeric G Protein-coupled Receptor
Signaling in Yeast Mating Pheromone Response. *The Journal of biological*
chemistry. 291:7788-7795.

Apanovitch, D.M., K.C. Slep, P.B. Sigler, and H.G. Dohlman. 1998. Sst2 is a GTPase-
activating protein for Gpa1: purification and characterization of a cognate RGS-
Galpha protein pair in yeast. *Biochemistry*. 37:4815-4822.

Azimi, M., Y. Jamali, and M.R. Mofrad. 2011. Accounting for diffusion in agent based
models of reaction-diffusion systems with application to cytoskeletal diffusion.
PLoS One. 6:e25306.

Bennet, M.A., P.R. Richardson, J. Arlt, A. McCarthy, G.S. Buller, and A.C. Jones. 2011.
Optically trapped microsensors for microfluidic temperature measurement by
fluorescence lifetime imaging microscopy. *Lab on a Chip*. 11:3821-3828.

Bharucha, J.P., J.R. Larson, J.B. Konopka, and K. Tatchell. 2008. Saccharomyces
cerevisiae Afr1 protein is a protein phosphatase 1/Glc7-targeting subunit that
regulates the septin cytoskeleton during mating. *Eukaryot Cell*. 7:1246-1255.

Brennwald, P. 2013. Membrane traffic: the exocyst meets the cell cycle. *Current biology*
: CB. 23:R838-840.

Bridges, A.A., and A.S. Gladfelter. 2015. Septin Form and Function at the Cell Cortex.
The Journal of biological chemistry. 290:17173-17180.

Burke, D., D. Dawson, and T. Stearns. 2000. Methods in yeast genetics: a Cold Spring
Harbor Laboratory course manual (2000 edition). *Plainview, NY: Cold Spring*
Harbor Laboratory Press.[Google Scholar].

Carrillo, L., B. Cucu, V. Bandmann, U. Homann, B. Hertel, S. Hillmer, G. Thiel, and A.
Bertl. 2015. High-resolution membrane capacitance measurements for studying
endocytosis and exocytosis in yeast. *Traffic*. 16:760-772.

Changwei, Z., X. Mingyong, and W. Ranran. 2007. Afr1p has a role in regulating the
localization of Mpk1p at the shmoo tip inSaccharomyces cerevisiae. *FEBS*
Letters. 581:2670-2674.

Chee, M.K., and S.B. Haase. 2012. New and redesigned pRS plasmid shuttle vectors
for genetic manipulation of Saccharomyces cerevisiae. *G3: Genes| Genomes|*
Genetics. 2:515-526.

Chen, H., S. Fre, V.I. Slepnev, M.R. Capua, K. Takei, M.H. Butler, P.P. Di Fiore, and P.
De Camilli. 1998. Epsin is an EH-domain-binding protein implicated in clathrin-
mediated endocytosis. *Nature*. 394:793-797.

Coon, B.G., J. Burgner, J.H. Camonis, and R.C. Aguilar. 2010. The epsin family of
endocytic adaptors promotes fibrosarcoma migration and invasion. *J Biol Chem*.
285:33073-33081.

de Godoy, L.M., J.V. Olsen, J. Cox, M.L. Nielsen, N.C. Hubner, F. Frohlich, T.C.
Walther, and M. Mann. 2008. Comprehensive mass-spectrometry-based
proteome quantification of haploid versus diploid yeast. *Nature*. 455:1251-1254.

DiBello, P.R., T.R. Garrison, D.M. Apanovitch, G. Hoffman, D.J. Shuey, K. Mason, M.I.
Cockett, and H.G. Dohlman. 1998. Selective uncoupling of RGS action by a
single point mutation in the G protein alpha-subunit. *The Journal of biological*
chemistry. 273:5780-5784.

Dixit, G., R. Baker, C.M. Sacks, M.P. Torres, and H.G. Dohlman. 2014a. Guanine
Nucleotide-binding Protein (Gα) Endocytosis by a Cascade of Ubiquitin Binding
Domain Proteins Is Required for Sustained Morphogenesis and Proper Mating in
Yeast. *Journal of Biological Chemistry*. 289:15052-15063.

Dixit, G., J.B. Kelley, J.R. Houser, T.C. Elston, and H.G. Dohlman. 2014b. Cellular noise
suppression by the regulator of G protein signaling Sst2. *Molecular cell*. 55:85-
96.

Dohlman, H.G., J. Song, D. Ma, W.E. Courchesne, and J. Thorner. 1996. Sst2, a
negative regulator of pheromone signaling in the yeast Saccharomyces
cerevisiae: expression, localization, and genetic interaction and physical
association with Gpa1 (the G-protein alpha subunit). *Molecular and cellular*
biology. 16:5194-5209.

Dores, M.R., J.D. Schnell, L. Maldonado-Baez, B. Wendland, and L. Hicke. 2010. The
function of yeast epsin and Ede1 ubiquitin-binding domains during receptor
internalization. *Traffic*. 11:151-160.

Dunn, R., and L. Hicke. 2001. Domains of the Rsp5 ubiquitin-protein ligase required for
receptor-mediated and fluid-phase endocytosis. *Mol Biol Cell*. 12:421-435.

Giot, L., and J. Konopka. 1997. Functional analysis of the interaction between Afr1p and
the Cdc12p septin, two proteins involved in pheromone-induced morphogenesis.
Mol Biol Cell. 8:987-998.

Gladfelter, A.S., I. Bose, T.R. Zyla, E.S. Bardes, and D.J. Lew. 2002. Septin ring
assembly involves cycles of GTP loading and hydrolysis by Cdc42p. *J Cell Biol*.
156:315-326.

Gladfelter, A.S., L. Kozubowski, T.R. Zyla, and D.J. Lew. 2005. Interplay between septin
organization, cell cycle and cell shape in yeast. *J Cell Sci*. 118:1617-1628.

Hicke, L., and H. Riezman. 1996. Ubiquitination of a yeast plasma membrane receptor
signals its ligand-stimulated endocytosis. *Cell*. 84:277-287.

Hicke, L., B. Zanolari, and H. Riezman. 1998. Cytoplasmic tail phosphorylation of the
alpha-factor receptor is required for its ubiquitination and internalization. *The*
Journal of cell biology. 141:349-358.

Huh, W.-K., J.V. Falvo, L.C. Gerke, A.S. Carroll, R.W. Howson, J.S. Weissman, and
E.K. O'Shea. 2003. Global analysis of protein localization in budding yeast.
Nature. 425:686-691.

Iwase, M., J. Luo, S. Nagaraj, M. Longtine, H.B. Kim, B.K. Haarer, C. Caruso, Z. Tong,
J.R. Pringle, and E. Bi. 2006. Role of a Cdc42p effector pathway in recruitment of
the yeast septins to the presumptive bud site. *Mol Biol Cell.* 17:1110-1125.

Kelley, J.B., G. Dixit, J.B. Sheetz, S.P. Venkatapurapu, T.C. Elston, and H.G. Dohlman.
2015. RGS proteins and septins cooperate to promote chemotropism by
regulating polar cap mobility. *Current Biology.* 25:275-285.

Khan, A., M. McQuilken, and A.S. Gladfelter. 2015. Septins and Generation of
Asymmetries in Fungal Cells. *Annu Rev Microbiol.* 69:487-503.

Konopka, J.B., D.D. Jenness, and L.H. Hartwell. 1988. The C-terminus of the S.
cerevisiae α-pheromone receptor mediates an adaptive response to pheromone.
Cell. 54:609-620.

Layton, A.T., N.S. Savage, A.S. Howell, S.Y. Carroll, D.G. Drubin, and D.J. Lew. 2011.
Modeling vesicle traffic reveals unexpected consequences for Cdc42p-mediated
polarity establishment. *Current biology : CB.* 21:184-194.

Longtine, M., H. Fares, and J. Pringle. 1998. Role of the yeast Gin4p protein kinase in
septin assembly and the relationship between septin assembly and septin
function. *The Journal of cell biology.* 143:719-736.

Lu, R., D.G. Drubin, and Y. Sun. 2016. Clathrin-mediated endocytosis in budding yeast
at a glance. *J Cell Sci.* 129:1531-1536.

Marco, E., R. Wedlich-Soldner, R. Li, S.J. Altschuler, and L.F. Wu. 2007. Endocytosis
optimizes the dynamic localization of membrane proteins that regulate cortical
polarity. *Cell.* 129:411-422.

Matheos, D., M. Metodiev, E. Muller, D. Stone, and M.D. Rose. 2004. Pheromone-
induced polarization is dependent on the Fus3p MAPK acting through the formin
Bni1p. *The Journal of cell biology.* 165:99-109.

McMurray, M., A. Bertin, G. rd, L. Lam, E. Nogales, and J. Thorner. 2011. Septin
filament formation is essential in budding yeast. *Developmental cell.* 20:540-549.

Metodiev, M.V., D. Matheos, M.D. Rose, and D.E. Stone. 2002. Regulation of MAPK
function by direct interaction with the mating-specific Gα in yeast. *Science.*
296:1483-1486.

Mukherjee, D., B.G. Coon, D.F. Edwards, C.B. Hanna, S.A. Longhi, J.M. McCaffery, B.
Wendland, L.A. Retegui, E. Bi, and R.C. Aguilar. 2009. The yeast endocytic
protein Epsin 2 functions in a cell-division signaling pathway. *Journal of Cell
Science.* 122:2453-2463.

Okada, S., M. Leda, J. Hanna, N.S. Savage, E. Bi, and A.B. Goryachev. 2013. Daughter
cell identity emerges from the interplay of Cdc42, septins, and exocytosis. *Dev
Cell.* 26:148-161.

Pedersen, R.T.A., J.E. Hassinger, P. Marchando, and D.G. Drubin. 2020. Spatial
regulation of clathrin-mediated endocytosis through position-dependent site
maturation. *J Cell Biol*. 219.

Pryciak, P.M., and F.A. Huntress. 1998. Membrane recruitment of the kinase cascade
scaffold protein Ste5 by the Gbetagamma complex underlies activation of the
yeast pheromone response pathway. *Genes Dev*. 12:2684-2697.

Sadian, Y., C. Gatsogiannis, C. Patasi, O. Hofnagel, R.S. Goody, M. Farkašovský, and
S. Raunser. 2013. The role of Cdc42 and Gic1 in the regulation of septin filament
formation and dissociation. *Elife*. 2:e01085.

Segall, J.E. 1993. Polarization of yeast cells in spatial gradients of alpha mating factor.
Proceedings of the National Academy of Sciences. 90:8332-8336.

Shenoy, S.K., and R.J. Lefkowitz. 2003. Multifaceted roles of beta-arrestins in the
regulation of seven-membrane-spanning receptor trafficking and signalling.
Biochem J. 375:503-515.

Simke, W.C., C.P. Johnson, A.J. Hart, S. Mayhue, P.L. Craig, S. Sojka, and J.B. Kelley.
2022. Phosphorylation of RGS regulates MAP kinase localization and promotes
completion of cytokinesis. *Life Sci Alliance*. 5.

Slessareva, J., and H. Dohlman. 2006. G protein signaling in yeast: new components,
new connections, new compartments. *Science (New York, N.Y.)*. 314:1412-1413.

Smith, G.R., S.A. Givan, P. Cullen, and G.F. Sprague, Jr. 2002. GTPase-activating
proteins for Cdc42. *Eukaryot Cell*. 1:469-480.

Smith, J.S., T.F. Pack, A. Inoue, C. Lee, K. Zheng, I. Choi, D.S. Eiger, A. Warman, X.
Xiong, Z. Ma, G. Viswanathan, I.M. Levitan, L.K. Rochelle, D.P. Staus, J.C.
Snyder, A.W. Kahsai, M.G. Caron, and S. Rajagopal. 2021. Noncanonical
scaffolding of G(alphai) and beta-arrestin by G protein-coupled receptors.
Science. 371.

Spiliotis, E.T., and M.A. McMurray. 2020. Masters of asymmetry - lessons and
perspectives from 50 years of septins. *Mol Biol Cell*. 31:2289-2297.

Suzuki, S.K., J.B. Kelley, T.C. Elston, and H.G. Dohlman. 2021. Gradient tracking by
yeast GPCRs in a microfluidics chamber. *In* G Protein-Coupled Receptor
Screening Assays. Springer. 275-287.

Valdez-Taubas, J., and H. Pelham. 2003. Slow diffusion of proteins in the yeast plasma
membrane allows polarity to be maintained by endocytic cycling. *Current biology
: CB*. 13:1636-1640.

Wang, Y., L.A. Marotti, Jr., M.J. Lee, and H.G. Dohlman. 2005a. Differential Regulation
of G Protein α Subunit Trafficking by Mono- and Polyubiquitination *.
Journal of Biological Chemistry. 280:284-291.

Wang, Y., L.A. Marotti, Jr., M.J. Lee, and H.G. Dohlman. 2005b. Differential regulation
of G protein alpha subunit trafficking by mono- and polyubiquitination. *The
Journal of biological chemistry*. 280:284-291.

Wendland, B. 2002. Epsins: adaptors in endocytosis? *Nat Rev Mol Cell Biol*. 3:971-977.

Figure 1

Figure 2

Figure 3

A

B Single Time Point at 200x

gpa1^{G302S} rga1Δ

gpa1^{G302S} rga2Δ

gpa1^{G302S} bem3Δ

C *gpa1^{G302S} rga1Δ*

D *gpa1^{G302S} rga2Δ*

E *gpa1^{G302S} bem3Δ*

F Septin Localization

Figure 4

Figure 5

Stochastic Agent-Based Model of Protein Trafficking

Figure 6

Cargo Simulations

A)

All Cargo

1 X Phosphorylation   5 X Phosphorylation   25 X Phosphorylation

B)

Ubiquitinated Cargo

1 X Phosphorylation   5 X Phosphorylation   25 X Phosphorylation

C)

Effect of Phosphorylation Rate on Location of Endocytosis

1 X Phosphorylation   5 X Phosphorylation   25 X Phosphorylation

Figure 7

A) Simulations of high and low endocytosis competency rate are able to capture the shape of septin distributions measured in WT and hyperactive Gpa1 at intermediate doses of pheromone

B) Model simulations of an intermediate endocytosis competency rate are able to predict the septin distribution in a high dose of pheromone with a wild-type Gpa1.

Figure 8

Figure 9

A

Time & Space

Protein travel distance is related to time on membrane

B

Proposed mechanism for asymmetric septin organization

Table 1: Strain List

Strain	Parent	Description
BEM1-GFP CDC3MCHERRY	BY4741	BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
RGA1Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	rga1Δ::KanMX4 Gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
RGA2Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	rga2Δ::KanMX4 Gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
BEM3Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	bem3Δ::KanMX4 Gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
ENT1Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	ent1Δ::KanMX4 gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2
ENT2Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	ent2Δ::KanMX4 gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2

GIC1Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	*gic1Δ::KanMX4 Gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2*
GIC2Δ GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	*gic2Δ::KanMX4 Gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY:LEU2*
Ste2^{T326} GPA1^{G302S} BEM1-GFP CDC3MCHERRY	BY4741	*ste2^{T326}::KanMX4 gpa1^{G302S}::URA3 BEM1-GFP::HIS3 CDC3MCHERRY::LEU2*
GPA1$^{G302S-ΔUD}$ BEM1-GFP CDC3MCHERRY	BY4741	*gpa1$^{G302S-ΔUD}$::URA3 BEM1-GFP::HIS3 CDC3MCHERRY:LEU2*
ENT1-GFP	BY4741	*ENT1-GFP::HIS3*
ENT2-GFP	BY4741	*ENT2-GFP::HIS3*

Table 2: Plasmid List

Plasmid Name	Vector Backbone	Description
pRSII416-Gpa1-3xFLAG	pRSII416	CEN ADHp-Gpa1-3xFLAG::LEU2 vector
pRSII406-GPA1^{G302S}	pRSII406	Integrating gpa1^{G302S}::URA3 vector
pRSII406-GPA1$^{G302S\text{-}\Delta UD}$	pRSII406	Integrating gpa1$^{G302S\text{-}\Delta UD}$::URA3 vector
pRSII416-GAL-Ste5CTM	pRSII416	Galactose-inducible Ste5-CTM

Table 3: Primer List

Primer Name	Sequence	Description
AHM-14	5'GGGCCGTATAAAGACTACAGACATTACAGAAACCGAATTTA 3'	Quickchange primer for G302S mutation in Gpa1
AHM-15	5' TAAATTCGGTTTCTGTAATGTCTGTAGTCTTTATAC GGCCC 3'	Quickchange primer for G302S mutation in Gpa1
AHM-44	5' GCGGCCGTTTTATAAAGAACAA 3'	Replicate deletion of Bem3
AHM-45	5' AGAAGTGGTGGAAGATGGTGAT 3'	Replicate deletion of Bem3
AHM-46	5' TTACAGCTTTGCCACATATCGG 3'	Verification of Bem3 del
AHM-47	5' AACAAAACTTTCAATCGCACGG 3'	Verification of Bem3 del
AHM-50	5' GATGAGAAGGTGTAAAGTGCCC 3'	Replicate deletion of Rga1
AHM-51	5' GCCCGTAAAAGGTTCTTGTCTA 3'	Replicate deletion of Rga1
AHM-52	5' AGTGTACCCTTTGAACCCTCTT 3'	Verify Replication of Rga1 del
AHM-53	5' ATCTCTGTCGCTCAAAAGTTGG 3'	Verify Replication of Rga1 del
AHM-54	5' AATCGTTGCATCGCTTCAGTAA 3'	Replicate deletion of Rga2
AHM-55	5' CCGCACAAAAGTACGACAAATG 3'	Replicate deletion of Rga2
AHM-56	5' CTCTGTAGAAGCTTTTCCGGAC 3'	Verify Replication of Rga2 del
AHM-57	5' CTTTACAGAAGTCGCGTCAACT 3'	Verify Replication of Rga2 del

CJM-3	5' CAATGTGGGCCACGGCTGCTAA 3'	Verify Ste2 C-terminal del
CJM-4	5' ACAGCGTACCTTTAGACACGTGGG 3'	Verify Ste2 C-terminal del
CJM-5	5' CATCCACAGATAGGTTTTATCCAGGCACGCT GTCTAGCGTTTAGCTTGCCTCGTCCCCGC 3'	Amplify pFA6-KanMX with Ste2 T-326 homology
CJM-6	5' AGGTCACGAAATTACTTTTTCAAAGCCGTAA ATTTTGACGTTTAAACTGGATGGCGGCGT 3'	Amplify pFA6-KanMX with Ste2 T-326 homology
SLM-3	5' TAGGCTTCTTACCGGCAGAG 3'	Amplify Ent1 del
SLM-4	5' AGGACATGAGTACAGAGCACA 3'	Amplify Ent1 del
SLM-5	5' AGATAGATGGCCGAACGTGG 3'	Amplify Ent2 del
SLM-6	5' GACTCCAAAGGTGAATCTGGC3'	Amplify Ent2 del
SLM-7	5' GGGAGAAGGAGTACCTTCTG 3'	Verify Ent1 del
SLM-8	5' CCAGAATTATCTTTCGGGCC 3'	Verify Ent1 del
SLM-9	5' ATGGCTCAAATTCCACGGTCC 3'	Verify Ent2 del
SLM-10	5' CGAAGATGGCCGAATGATTGG 3'	Verify Ent2 del
JKM-54	5' GACTACAAAGACCATGACGGTGATTATAAAGAT CATGACATCGACTACAAGGATGACGATGACAAGTAG 3'	Forward 3X Flag + stop
JKM-55	5' CTACTTGTCATCGTCATCCTTGTAGTCGATGT CATGATCTTTATAATCACCGTCATGGTCTTTGTAGTC 3'	Reverse 3X Flag + stop
JKM-56	5' tcgaggtcgacggtatcgatAAGGTGAGACGCGCATAAC 3'	Forward ADH promoter for Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag
JKM-57	5' tacaccccatTGTATATGAGATAGTTGA	reverse ADH promoter for

	TTGTATGCTTG 3'	Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag
JKM-58	5'ctcatatacaATGGGGTGTACAGTGAGTAC 3'	forward GPA1 for Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag
JKM-59	5' ctttgtagtcTATAATACCAATTTTTTTAAG GTTTTGCTG 3'	reverse GPA1 for Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag
JKM-60	5' tggtattataGACTACAAAGACCATGAC 3'	forward 3x Flag for Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag, uses JKM-54 and 55 as template
JKM-61	5' gcggccgctctagaactagtCTACTTGTCATCGTCATC 3'	reverse 3x Flag for Gibson assembly of pRSII416-ADHp-Gpa1-3XFlag, uses JKM-54 and 55 as template