**Neuronal modeling of magnetoencephalography responses in auditory cortex to**

**auditory and visual stimuli**

*Kaisu Lankinen[1,2], Jyrki Ahveninen[1,2], Mainak Jas[1,2], Tommi Raij[1,2], Seppo P. Ahlfors[1,2]*

*[1] Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital,*

*Charlestown, MA 02129*

*[2] Department of Radiology, Harvard Medical School, Boston, MA 02115*

Corresponding author: Kaisu Lankinen, klankinen@mgh.harvard.edu

Abbreviated title: Modeling cross-sensory MEG responses

Number of pages: 38

Number of figures: 7

Number of tables: 1

Number of words for abstract: 230

Number of words for introduction: 571

Number of words for discussion: 1517

Conflict of interest: The authors declare no competing financial interests.

Acknowledgments: Supported by R01DC016765, R01DC016915, R01DC017991,

P41EB030006, P41EB015896. We thank Dr. Stephanie Jones for useful discussions.

**ABSTRACT**

Previous studies have demonstrated that auditory cortex activity can be influenced by cross-

sensory visual inputs. Intracortical recordings in non-human primates (NHP) have suggested a

bottom-up feedforward (FF) type laminar profile for auditory evoked but top-down feedback (FB)

type for cross-sensory visual evoked activity in the auditory cortex. To test whether this principle

applies also to humans, we analyzed magnetoencephalography (MEG) responses from eight

human subjects (six females) evoked by simple auditory or visual stimuli. In the estimated MEG

source waveforms for auditory cortex region of interest, auditory evoked responses showed peaks

at 37 and 90 ms and cross-sensory visual responses at 125 ms. The inputs to the auditory cortex

were then modeled through FF and FB type connections targeting different cortical layers using

the Human Neocortical Neurosolver (HNN), which consists of a neocortical circuit model linking

the cellular- and circuit-level mechanisms to MEG. The HNN models suggested that the measured

auditory response could be explained by an FF input followed by an FB input, and the cross-

sensory visual response by an FB input. Thus, the combined MEG and HNN results support the

hypothesis that cross-sensory visual input in the auditory cortex is of FB type. The results also

illustrate how the dynamic patterns of the estimated MEG/EEG source activity can provide

information about the characteristics of the input into a cortical area in terms of the hierarchical

organization among areas.

**SIGNIFICANCE STATEMENT**

Laminar intracortical profiles of activity characterize feedforward- and feedback-type influences in

the inputs to a cortical area. By combining magnetoencephalography (MEG) and biophysical

computational neural modeling, we obtained evidence of cross-sensory visual evoked activity in

human auditory cortex being of feedback type. The finding is consistent with previous intracortical

recordings in non-human primates. The results illustrate how patterns of MEG source activity can

be interpreted in the context of the hierarchical organization among cortical areas.

## INTRODUCTION

Activity in sensory cortices is influenced by feedforward (FF) and feedback (FB) connections

between cortical layers and brain regions, following a hierarchical organization (Rockland and

Pandya, 1979; Felleman and Van Essen, 1991; Zeki, 2018). In the auditory cortex of non-human

primates (NHPs), the laminar profile of early auditory evoked responses has FF type

characteristics, whereas cross-sensory visual or somatosensory evoked activity are of FB type

(for reviews see, e.g., Foxe and Schroeder, 2005; Schroeder and Foxe, 2005; Ghazanfar and

Schroeder, 2006; Kayser and Logothetis, 2007). Human magneto- and electroencephalography

(MEG/EEG) studies have revealed that cross-sensory activations and multisensory interactions

can occur in low-order sensory areas very early, within a few tens of milliseconds from the

stimulus onset (Giard and Peronnet, 1999; Foxe et al., 2000; Molholm et al., 2002; Teder-Sälejärvi

et al., 2002; Molholm et al., 2004; Lakatos et al., 2007; Talsma et al., 2007; Raij et al., 2010). In

line with evidence from studies in other cognitive domains (Polimeni et al., 2010; Muckli et al.,

2015; Kok et al., 2016; Fracasso et al., 2018; Klein et al., 2018; Finn et al., 2019; Lawrence et al.,

2019a; Norris and Polimeni, 2019), recent high-field fMRI studies have provided evidence of FF-

and FB-like intracortical depth profiles in auditory cortex BOLD signals (De Martino et al., 2015;

Ahveninen et al., 2016; Moerel et al., 2018; Wu et al., 2018; Moerel et al., 2019; Gau et al., 2020;

Chai et al., 2021; Lankinen et al., 2022). However, detailed neurophysiological analysis or

computational modeling of such effects has not been done in humans.

Previous studies have suggested that early components of evoked responses are related to

FF processes, whereas later components reflect FB influences in activity evoked by auditory (Inui

et al., 2006; Kohl et al., 2022), visual (Aine et al., 2003; Inui and Kakigi, 2006), and somatosensory

(Cauller and Kulics, 1991; Inui et al., 2004; Jones et al., 2007) stimuli.

Biophysically realistic computational models have been used to investigate laminar

connections and cellular and circuit level processes of the neurons in detail, and they can also be

used to simulate MEG/EEG signals (Jones et al., 2007; Neymotin et al., 2020). The Human

Neocortical Neurosolver (HNN) (Neymotin et al., 2020) provides a cortical column model with FF-

and FB-type inputs targeting different layers. With HNN, the cellular and network contributions to

MEG/EEG signals from a source-localized region of interest can be modeled and compared to

the measured signals. Previously, HNN has been used to interpret mechanisms of sensory

evoked responses and oscillations in healthy and clinical populations (Jones et al., 2007; Jones

et al., 2009; Ziegler et al., 2010; Lee and Jones, 2013; Khan et al., 2015; Sherman et al., 2016;

Pinotsis et al., 2017; Sliva et al., 2018; Bonaiuto et al., 2021; Kohl et al., 2022; Law et al., 2022).

Kohl et al. (2022) showed that auditory responses in the auditory cortex could be modeled by

activating the neocortical circuit through a layer-specific sequence of FF-FB-FF inputs, similar to

a prior simulation of somatosensory evoked responses (Jones et al., 2007).

In the present study, we investigated auditory vs. cross-sensory visual evoked responses in

the auditory cortex by comparing the measured MEG responses with simulated source waveforms

from a computational model (HNN). We hypothesized that the auditory evoked responses

observed with MEG can be explained by a sequence of FF and FB influences, whereas FB-type

input is adequate to explain the cross-sensory visual evoked response.

## MATERIAL AND METHODS

### Subjects

Eight healthy right-handed subjects participated (six females, age 22–30 years). All subjects gave written informed consent, and the study protocol was approved by the Massachusetts General Hospital institutional review board and followed the guidelines of the Declaration of Helsinki.

### Stimuli and task

The subjects were presented with *Noise/Checkerboard* and *Letter* stimuli in separate runs while MEG was recorded. Data for the *Noise/Checkerboard* stimuli were used in our earlier publication (Raij et al., 2010). Here we re-analyzed data from the *Noise/Checkerboard* experiment, together with the previously unpublished data from the *Letter* experiment. Equiprobable 300-ms auditory, visual, and audiovisual (simultaneous auditory and visual) stimuli were delivered in an event-related design with pseudorandom order. The auditory *Noise* stimuli were white noise bursts (15 ms rise and decay) and the visual *Checkerboard* stimuli static checkerboard patterns (visual angle 3.5°×3.5° and contrast 100%, with a peripheral fixation crosshair). The *Letter* stimuli were spoken and written letters of Roman alphabet ('A', 'B', 'C', etc.). The subjects' task was to respond to rare (10%) auditory, visual, or audiovisual target stimuli with the right index finger movement as quickly as possible. In the *Noise/Checkerboard* experiment, the target stimulus was a tone pip, a checkerboard with a gray diamond pattern in the middle, or a combination of the two. In the *Letter* task, the target stimulus was the letter 'K', spoken and/or written. Data were recorded in three runs with different stimulus onset asynchrony (SOA, mean 1.5, 3.1, or 6.1 s, all jittered at 1.15 s). There were 375 stimuli per category (auditory, visual, and audiovisual): 150 in the short, 125 in the intermediate, and 100 in the long SOA runs. All subjects were presented with the same order of tasks and stimuli. The auditory stimuli were presented with MEG-compatible headphones, with the intensity adjusted to be as high as the subject could comfortably listen to. The visual stimuli

were projected onto a translucent screen. The stimuli were controlled using Presentation 9.20

(Neurobehavioral Systems Inc, Albany, CA, USA).

***MEG and MRI acquisition and co-registration***

MEG was recorded with a 306-channel instrument with 204 planar gradiometer and 102

magnetometer sensors (VectorView; MEGIN, Finland) inside a magnetically shielded room

(Cohen et al., 2002). Simultaneous horizontal and vertical electro-oculograms (EOG) were also

recorded. All signals were bandpass-filtered to 0.03–200 Hz and sampled at 600 Hz.

Structural T1-weighted MRIs of the subjects were acquired with a 1.5 T Siemens Avanto

scanner (Siemens Medical Solutions, Erlangen, Germany) and a head coil using a standard

MPRAGE sequence. Cortical surfaces were reconstructed using the FreeSurfer software

(http://www.surfer.nmr.mgh.harvard.edu, (Fischl, 2012).

Prior to the MEG recording, the locations of four small head position indicator coils attached to

the scalp and several additional scalp surface points were determined with respect to the fiducial

landmarks (nasion and two preauricular points) using a 3-D digitizer (Fastrak Polhemus, VT,

USA). For the MRI–MEG coordinate system alignment, the fiduciary points were first identified

from the structural MRIs, and then this initial co-registration was refined using an iterative closest-

point search algorithm for the scalp surface locations using the MNE Suite software (Gramfort et

al., 2014, http://www.martinos.org/mne/).

***MEG preprocessing and source estimation***

The MEG data were analyzed using MNE-Python (Gramfort et al., 2013). After excluding channels

and time segments with excessive noise, independent component analysis (ICA) was used to

identify and remove artifacts related to eye blinks, eye movements, and cardiac activity. The

signals were then lowpass filtered at 40 Hz, and event-related responses were averaged

separately for the auditory and visual trials, combining the long, intermediate, and short SOA runs.

After exclusion of artifactual time segments an average of 369.9 (std 6.5) epochs per subject

remained in response to auditory, and 370.2 (std 5.1) to visual stimulation. In the present study

we did not analyze the audiovisual or target trials. The zero level in each channel was defined as

the mean signal over the 200-ms prestimulus baseline period.

Source activity was estimated at 4098 discrete locations per hemisphere on the cortical

surface, with an average separation of the source elements being about 4.9 mm. For the forward

solution, a single-compartment boundary element model was used. Forward solutions were first

computed separately for the three runs with different SOAs and then averaged (Uutela et al.,

2001). Minimum-norm estimates (MNE, (Hamalainen and Ilmoniemi, 1994)) for the cortical source

currents were calculated. Both the gradiometer and the magnetometer channels were included in

the source estimation. We used fixed source orientation normal to the cortical surface and depth

weighting 0.8 to reduce bias towards superficial currents. For region-of-interest (ROI) selection,

the MNE values were noise-normalized to obtain dynamic statistical parametric maps (dSPM;

Dale et al., 2000).

***Regions-of-interest and source time courses***

Auditory evoked potentials and magnetic fields typically have three main deflections: P50-N100-

P200 (or P50m-N100m-P200m for MEG), peaking approximately at 50, 100 and 180 ms,

respectively, after the auditory stimulus onset (Picton et al., 1974; Hari et al., 1980; Hämäläinen

et al., 1993; Jones et al., 2007; Ahlfors et al., 2015). The ROIs were determined based on the

auditory N100m response, because the SNR of the visual evoked response over the auditory

cortex was too low to reliably determine auditory cortex ROIs from the visual evoked data in the

presence of partially coinciding strong occipital visual cortex activity. We identified functional ROIs

for the auditory cortex in each hemisphere, separately for each subject, based on the N100m

peak of the auditory evoked response. First, anatomically defined regions were selected using

the Destrieux atlas parcellation from Freesurfer (Fischl et al., 2004; Destrieux et al., 2010):

Heschl's gyrus, Heschl's sulcus, and the lower part of planum temporale (masked with

supramarginal gyrus) were combined to cover the primary auditory areas. Then, from these

regions the source element with the largest negative deflection between 60–110 ms (except for

manually set 105 ms in one subject) in the dSPM source time course was identified. Using that

source element as a seed point, all source elements that had a magnitude of 30% or more of the

peak dSPM value and formed a continuous area around the seed point were selected. The

average number of selected elements across subjects, hemispheres and experiments for the

auditory cortex ROIs was 19 (standard deviation 8.7, range 3–38). The same procedure was used

to determine also additional control ROIs in the occipital cortex (V1, V2, and MT based on the

FreeSurfer atlas (Fischl et al., 2008). The source waveform for an ROI was defined as the sum of

the MNE time courses over those selected source elements. Note that the magnitude of the

response depended on the number of the vertices that were included in the ROI, and thus was

expected to give a smaller amplitude than would be found by the use of a single equivalent current

dipole to represent the auditory cortex activity (as used, *e.g.*, by Kohl et al. (2022)). Although

equivalent current dipoles are in general well suited to describe auditory evoked responses, here

it was more convenient to use a distributed source model (MNE) for wide-spread visual evoked

response, to extract cross-sensory responses in the auditory cortex.

***Neural modeling with Human Neocortical Neurosolver (HNN)***

Activity in the auditory cortex evoked by the auditory and visual stimuli was modeled using HNN

(https://jonescompneurolab.github.io/hnn-core/) (Neymotin et al., 2020). HNN is a software for

simulating neocortical circuits and linking cellular- and circuit-level physiology to the electrical

source currents measured by MEG and EEG. Thus, HNN provides a tool to develop and test

hypotheses on the neural origins of MEG/EEG signals. The neural currents contributing to the

MEG/EEG signals from a source region are modeled in terms of the local network dynamics driven

by layer-specific inputs (see **Fig. 1**). Simulated MEG/EEG source currents are represented as

current dipole waveforms calculated from the distribution of intracellular currents in the dendrites

of the pyramidal cells. MEG/EEG signals originate mostly from postsynaptic currents in cortical

pyramidal neurons (Hämäläinen et al., 1993; Okada et al., 1997), and the magnitude and direction

of the source current depends on the type of the synaptic input and its dendritic location (Allison

et al., 2002; Jones et al., 2007; Linden et al., 2010; Lopes da Silva, 2010; Ahlfors et al., 2015;

Ahlfors and Wreh, 2015), providing a link between the laminar distribution of synaptic inputs and

the MEG/EEG source waveforms.

***Figure 1.*** *Schematic illustration of the HNN model. (A) A network of neurons in a local cortical*

*area generates an evoked response. (B) Local network structure with pyramidal cells (blue) and*

*interneurons (orange). Excitatory and inhibitory coupling is indicated by a black circle and bar,*

*respectively. The network is activated by proximal (red) and distal (green) drives by input spike*

*trains. Modified from Neymotin et al. (2020).*

In HNN, the model for a local cortical circuit has a layered structure with pyramidal neurons

whose somata are in the supragranular (layer 2/3) or infragranular (layer 5) layers and whose

dendrites span across the layers. The model also includes inhibitory interneurons. External input

to the circuit arrives through characteristic layer-specific FF and FB type connections. FF type

inputs consist of proximal drives to the basal dendrites of the pyramidal cells (assumed to arrive

via the middle cortical layer), whereas FB inputs are represented by distal drive to the apical

dendrites of the pyramidal cells. The model has 100 pyramidal neurons in each of layers 2/3 and

5; a scaling factor is used to match the simulated dipole to the magnitude of the recorded evoked

response. The parameters of the HNN model originate from known anatomical and physiological

cell properties, and the local connectivity within and between cortical layers is based on a large

body of literature from animal studies (Jones et al., 2007; Neymotin et al., 2020).

We used HNN to test the hypothesis that the differences in the MEG responses to auditory

and visual stimuli can be explained by a different sequence of FF and FB inputs to the auditory

cortex. This hypothesis is based on neurophysiological evidence from animal studies (Schroeder

and Foxe, 2002). Underlying mechanisms of auditory responses in humans have been previously

described using HNN (Kohl et al., 2022). Our specific hypothesis was that the auditory response

can be explained by an initial FF input followed by an FB input, but the visual response just by an

FB input.

We created two main HNN models for event-related activity in the auditory cortex: one for the

response to auditory stimuli and one for the response to visual stimuli.  The grand average MEG

source waveforms (averaged across subjects, hemispheres, and experiments) were modeled

using HNN. As a starting point, we used the auditory cortex model by Kohl et al. (2022) for activity

in the right hemisphere evoked by auditory stimuli presented to the left ear. Because HNN has a

large number of user-defined parameters, we made the following assumptions to limit the

parameter space: a) Only the timing parameters of the FF/FB spike-train inputs (mean $\mu$ and

standard deviation σ of a Gaussian distribution) were adjusted, in addition to an overall scaling

factor for the simulated source waveforms; all the other parameters were kept unchanged. b)

These other, internal, model parameters were assumed to be the same for the responses to visual

and auditory stimuli. c) The simulations were limited to the time window of 0–150 ms for the

auditory and 0–170 ms for the visual response, in order to focus on the early part of the responses.

HNN model parameters were determined by minimizing the root mean square error (RMSE)

between the simulated and experimentally observed MEG source waveforms. To improve the

SNR of the experimental data, we averaged MEG source waveforms over subjects, hemispheres,

and the two experiments. The simulated HNN waveforms were smoothed in the default 30-ms

window (Hamming window convolution).

We first manually adjusted the start time of the FF/FB inputs and scaling of the response to

achieve a close initial fit to the MEG responses. An optimal scaling factor was determined by

minimizing the RMSE between the average of 10 simulation runs and the MEG waveform over

the specified time windows. Thereafter, we further tuned the model parameters using Bayesian

optimization          implemented          in          scikit-optimize          (Head,          2020)

(http://doi.org/10.5281/zenodo.1207017) for estimating $\mu$ (mean input spike timing) and $\sigma$

(temporal distribution of input spikes) for each model by minimizing the RMSE between the

simulated and the measured signal. We used "expected improvement" as the acquisition function.

The initial parameters were defined from the manual fit and the bounds for the search space were

($\mu_{FF}$: 20…50, $\mu_{FB}$: 55…95, $\mu_{FF2}$: 90…130, $\sigma_{FF}$: 1…5, $\sigma_{FB}$: 5…20, $\sigma_{FF2}$: 5…20).

As HNN has a large number of parameters, it is possible that even after optimizing our main

models, some other combination of parameter values could explain the waveforms equally well

or better. Therefore, we formed alternative models by varying the number and timing of the FF

and FB inputs. We focused on the comparison of FF + FB *vs.* FB models for explaining the early

part of the MEG activity evoked by auditory and visual stimuli.

***Statistical analyses***

To evaluate whether the magnitudes of the estimated MEG source waveforms (averaged

across tasks and hemispheres) were significantly different from zero, we used *t*-tests with a

threshold $p < 0.05$ in each of the 150 time points in the 0–250 ms window. The *p*-values were

Bonferroni adjusted for the two stimulus types and 150 time points. To evaluate between-subject

consistency of the magnitudes of the largest defections in the evoked responses in each

hemisphere and experiment, the average value over time points within ±10 ms windows around

the peak latencies were calculated for each subject and submitted to *t*-tests with False discovery

rate (*fdr)* adjustment for 12 tests.

For the HNN models, a non-parametric resampling approach was used to test whether the

alternative models could provide a significantly better fit than our main models. First, the MEG

source waveforms for auditory and visual evoked responses were resampled by drawing from 32

signals (8 subjects x 2 hemispheres x 2 experiments) 500 times with replacement. The same was

repeated for 32 simulation runs for each of the models (FF + FB and FB). Next, the root-mean-

square error (RMSE) between each of the 500 resampled MEG signals and 500 resampled

simulations for each model was calculated, resulting in histograms of RMSE values within each

model. We tested whether the difference between the simulated source waveforms from the FB

*vs.* FF + FB models was significantly different from 0. The RMSE difference histograms were

normalized for each model between -1 and 1, as the ranges in the auditory and visual models

were different. To create a null-distribution, the signs of the waveforms were randomly flipped

10,000 times, an average of 500 resamplings was calculated. To assign a *p*-value for each model,

the mean RMSE value was compared with the null distribution, with the Bonferroni adjustment of

$n = 2$ (auditory and visual models). If the difference of the models (FF + FB vs. FB) was significant,

we concluded that including the first FF was necessary for the model.

**RESULTS**

***MEG source waveforms in auditory cortex in response to auditory and visual stimuli***

Estimated MEG source waveforms for auditory and visual evoked activity in the auditory cortex

ROIs, averaged over subjects, tasks, and hemispheres, are shown in **Fig. 2**. The auditory evoked

response showed a characteristic biphasic P50m-N100m waveform, with a positive peak at 37

and a negative peak at 90 ms after the onset of the auditory stimuli. These peak latencies are

similar to those reported previously for auditory noise burst stimuli (Hari et al., 1987). The cross-

sensory visual evoked response in the auditory cortex had a monophasic peak at 125 ms after

the appearance of the visual stimuli. The source magnitudes at the peak latencies were

significantly different from zero (*t*-test, $p < 0.05$, Bonferroni adjusted). The magnitude of the visual

evoked response was about 13% of the magnitude of the auditory N100m. The direction of the

source current for the  visual response was the same as that of the auditory N100m response,

pointing from the gray matter towards the white matter.

*Figure 2. MEG source activity in the auditory cortex. The estimated source waveforms in response to the auditory (orange) and visual (blue) stimuli (mean and standard deviation across subjects, hemispheres, and experiments). Negative values correspond to inward cortical currents, i.e., pointing from the gray matter towards the white matter. The gray shading indicates time points that differing significantly from zero (t-test, p < 0.05, Bonferroni adjusted).*

We examined the reproducibility of the estimated source waveforms across the experiments, hemispheres, and individual subjects. MEG source waveforms in the left and the right hemispheres in response to the *Noise/Checkerboard* and *Letter* stimuli  are illustrated in **Fig. 3**. The magnitude of the auditory N100m was larger for the *Letter* than for the *Noise* stimuli in the left hemisphere, but similar in the right hemisphere; this lateralization is expected for responses to phonetic vs. non-verbal stimuli (Gootjes et al., 1999; Parviainen et al., 2005). The anatomical overlap of ROIs across subjects (**Fig. 3**, middle panel)  suggested that the prominent auditory

evoked responses originated mostly in the Heschl's sulcus and the anterior part of the planum temporale. There were no clear differences in the location of the ROIs between the *Noise/Checkerboard* and *Letter* experiments; however, for the *Letter* stimuli, the location extended to the Heschl's gyrus in half of the subjects. The peak latencies of the auditory evoked responses were similar within a few milliseconds in both experiments. For the visual evoked response, a negative deflection with the peak latency ranging from 113 to 132 ms was seen in both experiments in both hemispheres.

***Figure 3.*** *MEG source waveforms in the left and right hemisphere auditory cortex in response to auditory and visual stimulation, shown separately for the Noise/Checkerboard and Letter experiments. The source waveforms were averaged over subjects. The locations of the functional ROIs morphed to common anatomical space ('fsaverage' from FreeSurfer) are shown in the middle; the color bar indicates how many subjects' individual ROIs overlapped at each cortical location. The black lines illustrate the Heschl's gyrus (anterior), Heschl's sulcus (middle) and part of planum temporale (posterior).*

To evaluate between-subject consistency of the largest defections in the evoked responses in each hemisphere in each experiment, we calculated for each subject the average value over time

points within ±10 ms windows around the peak latencies (black dots in **Fig. 4**). The auditory

N100m peak was statistically significant in all cases (*Noise*: left hemisphere $p$ = 0.027, right $p$ =

0.0045; *Letter*: left $p$ = 0.027, right $p$ = 0.027; $t$-test, False discovery rate (*fdr*) adjusted). For the

response to the visual stimuli, the negative peak was statistically significant in the right

hemisphere (*Checkerboard*: $p$ = 0.040; *Letter*: $p$ = 0.027) but not in the left hemisphere

(*Checkerboard*: $p$ = 0.19; *Letter*: $p$ = 0.). The auditory P50m peaks were not significant when

calculated separately for the different cases, but they were significant for the grand average

responses (see **Fig. 2**).

*Figure 4.* Variation of the estimated source waveforms among individual subjects. The p-values

indicate the significance of the response magnitudes at the peak latencies (t-test; fdr adjusted).

Continuous lines and shading: mean ± standard deviation across subjects; black dots: response

magnitudes for individual subjects, calculated as the average over ±10 ms time windows around

the peak latencies. LH: left hemisphere, RH: right hemisphere. * p < 0.05, ** p < 0.01.

The observed weak visual evoked activity in the auditory cortex partially coincided with strong

activity in occipital visual cortical regions (**Fig. 5**). The estimated auditory cortex source

waveforms could potentially reflect artefactual spread in the MEG source estimates due to activity

in other cortical regions responding to the visual stimuli. We examined this possibility in two ways.

First, we observed that the time course of the estimated sources for visual cortex ROIs had

prominent deflections for both the onset (with peak latencies at ~100 ms) and the offset (~400

ms) of the visual stimuli, whereas in the auditory cortex the response was seen mainly for the

onset only (**Fig. 5A**). If the onset and offset responses share a common spatial distribution in the

occipital cortex, then also the potential artefactual spreading to the auditory cortex is expected to

be the similar after the onset and the offset of the visual stimuli. However, this was not found in

the data. Second, the spatial maps of the source estimates for the visual evoked responses have

a gap between the weak auditory cortex activity and the large occipital cortex activity (**Fig. 5B)**.

Artificial spread would be expected to be spatially uniform rather than forming separate foci in the

auditory cortex. These observations argue against the possibility of the cross-sensory visual

evoked response in the auditory cortex to be  artefactually resulting from spread from visual cortex

in the source estimates.

**Figure 5.** *Evaluation of potential artefactual spatial spread in the estimated MEG source activity from visual cortex to the auditory ROIs. (A) Source time-courses (MNE, averaged across subjects and tasks) in response to visual stimuli for occipital areas V1, V2, MT (green) and the auditory cortices (V AC, blue). (B) Spatial maps of the MNE source estimate for the visual evoked activity at the time of the largest peak in the response to visual stimuli in the auditory cortex.*

### Neural modeling with HNN

The initial manual tuning values for the mean (and standard deviation) of the time distribution of the inputs were $\mu_{FF}$ = 35 ($\sigma_{FF}$ = 3.0) ms for the FF and $\mu_{FB}$ = 75 ($\sigma_{FB}$ = 13.3) ms for the FB input in the auditory model, and $\mu_{FB}$ = 105 ($\sigma_{FB}$ = 13.3) ms for the FB input in the visual model. The optimal scaling factor was found to be 53 for the auditory and 5 for the visual simulation. Fine-tuning with Bayesian hyperparameter optimization resulted in only small adjustments to the timing parameters. The optimized values were $\mu_{FF}$= 34 ($\sigma_{FF}$ = 1.0), $\mu_{FB}$= 74 ($\sigma_{FB}$ = 14.0) in the auditory

model, and $\mu_{FB}$ = 105 ($\sigma_{FB}$ = 17.5) in the visual model (**Table 1**). The temporal distributions of

the inputs are depicted in **Fig. 6B**. For both the auditory responses (P50m-N100m) and the visual

responses (peaking at 125 ms), the simulated source waveforms captured the main features of

the experimentally observed MEG results **(Fig. 6A).**

***Table 1.*** *Comparison of HNN parameters for auditory and visual models. The mean μ and*

*standard deviation σ (milliseconds) describe the temporal distribution of the inputs. Scaling factor*

*is used to match the simulated dipole to the measured evoked response waveform. RMSE is root-*

*mean-square error calculated between simulated and measured waveform. The main models are*

*highlighted.*

| Model | $\mu_{FF}$ ($\sigma_{FF}$) | $\mu_{FB}$ ($\sigma_{FB}$) | $\mu_{FF2}$ ($\sigma_{FF2}$) | Scaling | RMSE |
|---|---|---|---|---|---|
| A: FF + FB + FF (Kohl et al., 2022) | 47 (3.0) | 81 (13.3) | 151 (11.1) | 1500 | |
| A: FF + FB + FF | 35 (1.0) | 77 (15.0) | 90 (14.4) | 57 | 0.15 |
| A: FF + FB | 34 (1.0) | 74 (14.0) | - | 53 | 0.072 |
| A: FB | - | 78 (14.8) | - | 38 | 0.23 |
| A: FF | 20 (3) | - | - | 1 | 0.86 |
| V: FF + FB | 42 (1.0) | 99 (12.7) | - | 6 | 0.012 |
| V: FB | - | 105 (17.5) | - | 5 | 0.024 |
| V: FF | 1 (3.0) | - | - | 1 | 0.095 |

**Figure 6.** HNN simulations of the auditory cortex activity in response to auditory (left) and visual

(right) stimuli. **A:** Simulated source waveforms using the initial manual adjustments to the model

parameters (dashed gray lines), after parameter optimization (thick gray: average, thin gray: 10

Further insights to the generation of the source currents can be obtained by plotting separately

the contributions from layer-2/3 and in layer-5 pyramidal cells (**Fig. 6C**) and the sequences of the

spiking activity of the four cell types included in the HNN model (**Fig. 6D**). In the model for the

auditory evoked response, FF input was assumed to arrive to the auditory cortex through the

middle cortical layer and the excite the basal dendrites of the pyramidal cells in both layers 2/3

and 5 **(Fig. 6C**, left). The net result of the FF input was an initial upward (positive) peak. The

arrival of the FB input to the distal parts of the apical dendrites of the pyramidal cells resulted in

reversal of the net current to be downwards. In the model for the cross-sensory visual evoked

response, the FB input arriving distally drove the net source current downwards within the apical

dendrites of both layer 2/3 and layer 5 pyramidal cells **(Fig. 6C**, right)**.**

As HNN has a large number of parameters, it is possible that our chosen models are not the

only ones that can reproduce the experimentally observed MEG source waveforms. However,

HNN can serve us as a valuable hypothesis testing tool to test different models. Alternative

models with different combinations of FF and FB inputs are shown in **Fig. 7**, and the

corresponding optimized HNN parameters for these are listed in **Table 1**.

**Figure 7.** *Alternative models for auditory (**A**) and visual (**B**) responses. The main models (A: FF + FB and V: FB) are framed. The experimentally observed MEG source waveforms (orange: auditory stimulus, blue: visual stimulus) are overlayed with the simulated waveforms (thin gray: 10 individual simulation runs, thick gray: average of the individual runs. Histograms below the waveforms show the temporal distribution of FF (red) and FB (green) inputs to the HNN model of the auditory cortex neural circuit. FF only simulations are scaled to illustrate their waveforms compared with the MEG signal.*

For the auditory evoked responses, inclusion of a later second FF input to the model had only little effect on the simulated source waveforms within 0–150 ms (A: FF+FB+FF2 *vs.* A: FF+FB, **Fig. 7A**). Removing the first FF input, however, resulted in a notable difference in the early time window (30–80 ms), during which the first upward deflection was seen in the MEG data.

Interestingly, if the FB input was removed, the FF input alone could not produce response

waveforms similar to those observed empirically. As the optimal scaling factor for the FF only

model was 1, **Fig. 7** (right column) shows the model scaled up in order to illustrate how the

waveform looks like compared with the MEG response. Thus, the FB input seems to have an

essential role in the generation of the evoked responses studied here.

For the visual evoked response, the difference between models with and without an FF input

(V: FF+FB *vs.* V: FB) was most pronounced in the early part (30–80 ms) of the simulated source

waveforms **(Fig. 7B)**. However, although the V: FF+FB model slightly improved the fit to the

measured MEG signal in comparison with V: FB, considering the magnitude of the response with

the baseline noise level (see **Fig. 2**) suggests that the additional FF input in the model for the

response to the visual stimuli may be mostly explaining just noise in the data. Using a non-

parametric resampling approach, a significant difference between FF+FB *vs.* FB was found for

the auditory models ($p < 0.001$) but not for the visual models ($p = 0.39$). In other words, early FF

input did not significantly improve the model fit to the response to visual stimuli. Thus, these

results support our main hypothesis that the response to the auditory stimuli results from a

combination of FF and FB inputs to the auditory cortex, whereas the cross-sensory visual

response can be explained with just FB input to the auditory cortex.

**DISCUSSION**

The MEG data revealed a cross-sensory event-related response in the auditory cortex, peaking

at about 125 ms after the appearance of the visual stimuli. The direction of the estimated source

current for this response was the same as for the auditory N100m response, pointing from the

cortical gray matter towards the white matter. The main shape of the visual evoked response

waveform could be reproduced by an HNN model with FB-type input, whereas for the biphasic

P50m-N100m auditory evoked response both FF and FB inputs were needed. The experimental

and modeling results are consistent with the hypothesis that cross-sensory visual input to the

auditory cortex is of FB type (Schroeder and Foxe, 2002).

*Characterization of cross-sensory visual evoked activation in auditory cortex*

Recently, Kohl et al. presented an HNN model with a sequence of FF and FB inputs explaining

several properties of auditory evoked responses in the auditory cortex (Kohl et al., 2022). With

only minor adjustments to the input timings and the overall scaling, the model could be adapted

to explain the MEG source waveforms for the auditory evoked responses observed in the present

study. A sequence of FF-FB (and -FF) inputs has been shown to model well also somatosensory

responses in the somatosensory cortex (Jones et al., 2007). In contrast, to explain the early part

of the cross-sensory visual response in the auditory cortex, we found that a model with only an

FB input, without a preceding FF input, was adequate. The FB-type characteristics is consistent

with previous NHP electrophysiological studies (Schroeder and Foxe, 2002). Multi-contact

electrode recordings in the macaque have shown early activity in the granular (middle) layer of

auditory cortex in response to auditory stimuli, suggesting FF-type input, whereas cross-sensory

visual evoked activity appeared first in supra- and infragranular layers (Schroeder and Foxe,

2002). Similar laminar properties in the auditory cortex have also been seen in human fMRI

studies  (Gau et al., 2020; Chai et al., 2021; Lankinen et al., 2022). In the high-field laminar fMRI

study of Lankinen et al. (2022), which used the same stimuli as in the *Noise/Checkerboard*

experiment in the present MEG study, BOLD signal depth profiles in the auditory cortex showed

different curvature for auditory vs. visual stimuli, consistent with the hypothesized difference in

the FF vs. FB type inputs.

There are several possible neural pathways for the visual evoked activity to reach the auditory

cortex. The relatively long latency of the visual response observed here is consistent with what

would be expected from input from higher-order polysensory areas such as the superior temporal

sulcus (Foxe and Schroeder, 2005). However, the present analyses focusing on activity within

auditory cortex only do not reveal the origin of the inputs to the auditory cortex. That type of

information could be deduced, e.g., from Granger-causality measures between estimated source

waveforms in multiple cortical areas (Milde et al., 2011; Gow and Nied, 2014; Michalareas et al.,

2016).

Interestingly, NHP studies have shown different characteristics for visual and somatosensory

cross-sensory inputs to the auditory cortex: FB-type for visual but FF-type for somatosensory

(Schroeder and Foxe, 2002). The role of different types of cross-sensory inputs to the auditory

cortex may have important implications to theories of multisensory processing (Schroeder and

Foxe, 2005). There appear to be multiple ways how cross-sensory processes may be influenced

by the hierarchical organization among brain areas. FB-type inputs are commonly associated with

modulatory influences, whereas FF-type inputs are more directly related to sensory information

(Schroeder and Foxe, 2005).

***Complementary approaches to noninvasive detection of FF and FB processes***

The present approach of combining MEG and cellular-level computational modeling complements

other non-invasive methods for studying the organization of cortical processes in the human brain.

The millisecond-scale time resolution of MEG and EEG enables the investigation of fast dynamics

of the brain activity, which is not attainable with hemodynamic fMRI. High-field fMRI, however,

can provide laminar-level spatial resolution for making inferences about FF and FB activity (see

e.g., De Martino et al., 2018; Lawrence et al., 2019b; Norris and Polimeni, 2019). With certain

strong assumptions about the location and extent of the spatial distribution, layer-specific source

localization in MEG has also been demonstrated (Bonaiuto et al., 2018a; Bonaiuto et al., 2018b).

FF/FB influences can also be inferred from directed connectivity measures for MEG source

estimates at specific frequency bands (Michalareas et al., 2016).

The present results also support the view that the direction of MEG source waveforms can be

useful for inferring information about the hierarchical organization of cortical processing (Ahlfors

et al., 2015). In particular, FF-type input to the supragranular layer, with excitatory synaptic

connections to the distal part of the apical dendrites of pyramidal cells, is likely to be a major

contributor to the downward-directed MEG source currents (Lopes da Silva, 2010; Ahlfors and

Wreh, 2015). There was a general correspondence between the source direction and the type of

input in the HNN model: the outward directed source current during the auditory P50m response

was associated with FF input in HNN, whereas FB inputs were needed to model the inward source

currents during the auditory N100m and the visual response peaking at 125 ms. A close

relationship between the direction of MEG source currents and FF- vs. FB-type inputs has also

been found in HNN modeling of somatosensory response in the primary somatosensory cortex

(Jones et al., 2007). Furthermore, the direction of the MEG source currents in inferior

occipitotemporal cortex has been found to reverse between two experimental conditions for which

a cognitive neuroscience theory for visual object recognition predicted FF vs. FB inputs to the

area (Ahlfors et al., 2015).

***Limitations of the current study***

Localizing weak cross-sensory visual evoked activity in the auditory cortex is challenging because

of potential interference in the MEG source estimate from the partially coinciding occipital cortex

activity. However, both the shape of the time courses and the patterns in the spatial distributions

of the source estimates (see **Fig. 6**) suggested that it was unlikely that the visual evoked activity

in the auditory cortex was due to artefactual long-range crosstalk caused by spatial spread in the source estimates. Short-range spread in the source estimates can also confound the interpretation of the source waveforms. If the true location of the visual responses were not within the auditory cortex ROI, but, e.g., in the opposite side of the superior temporal gyrus, the source direction could become incorrectly identified. Combining MEG with high-resolution fMRI could help to confirm the location of the activity. It is also possible that there was simultaneous activity in multiple auditory areas in the supratemporal plane. Most of the individual subjects' ROIs were located directly at the primary auditory regions, at or near the at Heschl's sulcus, being thus slightly different than the auditory association area just posterior to primary auditory region studied by Schroeder and Foxe (2002). However, it has been shown in monkeys that such FF type patterns are typical throughout the core and belt regions of auditory cortex (Schroeder et al. 2001). Without further data, e.g., intracranial recordings, it is difficult to conclusively resolve the locations of the sources of the observed cross-sensory MEG response.

HNN, and biophysical computational neural modeling in general, has two challenges of opposite nature: the neural circuit model is complex, with a large number of adjustable parameters, and yet the model is a simplified representation of the cortical circuitry. We used neural circuit parameters of the pre-tuned model for auditory evoked responses in the auditory cortex by Kohl et al. (2022) and only adjusted a small number of selected parameters, focusing on the timing of the FF and FB inputs. Given the limited SNR of the experimental source waveforms, we did not attempt to vary the neural connectivity parameters. We cannot exclude the possibility that there could be some combinations within the high-dimensional parameter space that could explain the responses with a very different circuit model than the one reported here. Useful in future studies, it has been recently demonstrated that combining simulation-based inference (SBI) to HNN modeling can help in parameter estimation (Tolley et al., 2023).

We modeled only one local region (auditory cortex) receiving one-directional external inputs.

To determine where the inputs are arriving from and where the information will be sent, directional

connectivity analyses between multiple regions would be needed. Thus, further studies would be

necessary to connect other areas of interest to the network. Furthermore, combining MEG with

layer-specific fMRI could provide complementary information which could help to build a more

detailed picture of the FF/FB influences.

***Conclusions***

The combined MEG and HNN modeling results support the hypothesis that cross-sensory visual

input to the auditory cortex is of FB type. The results also illustrate how the dynamic patterns of

the estimated MEG/EEG source activity can provide information about the characteristics of the

input into the cortical areas in terms of hierarchical organization among the cortical areas.

Avenues for future research could include connecting other areas of interest to the network,

calculating directed (effective) connectivity measures between cortical areas specifically, and

combining complementary information from MEG data with layer-specific fMRI to build a more

detailed picture of the FF/FB influences.

**References**

Ahlfors SP, Wreh C, 2nd (2015) Modeling the effect of dendritic input location on MEG and EEG

source dipoles. Med Biol Eng Comput 53:879-887.

Ahlfors SP, Jones SR, Ahveninen J, Hämäläinen MS, Belliveau JW, Bar M (2015) Direction of

magnetoencephalography  sources  associated  with  feedback  and  feedforward

contributions in a visual object recognition task. Neurosci Lett 585:149-154.

Ahveninen J, Chang WT, Huang S, Keil B, Kopco N, Rossi S, Bonmassar G, Witzel T, Polimeni

JR (2016) Intracortical depth analyses of frequency-sensitive regions of human auditory

cortex using 7TfMRI. Neuroimage 143:116-127.

Aine CJ, Stephen JM, Christner R, Hudson D, Best E (2003) Task relevance enhances early

transient and late slow-wave activity of distributed cortical sources. J Comput Neurosci

15:203-221.

Allison T, Puce A, McCarthy G (2002) Category-sensitive excitatory and inhibitory processes in

human extrastriate cortex. J Neurophysiol 88:2864-2868.

Bonaiuto JJ, Little S, Neymotin SA, Jones SR, Barnes GR, Bestmann S (2021) Laminar dynamics

of high amplitude beta bursts in human motor cortex. Neuroimage 242:118479.

Bonaiuto JJ, Meyer SS, Little S, Rossiter H, Callaghan MF, Dick F, Barnes GR, Bestmann S

(2018a) Lamina-specific cortical dynamics in human visual and sensorimotor cortices.

Elife 7.

Bonaiuto JJ, Rossiter HE, Meyer SS, Adams N, Little S, Callaghan MF, Dick F, Bestmann S,

Barnes GR (2018b) Non-invasive laminar inference with MEG: Comparison of methods

and source inversion algorithms. Neuroimage 167:372-383.

Cauller LJ, Kulics AT (1991) The neural basis of the behaviorally relevant N1 component of the

somatosensory-evoked potential in SI cortex of awake monkeys: evidence that backward

cortical projections signal conscious touch sensation. Exp Brain Res 84:607-619.

Chai Y, Liu TT, Marrett S, Li L, Khojandi A, Handwerker DA, Alink A, Muckli L, Bandettini PA

(2021) Topographical and laminar distribution of audiovisual processing within human

planum temporale. Prog Neurobiol 205:102121.

Dale AM, Liu AK, Fischl BR, Buckner RL, Belliveau JW, Lewine JD, Halgren E (2000) Dynamic

statistical parametric mapping: combining fMRI and MEG for high-resolution imaging of

cortical activity. Neuron 26:55-67.

De Martino F, Moerel M, Xu J, van de Moortele PF, Ugurbil K, Goebel R, Yacoub E, Formisano

E (2015) High-Resolution Mapping of Myeloarchitecture In Vivo: Localization of Auditory

Areas in the Human Brain. Cereb Cortex 25:3394-3405.

De Martino F, Yacoub E, Kemper V, Moerel M, Uludag K, De Weerd P, Ugurbil K, Goebel R,

Formisano E (2018) The impact of ultra-high field MRI on cognitive and computational

neuroimaging. Neuroimage 168:366-382.

Destrieux C, Fischl B, Dale A, Halgren E (2010) Automatic parcellation of human cortical gyri and

sulci using standard anatomical nomenclature. Neuroimage 53:1-15.

Felleman DJ, Van Essen DC (1991) Distributed hierarchical processing in the primate cerebral

cortex. Cereb Cortex 1:1-47.

Finn ES, Huber L, Jangraw DC, Molfese PJ, Bandettini PA (2019) Layer-dependent activity in

human prefrontal cortex during working memory. Nat Neurosci 22:1687-1695.

Fischl B (2012) FreeSurfer. Neuroimage 62:774-781.

Fischl B, Rajendran N, Busa E, Augustinack J, Hinds O, Yeo BT, Mohlberg H, Amunts K, Zilles K

(2008) Cortical folding patterns and predicting cytoarchitecture. Cereb Cortex 18:1973-

1980.

Fischl B, van der Kouwe A, Destrieux C, Halgren E, Segonne F, Salat DH, Busa E, Seidman LJ,

Goldstein J, Kennedy D, Caviness V, Makris N, Rosen B, Dale AM (2004) Automatically

parcellating the human cerebral cortex. Cereb Cortex 14:11-22.

Foxe J, Morocz I, Murray M, Higgins B, Javitt D, Schroeder C (2000) Multisensory auditory-

somatosensory interactions in early cortical processing revealed by high-density electrical

mapping. Cognitive Brain Research 10:77-83.

Foxe JJ, Schroeder CE (2005) The case for feedforward multisensory convergence during early

cortical processing. Neuroreport 16:419-423.

Fracasso A, Luijten PR, Dumoulin SO, Petridou N (2018) Laminar imaging of positive and

negative BOLD in human visual cortex at 7T. Neuroimage 164:100-111.

Gau R, Bazin PL, Trampel R, Turner R, Noppeney U (2020) Resolving multisensory and

attentional influences across cortical depth in sensory cortices. Elife 9.

Ghazanfar AA, Schroeder CE (2006) Is neocortex essentially multisensory? Trends Cogn Sci

10:278-285.

Giard M, Peronnet F (1999) Auditory-visual integration during multimodal object recognition in

humans: a behavioral and electrophysiological study. Journal of Cognitive Neuroscience

11:473-490.

Gootjes L, Raij T, Salmelin R, Hari R (1999) Left-hemisphere dominance for processing of vowels:

a whole-scalp neuromagnetic study. Neuroreport 10:2987-2991.

Gow DW, Jr., Nied AC (2014) Rules from words: a dynamic neural basis for a lawful linguistic

process. PLoS One 9:e86212.

Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Parkkonen L,

Hamalainen MS (2014) MNE software for processing MEG and EEG data. Neuroimage

86:446-460.

Gramfort A, Luessi M, Larson E, Engemann DA, Strohmeier D, Brodbeck C, Goj R, Jas M, Brooks

T, Parkkonen L, Hamalainen M (2013) MEG and EEG data analysis with MNE-Python.

Front Neurosci 7:267.

Hämäläinen M, Hari R, Ilmoniemi R, Knuutila J, Lounasmaa O (1993) Magnetoencephalography

626            -theory, instrumentation, and applications to noninvasive studies of the working human

627            brain. Rev Mod Phys 65:413-497.

628     Hamalainen MS, Ilmoniemi RJ (1994) Interpreting magnetic fields of the brain: minimum norm

629            estimates. Med Biol Eng Comput 32:35-42.

630     Hari R, Aittoniemi K, Jarvinen ML, Katila T, Varpula T (1980) Auditory evoked transient and

631            sustained magnetic fields of the human brain. Localization of neural generators. Exp Brain

632            Res 40:237-240.

633     Hari R, Pelizzone M, Makela JP, Hallstrom J, Leinonen L, Lounasmaa OV (1987) Neuromagnetic

634            responses of the human auditory cortex to on- and offsets of noise bursts. Audiology

635            26:31-43.

636     Head TK, Manor; Nahrstaedt, Holger; Louppe, Gilles; Shcherbatyi, Iaroslav; (2020) scikit-

637            optimize. In.

638     Inui K, Kakigi R (2006) Temporal analysis of the flow from V1 to the extrastriate cortex in humans.

639            J Neurophysiol 96:775-784.

640     Inui K, Wang X, Tamura Y, Kaneoke Y, Kakigi R (2004) Serial processing in the human

641            somatosensory system. Cereb Cortex 14:851-857.

642     Inui K, Okamoto H, Miki K, Gunji A, Kakigi R (2006) Serial and parallel processing in the human

643            auditory cortex: a magnetoencephalographic study. Cereb Cortex 16:18-30.

644     Jones SR, Pritchett DL, Stufflebeam SM, Hamalainen M, Moore CI (2007) Neural correlates of

645            tactile detection: a combined magnetoencephalography and biophysically based

646            computational modeling study. J Neurosci 27:10751-10764.

647     Jones SR, Pritchett DL, Sikora MA, Stufflebeam SM, Hamalainen M, Moore CI (2009) Quantitative

648            analysis and biophysically realistic neural modeling of the MEG mu rhythm:

649            rhythmogenesis and modulation of sensory-evoked responses. J Neurophysiol 102:3554-

650            3572.

Kayser C, Logothetis NK (2007) Do early sensory cortices integrate cross-modal information?

Brain Struct Funct 212:121-132.

Khan S, Michmizos K, Tommerdahl M, Ganesan S, Kitzbichler MG, Zetino M, Garel KL, Herbert

MR, Hamalainen MS, Kenet T (2015) Somatosensory cortex functional connectivity

abnormalities in autism show opposite trends, depending on direction and spatial scale.

Brain 138:1394-1409.

Klein BP, Fracasso A, van Dijk JA, Paffen CLE, Te Pas SF, Dumoulin SO (2018) Cortical depth

dependent population receptive field attraction by spatial attention in human V1.

Neuroimage 176:301-312.

Kohl C, Parviainen T, Jones SR (2022) Neural Mechanisms Underlying Human Auditory Evoked

Responses Revealed By Human Neocortical Neurosolver. Brain Topogr 35:19-35.

Kok P, Bains LJ, van Mourik T, Norris DG, de Lange FP (2016) Selective Activation of the Deep

Layers of the Human Primary Visual Cortex by Top-Down Feedback. Curr Biol 26:371-

376.

Lakatos P, Chen CM, O'Connell MN, Mills A, Schroeder CE (2007) Neuronal oscillations and

multisensory interaction in primary auditory cortex. Neuron 53:279-292.

Lankinen K, Ahlfors SP, Mamashli F, Blazejewska AI, Raij T, Turpin T, Polimeni JR, Ahveninen J

(2022) Cortical depth profiles of auditory and visual 7 T functional MRI responses in human

superior temporal areas. Hum Brain Mapp.

Law RG, Pugliese S, Shin H, Sliva DD, Lee S, Neymotin S, Moore C, Jones SR (2022)

Thalamocortical Mechanisms Regulating the Relationship between Transient Beta Events

and Human Tactile Perception. Cereb Cortex 32:668-688.

Lawrence SJ, Norris DG, de Lange FP (2019a) Dissociable laminar profiles of concurrent bottom-

up and top-down modulation in the human visual cortex. Elife 8.

Lawrence SJD, Formisano E, Muckli L, de Lange FP (2019b) Laminar fMRI: Applications for

cognitive neuroscience. Neuroimage 197:785-791.

Lee S, Jones SR (2013) Distinguishing mechanisms of gamma frequency oscillations in human

current source signals using a computational model of a laminar neocortical network. Front

Hum Neurosci 7:869.

Linden H, Pettersen KH, Einevoll GT (2010) Intrinsic dendritic filtering gives low-pass power

spectra of local field potentials. J Comput Neurosci 29:423-444.

Lopes da Silva FH (2010) Electrophysiological basis of MEG signals. In: MEG: An introduction to

methods (Hansen PC, Kringelbach, M.L., Salmelin, R., ed), pp 1-23. New York, NewYork:

Oxford University Press.

Michalareas G, Vezoli J, van Pelt S, Schoffelen JM, Kennedy H, Fries P (2016) Alpha-Beta and

Gamma Rhythms Subserve Feedback and Feedforward Influences among Human Visual

Cortical Areas. Neuron 89:384-397.

Milde T, Putsche P, Schwab K, Wacker M, Eiselt M, Witte H (2011) Dynamics of directed

interactions between brain regions during interburst-burst EEG patterns in quiet sleep of

full-term neonates. Neurosci Lett 488:148-153.

Moerel M, De Martino F, Ugurbil K, Formisano E, Yacoub E (2018) Evaluating the Columnar

Stability of Acoustic Processing in the Human Auditory Cortex. J Neurosci 38:7822-7832.

Moerel M, De Martino F, Ugurbil K, Yacoub E, Formisano E (2019) Processing complexity

increases in superficial layers of human primary auditory cortex. Sci Rep 9:5502.

Molholm S, Ritter W, Javitt DC, Foxe JJ (2004) Multisensory visual-auditory object recognition in

humans: a high-density electrical mapping study. Cereb Cortex 14:452-465.

Molholm S, Ritter W, Murray M, Javitt D, Schroeder C, Foxe J (2002) Multisensory auditory-visual

interactions during early sensory processing in humans: a high-density electrical mapping

study. Cognitive Brain Research 14:115-128.

Muckli L, De Martino F, Vizioli L, Petro LS, Smith FW, Ugurbil K, Goebel R, Yacoub E (2015)

Contextual Feedback to Superficial Layers of V1. Curr Biol 25:2690-2695.

Neymotin SA, Daniels DS, Caldwell B, McDougal RA, Carnevale NT, Jas M, Moore CI, Hines ML,

Hamalainen M, Jones SR (2020) Human Neocortical Neurosolver (HNN), a new software

tool for interpreting the cellular and network origin of human MEG/EEG data. Elife 9.

Norris DG, Polimeni JR (2019) Laminar (f)MRI: A short history and future prospects. Neuroimage

197:643-649.

Okada YC, Wu J, Kyuhou S (1997) Genesis of MEG signals in a mammalian CNS structure.

Electroencephalogr Clin Neurophysiol 103:474-485.

Parviainen T, Helenius P, Salmelin R (2005) Cortical differentiation of speech and nonspeech

sounds at 100 ms: implications for dyslexia. Cereb Cortex 15:1054-1063.

Picton TW, Hillyard SA, Krausz HI, Galambos R (1974) Human auditory evoked potentials. I.

Evaluation of components. Electroencephalogr Clin Neurophysiol 36:179-190.

Pinotsis DA, Geerts JP, Pinto L, FitzGerald THB, Litvak V, Auksztulewicz R, Friston KJ (2017)

Linking canonical microcircuits and neuronal activity: Dynamic causal modelling of laminar

recordings. Neuroimage 146:355-366.

Polimeni JR, Fischl B, Greve DN, Wald LL (2010) Laminar analysis of 7T BOLD using an imposed

spatial activation pattern in human V1. Neuroimage 52:1334-1346.

Raij T, Ahveninen J, Lin FH, Witzel T, Jaaskelainen IP, Letham B, Israeli E, Sahyoun C, Vasios

C, Stufflebeam S, Hamalainen M, Belliveau JW (2010) Onset timing of cross-sensory

activations and multisensory interactions in auditory and visual sensory cortices. Eur J

Neurosci 31:1772-1782.

Rockland KS, Pandya DN (1979) Laminar origins and terminations of cortical connections of the

occipital lobe in the rhesus monkey. Brain Res 179:3-20.

Schroeder CE, Foxe JJ (2002) The timing and laminar profile of converging inputs to multisensory

areas of the macaque neocortex. Brain Res Cogn Brain Res 14:187-198.

Schroeder CE, Foxe J (2005) Multisensory contributions to low-level, 'unisensory' processing.

Curr Opin Neurobiol 15:454-458.

Sherman MA, Lee S, Law R, Haegens S, Thorn CA, Hamalainen MS, Moore CI, Jones SR (2016)

Neural mechanisms of transient neocortical beta rhythms: Converging evidence from

humans, computational modeling, monkeys, and mice. Proc Natl Acad Sci U S A

113:E4885-4894.

Sliva DD, Black CJ, Bowary P, Agrawal U, Santoyo JF, Philip NS, Greenberg BD, Moore CI,

Jones SR (2018) A Prospective Study of the Impact of Transcranial Alternating Current

Stimulation on EEG Correlates of Somatosensory Perception. Front Psychol 9:2117.

Talsma D, Doty TJ, Woldorff MG (2007) Selective attention and audiovisual integration: is

attending to both modalities a prerequisite for early integration? Cereb Cortex 17:679-690.

Teder-Sälejärvi W, McDonald J, Di Russo F, Hillyard S (2002) An analysis of audio-visual

crossmodal integration by means of event-related potential (ERP) recordings. Cognitive

Brain Research 14:106-114.

Tolley N, Rodrigues PLC, Gramfort A, Jones S (2023) Methods and considerations for estimating

parameters in biophysically detailed neural models with simulation based inference.

bioRxiv.

Uutela K, Taulu S, Hamalainen M (2001) Detecting and correcting for head movements in

neuromagnetic measurements. Neuroimage 14:1424-1431.

Wu PY, Chu YH, Lin JL, Kuo WJ, Lin FH (2018) Feature-dependent intrinsic functional

connectivity across cortical depths in the human auditory cortex. Sci Rep 8:13287.

Zeki S (2018) The Rough Seas of Cortical Cartography. Trends Neurosci 41:242-244.

Ziegler DA, Pritchett DL, Hosseini-Varnamkhasti P, Corkin S, Hamalainen M, Moore CI, Jones

SR (2010) Transformations in oscillatory activity and evoked responses in primary

somatosensory cortex in middle age: a combined computational neural modeling and

MEG study. Neuroimage 52:897-912.