Fusobacterium sphaericum sp. nov., **isolated from a human colon tumor, adheres to colonic**

epithelial cells and induces IL-8 secretion

Martha A. Zepeda-Rivera*[1,9], Yannick Eisele*[4,5], Alexander Baryiames[2], Hanrui Wu[2], Claudia

Mengoni[6], Gianmarco Piccinno[6], Elsa F. McMahon[1], Kaitlyn D. LaCourse[2], Dakota S. Jones[1],

Hans Hauner[5], Samuel S. Minot[3], Nicola Segata[6], Floyd E. Dewhirst[7,8], Christopher D.

Johnston[1,9#], Susan Bullman[2,10#]

[1]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle, Washington,
USA
[2]Human Biology Division, Fred Hutchinson Cancer Center, Seattle, Washington, USA
[3]Data Core, Shared Resources, Fred Hutchinson Cancer Center, Seattle, Washington, USA
[4]School of Medicine, Technical University of Munich, Munich, Germany
[5]Institute Technical University of Munich of Nutritional Medicine, School of Medicine and
Health, , Munich Germany
[6]Department of Computational, Cellular and Integrative Biology, University of Trento, Trento,
Italy
[7]Forsyth Institute, Cambridge, MA, USA
[8]Department of Oral Medicine, Infection and Immunity, Harvard School of Dental Medicine,
Boston, Massachusetts, USA
[9]Genomic Medicine, MD Anderson Cancer Center, Houston, Texas, USA
[10]Immunology, James P Allison Institute, MD Anderson Cancer Center, Houston, Texas, USA
22
23

*Contributed equally to this work.

Keywords: Fusobacterium, novel species, pangenome, comparative genomics, intestinal

microbiota, colorectal cancer

#**Address Correspondence to**:

Susan Bullman, PhD, sbullman@mdanderson.org

Christopher D. Johnston, PhD, cdjohnston@mdanderson.org

Abstract

Cancerous tissue is a largely unexplored microbial niche that provides a unique environment for

the colonization and growth of specific bacterial communities, and with it, the opportunity to

identify novel bacterial species. Here, we report distinct features of a novel *Fusobacterium*

species, *F. sphaericum sp. nov.* (*Fs*), isolated from primary colon adenocarcinoma tissue. We

acquire the complete closed genome and associated methylome of this organism and

phylogenetically confirm its classification into the *Fusobacterium* genus, with *F. perfoetens* as

its closest neighbor. *Fs* is phenotypically and genetically distinct, with morphological analysis

revealing its coccoid shape, that while similar to *F. perfoetens* is rare for most *Fusobacterium*

members. *Fs* displays a metabolic profile and antibiotic resistance repertoire consistent with

other *Fusobacterium* species. *In vitro, Fs* has adherent and immunomodulatory capabilities, as it

intimately associates with human colon cancer epithelial cells and promotes IL-8 secretion.

Analysis of the prevalence and abundance of *Fs* in >20,000 human metagenomic samples shows

that it is a low-prevalence member within human stool with variable relative abundance, found in

both healthy controls and patients with colorectal cancer (CRC). Our study sheds light on a novel

bacterial species isolated directly from the human CRC tumor niche, and given its interaction

with cancer epithelial cells suggests that its role in human health and disease warrants further

investigation.

Introduction

In patients with colorectal cancer (CRC), unbiased genomic analyses have revealed an enrichment of *Fusobacterium* in CRC tumors relative to noncancerous colorectal tissue[1-4]. Previous work from our group and others demonstrated that within patient CRC tumors, *Fusobacterium* colonizes distinct regions with immune and epithelial functions supportive of cancer progression[5], that it persists with the primary tumor during metastasis[6], and that antimicrobial agents targeting these and other anaerobic bacterial species decrease cancer cell proliferation and tumor growth[6,7]. Furthermore, high *Fusobacterium* loads are associated with high-grade dysplasia[1], advanced disease stage[8], and poorer patient prognosis[9]. As tissue-associated microbiomes are not readily amenable to metagenomic analysis, due to the overabundance of host nucleic acids, characterization of tumor-infiltrating microbial communities is often restricted to amplicon-based sequencing approaches with limited phylogenetic resolution. To overcome this challenge, microbiological culture-based approaches with patient tissue specimens have reemerged as valuable tools to isolate members of tissue-associated microbiomes.

Through a culturing-based approach, we isolated, characterized, and sequenced the genome of a novel *Fusobacterium* species from the resected tumor of a treatment naïve female patient with microsatellite stable (MSS), stage III, right-sided colon cancer. Phylogenetic appraisal of this organism places this bacterium within the *Fusobacterium* genus but demonstrates that it is distinct from currently known species. Rather than the typical spindle-like, fusiforme morphologies associated with most *Fusobacterium* species, phenotypic analysis of this novel organism revealed a coccobacilloid shape. Acknowledging this distinct spherical cell-

shape, we propose this species as *Fusobacterium sphaericum sp. nov.* (*Fs*). In this present study,

we find that while the chromosomal structure and gene content of *Fs* is predominantly unique to

this species, its metabolic profile and antibiotic resistance repertoire are consistent with other

Fusobacterium members. Furthermore, our analyses indicate that its predicted virulence factors

are related to metabolism, cell-cell adherence, and immunomodulation. Further, *in vitro* co-

culture with human colon cancer epithelial cells confirmed *Fs* adherence and indicated that its

presence significantly increases IL-8 secretion. Assessment of the prevalence and abundance of

Fs in 21,212 human metagenomic samples shows that it is a low-prevalence member in human

stool with variable relative abundance.

Results

Phylogenetic and morphologic characterization of a novel *Fusobacterium* species

Fusobacterium isolation was performed on a human colorectal cancer (CRC) tumor from

a treatment naïve female patient with microsatellite stable (MSS), stage III, right-sided colon

cancer. Using long-read s̲ingle-m̲olecule r̲eal-t̲ime (SMRT)[10] sequencing we generated a

complete and closed genome and methylome for isolate SB021, identifying two putative

chromosomes (1.69 Mb and 0.57 Mb) and two additional extrachromosomal elements (1.59 Kb

and 0.32 Kb) (**Fig. 1a, Supplementary Table 1, Supplementary Table 2**). BLASTn analysis of

the 16S rRNA gene of SB021 confirmed its classification into the *Fusobacterium* genus with a

top hit against *Fusobacterium perfoetens*, yet all related hits fell below the established 16S rRNA

gene species similarity threshold of 98.65%[11], suggesting that SB021 represents a novel

Fusobacterium species (**Supplementary Table 3**). Comparative phylogenetics of SB021 and

other *Fusobacterium spp.* genomes (**Table 1**), both by the 16S rRNA gene and a reference-free

whole-genome approach[12], further showed that SB021 forms a distinct branch within the same clade as *F. perfoetens* (**Fig. 1b-c**). To quantify the relatedness between SB021 and the publicly available *F. perfoetens* ATCC 29250 genome, we calculated the average nucleotide identity (ANI) and found it to be 75.27%, well below the established species threshold of 95-96%[11], supporting the classification of SB021 as a novel *Fusobacterium* species. (**Fig. 1d, Supplementary Fig. 1, Supplementary Table 4**). GTDB-tk[13] analysis further supported these observations, as it classified SB021 within the same clade as *F. perfoetens*, as part of "Fusobacterium_B_sp900541465" (**Supplementary Table 5**). This group is predominantly represented by metagenomically assembled genomes (MAGs) from uncultured samples, with a single uncharacterized isolate from a large-scale cultivation effort of the fecal microbiota in healthy Chinese individuals[14].

The cellular morphology and pH tolerance of *Fusobacterium* cells is known to vary[15-17]. As SB021 is phylogenetically closely related to *F. perfoetens*, we assessed whether like *F. perfoetens*, SB021 cells have a short, ovoid coccobacillus morphology. Macroscopically, SB021 cells form distinct, vaguely round, raised opaque colonies with central indentations following 48 hours of anaerobic culture. The colonies are viscous and malodorous and measure 3-5 millimeters in diameter (**Fig. 1e**). Microscopically, SB021 cells are short, gram-negative coccobacilli, with an average length of 1.60µm and an average width of 1.25µm (**Fig. 1f**). Transmission electron microscopy further revealed a spherical shape and a granulated cytoplasm (**Fig. 1g**). These results suggest that members within this *Fusobacterium* clade share similar cellular morphology. Assessment for preferential growth pH indicated that SB021 is sensitive to pH below 5 and demonstrates a growth preference between pH 8 and pH 10 (**Fig. 1h**). Given the

coccobacilloid morphology of SB021 cells, we propose this novel species, part of

"Fusobacterium_B_sp900541465" as *Fusobacterium sphaericum sp. nov.* (*Fs*).

***Fs* genomic attributes predict metabolic and virulence capabilities**

As significant genomic heterogeneity has been reported within the *Fusobacterium* genus,

with genome size, architecture, and content differing between species of varying pathogenic

capabilities[17,18], we sought to characterize the genomic content of *Fs*. To compare the genetic

content of *Fs* to other *Fusobacterium spp.*, we implemented the analysis and visualization

platform for 'omics data (Anvi'o) workflow for microbial pangenomics[19]. This genus-level

pangenomic analysis allows us to identify all genes present in the *Fusobacterium* genus

("pangenome") and discern between gene content that is shared amongst ≥95% of species ("core

genome"), is shared amongst subsets of species ("accessory genomes") or is unique to individual

members ("singletons"). Across eighteen representative *Fusobacterium* genomes (**Table 1**),

only 4.9% of gene clusters compose the core (576/11,710), indicative of the extreme genetic

heterogeneity across this genus (**Fig. 2a**). *Fs* harbored 18.7% of identified gene clusters

(2,192/11,710), but of these, 31.8% (697/2,192) were unique to *Fs*, while only 6.8% (150/2,192)

were shared with its closest phylogenetic neighbor, *F. perfoetens* (**Fig. 2b**).

As the genetic content of *Fs* is predominantly unique to this species, we assessed the

2,192 identified gene clusters. KEGG ortholog analysis[20] revealed that mapped gene clusters

(50.8%) were predominantly involved in metabolic functions and pathways (**Supplementary**

Table 6). Primary metabolic pathways assessment via gutSMASH[21] indicated shared predicted

metabolic capabilities between *Fs* and other *Fusobacterium* genomes (**Supplementary Fig. 2**).

In vitro metabolic profiling via the API 20A Gallery System showed that *Fs* ferments the

monosaccharides glucose and mannose, the disaccharide lactose, the trisaccharide raffinose, and

the coumarin glucoside esculin (**Supplementary Fig. 3a**). Glucose, mannose, and raffinose

metabolism were also observed via the Biolog Anaerobe Identification Test Panel (AN Plate)

(**Supplementary Fig. 4**). Enzymatic profiling via the API ZYM panel additionally showed

positive activity for alkaline phosphatase, C4-esterase, C8-esterase-lipase (weak), acid

phosphatase (weak), napthol-AS-BI-phosphohydrolase (weak) and α-Galactosidase (strong)

(**Supplementary Fig. 3b**). This strong α-Galactosidase activity may be a distinct feature of *Fs*,

as activity for this enzyme has only been observed weakly in *F. mortiferum* and *F. necrogenes*[22].

Notably, *Fs* harbors nineteen open reading frames encoding putative multidrug resistance

proteins (**Supplementary Table 1**) and further analysis with the comprehensive antibiotic

resistance database (CARD)[23] identified 183 putative antibiotic resistance genes

(**Supplementary Table 7**), suggesting that *Fs* is resistant to a suite of antibiotics. Our group has

previously demonstrated that antibiotic-mediated microbiota modulation, targeting

Fusobacterium and other anaerobes, decreases cancer cell proliferation and tumor growth[6]. As

such, we sought to characterize the antibiotic response of *Fs*. *In vitro* antibiotic susceptibility

testing demonstrated that *Fs* is susceptible to various classes of antibiotics, including

metronidazole (MIC 0.502 µg/ml; nitroimizadole class), colistin (MIC 0.875 µg/ml; polymyxin

class), and penicillin (MIC 0.079 µg/ml; beta-lactam class). Consistent with other *Fusobacterium*

spp., *Fs* is resistant to erythromycin (macrolide class) and vancomycin (glycopeptide class)

antibiotics. Curiously, *Fs* susceptibility to aminoglycosides varied, with susceptibility to

gentamycin (MIC 62.2 µg/ml) and resistance to streptomycin and kanamycin (**Table 2**).

Given that *Fs* was isolated directly from a patient CRC tumor, we next queried the

presence of virulence factors important for host-bacterial interactions. We found that *Fs* lacks

known *Fusobacterium* virulence factors[24-32] (**Fig. 2c**) but of the 2,192 identified gene clusters,

28.4% of them are homologous to known virulence factors in other bacterial pathogens[33]

(**Supplementary Table 6**). Of this subset, the majority function to modulate the immune system

(29.7%), to scavenge and metabolize nutritional sources (15.4%), or to adhere to host cells

(12.9%) (**Fig. 2d**). This suggests that *Fs* has genetic attributes consistent with facilitating

eukaryotic cell interactions and modulation of immune responses.

_Fs_ adheres to colon cancer epithelial cells and increases IL-8 secretion

To evaluate the interaction between *Fs* and colon epithelial cells, we co-cultured *Fs*

SB021 with cell lines derived from human colorectal adenocarcinomas, HCT116 and HT-29. As

Fusobacterium nucleatum subsp. *animalis* (*Fna*) has previously been reported to adhere to and

invade cancer epithelial cells[6], we used *Fna* SB010 as a positive control. Confocal laser scanning

microscopy of fluorescent immunocytochemically stained cells demonstrated that SB021

strongly adheres to epithelial cell surfaces in chain-like clusters with both HCT116 and HT-29

(**Fig. 3a**). However, we did not observe intracellular SB021, which suggests that, unlike SB010,

SB021 adheres to but does not invade colonic epithelial cells.

To investigate whether this physical interaction between *Fs* and colonic epithelial cells

modulates eukaryotic immune responses, we assessed chemokine and cytokine induction in

HCT116 cells post co-culturing. In the presence of SB021 cells, we observed an increase in

various pro-inflammatory chemokines, most notably IL-8, occurring with both live (2.59-fold

increase) and dead (2.29-fold increase) bacterial cells (**Fig. 3b, Supplementary Fig. 5**),

indicating that these immunomodulatory capabilities of SB021 are independent of viability. IL-8

expression is elevated in colonic tissue of patients with CRC[34], enhances colon cancer cell

growth and migration to the liver and lungs[35,36], and has previously been reported to be

stimulated by the presence of other Fusobacterium species including *Fna* and *F. nucleatum*

subsp. nucleatum (*Fnn*)[37]. Prior genetic analysis in *Fnn* indicated that the stimulation of IL-8

expression is dependent on Fap2-mediated cell adhesion[37]. To assess whether physical

interaction between *Fs* and colonic epithelial cells is required for IL-8 secretion, we co-incubated

SB021 cell-free supernatant with HCT116 cells and detected a 1.27-fold increase in IL-8 (**Fig.**

3b). These results suggest that unlike other Fusobacterium species, direct bacterial-eukaryotic

cell interactions with *Fs* are not required to stimulate secretion of IL-8 and other pro-

inflammatory chemokines and cytokines (**Fig. 3b, Supplementary Fig. 5**). Furthermore,

quantitative comparison of IL-8 production in the presence of SB021 or SB010 further showed

that co-incubation with SB021 stimulated higher IL-8 levels at both 4-hour (SB021: average

13,403 pg/ml; SB010: average 1,732 pg/ml) and 24-hour (SB021: average 12,004 pg/ml; SB010:

average 5,551 pg/ml) time points (**Fig. 3c**).

Fs is a low prevalence member in human stool

207   Given that *Fs* is a novel *Fusobacterium* species with adherent and immunomodulatory

capabilities, we next sought to determine its prevalence in human specimens from various body

sites. The presence of *Fs* SB021 and its nearest phylogenetic neighbor, *F. perfoetens* ATCC

29250, was assessed across 21,212 publicly available human metagenomic samples from the

human oral cavity, skin, vagina, airways, breastmilk, nasal passage, and stool (**Fig. 4a,**

Supplementary Table 8). Results indicated that SB021 could be detected in stool samples at a

low prevalence (147/21,382=0.687%), with a range of relative abundances from 0.0005%-

17.829% when detected (**Fig. 4b**). Of the 147 samples in which SB021 could be detected, 31

originated from cohorts comparing patients with CRC to healthy controls. Nevertheless, no

significant difference in SB021 prevalence (p=0.1936, two sample Z-test) (**Fig. 4c**) or abundance

(p=0.8124, Welch's T-test) (**Fig. 4d**) was observed based on disease status in those cohorts. In

contrast, ATCC 29250 was detected in only a single stool sample with a low relative abundance

of 0.00468% (**Supplementary Table 8**). Collectively, these findings shed light on a novel

bacterial species isolated from the CRC tumor niche, which is also detectable within the human

intestinal microbiota.

Discussion

Owing to the abundance of host-associated nucleic acids in complex specimens, tissue-

associated microbiome analysis is currently restricted to amplicon-based sequencing. As such,

applying traditional culturing-based approaches to tumor specimens provides a valuable

opportunity to expand our understanding and classification of tumor-infiltrating microbes[38].

Here, we report the isolation and characterization of a novel *Fusobacterium* species, *F.*

sphaericum sp. nov. (*Fs*), from a primary colon adenocarcinoma of a female patient.

Phylogenetic and pangenomic analyses reveal that *Fs* is most closely related to *F. perfoetens*

(ATCC 29250), but support that *Fs* is a distinct species with predominantly unique gene content

(**Fig. 1, Fig.2**).

Motivated by the putative cell-cell adherent and immunomodulatory capabilities of *Fs*

genetic content (**Fig. 2d**), we investigated the interactions between *Fs* and colonic epithelial cells

in vitro. Transmission electron microscopy and confocal microscopy of fluorescent

immunocytochemically stained cells from *Fs* co-culture with human colon epithelial cell lines

demonstrated that *Fs* adheres to eukaryotic epithelial cells but does not appear to have invasive

capabilities, in contrast to other *Fusobacterium* species, such as *F. nucleatum* subsp. *animalis*

(*Fna*) (**Fig. 3a**). *Fs* cells furthermore induced eukaryotic IL-8 secretion from epithelial cells at

higher levels than *Fna* (**Fig. 3b-c**). Recent studies have demonstrated increased secretion of IL-8

in colorectal cancer (CRC) tissue and liver metastases[34,35] with a significant upregulation in the

presence of other *Fusobacterium* species including *Fna* and *F. nucleatum* subsp. *nucleatum*

(Fnn)[37]. However, in *Fnn* induction of IL-8 secretion is dependent on Fap2-mediated adhesion to

host cells[37]. As *Fs* lacks Fap2 (**Fig. 2c**) and was not observed to invade colonic epithelial cells

(**Fig. 3a**), further studies to identify the *Fs* factors that induce IL-8 secretion are warranted. As

Fs induction of IL-8 secretion was comparable independent of bacterial cell viability, and to a

lower extent with cell-free supernatants (**Fig. 3b**), it is possible such factors are secreted or

present in outer membrane vesicles. Demonstrating that *Fs* induces IL-8 secretion adds a novel

species to a growing body of evidence for a distinct role of microbes contributing to a pro-

inflammatory environment within human tissues, including tumors.

We demonstrate that in human metagenomic samples (n=21,212) *Fs* is also a member of

the human stool microbiota in both healthy individuals and patients with CRC (**Fig. 4**). Although

no significant difference in prevalence or abundance based on health status was observed, this

could be due to the small number of Fs^+ samples (n=31). Future genomic studies including *Fs*

will be required to reveal its relationship with other microbial members and decipher its impact

on the human host. Whether the presence of *Fs* in colorectal patients may contribute to a model

of dysbiosis and chronic bacterial infection in the context of CRC requires further investigation

in animal models and within longitudinal studies of large human cohorts.

The closest phylogenetic neighbor of *Fs* SB021, *F. perfoetens* ATCC 29250, was only

detected in a single human stool sample. It is noteworthy that the first described isolate of *F.*

perfoetens, acquired in 1900 from the diarrhea of infants, was unfortunately lost to history. As a

result, subsequent isolates from a horse cecum (strain CC1) and piglet feces (strain ATCC

29250) were classified as the same species as the original 1900 isolate, but based only on cellular

morphology and metabolism, as sequencing technologies for phylogenetic classifications were

unavailable at the time[16]. Although this species underwent various name changes, ATCC 29250

was designated as the type strain for species *F. perfoetens* and remains the only available isolate

in existence[16]. The very low detection of *F. perfoetens* ATCC 29250 across human metagenomic

samples is likely reflective of its swine origin and perhaps suggests that the original 1900 human

stool isolate was a phylogenetically distinct microbe despite morphological similarities. With its

similar coccobacilloid morphology, it is possible that *Fs* SB021, isolated from a human colonic

tumor, is a phylogenetic representative of the original 1900 human isolate, distinct from *F.*

perfoetens ATCC 29250.

In conclusion, in this study we have emphasized and demonstrated the importance of

reductionist approaches, such as traditional microbial culturing, to isolate and identify members

of tissue-associated microbiota which are not yet amenable to metagenomics. Next generation

sequencing techniques, such as long-read SMRT-Sequencing[10], allows for the assembly of high-

quality, closed, and complete genomes from which accurate phylogenetic characterization,

putative functions and virulent capabilities can be predicted through comparative genomic

analyses. Deciphering the role of identified novel microbial species within tumor-associated

bacterial communities on disease outcomes remains crucial for informing future preventative and

therapeutic interventions.

Figures

Fig. 1: Taxonomic and Morphologic Analysis of *Fusobacterium sphaericum sp. nov.* (SB021)

a) Proksee circular genome architecture of *F. sphaericum sp. nov.* (*Fs*) SB021 contigs. From outwards to inwards, the first and second circles show annotated coding sequences on positive strand and negative strand, respectively. Third circle indicates average GC content, with areas higher than average GC content as outward projections and areas lower than average as inward

projections. Fourth circle shows GC skew, with above zero values in green and below zero

values in purple. **b)** A 16S rRNA gene-based dendrogram illustrating the phylogenetic

relationship between *Fs* (blue) and other *Fusobacterium spp*. **c)** kSNP[12] maximum-likelihood

whole-genome phylogenetic tree with *Fs* highlighted in blue. **d)** Clustered average nucleotide

identity (ANI) matrix with *Fs* highlighted in blue. ANI values are reported in **Supplementary**

Table 4. White to black scale indicates and ANI from 65% to 100%, respectively. **e)** Digital

photography of *Fs* SB021 colonies grown on fastidious anaerobe agar enriched with 10%

defibrinated horse blood. **f)** Bright-field microscopy of Gram-stained *Fs* SB021. Images acquired

at 100X magnification with oil immersion lens. Image scale bar is 10 µm. **g)** Transmission

electron micrographs of an *Fs* SB021 cell. Image scale bar is 500 µm. **h)** Plot indicates growth

activity as measured in Biolog PM10 plates for *Fs* SB021. Data is normalized to the maximum

value per replicate (n=3).

Fig. 2: *Fusobacterium sphaericum sp. nov.* **(SB021) Gene Content Analysis**

a) Anvi'o pangenomic analysis[19] of *F. sphaericum sp. nov.* (*Fs*) SB021 (blue) as compared to seventeen other *Fusobacterium spp.* genomes (**Table 1**). Each circle represents an individual genome, with identified gene clusters in blue (SB021) or black (other *Fusobacterium spp.*). Genomes are ordered by ANI clustering (**Fig. 1d**). **b)** Upset plot depicting the presence of gene clusters identified in *Fs* across other *Fusobacterium spp.* genomes. Bar height (top) indicates number of gene clusters shared between combinations of genomes (bottom). **c)** Presence (black) versus absence (white) plot of canonical virulence factors across *Fusobacterium spp.* genomes, ordered by a kSNP[12] maximum-likelihood whole-genome phylogenetic tree with *Fs* highlighted

313 in blue. **d)** *Fs* anvi'o-identified gene clusters were analyzed using the virulence factor database

314 (VFDB)[33]. Graph shows the number of gene clusters in each of the top functional categories of

315 putative virulence factors (**Supplementary Table 5**).

a

SB010 | SB021

Middle Cross Section | Middle Cross Section | Top View

HCT116

HT-29

Eukaryotic Stains: Actin DAPI   Bacterial Stains: Membrane Dye (FM4-64X) DAPI

b IL-8

Fold Change

Antibody Cocktail, Fna, Fs (live), Fs (dead), Fs (supernatant)

c 4-hour co-incubation

IL-8 (pg/ml)

p=0.0008
p=0.0008
p=0.0015
ns
ns

No bacteria control, Ec, Fna, Fs (live)

24-hour co-incubation

IL-8 (pg/ml)

p=0.0001
p=0.0001
p=0.0012
p=0.0022
p=0.0022

No bacteria control, Ec, Fna, Fs (live)

Fig. 3: *Fusobacterium sphaericum sp. nov.* **adheres to colonic epithelial cells and stimulates**

IL-8 secretion

Confocal laser scanning microscopy of colon cancer epithelial cell lines, HCT116 and HT-29,

co-incubated with *Fusobacterium nucleatum* subsp. *animalis* (*Fna*) SB010 or *F. sphaericum sp.*

nov. (*Fs*) SB021. Bacterial cell staining: FM4-64X (red) and DAPI (blue). Eukaryotic cell

staining: DAPI (blue), actin (green). **b)** Comparison of IL-8 secretion measured via the Multi-

Analyte ELISArray from HCT116 cells co-incubated with different *Fusobacterium* species or

antibody cocktail control, normalized to a no-bacteria control. Bacterial strains used included

SB010 and SB021 live cells or oxygen-exposed dead cells. An additional co-incubation with

SB021 supernatant alone was also performed. Additional cytokine and chemokine measurements

are shown in **Supplementary Fig. 5**. **c)** Quantitative levels of IL-8 secretion measured via the

Single-Analyte ELISArray from HCT116 cells incubated alone or co-incubated with *Escherichia*

coli TOP10, SB010, or SB021 at 4-hour and 24-hour time points. Statistical analysis performed

via a one-way ANOVA. For **a-c**, all co-incubations were at a multiplicity of infection (MOI) of

100.

Fig. 4: *Fusobacterium sphaericum sp. nov.* **is present in human stool.**

The relative abundance of *F. sphaericum sp. nov.* (*Fs*) SB021 was assessed in **a)** 21,212 metagenomic samples from a various human body sites. **b)** Graphs show *Fs* percent relative abundance in samples detected. Black line indicates mean. **c-d)** In datasets comparing patients with CRC to healthy controls **c)** graph shows proportion of samples in which *Fs* was detected. Statistical analysis performed via two-sample Z test. **d)** Graph shows *Fs* percent relative abundance in samples detected for patients with CRC or healthy controls. Black line indicates mean. Statistical analysis performed via Welch's T-test.

 Table Legends

345 **Table 1: Genomes used in this study.**
346 Table indicates each *Fusobacterium* genome used for comparative analyses in this study with
347 their corresponding genome accession numbers, genome length, GC content, and number of
348 contigs in the assembly noted.
349
350 **Table 2: *F. sphaericum sp. nov.* antibiotic resistance profile**
351 Table shows results for *F. sphaericum sp. nov.* susceptibility or resistance to nine tested
352 antibiotic compounds.
353

354   *Supplementary Figures*

Supplementary Fig. 1: Genomic architecture of *Fusobacterium sphaericum sp. nov.* SB021

as compared to *Fusobacterium perfoetens* ATCC 29250.

Visualization of fastANI alignment between *Fusobacterium sphaericum sp. nov.* SB021 and its

nearest phylogenetic neighbor, *Fusobacterium perfoetens* ATCC 29250, where each red line

segment denotes a reciprocal mapping of the 275 orthologous matches between the two

genomes, indicating evolutionarily conserved regions.

Supplementary Fig. 2: Primary metabolic pathways in *Fusobacterium sphaericum* SB021

Presence (black) versus absence (white) plot of primary metabolic pathways identified by gutSMASH[21] across *Fusobacterium spp.* genomes, ordered by a kSNP[12] maximum-likelihood whole-genome phylogenetic tree with *F. sphaericum sp. nov.* SB021 highlighted in blue.

Supplementary Fig. 3: API metabolic characterization of *Fusobacterium sphaericum sp.*

nov.

F. sphaericum sp. nov. (*Fs*) **a)** metabolic and **b)** chemotaxonomic characteristics assayed by API

20A and API ZYM detection kits, respectively. Images are representative results for *Fs* SB021.

Results for *Fs* SB021, *Fusobacterium nucleatum* subsp. *animalis* SB010, and *Bacteroides*

fragilis are summarized below each API strip, where (+) indicates a positive result, (-) a negative

result, and "*w*" a weakly positive result.

Substrate	OD$_{max}$ Normalized to Water Control	p-value
Water		
Urocanic acid		0.008217
Uridine-5'-mono-phosphate		0.692929
Uridine		0.021194
Turanose		0.004532
Thymidine-5'-mono-phosphate		0.135523
Thymidine		0.011693
Sucrose		0.008174
Succinic acid mono methyl ester		0.073138
Succinic acid		0.403025
Succinamic acid		0.667086
Stachyose		0.029293
Salicin		0.044489
Pyruvic acid methyl ester		0.001280
Pyruvic acid		0.102547
Propinic acid		0.704942
Palatinose		4.16E-05
N-acetyl-D-galactosamine		0.158245
N acetyl Beta D mannosamine		0.009274
N-acetyl-Beta-D-glucosamine		0.032927
Maltotriose		0.001400
Maltose		0.003071
m-Tartaric acid		0.117085
m-Inositol		0.168982
Lactulose		0.006336
L-valine + L-aspartic acid		0.447560
L-valine		0.120809
L-threonine		0.054017
L-serine		0.003536
L-rhamnose		0.138255
L phenylalanine		0.058592
L-methionine		1.54E-05
L-maliac acid		0.326006
L-lactic acid		0.081255
L glutamine		0.669195
L-glutamic acid		0.086694
L-fucose		0.000452
L-asparagine		0.414496
L-alanyl-L-threonine		0.015169
L-alanyl-L-histidine		0.008855
L alanyl L glutamine		0.014483
L-alanine		0.073740
Itaconic acid		0.001499
Inosine		0.088073
i-Erythritol		0.654506
Glyoxylic acid		0.008848
Glycyl-l -proline		0.662104
Glycyl-L-methionine		0.001890
Glycyl-L-glutamine		0.518042
Glycyl-L-aspartic acid		0.248276
Glycerol		0.070117
Gentiobiose		0.003946
Fumaric acid		0.251531
Formic acid		0.037217
Dulcitol		0.402789
Dextrine		0.004866
D,L-lactic acid		0.154663
D,L-alpha-glycerol phosphate		0.169483
D-trehalose		0.006014
D-sorbitol		0.008281
D saccharic acid		0.899614
D-raffinose		0.035048
D-melibiose		0.000855
D-melezitose		0.008173
D-mannose		0.033346
D-mannitol		0.353937
D-malic acid		0.419386
D-Lactic acid methyl ester		0.008196
D-glucose-6-phosphate		0.116122
D-glucosaminic acid		0.287102
D-gluconic acid		0.963731
D-Galacturonic acid		0.032048
D-galactose		0.002242
D-fructose		0.001149
D-cellobiose		0.000470
D-arabitol		0.056127
D-1-alpha-methyl-D-galactoside		0.059990
Beta-methyl-D-glucoside		0.008695
Beta methyl D galactoside		0.002321
Beta-hydroxybutyric acid		0.014789
beta-cyclodextrin		0.182571
Arbutin		0.024074
Amygdalin		0.049623
alpha-methyl-D-glucoside		0.037988
alpha-ketovaleric acid		0.003486
alpha-ketobutyric acid		0.058895
alpha hydroxybutyric acid		0.115743
alpha-D-lactose		3.44E-05
alpha-D-glucose-1-phosphate		0.243001
alpha-D-glucose		0.057454
alpha-cyclodextrin		0.010010
Alaninamide		0.358624
Adonitol		0.313745
Acetic acid		0.327904
3-methyl-D-glucose		0.019645
2-Deoxy Adenosine		0.011188

■ *Fusobacterium nucleatum subsp. animalis*   ■ *Fusobacterium sphaericum sp. nov.*

Supplementary Fig. 4: Biolog anaerobe identification test panel characterization of *F.

sphaericum sp. nov. substrate utilization**

F. sphaericum sp. nov. substrate utilization assayed by Biolog Anaerobe Identification Test

Panel (AN) assay plate. Graphs show results for *F. sphaericum sp. nov.* SB021 (blue) and

Fusobacterium nucleatum subsp. *animalis* (*Fna*) SB010 (black) normalized to internal water

control, in triplicate. Statistical analysis performed via a two-sided T-test. All p-values less than

0.05 are highlighted in red.

Supplementary Fig. 5: Cytokine and chemokine induction in HCT116 cells co-incubated

with *F*. *sphaericum sp. nov.*

Measurements of **a**) common chemokine and **b**) inflammatory cytokine levels measured via the

Multi-Analyte ELISArray from HCT116 cells co-incubated with different *Fusobacterium*

species at a multiplicity of infection (MOI) of 100 or antibody cocktail control, normalized to a

no-bacteria control. Bacterial species used included *F. nucleatum* subsp. *animalis* (*Fna*) SB010

and *F. sphaericum sp. nov.* (*Fs*) SB021 live cells or oxygen-exposed dead cells. An additional

co-incubation with *Fs* supernatant alone was also performed.

396
397

Supplementary Table Legends

Supplementary Table 1: RAST annotation of *F. sphaericum sp. nov.* SB021 genome
The *F. sphaericum sp. nov.* SB021 genome was annotated by the Rapid Annotation using Subsystem Technology (RAST) pipeline. Table indicates resulting genes for each contig, with their length and predicted product.

Supplementary Table 2: Methyl-modified motifs across *F. sphaericum sp. nov.* SB021 genome
SB021 methyl-modified nucleotides were analyzed using Single-Molecule Real Time Sequencing (SMRTSeq)[10] kinetics (Basemod analysis) and the predicted methyl-modified motif acquired via REBASE[39] analysis.

Supplementary Table 3: BLASTn analysis of *F. sphaericum sp. nov.* SB021 16S rRNA gene
Table shows BLASTn top 100 results for the *F. sphaericum sp. nov.* SB021 16S rRNA gene by decreasing percent identity.

Supplementary Table 4: Average nucleotide identity (ANI) scores between *Fusobacterium* genomes
Table shows the pairwise average nucleotide identity (ANI) scores between each *Fusobacterium* genome (Table 1).

Supplementary Table 5: GTDB-tk analysis of *F. sphaericum sp. nov.* SB021 genome
Table shows the assigned phylogenetic classification of SB021 as determined by GTDB-tk[13].

Supplementary Table 6: Analysis of *F. sphaericum sp. nov.* SB021 anvi'o-identified gene clusters
The *Fusobacterium* pangenome was characterized using the analysis and visualization platform for microbial 'omics (Anvi'o)[19]. For each gene cluster (GC) identified in *F. sphaericum sp. nov.*, the table shows its representative amino acid sequence, its mapped KO identifier by KEGG ortholog mapping via KofamKOALA[20], and any identified hits by the virulence factor database (VFDB)[33].

Supplementary Table 7: CARD analysis of *F. sphaericum sp. nov.* SB021 genome
Table shows results for the analysis of the *F. sphaericum sp. nov.* SB021 genome against the comprehensive antibiotic resistance database (CARD)[23].

Supplementary Table 8: SGB detection across human metagenomic samples
Table shows relative abundance for *F. sphaericum sp. nov.* SB021 (SGB59307_group) and *Fusobacterium perfoetens* ATCC 29250 (SGB29302) across 21,382 human metagenomic samples. For each sample, dataset accession number and body site sampled is included.

Methods

Bacterial isolation and culturing

Written informed consent was obtained from patient as approved by the Fred Hutchinson Cancer Center Institutional Review Board. Primary adenocarcinoma tumor tissue was collected by hospital staff. Tissue was stored in sterile, chilled RPMI growth medium (Corning, Corning NY, USA) on ice. A small cross section of the tumor was cut from the main tissue mass, lightly scored, and then spread onto a small section on fastidious anaerobe agar (FAA) plates (Oxoid, ThermoFisher Scientific, Waltham, MA, USA) supplemented with 10% defibrinated horse blood (DHB; Lampire Biological Laboratories, Fisher Scientific, Pipersville, PA, USA). Using a sterile loop, the section was spread on the plates for the isolation of single colonies and subsequently, plates were incubated at 37°C under anaerobic conditions. Every 48 hours the plates were checked for new growth and the identity of colonies was determined by colony PCR. For all subsequent experiments, *F. sphaericum sp. nov* SB021 was grown for 48 hours on FAA+10% DHB plates supplemented with josamycin (3 μg/ml), vancomycin (4 μg/ml), and norfloxacin (1μg/ml) (Sigma Aldrich, USA) and incubated at 37°C under anaerobic conditions (AnaeroGen Gas Generating Systems, Oxoid, ThermoFisher Scientific, USA).

16S rRNA bacterial colony PCR

Single colonies were collected with a 1μl loop and suspended in 20μl of nuclease free water (Promega, Madison WI, USA)**.** The same 1μl loop was then used to streak a FAA + 10% DHB plate to subculture and isolate pure colonies. The suspensions were placed in a thermocycler at 98°C for 20 minutes and then cooled to 4°C. 2μl of the lysate was subsequently used in a 20μL PCR reaction (Promega, Madison WI, USA) with forward primer 342F (5'- CTA CGG GGG GCA GCA G -3') and reverse primer 1492R (5'- TAC GGY TAC CTT GTT ACG ACT T -3')

at a final concentration of 1μM. PCR products were shipped for Sanger Sequencing and

sequencing results were analyzed using NCBI's BLASTn for species level identification.

Identified strains were stocked in 60% TSB with 40% glycerol (v/v) at -80°C.

High-molecular weight genomic DNA (gDNA) extraction

High molecular weight genomic DNA was extracted using the MasterPure^TM Gram Positive

DNA Purification Kit (Epicentre, Lucigen, USA). Cells from two plates were re-suspend in

1.5mL 1X PBS and harvested by centrifugation. Pellets were subjected to manufacturer

instructions modified by doubling all reagent volumes and removing vortexing steps to prevent

DNA shearing. HMW gDNA was quantified using a Qubit fluorometer (ThermoFisher

Scientific, USA).

PacBio Single-Molecule Real-Time Sequencing and Genome Assembly

Single molecule real-time sequencing (SMRT-Seq)[10] was carried out on a PacBio Sequel

instrument (Pacific Biosciences, USA) or a PacBio Sequel II instrument (Pacific Biosciencies,

USA) at the University of Minnesota Genomics Center. Sequencing reads were processed using

Microbial Assembly pipeline within Pacific Biosciences' SMRTAnalysis pipeline version

9.0.0.92188. Additional assembly was performed using Flye assembler version 2.8

(https://github.com/fenderglass/Flye). Functional predictions for genes was conducted with the

Rapid Annotations using Subsystems Technology (RAST)[40,41] tool (**Supplementary Table 1**).

Phylogenetic classification

The full-length 16S rRNA gene sequence for *F. sphaericum sp. nov.* SB021 was extracted from

its whole genome sequence and analyzed using NCBI BLASTn (**Supplementary Table 3**). For

available *Fusobacterium* type strains, the 16S rRNA gene sequence was downloaded and all

resulting sequences were aligned via MEGA X[42] using the MUSCLE clustering algorithm from

which a maximum-likelihood dendrogram was generated. kSNP3[12] with a kmer size of 17,

resulting in a fraction of core kmers (FCK) of 0.113, was used to generate a maximum-

likelihood whole-genome, reference-free phylogeny of these strains. Final tree were generated

using the interactive tree of life (iTOL) tool, version 5[43]. Average nucleotide identity was

calculated using the EZ BioCloud ANI calculator[44]. Further phylogenetic classification was

determined using GTDB-tk[13] (https://github.com/Ecogenomics/GTDBTk). For *Fusobacterium*

nucleatum, ANI thresholds have led to adoption of different naming schemes based on

recognition of the historical four subspecies as distinct species. In this work, we have

implemented the *F. nucleatum sensu latu* nomenclature, referring to each subgroup by their

historical subspecies name.

Pangenomic Analysis

Pangenome analysis was performed using the Analysis and Visualization platform for microbial

'omics (Anvi'o) workflow[19] with standard thresholds set to a minbit of 0.5 and an MCL of 2. All

F. sphaericum sp. *nov.* gene clusters were analyzed via KofamKOALA[20] for KEGG ortholog

mapping for their putative functions and analyzed against the Virulence Factor Database

(VFDB)[33] to identify putative virulence-associated genes.

Bacterial Gram Staining

Gram staining was performed on *F. sphaericum sp. nov.* SB021 uniform bacterial smears after

growth at 37°C for 48 hours followed by toxic oxygen exposure using a Gram Stain Kit (Remel,

Lenexia, KS, USA). The stains were imaged with TissueFAXS microscope system

(TissueGnostics, Vienna, Austria) with 100x oil immersion at Fred Hutchinson Cancer Center,

Seattle, WA, USA.

Biolog PM10 phenotype and anaerobe identification test panel microarray plates

F. sphaericum sp. nov. SB021 and *F. nucleatum* subsp. *animalis* SB010 were grown on FAA

plates (Oxoid, Thermo Fisher Scientific) supplemented with 10% DHB (Fisher Scientific)

incubated at 37°C in a Concept1000 anaerobic chamber (BakerRuskinn) for 24 h. PM10 and

AN plates with corresponding inoculating fluids (IF-0a/IF-10b, and AN-IF, correspondingly)

were pre-reduced, either at 4°C overnight (PM10) (AnaeroGen Gas Generating Systems, Oxoid,

Thermo Fisher Scientific) or by manufacturer (AN). PM10 and AN plates were brought to room

temperature under anaerobic conditions (AnaeroGen Gas Generating Systems, Oxoid, Thermo

Fisher Scientific). Bacterial suspensions were made and added to each plate under anaerobic

conditions at 37°C (BakerRuskinn). For PM10 plates, bacterial cells were resuspended in 2 ml

of pre-reduced IF-0a and normalized across all samples to an optical density at 600nm (OD_{600nm})

of 0.179 as recommended by Biolog. The final suspension was prepared by combining

0.75 ml of normalized bacterial suspension with 11.25 ml of mix B (100 ml pre-reduced IF-

10b with 1.2 ml dye mix D, and 11.18 ml pre-reduced sterile water) to a final volume of

12 ml. For AN plates, bacterial cells were resuspended in 3 ml of pre-reduced AN-IF and

normalized across all samples to an OD_{600nm} of 0.1 as recommended by Biolog. For both PM10

and AN plates, 100 μl of their respective final suspension was added to each well. All plates

were then equilibrated to aerobic conditions at room temperature for 10 min and then incubated

under anaerobic, hydrogen-free conditions for 24 h at 37°C (AnaeroGen Gas Generating

Systems, Oxoid, Thermo Fisher Scientific). All plates were imaged and absorbance at 590 nm

was quantified using a plate reader (Biotek).

API 20A and API Zym Metabolic Profiling

The metabolic profile of *F. sphaericum sp. nov.* SB021 was determined with the API 20A

Gallery System (bioMérieux, Marcy-l'Étoile, France) by utilization of colonies after 24h

anaerobic culture. The colonies were resuspended in phosphate-buffered saline (PBS, Corning,

Corning, NY, USA) to obtain an optical density at 600nm (OD_{600nm}) of 0.4-0.5 (=3 McFarland)

for inoculation of pre-coated cupules on the incubation strip. Results were read after 24 hours

and additional procedures were carried out as indicated by the manufacturer. The enzymatic

profile was investigated similarly with the API Zym Gallery System (bioMérieux, Marcy-

l'Étoile, France), except for an OD_{600nm} of 0.7 (=5-6 McFarland) and an incubation period of 4-

4.5 hours. Both API 20A and API Zym were performed with 3 replicates.

Antibiotic Sensitivity Testing

In vitro testing of antibiotic susceptibility of *F. sphaericum sp. nov.* SB021 was performed on

harvested bacteria after 24 hours incubation at 37°C under anaerobic conditions. Cells were

resuspended in PBS to an OD_{600nm} of 0.5 and used to seed bacterial lawns on fastidious anaerobe

agar (FAA) plates (Oxoid, ThermoFisher Scientific, Waltham, MA, USA) supplemented with

10% defibrinated horse blood (DHB; Lampire Biological Laboratories, Fisher Scientific,

Pipersville, PA, USA). Antibiotic reading strips (MIC test strips; Liofilchem Diagnostics,

Waltham, MA, USA) were placed on the lawns with sterile tweezers. All experiments were

performed under anaerobic conditions and in triplicates for each antibiotic agent including

negative controls (no bacteria). Results were read after 24 hours of incubation.

Mammalian Cell Culture

Mammalian cell culture was performed on the cell lines HCT116 (CCL-247; ATCC, Manassas,

VA, USA) and HT-29 (HTB-38; ATCC, Manassas, VA, USA). Mammalian cells were grown in

T75 culture flasks using McCoy's 5A culture medium (Iwakata and Grace Modification)

(Corning, Corning NY, USA) with 10% fetal bovine serum (FBS; Sigma-Aldrich, St. Louis,

MO, USA), and antibiotics (1x Pen/Strep: penicillin (5000 units/ml), streptomycin (5000 µg/ml),

Gibco, Carlsbad, CA, USA) and incubated constantly in a 37°C/5% CO_2 air humidified

incubator. Cells were checked microscopically for monitoring purposes (growth rate,

confluency) and split upon reaching confluency (>=80%). Trypsin-EDTA (Gibco, Carlsbad, CA,

USA) was added to the flask to dissociate cells from flask surface, followed by neutralization of

the dissociation agent with growth medium. Subsequently, the cell suspension was centrifuged

for 3 minutes at 302 RCF at room temperature. The cell pellet was resuspended in sterile

medium and seeded into new flasks. Passage numbers were recorded and did not exceed $n=10$.

Prior to co-culture experiments, cells were split as described above and resuspended in

antibiotic-free sterile growth medium, added to 6-well plates to a final concentration of 1.5×10^6

cells per well and incubated for 24 hours before experimental testing.

Scanning Transmission Electron Microscopy (S/TEM)

F. sphaericum sp. nov. SB021 was cultured as described above. Cell size of *F. sphaericum sp.*

nov. SB021 was determined via scanning transmission electron microscopy (S/TEM). We co-

incubated HCT116 human colorectal carcinoma cell line (CCL-247; ATCC, Manassas, VA,

USA) at low passage (<10) with *F. sphaericum sp. nov.* SB021 ($OD_{600nm} = 1.0$) for 4 hours

followed by fixation with ½ strength Karnovsky's fixative (2.5% glutaraldehyde and 2%

paraformaldehyde in 0.1M cacodylate buffer) and further sample preparation on the same day.

Specimens were viewed on a JEOL JEM 1400 transmission electron microscope (JEOL, Tokyo,

Japan) with an accelerating voltage of 120kV. Images were acquired with a Gatan Rio 4k x 4k

CMOS Camera (Gatan, AMETEK, Berwyn, PA, USA) at Fred Hutchinson Cancer Center,

Seattle, WA, USA.

Confocal Microscopy of *Fusobacterium* co-cultures with human cell lines

Fusobacterium strains, *F. sphaericum sp. nov.* SB021 and *F. nucleatum* subsp. *animalis* SB010,

were grown for 48 hours at 37°C in anaerobic conditions and resuspended in 1 mL of PBS at a

final concentration of $5x10^8$ cells/mL. Bacterial membranes were stained in this suspension with

50 µl of 100 mg/mL FM 4-64FX (Molecular Probes) and washed twice in PBS by centrifugation

for 7,000 x g at RT. Cells from human adenocarcinoma tumor cell lines, HCT116 and HT-29,

were grown and passaged as described above and resuspended in 1 mL of McCoys 5A + 10%

FBS. $6x10^4$ HCT116 or HT-29 cells were seeded with an MOI of 100 bacterial cells in 6-well

plates. Bacteria were centrifuged onto HCT 116 and HT-29 at 300 x g for 5 minutes at room

temperature (RT), and plates were immediately placed in a 37°C CO_2 incubator and cultured

together for 4 hours. Cell lines were then washed 3x with PBS (all washes are 5 min in duration

at RT) and then fixed in 4% paraformaldehyde in PBS for 30 min at RT. Following fixation,

wells were washed 3x in PBS and then permeabilized with 0.2% (*v/v*) Triton X-100 in PBS for 4

mins at RT. Cells were washed 3x in PBS and then stained for 20 min at RT with 2 drops/mL of

NucBlue Fixed Cell Strain ReadyProbes (Invitrogen, Carlsbad, CA, USA) and ActinGreen 488

Ready Probes (Invitrogen) to stain DNA and actin, respectively. Cells were washed a final 3x

with PBS and then the coverslips were mounted onto glass slides in ProLong Gold antifade

mounting medium (Invitrogen). Samples were viewed with a Leica SP8 confocal laser-scanning

microscope (Leica, Wetzlar, Germany) for image acquisition. Representative confocal

micrographs of 1024 x 1024 pixels (pixel size: 103.3 nm) were acquired and assembled using

Fiji (with Bio-Formats Plugin)[49].

Analysis of Chemokine and Cytokine Induction in HCT116

To determine the immunologic impact of *Fusobacterium* presence on HCT116 human colon

epithelial cells, we utilized the Multi-Analyte ELISArray kits protocol version 1.4 from

SABiosciences (Qiagen, Valencia, CA, USA) for common human chemokines and inflammatory

human cytokines according to the manufacturer's instructions. Tests were performed with the

following samples after a 4-hour co-incubation with HCT116 (low passage, <10): *F. sphaericum*

sp. nov. SB021 at MOI of 1:100 (live and oxygen-exposed), *F. sphaericum sp. nov.* SB021 cell-

free supernatant, *F. nucleatum* subsp. *animalis* SB010 at MOI 1:100 (live), a negative control

(HCT116 only) and a positive control (cocktail of antibodies). Supernatants were centrifuged at

1,000 x g for 10 minutes to remove any particulate material, filter sterilized and 50µl of each

experimental sample were added to the array with the specific capture antibodies (IL-1α, IL-1β,

IL-2, IL-4, IL-6, IL-8, IL-12, IL-17α, IFN-γ, TNF-α and GM CSF; IL-8, MCP-1, RANTES,

MIP-1a, IP-10, I-TAC, MIG, Eotaxin, TARC, MDC, GROα), followed by an incubation at room

temperature (RT) for 2 hours. Subsequently, the wells were buffer washed and 100µl of diluted

biotinylated detection antibodies were added for a 1h-incubation at RT in the dark. After another

washing procedure, the wells were filled with 100µl of Avidin-horseradish peroxidase (HRP)

and incubated at RT for 30 minutes, again in the dark. We then added development and stop

solutions to detect absorbance changes immediately after termination of the experiment with a

microplate reader (Synergy H4 Hybrid Reader, BioTek) at 570nm and 450nm, visualized with

Gen5 Synergy H4 software (Version 3.08, BioTek). Raw data of the absorbance reading were

normalized to the cell counts of HCT116 cells incubated without bacteria. This experiment was

executed in duplicates.

For quantification of immunologic effects, we utilized the Single-Analyte ELISArray kit

protocol version 1.4. from SABiosciences (Qiagen, Valencia, CA, USA) for the chemokine IL-8

according to the manufacturer's instructions. Co-cultures were set up as described above with *F.*

sphaericum sp. nov. SB021 at MOI of 1:100 (live), *F. sphaericum sp. nov.* SB021 cell-free

supernatant, *F. nucleatum* subsp. *animalis* SB010 at MOI 1:100 (live), *Escherichia coli* TOP10,

or a positive control (HCT116 only) and incubated for 4-hours or 24-hours. Supernatants were

centrifuged at 1000 x g for 10 minutes and filter sterilized subsequently. A serial dilution of the

antigen standard was prepared and 50μl of each sample was added to the array, followed by an

incubation for 2 hours at RT. Similar to the Multi-Analyte kit, the array was then incubated with

detection antibodies and subsequently Avidin-HRP to determine the attachment of detection

antibodies to the chemokine. After the color development, changes of absorbance were read at

570nm and 450nm and raw data were equally normalized. This experiment was executed in

duplicates. Statistical comparison was calculated by applying a one-way ANOVA using

GraphPad Prism v7.0 Software (GraphPad Software).

Analysis within Human Metagenomic Specimens

Human metagenomic samples were profiled at SGB-level resolution with MetaPhlAn 4

(v4.beta.1) using the vJan21 markers database[45] SGB assignment of *Fusobacterium sphaericum*

sp. *nov.* SB021 (SGB6028, part of SGB59307_group) and *Fusobacterium perfoetens* ATCC

29250 (SGB29032) was performed with 'phylophlan_metagenomic' subroutine of PhyloPhlAn 3

(v3.0)[46].

Data Availability

F. sphaericum sp. nov. strain SB021 whole genome sequence is available in GenBank under the

accession number SAMN15202580 and methylome is available in the Restriction Enzyme

Database (REBASE) under organism number 39739.

Acknowledgements

We would like to thank the Cellular Imaging Shared Resource of the Fred Hutchinson Cancer

Center/University of Washington Cancer Consortium (P30 CA015704). Scientific Computing

Infrastructure at Fred Hutchinson Cancer Center was funded by ORIP grant S10OD028685.

Funding

Research reported in this publication was supported by the National Institute of Dental and

Craniofacial Research of the National Institutes of Health under award number R01 DE027850

(to C.D.J.), the National Cancer Institute under award number R00 CA229984-03 (to S.B.), start-

up funds provided by the FHCC (to S.B and C.D.J), a study scholarship from the Konrad

Adenauer Foundation (Y.E.), and the Washington Research Foundation Postdoctoral Fellowship

(to M.Z.R).

Contributions

Y.E., H.H., C.D.J. and S.B. designed the study. M.Z.R, Y.E., F.E.D, C.D.J., and S.B. wrote the

paper. A.B. and K.D.L. processed patient tissue specimen. A.B. and K.D.L. carried out

microbial isolation and culture. M.Z.R., D.S.J. and C.D.J. performed DNA extraction, genome

sequencing, and assembly. M.Z.R., Y.E., A.B., H.W., C.M., G.P., and S.S.M. performed

computational and statistical analysis. Y.E., A.B., E.F.M., and K.D.L. carried out *in vivo* and *in*

vitro assays. N.S., C.D.J., and S.B. obtained funding and supervised computational and wet lab

experiments. All authors read and provided edits to the paper and contributed to the final version.

Competing Interests

678
679 ***References***

680
1   Flanagan, L. *et al.* *Fusobacterium nucleatum* associates with stages of colorectal
neoplasia development, colorectal cancer and disease outcome. *Eur J Clin Microbiol*
Infect Dis **33**, 1381-1390 (2014). https://doi.org:10.1007/s10096-014-2081-3
2   Kostic, A. D. *et al.* Genomic analysis identifies association of *Fusobacterium* with
colorectal carcinoma. *Genome Res* **22**, 292-298 (2012).
https://doi.org:10.1101/gr.126573.111
3   Castellarin, M. *et al.* *Fusobacterium nucleatum* infection is prevalent in human colorectal
carcinoma. *Genome Res* **22**, 299-306 (2012). https://doi.org:10.1101/gr.126516.111
4   Viljoen, K. S., Dakshinamurthy, A., Goldberg, P. & Blackburn, J. M. Quantitative
profiling of colorectal cancer-associated bacteria reveals associations between
Fusobacterium spp., enterotoxigenic *Bacteroides fragilis* (ETBF) and clinicopathological
features of colorectal cancer. *PLoS One* **10**, e0119462 (2015).
https://doi.org:10.1371/journal.pone.0119462
5   Galeano Nino, J. L. *et al.* Effect of the intratumoral microbiota on spatial and cellular
heterogeneity in cancer. *Nature* **611**, 810-817 (2022). https://doi.org:10.1038/s41586-
022-05435-0
6   Bullman, S. *et al.* Analysis of *Fusobacterium* persistence and antibiotic response in
colorectal cancer. *Science* **358**, 1443-1448 (2017).
https://doi.org:10.1126/science.aal5240
7   LaCourse, K. D. *et al.* The cancer chemotherapeutic 5-fluorouracil is a potent
Fusobacterium nucleatum inhibitor and its activity is modified by intratumoral
microbiota. *Cell Rep* **41**, 111625 (2022). https://doi.org:10.1016/j.celrep.2022.111625
8   Mima, K. *et al.* *Fusobacterium nucleatum* in Colorectal Carcinoma Tissue According to
Tumor Location. *Clin Transl Gastroenterol* **7**, e200 (2016).
https://doi.org:10.1038/ctg.2016.53

9   Mima, K. *et al. Fusobacterium nucleatum* in colorectal carcinoma tissue and patient
prognosis. *Gut* **65**, 1973-1980 (2016). https://doi.org:10.1136/gutjnl-2015-310101
10   Eid, J. *et al.* Real-time DNA sequencing from single polymerase molecules. *Science* **323**,
133-138 (2009). https://doi.org:10.1126/science.1162986
11   Kim, M., Oh, H. S., Park, S. C. & Chun, J. Towards a taxonomic coherence between
average nucleotide identity and 16S rRNA gene sequence similarity for species
demarcation of prokaryotes. *Int J Syst Evol Microbiol* **64**, 346-351 (2014).
https://doi.org:10.1099/ijs.0.059774-0
12   Gardner, S. N., Slezak, T. & Hall, B. G. kSNP3.0: SNP detection and phylogenetic
analysis of genomes without genome alignment or reference genome. *Bioinformatics* **31**,
2877-2878 (2015). https://doi.org:10.1093/bioinformatics/btv271
13   Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925-1927
(2019). https://doi.org:10.1093/bioinformatics/btz848
14   Lin, X. *et al.* The genomic landscape of reference genomes of cultivated human gut
bacteria. *Nat Commun* **14**, 1663 (2023). https://doi.org:10.1038/s41467-023-37396-x
15   Brennan, C. A. & Garrett, W. S. *Fusobacterium nucleatum* - symbiont, opportunist and
oncobacterium. *Nat Rev Microbiol* **17**, 156-166 (2019). https://doi.org:10.1038/s41579-
018-0129-6
16   Van Assche, P. F. W., A.T. *Fusobacterium perfoetens* (Tissier) Moore and Holdeman
1973: Description and Proposed Neotype Strain. *Journal of Systematic Bacteriology* **27**,
1-5 (1977).
17   Zepeda-Rivera, M. *et al.* A distinct *Fusobacterium nucleatum* clade dominates the
colorectal cancer niche. *Nature* **628**, 424-432 (2024). https://doi.org:10.1038/s41586-024-
07182-w
18   Manson McGuire, A. *et al.* Evolution of invasion in a diverse set of *Fusobacterium*
species. *mBio* **5**, e01864 (2014). https://doi.org:10.1128/mBio.01864-14
19   Eren, A. M. *et al.* Anvi'o: an advanced analysis and visualization platform for 'omics
data. *PeerJ* **3**, e1319 (2015). https://doi.org:10.7717/peerj.1319
20   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM
and adaptive score threshold. *Bioinformatics* **36**, 2251-2252 (2020).
https://doi.org:10.1093/bioinformatics/btz859
21   Pascal Andreu, V. *et al.* gutSMASH predicts specialized primary metabolic pathways
from the human gut microbiota. *Nat Biotechnol* **41**, 1416-1423 (2023).
https://doi.org:10.1038/s41587-023-01675-1
22   De Witte, C. *et al.* Detection, isolation and characterization of *Fusobacterium gastrosuis*
sp. nov. colonizing the stomach of pigs. *Syst Appl Microbiol* **40**, 42-50 (2017).
https://doi.org:10.1016/j.syapm.2016.10.001
23   Alcock, B. P. *et al.* CARD 2023: expanded curation, support for machine learning, and
resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids*
Res **51**, D690-D699 (2023). https://doi.org:10.1093/nar/gkac920
24   Bachrach, G., Rosen, G., Bellalou, M., Naor, R. & Sela, M. N. Identification of a
Fusobacterium nucleatum 65 kDa serine protease. *Oral Microbiol Immunol* **19**, 155-159
(2004). https://doi.org:10.1111/j.0902-0055.2004.00132.x

25 Ogawa, A. T., Brasil de Souza Tde, A., de Uzeda, M., Jankevicius, J. V. & Jankevicius,
S. I. Characterization of proteolytic activities of *Fusobacterium nucleatum*. *J Endod* **32**,
521-523 (2006). https://doi.org:10.1016/j.joen.2005.10.045

26 Doron, L. *et al.* Identification and characterization of fusolisin, the *Fusobacterium*
nucleatum autotransporter serine protease. *PLoS One* **9**, e111329 (2014).
https://doi.org:10.1371/journal.pone.0111329

27 Coppenhagen-Glazer, S. *et al.* Fap2 of *Fusobacterium nucleatum* is a galactose-
inhibitable adhesin involved in coaggregation, cell adhesion, and preterm birth. *Infect*
Immun **83**, 1104-1113 (2015). https://doi.org:10.1128/IAI.02838-14

28 Kaplan, C. W. *et al. Fusobacterium nucleatum* apoptosis-inducing outer membrane
protein. *J Dent Res* **84**, 700-704 (2005). https://doi.org:10.1177/154405910508400803

29 Kaplan, C. W., Lux, R., Haake, S. K. & Shi, W. The *Fusobacterium nucleatum* outer
membrane protein RadD is an arginine-inhibitable adhesin required for inter-species
adherence and the structured architecture of multispecies biofilm. *Mol Microbiol* **71**, 35-
47 (2009). https://doi.org:10.1111/j.1365-2958.2008.06503.x

30 Lima, B. P., Shi, W. & Lux, R. Identification and characterization of a novel
Fusobacterium nucleatum adhesin involved in physical interaction and biofilm formation
with Streptococcus gordonii. *Microbiologyopen* **6** (2017).
https://doi.org:10.1002/mbo3.444

31 Han, Y. W. *et al.* Identification and characterization of a novel adhesin unique to oral
fusobacteria. *J Bacteriol* **187**, 5330-5340 (2005). https://doi.org:10.1128/JB.187.15.5330-
5340.2005

32 Umana, A. *et al.* Utilizing Whole *Fusobacterium* Genomes To Identify, Correct, and
Characterize Potential Virulence Protein Families. *J Bacteriol* **201** (2019).
https://doi.org:10.1128/JB.00273-19

33 Liu, B., Zheng, D., Zhou, S., Chen, L. & Yang, J. VFDB 2022: a general classification
scheme for bacterial virulence factors. *Nucleic Acids Res* **50**, D912-D917 (2022).
https://doi.org:10.1093/nar/gkab1107

34 Chen, L. C. *et al.* Alteration of gene expression in normal-appearing colon mucosa of
APC(min) mice and human cancer patients. *Cancer Res* **64**, 3694-3700 (2004).
https://doi.org:10.1158/0008-5472.CAN-03-3264

35 Rubie, C. *et al.* Correlation of IL-8 with induction, progression and metastatic potential
of colorectal cancer. *World J Gastroenterol* **13**, 4996-5002 (2007).
https://doi.org:10.3748/wjg.v13.i37.4996

36 Lee, Y. S. *et al.* Interleukin-8 and its receptor CXCR2 in the tumour microenvironment
promote colon cancer growth, progression and metastasis. *Br J Cancer* **106**, 1833-1841
(2012). https://doi.org:10.1038/bjc.2012.177

37 Casasanta, M. A. *et al. Fusobacterium nucleatum* host-cell binding and invasion induces
IL-8 and CXCL1 secretion that drives colorectal cancer cell migration. *Sci Signal* **13**
(2020). https://doi.org:10.1126/scisignal.aba9157

38 Lagier, J. C. *et al.* The rebirth of culture in microbiology through the example of
culturomics to study human gut microbiota. *Clin Microbiol Rev* **28**, 237-264 (2015).
https://doi.org:10.1128/CMR.00014-14

39 Roberts, R. J., Vincze, T., Posfai, J. & Macelis, D. REBASE: a database for DNA
restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res* **51**, D629-
D630 (2023). https://doi.org:10.1093/nar/gkac975

40 Aziz, R. K. *et al.* The RAST Server: rapid annotations using subsystems technology.
BMC Genomics **9**, 75 (2008). https://doi.org:10.1186/1471-2164-9-75
41 Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial genomes using
Subsystems Technology (RAST). *Nucleic Acids Res* **42**, D206-214 (2014).
https://doi.org:10.1093/nar/gkt1226
42 Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**, 1547-
1549 (2018). https://doi.org:10.1093/molbev/msy096
43 Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. *Nucleic Acids Res* **49**, W293-W296 (2021).
https://doi.org:10.1093/nar/gkab301
44 Yoon, S. H., Ha, S. M., Lim, J., Kwon, S. & Chun, J. A large-scale evaluation of
algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* **110**, 1281-
1286 (2017). https://doi.org:10.1007/s10482-017-0844-4
45 Blanco-Miguez, A. *et al.* Extending and improving metagenomic taxonomic profiling
with uncharacterized species using MetaPhlAn 4. *Nat Biotechnol* **41**, 1633-1644 (2023).
https://doi.org:10.1038/s41587-023-01688-w
46 Asnicar, F. *et al.* Precise phylogenetic analysis of microbial isolates and genomes from
metagenomes using PhyloPhlAn 3.0. *Nat Commun* **11**, 2500 (2020).
https://doi.org:10.1038/s41467-020-16366-7
816