Single-cell sequencing of tumor-associated macrophages in a *Drosophila* model

Dilan Khalili[#1], Mubasher Mohammed[#1], Martin Kunc[#1, 2], Johan Ankarklev[1], Ulrich
Theopold[1*]

[1]Department of Molecular Biosciences, The Wenner-Gren Institute, Stockholm University,
Stockholm, Sweden

[2]Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

* Correspondence:
Ulrich Theopold
uli.theopold@su.se

[#]**equal contribution**

Keywords: *Drosophila melanogaster,* macrophages, tumor model, hemocyte, single-cell
transcriptomics, scRNA-seq

Abstract

Introduction: Tumor-associated macrophages may act to either limit or promote tumor growth, yet
the molecular basis for either path is poorly characterized.

Methods: We use a larval *Drosophila* model that expresses a dominant-active version of the Ras-
oncogene (Ras[V12]) to study dysplastic growth during early tumor progression. We performed single-
cell RNA-sequencing of macrophage-like hemocytes to characterize these cells in tumor- compared to
wild type larvae. Hemocytes included manually extracted tumor-associated- as well as circulating cells.

Results and discussion: We identified 5 distinct hemocyte clusters. In addition to Ras[V12] larvae we
included a tumor model where the activation of effector caspases was inhibited, mimicking an
apoptosis-resistant setting. Circulating hemocytes from both tumor models differ qualitatively from
control wild-type cells – they display an enrichment for genes involved in cell division, which was
confirmed using proliferation assays. Split analysis of the tumor models further reveals that
proliferation is strongest in the caspase-deficient setting. Similarly, depending on the tumor model,
hemocytes that attach to tumors activate different sets of immune effectors – antimicrobial peptides
dominate the response against the tumor alone, while caspase inhibition induces a shift toward
members of proteolytic cascades. Finally, we provide evidence for transcript transfer between
hemocytes and possibly other tissues. Taken together, our data support the usefulness of *Drosophila*
to study the response against tumors at the organismic level.

1   Introduction

When cellular homeostasis is impaired, affected cells may limit the damage by inducing either cellular
arrest or cell death. Best characterized in this context is the tumor suppressor p53, which – depending
on the amount of DNA damage or other forms of cellular stress, induces cellular senescence, apoptosis,
or alternative forms of cell death (1–3). Collectively or individually, these responses will prevent
malfunction at the tissue/organ and organismic levels. Notably, this may also prevent the transition
from benign to more aggressive forms of tumor growth by eliminating or silencing damaged cells
locally in the tumor microenvironment (TME) and at an early stage of tumorigenesis (1). Cell cycle
arrest is usually expected to prevent this transition on its own while apoptosis involves the clearance
of apoptotic bodies by phagocytic neighboring cells or professional phagocytes such as macrophages,
which migrate into the TME (2,3).

While apoptosis thus acts as a tumor suppressor mechanism, tumor cells have been found to evade cell
death and in several tumors with poor prognosis, high levels of apoptosis are detected (4). This raises
the question of whether apoptosis, both when occurring naturally or as part of anti-cancer treatments
is fully beneficial (4).

Tumor progression depends on factors that act both locally in the tumor microenvironment as well as
on the communication with tissues in trans, which modulate physiological and immune responses
towards tumors. Key actors in the TME are tumor-associated macrophages (TAMs), which may affect
tumor progression both positively or negatively. In *vitro* activated macrophages have been roughly
classified as either classically activated (M1-type macrophages) or alternative (M2 macrophages, (5)).
M1 macrophages are pro-inflammatory and are involved in anti-viral, anti-bacterial, and anti-tumor
responses, whereas M2 macrophages are anti-inflammatory, contribute to anti-helminth and tissue
repair responses, and are considered pro-tumoral (6,7). Additional *in vitro* and *in vivo* data led to further
subdivision, in particular of M2 macrophages, and show that their activation state is only partially
reflected by the M1/M2 classification (6,8).

Facilitating the communication within the tumor microenvironment, which includes tumor cells, non-
tumor stroma, and TAMs, extracellular vesicles (EVs) have gained increasing attention in recent years
(4). EVs include (a) tumor-derived apoptotic fragments, (b) microparticles, which are shed from the
cell surface, and (c) exosomes, which are released through the fusion of multivesicular bodies with the
cell surface (9). EVs may have tumor and metastasis-promoting capacity both within the TME (4) and
systemically. Apoptotic bodies contain nuclear fragments (including DNA and primary transcripts)
while the content of microparticles and exosomes is derived from the cytosol and therefore contains
mature RNAs. Potential intermediaries of EV-mediated communication include proteins, lipids, and
different types of RNA, including long non-coding RNAs (lnc RNAs) (9,10).

Despite lacking adaptive immunity in a mammalian sense, insects possess a highly effective innate
immune system. This comprises both humoral and cellular elements. Humoral components are secreted
into the hemolymph – the insect equivalent of blood – primarily from the fat body, which fulfills
functions of both the mammalian fat body and liver. In *Drosophila melanogaster*, one of the major
models for insect immunity, the cellular branch comprises three classes of cells collectively called
hemocytes (11): (1) plasmatocytes, which are functionally equivalent to both mammalian macrophages
and white blood cells (2) crystal cells which contain prophenoloxidase, the precursor for a key enzyme
involved in antimicrobial and wound responses and (3) lamellocytes, which are absent in naive animals
but produced in response to wounding and invasion by large intruders such as parasitoid eggs. Recent
hemocyte transcriptome profiling at the single-cell level revealed a much higher diversity, in particular
among plasmatocytes (11–14).

The interplay between fly tumors and hemocytes was first studied in pioneering work by Pastor Pareja et al (15), where it was shown that upon recognition of damage to the basement membrane, hemocytes are recruited to tumor tissue with tumor-limiting effects. Conversely when in subsequent work tumors were induced in a background expressing a dominant-active version of the Ras oncogene (Ras^{V12}) tumor-associated hemocytes (TAHs) were instead involved in a positive feedback loop that involved JNK signaling and promoted tumor growth (16,17). Thus, similar to what is found in mammals, the effects of TAHs appear to be in many cases tissue- and tumor stage-dependent. In previous work, we found that expression of dominant active Ras^{V12} in larval *Drosophila* salivary glands (SGs) induced ductal hypertrophy (18) similar to what is observed in ductal tumors in humans. In SGs, this led to (1) the loss of cellular integrity; (2) nuclear disintegration and caspase activation (3) loss of the SG lumen and of secretory activity (4) damage to the basement membrane (5) induction of fibrotic lesions including activation of the flies' coagulation system and (6) recruitment of TAHs (18–20). Despite the presence of hallmarks of apoptosis and activation of JNK signaling, SG cells were not eliminated (18,20) although SG cell fragments were released into the hemolymph (18). Surprisingly, forced expression of the antimicrobial peptide Drosomycin (Drs) across whole SGs reverted the majority of tumor-associated phenotypes through negative regulation of JNK signaling (19,20).

Here, to identify genes that are differentially expressed in hemocytes from tumor larvae, we profiled the transcriptome of circulating single-cell hemocytes from wild-type and tumor larvae, as well as from tumor-associated hemocytes (TAHs), which were extracted manually from tumorous salivary glands (Fig 1A). Since we had previously observed strong activation of caspases in Ras^{V12}-expressing SGs (20), and in light of the bi-edged nature of caspase activation and apoptosis in cancer (see above), we included larvae where effector caspases were inhibited by the expression of the specific inhibitor p35.

2   Material and Methods

2.1   Fly strains and sample preparation

w^{1118}, *Beadex^{MS1096}-Gal4* (Referred here as *Bx*: 8860/Bl), *Beadex^{MS1096}-Gal4; EaterDsRed, w^{1118}*, *w^{1118};p35* (5072/Bl), *w^{1118};;UAS-Ras^{V12}*(4847/Bl), *w^{1118};p35;UAS-Ras^{V12}* flies were used in the experiments. Flies were cultured in a 25°C, 12-h dark/light cycle room. Female virgins were collected for 5 days and crossed with respective males on day 7. Progeny larvae were kept as described in (20). Twenty to thirty salivary gland pairs were fixed in 4% PFA for 20 min. Samples for extracellular staining were washed 3 x 10 min with 1x phosphate-buffered saline (PBS).

2.2   Immunohistochemistry

Antibodies against P1 (Plasmatocytes, 1:50), L1 (Lamellocytes, 1:50), C4 (Crystal cells, 1:50) were incubated for 1 h at room temperature in PBS and subsequently, washed 3 x 10 min in PBS. The samples were incubated with secondary antibody Anti-Mouse-546 (1:500, ThermoFisher #A11030), DAPI (1:500, Sigma-Aldrich D9542) for 1 hour at room temperature and washed 3 x 10 min with PBS before mounting in Fluoromount-G (ThermoFisher, 00-4958-02).

2.3   Proliferation assay

Drosophila larva at 120 hours AED was bled into 10 µl of PBS and the sample was incubated for 5 min at room temperature. The sample was fixed in 4% PFA for 20 min and washed 3 x 5 min with PBS. The sample was stained with antibody against pH3 (phosphorylated histone 3, 1:800, Sigma-

Aldrich 06-570) and H2 (Hemese, 1:5 kindly provided by Istvan Ando, Szeged) overnight in 4°C. After
incubation, the sample was washed 3 x 5 min with PBS and stained with secondary antibodies Anti-
Rabbit-568 (1:500, ThermoFisher A-21069), Anti-Mouse-488 (1:250, ThermoFisher A-11001) and
DAPI (1:500, Sigma-Aldrich D9542) for 2 h in room temperature. Subsequently, the sample was
washed 3 x 5 min with PBS and mounted in Fluoromount-G (ThermoFisher, 00-4958-02). 15 samples
per genotype were prepared and photographed using a tile scan at a confocal microscope Zeiss LSM780
(Zeiss, Germany), and the count of cells was evaluated in ImageJ (version 1.53t). Statistics and graph
plotting was performed in Prism software (GraphPad Software, 9.3.0, USA).

126 2.4 Image acquisition and analysis

127 Whole salivary glands were photographed using Zeiss Axioscope II (Zeiss, Germany) microscope and
128 images were exported as .tiff files. The intensity was measured using ImageJ (version 1.53t) and
129 representative pictures were obtained from a confocal Zeiss LSM780 (Zeiss, Germany) microscope.
130 Statistics were performed in Prism software (GraphPad Software, 9.3.0, USA).

131 2.5 Micro-manipulation of attached hemocytes

Ten pairs of salivary glands per genotype were dissected into individual droplets of 200 µl PBS. A
fluorescent microscope Leica DMi8 (Leica, Germany) was used to locate the attached hemocytes at
the surface of the salivary gland. The hemocytes were collected using micromanipulator TransferMan
NK2 (Eppendorf, Germany) with ø 7 µm collection capillary and separated by piezo-vibrator
PiezoXpert (Eppendorf, Germany). Single-cell state hemocytes were individually transferred to 2.3 µl
of lysis buffer (38) and each sample library was prepared as described below.

138 2.6 cDNA synthesis and library preparation for attached hemocytes

cDNA libraries of single attached hemocytes were generated using a modified version of the Smart-
seq2 protocol (38). In short, cDNA synthesis was performed using universal primers, and PCR
amplification was carried out over 24 cycles. cDNA products were subsequently purified using CA
beads (Sigma; catalog no. 81260) for size selection using 8.8% polyethylene glycol 6000 (PEG 6000)
to exclude primer-dimers and nonspecific amplicons with sizes less than 150 bp. Combinatorial
indexing via tagmentation was carried out in 96-well plates using 200 pg (measured in a Qubit
fluorometer) of amplified cDNA, for a final volume of 10 mL/well. cDNA fragmentation using Tn5
transposase was carried out for 20 min on ice using the Illumina Nextra XT DNA sample preparation
kit. Ligation and amplification of adaptors were carried out over 15 cycles in a final volume of 25
mL/well. Primer indices were used in the reaction from Illumina (Nextera index primers i7 and i5,
catalog no. FC-131-1001). Tagmented and barcoded amplicons were then purified using CA beads for
size selection. Quality control and fragment size distribution of the cDNA libraries were performed on
a Bioanalyzer with the Agilent high-sensitivity DNA chip (catalog no. 5067-4626). Concentrations of
each sample of cDNA libraries were measured on a PicoGreen 96-well plate NucleoScan fluorometer
using a high-sensitivity double-stranded DNA (dsDNA) (HS assay kit; catalog no. Q32851). To
perform library dilutions, the average fragment sizes of all cDNA libraries were measured for a final
concentration of 2 nM in each sample. Finally, cDNA libraries were pooled and sequenced using
Illumina NextSeq with 75-bp paired-end reads.

157 2.7 Cell sorting and cDNA library preparation for circulating hemocytes

Ten larvae per genotype were bled into 500 µl of PBS and the cells were sorted. The sorting was
performed with a MoFlo Astrios EQ (Beckman Coulter, USA) cell sorter using a 488 and 532 nm laser

for excitation, 100 µm nozzle, sheath pressure of 25 psi, and 0.1 µm sterile-filtered 1 x PBS as sheath
fluid. Flow sorting data was interpreted and displayed using the associated software, Summit v 6.3.1.

To test the precision of the adjustments made to center the drop in each well, a colorimetric test
mimicking the sort was done based on (39). A 1.5 mg/µl solution of HRP (cat no 31490, ThermoFisher
Scientific) with 1 drop of flow check beads (Beckman Coulter, USA) was sorted into each well of an
Eppendorf 384-well plate (Cat no 34028, ThermoFisher Scientific). A color change after sorting
indicated that the drop hit the sort buffer and that the precision was adequate.

Single hemocytes were sorted directly into a 384-well plate containing 2.3 µl of lysis buffer (Eppendorf
twin.tecTM PCR plates) using a CyCloneTM robotic arm and at highly stringent single cell sort settings
(single mode, 0.5 drop envelope) and cDNA libraries were generated by the Eukaryotic Single Cell
Genomics Facility at SciLifeLab, Stockholm using a slightly modified version of Smart-seq2 as
previously described (38), but where we used 20 cycles for cDNA amplification. The plate and sample
holder were always kept at 4 ºC during the sort. After the sort, the plates were immediately spun down
and put on dry ice.

174 2.8   Single-cell RNA sequencing

175 Single-cell libraries were sequenced at the National Genomics Infrastructure, SciLifeLab Stockholm,
176 using the HiSeq2500 platform (Illumina) for 56 bps single-end sequencing. We sequenced a total of
177 463 (negative controls n=2, per plate).

178 2.9   Mapping, annotation and filtering of low-quality cells

A reference genome *Drosophila melanogaster* (Dm6 v r6.37) was indexed and raw fastq files were
used in mapping to the genome using STAR v 2.7.2 (40), and gene expression was measured using
featureCounts v 2.0.0 (41), using default settings. expression matrix was filtered to obtain high-quality
cells using the following criteria: cells with >5% mitochondrial transcripts (stressed/dead/dying cells),
<200 genes (low-quality cells), those expressing more than 4000 features (genes) (potential doublets
or triplets) were removed in each replicate and the remaining cells were subjected to subsequent
computational analysis.

186 2.10   Normalization, dimensionality reduction and clustering

The main computational analysis of read-count matrices was performed using the Seurat package (v
4.0.3) (42) in R (v 4.1.0). The complete R workflow can be assessed and reproduced in the R markdown
(see code availability section). we used the default processing pipeline
https://satijalab.org/seurat/v3.2/pbmc3k_tutorial.html. First, count matrices and metadata were loaded.
A mitochondrial gene count above 10 % was filtered. Quality filtering was performed and cells with a
minimum of 200 genes expressed were kept for further processing. Subsequently, reads were
normalized for sequencing depth using the "NormalizeData" function in the Seurat toolkit, selecting
the top 2000 variable genes. Thereafter, dimensionality reduction was performed using PCA
computing the first 50 PCs. The first 10 PCs from the analysis were then subjected to shared-nearest-
neighbor (SNN) inspired graph-based clustering via the "FindNeighbors" and "FindClusters"
functions. For modularity optimization, the Louvain algorithm was used, and clustering was performed
at a resolution of 0.8 for clustering granularity, resulting in 9 clusters. After clustering, a UMAP
dimensionality reduction was performed using the first 10 dimensions of the PCA.

200 2.11   Differential gene expression analysis (DGEA)

Differential gene expression analysis (DGEA) of genes in identified clusters was performed using the
function "FindAllMarkers" from the Seurat package (v. 4.0.3). Following the default option of the
method, differentially expressed genes for each cluster were identified using a non-parametric
Wilcoxon rank sum test. Differentially expressed genes in a cluster were defined by setting initial
thresholds above a logarithmic fold-change of 0.5 and being present in at least 25% of the cells
belonging to the same cluster. Representative marker genes with an adjusted *p*-value below 0.05 for
each cluster were further selected. *p*-values were adjusted using a Bonferroni correction including all
genes in the dataset. To find representative marker genes with elevated expression in comparison to
the remaining clusters, only positive log fold-changes were considered. For individual analyses such
as gene enrichment analysis (see "Gene set enrichment analysis (gsea)"), threshold values for
differential gene expressions were modified and will be described in detail in the respective sections
of the materials and methods and results. To identify DEGs between specific clusters of interest, the
"FindMarkers" function in Seurat was used and the identities were set to the respective clusters of
interest. The same thresholds as stated above were used to define DEGs.

2.12 Biological pathways and GSEA

To track tests for top functional class enrichment among the global clusters, we used "ClusterProfiler"
package v 3.16 (43) tool to conclude the enriched ontology terms as previously mentioned specifying
the database "Org.Dm.db" to calculate the top 5 biological pathway enrichment. The gene set
enrichment analysis (GSEA) was performed on top differentially expressed genes over the identified
clusters in regards to gene expression profiles "Log2FC" as input in cluster profiler v 3.18.1 and
ggupset package v 0.3.1 with a *P value* cutoff of 0.05, minGSSsize of 3, maxGSSize of 800, and
scoreType of "pos" to estimate for biological process ontology across clusters.

2.13 Computational summary

The read alignment and gene count matrix generation were performed as previously described
(Methods). The single-cell gene count matrix cells with fewer than 250 UMIs, more than 10,000 UMIs,
reads mapping to more than 7000 genes, or more than 10% of read counts mapping to ribosomal genes
were excluded. Each single cell transcriptome was mapped to its original time window from which it
was extracted by using the RT barcode. We performed standard processing of the data split by
experiments as recommended by Seurat v4 documentation including NormalizeData,
FindVariableFeatures (with the method set to 'vst'), ScaleData, RunPCA, RunUMAP (with dims set
to 1:50 and n.components to 2), FindNeighbors (with reduction set to UMAP and dims set to 1:2),
FindClusters. This version uses the first 10 principal components, along with pN of 0.2 and pK of
0.005. Dimensionality reduction, clustering, and identifying cluster-specific marker genes. The
standard Seurat processing pipeline, as described in the previous section, was performed on each non-
overlapping inferred age window separately. We found that the default Seurat clustering resolution
parameter did not capture the dynamics of the presence of different cell types. For each batch, we
clustered the data with a variety of resolutions (from 0.1 to 1.5 in increments of 0.3) and then computed
the within-cluster sum of squares (WSS). Visualizing WSS across increasing resolutions we chose the
resolution in which there was a visible plateau in the decrease of WSS. Finally, we integrated and
clustered the data and from these clusters, we used Seurat's FindMarkers to iteratively loop through all
clusters and identify marker genes.

2.14 RNA velocity and lineage interference

To interpret the global transcriptional progression of hemocytes and their cell fate decision, we
established the cell continuum of cell differentiation, data layers of unspliced and spliced mRNA for

the entire data generated with Velocyto CLI (v.0.17.17) according to the CLI usage guide (Velocyto
run-smartseq2). The output loom files were combined using "loompy". The merged loom file was
imported into the scVelo package (v1.0.6) (44,45). The unspliced and spliced mRNA counts of cells
from clusters C0-C4 were extracted. We used the "merged.utilis" function in the scVelo pipeline,
where cells with low pre-mRNA counts were removed as part of the filtering. In short, the gene-specific
velocities are obtained by fitting a ratio between unspliced and spliced mRNA abundances and then
computing how the observed abundances change from those observed in a steady state. The ratio of
'spliced', 'unspliced', and 'ambiguous' transcripts were calculated and data were pre-processed using
functions for detection of minimum count number, filtering and normalization using
"scv.pp.filter_and_normalise" and followed by "scv.pp.moments" function. The gene-specific
velocities were then calculated using "scv.tl.velocity" with mode set to "deterministic" and
"scv.tl.velocity_graph" function to generate velocity graph, and visualization using
"scv.pl.velocity_graph" function. In addition, we used the "scv.tl.recover_latent_time" function to
infer a shared latent time from splicing dynamics and plotted the genes along the time axis.

2.15 Data availability

The raw processed data generated for this study have been deposited on Zenodo repository and can be
accessed via [https://zenodo.org/deposit/7997643] and interactively on [https://mubasher-
mohammed.shinyapps.io/Sc-drosophila/]. The custom code scripts used to analyze data for this study
are available at [https://github.com/ANKARKLEVLAB/Single-cell].

3   Results

3.1   Single-cell profiling of tumor-associated hemocytes reveals transcriptional heterogeneity

We initially characterized the population of tumor-associated hemocytes (TAHs) that attach to Ras[V12]
salivary glands (SGs) and found they consist mainly of plasmatocytes and the occasional lamellocyte
(**Figure 1A-B**). To identify genes that are differentially regulated in hemocytes (DEGs) from tumor
larvae we profiled the transcriptome of single circulating hemocytes from wild-type and tumor larvae
as well as from tumor-associated hemocytes (TAHs), which were extracted manually from tumorous
SGs. Additionally, to characterize the contribution of effector caspases to tumor progression we
analyzed larvae in which the effector caspase inhibitor p35 was co-expressed with Ras[V12]: p35;Ras.
We had previously shown that – similar to Drs - expression of p35 in Ras-SGs restores the spherical
nuclear shape that is disturbed upon sole Ras-expression although SGs from both combinations show
the same size increase as Ras[V12] larvae (20) and (**Figure 2A-D, quantified in E**). In contrast to Drs
(20), co-expression of p35 still led to hemocyte recruitment (**Figure 2A'-D' quantified in F**). This
allowed us to compare tumor-associated hemocytes (TAHs) both between wild-type and tumor larvae
as well as in settings with and without active caspases downstream of JNK-signaling. TAHs as well as
circulating hemocytes were selected for expression of the plasmatocyte-specific marker Eater (**Figure
3A**, see Material & Methods).

Collectively, hemocyte transcriptomes displayed additional plasmatocyte markers indicating
successful purification of these macrophage-related cells (**Figure 3B-H**). Plasmatocyte markers
include Nimrod C1 (NimC1), serpent (srp), Hemolectin (Hml), and spätzle (spz), several of which have
been used as pan-plasmatocyte markers and more recently as markers for specific plasmatocyte
subpopulations (21). As expected, the crystal cells and lamellocytes markers lozenge (lz) and Atilla,

respectively; were not detected (**Figure 3C-D**). Crystal cell prophenoloxidase 1 (PPO1) was detected
in a few cells (**Figure 3H**) see also below.

Clustering analysis revealed true variation between the single cell transcriptomes for the identified 5
clusters (at p<0.05 and a difference in expression >2 if not stated otherwise; (**Figure 4A-C, middle**
panel), three of which included tumor cells (clusters C0, C2 and C3 (**Figure 4C-D**). Cluster C2
overlaps largely with TAHs (**Figure 4A**). Gene Ontology analysis for biological processes of up-
regulated genes in the clusters highlighted several gene sets that overlapped between clusters but also
some cluster-specific sets (**Figure 4E**). The latter category included genes involved in cell motility
(cluster C1), translation (C0) protein folding (C4) and immune response genes for cluster 2 (attached
cells) and several categories specific for cell division (C3, circulating cells from both tumor models).
Taken together, despite the technically limited number of cells compared to other studies (21), we
identified subpopulations of plasmatocytes whose signatures differed significantly depending on
genotype (tumor versus wild type) and cell status (attached versus circulating).

3.2   The expression of signature genes for the different clusters differs between tumor models

To obtain more detailed insight into the gene categories that specified the clusters, we identified the
genes whose expression significantly specified the clusters (**Tables S1-S5**). In parallel, we compared
expression strengths between genotypes focusing on the three populations that comprised cells from
the tumor models (clusters 0, 2, and 3 in **Figure 4D and Figure 5A-C**). For attached hemocytes (C2),
the list of the most strongly expressed genes varied between the two tumor models (**Figure 5A**). To
follow this up we scanned cluster 2-specific genes for gene-set enrichment and identified two enriched
categories: (1) pupal adhesion and (2) immune response (**Figure S1A-B**). Pupal adhesion was due to
the presence of genes that code for salivary gland secretions (**Figure S1B**). Since these genes are hardly
expressed outside salivary glands (FlyAtlas: (22)), we interpret their presence in hemocytes as being
passenger transcripts (23), i.e., transcripts that have been taken up by TAHs by way of cellular
fragments released from the glands. Alternatively, SG transcripts may have been co-purified together
with TAHs during extraction. This latter explanation implies that expression of p35 partially prevented
the release of SG fragments from Ras^{V12} SGs in line with its anti-apoptotic function. The second
category (immune genes) contains genes that are most strongly (although not mutually exclusively)
expressed in either tumor model (**Figure S1A**). Taken together, we show that hemocytes in both tumor
models differ qualitatively from hemocytes in control larvae and that inhibition of effector caspases in
tumors reveals additional quantitative differences.

3.3   The response towards tumors differs between the two models

Generally, TAHs in the Ras model show strong expression of AMPs including Dpt(Diptericin)A and
B, Att(Attacin)C, and Cec(Cecropin)C. In contrast p35;Ras^{V12} TAHs show higher levels of all three
phenoloxidases PPO1-3 (**Figure S1A**). This is in line with the previously identified function of the
TGF-like protein dawdle (*daw*), which promotes AMP activation (24) and indeed, *daw* clusters with
AMP expression in our hands (**Figure S1A**). A second TGF member (*dpp*) with immune-regulatory
function (24) is less expressed in tumors compared to wild-type hemocytes. Similarly, the M1 marker
iNOS is more strongly expressed in p35;Ras^{V12} TAHs while the M2 marker Arginase (Arg) peaks in
Ras TAHs.

For both clusters 0 and 3 (**Figure 5B-C**), the same set of genes was more strongly induced in circulating
tumor compared with circulating wild-type hemocytes although the majority showed stronger
expression when effector caspases were repressed (p35;Ras^{V12}) compared to the tumor-only (Ras)
model. Gene set enrichment analysis fully confirmed the enrichment for genes involved in cell division

(**Figure 4E**) in cluster 3 (circulating cells from both tumor models) but failed to deliver significant
returns for the category "Biological process" for cluster 0. In line, the quantification of hemocyte
proliferation fully confirmed an increase in cell divisions in both tumor models including the additional
increase in the p35;Ras^{V12} model (**Figure 5D**). This may contribute more to the differentiation of TAHs
rather than circulating cells, which show similar counts (Hauling et al., 2014).

Finally, when searching for pathways compatible with expression in the five clusters using the
Reactome database (https://reactome.org), an enrichment for genes involved in neutrophil
degranulation was found for cluster 0 (**Figure S1C**), which when using a split analysis also indicated
that this pathway was more strongly expressed in p35;Ras^{V12} hemocytes. Pseudotime analysis using
RNA Velocity (**Figure 6A-D**) indicated that TAHs and circulating hemocytes from both tumor models
follow distinct pathways although due to the restricted number of TAHs, more rare intermediate
population may have been missed.

3.4   Transcript transfer between hemocytes and other tissues

Recently, we and others have characterized the activation of crystal cells and the subsequent release of
leaderless PPO2 into the extracellular environment via cell rupture (25,26). Functionally, crystal cell
activation bears similarities to pyroptotic cell death in mammals, including its dependence on caspase
activity, which can be inhibited by p35 (27). While our findings explained how PPO2 was released
into the hemolymph, the mechanism for secretion of PPO1, which also lacks a signal peptide remained
obscure. Since we found that PPO2 was enriched in TAHs (**Figure 7A, S1A**), we wondered which
secretion mechanism led to their spread to other plasmatocytes. In line with crystal cell rupture and
similar to passenger transcripts from SGs, the mRNA encoding PPO2 was detected in plasmatocytes
both as a mature (spliced) form and as a non-spliced (nuclear) immature mRNA (**Figure 7C, D**). This
indicates that phagocytic hemocytes had access to both the nuclear and the cytosolic fraction of
ruptured crystal cells. In contrast, PPO1 transcripts were only detected in their mature form (**Figure
7B**). We conclude that PPO2 transcripts in TAHs originate either from pyroptotic crystal cells or TAHs
which have acquired crystal cell characteristics, while PPO1 mRNA originates from live crystal cells.

TAHs also contained transcripts that – according to FlyAtlas – are strongly expressed in fat bodies.
When checked for the presence of spliced and unspliced versions we found most of them displayed
both, indicating genuine expression in hemocytes, which was strengthened by the data from Tattikota
et al who detected them in hemocytes too (21).

In addition to PPO1, 3 transcripts (UDP-galactosyltransferase activity, CG 13962, and the transcription
factor Relish) were detected only in their mature form indicating a possible origin outside TAHs.

4   Discussion

To our best knowledge, we provide the first transcript profile of tumor-associated hemocytes (TAHs)
– the invertebrate equivalent of mammalian tumor-associated macrophages (TAMs). Relying on a
molecularly induced early stage of tumor progression that affects the salivary glands (18,20), we find
that TAHs display some features of mammalian M2-like macrophages, which have been implicated in
regenerative processes. These include the presence of members of the chitinase-like proteins (IDGFs
in *Drosophila*), which are amongst the most abundant proteins in activated macrophages and are used
as markers for M2 macrophages (28) (**Figure S1C**). Notably, TAHs show a clear signature of immune
activation, which includes several antimicrobial peptides as well as members of clotting systems

including IDGF3, Fondue, and phenoloxidases (**Figure S1A**). Although both effector branches are activated in the TAHs from both tumor models, there are differences (**Figure S1A**): AMPs appear to dominate the TAH response in Ras-SGs whereas clotting factors are more strongly expressed in p35;Ras^{V12} SGs. AMP induction may be in line with the presence of SG (passenger) transcripts in the Ras-alone model **(Figure 8, right part of SGs)** and may serve to degrade tumor fragments during efferocytosis including cytosolic and nuclear parts. In contrast, inhibition of caspases appears to activate immune reactions that are more akin to the formation of mammalian granulomas (**Figure 8 left part).** Bifurcation in hemocyte differentiation is reminiscent of the division of labor between *Drosophila* hemocytes shown previously which depend on two members of the TGF family (the BMP-like member Dpp and the Activin-like member Dawdle, (24)). In line with Dpp's role in suppressing antimicrobial responses, we find that two receptors for BMP-like TGFs (thickveins; tkv and saxophone; sax) are strongly expressed in p35:Ras^{V12} hemocytes but not in Ras or control hemocytes (**Figure S2**). While the difference in responses between Ras larvae and larvae that co-express the caspase inhibitor p35 may be explained by a lack of apoptosis downstream of caspases, we prefer alternative explanations that include non-apoptotic form of cell death or non-apoptotic functions of caspases (29).

Taken together, our results identify targets for modifier screens that address the contribution of TAH-DEGs to tumor development using both classical and molecularly induced tumor mutants.

Additionally, our work points toward a potential communication network that links TAHs, SG tumors, and non-tumor tissues (**Figure 8**). This likely includes efferocytosis of apoptotic bodies derived from hyperplastic SGs (18), leading to an extended presence of SG transcripts (passenger transcripts) in TAHs before their degradation. While most of the DEGs in TAHs are genuine TAH transcripts, we find exceptions. These include two prophenoloxidases (26), which are known to be exclusively expressed in crystal cells (30) and not in the plasmatocytes we sequenced. Notably, transcripts of potential non-plasmatocyte origin also include the NF-kappaB factor Relish, a key regulator of the Imd pathway (31). While the mechanism for this cell's non-autonomous presence of transcripts in TAHs is unknown, we hypothesize that exosomes may be likely candidates (**Figure 8 solid arrows**). Alternatively, plasmatocytes may transdifferentiate and express transcripts that are more specific for crystal cells and lamellocytes (32,33) (**Figure 8, dashed arrows).** Supporting an exosome origin **(Figure 8)**, two long non-coding RNAs (CR 34335 and R 40469) we found in TAHs have been identified among the 10 most abundant non-ribosomal RNAs in exosomes released from *Drosophila* cell lines (34). Additionally, in the same study, *Arc1* was shown to be amongst the most abundant mRNAs in exosomes from one of the two cell lines used (34). Notably, in the flies' nervous system, Arc1 is involved in the formation of capsid-like vesicles, which also contain Arc1 transcripts that are recruited through the binding of Arc1 protein to the 3' end of the *Arc1* transcript (35). Both Arc1 and its human equivalent derive from retroviral Gag proteins (35–37) and mediate neuronal plasticity. A function for *Drosophila* Arc1 in immunity has so far not been suggested although it is expressed in non-neural tissues (FlyAtlas: (22)) and we find it is enriched in TAHs **(Figure 5)**. Of note, similar to tumor-associated macrophages, extracellular vesicles have been shown to display pro-tumor potential. Taken together, our findings on macrophage-like cells from an invertebrate provide targets that may turn out useful to steer tumor therapy even in humans (2).

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6 Author Contribution

UT conceived the study. **JA** and **UT** supervised the study. **DK** experimental design, investigation, methodology and visualization **MK** performed immunohistochemistry analyses and proliferation assay as well as *Drosophila* genetics. **JA** and **DK** performed sorting and scSeq. **MM** performed mapping and computational analyses of the data. **MM**, **DK** and **MK** generated figures. **JA** supervised the computational analysis. **UT, MM** wrote the manuscript. All authors read and reviewed the manuscript.

7 Funding

This work was supported by grants from the Swedish Cancer Foundation (CAN 2015-546 to UT), the Wenner-Gren Foundation (UPD2020-0094 and UPD2021-0095 to MK), the Swedish Research Council (VR 2016-04077 and VR 2021-04841 to UT), the Carl-Tryggers Foundation (CTS 21:1263 to UT), the Swedish Research Council (VR 2021-05057 to JA), and the Swedish Society for Medical Research (SSMF) to JA.

8 Acknowledgements

We thank the Microbial Single Cell Genomics and the Eukaryotic Single Cell Genomics facilities at Science for Life Laboratory (SciLifeLab), Sweden, for cell sorting as well as smart-seq2 library preparations, respectively, and the National Genomics Infrastructure, SciLifeLab for sequencing. The computations were performed using resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX). We also thank Jiwon Shim (Hanyang University, Korea) for useful comments on the manuscript.

9 Data availability statement

The raw processed data generated for this study have been deposited on Zenodo repository and can be accessed via [https://zenodo.org/deposit/7997643] and interactively on [https://mubasher-mohammed.shinyapps.io/Sc-drosophila/]. The custom code scripts used to analyze data for this study are available at [https://github.com/ANKARKLEVLAB/Single-cell].

10 References

1. Dhokia V, Moss JAY, Macip S, Fox JL. At the Crossroads of Life and Death: The Proteins That Influence Cell Fate Decisions. Cancers. 2022 May 31;14(11):2745.

2. Gregory CD, Paterson M. An apoptosis-driven "onco-regenerative niche": roles of tumour-associated macrophages and extracellular vesicles. Philos Trans R Soc Lond B Biol Sci. 2018 Jan 5;373(1737):20170003.

3. Lees A, Sessler T, McDade S. Dying to Survive-The p53 Paradox. Cancers. 2021 Jun
29;13(13):3257.

4. Morana O, Wood W, Gregory CD. The Apoptosis Paradox in Cancer. Int J Mol Sci. 2022 Jan
25;23(3):1328.

5. Viola A, Munari F, Sánchez-Rodríguez R, Scolaro T, Castegna A. The Metabolic Signature of
Macrophage Responses. Front Immunol. 2019;10:1462.

6. Cassetta L, Pollard JW. Tumor-associated macrophages. Curr Biol CB. 2020 Mar
23;30(6):R246–8.

7. Kelly B, O'Neill LAJ. Metabolic reprogramming in macrophages and dendritic cells in innate
immunity. Cell Res. 2015 Jul;25(7):771–84.

8. Kolliniati O, Ieronymaki E, Vergadi E, Tsatsanis C. Metabolic Regulation of Macrophage
Activation. J Innate Immun. 2022;14(1):51–68.

9. Soltész B, Buglyó G, Németh N, Szilágyi M, Pös O, Szemes T, et al. The Role of Exosomes in
Cancer Progression. Int J Mol Sci. 2021 Dec 21;23(1):8.

10. Zhang WL, Liu Y, Jiang J, Tang YJ, Tang YL, Liang XH. Extracellular vesicle long non-coding
RNA-mediated crosstalk in the tumor microenvironment: Tiny molecules, huge roles. Cancer
Sci. 2020 Aug;111(8):2726–35.

11. Bilder D, Ong K, Hsi TC, Adiga K, Kim J. Tumour-host interactions through the lens of
Drosophila. Nat Rev Cancer. 2021 Nov;21(11):687–700.

12. Cho B, Yoon SH, Lee D, Koranteng F, Tattikota SG, Cha N, et al. Single-cell transcriptome
maps of myeloid blood cell lineages in Drosophila. Nat Commun. 2020 Sep 8;11(1):4483.

13. Fu Y, Huang X, Zhang P, van de Leemput J, Han Z. Single-cell RNA sequencing identifies novel
cell types in Drosophila blood. J Genet Genomics Yi Chuan Xue Bao. 2020 Apr 20;47(4):175–
86.

14. Leitão AB, Arunkumar R, Day JP, Geldman EM, Morin-Poulard I, Crozatier M, et al.
Constitutive activation of cellular immunity underlies the evolution of resistance to infection in
Drosophila. eLife. 2020 Dec 24;9:e59095.

15. Pastor-Pareja JC, Wu M, Xu T. An innate immune response of blood cells to tumors and tissue
damage in Drosophila. Dis Model Mech. 2008;1(2–3):144–54; discussion 153.

16. Fogarty CE, Diwanji N, Lindblad JL, Tare M, Amcheslavsky A, Makhijani K, et al. Extracellular
Reactive Oxygen Species Drive Apoptosis-Induced Proliferation via Drosophila Macrophages.
Curr Biol CB. 2016 Mar 7;26(5):575–84.

17. Fogarty CE, Bergmann A. Killers creating new life: caspases drive apoptosis-induced
proliferation in tissue repair and disease. Cell Death Differ. 2017 Aug;24(8):1390–400.

18. Hauling T, Krautz R, Markus R, Volkenhoff A, Kucerova L, Theopold U. A Drosophila immune
response against Ras-induced overgrowth. Biol Open. 2014 Apr 15;3(4):250–60.

19. Khalili D, Kalcher C, Baumgartner S, Theopold U. Anti-Fibrotic Activity of an Antimicrobial
Peptide in a Drosophila Model. J Innate Immun. 2021;13(6):376–90.

20. Krautz R, Khalili D, Theopold U. Tissue-autonomous immune response regulates stress signaling
during hypertrophy. eLife. 2020 Dec 30;9:e64919.

21. Tattikota SG, Cho B, Liu Y, Hu Y, Barrera V, Steinbaugh MJ, et al. A single-cell survey of
Drosophila blood. eLife. 2020 May 12;9:e54818.

22. Leader DP, Krause SA, Pandit A, Davies SA, Dow JAT. FlyAtlas 2: a new version of the
Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.
Nucleic Acids Res. 2018 Jan 4;46(D1):D809–15.

23. Lantz C, Radmanesh B, Liu E, Thorp EB, Lin J. Single-cell RNA sequencing uncovers
heterogenous transcriptional signatures in macrophages during efferocytosis. Sci Rep. 2020 Aug
31;10(1):14333.

24. Clark RI, Tan SWS, Péan CB, Roostalu U, Vivancos V, Bronda K, et al. MEF2 is an in vivo
immune-metabolic switch. Cell. 2013 Oct 10;155(2):435–47.

25. Myers AL, Harris CM, Choe KM, Brennan CA. Inflammatory production of reactive oxygen
species by Drosophila hemocytes activates cellular immune defenses. Biochem Biophys Res
Commun. 2018 Nov 2;505(3):726–32.

26. Schmid MR, Dziedziech A, Arefin B, Kienzle T, Wang Z, Akhter M, et al. Insect hemolymph
coagulation: Kinetics of classically and non-classically secreted clotting factors. Insect Biochem
Mol Biol. 2019 Jun;109:63–71.

27. Dziedziech A, Theopold U. Proto-pyroptosis: An Ancestral Origin for Mammalian Inflammatory
Cell Death Mechanism in Drosophila melanogaster. J Mol Biol. 2022 Feb 28;434(4):167333.

28. Kang Q, Li L, Pang Y, Zhu W, Meng L. An update on Ym1 and its immunoregulatory role in
diseases. Front Immunol. 2022;13:891220.

29. Xu DC, Wang L, Yamada KM, Baena-Lopez LA. Non-apoptotic activation of Drosophila
caspase-2/9 modulates JNK signaling, the tumor microenvironment, and growth of wound-like
tumors. Cell Rep. 2022 Apr 19;39(3):110718.

30. Dudzic JP, Kondo S, Ueda R, Bergman CM, Lemaitre B. Drosophila innate immunity: regional
and functional specialization of prophenoloxidases. BMC Biol. 2015 Oct 1;13:81.

31. Hedengren M, Asling B, Dushay MS, Ando I, Ekengren S, Wihlborg M, et al. Relish, a central
factor in the control of humoral but not cellular immunity in Drosophila. Mol Cell. 1999
Nov;4(5):827–37.

32. Anderl I, Vesala L, Ihalainen TO, Vanha-Aho LM, Andó I, Rämet M, et al. Transdifferentiation
and Proliferation in Two Distinct Hemocyte Lineages in Drosophila melanogaster Larvae after
Wasp Infection. PLoS Pathog. 2016 Jul;12(7):e1005746.

33. Leitão AB, Sucena É. Drosophila sessile hemocyte clusters are true hematopoietic tissues that

regulate larval blood cell differentiation. eLife. 2015 Feb 4;4:e06166.

34. Lefebvre FA, Benoit Bouvrette LP, Perras L, Blanchet-Cohen A, Garnier D, Rak J, et al.
Comparative transcriptomic analysis of human and Drosophila extracellular vesicles. Sci Rep.
2016 Jun 10;6:27680.

35. Ashley J, Cordy B, Lucia D, Fradkin LG, Budnik V, Thomson T. Retrovirus-like Gag Protein
Arc1 Binds RNA and Traffics across Synaptic Boutons. Cell. 2018 Jan 11;172(1–2):262-274.e11.

36. Cottee MA, Letham SC, Young GR, Stoye JP, Taylor IA. Structure of Drosophila melanogaster
ARC1 reveals a repurposed molecule with characteristics of retroviral Gag. Sci Adv. 2020
Jan;6(1):eaay6354.

37. Pastuzyn ED, Day CE, Kearns RB, Kyrke-Smith M, Taibi AV, McCormick J, et al. The
Neuronal Gene Arc Encodes a Repurposed Retrotransposon Gag Protein that Mediates
Intercellular RNA Transfer. Cell. 2018 Mar 22;173(1):275.

38. Picelli S, Faridani OR, Björklund AK, Winberg G, Sagasser S, Sandberg R. Full-length RNA-seq
from single cells using Smart-seq2. Nat Protoc. 2014 Jan;9(1):171–81.

39. Rodrigues OR, Monard S. A rapid method to verify single-cell deposition setup for cell sorters.
Cytom Part J Int Soc Anal Cytol. 2016 Jun;89(6):594–600.

40. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq aligner. Bioinforma Oxf Engl. 2013 Jan 1;29(1):15–21.

41. Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput
sequencing data. Bioinformatics. 2015 Jan 15;31(2):166–9.

42. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of
multimodal single-cell data. Cell. 2021 Jun 24;184(13):3573-3587.e29.

43. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool
for interpreting omics data. The Innovation. 2021 Aug 28;2(3):100141.

44. Bergen V, Lange M, Peidli S, Wolf FA, Theis FJ. Generalizing RNA velocity to transient cell
states through dynamical modeling. Nat Biotechnol. 2020 Dec;38(12):1408–14.

45. Svensson V, Natarajan KN, Ly LH, Miragaia RJ, Labalette C, Macaulay IC, et al. Power analysis
of single-cell RNA-sequencing experiments. Nat Methods. 2017 Apr;14(4):381–7.

11   Figure captions

Figure 1
Classification of Tumor-associated hemocytes
(A) Tumor-associated hemocytes were labeled using plasmatocyte (P1), crystal cell (C4), and
lamellocytes (L1)-specific antibodies. **(B)** The frequency of the three hemocyte classes is shown in the
lower part for wild type and Ras^{V12}-SGs.

Figure 2

Characterization of SGs from the used genotypes

(**A-D**) SGs from wild type (w[1118]), p35 expressing larvae, Ras[V12] larvae and p35;Ras[V12] larvae were stained with phalloidin and a plasmatocyte-specific antibody (Hemese, H2, **A'-D'**). SG size for the different genotypes was quantified shown in **E** and hemocyte attachment in **F**. Whisker length min to max, bar represent median. P-value quantified with ANOVA.

Figure 3

Extraction and initial characterization of circulating hemocytes and TAHs from wild type and tumor larvae

(**A**) Upper panel: circulating hemocytes were extracted from larvae and individualized using cells sorting. Lower panel: TAHs were extracted from salivary glands using capillary suction, individualized and processed for single-cell RNA sequencing. See text for further details. (**B-H**) Classification of single hemocytes confirms their plasmatocyte identity and the absence of lamellocytes (Atilla) or crystal cell (Lozenge) markers.

Figure 4

Clustering analysis of hemocytes

Circulating single cells (Genotype: w[1118], Ras[V12] and p35;Ras[V12]) were processed using Smart-seq2. For manually extracted TAHs (Genotype: Ras[V12], p35;Ras[V12]) libraries were generated according to the Smart-seq2 protocol. (**A**) UMAP projection of single-cell transcriptome integrated datasets overlaid with batch labels. (**B**) Louvain clustering analysis based on the first 10 PCs shows the biological variation of the data and cell communities' assignments. (**C**) UMAP projection overlaid with single-cell specific genotypes. The global transcriptome similarities and differences were assessed based on k-nearest neighbors (kNN) force-directed graph with a true signal variation of the single-cell transcriptome on the integrated datasets. (**D**) Relative frequency of w[1118], Ras[V12], and p35Ras[V12] hemocytes within the five identified clusters. (**E**) Gene set enrichment analysis (GSEA) (Biological processes) of up-regulated genes in a minimum of 25 % of the cells in each cell community with a cut-off 0.25 log fold change threshold, comparing different cell communities (clusters) and highlighting the various biological processes' heterogeneity. Color scale indicating the corrected p-values where blue is less significant, and red is highly significant. Black circular dots indicate the gene ratio in comparison to the universal background gene list. Numbers in brackets indicate the gene numbers overlapping the ontology terms for a specific cluster.

Figure 5

Contribution of the three genotypes to clusters 2 (A), 3 (B) and 0 (C)

Expression intensity was analyzed for the most significantly enriched genes in the respective clusters using the visualization tool created for this project (see: https://mubasher-mohammed.shinyapps.io/Sc-drosophila/). (**D**) The enrichment of genes involved in cell division in cluster 3 (see also Figure 4E) was functionally confirmed using anti-phospho Histone 3 labeling to detect proliferating cells. Whisker length min to max, bar represent median. P-value quantified with ANOVA.

Figure 6

Delineating splicing kinetics through generalizing RNA velocity to cell population clusters

(**A**) Projecting velocities across the wild type, Ras[V12], and p35;Ras[V12] hemocytes. (**B**) Across conditions (attached and circulating). (**C**) Identified Seurat clusters. (**D**) Pseudotime inference of integrated datasets shows the root cells composed mainly of cluster 3 cells (mixture of p35;Ras[V12] and Ras[V12] larvae) and trajectory depicting pseudotime units assignment with terminal state composed mainly of cluster 1 (wild type and Ras[V12] larvae).

609
Figure 7
Non-autonomous distribution of crystal cell derived prophenoloxidase transcripts
(A) UMAP visualizes transcripts analysis showing the strongest presence of mature (spliced) mRNA
of crystal cell-derived prophenoloxidase 1 (PPO1) across attached hemocyte TAHs condition
corresponding to Ras^{V12} larvae. **(B)** scatter plot shows the absence of immature transcripts (unspliced)
reads across conditions (attached and circulating). **(C)** UMAP visualizes transcripts depicting cell-
derived prophenoloxidase 2 (PPO2) in attached hemocyte TAHs condition. **(D)** scatter plot indicates
the exclusive presence of mature and immature (spliced vs unspliced) PPO2 transcripts in
plasmatocytes (higher ratio of unspliced across circulating condition), in contrast to PPO2 which is
released from crystal cells through cell rupture and subsequently taken up by plasmatocytes most likely
through phagocytosis.
621
622 **Figure 8**
623 **A proposed model for the differentiation and distribution of hemocyte transcriptomes**
624 Hemocyte reactions towards SG dysplasia are schematically depicted for the Ras model (right part)
625 and after inhibition of effector caspases by p35 (left part). Communication via exosomes (broken
626 arrows) and transdifferentiation (solid arrows) are indicated. See text for further details.

A Genotype

● p35;Ras^{V12} ● Ras^{V12} ● w^{1118}

B Condition

● Circulating   ● Attached

C Seurat clusters

● Cluster 0   ● Cluster 1   ● Cluster 2
● Cluster 3   ● Cluster 4

D Pseudotime analysis

Figure 8

A

B

C