1 **The Janelia Atalanta plasmids provide a simple and efficient CRISPR/Cas9-mediated**

2 **homology directed repair platform for *Drosophila***

4 David L. Stern, Elizabeth Kim & Emily L. Behrman

6 Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, 20147 USA

Abstract

Homology-directed repair (HDR) is a powerful tool for modifying genomes in precise ways to address many biological questions. Use of Clustered Regularly Interspersed Short Palindromic Repeats (CRISPR)-Cas9 induced targeted DNA double-strand breakage has substantially simplified use of homology-directed repair to introduce specific perturbations in *Drosophila*, but existing platforms for CRISPR-Cas9-mediated HDR in *Drosophila* involve multiple cloning steps and have low efficiency. To simplify cloning of HDR plasmids, we designed a new plasmid platform, the Janelia Atalanta (pJAT) series, that exploits recent advances in dsDNA synthesis to facilitate Gateway cloning of gRNA sequences and homology arms in one step. Surprisingly, the pJAT plasmids yielded considerably higher HDR efficiency (approximately 25%) than we have observed with other approaches. pJAT plasmids work in multiple *Drosophila* species and exhibited such high efficiency that previously impossible experiments in *Drosophila*, such as driving targeted chromosomal inversions, were made possible. We provide pJAT plasmids for a range of commonly performed experiments including targeted insertional mutagenesis, insertion of phiC31-mediated attP landing sites, generation of strains carrying a germ-line source of Cas9, and induction of chromosomal rearrangements. We also provide "empty" pJAT plasmids with multiple cloning sites to simplify construction of plasmids with new functionality. The pJAT platform is generic and may facilitate improved efficiency CRISPR-Cas9 HDR in a wide range of model and non-model organisms.

Introduction

CRISPR-Cas9 induced mutagenesis has revolutionized experimental approaches in model and non-model organisms. DNA cuts induced by Cas9 or other enzymes[1,2] can be repaired by the non-homologous end-joining pathway to induce small deletions or cuts may be repaired by HDR if a homologous DNA template is present[3]. HDR provides the greatest power to manipulate gene function and to introduce new reagents into the germ line of experimental organisms, but CRISPR-Cas9 induced HDR works at low efficiency in many species.

HDR is normally implemented by introduction of three separate elements: Cas9 (or analogous enzyme), guide RNA, and homologous DNA. Usually, these three reagents are introduced as distinct components. In the most efficient systems, a source of gRNA and the homology arms are typically cloned into separate plasmids. Usually, plasmids containing homology arms are constructed by polymerase chain reaction (PCR) of homology arms from genomic DNA followed by cloning of arms together, often with a "payload" located between flanking homology arms. Generation of homology arms with PCR can be difficult for some target regions and cloning the homology arms with the payload also may have low efficiency, particularly when attempting to assemble multiple fragments.

We designed a new series of plasmids to simplify cloning for CRISPR-HDR experiments. The plasmid design includes two Gateway cloning sites for simultaneous introduction of long dsDNA fragments produced using new cost-effective DNA synthesis methods (Figure 1). The synthesized fragments incorporate homology arms between the *attL* Gateway cloning sites and one or both fragments includes a tRNA-sgRNA-tRNA array. We found that plasmids containing multiple copies of the standard Gateway negative selection marker, *ccdB*, were unstable[4] and we therefore introduced a novel dual selection system to ensure efficient simultaneous cloning of two attL cassettes. We placed a U6 promoter sequence flanking one or both Gateway cloning sites to drive transcription of the tRNA-sgRNA-tRNA array, followed by a U6 terminator; this combination provides efficient precise release of sgRNA molecules[5]. Any payload can then be placed between the Gateway cloning sites, and we have generated a series of plasmids carrying some commonly used payloads.

While we designed this plasmid for efficient cloning, we found that the plasmids yielded far higher HDR integration efficiencies on average than has been observed by us and others using other methods[3,6,7]. We have therefore named these plasmids the Janelia Atalanta (pJAT) series, after the mythological Greek heroine and exceptional archer (Supplementary Figure 1). The high efficiency of HDR with pJAT plasmids inspired us to attempt new kinds of experiments that have so far not been feasible in *Drosophila*, including introduction of chromosomal inversions.

The pJAT plasmids are modular and generic. One or two U6 promoter from any species can be introduced outside the Gateway cloning cassettes and any payload can be introduced between the cloning cassettes. pJAT plasmids may therefore be of value for HDR experiments in many model and non-model organisms.

Results

The pJAT *series dual-Gateway cloning platform*

Our initial aim was to build a plasmid platform that could utilize recent advances in DNA synthesis technology to simplify the cloning of reagents for CRISPR-Cas9 induced HDR. We found that some companies could synthesize long dsDNA fragments containing repetitive DNA attL sequences at the ends at a reasonable price (for example, 7 cents per bp from Twist Bioscience). This inspired us to design a dual-Gateway cloning platform for direct cloning of two fragments that each contain a homology arm (Gateway Cassettes 1 and 2; Figure 1A-C). To further reduce the number of plasmids that were required for experiments, we incorporated a tRNA-gRNA-tRNA design[5] directly into the synthesized fragments (Figure 1B, D). Our initial attempts at building a dual-Gateway plasmid followed earlier examples and simply duplicated the standard *ccdB* negative selection gene in two separate attR-attR cassettes[8]. We found, however, that every plasmid recovered during cloning carried a mutation in one of the *ccdB* genes, suggesting that high *ccdB* dosage is deleterious to *E. coli*[4]. We therefore replaced one of

the *ccdB* genes with a *sacB* gene, which allows negative selection by growing *E. coli* on plates containing high levels of sucrose[9].

The pJAT plasmids contain one or two U6 promoters flanking the Gateway cloning cassettes or restriction sites that can be used to introduce new U6 promoters. We have generated pJAT plasmids containing multiple kinds of payloads, including (1) a germline source of Cas9, (2) attP landing sites, (3) marker genes, and (4) a split-GFP system to report on simultaneous HDR of two plasmids (Figure 2). All of the current pJAT plasmids utilize fluorescent reporter genes expressed in different anatomical regions[10], which allow recombination of multiple reagents into a single wild-type fly without the use of balancer chromosomes. To facilitate introduction of novel payloads, we have generated one pJAT plasmid that contains a multiple cloning site for both standard ligation based cloning and Golden Gate cloning with BsaI or BbsI sites (Figure 2).

For plasmid construction, homology arms were synthesized by Twist Bioscience with compatible attL sites and the company's adaptor sequences are left on for three reasons: (1) synthesis sometimes fails without the adaptor sequences, probably because the repetitive attL sites are then located at the ends of the synthesized fragments, (2) keeping adaptors reduces synthesis cost; and (3) the adaptor sequences do not conflict with Gateway cloning. Synthesis of arms confers many benefits compared with PCR based construction of HDR plasmids, including the ability to include exogenous sequences, such as the tRNA-gRNA-tRNA array and other elements, as described further below, and the ability to introduce single base pair modifications. Synthesis by Twist Biosciences currently faces several limitations: (1) short homology arms are difficult to synthesize, probably because the repetitive attL sites are then positioned too close together; (2) the sequence cannot include more than 10 consecutive bp of the same nucleotide; (3) there are limitations on regions containing strong AT/GC bias and on repetitive sequences, which has prevented inclusion of multiple sgRNAs in a single synthesized fragment. Some of these synthesis challenges can be overcome by introducing innocuous point mutations, since HDR appears to be robust to considerable sequence variation relative to the target sequence. Cloning into pJAT plasmids requires standard Gateway cloning reagents and

little hands-on time. A full protocol describing design of homology arms and all steps of cloning

is provided at protocols.io.

pJAT *plasmids yield high efficiency HDR*

Our initial experiments with pJAT plasmids yielded efficiencies of HDR of approximately

40-50%. We had previously only rarely observed such high efficiencies of HDR in *Drosophila* and

so we systematically targeted many loci in *D. melanogaster* and found that we could

successfully target all loci. We confirmed that all HDR products were integrated into the correct

locations using TagMap[11] with pJAT-specific primers (Materials and Methods). The average

integration efficiency—that is, the fraction of fertile G0 injected eggs that yielded at least one

positive event in the next generation—was approximately 25% for insertions into genes in

which null mutations are homozygous viable and was more variable and lower on average for

genes in which null mutations were homozygous lethal (Figure 3).

Seven of the injections into lethal genes (shown as separate data points in Figure 3)

yielded no events, but these were repeated injections using into only two loci. Despite these

failures, other injections into these two loci yielded some events, though at low frequency.

Thus, all genes were targetable with pJAT plasmids, but a few genes, and specifically a subset of

those that are homozygous lethal, showed low efficiency. This low efficiency did not appear to

result from the specific gRNA sequences used (data not shown) or the homology arm lengths

(Supplementary Table 1). One possibility, which requires further investigation, is that the high

efficiency of pJAT plasmids may induce homozygous null genotypes in many cells of the

developing embryo, leading to lethality or sterility.

Multiple features contribute to high efficiency of pJAT *plasmids*

To determine possible reasons for the high efficiency of HDR with pJAT plasmids, we

systematically varied sections of the synthesized fragments. We identified two features that

improved HDR efficiency. First, as has been documented previously[5], flanking the sgRNA with

tRNAs increased efficiency over co-injection with a sgRNA (Supplementary Figure 2).

Surprisingly, a pJAT plasmid containing the sgRNA, but no flanking tRNAs, had similar efficiency

to co-injecting with prepared sgRNA (Supplementary Figure 2), indicating that Cas9 can utilize a

sgRNA even when it is embedded within flanking irrelevant RNA sequences, similar to results

observed by others[7]. Second, use of intact dsDNA plasmids yielded higher efficiency HDR than

plasmids that contained homology arms flanked by CRISPR-Cas9 cut sites (Supplementary

Figure 3), which should linearize the plasmid DNA *in vivo* and which has been recommended

previously[12]. It is possible that the relatively long homology arms used here improved

efficiency, but we found that arms as short as 250 bp yielded similar efficiency as longer arms

(Supplementary Table 1). We did not test shorter arms, since it is challenging to synthesize

short arms with the long repeated attR sequences on both ends. The concentration of sgRNA

produced from a single U6 promoter does not appear to be limiting, since synthesis of the same

gRNA from two U6 promoters did not increase HDR efficiency (Supplementary Figure 4).

While pJAT plasmids provide sufficient gRNA for efficient HDR, we hypothesized that the

quantity of Cas9 protein may be limiting when Cas9 is introduced exogenously, either as mRNA

or protein. To test this idea, we built a pJAT plasmid that drives expression of a *Drosophila*

codon optimized Cas9 mRNA from a germline specific *nanos* promoter with a *nanos* 3'UTR to

localize Cas9 mRNA to the developing oocytes (Figure 2) and found that it increased HDR

efficiency approximately 2-3 fold compared to co-injection with *Cas9* mRNA (Supplementary

Table 2).

All of the experiments described above were performed by incubating injected embryos

at 22°C after injection. Given the known temperature sensitivity of Cas9[13], we performed a

small test of incubation temperature by placing eggs at either 20°C, 22°C, or 25°C for 4 h after

injection. We observed an increase in integration efficiency at higher temperatures

(Supplementary Table 3), suggesting that the effect of temperature on CRISPR-Cas9 HDR in

Drosophila should be investigated further.

Note that the efficiencies reported for many experiments shown in Figure 3 resulted

from injections using Cas9 mRNA and with incubation at 22°C, which we now know are sub-

optimal conditions. Thus, the average efficiencies reported here may underestimate the

average efficiency using optimal conditions. The availability of a high-efficiency CRISPR-HDR

platform that works in multiple species invites new kinds of experiments. We next illustrate two

kinds of experiments that have been challenging to implement with previous technology.

Scarless mutagenesis with pJAT *plasmids*

The use of synthesized homology arms and Gateway cloning can simplify the

performance of scarless mutagenesis. One approach, which we have not explored, would be to

insert a selectable marker at a target genome location and then, in a second set of injections, to

replace the selectable marker with the intended genomic changes[14]. A second approach, which

we illustrate here, is to place a selectable fluorescent marker between *piggyBac* transposon

arms, which are included in the synthesized DNA fragments adjacent to homology arms that

include the intended genomic changes[15]. In cases where the synthesized homology arms carry

sites that differ from the genomic target, we have found that variable lengths of the

synthesized homology arms, starting from the sites closest to the CRISPR cut site, are integrated

into the genome. It is therefore best to place the intended genomic changes close to the *pBac*

arms and to sequence the target region in HDR integrants to identify flies that carry the

intended changes. The selectable marker, together with the *piggyBac* arms, is then removed by

crossing flies to a strain expressing the *piggyBac* transposase gene (Figure 4). In the experiment

performed here, we observed 58% integration efficiency (7 of 12 fertile injected G0 yielded

integrants) into a site in *D. yakuba* and the *pBac* sequence was efficiently removed from

multiple lines. This approach allowed efficient recovery of two scarless individual nucleotide

changes at the targeted locus (Supplementary Figure 5).

pJAT *plasmids can direct one-step targeted chromosomal inversions*

The HDR efficiency of pJAT plasmids was considerably higher than we have observed

using other approaches and this high efficiency emboldened us to try more complex

experiments. One experiment that would be of considerable value, especially for studies of

non-*melanogaster Drosophila* species, would be the ability to generate chromosomal inversions

that could subsequently be used to balance deleterious alleles[16]. This experiment requires

efficient cutting at two chromosomal sites followed by inversion of the intervening sequence

(Figure 5A). To encourage production of the inversion event, we designed a pair of pJAT

plasmids where each plasmid carried homology arms oriented in the same direction as the

intended inverted chromosomal region (Figure 5B). To simplify detection of simultaneous

integration of both plasmids, potentially indicative of an inversion, we built three pairs of pJAT

plasmids expressing split-GFP in either the eyes, the thorax, or the abdomen (Figure 2).

We targeted three inversions of approximately 58 kb to 3 Mbp in *Drosophila yakuba* to

provide potential balancer chromosomes for reagents that we had previously built at the

fruitless and *doublesex* genes in this species (Figure 5c). We targeted breaks within genes that

are expected to be homozygous deleterious or lethal. We detected multiple events for each

targeted inversion, at efficiencies of 8-23% (Supplementary Table 4).

We PCR amplified fragments corresponding to the expected inversion event breakpoints

(Supplementary Figure 6) and confirmed that the sequences of both breakpoints were oriented

as expected (Supplementary Figure 7). We crossed the three *D. yakuba* inversions to null alleles

of either *dsx* or *fru*, which they were designed to balance, and found that they all balanced the

targeted alleles for multiple generations. Thus, single inversions can be quickly introduced to

balance specific alleles in *Drosophila* species. This technology will substantially simplify the

maintenance of transgenic reagents and mutants in non-*melanogaster Drosophila* species. It is

possible that multiple inversions can be introduced serially onto a single chromosome arm to

generate balancer chromosomes of even greater utility[10], and we provide three sets of split-

GFP reagents that may enable these experiments (Figure 2).

Discussion

We have developed a series of new plasmids that provide a simple cloning platform and

enable high efficiency HDR in multiple *Drosophila* species. The plasmid design is flexible and

allows introduction of any payload of interest. The efficiency of pJAT plasmids is so high that it

is almost certainly preferable to random integration via transposable elements for most

purposes and may even be more efficient than the most widely used method of transgenesis in

Drosophila at the moment, site-specific recombination into attP sites[17]. We have reported

previously that some strains and some species appear to be resistant to specific transposable

elements and to *attP-attB* integration[18]. In fact, we previously tried multiple times to introduce

a *nos-Cas9* transgene using transposable elements into the specific *D. simulans* strain used in

this study without success, but we identified multiple independent events with a single

injection of a pJAT plasmid carrying *nos-Cas9*. Thus, pJAT plasmids are probably a preferable

method of transgenesis to random integration using transposable elements, especially for

studies of non-*melanogaster Drosophila* species.

We designed the pJAT plasmids for ease of cloning using synthesized homology arms

that incorporate tRNA-sgRNA-tRNA arrays. Because these arms are synthesized, any

modifications to the native sequence can be introduced during synthesis and then integrated

into the genome. We illustrated the power of this approach by including mobilization

sequences of the *piggyBac* transposon into the synthesized arms. This allows efficient scarless

removal of the transposon and reporter gene, leaving behind substitutions introduced into the

genome (Figure 4). An additional illustration of the utility of this approach is that additional

CRISPR target sites can be introduced flanking the homology arms, allowing the homology arms

and payload to be linearized *in vivo* (Supplementary Figure 3). This does not appear to increase

HDR efficiency using pJAT plasmids in *Drosophila* (Supplementary Figure 3), which is consistent

with reports of HDR induced by zinc-finger nucleases in *Drosophila* that showed that circular

donor DNA was more efficient than linearized DNA[2]. However, linearization may increase

efficiency of HDR or of homology-mediated end joining in other systems[19,20].

We initially attempted to build the double-Gateway system using multiple *ccdB*

containing cassettes, as has been demonstrated by others[8]. However, we discovered during

subsequent cloning that in every case one of the two *ccdB* genes had been inactivated. This

may result from the dosage sensitivity of *E. coli* to *ccdB* copy number[4]. We therefore developed

a novel Gateway-like second cloning cassette that exploits the well-established strong negative

selective capability of the *SacB* gene[21]. When *E. coli* are grown on high levels of sucrose,

expression of the *SacB* gene causes lethality. Thus, this stabilized double-Gateway system

dramatically improved cloning efficiency of two synthesized fragments containing

complementary *attL* sites over a design that utilizes two cassettes both using the *ccdB* gene.

While we designed the Janelia Atalanta series to simplify cloning, we found that these

plasmids also increased HDR efficiency compared with other methods that we have tested and

that have been reported. Often, HDR in many genes fails[3,6] and, when it succeeds and has been

reported efficiency is approximately 1-10% on average. Although difficult to quantify,

publication bias may mean these reported efficiencies are over-estimates of true efficiencies,

since we are aware of multiple examples of failed CRISPR-HDR attempts in *Drosophila* species

performed by other research groups. We so far have been able to target every locus attempted

and on average targeting genes or genomic regions that have non-lethal mutant phenotypes

results in approximately 25% of fertile injected animals yielding correct insertion events. We

performed a series of experiments to try to determine what might contribute to this increased

efficiency and it appears that multiple factors contribute to higher efficiency HDR. Some of the

most important factors appear to be (1) production of sgRNA from a tRNA-sgRNA-tRNA array,

(2) injection into embryos with a germline source of Cas9 protein, (3) homology arms of at least

250 bp long, (4) circular plasmid DNA, and possibly (5) increasing the incubation temperature to

25°C for a time after injection.

Despite the overall high efficiency of HDR with pJAT plasmids, we identified a strikingly

lower efficiency on average with plasmids targeting genes with lethal null phenotypes. A subset

of these "lethal" genes did show high efficiencies, however. In limited experiments, we found

no evidence that lower HDR efficiencies resulted from the specific gRNA or homology arms

used, but we cannot rule out the possibility that these variables influenced efficiency. One

possibility which we have not tested is that the pJAT plasmids are so efficient that when the

reagents work at one targeted allele in an individual, the second allele is also hit at high

frequency. This would cause injected animals to be mosaic for homozygous null mutations,

which may lead to lethality of sterility of many flies carrying HDR events.

The most powerful demonstration of the high efficiency of pJAT plasmids is that they

could be used to introduce two simultaneous breaks in a single chromosome and facilitate

inversion of large chromosomal regions through HDR (Figure 5, Supplementary Figures 5 & 6).

Three independent experiments, using two separate injection companies, yielded inversions

(Supplementary Table 4). These inversions provide practical benefits, since they can be used to

balance deleterious alleles and will simplify stock maintenance. It may also be possible to use

pJAT plasmids to generate multiply inverted balancer chromosomes with wide applicability[16]. It

has not escaped our attention that this approach may also allow reversion of naturally

occurring chromosomal inversions[22], which often become hot-spots of evolutionary

divergence[23]. Such experiments may allow, for the first time, detailed recombination-based

identification of loci "locked" within non-recombining inversions that have contributed to

phenotypic divergence.

The pJAT plasmids were designed to provide a generic platform for introducing many

variant elements easily and it is possible that pJAT plasmids may provide an improved CRISPR-

HDR platform for other species. One potentially species-specific component of this first set of

pJAT plasmids are the U6 promoters. We used the *D. melanogaster* U6-3 promoter for all of the

experiments described in this paper and we observed similar HDR efficiencies in *D.*

melanogaster, *D. simulans*, and *D. yakuba* (Figure 3), suggesting that this U6-3 promoter works

well for these species. This U6 promoter could be easily swapped with a different species-

specific or inducible promoter. We have therefore generated pJAT77, where cloning sites can

allow any promoter to be placed on either or both sides of the Gateway cloning cassettes.

Some pJAT plasmids presented in this paper (Figure 2_ contain other species-specific

components that may not work efficiently in other species. For example, some of these

plasmids include germ-line specific promoters and 3'UTRs that were derived from the *D.*

melanogaster nanos and *vasa* genes. It is possible that these will not work efficiently in other

species. But the pJAT plasmids with multiple cloning sites allow efficient cloning of new

payloads with other species-specific components. It will be interesting to determine whether

pJAT plasmids increase HDR efficiency in organisms other than *Drosophila*.

Methods

*Construction of Janelia Atalanta (*pJAT*) plasmids*

pJAT plasmids were assembled by Golden Gate cloning[24] of PCR amplified and synthesized fragments. All plasmids reported here are available from Addgene under plasmid numbers 204289-204312.

Gateway compatible homology arms were synthesized by Twist Bioscience with Adaptors On. Because these sequences contain almost identical attL sites at either end, synthesis is inefficient and often fails with Adaptors Off. In addition, since these fragments are cloned into the pJAT plasmids using Golden Gate cloning, there is no need to remove the Twist synthesis adaptors. Golden Gate cloning was performed as follows.

pJAT plasmid (~50 ng/uL)	1 uL
Homology arm left (50 ng/uL)	0.5 uL
Homology arm right(50 ng/uL)	0.5 uL
LR Clonase II	0.25 uL

Reaction was incubated at 25°C in a thermocycler for 2-12 hours. Then 0.125 uL of Proteinase K (provided with LR clonase kit) was added and sample was incubated in a thermocycler at 37°C for 20 minutes. The entire reaction volume was gently mixed with at least 25 uL of Zymo Mix&Go Competent Cells thawed on ice. The mixture was incubated for 30 minutes on ice, heat shocked for 30 seconds at 42°C, and returned to ice. 100 uL of SOC was added and the mixture was shaken at 200 RPM at 37°C for 1 hour. The entire mixture was plated on Tryptone-Yeast + Sucrose plates + Spectinomycin plates. The entire protocol, including a guide for designing and ordering homology arms, is provided at Protocols.io: dx.doi.org/10.17504/protocols.io.bp2l6bjokgqe/v1.

Injection of Drosophila *embryos*

Most pJAT plasmids were injected into Drosophila embryos by Rainbow Transgenics and

some were injected by Genetivision. Initially, plasmids were co-injected with Cas9 RNA

generated by *in vitro* transcription from plasmid encoding Cas9[10]. After we generated fly lines

carrying a *nanos-Cas9* transgene (*pJAT17*), plasmids were injected into flies homozygous for

pJAT17 and fertile adults emerging from injected eggs were outcrossed to a wild-type strain

and offspring were screened for fluorescent reporter genes. The genomic insertion location of

all integrants were determined using TagMap[11] with the following TagMap PCR primers, where

the underlined sequence represents the Tn5 MEB adaptor and the non-underlined sequence

represents the sequence specific to each pJAT plasmid.

Primer	Sequence
Tn5 ME B adaptor-attB4	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>AGCTGGACAACTTTGTATAGAAAAG
Tn5 ME B adaptor-P767_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>CGTCTCACAACTTTGTATACAAAAG
Tn5 ME B adaptor-P776_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>CGAGTATAATCAACTTTGTATACAAAAG
Tn5 ME B adaptor-P777_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>AAGCAACCCAACTTTGTATACAAAAG
Tn5 ME B adaptor-P795_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>AACAACTAGTCAACTTTGTATACAAAAG
Tn5 ME B adaptor-P832_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>CTCTCTAGTCAACTTTGTATACAAAAG
Tn5 ME B adaptor-P859_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>CTCACCATCAACTTTGTATACAAAAG
Tn5 ME B adaptor-P887_attB5	<u>GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAG</u>CTCGGCAACTTTGTATACAAAAG

Design and generation of scarless site specific mutations

Gateway compatible fragments were designed to include the 5' and 3' pBac sequences

required for transposition internal to the targeting homology arms (Figure 4). Fragments were

synthesized by Twist Biosciences and Gateway cloned into pJAT10 as described above.

Design and generation of targeted inversions

Gateway compatible fragments were designed with homology arms configured to

promote chromosomal inversions (Figure 5). Fragments were synthesized by Twist Biosciences

and Gateway cloned into pJAT21 and pJAT25 as described above. Both plasmids were injected

simultaneously into flies carrying a nos-Cas9 transgene.

Acknowledgements

We thank Carlos Machado, Thomas Ravenscroft, and Gerry Rubin for helpful comments on the

manuscript. Elements of Figures 1, 4, and 5 and Supplementary Figures 2, 3, and 4 were created

with BioRender.com. We especially thank Justine Ayelet Stern for designing and creating

Supplementary Figure 1 and for recommending the name Atalanta for these plasmids.

Author Contributions

D.L.S. conceptualized, designed, and constructed most of the plasmids and performed much of

the fly work. E.K. assisted with construction and quality control of plasmids and with fly work.

E.L.B. constructed pJAT17 and confirmed its utility. The paper was written by D.L.S. with input

from E.K. and E.L.B.

Ethics Declarations

HHMI has filed a provisional patent, number 63/507,335, for the inventor D.L.S. covering

vectors and methods for efficient cloning and homology directed repair.

References

1. Joung, J.K., and Sander, J.D. (2013). TALENs: a widely applicable technology for targeted genome editing. Nat. Rev. Mol. Cell Biol. *14*, 49–55. 10.1038/nrm3486.

2. Beumer, K.J., Trautman, J.K., Bozas, A., Liu, J.L., Rutter, J., Gall, J.G., and Carroll, D. (2008). Efficient gene targeting in Drosophila by direct embryo injection with zinc-finger nucleases. Proc Natl Acad Sci U A *105*, 19821–19826. 10.1073/pnas.0810475105.

3. Gratz, S.J., Ukken, F.P., Rubinstein, C.D., Thiede, G., Donohue, L.K., Cummings, A.M., and Oconnor-Giles, K.M. (2014). Highly specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in Drosophila. Genetics *196*, 961–971. 10.1534/genetics.113.160713.

4. Afif, H., Allali, N., Couturier, M., and Melderen, L.V. (2001). The ratio between CcdA and CcdB modulates the transcriptional repression of the ccd poison–antidote system. Mol. Microbiol. *41*, 73–82. https://doi.org/10.1046/j.1365-2958.2001.02492.x.

5. Port, F., and Bullock, S.L. (2016). Augmenting CRISPR applications in Drosophila with tRNA-flanked sgRNAs. Nat. Methods *13*, 852–854. 10.1038/nmeth.3972.

6. Ding, Y., Berrocal, A., Morita, T., Longden, K.D.K.D., and Stern, D.L.D.L. (2016). Natural courtship song variation caused by an intronic retroelement in an ion channel gene. Nature *536*, 329–332. 10.1038/nature19093.

7. Kanca, O., Zirin, J., Hu, Y., Tepe, B., Dutta, D., Lin, W.-W., Ma, L., Ge, M., Zuo, Z., Liu, L.-P., et al. (2022). An expanded toolkit for Drosophila gene tagging using synthesized homology donor constructs for CRISPR-mediated homologous recombination. eLife *11*, e76077. 10.7554/eLife.76077.

8. Aboulela, M., Tanaka, Y., Nishimura, K., Mano, S., Kimura, T., and Nakagawa, T. (2017). A dual-site gateway cloning system for simultaneous cloning of two genes for plant transformation. Plasmid *92*, 1–11. 10.1016/j.plasmid.2017.05.001.

9. Li, X., Thomason, L.C., Sawitzke, J.A., Costantino, N., and Court, D.L. (2013). Positive and negative selection using the tetA-sacB cassette: recombineering and P1 transduction in Escherichia coli. Nucleic Acids Res. *41*, e204. 10.1093/nar/gkt1075.

10. Stern, D.L. (2022). Transgenic tools for targeted chromosome rearrangements allow construction of balancer chromosomes in non- *melanogaster Drosophila* species. G3 GenesGenomesGenetics *12*, jkac030. 10.1093/g3journal/jkac030.

11. Stern, D.L. (2016). Tagmentation-Based Mapping (TagMap) of Mobile DNA Genomic Insertion Sites. bioRxiv, 1–7. 10.1101/037762.

12. Kanca, O., Zirin, J., Garcia-Marques, J., Knight, S.M., Yang-Zhou, D., Amador, G., Chung, H., Zuo, Z., Ma, L., He, Y., et al. (2019). An efficient CRISPR-based strategy to insert small and large fragments of DNA using short homology arms. eLife *8*, e51539. 10.7554/eLife.51539.

13. Xiang, G., Zhang, X., An, C., Cheng, C., and Wang, H. (2017). Temperature effect on CRISPR-Cas9 mediated genome editing. J. Genet. Genomics *44*, 199–205. 10.1016/j.jgg.2017.03.004.

14. Lamb, A.M., Walker, E.A., and Wittkopp, P.J. (2016). Tools and strategies for scarless allele replacement in Drosophila using CRISPR/Cas9. Fly (Austin) *0*, 1–12. 10.1080/19336934.2016.1220463.

15. Nyberg, K.G., and Carthew, R.W. (2022). CRISPR-/Cas9-Mediated Precise and Efficient Genome Editing in Drosophila. Methods Mol. Biol. Clifton NJ *2540*, 135. 10.1007/978-1-0716-2541-5_6.

16. Miller, D.E., Cook, K.R., and Hawley, R.S. (2019). The joy of balancers. PLoS Genet. *15*, e1008421. 10.1371/journal.pgen.1008421.

17. Groth, A.C., Fish, M., Nusse, R., and Calos, M.P. (2004). Construction of Transgenic Drosophila by Using the Site-Specific Integrase From Phage {phi}C31. Genetics *166*, 1775–1782.

18. Stern, D.L., Crocker, J., Ding, Y., Frankel, N., Kappes, G., Kim, E., Kuzmickas, R., Lemire, A., Mast, J.D., and Picard, S. (2017). Genetic and Transgenic Reagents for Drosophila simulans, D. mauritiana , D. yakuba , D. santomea, and D. virilis. G3, 1–10. 10.1534/g3.116.038885/-/DC1.1.

19. Bai, H., Liu, L., An, K., Lu, X., Harrison, M., Zhao, Y., Yan, R., Lu, Z., Li, S., Lin, S., et al. (2020). CRISPR/Cas9-mediated precise genome modification by a long ssDNA template in zebrafish. BMC Genomics *21*, 67. 10.1186/s12864-020-6493-4.

20. Wierson, W.A., Welker, J.M., Almeida, M.P., Mann, C.M., Webster, D.A., Torrie, M.E., Weiss, T.J., Kambakam, S., Vollbrecht, M.K., Lan, M., et al. (2020). Efficient targeted integration directed by short homology in zebrafish and mammalian cells. eLife *9*, e53968. 10.7554/eLife.53968.

21. Reyrat, J.-M., Pelicic, V., Gicquel, B., and Rappuoli, R. (1998). Counterselectable Markers: Untapped Tools for Bacterial Genetics and Pathogenesis. Infect. Immun. *66*, 4011–4017. 10.1128/IAI.66.9.4011-4017.1998.

22. Dobzhansky, T., and Sturtevant, A.H. (1937). Inversions in the chromosomes of Drosophila pseudoobscura. Genetics *23*, 28–64.

23. Joron, M., Frezal, L., Jones, R.T., Chamberlain, N.L., Lee, S.F., Haag, C.R., Whibley, A., Becuwe, M., Baxter, S.W., Ferguson, L., et al. (2011). Chromosomal rearrangements

453   maintain a polymorphic supergene controlling butterfly mimicry. Nature *477*, 203–206.
454   10.1038/nature10341.

455 24. Engler, C., Kandzia, R., and Marillonnet, S. (2008). A One Pot, One Step, Precision Cloning
456   Method with High Throughput Capability. PLoS ONE *3*, e3647.
457   10.1371/journal.pone.0003647.

Figure Legends

Figure 1 – Design and use of the Janelia Atalanta plasmids. (A) Generic design of the Janelia Atalanta series. Each plasmid contains two Gateway compatible cloning sites, flanked on the left and right, respectively, by attR3-attR4 and attR5-attR6 sites. Each Gateway compatible cassette contains a different positive and negative selection marker. The negative selection markers *ccdB* and *SacB* can be selected simultaneously by cloning into *ccdB* sensitive cells and plating on minimal media containing a high concentration of sucrose. Each Atalanta plasmid contains one or two U6 promoters flanking the Gateway cassettes and a payload between the Gateway cassettes. (B) DNA fragments or plasmids containing paired attL3-attL4 and attL5-attL6 sites can be cloned into the based Atalanta plasmid simultaneously. In this example, synthetic DNA fragments containing left and right homology arms and a tRNA-sgRNA-tRNA array are cloned into the Gateway sites. The plasmid thus provides a source both of sgRNA and of homology arms for homology directed repair. (C) An example of an Atalanta plasmid after cloning of Gateway-compatible arms that is ready for injection into *Drosophila* embryos. Cas9 is provided either through co-injection of Cas9 mRNA or protein or by injection into embryos expressing Cas9 in the germline. (D) Detail of the components of a generic tRNA-sgRNA-tRNA region cloned downstream of the U6 promoter.

Figure 2 – Payloads of Janelia Atalanta plasmids. (A) A set of plasmids containing fluorescent protein markers, attP sites, and removable fluorescent protein markers useful for targeted mutagenesis and generation of attP landing sites. (B) Plasmids to facilitate cloning of novel payloads. MCS=Multiple Cloning Site. (C) Plasmids carrying various effectors, including *nanos-Cas9* (pJAT17), B3 integrase under heat shock control (pJAT 43), and two plasmids expressing *vasa*-phiC31 integrase with different fluorescent protein markers (pJAT 56, pJAT 62). (D) Plasmids carrying split-*GFP* constructs under three alternative promoter systems for construction of chromosomal inversions.

Figure 3 – Integration efficiencies of Atalanta homology directed repair payloads into

Drosophila species. The percent of positive events is plotted against the number of viable

crosses screened. (A) Approximately 25% of injected animals producing fertile adults yielded

integration events into genes categorized as "viable" on Flybase or into non-genic regions. No

obvious difference was observed in integration efficiencies between the three *Drosophila*

species *D. melanogaster*, *D. simulans*, and *D. yakuba*. (B,) Integration efficiency for payloads

targeting genes categorized as "lethal" on Flybase (B).

Figure 4 – Schematic representation of scarless genome modification using pJAT plasmid. (A, B)

Diagram of a pJAT plasmid illustrating the location of the synthesized *pBac* transposon 5' and 3'

arms internal to the synthesized homology arms (A). The gRNA, included within the tRNA-gRNA-

tRNA array on the left homology arm, targets a genomic site that includes a TTAA motif (B), the

native target site for the *pBac* transposon. The synthesized right homology arm includes an

additional TTAA sequence between the 3' *pBac* sequence and the right homology arm.

Intended modifications to the genomic sequence can be included in either or both homology

arms. (C) Representation of the genomic sequence after integration of the pJAT plasmid. Flies

containing the intended genomic modification(s) can be identified by PCR and sequencing. (D)

After exposure to a source of *pBac* transposase, flies can be recovered that have lost the

reporter gene and contain the intended scarless genome modification.

Figure 5 – Design of pJAT plasmids for targeted inversions. (A) Schematic illustrating the

position and orientation of DNA sequences flanking two cut sites before (top) and after

(bottom) a chromosomal inversion. (B) Illustration of two pJAT plasmids, each containing one

half of a split-GFP reporter and a gRNA targeting one of the two intended cut sites. Each

plasmid includes synthesized arms oriented in the direction of the intended inversion so that

homology directed repair at both sites simultaneously will tend to drive an inversion. (C)

Illustration of three inversions targeted on the right arm of chromosome 3 in *D. yakuba*. The

positions of the *fruitless* (*fru*) and *Doublesex* (*dsx*) genes are shown in green. The locations of

target gRNA sites in the genes are shown in magenta. The three targeted inversions are shown

in beige.

Supplementary Figure Legends

Supplementary Figure 1 - Icon for the Janelia Atalanta series of plasmids. Representation of the

Greek goddess Atalanta practicing archery (artwork by Justine Ayelet Stern).

Supplementary Figure 2 - Effect of tRNA-sgRNA-tRNA array on CRISPR-Cas9 HDR efficiency.

Three plasmids were created where the left homology arm contained either no tRNA and

sgRNA (A), just the sgRNA (B), or the full tRNA-sgRNA-tRNA array (C). Injections of the plasmid

without a sgRNA were supplemented with *in vitro* transcribed sgRNA. All plasmids targeted the

CG42402 gene. The table shows the HDR efficiency for each construct type.

Supplementary Figure 3 – To test the hypothesis that *in vivo* linearization of the template DNA

increases the efficiency of HDR, plasmids targeting two loci were constructed either without (A)

or with (B) CRISPR target sites for the sgRNA encoded in the tRNA-sgRNA-tRNA array. Results

are shown in the table, which illustrates that plasmids containing the flanking CRISPR sites did

not increase HDR efficiency for two genes.

Supplementary Figure 4 – To test the hypothesis that sgRNA concentration limited HDR,

plasmids including either one (A) or two (B) sources of sgRNA were constructed. Results are

shown in the table, which illustrates that plasmids containing two sources of sgRNA did not

increase HDR efficiency for two genes on average.

Supplementary Figure 5 – Sequence confirmation of scarless genome mutagenesis. (A)

Sequencing of genomic DNA containing inserted pBac transposable element with internal MHC-

DsRed reporter. Left homology arm includes intended 2 bp changes. PCR fragments were

generated with primers external to homology arms (green) and internal to the pBac transposon

and Sanger sequenced with internal primers (magenta). Sanger sequencing products were

aligned to the original genome sequence and mismatches are shown in the Identity track and

below the Sanger sequencing read. (B) A PCR product of the final scarless allele generated with

primers shown (green) was Nanopore sequenced and aligned to the original genome sequence

and mismatches are shown in the Identity track and within the consensus Nanopore sequencing

read.

Supplementary Figure 6 – Detailed maps from the *D. yakuba* genome of the gRNA sites

(magenta), homology arms (yellow), intended inversion ends (brown), and PCR primers for the

genes *VMS100-2* (A), *fray* (B), *Os-C* (C), and *Ids* (D).

Supplementary Figure 7 – Sequences of PCR products generated using primers illustrated in

Supplementary Figure 6 on strains carrying the putative inversions. Three inversions were

generated, one between *VMS100-2* and *fray* (A, B), one between *Osc* and *Ids* (C, D), and one

between *VMS100-2* and *Ids* (E, F).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

Site1   Site2

B

attB3   Site 1   Site 1 Homology   attB4   enhancer   EGFPn   attB5   Site 2 Homology   attB6
        tRNA-sgRNA   Arm Left   Arm Left

d-U6-3

attB3   Site 1   Site 1 Homology   attB4   enhancer   EGFPc   attB5   Site 2 Homology   attB6
        tRNA-sgRNA   Arm Right   Arm Right

d-U6-3

C

4,970,782   5,470,782   5,970,782   6,470,782   6,970,782   7,470,782   7,970,782   8,470,782

VMS100-2 fray   Os-C lds
fru   dsx

In (3R:4.8-8.4)

In (3R:4.8-5.0)   In (3R:8.3-8.4)

Supplementary Figure 1

Construct form	Construct	Number injected	Number Fertile	Number Positive	Efficiency (%)
A	CG42402 + gRNA	250	17	3	18
B	gRNA-CG42402	285	17	3	18
C	tRNA-gRNA-tRNA-CG42402	283	25	12	46

Construct Form	Construct	Left Homology Arm Length	Right Homology Arm Length	Number injected	Number Fertile	Number Positive	Efficiency (%)
A	l(2)gl	250	250	265	65	1	2
A	l(2)gl	1000	1000	265	45	0	0
B	CRISPR-l(2)gl-CRISPR	250	250	250	37	0	0
B	CRISPR-l(2)gl-CRISPR	1000	1000	280	61	0	0
A	Ir41a	250	250	260	64	25	39
A	Ir41a	600	600	260	46	12	26
B	CRISPR-Ir41a-CRISPR	250	250	255	53	10	19
B	CRISPR-Ir41a-CRISPR	250	600	280	47	4	9

Construct form	Construct	Number injected	Number Fertile	Number Positive	Efficiency (%)
A	dU6-3-Tm1	240	27	13	48
B	dU6-3-Tm1-dU6-1	260	31	8	26
A	dU6-3-Osi2	250	49	26	53
A	dU6-3-Osi2	250	72	10	36
A	dU6-3-Osi2	275	82	10	14
B	dU6-3-Osi2-dU6-1	275	44	19	43

Supplementary Figure 5

A

B

Supplementary Figure 6

Supplementary Figure 7