1 # First Chromosome-scale Genome Assembly of *Arundo donax*

2 **Authors & Affiliations**

3 Mengmeng Ren[1,+,*], Fupeng Liu[1], Xiaohong Han[1]

4 [1]Wuhan Rundo Biotechnology Co., Ltd. Wuhan, Hubei Province, 430300, China

5 [+,*]First author and corresponding author: Mengmeng Ren (ren_mm1991@163.com)

Abstract

   Arundo donax L (*A. donax*). is a promising energy crop with high biomass and wide adaptability, while lack of reference genome limiting the genetic improvement of this crop. Here, we report two chromosome-scale assemblies of *A. donax* genome (0004 and 0408) using Pacbio SMRT sequencing and Hi-C technology. The genome size of these two assemblies is 1.30 Gb / 2.86 Gb with contig N50 33.15 Mb / 34.62 Mb respectively. A total of 74,403 / 167,586 gene models were predicted in 0004 and 0408 genome, of which over 90% genes were functionally annotated. Our result will enhance genetic understanding and promote the genetic improvement of *A. donax*.

Keywords: *A. donax*, genome assembly, genome annotation

1. Introduction

Arundo donax L. (*A. donax*) is a tall perennial grass that is commonly known as giant reed. *A. donax* is a widely utilized plant species with various applications in bioenergy, papermaking, and even as an ornamental lawn grass due to its fast growth and high biomass production [1,2,3,4]. One of the significant characteristics of *A. donax* is its high productivity and fast growth rate. The plant matures rapidly and is capable of producing large quantities of biomass in a short time, which makes it an attractive crop for bioenergy production [5,6,7]. Additionally, *A. donax* is highly tolerant of a broad range of environmental conditions, including drought, excess water, and high salinity [8,9]. These attributes make the plant an excellent candidate for sustainable agriculture and bioenergy development.

A. donax is also known for its environmental and ecological benefits. It is considered to be an excellent tool for phytoremediation, which could be used to clean up contaminated soil and water [10,11]. *A. donax* is effective in removing heavy metals and other pollutants from contaminated sites, making it a valuable resource in environmental restoration [12]. In recent years, there has been increased interest in the study and utilization of *A. donax*, which has led to a number of research efforts aimed at understanding its biological properties.

The sterile nature limits the genetic variability and delays the genetic improvement of *A. donax* [13,14,15,16]. However, the fundamental research of *A. donax* genome encountered great difficulties. The ploidy of *A. donax* is still unclear because of the high chromosome number and small chromosome size [17,18]. Similar as another aquatic plant *Phragmites australis* [19], *A. donax* showed elastic chromosome number with variable ploidy. Different chromosome number of *A. donax* were reported by several authors, including 110 [20,21], 108 [22] and 84 chromosomes [23]. The complexity of *A. donax* genome hinders further understanding and the genetic improvement of this valuable energy crop.

The large and complex nature of plant genomes, coupled with varying degrees of ploidy, has long posed challenges for genome assembly in plants. Fortunately, advancements in sequencing technologies, such as third-generation long-read sequencing, particularly High Fidelity reads (Hifi) sequencing, along with new assembly algorithms serve as the foundation for the rapid advancements in plant genome sequencing [24]. Hifi sequencing allows for the generation of long-read, high-quality data that can be used to accurately assemble and annotate complex genomes. It has already proven to be a valuable tool in multiple complex plant genomes, such as allohexaploid oat [25], tetraploid potato [26], and autotetraploid alfalfa [27].

Here, we assembled the first two chromosome-scale *A. donax* genome by combining Pacbio SMRT sequencing and high-throughput chromosome conformation capture (Hi-C) technology. These two assemblies will facilitate the better understanding of *A. donax*

genomics and better utilization of *A. donax* resourses.

2. Results

2.1. Cytological analysis of A. donax

In the process of collecting domestic *A. donax* germplasm resources, we found that *A.*

donax in China can be divided into three main types according to the leaf color, i.e. common,

versicolor and emerald green (emerald here after) (Figure 1). Among them, there is little

difference in plant architecture, growth period and other aspects between common and

versicolor varieties. In contrast, obvious differences can be observed between common

and emerald *A. donax* in plant architecture, panicle architecture and growth period.

Therefore, we selected a common variety 0004 and an emerald variety 0408 for genome

assembly.

We first analyzed the karyotype of the two varieties. DAPI staining *in situ* hybridization

of telomere repeat sequences showed that 0004 had 112 (Figure 2A-B) chromosomes,

while 0408 had 62 chromosomes (Figure 2D-E), in consistent with the fact that

chromosome number in *A. donax* is variable. Moreover, rDNA fluorescence *in situ*

hybridization revealed that 6/6 chromosomes in 0004 showed strong hybridization signals

of 5SrDNA and 18SrDNA (Figure 2C). In contrast, only 2/2 chromosomes in 0408 showed

hybridization signals of 5SrDNA and 18SrDNA (one strong signal and one weak signal)

(Figure 2F). These results indicated that 0004 and 0008 may have different chromosome

basic number and ploidy.

2.2. High-quality genome assembly of A. donax

Before the genome *de novo* assembly, we conducted k-mer analysis using second

generation sequencing (MGI-2000 platform). We first checked for sequence contamination

by randomly selecting 100,000 reads and matched against sequences in the nucleotide

sequence database. About 79.8% and 84.5% reads of 0004 and 0408 were unmapped in

the database, confirming the low contamination of MGI data. Besides, over 2000 read were

mapped with *Setaria viridis* and *Digitaria exilis*, indicating the homology of *A. donax* and

these two species (Supplementary Figure S1A-B). The estimated genome size of 0004

and 0408 was 1.46 Gb and 3.14 Gb respectively. In addition, we estimated the genome

heterozygosity using simulated data of *Arabidopsis*. The genome heterozygosity of 0004

and 0408 is 0.8% and 0.3 % respectively (Supplementary Figure S1C-D).

The genome assembly were performed using long reads generated by PacBio platform.

After quality control, we obtained 152.30 Gb reads with N50 16.92 kb for 0004 and 133.18

Gb reads with N50 21.37 kb for 0408. The preliminary assembly of 0408 was 2.97 Gb, in

consistent with 3.14 Gb estimated with genome survey. However, assembly of 0004 was

2.24 Gb, much larger than 1.46 Gb. Consider the high heterozygosity of 0004 genome

(0.8%), we used Purge_Dups (-f .9) [28] to obtain the 1.41 Gb non-redundant genome. The

GC depth analysis showed improvement of genome quality for 0408 after redundancy reduction (Supplementary Figure S2).

We use Benchmarking Universal Single-Copy Orthologs (BUSCO) and CEGMA to assess the completeness of genome assembly. The percentage of complete BUSCOs was 99.57% and 99.69% in 0004 and 0408 genome (Supplementary Figure S3A-B), and the percentage of complete core gene was 92.74% and 91.53% in 0004 and 0408 genome (Supplementary Figure S3C-D), confirming the completeness of two genome assembly. Besides, to determine the accuracy of the genome assembly, we mapped the MGI reads to the assemblies. The map rate is 99.60% and 99.96% for 0004 and 0408 respectively, proving the accuracy of the genome assemblies.

2.3. Hi-C scaffolding

Since the chromosome basic number of *A. donax* is still unclear, we first assumed that the two *A. donax* varieties are diploid, thus the chromosome basic number is 56 and 31 for 0004 and 0408 respectively. We performed Hi-C scaffolding and the Hi-C interaction heatmap showed that both 0004 and 0408 are not diploid (Supplementary Figure S4). The number of Hi-C interaction signals for 0004 and 0408 is around 40 and 70 respectively. A hypothesis is that ancestor of all *Arundo* species is *A. plinii*, whose haploid chromosome number is 36 (n=36) [29]. Therefore, we redid Hi-C scaffolding with n = 36 for 0004 (Figure 3A). Overall, about 99.78% of total sequences were anchored into 36 pseudochromosomes with sizes ranging from 18.58 Mb to 55.91 Mb. The final genome assembly is 1.30 G with Scaffold N50 37.31 Mb (Table 1 and Figure 3C). Based on these results, we speculated common *A. donax* is triploid (2n = 3x = 108), while 0004 is an aneuploid with four extra chromosomes (3x + 4 =112). The LAI score of assembly constructed by Hi-C scaffolding was 12.63, reaching to the standard of reference quality.

However, for the 0408 genome assembly, the chromosome basic number is obscure. The number of Hi-C interaction signals (about 70) is more than the total chromosome number (62). Therefore, we assumed that 0408 is haploid and redid Hi-C scaffolding with n = 62 for 0408 (Figure 3B). About 97.00% of total sequences were anchored into 62 pseudochromosomes with sizes ranging from 24.80 Mb to 97.92 Mb. The final genome size is 2.86 Gb with scaffold N50 44.86 Mb (Figure 3D). The LAI score of assembly constructed by Hi-C scaffolding was 8.94 for 0408, approaching to the standard of reference quality.

To clarify the relationship between 0004 and 0408, we compared genomes of 0004 and 0408. We identified some chromosome structure variations of the two genomes. More interestingly, the two genomes showed an obvious 1:2 pattern of syntenic depth between 0004 and 0408 (Figure 3E). Therefore, we speculated that 0408 is an allodiploid.

189 Table 1. Statistics of genome assemblies.

	0004	0408
Assembly features		
Number of scaffolds	65	162
Total size of scaffolds	1299.92 Mb	2770.95 Mb
Longest scaffold	55.91 Mb	97.92 Mb
Shortest scaffold	18.58 Mb	24.80 Mb
Mean scaffold size	20.00 Mb	17.10 Mb
N50 scaffold length	37.31 Mb	44.86 Mb
L50 scaffold count	7	9
Scaffold GC content	44.07%	44.05%
Scaffold N content	0.0002%	0.0003%
Percentage of assembly in scaffolded contigs	99.78%	97.96%
Average number of contigs per scaffold	1.46	1.12
BUSCO (complete)	99.57%	99.69%
LTR Assembly Index (LAI)	12.63	8.94
Gene models		
Number of gene models	74,403	167,586
Mean coding sequence length	1192.62	1,138.43
Mean number of exons per gene	5.31	5.02
Mean exon length	224.79	226.57
Mean intron length	537.64	518.93
Non-protein-coding RNA		
Number of rRNA	2,320	6,305
Number of sRNA	3,118	6,702
Number of regulatory	19	29
Number of tRNA	1,392	3,175

2.4. Repeat elements analysis and gene model prediction

The total length of interspersed repeats is 711.24 Mb (54.71% of the genome) for 0004

and 1627.27 Mb (56.96% of the genome) for 0408. To be specific, a total of 70,102 (0.07%

of the genome) and 149,490 (0.07% of the genome) simple repeat sequence (SSR) were

identified in 0004 and 0408 genome; a total of 89,378 (0.71% of the genome) and 181,468

(0.78% of the genome) tandem repeat sequences were identified in 0004 and 0408

genome; a total of 1,400,366 (51.54% of the genome) and 2,937,325 (52.91% of the

genome) transposable elements (TEs) were identified in 0004 and 0408 genome. The

detailed statistics of TEs were listed in Supplementary file (TE).

We performed gene structure prediction by combining transcriptome prediction,

homologous protein prediction and ab initio prediction. Firstly, we found that the mapped

rate of RNA-seq data to the genome in four tissues were all over 90% (Supplementary

table 1), and the mapped rate of Pacbio Isoseq to the genome is 99.86% for 0004 and

99.87% for 0408, further confirming the accuracy of transcriptome data and the genome assemblies. The RNA-seq and Pacbio transcripts were used for gene prediction, resulting in 49,524 / 80,249 predicted genes for 0004 and 0408 respectively. Secondly, we selected five Poaceae plants, including *Saccharum spontaneum*, *Sorghum bicolor*, *Zea mays*, *Triticum aestivum* and *Oryza sativa* for homologous protein prediction. A total of 94,613 / 172,436 genes for 0004 and 0408 were predicted. Thirdly, we performed ab initio prediction, and 78,102 / 174,926 gene models for 0004 and 0408 were predicted. The final gene set were obtained by integrating the above results. In total, 74,403 / 167,586 gene models with average gene length 3,507.41 / 3,226.92 bp, average CDS length 1192.62 / 1,138.43 bp, average exon length 224.79 / 226.57 bp, average intron length 537.64 / 518.93 bp for 0004 and 0408 respectively (Supplementary file (Gene prediction)). Except the gene model, we also predicted non-coding RNA. In total, 2,320 / 6,305 rRNA, 3,118 / 6702 small RNA, 1,392 / 3,175 tRNA were identified in 0004 and 0408 genome. The parameters of the two assemblies were listed in Supplementary file (ncRNA).

2.5. Gene function annotation and evaluation of genome annotation

We predicted the gene function based on five databases, including Non-Reduntant Protein Database (NR), Kyoto Encyclopedia of Gene and Genomes (KEGG), Eukaryotic Orthologous Groups of protein (KOG), GO and Swissprot (Supplementary Figure S5). In total, 67,377 / 152,275 genes were annotated, account for 90.56% / 90.86 % of the 0004 and 0408 genome (Supplementary file (Gene prediction)). The co-annotated gene number is 16,877 / 37,776 for 0004 and 0408 genome (Supplementary Figure S6).

The annotated gene sets were evaluated using BUSCO. Among the 1,614 BUSCO groups, about 98.45% / 97.96% of complete gene elements can be found in the annotated gene set for 0004 and 0408 respectively, indicating that the majority of conservative gene predictions are relatively complete, indicating the high reliability of the gene prediction result. Besides, the proportion of expressed genes in four tissues ranged from 71.09% to 80.34% in 0004 and 67.01% to 73.75% in 0408. Total expressed genes account for 86.60% and 82.53% of the whole gene sets of 0004 and 0408 respectively (Supplementary file1 (Transcripts)). The gene structure in 0004 and 0408 genome showed similar distribution trend with other Poaceae plants, including gene length, CDS length, exon length, exon number, intron length and intron number (Supplementary Figure S7), demonstrating the reliability of the genome annotations.

3. Materials and Methods

3.1. Plant materials

The two *A. donax* plants were collected from Wuhan, Hubei (0004) and Yueyang, Hunan (0408) respectively. About 3 cm stem with a tiller bud were used as explant to propagate with tissue culture. Seedlings were then acclimatized outside and transplanted in the field.

3.2. Karyotype analysis

Karyotype analysis were conducted followed a previously report [30,31]. Briefly, root tips about 2 cm in length were placed in Nitrous Oxide (0.9-1.0 Mpa) for two hours, and fixed with pre-cold glacial acetic acid. After that, Root tips were digested with cellulase and pectinase at 37°C for 1 hour and crushed using a dissecting needle. Cells suspension were observed under an optical microscope to get the suitable chromosome specimens. Fluorescent *in situ* hybridization probes were prepared using nick translation method. Probes for telomere, 5SrDNA and 18SrDNA are fluorescently-labeled $(TTTAGGG)_6$ oligonucleotides, pTa794 and pTa71 plasmids respectively. After hybridization, slides were photographed under a fluorescence microscope.

3.3. Flow cytometry analysis

Karyotype analysis were conducted followed a previously report [32]. Samples were immersed into MG^b dissociation solution (45 mM $MgCl_2$ ·$6H_2O$, 20 mM MOPS, 30 mM sodium citrate, 1% (W/V) PVP, 0.2% (v/v) Tritonx-100, 10 mM Na 2 EDTA, 20 µL/mL β-mercaptoethanol pH 7.5), chopped with blade and filtered to get the nuclear suspension. The suspension was then incubated in propidium iodide and RNase for 1 h and detected with flow cytometry (BD FACScalibur). Tomato and barley were used as internal reference.

3.4. DNA extraction, Pacbio library preparation and sequencing

High molecular weight genomic DNA was prepared with QIAGEN® Genomic kit (13343) according to the manufacturer's instructions. SMRTbell libraries were constructed PacBio's standard protocol. Briefly, the quality qualified DNA was sheared by g-TUBEs (Covaris, USA), damage repaired and 20-50 kb DNA fragments were screened by the BluePippin (Sage Science, USA) and purified using AMPure PB beads (Agilent technologies, USA). Sequencing was performed on a PacBio Sequel instrument with Sequel II Sequencing Kit 2.0. Raw sequencing data were filtered through the SMRTlink 8.0.

3.5. MGISEQ-2000 library preparation and sequencing

Genomic DNA was extracted using CTAB method and subjected to sonication by Covaris. The DNA fragments range from 200-400 bp were selected using Agencourt AMPure XP-Medium kit. After end-repair, 3' adenylated, adapters-ligation and PCR Amplifying, the libraries were prepared using Hieff NGS® Fast-Pace DNA Cyclization Kit (Yeasen, 13341ES96) and sequencing on MGISEQ-2000 platform.

To facilitate the annotation of the genome, we performed mRNA sequencing on four tissues of the plant including leaf, root, panicle and lateral bud. We purified up to 400 µg of total RNA per sample using the TRIzol-based method (Invitrogen, USA) and subsequently treated it with DNase I. PolyA mRNA was enriched using the protocol of NEBNext ® Poly(A) mRNA Magnetic Isolation Module (New England Biolabs #7490 S, USA). RNA sequencing libraries were then prepared using Hieff NGS® Ultima Dual-mode mRNA Library Prep Kit

(Yeasen, 12309ES96) following the manufacturer's instructions. The RNA library concentration was determined using the Qubit ® 3.0 Fluorometer, and the libraries were then subjected to 150 bp paired-end sequencing on a MGISEQ-2000 platform. In total, we generated 55.8 Gb / 51.9 Gb raw data for the four tissue samples of 0004 and 0408 respectively.

3.6. Estimation of genome size and heterozygosity

The genome size of *A. donax* was estimated based on k-mer analysis using KMC 3 program [33]. The GCE and FindGSE software were used to analyze the 17-mer frequency distribution and perform genome size estimation [34]. We estimated the genome heterozygosity by combining simulated data with different heterozygosity in *Arabidopsis* genome and the distribution of 17 k-mer data of *A. donax*.

3.7. De novo assembly

The quality-filtered HiFi reads were used for genome assembly with hifiasm (v.0.12) [35] to obtain the preliminary assembly. The completeness of genome was assessed using BUSCO (v.4.0.5) [36] and CEGMA (v.2) [37]. To assess the accuracy of the assembly, all the MGISEQ-2000 reads were mapped to the genome using BWA [38], and SAMtools [39] and bcftools [40] were used to calculate the error rate of genome assembly. Beisides, LTR Assembly Index (LAI) was calculated to assess the assembly quality [41].

3.8. Hi-C scaffolding

To attach the scaffolds to chromosomes, we extracted genomic DNA using young leaves for the Hi-C library, and the sequencing data was obtained via the MGISEQ-2000 platform. To elaborate, we first cut freshly harvested leaves into 2 cm pieces and vacuum infiltrated them with nuclei isolation buffer supplemented with 2% formaldehyde. Crosslinking was stopped by adding glycine and performing additional vacuum infiltration. We then converted the fixed tissue into powder and re-suspended it in nuclei isolation buffer to obtain a nuclei suspension. D*pn*II was used to digest the purified nuclei with 100 units, and after marking with biotin-14-dCTP, biotin-14-dCTP from non-ligated DNA ends was removed through the exonuclease activity of T4 DNA polymerase. The ligated DNA was sheared into 300−600 bp fragments, followed by blunt-end repair and A-tailing before purification through biotin-streptavidin-mediated pull down. Finally, the Hi-C libraries were quantified and sequenced using the MGI-2000 platform.

In total, 375 Gb and 347 Gb paired-end reads were generated for 0004 and 0408. To ensure the quality of the Hi-C raw data, we employed HiC-Pro (v2.8.1), a previously established quality control tool [42]. We first filtered out low-quality sequences (quality scores < 20), adapter sequences, and sequences shorter than 30 bp using fastp [43]. The clean paired-end reads were then mapped to the draft assembled sequence using bowtie2 (v2.3.2) with the parameters "-end-to-end --very-sensitive -L 30" to obtain the unique

mapped paired-end reads [44]. We identified and retained valid interaction paired reads
using HiC-Pro from the unique mapped paired-end reads for further analysis. Additionally,
we filtered out invalid read pairs, including dangling-end, self-cycle, re-ligation, and
dumped products, using HiC-Pro. We further clustered, ordered, and oriented the scaffolds
onto chromosomes using LACHESIS [45], with parameters
CLUSTER_MIN_RE_SITES=100, CLUSTER_MAX_LINK_DENSITY=2.5, CLUSTER
NONINFORMATIVE RATIO = 1.4, ORDER MIN N RES IN TRUNK=60, ORDER MIN N
RES IN SHREDS=60. Finally, we manually adjusted any placement and orientation errors
exhibiting obvious discrete chromatin interaction patterns.

3.9. Repeat Annotation
To identify tandem repeats and transposable elements in the genome, we used GMATA
[46] and Tandem Repeats Finder (TRF) software [47]. GMATA identified simple repeat
sequences (SSRs), while TRF recognized all tandem repeat elements in the genome. To
identify transposable elements, we used a combination of ab initio and homology-based
methods. An ab initio repeat library for the genome was predicted using MITE-hunter [48]
and RepeatModeler with default parameters, including LTR_FINDER [49], LTRharvest [50],
and LTR_retriever [51] for plant genome. The resulting library was aligned to TEclass
Repbase (http://www.girinst.org/repbase) to classify each repeat family's type. To identify
repeats throughout the genome, we applied RepeatMasker [52] to search for known and
novel TEs using the *de novo* repeat library and Repbase TE library. Finally, we collated
and combined overlapping transposable elements belonging to the same repeat class.

3.10. Gene Prediction
To predict genes in a repeat-masked genome, we employed three distinct approaches:
ab initio prediction, homology search, and reference-guided transcriptome assembly.
Homolog prediction was performed using GeMoMa by aligning homologous peptides from
related species to the assembly to obtain gene structure information [53]. For RNAseq-
based gene prediction, we aligned filtered MGISEQ-2000 reads to the reference genome
using STAR [54] (default settings) and then assembled the resulting transcripts with
stringtie2 [55]. Open reading frames (ORFs) were predicted using PASA [56]. Ab initio gene
prediction was performed with Augustus [57], utilizing the default parameters with the
training set. We then employed EVidenceModeler (EVM) [58] to produce an integrated
gene set, removing genes with TE using the TransposonPSI package
(http://transposonpsi.sourceforge.net/) and filtering miscoded genes. Untranslated regions
(UTRs) and alternative splicing regions were identified using PASA based on RNA-seq
assemblies. We retained the longest transcripts for each locus, while regions outside of
the ORFs were designated as UTRs.

3.11. Functional annotation of gene models

Gene function information, as well as motifs and domains of their proteins, were
assigned by comparing them with various publicly available databases, including
SwissProt, NR, KEGG, KOG, and Gene Ontology. We utilized the InterProScan program
with default parameters to identify the putative domains and GO terms of the genes [59].
We also employed BLASTp to compare the EvidenceModeler-integrated protein
sequences against four well-known public protein databases, using an E-value cutoff of
1e-05. We selected the results with the lowest E-value. Results from the searches of these
five databases were then concatenated to obtain a comprehensive gene function
annotation.

3.12. Annotation of non-coding RNAs (ncRNAs)
For the identification of ncRNA, we utilized two main strategies: searching against
databases and prediction with models. To predict transfer RNAs (tRNAs), we applied
tRNAscan-SE with eukaryote parameters [60]. We employed Infernal cmscan [61] to
search the Rfam database for microRNA, rRNA, small nuclear RNA, and small nucleolar
RNA detection. Additionally, we utilized RNAmmer to predict the rRNAs and their subunits
[62].

3.13. Collinearity analysis
Pairwise alignment between 0004 and 0408 genome were performed using Minimap2
software [63] with parameters: -cx asm10. The paf file were used to obtain the dotplot using
R package dotPlotly (https://github.com/tpoorten/dotPlotly/).

4. Discussion
4.1 A. donax is a promising energy crop with complex genome
The rapidly deteriorating environment in earth has made an urgent claim for carbon
emission reduction [64]. Using bioenergy to replace part of fossil energy can make
remarkable contributions to reducing carbon emissions [65]. However, the land-intensive
bioenergy is limited by the finite land resource. Especially in China, because of the huge
population size, croplands are inviolable to guarantee the food security. However, except
for the cropland, there are about 78 million hectares of marginal lands [66]. Due to climate,
terrain, soil and other limiting factors, marginal lands cannot be planted with food crops,
while energy crops can often grow on these lands owing to their great environmental
adaptability. Of all the energy crops, the high yield, extensive adaptability and low inputs
make *A. donax* an ideal energy crop.
Here, we generated the first two chromosome-scale assemblies of the *A. donax* genome
using HiFi reads and Hi-C technology. According to the LAI score of the assemblies, the
assembly of 0004 reached to the criterion of reference quality, and assembly of 0408 also
approached to the reference quality (Table 1). Based on the genome, we speculated that
common *A. donax* is a triploid (2n = 3x = 108), and the aneuploid 0004 were generated

because of the abnormal replication of four chromosomes (3x + 4 = 112). In contrast, the genome of emerald *A. donax* is far more complicated. The chromosome base number (about 70) is more than the total chromosome number 62 (Figure 2), and most chromosomes of 0004 can aligned with two chromosomes of 0408. Therefore, we guess emerald *A. donax* is allodiploid (2n = 2x =72), and 0408 is an aneuploid with abnormal chromosome deletion (2x -10 = 62).

4.2 The reference genome will contribute to genetic improvement of A. donax

Clonal propagation crops like potato often showed small genetic gains. Recently, using precise genome design, vigorous F_1 hybrids potatoes has been developed [67]. Similar to potato, *A. donax* is completely agamic and showed a narrow genetic variability. To accelerate genetic improvement of *A. donax* in future, we need to elucidate the mechanism of infertility of *A. donax*. In addition, genetic transformation of *A. donax* encountered difficulties, and no transgenic *A. donax* reported until now. We speculate the difficulty of genetic transformation of *A. donax* lied in the complexity of *A. donax* genome. Several developmental regulators, including BBM, WUS, GRF4 and GIF were reported to improve transformation efficiency in crops like corn and wheat [68]. Therefore, cloning key developmental regulators in *A. donax* may help to break down technical barriers in genetic transformation of *A. donax*. To overcome the above two difficulties, a reference genome is indispensable.

Until now, little is known about the *A. donax* genome, except a few transcriptome studies [69,70,71,72]. Our result showed that *A. donax* has a complex genome that contains a large number of genes, many of which are involved in important biological processes such as growth, development, and stress response. Mining important genes from the reference genome will undoubtedly contribute to genetic improvement of *A. donax*.

In conclusion, *A. donax* is a valuable plant species with numerous characteristics that make it an attractive resource for various applications. Its fast growth, high biomass production, and environmental tolerance make it ideal for bioenergy production and environmental remediation. The plant's genetic diversity also offers many opportunities for genetic improvement and crop breeding. Research on *A. donax* continues to uncover new insights into the plant's biological properties and the potential for its utilization in different industries. Future studies will undoubtedly uncover more valuable information and offer even more exciting opportunities to take advantage of this remarkable plant species.

Authorship contribution statement

Mengmeng Ren: Data analysis and writing. Fupeng Liu and Xiaohong Han: Plant material collection.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

Not applicable.

Institutional Review Board Statement

Not applicable.

Data Availability

Informed Consent Statement

Not applicable.

Conflicts of Interest

The authors declare no conflict of interest.

Reference

1. Corno, L.; Pilu, R.; Adani, F. *Arundo donax* L.: a non-food crop for bioenergy and bio-compound production. *Biotechnol. Adv*. **2014**, *32*, 1535–1549.
2. Jámbor A.; Török, A. The Economics of *Arundo donax*—A Systematic Literature Review. *Sustainability* **2019**, *11*, 4225.
3. Pilu, R.; Manca, A.; Landoni, M. *Arundo donax* as an energy crop: pros and cons of the utilization of this perennial plant. *Maydica* **2013**, *58*.
4. Zhang, J.; Li, Y.; Zhang, C.; Jing, Y. Adsorption of malachite green from aqueous solution onto carbon prepared from *Arundo donax* root. *J. Hazard. Mater*. **2008**, *150*, 774–782.
5. Angelini, L.G., Ceccarini, L.; Bonari, E. Biomass yield and energy balance of giant reed (*Arundo donax* L.) cropped in central Italy as related to different management practices. *Eur. J. Agron*. **2005**, *22*, 375–389.
6. Angelini, L.G.; Ceccarini, L.; Nasso, N.; Bonari, E. Comparison of *Arundo donax* L. and *Miscanthus* x *giganteus* in a long-term field experiment in Central Italy: Analysis of productive characteristics and energy balance. *Biomass & Bioenergy* **2009**, *33*, 635–643.
7. Nasso, N.N.O.D.; Roncucci, N.; Bonari, E. Seasonal Dynamics of Aboveground and Belowground Biomass and Nutrient Accumulation and Remobilization in Giant Reed (*Arundo donax* L.): A Three-Year Study on Marginal Land. *Bioenergy Research* **2013**, *6*, 725–736.
8. Nackley, L.L.; Kim, S.H. A salt on the bioenergy and biological invasions debate: salinity tolerance of the invasive biomass feedstock *Arundo donax*. *GCB Bioenergy* **2015**, *7*, 752–762.
9. Sánchez, E.; Scordia, D.; Lino, G. Salinity and Water Stress Effects on Biomass Production in Different *Arundo donax* L. Clones. *BioEnergy Res*. **2015**, *8*, 1461–1479.
10. Mirza, N.; Mahmood, Q.; Pervez, A.; Ahmad, R.; Farooq, R.; Shah, M.M.; Azim, M.R. Phytoremediation potential of *Arundo donax* in arsenic-contaminated synthetic wastewater. *Bioresource Technol*. **2010**, *101*, 5815–5819.
11. Mirza, N.; Pervez, A.; Mahmood, Q.; Shah, M.M.; Shafqat, M.N. Ecological restoration of arsenic contaminated soil by *Arundo donax* L. *Ecol. Eng*. **2011**, *37*, 1949–1956.
12. Papazoglou, E.G.; Karantounias, G.A.; Vemmos, S.N.; Bouranis D.L. Photosynthesis and growth responses of giant reed (*Arundo donax* L.) to the heavy metals Cd and Ni. *Environ. Int*. **2005**, *31*, 243–249.
13. Ahmad, R.; Liow, P.S.; Spencer. D.F.; Jasieniuk, M. Molecular evidence for a single genetic clone of invasive *Arundo donax* in the United States. *Aquat. Bot*. **2008**, *88*, 113–120.
14. Danelli, T.; Laura, M.; Savona, M.; Landon, M.; Adani, F.; Pilu, R. Genetic Improvement of *Arundo donax* L.: Opportunities and Challenges. *Plants* **2020**, *9*, 1584.
15. Malone, J.M.; Virtue, J.G.; Williams, C.; Preston, C. Genetic diversity of giant reed (*Arundo donax*) in Australia. *Weed Biol. Manag*. **2017**, *17*.
16. Tarin, D.; Pepper, A.E.; Goolsby, J.A.; Moran, P.J.; Arquieta, A.C.; Kirk, A.E.; Manhart, J.R. Microsatellites Uncover Multiple Introductions of Clonal Giant Reed (*Arundo donax*). Invas. *Plant Sci. Mana*. **2013**, *6*, 328–338.

17. Bucci, A.; Cassani, E.; Landoni, M.; Cantaluppi, E.; Pilu, R. Analysis of chromosome
number and speculations on the origin of *Arundo donax* L. (Giant Reed). *Cytolo. Genet.*
2013, *47*, 237–241.
18. Mariani, C.; Cabrini, R.; Danin, A.; Piffanelli, P.; Fricano, A.; Gomarasca, S.; Dicandilo,
M.; Grassi, F.; Soave, S. Origin, diffusion and reproduction of the giant reed (*Arundo*
donax L.): a promising weedy energy crop. *Ann. Appl. Biol*. **2010**, *157*, 191–202.
19. Clevering, O.A.; Lissner, J. Taxonomy, chromosome numbers, clonal diversity and
population dynamics of *Phragmites australis*. *Aquat. Bot*. **1999**, *66*, 249–250.
20. Hunter, A.W.S. A Karyosystematic investigation in Gramineae. *Can. J. Res*. **1934**, *11*,
213–241.
21. Pizzolongo, P. Osservazioni cariologiche su *Arundo donax* e *Arundo plinii*. *Annuali Bot.*
1962, *27*, 173–187.
22. Christopher, J.; Abraham, A. Studies on the cytology and phylogeny of South Indian
grasses I. Subfamilies Bambusoideae, Oryzoideae, Arundinoideae and Festucoideae.
Cytologia **1971**, *36*, 579–594.
23. Haddadchi, A.; Gross, C.L.; Fatemi, M. The expansion of sterile *Arundo donax*
(Poaceae) in southeastern Australia is accompanied by genotypic variation. *Aquat.*
Bot. **2013**, *104*, 153–161.
24. Wang, Y.; Yu, J.; Jiang, M.; Lei, W.; Zhang, X.; Tang, H. Sequencing and Assembly of
Polyploid Genomes. *Methods Mol. Biol.* **2023**, *2545*, 429–458.
25. Peng, Y., Yan, H., Guo, L., Deng, C.; Wang, C., Wang, Y., Kan, L.; Zhou, P.; Y, K.;
Dong, X.; Liu, X.; Su, Z.; Peng, Y.; Zhao, J.; Deng, D.; Xu, Y.; Li, Y.; Jiang, Q.; Li, Y.;
Wei, L.; Wang, J.; Ma, J.; Hao, M.; Li, W.; Kang, H.; Peng, Z.; Liu, D.; Jia, J.; Zheng,
Y.; Ma, T.; Wei, Y.; Lu, F.; Ren, C. Reference genome assemblies reveal the origin and
evolution of allohexaploid oat. *Nat. Genet.* **2022**, *54*, 1248–1258.
26. Sun, H.; Jiao, W.B.; Krause, K.; Campoy, J.A.; Goel, M.; Folz-Donahu, K.; Kukat, C.;
Huettel, B.; Schneeberger, K. Chromosome-scale and haplotype-resolved genome
assembly of a tetraploid potato cultivar. *Nat. Genet.* **2022**, *54*, 342–348.
27. Chen, H.; Zeng, Y.; Yang, Y.; Huang, L.; Tang, B.; Zhang, H.; Hao, F.; Liu, W.; Li, Y.;
Liu, Y.; Zhang, X.; Zhang, R.; Zhang, Y.; Li, Y.; Wang, K.; He, H.; Wang, Z.; Fan, G.;
Yang, H.; Bao, A.; Shang, Z.; Chen, J.; Wang, W.; Qiu Q. Allele-aware chromosome-
level genome assembly and efficient transgene-free genome editing for the
autotetraploid cultivated alfalfa. *Nat. Commun.* **2020**, *19*, 2494.
28. Guan, D.; McCarthy, S.A.; Wood, J.; Howe, K.; Wang, Y.; Durbin, R. Identifying and
removing haplotypic duplication in primary genome assemblies. *Bioinformatics* **2020**,
36, 2896-2898.
29. Jike, W.; Li, M.; Zadra, L.; Barbaro, E.; Sablok, G.; Bertorelle, G.; Rota-Stabelli, O.;
Varotto, C. Phylogenomic proof of Recurrent Demipolyploidization and Evolutionary
Stalling of the "Triploid Bridge" in *Arundo* (Poaceae). *Int. J. Mol. Sci.* **2020**, *21*, 5247.
30. Bayani, J.; Squire, J.A. Fluorescence *in situ* Hybridization (FISH). *Current Protocols*
in Cell Biology **2004**, Chapter 22.
31. Jiang, J. Fluorescence *in situ* hybridization in plants: recent developments and future
applications. *Chromosome Res.* **2019**, *27*, 153–165.
32. Dolezel, J.; Greilhuber, J.; Suda, J. Estimation of nuclear DNA content in plants using

flow cytometry. *Nat. Protoc.* **2007**, *2*, 2233–2244.

33. Kokot, M.; Dlugosz, M.; Deorowicz, S. KMC 3: counting and manipulating k-mer statistics. *Bioinformatics* **2017**, *33*, 2759-2761.

34. Sun, H.; Ding, J.; Piednoël, M.; Schneeberge, K. FindGSE: Estimating genome size variation within human and Arabidopsis using k -mer frequencies. *Bioinformatics* **2018**, *34*: 550–557.

35. Cheng, H., Concepcion, G.T.; Feng X.; Zhang, H.; Li, H. Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm. *Nat. Methods* **2021**, *18*, 1–6.

36. Simão, F.A.; Waterhouse, R.M.; Ioannidis, P.; Kriventseva, E.V.; Zdobnov, E.M. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **2015**, *31*, 3210–3212.

37. Parra, G.; Bradnam, K.; Korf, I. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* **2007**, *23*, 1061–1067.

38. Li, H.; Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **2010**, *26*, 589-595.

39. Li, H.; Handsaker, B. Wysoker A, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **2009**, *25*: 2078-2079.

40. Danecek, P.; McCarthy, S.A. BCFtools/csq: haplotype-aware variant consequences. *Bioinformatics* **2017**, *33*, 2037–2039.

41. Ou, S.; Chen, J.; Ning, J. Assessing genome assembly quality using the LTR Assembly Index (LAI). *Nucleic Acids Res*. **2018**, *46*, e126.

42. Servant, N.; Varoquaux, N.; Lajoie, B.R.; Viara, E.; Chen, C.; Vert, J.P.; Heard, E.; Dekker, J.; Barillot, E. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol*. **2015**, *16*, 259.

43. Chen, S.; Zhou, Y.; Chen, Y.; Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **2018**, *34*, i884–i890.

44. Langmead, B.; Salzberg, S. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **2012**, *9*, 357-359.

45. Burton, J.N.; Adey, A.; Patwardhan, R.P.; Qiu, R.; Kitzman, J.O.; Shendure, J. Chromosome-scale scaffolding of *de novo* genome assemblies based on chromatin interactions. *Nat. Biotechnol*. **2013**, *31*, 1119–1125.

46. Wang, X.; Wang, L. GMATA: An Integrated Software Package for Genome-Scale SSR Mining, Marker Development and Viewing. *Front. Plant Sci*. **2016**, *7*, 1350.

47. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res*. **1999**, *27*, 573–580.

48. Han, Y.; Wessler, S.R. MITE-Hunter: a program for discovering miniature inverted-repeat transposable elements from genomic sequences. *Nucleic Acids Res.* **2010**, *38*: e199.

49. Xu, Z.; Wang, H. LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. *Nucleic Acids Res.* **2007**, *35* (Web Server issue), W265–W268.

50. Ellinghaus, D.; Kurtz, S.; Willhoeft, U. LTRharvest, an efficient and flexible software for *de novo* detection of LTR retrotransposons. *BMC Bioinformatics* **2008**, *9*, 18.

50. Ou, S.; Jiang, N. LTR_retriever: A Highly Accurate and Sensitive Program for Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol.* **2018**, *176*,

597   1410–1422.

51. Chen, N.S. Using RepeatMasker to identify repetitive elements in genomic sequences. *Curr. Protoc. Bioinformatics* **2004**, *4*, Unit 4.10.

52. Keilwagen, J.; Wenk, M.; Erickson, J.L.; Schattat, M.H.; Grau, J.; Hartung, F. Using intron position conservation for homology-based gene prediction. *Nucleic Acids Res.* **2016**, *44*, e89.

53. Dobin, A.; Davis, C.A.; Schlesinger, F., Jorg, Drenkow.; Zaleski, C.; Jha, S.; Batut, P.; Chaisson, M.; Gingeras, T.R. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **2013**, *29*, 15–21.

54. Kovaka, S.; Zimin, A.V.; Pertea, G.M.; Razaghi, R.; Salzberg, S.L.; Pertea, M. Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome Biol.* **2019**, *20*, 278.

55. Haas, B.J.; Salzberg, S.L.; Zhu, W.; Pertea, W.; Allen, J.E.; Orvis, J.; White, O.; Buell, C.R.; Wortman, J.R. Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.* **2008**, *9*, R7.

56. Mario, S.; Burkhard, M. AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Res.* **2005**, *33* (Web Server issue), W465–W467.

57. Haas, B.J.; Salzberg, S.L.; Zhu, W.; Pertea, M.; Allen, J.E; Orvis, J.; White, O.; Buell, C.R.; Wortman, J.R. Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome. Biol.* **2008**, *R7*.

58. Zdobnov, E.M.; Apweiler, R. InterProScan--an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* **2001**, *17*, 847–848.

59. Chan, P.P.; Lin, B.Y.; Mak, A.J. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Nucleic Acids Res. 1997, 25, 955–964.

60. Nawrocki, E.P.; Eddy, S.R. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* **2013**, *29*, 2933-2935.

61. Lagesen, K.; Hallin, P.; Rødland, E.A.; Stærfeldt, H.H.; Rognes, T.; Ussery, D.W. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* **2007**, *35*, 3100-3108.

62. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **2018**, *34*, 3094–3100.

63. Patz, J.A.; Frumkin, H.; Holloway, T.; Vimont, D.J.; Haines, A. Climate Change:challenges and opportunities for global health. *JAMA* **2014**, *312*, 1565-1580.

64. Walter, V.R.; Mariam, K.A.; Christopher B.F. The future of bioenergy. *Glob Chang Biol.* **2020**, *26*, 274–286.

65. Tang, Y.; Xie, J.S.; Geng, S. Marginal Land-based Biomass Energy Production in China. *J. Integr. Plant Biol.* **2010**, *52*, 112–21.

66. Zhang, C.; Yang, Z.; Tang, D.; Zhu, Y.; Wang, P.; Li, D.; Zhu, G.; Xiong, X.; Shang, Y.; Li, C.; Huang, S. Genome design of hybrid potato. Cell 2021, 184, 3873–3883.e12.

67. Chen, Z.; Debernardi, J.M.; Dubcovsky, J.; Gallavotti, A. Recent advances in crop transformation technologies. *Nat. Plants* **2022**, *8*, 1343–1351

68. Evangelistella, C.; Valentini, A.; Ludovisi, R.; Firrincieli, A.; Fabbrini, F.; Scalabrin, S.;
Cattonaro, F.; Morgante, M.; Mugnozza, G.S.; Keurentjes, J.J.B.; Harfouche, A. *De*
novo assembly, functional annotation, and analysis of the giant reed (*Arundo donax*
L.) leaf transcriptome provide tools for the development of a biofuel feedstock.
Biotechnol. *Biofuels* **2017**, *10*, 138.
69. Fu, Y.; Poli, M.; Sablok, G.; Wang, B.; Liang, Y.; Porta, N.L.; Velikova, V.; Loreto, F., Li,
M., Varotto, C. Dissection of early transcriptional responses to water stress in *Arundo*
donax L. by unigene-based RNA-seq. *Biotechnol. Biofuels* **2016**, *9*, 54.
70. Sablok, G.; Fu, Y.; Bobbio, V.; Laura, M.; Rotino, G.L.; Bagnaresi, P.; Allavena, A.;
Velikova, V.; Viola, R.; Loreto, F.; Li, M.; Varotto, C. Fuelling genetic and metabolic
exploration of C3 bioenergy crops through the first reference transcriptome of *Arundo*
donax L.. *Plant Biotechnol. J*. **2014**, *12*, 554–567.
71. Sicilia, A.; Testa, G.; Santoro, D.F.; Cosentino, S.L.; Piero, A.R.L. RNASeq analysis of
giant cane reveals the leaf transcriptome dynamics under long-term salt stress. *BMC*
Plant Biol. **2019**, *19*.

682 **Figure legend**

Common    Emerald    Versicolor

683
684 **Figure 1.** Morphology of the common, emerald and versicolor *A. donax*. Pictures were
685 taken in September 2022 in Shishou, Hubei province (E112.40°, N29.73°). Giant panicles
686 can be observed in common and versicolor *A. donax*, while no panicle was observed in
687 emerald *A. donax*. Bar = 50 cm.

688
689 **Figure 2.** Karyotype analysis of 0004 and 0408. (A, D) DAPI staining of chromosomes of
690 0004 (A) and 0408 (D). (B, E) *In situ* hybridization of telomere repeat sequences of 0004
691 (B) and 0408 (E). (C, F) rDNA fluorescence *in situ* hybridization of chromosomes of 0004
692 (C) and 0408 (F). Red and green arrows denote the 5SrDNA and 18SrDNA hybridization
693 signals. Bar = 50 μm.

were obtained in 10 Kb successively window.

Supplementary Figure S3. Assessment of genome assembly completeness. (A-B)
Genome assembly completeness was assessed using BUSCO for 0004 (A) and 0408 (B).
(C-D) Conserved gene was assessed using CEGMA for 0004 (C) and 0408 (D).

Supplementary Figure S4. Heatmap of primary Hi-C scaffolding of 0004 (left) and 0408
(right). The chromosome basic number was assumed to 56 for 0004 and 31 for 0408. The
heatmap coordinates represent chromosomes, and the color of each point represents the
log value of the corresponding genome bin pair interaction intensity, which increases
sequentially from yellow to black.

723
Supplementary Figure S5. Genome annotation based on public databases. (A-B)
Species distribution for NR-based annotation for 0004 (A) and 0408 genome (B). (C-D)
KEGG classification of 0004 (C) and 0408 genome (D) annotation. (E-F) KOG function
classification of 0004 (E) and 0408 (F) genome annotation. (G-H) GO domain of 0004 (G)
and 0408 (H) genome annotation.

Supplementary Figure S6. Statistics of genome annotation for 0004 (left) and 0408
genome based on five databases, including Non-Reduntant Protein Database (NR), Kyoto
Encyclopedia of Gene and Genomes (KEGG), Eukaryotic Orthologous Groups of protein
(KOG), GO and Swissprot.

Supplementary Figure S7. Comparative statistics of gene information of related species.
Comparison of gene information of 0004 (left column) and 0408 (right column) with five
Poaceae plants (*Oryza sativa*, *Saccharum spontaneum*, *Sorghum bicolor*, *Zea mays* and
Triticum aestivum) including distribution of gene length (A-B), CDS length (C-D), exon
length (E-F), exon number (G-H), intron length (I-J) and intron number (K-L).
740

741 **Supplementary table 1** Statistics of RNA-seq data

Sample name	Total reads	Total mapped	Unique mapped	Multiple mapped
0004-root	75,570,094	69,694,640 (92.23%)	67,979,118 (89.96%)	1,715,522 (2.27%)
0004-leaf	73,310,846	68,078,324 (92.86%)	66,040,232 (90.08%)	2,038,092 (2.78%)
0004-panicle	121,085,174	109,619,650 (90.53%)	107,277,684 (88.60%)	2,341,966 (1.93%)
0004-bud	102,532,136	95,570,548 (93.21%)	93,230,426 (90.93%)	2,340,122 (2.28%)
0408-root	112,487,910	110,342,744 (98.09%)	95,485,032 (84.88%)	14,857,712 (13.21%)
0408-leaf	81,606,088	79,939,116 (97.96%)	68,384,754 (83.80%)	11,554,362 (14.16%)
0408-panicle	62,558,910	59,462,320 (95.06%)	50,570,542 (80.84%)	8,891,778 (14.22%)
0408-bud	89,793,602	87,915,064 (97.91%)	75,257,540 (83.81%)	12,657,524 (14.10%)

742