Dating the bacterial tree of life based on ancient symbiosis

Sishuo Wang[1,2], Haiwei Luo[1,3*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory
of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]Department of Microbiology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[3]Institute of Environment, Energy and Sustainability, The Chinese University of Hong Kong,
Shatin, Hong Kong SAR

[*]**Corresponding author:**

Haiwei Luo
School of Life Sciences, The Chinese University of Hong Kong
Hong Kong SAR
Phone: (+852) 39436121
E-mail: haiweiluo@cuhk.edu.hk

Key words: molecular clock, bacterial tree of life, timetree, symbiosis, relative time

constraint, early life evolution

Abstract

Obtaining a timescale for bacterial evolution is crucial to understand early life evolution but is difficult owing to the scarcity of bacterial fossils and absence of maximum age constraints of the available fossils. Here, we introduce multiple new time constraints to calibrate bacterial evolution based on ancient symbiosis. This idea is implemented using a bacterial tree constructed with mitochondria-originated genes where the mitochondrial lineage representing eukaryotes is embedded within Proteobacteria, such that the date constraints of eukaryotes established by their abundant fossils are propagated to ancient co-evolving bacterial symbionts and across the bacterial tree of life. Importantly, we formulate a new probabilistic framework that considers uncertainty in inference of the ancestral lifestyle of modern symbionts to apply 19 relative time constraints (RTC) each informed by host-symbiont association to constrain bacterial symbionts no older than their eukaryotic host. Moreover, we develop an approach to incorporating substitution mixture models that better accommodate substitutional saturation and compositional heterogeneity for dating deep phylogenies. Our analysis estimates that the last bacterial common ancestor (LBCA) occurred approximately 4.0-3.5 billion years ago (Ga), followed by rapid divergence of major bacterial clades. It is robust to alternative root ages, root positions, tree topologies, fossil ages, ancestral lifestyle reconstruction, gene sets, among other factors. We further obtain a taxonomically high-resolution bacterial timetree, which serves as a foundation for testing hypotheses regarding bacterial diversification and its correlation with geobiological events across different timescales.

Introduction

Bacteria are the dominating form of life in the first two thirds of the history of our planet, and their evolution and diversification have greatly shaped the geosphere of Earth (Falkowski et al. 2008). Many hypotheses have been proposed to interpret outstanding questions like how bacteria affected the evolution of biogeochemical cycles, and when major innovations of bacteria such as photosynthesis occurred and why they did not appear earlier (Battistuzzi et al. 2004; Battistuzzi and Hedges 2009; Louca et al. 2018). Interpreting these questions depends on temporal links between bacterial evolution and the corresponding geobiological events. However, efforts to date the early evolution of bacteria based on relaxed molecular clock (Zuckerkandl and Pauling 1965; Drummond et al. 2006; Yang and Rannala 2006) often fail to reach convergence. A major challenge is that bacteria have rarely left calibrations well suited for molecular dating, i.e., fossils attributable to or biomarkers diagnostic for certain clades (Battistuzzi and Hedges 2009). Further, because fossils or biomarkers by themselves provide only a lower minimum bound (Benton and Donoghue 2007; Marshall 2019), there are no reliable maximum time constraints to calibrate bacterial evolution. Accordingly, the time estimates heavily depend on the maximum age assigned to the root, i.e., the LBCA, whose origin time involves even more debate (Wang and Luo 2021; Moody et al. 2022). Moreover, previous studies in dating the early evolution of bacteria (Battistuzzi et al. 2004; Battistuzzi and Hedges 2009; Wang and Luo 2021) are mostly based on site-homogeneous amino acid substitution models that do not well account for among-site compositional heterogeneity and substitutional saturation (Gouy et al. 2015). In deep-time phylogenetics, this may lead to an underestimate of branch length (expected number of substitutions per site) (Moody et al. 2022), which potentially affects time estimation.

Recent studies have used eukaryotic fossils to date the evolution of Proteobacteria (Wang and Luo 2021) and Cyanobacteria (Sánchez-Baracaldo et al. 2017; Shih et al. 2017)

where mitochondrial and plastid-endosymbiosis are believed to have occurred, respectively. These successes imply the value of the abundant eukaryotic fossils in timing bacterial evolution by molecular clock, but more temporal information of eukaryote evolution is waiting to be harvested. One such source is bacteria-eukaryote symbiosis: as such symbioses occurred in multiple ancient bacterial lineages (Toft and Andersson 2010; Perreau and Moran 2021), it is tempting to fit it into a recently developed framework called relative time constraint (RTC) dating (Davín et al. 2018; Magnabosco et al. 2018) which constrains the order of divergence by assuming the age of some nodes of the tree to be younger than others. Despite some encouraging progress, the current implementation of this method treats all RTCs as if they are known with full certainty. However, this assumption might be oversimplified because of the uncertainty in identifying the relative time order based on horizontal gene transfer (HGT) (Ravenhall et al. 2015) or other evolutionary events. Here, we addressed the above challenges by leveraging combined evidence of sequences, fossils, and ancient symbiosis to calibrate an integrative timescale of bacterial evolution. We developed and implemented a framework to propagate date constraints of eukaryotes established by abundant fossils to ancient co-evolving bacterial symbionts and other bacteria by requiring that the origin of symbionts postdates their hosts' while accommodating the uncertainty in ancestral host reconstruction. We established timelines of bacterial evolution at different taxonomy resolutions and discussed how they may affect our views of the evolution of early life and habitat.

Results

A new framework for dating deep-time bacterial evolution informed by host-symbiont association

We overcame the paucity of bacterial fossils in dating the bacterial tree by leveraging the

temporal information of eukaryotes recorded in their abundant fossils. Briefly, by

incorporating eukaryotes into bacterial phylogeny as mitochondrial lineages affiliated with α-

proteobacteria based on mitochondrial endosymbiosis (Martijn et al. 2018; Muñoz-Gómez et

al. 2022), it is tempting to co-estimate the ages of both the eukaryotic host and bacterial

symbiont (Wang and Luo 2021) while applying the recently developed RTC-based molecular

dating (Davín et al. 2018; Magnabosco et al. 2018; Szöllõsi et al. 2022). In the context of

host-symbiont association, this means constraining the origin time of the symbiont's crown

group to be no earlier than that of the host's total group, thereby providing maximum time

constraints to symbionts whose temporal information may further be propagated across

bacteria.

It is well possible that the ancestor of modern symbionts adapted to a different host or

even a free-living lifestyle. We formulate a three-step approach accounting for the possibility

of host shift of symbionts through ancestral state reconstruction (ASR) and fit it into RTC-

based dating in a probabilistic framework, which we call probability-based RTC (pRTC; see

Methods for details). First, ASR is performed to estimate the probabilities that the ancestral

nodes of modern symbionts are associated with different hosts or are free-living. Second,

MCMC molecular dating is performed using mitochondria-originated genes with fossil

calibrations but without RTC to generate posterior samples of timetrees where the ages of

symbionts and eukaryotic hosts, which are placed as the mitochondrial clade sister to α-

proteobacteria (Martijn et al. 2018; Wang and Luo 2021; Muñoz-Gómez et al. 2022), are co-

estimated. Third, the posterior samples of timetrees collected in Step 2 are examined to

remove those where the host's age is younger than the symbiont's, weighted by the

probability of different states (hosts) taken by the ancestor of modern symbionts estimated in

Step 1, using a rejection sampling approach. The idea is further illustrated in Fig. 1A.

Assume that ASR infers the ancestral lifestyles of "symbiont1" as $P(S_1 = mammal) = 0.9$

and $P(S_1 = freeliving) = 0.1$, and for "symbiont2" as $P(S_1 = mammal) = 0.5$

and $P(S_1 = freeliving) = 0.5$. For the two timetrees in the posterior samples shown in Fig.

1A, the probabilities of acceptance are $(0.9 \times 1 + 0.1 \times 1) \times (0.5 \times 1 + 0.5 \times 1) = 1$, and

$(0.9 \times 1 + 0.1 \times 1) \times (0.5 \times 0 + 0.5 \times 1) = 0.5$, respectively (see Eqs. 3-5 in Methods). If

the above procedure is performed for all posterior timetrees collected from Step 2, the

posterior dates of all bacterial lineages, particularly the symbiont, might shift toward the

present due to the use of RTC informed by host-symbiont association (Fig. 1A). As ancient

bacterial symbionts are distributed in different deep-branching lineages (Toft and Andersson

2010; McCutcheon and Moran 2012), this method has a great potential to date the bacterial

tree of life.

Integrating eukaryote temporal information by a Bayesian sequential approach

We reconstructed the phylogeny of bacteria using the 265 genomes and 60 genes carefully

selected in a comprehensive phylogenomics study (Coleman et al. 2021) using the best-fit

site-heterogeneous profile mixture model LG+G+C60 (Table S1A; Note S1). We applied the

time non-reversible model (Naser-Khdour et al. 2022) which infers the root of the bacterial

tree as part of maximum-likelihood phylogenetic reconstruction where different root

positions may have different likelihoods. This method does not depend on any outgroup,

thereby avoiding long branch attraction artefacts caused by using highly divergent sequences

to root deep phylogenies. Our analyses rooted the bacterial phylogeny in the neighbourhood

of a clade consisting of Fusobacteriota and DST (Deinococcota, Synergistota, and

Thermotogota). The analyses also supported the division of most bacterial lineages into

Gracilicutes and Terrabacteria, and placed the CPR (Patescibacteria) clade (Fig. S1), initially

suggested as the earliest-split bacterial group (Hug et al. 2016; Zhu et al. 2019), as a late-split

clade sister to Chloroflexi (Fig. S1; Table S2; Note S2). This is generally consistent with a

recent study which roots the bacterial tree using an independent method based on gene-

species tree reconciliation (Coleman et al. 2021).

To apply pRTC based on host-symbiont co-evolution, we integrated eukaryotes and

bacteria in a unified framework to co-estimate their times using the mitochondrial

endosymbiosis-based dating strategy (Wang and Luo 2021). Apparently, the ages of

symbionts are most strongly constrained if the host date is inferred with a high precision, thus

less uncertainty. We therefore took the advantages of the genome-scale data of eukaryotes by

employing a Bayesian sequential approach in which the posterior dates of a small number of

organisms with many genes are estimated to serve as the time priors of a second analysis with

many organisms but a few genes (dos Reis et al. 2012; Álvarez-Carretero et al. 2022). Note

that sequential Bayesian molecular dating is conceptually distinct from secondary calibrations

(using time estimates from other studies as calibrations) by accommodating the uncertainty in

time estimation in the first-step analysis within a Bayesian framework (Note S1.2). We

estimated the divergence times of 29 representative eukaryotes with 11 fossil calibrations

(Figs. S2-S3) and 320 orthologs conserved across eukaryotes (Strassert et al. 2021). The

posterior dates of eukaryote nodes were in agreement with most studies (Parfrey et al. 2011;

Betts et al. 2018; Gibson et al. 2018; Wang and Luo 2021). Next, we approximated the

posterior times of each node in the eukaryote tree by a best-fit probability distribution (Fig.

S4; Note S1.2.3), which were further used to calibrate the eukaryote nodes for dating the

bacterial tree where eukaryotes were placed as a subtree sister to α-proteobacteria (Martijn et

al. 2018; Wang and Luo 2021; Muñoz-Gómez et al. 2022).

A time-calibrated bacterial phylogeny based on ancient symbiosis

To apply the pRTC dating approach based on ancient symbiosis, we first performed

comprehensive MCMC-based ASR to infer the ancestral lifestyles of seven well-known

ancient clades where symbionts are widely distributed, four at a phylum level (Spirochaetota,

Tenericutes, Chlamydiae, and Elusimicrobiota) and three at an order level (Rickettsiales,

Holosporales, and Legionellales). The ASR was performed using the 16S rRNA gene, which

best captures bacterial lifestyle diversity by including environmental and uncultured samples,

based on the lifestyle information of modern members within each of the ancient bacterial

clades mainly consisting of symbionts (Fig. S5). We additionally included two genus-level

symbionts (*Buchnera* and *Blattabacterium*) which are classical host-symbiont co-evolution

examples (Perreau and Moran 2021). A total of 19 ancestral nodes consisting of 28 symbiont

lineages from the above nine symbiont clades were selected to represent the host-symbiont

association-based RTCs in subsequent analyses (Figs. 2, S6; Note S3.1).

Further, we identified 19 genes well conserved across bacteria (Figs. S7-S8) from 108

genes with a mitochondrial origin reported in a recent study (Muñoz-Gómez et al. 2022). We

performed molecular clock analysis with these 19 genes on a 148-genome focal dataset

consisting of 29 eukaryotes, 28 symbionts, and 91 non-symbionts. The phylogeny was based

on the tree reconstructed above using the non-reversible model and with eukaryotes placed as

the sister to α-Proteobacteria. The posterior dates of eukaryotic lineages inferred in the first-

step sequential analysis above, and five lineage-specific fossil- or biomarker-calibrations

within bacteria (see Methods and Note S3.2), were used as time calibrations. We employed

MCMCTree's approximate likelihood method, which obtains a 1000× speed-up without loss

of accuracy compared with the exact likelihood method (dos Reis and Yang 2011), well-

suited for our large-scale molecular clock analysis. However, the amino acid substitution

models (e.g. LG+G) used to calculate branch lengths in MCMCTree (and many other

molecular dating software such as BEAST and MrBayes) do not consider across-site

compositional heterogeneity and often show poor model fit in reconstructing deep

phylogenies (Gouy et al. 2015; Moody et al. 2022).

We therefore developed a bootstrap-based approach to incorperating branch length

estimates inferred under the site-heterogeneous profile mixture substitution model

LG+G+C60, which better accommodates issues in deep-time phylogenetics like substitution

saturation (Moody et al. 2022), into MCMCTree's approximate likelihood method (Fig. 1B;

see also Methods). Briefly, maximum-likelihood trees were built for 1000 bootstrap

alignments, and the inverse of the bootstrap covariance matrix of their branch lengths were

calculated to approximate the hessian matrix which is necessary for MCMCTree's

approximate method. Simulations showed considerable improvements in the estimates of

both ages and absolute rates by integrating site-heterogeneous models with this approach,

partcularly in the absence of internal calibrations or if the divergence times are older than 3.0

Ga (Figs. S9-S10; see Note S4 for details).

Next, we followed Eq. (5) (Methods) to apply a rejection sampling approach to filter out

the posterior timetrees obtained without applying RTC from Step 2, based on the probabilities

of the ancestor of modern symbionts being associated with different eukaryotic hosts or free-

living inferred from Step 1 (Figs. 1A, S8C). Collectively, using a carefully selected focal

dating scheme (Data S3), we estimated that the LBCA occurred at around 3648 Ma (95%

HPD [highest posterior density] 4111-3162 Ma) (Fig. 2), followed by the split between

Terrabacteria and Gracilicutes estimated to be 3629 Ma (95% HPD 4103-3167 Ma). Most

major clades, many of which are beyond a phylum level, diverged during 3500-3000 Ma

(Fig. 2). All selected symbiotic bacteria lineages (purple nodes in Fig. 2) originated no earlier

than 1700 Ma, approximately the appearance time of the LCA of eukaryotes.

We gradually included eukaryotic temporal constraints, the LG+G+C60 substitution

model, and host-symbiont association-based RTCs to improve date estimates of bacteria

(lower triangle in Fig. 3A). The inclusion of eukaryote's timing (*Strategy2*) led to a ~13%

decrease in the posterior dates compared with *Strategy1* where the five bacterial calibrations

were the only constraints (*Strategy1*). When the improved branch length estimates based on

the LG+G+C60 model was additionally included (*Strategy3*), those nodes between 3000 and

2000 Ma were estimated to be older by 7% (*Strategy2* vs. *Strategy3*). Further, when pRTC

was applied on the basis of *Strategy3*, the posterior dates of almost all nodes systematically

decreased ~5% (*Strategy3* vs. *Focal* in Fig. 3A), suggesting that the constraint on the time

order imposed by host-symbiont co-evolution was propagated through the bacterial tree of

life. Collectively, the combination of the use of time constraints provided by eukaryote

temporal information, site-heterogeneous profile mixture models, and host-symbiont

association-based RTCs estimated the divergence times of analysed bacteria ~15% younger

than those obtained without any of these settings (*Strategy1* vs. *Focal* in Fig. 3A; this

represents a net reduction and some effects cancelled out as illustrated above). As to the

posterior rate for each branch, the use of C60 led to an almost doubling of rate estimate

(upper triangle in Fig. 3A; *Strategy2* vs. *Strategy3*), indicating that the absolute rates are

largely underestimated by the compositionally homogeneous LG+G model. Moreover, when

we successively increased the maximum time prior of the root from 4500 Ma to 6000 Ma, the

posterior dates of bacterial lineages increased by more than 20% with *Strategy1*, but

remained almost unchanged with the focal scheme (Figs. 3B, S11A), indicating that the

(maximum) time constraints imposed by eukaryotes effectively constrained the timing of

bacterial evolution, such that their time estimates were robust to the root calibration where

much debate has arisen (Nisbet and Sleep 2001; Moody et al. 2022).

Accommodating the uncertainty in divergence time estimation

We evaluated the influence of multiple sources of uncertainties in molecular clock analysis

on the posterior dates. First, fossil-based calibrations are known to play a major role in

molecular dating (Ho and Duchêne 2014). The absolute time calibrations consisted of 11

eukaryotic ones, five from within bacteria, and one at the root (LBCA). Replacing the 1.047

Ga-old fossil of *Bangiomorpha*, the most widely recognized oldest crown-group eukaryote

(Betts et al. 2018; Gibson et al. 2018), by the recently discovered 1.6 Ga-old *Rafatazmia*

(Bengtson et al. 2017), pushed the LCA of eukaryotes from ~1600 Ma to ~2100 Ma (Fig.

S3), but had little impact on the time estimates of bacteria (*Euk1* in Fig. 3C). A similar

pattern was observed when the recently discovered 0.89 Ga-old sponge fossil instead of the

0.55 Ga-old one (Note S3.2) were alternatively used as the minimum age of the total-group

animals (*Euk2* in Fig. 3C). As to the five bacterial calibrations, the two cyanobacteria fossils

and the two biomarkers displayed considerable impacts on the posterior dates as their

removal led to a decrease of ~7% in time estimates (*noCyanoFossil* and *noBiomarker* in Fig.

3C). Replacing the 1.6 Ga-old Nostocales fossil calibration (Tomitani et al. 2006) by a more

disputable estimate based on the 2.0 Ga-old fossil (Amard and Bertrand-Sarfati 1997) led to

an increase in posterior dates by more than ~10% (Fig. S11B). Except the above, the use of

all alternative fossil calibrations returned similar time estimates of both eukaryotes and

bacteria (Figs. S11B, S12).

Second, another source of time constraint is the use of ASR-based RTC. In the above

analysis, the ancestral states (hosts) between nodes within the same tree were considered

independent with each other such that their joint probability can be approximated by the

multiplied marginal probability (Fig. S6B). To check if this approximation affected the final

time estimates, we directly estimated the joint probability of their ancestral states using a

stochastic character mapping (SCM) method (see Methods). This did not change the posterior

dates (*joint_prob* in Fig. 3C). Further, while we have conducted comprehensive work in

classifying the traits into different hosts or free-living, there exist different interpretations of

the phenotypic traits particularly free-living, which might affect ASR. Additionally, we tried

different cut-offs of 16S OTU and a different algorithm (maximum-likelihood) in ASR, all of

which resulted in highly consistent time estimates (Fig. 3C). Including 13 more symbiont

lineages returned similar time estimates (*more_RTCs* in Fig. S12), suggesting that adding

host association-based RTCs would not lead to further change in the posterior ages of

bacteria.

Third, depencence of the results on the phylogenomic tree topology was investigated.

When molecular clock analysis was performed with three alternative root positions, two near

the root (*root0*) used in the focal analysis (*root1* and *root2* in Fig. 3D; Fig. S1), and one

(*rootCPR*) at the CPR clade as suggested by earlier studies (Hug et al. 2016; Zhu et al. 2019),

the posterior dates remained similar for most clades, except those that were alternatively

placed as an early-split phylogenetic position, particularly the CPR. Alternative phylogenetic

positions of mitochondria or between eukaryotic groups (Fig. S8D-S8E) displayed minor

impacts on the posterior dates of bacteria (Fig. S11C).

Fourth, the use of different genes in dating may lead to different divergence time

estimates. Mitochondrial genes tend to evolve faster than their orthologs in bacterial genomes

(Martijn et al. 2018; Wang and Luo 2021). In theory, the issue can be accommodated by

relaxed molecular clock models, but this is not guaranteed (Drummond et al. 2006; Yang and

Rannala 2006). Also, MCMCTree assumes a single tree topology for all genes, which might

not always hold true due to HGT, unresolved paralogy, or limited phylogenetic signal of

individual genes. To minimize the bias caused by gene selection, we ranked the 19

mitochondria-originated genes by i) the relative difference in their estimated evolutionary

rates between eukaryotic and bacterial lineages, and ii) $\triangle$LL, an index to measure the

species- and gene-tree incongruence (Moody et al. 2022). We selected the 5, 10, and 15 top-

ranking (best) genes using each of the above two measures and repeated molecular clock

analyses (Fig. 3D). Additionally, we analysed the full set of 32 mitochondria-originated

genes (*all_genes* in Fig. 3D). Most analyses showed consistent results compared with using

the 19-gene dataset; the schemes using only the 5 top-ranking genes (schemes *top5*)

displayed larger uncertainty in the date estimates of certain nodes (e.g., MBDD), but this is

likely due to their small number of genes.

Fifth, other factors potentially affecting the dating analysis were examined. The largest

change in posterior dates were observed when the autocorrelated rates (AR) model, which

was consistently supported as the preferred model (Fig. S13), was replaced by the poorly

supported independent rates (IR) model: the divergence times of most lineages shifted toward

the present by ~20% (*IR* in Fig. 3C). The priors on the birth-death model which provides time

priors for nodes without time calibration and on the lineage-specific rate showed limited

impact (Fig. S11B). In addition, analysis with an independently sampled and expanded

dataset with 285 bacterial genomes (Coleman et al. 2021) showed similar time estimates

(*Secondary* in Fig. S12), suggesting that the time estimates presented in the main analysis

were not dependent on a specific taxon sampling. Note that the overall acceptance rate of the

rejection sampling was less than 5%, meaning that more than 95% posterior were removed in

Step 3. Because running time of MCMCTree is roughly proportional to the number of

partitions, to save time, all analyses were performed on a single partitioned alignment, but

similar patterns were found when two partitions were used (Fig. S14).

A taxonomically high-resolution timetree of bacteria

Based on the time estimates from the above analysis where most lineages were sampled at a

phylum level, we proceeded with a sequential approach to date a taxonomically higher-

resolution timetree of bacteria. Due to the large computational burden associated with

Bayesian-based molecular clock methods, we applied the fast dating program RelTime.

Spefically, we used lineages included in the 148-genome dataset as the backbone, and used

their posterior dates estimated in the above analysis as calibrations to further infer a full-scale

bacterial timetree comprising 8898 tips each representing a genus based on GDTB (r95)

classification (Figs. 4A, S15; Data S4). The three major clades of Proteobacteria, α-, β-, and

γ-Proteobacteria, were estimated to originate ~2.1, ~1.6, and ~1.9 Ga, respectively, greatly

postdating the origin of those previously known as δ-Proteobacteria whose origin time was

estimated no later than 3.0 Ga (Fig. 4A). This supports the recent proposal to exclude δ-

Proteobacteria from the phylum Proteobacteria (Waite et al. 2020), which was further

renamed as the MBDD group (Myxococcota, Bdellovibrionota, Desulfomonadota, and

Desulfobacterota) (Coleman et al. 2021). Firmicutes as a whole was the oldest phylum

concerning the crown group, originating at ~3.3 Ga. Family-, order-, class, and phylum-level

taxonomy groups were estimated to diverge at 0.90±0.31, 1.4±0.36, 2.1±0.46, 2.5±0.56 Ga

(mean ± standard deviation; Fig. 4B). Further, we found a significant correlation between

substitution rate of 16S rRNA gene and divergence time at the family level ($p = 1.98E-44$, F-

test; Fig. 4C). A simple linear regression analysis suggests an average substitution rate of

~0.015 nucleotides/site/100 Ma for 16S rRNA gene, but there are great differences between

lineages (Fig. 4C). In other words, the divergence time between bacteria may be implied by

comparing the sequence of 16S rRNA gene where 1% nucleotide difference roughly

corresponds to divergence time of 1/0.015=67 Ma. This indicates a longer time span for 1%

nucleotide difference than previously estimated, likely because the bacteria analysed in prior

studies are mainly fast-evolving endosymbionts (Moran et al. 1993; Ochman et al. 1999; Kuo

and Ochman 2009). The correlation was robust to linear mixed models where the rate

differnece of 16S rRNA gene evolution in different lineages was controlled for (Note S1.4;

Data S5).

Discussion

A major challenge in dating bacterial evolution is the paucity of credible calibrations (Wang

and Luo 2021; Moody et al. 2022), particularly the maximum time constraints. Previously,

this was usually done by assigning a somewhat arbitrary maximum age to the root (LBCA),

which greatly varies between studies due to the uncertainty in palaeobiological evidence and

phylogenetic relationship around the origin of life (Knoll 2015; Moody et al. 2022). We

addressed this problem by leveraging the timing information of eukaryotes based on ancient

symbiosis. This provides two additional layers of timing information besides bacterial

calibrations. First, we linked the evolution of bacteria and eukaryotes by mitochondrial

endosymbiosis. Because several eukaryotic calibrations have maximum time bounds

achieved based on the time estimates of the most recent geological formation that ought to

haven fossils of the clade of interest but that does not, this introduces many reliable

(maximum) time bounds to indirectly constrain bacterial evolution by informing the

substitution rates. Second, based on RTCs informed by host-symbiont association, the timing

of eukaryote evolution directly constrains the origin time of symbionts to be no earlier than

their eukaryotic host, which is further propagated across the bacterial phylogeny.

The idea of RTC in molecular dating has recently caught much attention (Davín et al.

2018; Magnabosco et al. 2018; Fournier et al. 2021). However, to our knowledge, none of the

available RTC applications has considered the uncertainty in the inference of RTC. In other

words, previously the relative time order established by horizontal gene transfer (i.e., gene

recipient lineage cannot be older than gene donor lineage) or any other forms of RTC was

assumed to be known with certainty, which might be oversimplified due to the uncertainties

in identifying these evolutionary events. Our new approach removes this assumption by

weighting the different ancestral hosts by their probability established in ASR through a

rejection sampling framework. This greatly extends the RTCs based on host-symbiont co-

evolution to dating the bacterial tree, which involves many ancient symbioses and large

uncertainty in assigning the ancestral host. In addition, prior studies that have applied the idea

of host-symbiont association in molecular dating typically used time estimates of the host

from other studies (i.e., secondary calibrations) to calibrate symbiont evolution, which may

cause wrong time estimates by propagating errors associated with the original time estimates

(Schenk 2016). For example, the estimated origin times of animals range from 1355 to 650

Ma according to 11 published recorded in the TimeTree database (Kumar et al. 2017).

Apparently, calibrating with different times of animals can lead to dramatically different time

estimates of the symbionts. Our strategy co-estimates the origin times of hosts and symbionts

based on mitochondrial endosymbiosis (Wang and Luo 2021), avoiding methodological and

data differences in comparing times estimates from different studies.

Great care should be taken when interpreting the estimated timeline of the bacterial tree

of life and reconstructed evolutionary events taking place anciently. Nonetheless, it is

remarkable to note the congruence between time estimates obtained using different time

constraints, different root positions, independent sets of genomes, and different settings in

molecular clock analysis. Our time estimate of LBCA is ~0.5 Ga younger than estimated in

some studies (Battistuzzi and Hedges 2009; Marin et al. 2017). As evident in Fig. 3A, this

might be due to the lack of enough maximum time constraints and the use of poorly fit

substitution models for inferring deep phylogenies in these early efforts. Our study also

improves molecular clock inferences by using a most-updated phylogeny of the bacterial tree

of life. For example, some groups once thought basal in the bacterial tree, e.g., Aquificae

(Battistuzzi and Hedges 2009) and CPR (Hug et al. 2016), are suggested as later-split

lineages in recent (Coleman et al. 2021; Moody et al. 2022) and the present study based on

improved phylogenetics methods, leading to different time estimates of corresponding

lineages.

The estimated timescale of bacterial evolution allows testing hypotheses involving early

life evolution. For example, It is generally believed that the late heavy bombardment likely

occurred on early Earth ~4.1-3.8 Ga, during which large asteroids intensely pummelled Earth,

melted most parts of its surface, and killed off any forms of life that might have begun to

emerge (Abramov and Mojzsis 2009; Boehnke and Harrison 2016). Because primordial form

of life likely had existed before LBCA, our estimates for the age of LBCA at 4.0-3.5 Ga hint

that life had survived the cataclysm, consistent with modelling studies showing that much of

Earth's crust and the microbes living therein were unlikely affected by the intense

bombardment (Abramov and Mojzsis 2009). The robustness of the posterior age of the LBCA

to the maximum time prior assigned to it (Fig. 3B) provides support to reject the hypothesis

that life originated extra-terrestrially (Nisbet and Sleep 2001), where the LBCA should be

much older than Earth' age. Further, under the most parsimonious hypothesis that oxygenic

cyanobacteria arose in the cyanobacterial stem lineage (Sánchez-Baracaldo and Cardona

2020; Fournier et al. 2021), our analysis placed the appearance of the first oxygenic

cyanobacteria between 3 Ga (total-group Cyanobacteria) and 2.5 Ga (crown-group

Cyanobacteria) (Fig. 2). This suggests that one cannot link stromatolites found in ancient

sediments dated ~3.5 Ga to oxygenic cyanobacteria despite their dominance in stromatolites

nowadays (Hofmann et al. 1999; Allwood et al. 2006). This is supported by recent evidence

showing that stromatolites preserved in Archaean rocks neither required oxygen-producing

cyanobacteria for formation (Schopf 2011) nor were necessarily biogenic (McLoughlin et al.

2008).

The new bacterial timetree also makes it possible to infer the mechanisms that drive the

origin of enzymes that involve oxygen. Accumulating evidence from isotope analyses

suggests the presence of oxygen on Earth back to 3.1 Ga (Crowe et al. 2013; Planavsky et al.

2014; Wang et al. 2018), which could be produced by cyanobacteria as mentioned above,

and/or abiotically from H_2O_2 (He et al. 2021), potentially leading to the formation of "oxygen

oasis" thus extracellular access to O_2. A recent study (Jabłońska and Tawfik 2021) tracked

the origin of most oxygen-utilizing and -producing enzymes families to the Gracilicutes-

Terrabacteria split, which according to our estimate is ~3.6 Ga, predating the above events

for ~0.5 Ga (Fig. 2). This hints that life might have access to oxygen earlier than the GOE

(Jabłońska and Tawfik 2021), either extra- or intracellularly [possibly under anaerobic

conditions using alternative electron acceptors as observed in Thaumarchaea (Kraft et al.

2022) and the NC10 group (Ettwig et al. 2010)] but more likely the latter.

Our work shows that there is much more information beyond bacterial fossils for

calibrating bacterial evolution. The estimated timeline of bacterial evolution may well serve

as a foundation for studying the evolution of bacteria, their diversification rate, and its

correlation with Earth's geochemistry. It is hopeful that some of these questions can be

answered, and competing hypothesis can be reconciled by future molecular clock studies in

combination with modern phylogenetics and comparative genomics methods.

Methods

Phylogenomic reconstruction

A total of 62 marker genes considered conserved across bacteria were retrieved from a recent

phylogenomics study of the bacterial tree (Coleman et al. 2021). Their sequences were

aligned using MAFFT v7.22261 (Katoh and Standley 2013) and trimmed using TrimAl v1.4

(settings: "-resoverlap 0.55 -seqoverlap 60") (Capella-Gutiérrez et al. 2009). Unrooted

phylogenomic trees were constructed based on the concatenated alignment with time-

reversible Markov model and each of the three empirical site-heterogeneous profile mixture

models C20, C40, and C60 using a PMSF approximation (guide tree built by LG4M+G+I)

with 1000 ultrafast bootstraps (Minh et al. 2020). The model LG+G+C60 was selected as the

best-fit model according to AIC and BIC, and the tree constructed by this model was used in

the main analysis. Rooted bacterial trees were constructed using the recently developed time-

non-reversible model implemented in IQ-Tree (Naser-Khdour et al. 2022) that took the

unrooted tree as either a fixed tree or a start tree (see Note S2 for details).

Identification of mitochondria-originated genes conserved across the bacterial tree

We retrieved the sequences of the 108 genes identified to have a mitochondrial origin from

the study (Muñoz-Gómez et al. 2022). We identified their homologs in eukaryotes encoded

by either the mitochondrial or nuclear genome based on BLAST with an e-value cut-off of

1e-30. Next, we followed (Liao et al. 2022) to keep 32 genes present in at least one of the

following three gene sets namely bac120 (120 genes) (Parks et al. 2017), Battistuzzi2009 (25

genes) (Battistuzzi and Hedges 2009), and Coleman2021 (62 genes) (Coleman et al. 2021),

which include orthologs well conserved and likely vertically transmitted across bacteria. Note

that two of the 62 orthologs originally included in Coleman2021 correspond to the same id

(K04485), and were both discarded. For each of the remaining 32 genes, we built a gene tree

using IQ-Tree, based on which we further removed i) two genes that did not show monophyly

between mitochondria and α-proteobacteria, indicating unresolved paralogy, and ii) 11 genes

where mitochondria/α-proteobacteria were placed at a basal position to all other bacteria with

different rooting approaches, indicating fast-evolving genes. This returned 19 mitochondria-

originated genes conserved across the bacterial tree, which were used in the main molecular

clock analysis (Table S3; Fig. S7).

Ancestral lifestyle reconstruction

The 16S rRNA genes of each interested symbiont group were collected from the NCBI

Entrez Nucleotide database with the keywords "taxon_name[ORGN] AND 16S[TITL] AND

rrna AND 1000:2000[SLEN] NOT spacer NOT trna" (April 2022). We also retrieved 16S

rRNA genes predicted by RNAmmer v1.2 (Lagesen et al. 2007) from genomes deposited in

Genbank (August 2021). Those without any information of host or isolation source in the

metadata were removed (see SI). All remaining sequences were clustered using CD-HIT v4.7

(Li and Godzik 2006) with 98.7% as the identity cut-off (Yarza et al. 2014) into operational

taxonomic units (OTUs), and one representative was chosen from each OTU (see SI for

alternative strategies in sequence selection). We applied the stochastic character mapping

(SCM) (Huelsenbeck et al. 2003) to directly sample from the joint posterior distribution,

using *make.simmap* from the R package "phytools" v0.2-40 (Revell 2012) to estimate

ancestral hosts (or lifestyles) of modern symbionts. To account for phylogenetic uncertainty,

we applied the so-called most-recent-common-ancestor approach (Pagel et al. 2004) by

identifying the LCA of the group of species of interest, reconstructing the state of that node,

and then summarizing the information across 100 ultrafast bootstrap (UFB) trees

reconstructed by IQ-Tree (settings: -m GTR+G). For each of the 100 UFB trees, 100

simulated stochastic character maps were generated using the MCMC algorithm implemented

in phytools under the "all-rates different" (ARD) model which allows transition rates at all

directions to be different, totalling $100 \times 100 = 10000$ simulations. All ASR analyses were

performed on phylogram—a phylogeny where branch lengths indicate amount of

evolutionary change, built by the 16S rRNA gene, as done in previous studies (Wang et al.

2020; Wang and Luo 2021).

488   Specifically, let S_i be the inferred lifestyle (or state) of the i^{th} internal node of symbionts,

489   which can take any possible hosts observed in the group of modern symbionts (including

490   free-living), denoted as s_i. By SCM, the marginal probability that the i^{th} internal node of

491   symbionts takes the state s_i may be approximated by

492
$$P(S_i = s_i) \approx \frac{n_{\{S_i=s_i\}}}{n},$$
(1)

493   where $n_{\{S_1=s_1\}}$ is the number that node i takes the state s_i during n rounds of SCMs.

494   Likewise, the same approach can be extended to estimate the joint probability that internal

495   nodes 1 to k take states $s_1, \ldots, s_k$ respectively by

496
$$P(S_1 = s_1, \ldots, S_k = s_k) \approx \frac{n_{\{S_1=s_1,\ldots,S_k=s_k\}}}{n},$$
(2)

where $n_{\{S_1=s_1,\ldots,S_k=s_k\}}$ is the number that the event $\{S_1 = s_1, \ldots, S_k = s_k\}$ happens during n

rounds of SCMs.

pRTC informed by host-symbiont association

Denote the data (i.e., sequence alignments) for molecular dating as D, and the divergence

times as $t = (t_1, \ldots, t_k)$ where $t_1, \ldots, t_k$ indicate the divergence time of the k ancestral nodes

of modern symbionts. Let S_i be the inferred lifestyle (or state) of the i^{th} ancestral node of

symbionts, which can take any possible hosts observed in the group of modern symbionts or

free-living, denoted as s_i. Let A indicate the event that the origin time of all symbionts is no

earlier than that of their host, i.e., $A = \{t_1 \leq t_{S_1}, \ldots, t_k \leq t_{S_k}\}$. It follows that

507
$$f(t|D, A) \propto f(t|D)P(A|t, D)$$

508
$$= f(t|D)P(A|t)$$
(3)

where $f(t|D, A)$ is the posterior of divergence times t given sequence data D and the

assumption A, $f(t|D)$ is the posterior of divergence times t given only sequence data D, and

$P(A|t)$ indicates the probability A is satisfied given time t. We further assume that the

inference of ancestral lifestyles of modern symbionts depends on only the phylogenetic

comparative data estimated by SCM and is independent of time. The second term on the

right-hand side of Eq. (3) can be rewritten as

$P(A|t) = P\left(t_1 \leq t_{S_1}, \ldots, t_k \leq t_{S_k} \middle| t\right)$

$= \sum_{s_1,\ldots,s_k} P\left(t_1 \leq t_{S_1}, \ldots, t_k \leq t_{S_k} \middle| t, S_1 = s_1, \ldots, S_k = s_k\right) P(S_1 = s_1, \ldots, S_k = s_k | t)$

$= \sum_{s_1,\ldots,s_k} P\left(t_1 \leq t_{S_1}, \ldots, t_k \leq t_{S_k} \middle| t, s_1, \ldots, s_k\right) P(S_1 = s_1, \ldots, S_k = s_k)$

$= \sum_{s_1,\ldots,s_k} \left(\prod_{i=1}^{k} I\left(t_i, t_{s_i}\right)\right) P(S_1 = s_1, \ldots, S_k = s_k),$

where $I\left(t_i, t_{s_i}\right)$ is defined as

$$I\left(t_i, t_{s_i}\right) = \begin{cases} 1, t_i \leq t_{s_i} \\ 0, t_i > t_{s_i} \end{cases},$$

Particularly, we define $I\left(t_i, t_{fl}\right) = 1$ (free-living ancestral state) since the RTC that

symbiont's age is no older than the host's is automatically satisfied for free-living bacteria.

Further, assuming that the states of the ancestral nodes of symbiotic bacteria are

independent of each other so that $P(S_1 = s_1, \ldots, S_k = s_k) = \prod_{i=1}^{k} P(S_i = s_i)$, which

generally holds if the ancestral nodes are not phylogenetically too closely related (see Fig.

S6B for details), the above can be rewritten as

$$P(A|t) = \prod_{i=1}^{k} \sum_{s_i} I\left(t_i, t_{s_i}\right) P(S_i = s_i). \tag{4}$$

Collectively, by introducing Eq. (4) into Eq. (3), we have

$$f(t|D, A) \propto f(t|D) \left[\prod_{i=1}^{k} \sum_{s_i} I\left(t_i, t_{s_i}\right) P(S_i = s_i)\right]. \tag{5}$$

The above indicates that estimating the posterior dates under the assumption that the

origin time of all symbionts is no earlier than their hosts' while considering the uncertainty in

the inference of ancestral hosts can be obtained by:

i)   Perform ASR to obtain the probability of each ancestral state for selected nodes in the

group of modern symbionts of interest.

ii) Perform MCMC molecular clock analysis with fossil calibrations but without any RTC.

iii) For each MCMC posterior sample collected from Step ii), perform a rejection sampling

by accepting it with a probability according to Eq. (5) (based on the ASR results obtained

from Step i).

Obviously, Step 2 is independent of Step 1, and Step 3 is based on Step 1 and Step 2.

Sequential Bayesian approach to dating the bacterial tree of life

We applied a sequential Bayesian approach (dos Reis et al. 2012; Álvarez-Carretero et al.

2022) to establish a timeline of the bacterial tree, which allows using the genome-scale data

and abundant fossil-based calibrations of eukaryotes to better constrain symbionts' ages

through the pRTC framework (see Supplementary Note 1.3.2 for details). Sequential

Bayesian molecular dating contains two steps: i) run a dating analysis of the first subset of

data and collect the posterior samples of times and rates, ii) approximate the posteriors of

each node in the first-step analysis by a parametric probability distribution and use it as the

prior in a subsequent analysis on a second subset of data (see SI for details). We first used

MCMCTree from PAML v4.10.0 (Yang 2007) (see Note S1.2 for details) to date an

eukaryote tree with 29 selected species based on our previous study (Wang and Luo 2021)

and 320 orthologs (Strassert et al. 2021) with 11 fossil-based calibrations (Note S3.2), then

approximated the posterior samples of the dates obtained in the first-step analysis using a

best-fit probability distribution available in MCMCTree. The approximated dates were used

as the priors to calibrate the bacterial tree using the abovementioned 19 mitochondria-

originated genes conserved across bacteria (no overlap in genes between the first- and the

second-step analysis).

The posterior dates obtained in the above first-step sequential analysis were used

together with the following bacterial calibrations to date the bacterial tree (Note S3.2). The

calibrations from within bacteria were the 1.6 Ga-old akinete fossils for the filamentous

cyanobacteria Nostocales (Tomitani et al. 2006), the 1.7 Ga-old microfossils for the coccoid

cyanobacteria order Pleurocapsales (Zhang and Golubic 1987), the GOE ~2.32 Ga for

photosynthetic cyanobacteria (Luo et al. 2016), and the hydrocarbon biomarkers for green

and purple sulphur bacteria both dated at 1.64 Ga (Brocks et al. 2005). The root minimum

was given as 2.32 Ga, the oldest among the above five. A maximum was set as the age of

Earth (4.5 Ga) for all the above calibrations within bacteria. If not otherwise specified, in all

molecular clock analyses we followed Betts et al. 2018 to apply a uniform distribution as the

prior: a soft tail was applied as the upper calibration bound, which means a small probability

(by default 2.5%) where the age is beyond the bound, and a hard bound was set as the

minimum age meaning that the age cannot be lower than the bound. This makes sense as

minimum bounds are based on solid fossil or biomarker records while the maxima are often

based on indirect evidence (Benton and Donoghue 2007; Marshall 2019).

574   A bootstrap-based approach to improving MCMCTree branch length estimate by

575   incorporating the mixture substitution model

We applied MCMCTree's approximate likelihood method (Thorne et al. 1998; dos Reis and

Yang 2011) which estimated the branch lengths (expected number of amino acid substitutions

per site) in the first step, and in the second step, the branch length estimates, the gradient and

the Hessian matrix are used to calculate the likelihood by the second-order Taylor

approximation to the likelihood function. Different from previous studies which often use the

LG+G substitution model, we applied the bootstrap-based method developed in the present

study to integrate the site-heterogeneous profile mixture model LG+G+C60 (Quang et al.

2008) in MCMCTree analysis. We based our method on the assumption that the maximum

likelihood estimates of branch length follow a multivariate normal (MVN) distribution,

according to the maximum likelihood (ML) estimation theory stating that ML estimators are

asymptotically multivariate normally distributed (Pawitan 2001). The idea was originally

proposed by Thorne et al. 1998 and later implemented in MCMCTree as the approximate

likelihood method (dos Reis and Yang 2011). The approximate likelihood method (dos Reis

and Yang 2011) first obtains the maximum-likelihood estimates (MLEs) of the branch length

$\widehat{\boldsymbol{\theta}} = (\widehat{\theta}_1, \dots, \widehat{\theta}_n)^T$, plus the gradient $\mathbf{g}$ and the Hessian matrix $\mathbf{H}$, which are the first and

second derivatives of the log-likelihood evaluated at $\widehat{\boldsymbol{\theta}}$ respectively. Next, according to Eq.

(1) in the study (dos Reis and Yang 2011), Taylor expansion at the second order derivative

was applied to approximate the log-likelihood

$$\ell(\boldsymbol{\theta}) \approx \ell(\widehat{\boldsymbol{\theta}}) + \mathbf{g}^T \Delta\boldsymbol{\theta} + \frac{1}{2}\Delta\boldsymbol{\theta}^T \mathbf{H}\Delta\boldsymbol{\theta} \tag{6}$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots, \theta_n)^T$ denotes the parameters to estimate (branch length) and $\ell(\boldsymbol{\theta})$ is the

log-likelihood as a function of branch lengths.

While the gradient is zero at the maximum likelihood estimate ($\mathbf{g} = \mathbf{0}$), directly

calculating $\mathbf{H}$ under IQ-Tree's profile mixture and other complex substitution model is

difficult and has not been available. However, note that when the sample size is large enough,

the Hessian matrix evaluated at $\widehat{\boldsymbol{\theta}}$ is approximately the negative inverse matrix of the

covariance matrix of the MLEs $\mathbf{V}(\widehat{\boldsymbol{\theta}})$ (Pawitan 2001):

$$\mathbf{H} \approx -\mathbf{V}(\widehat{\boldsymbol{\theta}})^{-1}. \tag{7}$$

603 Hence, we instead applied a bootstrap method to estimate $\mathbf{V}(\hat{\boldsymbol{\theta}})$ by the bootstrap covariance

604 matrix $\hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}^*)$ (Efron 1982) by taking B bootstrap alignments and estimating the MLEs of

605 branch lengths $\hat{\boldsymbol{\theta}}^*$:

$$\mathbf{V}(\hat{\boldsymbol{\theta}}) \approx \hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}^*) = \frac{1}{B-1} \sum_{b=1}^{B} (\hat{\boldsymbol{\theta}}^{(b)} - \overline{\hat{\boldsymbol{\theta}}^*})(\hat{\boldsymbol{\theta}}^{(b)} - \overline{\hat{\boldsymbol{\theta}}^*})^T \tag{8}$$

where $\hat{\boldsymbol{\theta}}^{(b)}$ indicate the MLEs of the branch lengths in the b^{th} bootstrap sample. Specifically,

1000 bootstraps ($B = 1000$) were performed using the site-heterogeneous profile mixture

model LG+G+Cxx with a fixed tree topology by IQ-Tree, therefore generating 1000

bootstrap MLEs for each branch length and allowing calculating the bootstrap covariance

matrix $\hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}^*)$. Accordingly, the Hessian matrix can be approximated by solving the inverse

matrix of $-\hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}^*)$. We imported the information of branch length estimates, i.e. the MLEs of

branch length, the gradient as a zero matrix, and the Hessian matrix approximated by the

above procedure, into a single file *in.BV*, which was used for time estimation using

MCMCTree's approximate likelihood method (dos Reis and Yang 2011). A series of

simulations were performed to show the feasibility of this approach to integrating profile

mixture substitution model into MCMCTree analysis (see Note S4).

A taxonomically high-resolution timeline of the bacterial tree

A representative genome was downloaded from each genus deposited in the Genome

Taxonomy Database (GTDB; version r95) (Parks et al. 2022). We selected 14 out of the 25

orthologs conserved across prokaryotes identified by a prior study (Battistuzzi and Hedges

2009), the other 11 being removed to avoid squaring of likelihood using the sequential

approach [otherwise the same genes are used twice; see Note S1.2.2 and the studies (dos Reis

et al. 2012; Álvarez-Carretero et al. 2022)]. These genes were then used to construct a

phylogenomic tree based on the first two codon positions of coding sequences using GTR+G

model and the "-fast" mode with IQ-Tree to reduce computational burden. We applied

RelTime implemented in MEGA X (Kumar et al. 2018), which is based on estimating the

relative rates (Tamura et al. 2012), to establish an evolutionary timescale for bacteria at the

genus-level resolution. The posterior dates obtained in the 148-genome dataset were

approximated by a normal (shown in Fig. 4) or uniform (in alternative analysis) distribution.

Because RelTime calculates divergence time by measuring the relative rate between

neighbouring branches (Tamura et al. 2012) which is insensitive to overlaps in time priors

between parent and child nodes (also known as "truncation effect"; see Note S1.2.2), the age

estimates using RelTime may not change much from the specified calibrations. However, to

minimize the impact of the overlap in the calibrations between parent and child nodes, we

alternatively removed those calibrations where more than 10% overlap in the specified

distributions between the parent and child nodes to keep a maximum number of calibrations

without significant overlap with each other (Fig. S15D), and re-performed RelTime analysis

(Fig. S15A-S15C). Taxonomic classification was according to GTDB version r95 (Parks et

al. 2020). Genetic divergence (substitutions/site) between 16S rRNA sequences was

calculated using IQ-Tree under GTR+G+I model.

Acknowledgements

We are particularly grateful to Jianhao Lv for the critical comments on the mathematical

derivation of the formulae, and the professional insights in statistics. We also thank

Daniel Mills and Philip Donoghue for insights in the interpretation of eukaryotic fossils,

Mario dos Reis, José Barba-Montoya and Suha Naser for technical support in phylogenetics

analysis, and Andrew Meade, Stilianos Louca and Liam Revell for guidance in ancestral state

reconstruction. We thank László Nagy, Torda Varga and Tomáš Flouris for kindly sharing

with us software. We also thank Luo Lab members Tianhua Liao and Hao Zhang for

discussion, and PhyloPic (https://www.phylopic.org/) for the free images of organisms. The

work is supported by the Hong Kong Research Grants Council General Research Fund

(14116922), the Natural Science Foundation of China (42293294), and the Hong Kong

Research Grants Council Area of Excellence Scheme (AoE/M-403/16).

References

Abramov O, Mojzsis SJ. 2009. Microbial habitability of the Hadean Earth during the late

heavy bombardment. *Nature* 459:419–422.

Allwood AC, Walter MR, Kamber BS, Marshall CP, Burch IW. 2006. Stromatolite reef from

the Early Archaean era of Australia. *Nature* 441:714–718.

Álvarez-Carretero S, Tamuri AU, Battini M, Nascimento FF, Carlisle E, Asher RJ, Yang Z,

Donoghue PCJ, dos Reis M. 2022. A species-level timeline of mammal evolution

integrating phylogenomic data. *Nature* 602:263–267.

Amard B, Bertrand-Sarfati J. 1997. Microfossils in 2000 Ma old cherty stromatolites of the

Franceville Group, Gabon. *Precambrian Res.* 81:197–221.

Battistuzzi FU, Feijao A, Hedges SB. 2004. A genomic timescale of prokaryote evolution:

Insights into the origin of methanogenesis, phototrophy, and the colonization of land.

BMC Evol. Biol. 4:44.

Battistuzzi FU, Hedges SB. 2009. A major clade of prokaryotes with ancient adaptations to

life on land. *Mol. Biol. Evol.* 26:335–343.

Bengtson S, Sallstedt T, Belivanova V, Whitehouse M. 2017. Three-dimensional preservation

of cellular and subcellular structures suggests 1.6 billion-year-old crown-group red

algae. *PLoS Biol.* 15:e2000735.

Benton MJ, Donoghue PCJ. 2007. Paleontological evidence to date the tree of life. *Mol. Biol.*

Evol. 24:26–53.

Betts HC, Puttick MN, Clark JW, Williams TA, Donoghue PCJ, Pisani D. 2018. Integrated

genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nat.*

Ecol. Evol. 2:1556–1562.

Boehnke P, Harrison TM. 2016. Illusory late heavy bombardments. *Proc. Natl. Acad. Sci. U.*

S. A. 113:10802–10806.

Brocks JJ, Love GD, Summons RE, Knoll AH, Logan GA, Bowden SA. 2005. Biomarker

evidence for green and purple sulphur bacteria in a stratified Palaeoproterozoic sea.

Nature 437:866–870.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25:1972–1973.

Coleman GA, Davín AA, Mahendrarajah TA, Szánthó LL, Spang A, Hugenholtz P, Szöllsi

GJ, Williams TA. 2021. A rooted phylogeny resolves early bacterial evolution. *Science*

(80-.). 372.

Crowe SA, Døssing LN, Beukes NJ, Bau M, Kruger SJ, Frei R, Canfield DE. 2013.

Atmospheric oxygenation three billion years ago. *Nature* 501:535–538.

Davín AA, Tannier E, Williams TA, Boussau B, Daubin V, Szöllosi GJ. 2018. Gene transfers

can date the tree of life. *Nat. Ecol. Evol.* 2:904–909.

Drummond AJ, Ho SYW, Phillips MJ, Rambaut A. 2006. Relaxed phylogenetics and dating

with confidence. *PLoS Biol.* 4:699–710.

Efron B. 1982. The Bootstrap. In: The Jackknife, the Bootstrap and Other Resampling Plans.

p. 36. Available from: https://epubs.siam.org/doi/book/10.1137/1.9781611970319

Ettwig KF, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM, Schreiber F,

Dutilh BE, Zedelius J, De Beer D, et al. 2010. Nitrite-driven anaerobic methane

oxidation by oxygenic bacteria. *Nature*.

Falkowski PG, Fenchel T, Delong EF. 2008. The microbial engines that drive earth's

biogeochemical cycles. *Science (80-.).* 320:1034–1039.

Fournier GP, Moore KR, Rangel LT, Payette JG, Momper L, Bosak T. 2021. The Archean

origin of oxygenic photosynthesis and extant cyanobacterial lineages. *Proc. R. Soc. B*

Biol. Sci. 288:20210675.

Gibson TM, Shih PM, Cumming VM, Fischer WW, Crockford PW, Hodgskiss MSW,

Wörndle S, Creaser RA, Rainbird RH, Skulski TM, et al. 2018. Precise age of

Bangiomorpha pubescens dates the origin of eukaryotic photosynthesis. *Geology*

46:135–138.

Gouy R, Baurain D, Philippe H. 2015. Rooting the tree of life: The phylogenetic jury is still

out. *Philos. Trans. R. Soc. B Biol. Sci.* 370.

He H, Wu X, Xian H, Zhu J, Yang Y, Lv Y, Li Y, Konhauser KO. 2021. An abiotic source of

Archean hydrogen peroxide and oxygen that pre-dates oxygenic photosynthesis. *Nat.*

Commun.

Ho SYW, Duchêne S. 2014. Molecular-clock methods for estimating evolutionary rates and

timescales. *Mol. Ecol.* 23:5947–5965.

Hofmann HJ, Grey K, Hickman AH, Thorpe RI. 1999. Origin of 3.45 Ga coniform

stromatolites in Warrawoona Group, Western Australia. *Bull. Geol. Soc. Am.* 111:1256–

1262.

Huelsenbeck JP, Nielsen R, Bollback JP. 2003. Stochastic Mapping of Morphological

Characters. *Syst. Biol.* 52:131–158.

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN,

Hernsdorf AW, Amano Y, Ise K, et al. 2016. A new view of the tree of life. *Nat.*

Microbiol. 1.

Jabłońska J, Tawfik DS. 2021. The evolution of oxygen-utilizing enzymes suggests early

biosphere oxygenation. *Nat. Ecol. Evol.* 5:442–448.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Mol. Biol. Evol.* 30:772–780.

Knoll AH. 2015. Paleobiological perspectives on early microbial evolution. *Cold Spring*

Harb. Perspect. Biol. 7:1–17.

Kraft B, Jehmlich N, Larsen M, Bristow LA, Könneke M, Thamdrup B, Canfield DE. 2022.

Oxygen and nitrogen production by an ammonia-oxidizing archaeon. *Science (80-.).*

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular evolutionary

genetics analysis across computing platforms. *Mol. Biol. Evol.* 35:1547–1549.

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. *Mol. Biol. Evol.* 34:1812–1819.

Kuo CH, Ochman H. 2009. Inferring clocks when lacking rocks: The variable rates of

molecular evolution in bacteria. *Biol. Direct* 4:35.

Lagesen K, Hallin P, Rødland EA, Stærfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer:

Consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35:3100–

3108.

Li WZ, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of

protein or nucleotide sequences. *Bioinformatics* 22:1658–1659.

Liao T, Wang S, Stüeken EE, Luo H. 2022. Phylogenomic Evidence for the Origin of

Obligate Anaerobic Anammox Bacteria Around the Great Oxidation Event. *Mol. Biol.*

Evol. 39:msac170.

Louca S, Shih PM, Pennell MW, Fischer WW, Parfrey LW, Doebeli M. 2018. Bacterial

diversification through geological time. *Nat. Ecol. Evol.* 2:1458–1467.

Luo G, Ono S, Beukes NJ, Wang DT, Xie S, Summons RE. 2016. Rapid oxygenation of

Earth's atmosphere 2.33 billion years ago. *Sci. Adv.* 2.

Magnabosco C, Moore KR, Wolfe JM, Fournier GP. 2018. Dating phototrophic microbial

lineages with reticulate gene histories. *Geobiology* 16:179–189.

Marin J, Battistuzzi FU, Brown AC, Hedges SB. 2017. The Timetree of Prokaryotes: New

Insights into Their Evolution and Speciation. *Mol. Biol. Evol.* 34:437–446.

Marshall CR. 2019. Using the Fossil Record to Evaluate Timetree Timescales. *Front. Genet.*

10:1049.

Martijn J, Vosseberg J, Guy L, Offre P, Ettema TJG. 2018. Deep mitochondrial origin outside

the sampled alphaproteobacteria. *Nature* 557:101–105.

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. *Nat.*

Rev. Microbiol. 10:13–26.

McLoughlin N, Wilson LA, Brasier MD. 2008. Growth of synthetic stromatolites and wrinkle

structures in the absence of microbes - Implications for the early fossil record.

Geobiology 6:95–105.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,

Lanfear R, Teeling E. 2020. IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* 37:1530–1534.

Moody ERR, Mahendrarajah TA, Dombrowski N, Clark JW, Petitjean C, Offre P, Szöllősi

GJ, Spang A, Williams TA. 2022. An estimate of the deepest branches of the tree of life

from ancient vertically evolving genes. *Elife* 11:e66695.

Moran NA, Munson MA, Baumann P, Ishikawa H. 1993. A molecular clock in

endosymbiotic bacteria is calibrated using the insect hosts. *Proc. R. Soc. B Biol. Sci.*

253:167–171.

Muñoz-Gómez SA, Susko E, Williamson K, Eme L, Slamovits CH, Moreira D, López-García

P, Roger AJ. 2022. Site-and-branch-heterogeneous analyses of an expanded dataset

favour mitochondria as sister to known Alphaproteobacteria. *Nat. Ecol. Evol.* 6:253–

262.

Naser-Khdour S, Quang Minh B, Lanfear R. 2022. Assessing Confidence in Root Placement

on Phylogenies: An Empirical Study Using Nonreversible Models for Mammals. *Syst.*

Biol. 71:959–972.

Nisbet EG, Sleep NH. 2001. The habitat and nature of early life. *Nature* 409:1083–1091.

Ochman H, Elwyn S, Moran NA. 1999. Calibrating bacterial evolution. *Proc. Natl. Acad. Sci.*

U. S. A. 96:12638–12643.

Pagel M, Meade A, Barker D. 2004. Bayesian estimation of ancestral character states on

phylogenies. *Syst. Biol.* 53:673–684.

Parfrey LW, Lahr DJG, Knoll AH, Katz LA. 2011. Estimating the timing of early eukaryotic

diversification with multigene molecular clocks. *Proc. Natl. Acad. Sci. U. S. A.*

108:13624–13629.

Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ, Hugenholtz P. 2020. A

complete domain-to-species taxonomy for Bacteria and Archaea. *Nat. Biotechnol.*

Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil PA, Hugenholtz P. 2022. GTDB:

An ongoing census of bacterial and archaeal diversity through a phylogenetically

consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Res.*

50:D785–D794.

Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, Hugenholtz P,

Tyson GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes

substantially expands the tree of life. *Nat. Microbiol.* 2:1533–1542.

Pawitan Y. 2001. In all likelihood: statistical modelling and inference using likelihood.

Oxford University Press

Perreau J, Moran NA. 2021. Genetic innovations in animal–microbe symbioses. *Nat. Rev.*

Genet.

Planavsky NJ, Asael D, Hofmann A, Reinhard CT, Lalonde S V, Knudsen A, Wang X, Ossa

Ossa F, Pecoits E, Smith AJB, et al. 2014. Evidence for oxygenic photosynthesis half a

billion years before the Great Oxidation Event. *Nat. Geosci.* 7:283–286.

Quang LS, Gascuel O, Lartillot N. 2008. Empirical profile mixture models for phylogenetic

reconstruction. *Bioinformatics* 24:2317–2323.

Ravenhall M, Škunca N, Lassalle F, Dessimoz C. 2015. Inferring Horizontal Gene Transfer.

PLoS Comput. Biol. 11:e1004095.

dos Reis M, Inoue J, Hasegawa M, Asher RJ, Donoghue PCJ, Yang Z. 2012. Phylogenomic

datasets provide both precision and accuracy in estimating the timescale of placental

mammal phylogeny. *Proc. R. Soc. B Biol. Sci.* 279:3491–3500.

dos Reis M, Yang Z. 2011. Approximate likelihood calculation on a phylogeny for Bayesian

Estimation of Divergence Times. *Mol. Biol. Evol.* 28:2161–2172.

Revell LJ. 2012. phytools: An R package for phylogenetic comparative biology (and other

things). *Methods Ecol. Evol.* 3:217–223.

Sánchez-Baracaldo P, Cardona T. 2020. On the origin of oxygenic photosynthesis and

Cyanobacteria. *New Phytol.* 225:1440–1446.

Sánchez-Baracaldo P, Raven JA, Pisani D, Knoll AH. 2017. Early photosynthetic eukaryotes

inhabited low-salinity habitats. *Proc. Natl. Acad. Sci.*

Schenk JJ. 2016. Consequences of secondary calibrations on divergence time estimates. *PLoS*

One 11:e0148228.

Schopf JW. 2011. The paleobiological record of photosynthesis. *Photosynth. Res.* 107:87–

101.

Shih PM, Hemp J, Ward LM, Matzke NJ, Fischer WW. 2017. Crown group

Oxyphotobacteria postdate the rise of oxygen. *Geobiology* 15:19–29.

Strassert JFH, Irisarri I, Williams TA, Burki F. 2021. A molecular timescale for eukaryote

evolution with implications for the origin of red algal-derived plastids. *Nat. Commun.*

12:1879.

Szöllõsi GJ, Höhna S, Williams TA, Schrempf D, Daubin V, Boussau B. 2022. Relative Time

Constraints Improve Molecular Dating. *Syst. Biol.* 71:797–809.

Tamura K, Battistuzzi FU, Billing-Ross P, Murillo O, Filipski A, Kumar S. 2012. Estimating

divergence times in large molecular phylogenies. *Proc. Natl. Acad. Sci. U. S. A.*

Thorne JL, Kishino H, Painter IS. 1998. Estimating the rate of evolution of the rate of

molecular evolution. *Mol. Biol. Evol.* 15:1647–1657.

Toft C, Andersson SGE. 2010. Evolutionary microbial genomics: Insights into bacterial host

adaptation. *Nat. Rev. Genet.* 11:465–475.

Tomitani A, Knoll AH, Cavanaugh CM, Ohno T. 2006. The evolutionary diversification of

cyanobacteria: Molecular-phylogenetic and paleontological perspectives. *Proc. Natl.*

Acad. Sci. U. S. A. 103:5442–5447.

Waite DW, Chuvochina M, Pelikan C, Parks DH, Yilmaz P, Wagner M, Loy A, Naganuma

T, Nakai R, Whitman WB, et al. 2020. Proposal to reclassify the proteobacterial classes

deltaproteobacteria and oligoflexia, and the phylum thermodesulfobacteria into four

phyla reflecting major functional capabilities. *Int. J. Syst. Evol. Microbiol.* 70:5972–

6016.

Wang S, Luo H. 2021. Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nat.*

Commun. 12:3324.

Wang S, Meade A, Lam H-M, Luo H. 2020. Evolutionary Timeline and Genomic Plasticity

Underlying the Lifestyle Diversity in Rhizobiales. *mSystems* 5:e00438-20.

Wang X, Planavsky NJ, Hofmann A, Saupe EE, De Corte BP, Philippot P, LaLonde S V.,

Jemison NE, Zou H, Ossa FO, et al. 2018. A Mesoarchean shift in uranium isotope

systematics. *Geochim. Cosmochim. Acta* 238:438–452.

Yang Z. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.*

24:1586–1591.

Yang Z, Rannala B. 2006. Bayesian estimation of species divergence times under a molecular

clock using multiple fossil calibrations with soft bounds. *Mol. Biol. Evol.*

Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer KH, Whitman WB,

Euzéby J, Amann R, Rosselló-Móra R. 2014. Uniting the classification of cultured and

857 uncultured bacteria and archaea using 16S rRNA gene sequences. *Nat. Rev. Microbiol.*

858 12:635–645.

859 Zhang Y, Golubic S. 1987. Endolithic microfossils (cyanophyta) from early Proterozoic

860 stromatolites, Hebei, China. *Acta Micropalaeontologica Sin.* 4:1–3.

861 Zhu Q, Mai U, Pfeiffer W, Janssen S, Asnicar F, Sanders JG, Belda-Ferre P, Al-Ghalith GA,

862 Kopylova E, McDonald D, et al. 2019. Phylogenomics of 10,575 genomes reveals

863 evolutionary proximity between domains Bacteria and Archaea. *Nat. Commun.* 10.

864 Zuckerkandl E, Pauling L. 1965. Evolutionary Divergence and Convergence in Proteins. In:

865 Evolving Genes and Proteins. p. 97–166.

 Figure legends

Figure 1. Graphical illustration of the two new methods in molecular dating developed in the

present study. (A) Procedure of the pRTC dating informed by host-symbiont association.

Suppose that there are two symbiotic lineages, Symbiont1 from non-Proteobacteria (blue) and

Symbiont2 from α-Proteobacteria (purple), and most of their members are mammal

symbionts (orange) but some are free-living (yellow). First, ASR is performed on the

phylogram of 16S rRNA tree to obtain the probability of each ancestral state taken by the

LCA of the two "symbiont" clades respectively, i.e., $P(S_1)$ and $P(S_2)$. Second, a phylogeny

consisting of symbionts and mammals is constructed by mitochondria-originated genes, and

MCMCTree is run with fossil calibrations but without RTCs. Third, rejection sampling is

performed to remove posterior timetrees obtained in the previous step based on the

requirement that symbiont's origin must postdate the host. Specifically, we require that the

age of the crown group of the symbionts (node1 and node2) no older than that of the total-

group mammals (node3), weighted by $P(S_i = mammal)$. The use of mammals' total group

is because there could be stem-group mammals that are unsampled or go extinct but that have

Symbiont1 or Symbiont2 as symbiont. For the free-living state, we define $I(t_i, t_{fl}) = 1$ as

the RTC is automatically satisfied for free-living bacteria (see Methods). In the final timetree,

the divergence time of symbiont lineages becomes younger than that inferred before host-

association is considered, suggesting that incorporating RTC informed by host-association

improves the molecular clock inferences. (B) Overflow of the bootstrap-based approach to

incorperating branch length estimates inferred under site-heterogeneous profile mixture

substitution models (LG+G+Cxx). $\mathbf{\theta} = (\theta_1, \theta_2, ..., \theta_n)^T$ denotes the parameters to estimate

(branch length, i.e., expected number of amino acid substitutions per site) and $\ell(\mathbf{\theta})$ is the log-

likelihood as a function of branch lengths. Briefly, the maximum-likelihood estimate (MLE)

of branch length $\widehat{\mathbf{\theta}}$ is obtained by LG+G+Cxx, and the bootstrap covariance matrix of branch

lengths $\mathbf{V}(\hat{\boldsymbol{\theta}}^*)$ is calculated to approximate the hessian matrix (second-order derivative of the

log-likelihood of branch length), which are further used for time estimation by MCMCTree.

Figure 2. An evolutionary timeline of the bacterial tree of life established by the three-step

pRTC procedure. The root position is determined using IQ-Tree's non-reversible substitution

model in maximum-likelihood phylogenomic tree reconstruction using the 19 mitochondria-

originated genes. Node bars indicate the 95% HPD interval of age estimates. Nodes with

orange circles indicate the bacterial calibration points (including the root). Eukaryotic

calibrations obtained from sequential Bayesian dating are given in Fig. S4. RTCs of the 28

bacterial symbionts and five eukaryotic hosts (LCA of eukaryotes, total-group animals, total-

group ciliates, total-group insects, crown-group fungi) are denoted by pink and yellow

circles, respectively. Because the 28 symbionts are selected from "symbiont" clades, a total

of 28-9=19 internal nodes serve as the RTCs for bacteria. The organism images are placed at

the origin times of eukaryotes of interest co-estimated with bacterial lineages. Detailed

posterior ages for clades of interest are summarised in Table S4. DST: Deinococcota,

Synergistota, and Thermotogota; Armat-Eremi: Armatimonadetes and Eremiobacteraeota.

CPR: Candidate phyla radiation; FCB: Fibrobacterota, Chlorobiota, Bacteroidota; PVC:

Planctomycetota, Verrucomicrobiota, Chlamydiota, and related lineages; ACD: Aquificota,

Campylobacterota, and Deferribacterota. MBDD (δ-Proteobacteria): Myxococcota,

Bdellovibrionota, Desulfomonadota, and Desulfobacterota. The images of the choanozoan,

and the cyanobacteria and mitochondria are credited to Urutseg and Kevin Song respectively

under CC BY-SA 3.0 (https://creativecommons.org/licenses/by-sa/3.0/) with slight

modifications. All other organism images are distributed under CC0.

Figure 3. Assessing the uncertainty in Bayesian molecular clock analysis. (A) Comparison of

the posterior mean ages (lower triangle; unit: Ma) and rates (upper triangle; unit: number of

substitutions per site per Ga) of bacteria estimated with different strategies. *Strategy 1*

(traditional strategy used in prior studies): five bacterial calibrations with hard minimum

bounds and a soft maximum bound <4.5 Ga at LBCA, no eukaryote timing information,

substitution model LG+G, no RTCs. *Strategy 2*: bacterial calibrations, eukaryote times by

sequential molecular dating, substitution model LG+G, no RTCs. *Strategy 3*: bacterial

calibrations, eukaryote times by sequential molecular dating, substitution model LG+G+C60,

no RTCs. Focal: bacterial calibrations, eukaryote times by sequential molecular dating,

substitution model LG+G+C60, with RTCs. (B) Posterior mean ages of the selected clades

calculated under root maximum prior ages from 4500 to 6000 Ma with the focal strategy

(dashed line) and Strategy 1 (solid line). (C) Divergence times of bacteria estimated by

alternative schemes (*y*-axis; Data S3) versus the one in the focal strategy (*x*-axis). The blue

bars denote the 95% HPD. *Euk1*: the minimum time bound of crown-group red algae

alternatively set as 1.6 Ga based on *Rafatazmia*; *Euk2*: the minimum time bound of animal

crown group alternatively set as 0.89 Ga (Note S3.2); *noCyanoFossil*: the two cyanobacteria

minimum time bounds (total-group Nostocales and Pleurocapsales) removed; *noBiomarker*:

the two bacterial biomarker minimum time bounds (total-group Chromatiaceae and Chlorobi)

removed; *noGOE*: the minimum time bound based on GOE removed. *Joint_prob*: rejection

sampling based on calculating the joint probability (see Fig. S6B) in Eq. (2). *Alt_lifestyle*:

ASR inferred with alternative classification of lifestyles of modern symbionts (Note S3.1).

OTU97: ASR performed with OTU at a cut-off of 97% sequence identity. *ASR_ML*:

maximum likelihood algorithm instead of MCMC used in ASR. *IR*: the independent rate (IR)

model is used instead of the auto-correlated rate (AR) model. (D) Changes in the posterior

dates shown as 95% HPD interval resulting from different root positions of the

phylogenomics tree (*root1*, *root2*, and *rootCPR*), and the use of different genes. *rate_top* and

deltaLL_top: the 5, 10, and 15 (best) top-ranking genes according to relative rate difference

between mitochondrial and bacterial lineages or △LL which measures the the degree of

species- and gene-tree incongruence (Note S1.2.4). *all_genes*: molecular dating performed on

all 32 mitochondria-originated genes conserved across bacteria. The centre is the posterior

mean age. FA: Firmicutes and Actinomycetes. MBDD (δ-Proteobacteria): Myxococcota,

Bdellovibrionota, Desulfomonadota, and Desulfobacterota.

Figure 4. A high-taxonomy resolution timetree of the bacterial tree of life. (A) Divergence

times of the 8898-tip bacterial tree, consisting of a randomly selected representative from all

genera according to GTDB r95, estimated using RelTime and 118 calibrations (black dots),

i.e., posterior dates calculated from the 148-genome molecular clock analysis. Ma: million

years ago. (B) The divergence times at a family-, order-, class-, and phylum level inferred

from the current study. Only families, orders, classes, and phyla with at least five members

are considered. (C) A simple linear regression analysis of the relationship between 16S rRNA

divergence (number of substitutions per site; *y*-axis) and absolute divergence time (*x*-axis) at

a family level. The intercept is fixed to zero. Shaded areas indicate the 95% confidence

interval of the estimated slope. For each family with at least five members, the largest

difference in 16S sequence between any of its members is considered the 16S divergence of

the family. The analysis is not performed at a higher taxonomic level as 16S sequence likely

reaches saturation due to the large divergence time. Detailed results of this simple linear

regression and alternative linear models are provided in Data S5 and Note S1.4.

A

Step 1

Reconstruction of ancestral lifestyle (host)

$P(S_1 = mammal) = 0.9$ $P(S_1 = fl) = 0.1$ $P(S_2 = mammal) = P(S_2 = fl) = 0.5$

Step 2

- ■ symbiont1   ■ symbiont2
- ■ other bacteria   ■ free-living (fl)
- ■ mammal   ■ bird

Proteobacteria
Fossil

time

posterior timetrees

time

Step 3

$$\prod_{i=1}^{k} \sum_{s_i} I(t_i, t_{s_i}) \, P(S_i = s_i)$$

$\overset{\text{Pr(accept)}}{}$

$$I(t_i, t_{s_i}) = \begin{cases} 1, t_i \leq t_{s_i} \\ 0, t_i > t_{s_i} \end{cases}$$

(0.9*1+0.1*1)*
(0.5*1+0.5*1)=1

(0.9*1+0.1*1)*
(0.5*0+0.5*1)=0.5

time

final timetrees

time

B

original
alignment

bootstrapped
alignment

bootstrap MLEs of
branch lengths $\hat{\boldsymbol{\theta}}^*$
by Cxx substitution model

$\hat{\boldsymbol{\theta}}^{(1)}$

$\hat{\boldsymbol{\theta}}^{(2)}$

$\hat{\boldsymbol{\theta}}^{(B)}$

bootstrap
estimate of $\mathbf{v}(\hat{\boldsymbol{\theta}})$

$$\mathbf{v}(\hat{\boldsymbol{\theta}}) \approx \frac{\sum_{b=1}^{B}(\hat{\boldsymbol{\theta}}^{(b)} - \overline{\hat{\boldsymbol{\theta}}^*})(\hat{\boldsymbol{\theta}}^{(b)} - \overline{\hat{\boldsymbol{\theta}}^*})^T}{B-1}$$

MLEs of branch
lengths $\hat{\boldsymbol{\theta}}$ by Cxx
substitution model

$\mathbf{H} \approx -\mathbf{V}(\hat{\boldsymbol{\theta}})^{-1}$

$\mathbf{g} = \mathbf{0}$

$$\ell(\boldsymbol{\theta}) \approx \ell(\hat{\boldsymbol{\theta}}) + \mathbf{g}^T \Delta\boldsymbol{\theta} + \frac{1}{2}\Delta\boldsymbol{\theta}^T \mathbf{H} \Delta\boldsymbol{\theta}$$

MCMCTree
time estimate

time